1

Title: Enhanced protein synthesis is a defining requirement for neonatal B cell

development

Running title: Early-life B cell output depends on enhanced protein synthesis

Hugo Åkerstrand*, Elena Boldrin*, Giorgia Montano*, Stijn Vanhee*, Karin Olsson*, Niklas

Krausse*, Stefano Vergani*, Maciej Cieśla†, Cristian Bellodi†, Joan Yuan*

* Developmental Immunology Unit, Department of Molecular Hematology, Lund Stem Cell

Center, Lund University, Lund, Sweden

† RNA and Stem Cell Biology Unit, Department of Molecular Hematology, Lund Stem Cell

Center, Lund University, Lund, Sweden

Corresponding author: +46462220430, joan.yuan@med.lu.se

Abstract

Protein synthesis is a stringently regulated process that fundamentally dictates capacity for

cellular growth. Rapid fluctuations in cell size and proliferation characterize B cell

development. However, little is known about how ribosomal control may impact this process.

Here, we identify elevated protein synthesis as a defining feature of early-life B cell

development and a determinant for the output of self-reactive CD5+ B-1 cells in mice.

Interactome analyses in primary B cell precursors showed direct binding of the heterochronic

RNA binding protein LIN28B to ribosomal protein transcripts, which correlated with their

elevated protein expression *in vivo*. Genetic perturbations demonstrated that LIN28B is both

necessary and sufficient to elevate protein synthesis during the small Pre-B and immature B

cell stages. Interestingly, IL-7 signaling was the dominant driver of the c-MYC/ribosome axis

in earlier stages and masked LIN28B induced protein synthesis in Pro-B cells. Finally, we

demonstrated that subdued protein synthesis was detrimental for neonatal B cell development

and the output of self-reactive B-1 cells, without affecting the adult. Thus, LIN28B elevates

ribosomal capacity to shape the naïve B cell repertoire in neonatal life.

Introduction

B-cell development is centered around alternating stages of cellular proliferation and V(D)J recombination at the immunoglobulin heavy and light chain loci, with the goal of producing a diverse yet self-tolerant B cell repertoire. During the Pro-B cell stage, productive IgH rearrangement results in the expression of a pre-BCR complex and differentiation into the large Pre-B cell stage. The latter is marked by a transient burst of clonal expansion. Cell cycle exit is accompanied by the initiation of light chain recombination and entry into the small Pre-B cell stage. IL-7 signaling is an essential trophic factor during the Pro- and large Pre-B cell stages, promoting cell survival, proliferation, and differentiation, through distinct effects mediated by the Stat5 and mTORC1/c-MYC signaling axes (1, 2). IL-7 signaling wanes as large Pre-B cells move away from IL-7 producing stromal cells in vivo and its attenuation is required for the developmental progression into the small Pre-B cell stage and the onset of IgK light chain recombination (3). Generation of a productive immunoglobulin heavy and light chain complex leads to membrane IgM expression and developmental progression into the immature B (Imm-B) cell stage. Here, cells carrying innocuous BCRs are allowed to mature while self-reactive specificities are subject to central tolerance mechanisms including receptor editing and apoptosis to remove potentially harmful recognition of "self" (4). Progression through these stages is accompanied by dramatic fluctuations in bioenergetic states. Although metabolic capacity has been proposed to modulate the central tolerance checkpoint (5), this theory has not been empirically proven.

Much of what we know about B cell development comes from studies in adult mice, which differs significantly from fetal and neonatal life. For example, while B cell development in adult bone marrow strictly requires IL-7 mediated signaling, B cell output can take place in neonatal mice lacking IL-7Ra until around two weeks of age (6). Furthermore, both the pre-

BCR and mature BCR checkpoints are less stringent during neonatal life allowing for the

positive selection of poly and self-reactive specificities in mice and men (7-9). This correlates

with the output of CD5+ B-1 cells in mice, which secrete natural antibodies important for the

clearance of cellular debris and act as a first line of defense against pathogens (10). Despite

these differences, the basis for the developmental switch in B cell output is not well understood.

One molecular program that distinguishes early life lymphopoiesis is centered around the post-

transcriptional regulator LIN28B. It is one of two mammalian homologs of *lin-28* first

identified in *Caenorhabditis elegans* to encode a heterochronic RNA binding protein in control

of the timing of larval development (11). In the mammalian hematopoietic system, *Lin28b*

expression is abundant during fetal life and gradually silenced within the first two to three

weeks of life, critically controlling a neonatal switch in hematopoietic stem cell function and

lineage output patterns (12). Like its mammalian paralog *Lin28a*, *Lin28b* plays an established

role in regulating organismal growth and cellular metabolism, most famously through the direct

inhibition of the biogenesis of the evolutionarily conserved Let-7 family of microRNAs.

LIN28A/B expression de-represses Let-7 targets which include the proto-oncogenes *Myc*,

Arid3a, *Hmga2* and *Igf2bp1-3* (13-15). We and others have previously shown that ectopic

LIN28A/B expression in the adult promotes key aspects of fetal-like lymphopoiesis, including

the efficient production of CD5+ B-1 cells (13, 16-18). Furthermore, studies from both BCR

poly-clonal and transgenic mouse models make clear that LIN28B relaxes the central tolerance

checkpoint to allow for the positive selection of self-reactive BCR specificities and thereby

uniquely shapes the neonatal B cell repertoire (19, 20). The mechanism for this effect is,

although not fully understood, at least in part linked to augmentation of c-MYC protein levels

in Imm-B cells and the CD19/PI3K/c-MYC positive feedback loop (19) known to promote B

cell positive selection (21).

A less understood mechanism of LIN28B action is through the direct binding of coding

mRNAs to either positively or negatively regulate transcript stability and protein translation,

in a context dependent manner (22-31). In particular, several studies have demonstrated the

ability of LIN28B to promote the translation of ribosomal proteins, through Let-7 independent

direct binding to their transcripts in human ES cells (32), transformed human kidney (22) and

neuroblastoma (31) cell lines. However, the role of the protein synthesis pathway in LIN28B

dependent B cell positive selection is not known. Here we demonstrate that LIN28B directly

interacts with ribosomal protein transcripts in primary B cell precursors. Importantly, LIN28B

expression is necessary and sufficient to elevate protein synthesis specifically during the later

stages of IL-7 independent bone marrow B cell maturation. Finally, we show that an augmented

mode of protein synthesis is a hallmark of neonatal B cell development and a pre-requisite for

the positive selection of self-reactive Imm-B cells and the output of CD5+ B-1 cells early in

life. We propose that the post-transcriptional actions of LIN28B cooperates with the

transcriptional regulation of ribosomal components by c-MYC to augment protein synthesis

and support neonatal B lymphopoiesis.

Materials and methods

Mice

tet-LIN28B mice were generated by intercrossing $Col1a^{tetO-LIN28B}$ (The Jackson Laboratory,

#023911) to $Rosa26^{rtTA*m2}$ (The Jackson Laboratory, #006965) mice and were originally

obtained from the laboratory of George Daley (Harvard Medical School) (33).

Transheterozygous mice were used for experiments. $Rosa26^{rtTA*m2}$ heterozygous littermates

were used as control mice. Doxycycline chow (200 mg/kg, Ssnif, cat#A112D70203) was fed

to tet-LIN28B and littermates for 10 days before analysis. B6.Cg-$Lin28b^{tm1.1Gqda}$/J ($Lin28b$ KO)

(The Jackson Laboratory, #023917) and C57BLKS-$Rpl24^{Bst}$/J (Bst) (The Jackson Laboratory,

#000516) were originally purchased from the Jackson laboratory. Wildtype mice were

originally purchased from Taconic (B6NTAC, Taconic). The $Lin28b$-eGFP knock-in strain was

generated as previously described (34). Adult mice were used at 10 to 16 weeks of age,

neonates were analyzed at the indicated days post-birth. Male and female mice were used

interchangeably throughout the experiments. Animal husbandry and experimental procedures

were performed in accordance with ethical permits issued by the Swedish Board of Agriculture.

Bone marrow cultures

Adult bone marrow B cell precursors were expanded and enriched *ex vivo*, as previously

described (35). Briefly, single cell suspension of red blood cell lysed bone marrow was plated

in pre-warmed RPMI-1640 (Fischer Scientific, cat#21875091) and incubated for 15 min at 37

ºC. The non-adherent cells were isolated and seeded at a cell density of 0.75-1.5 million/mL

using fully supplemented RPMI-1640, IL-7 (20 ng/mL, Peprotech, cat#217-17), and

doxycycline (0.1 µg/ml, Sigma-Aldrich, cat#D3072). The cell culture medium was refreshed

every second day.

RNA Immunoprecipitation Sequencing (RIP-seq)

Cells from day 5 bone marrow cultures from tet-LIN28B or littermate control mice were used

for lysis and RIP using Magna RIP RNA-Binding Protein Immunoprecipitation kit (Millipore,

cat#17-700) according to the manufacturer's instructions. Briefly, 10% of the total lysate was

kept as input sample. 4 μg of anti-Flag antibody (Sigma) and 50 μl of A/G beads (Millipore,

cat#16-663) were added to the lysate and incubated for 3 hours. Beads were washed 6 times

with ice-cold RIP wash buffer (Millipore, cat#17-700). RNA was extracted using RNAzol

(Sigma-Aldrich, cat#R4533) and Direct-zol RNA MicroPrep kit (Zymo Research, cat#R2060).

For RNA quantification and quality control respectively, Qubit with RNA HS kit

(Thermofisher Scientific, cat# Q32852) and Agilent Bioanalyzer with RNA Pico kit (Agilent,

cat#5067-1513) were used according to the manufacturer's instructions. 20ng of total RNA

was used for the cDNA synthesis and amplification with the SMART-Seq v4 Ultra Low Input

RNA Kit for Sequencing (Takara, cat#634889) according to instructions. Library tagmentation

and indexing was performed using Nextera XT DNA Kit (Illumina, cat#FC-121-1030) and

Nextera XT v2 set A Indexing kit (Illumina, cat#15052163). Library quantification and library

size were measured using Qubit DNA HS kit (Thermofisher Scientific, cat#Q32851) and

BioAnalyzer DNA HS kit (Agilent, cat#5067-4626). The samples were run in NextSeq 550

using NextSeq High output 150 cycles 2x76bp. Reads were aligned using STAR aligner v2.7.1

to mm10 and counted using RSEM v1.28. Genes were filtered for an FPKM of at least 50 in

total tet-LIN28B input samples and following LIN28B immunoprecipitation. Enrichment was

calculated as IP over Input FPKM for each of the two tet-LIN28B samples individually. 238

transcripts had an average enrichment of at least 2-fold and were considered for subsequent

analyses. Littermate control sample was not considered when generating this list. Gene

ontology analysis was done using the Panther database (pantherdb.org, release 17.0). Data is

accessible through (GEO number, data is submitted and being processed).

RNA sequencing

RNA sequencing was performed on sorted adult bone marrow large Pro-B, large Pre-B, small

Pre-B, or Imm-B cell subsets. Data is accessible through (GEO number, data is submitted and

being processed) except for Imm-B that was previously generated (19) and is accessible

through GSE135603. All data was processed and analyzed as previously described (19). GSEA

used Wald's stat as ranking metric.

Metabolic labeling of nascent proteins

Cells were depleted of endogenous methionine by pre-culturing them in methionine free RPMI

(Fisher Scientific, cat#A1451701). After 45 minutes, the medium was supplemented with L-

azidhomoalanine (AHA, 1 mM final concentration, Fisher Scientific, cat#C10102) and labeled

for one hour. AHA uptake was visualized using Click-iT Plus Alexa Fluor 555 Picolyl Azide

Toolkit (Fisher Scientific, cat#C10642), following the manufacturer's protocol. Briefly, up to

3×10^6 cells were fixed in 400 µL of 4% formaldehyde (Fischer Scientific, cat#28906) for 20

minutes at room temperature, washed, and then permeabilized in equal amounts of 0.1%

saponin (Sigma, cat#S4521) for 20 minutes. Alexa Fluor 488 Alkyne (Fisher Scientific,

cat#A10267) was used in place of Alexa Fluor 555 for FACS analysis.

Flow cytometry

Bone marrow cells were extracted by crushing bones from hind- and front limbs, hips, and

sternum, using mortar and pestle. Peritoneal cavity was isolated by flushing adult mice with 8

mL of FACS buffer (Hank's Balanced Salt Solution (Gibco, cat#14175-053) supplemented

with 0.5% Bovine Serum Albumin (Sigma, cat#A9647) and 2 mM EDTA (Invitrogen,

cat#15575-038)), or neonates 1-2 mL depending on the age. Red blood cell lysis was performed

on bone marrow and spleens using ACK lysis buffer (Fischer Scientific, cat#A1049201), only

for mice that were at least 19 days old. Lineage depletion was carried out before cell sorting by

MACS Cell Separation and LS columns (Miltenyi Biotec, cat#130-042-401) to deplete CD3+,

TER119+, or Gr-1+ cells, according to manufacturer's instructions.

Antibody staining of surface antigen was performed using FACS buffer, at a cell density of 1

x 10^7 cells per 100 µL of FACS buffer with antibodies, for 30 minutes at 4 °C. Analysis of

intracellular antigens was done by using BD Cytofix/Cytoperm fixation/permeabilization kit

(BD Biosciences, cat#554714), using a maximum of 3 x 10^6 cells per reaction. All flow

cytometry experiments were gated on live CD19+ B cells and the following

immunophenotypes: Pro-B cells (CD93+IGM–cKIT+), Pre-B cells (CD93+IGM–cKIT– and

FSC-A to separate large from small), Imm-B cells (CD93+IGM+), peritoneal cavity B-1a B

cells (CD5+CD23–B220low and CD43+), and peritoneal cavity B-2 B cells (B220high CD23+

and CD43–). All flow cytometry experiments were performed at the Lund Stem Cell Center

FACS Core Facility (Lund University) using BD Fortessa, BD Fortessa X20, BD FACS Aria

III, or BD FACS Aria II instruments. Cell sorting was performed using a 70 µm nozzle, 0.32.0

precision mask, and a flow rate below 6000 events/s. Analysis was performed using FlowJo

version 10. Antibodies are listed in Supplemental table S1.

In vivo treatment with Rapamycin

Rapamycin (Enzo LifeSciences, cat#BML-A275-0005) was dissolved in Ethanol (99.5%,

Solveco, cat#11594139) and stocked at a concentration of 10 mg/mL. Further dilution was

done in sterile DPBS (Cytiva, cat#SH30028.02) before administering a single intraperitoneal

injection of Rapamycin (2.5 mg/kg) or vehicle control (36).

Western Blot

194
Sorted B cells were lysed in RIPA buffer (CST, cat#9806S), protease inhibitor (Roche,
cat#4693132001), and PMSF (CST, cat#8553S). Loading samples were prepared with
Laemmli buffer (Bio-Rad, cat#1610747) with β-mercaptoethanol (Bio-Rad, cat#1610710).
Proteins were separated by SDS–polyacrylamide gel electrophoresis (Any kD Mini-Protean
TGX gel, Bio-Rad, cat#4568125) and transferred to a nitrocellulose membrane (Trans-Blot
Turbo Mini Nitrocellulose Transfer Packs, Bio-Rad, cat#1704158). Membranes were blocked
for 1 hour at RT in 5% skimmed milk. Antibodies are listed in Supplemental table S2. The
ECL Select kit (GE Healthcare, cat#RPN2235) was used to detect protein bands with a
ChemiDoc XRS+ system (Bio-Rad).

For RIP validation western blot of Pro-B cells cultured ex vivo from bone marrow, 10% of the
total protein extract was collected to validate the immunoprecipitation efficiency. Proteins were
separated by SDS–polyacrylamide gel electrophoresis (12% Mini-Protean TGX gel, Bio-Rad,
cat#4561043) and transferred to a nitrocellulose membrane (Trans-Blot Turbo Mini
Nitrocellulose Transfer Packs, Bio-Rad, cat#1704158). Membranes were blocked for 1 hour at
RT in 5% skimmed milk. Antibodies are listed in Supplemental table S2.

qRT-PCR

212
50.000 sorted cells were lysed in RNAzol (Sigma-Aldrich, cat#R4533) and total RNA was
extracted using Direct-zol RNA MicroPrep kit (Zymo Research, cat#R2060) according to the
manufacturer's instructions. Total RNA (8 μl) was used in random primed cDNA synthesis
with TaqMan Reverse Transcription Reagents (Applied Biosystem, cat#N8080234). Diluted
cDNA 1:2 (4 μl) was used in every qPCR reaction using KAPA probe Fast qPCR kit (Kapa
Biosystem, cat#KK4706) with primers targeting 45S (Thermofisher Scientific, cat#

Mm03985792_s1) and 18S (Thermofisher Scientific, cat# 4333760T) rRNA transcripts. The

relative expression levels of rRNA were evaluated using log2 (2ΔCt) calculation.

Statistical analysis

All statistical analysis was carried out in R (version 4.1.2) and RStudio (version 1.4.1717),

using functions from the following packages: tidyverse (version 1.3.1), DESeq2 (version

1.32.0), rstatix (version 0.7.0), and ggpubr (version 0.4.0).

Antibodies

Antibodies are listed in Supplemental table 1 (Flow cytometry) and 2 (Western blot and

Immunoprecipitation).

Results

The LIN28B mRNA interactome in primary B cell precursors is enriched for ribosomal

protein transcripts

To identify direct mRNA targets of LIN28B that could be important during normal B cell

development, we took advantage of tet-LIN28B mice which express ectopic FLAG tagged

human LIN28B upon doxycycline (Dox) treatment (33). To obtain sufficient cell numbers, B

cell precursors from tet-LIN28B and control mice were expanded for 5 days *ex vivo* from whole

bone marrow cultures in the presence of IL-7 and Dox (Supplemental figure 1A). Cultured

cells were subsequently subjected to RNA co-immunoprecipitation (IP) coupled with poly-A

selected RNA-sequencing (RIP-seq) (Figure 1A). Immunoprecipitation efficiency of LIN28B

was validated by western blot analysis (Supplemental figure 1B). Total RNA sequencing of

corresponding input samples was used for calculating RIP-seq enrichment. RIP/input ratio of

the technical replicates showed a high degree of correlation (R=0.93, p < 2.2e-16) (Figure 1B)

and were averaged for downstream analyses. 238 transcripts displayed a higher than two-fold

RIP/input enrichment in tet-LIN28B samples but showed no trend in the control (Supplemental

Table 3 and Figure 1C). Gene ontology (GO) analysis revealed a prominent enrichment for

ribosomal protein coding transcripts (Figure 1D), suggesting that ribosomal function might be

a core target pathway of LIN28B during B cell development. Interestingly, similar observations

were previously made in other cell types, including 293T cells (22) and BE(2)-C neuroblastoma

cells (31). Indeed, cross comparison of our identified targets with these studies yielded

cytoplasmatic translation as the predominant overlapping GO term (Figure 1E-F). Our analysis

also identified other GO terms involved in RNA splicing and oxidative phosphorylation,

consistent with previous LIN28A/B interactome studies in various cell types (23, 37). We

conclude that the direct LIN28B mRNA regulon in the context of primary bone marrow B cell

progenitor cells is dominated by transcripts implicated in universal biosynthetic pathways,

which are shared across a wide range of cell types. In particular, our findings implicate the

ribosome as a potential target for LIN28B regulation in B cell progenitors.

LIN28B enhances protein synthesis in the later stages of B cell development

LIN28B bound mRNA targets have been linked to both translational increase and decrease in

a context dependent manner (22-31). In the case of ribosomal protein transcripts, enhanced

protein translation was shown by stable isotope labeling using amino acid in culture (SILAC)

(22) and polysome profiling (31). To assess ribosomal protein levels during B cell

development, we performed western blot analysis on 50,000 sorted B cell precursors from tet-

LIN28B and control mice following 10 days of Dox induction *in vivo* (Supplemental figure

2A). As expected, based on the dynamic changes in bioenergetic states during B cell

development, the expression of ribosomal proteins rise and fall in the large and small Pre-B

cell stages, respectively (Figure 2A). Consistent with enhanced translation, we found increased

levels of ribosomal proteins upon LIN28B induction throughout B cell development.

Given the increase in ribosomal proteins, we expected a corresponding increase in ribosomal

RNA (rRNA) content. To this end, we measured the 18S rRNAs by qRT-PCR of FACS sorted

B cell precursors. To our surprise, we detected no LIN28B dependent differences in Pro-B or

large Pre-B cells. Instead, rRNA content first began to increase in a LIN28B dependent manner

in the small Pre-B cell stage reaching statistical significance in Imm-B cells (Figure 2B).

Additional analysis of the 45S precursor rRNA followed the same trend (Figure 2C). Thus,

increased ribosomal protein levels precede increased ribosome biogenesis during LIN28B

induced B cell development *in vivo*. To evaluate whether this observation correlates with

changes in global protein synthesis rates we measured newly synthesized proteins in freshly

isolated B cell precursors using metabolic labelling with the methionine analog L-

azidohomoalanine (AHA) (38). As expected, FACS readout of AHA uptake from Dox induced

control and tet-LIN28B mice showed highly dynamic protein synthesis levels during B cell

development (Figure 2D). Pro-B cells exhibited a bimodal profile with respect to protein

synthesis that correlated with cell size (AHA^{high} or AHA^{low}, Supplemental figure 2B). Although

there was a slight tendency towards more AHA^{high} Pro-B cells in tet-LIN28B induced mice, no

statistical significance was detected in either the median fluorescence intensity or frequency of

the two subsets (Figure 2E, Supplemental figure 2C). Instead, LIN28B induced a progressive

increase in global protein synthesis after the Pro-B cell stage, peaking in Imm-B cells. Protein

synthesis in mature B-1 and B-2 cells were not affected by LIN28B expression (Figure 2F).

Taken together, we conclude that LIN28B enhances ribosome content and global protein

synthesis levels during the later stages of bone marrow B cell development.

IL-7 responsiveness dictates the stage specific effects of LIN28B during B cell

development

Elevated ribosomal biogenesis is a hallmark of c-MYC activity (39). We have previously

shown that LIN28B upregulates c-MYC protein expression in Imm-B cells (19). However, the

status of LIN28B induced c-MYC in earlier B cell precursor stages is not known. Given our

insight on ribosomal content and protein synthesis, we analyzed c-MYC protein levels

throughout B cell development using intracellular FACS. Consistent with its known IL-7

driven expression in Pro-B cells (2, 40), we found that c-MYC peaked in a subset of Pro-B

cells regardless of genotype. The impact of LIN28B on c-MYC expression began after the Pro-

B cell stage and culminated in Imm-B cells (Figure 3A-B), mirroring the trend observed for

protein synthesis. These findings are also observed at the transcriptome level, where gene set

enrichment analysis (GSEA) identified a LIN28B induced c-MYC target gene signature

succeeding the IL-7 dependent Pro-B and large Pre-B stages (Figure 3C). The same trend was

observed for LIN28B RIP-seq targets, consistent with c-MYC driving ribosome biogenesis. In

contrast, LIN28B induced de-repression of Let-7 targets was most prominent during the IL-7

dependent stages of B cell development and waned upon entry into the small Pre-B cell stage.

These findings demonstrate that Let-7 dependent and independent targets of the LIN28B

molecular program alternates along the B cell differentiation axis. We propose that the LIN28B

induced increase in ribosomal protein levels may cooperate with its effects on c-MYC

expression to elevate global protein synthesis during the later stages of bone marrow B cell

development.

IL-7 is a major trophic factor driving c-MYC expression prior to the Pre-BCR checkpoint (2,

40). We therefore hypothesized that its potent effects on Pro-B cells might mask any LIN28B

mediated c-MYC up-regulation. To test this, we expanded B cell precursors from control and

tet-LIN28B bone marrow under saturating IL-7 conditions for five days, before replating the

cells under limiting doses of IL-7 (Figure 3D) (Mandal 2019). As expected, IL-7 withdrawal

led to a drastic decrease in c-MYC and a collapse in protein synthesis of Pro-B cells (Figure

3E-F). This decrease in c-MYC protein was substantially cushioned by tet-LIN28B (Figure 3E

left). The same trend was also observed for protein synthesis (Figure 3E right), although it was

not statistically significant (Figure 3F right). Consistent with the lack of IL-7Ra expression in

Imm-B cells, no effect of IL-7 titration was observed (Figure 3G). Taken together, we

demonstrate that LIN28B exerts an effect on the MYC/ribosome axis and that this effect is

masked by IL-7 signaling during the Pro-B and large Pre-B cell stages. These findings provide

an explanation to the stage specific effects of LIN28B on the c-MYC/ribosome axis during B

cell development (Figure 3H).

Neonatal B cell development relies on a heightened level of protein synthesis

Endogenous LIN28B expression is developmentally limited to fetal and neonatal life. FACS analysis of LIN28B-eGFP reporter mice (34) captured its window of expression in B cell precursors as it gradually diminished to background levels after day 10 (Figure 4A). Following the same kinetics, neonatal B cell development exhibits a transient elevation of protein synthesis rates as measured by AHA uptake (Figure 4B). This effect was LIN28B dependent during the small Pre-B and Imm-B cell stages and fully abolished in 3-day-old neonates that lacked endogenous *Lin28b* (Figure 4C). In contrast, *Lin28b* deficiency does not ablate increased protein synthesis in neonatal Pro-B cells. These findings establish increased protein synthesis as a hallmark of early life B cell development and confirm our observations from tet-LIN28B mice in the context of wildtype neonates.

To investigate the importance of protein synthesis for neonatal B cell development, we took advantage of the ribosomal protein hypomorphic $Rpl24^{WT/Bst}$ (Bst) mouse model. This model carries a mutation that disrupts $Rpl24$ mRNA splicing and thereby ribosome biogenesis, exhibiting a 30% reduction in overall protein synthesis (41). This reduction in protein synthesis does not affect normal adult B lymphopoiesis (38), but is sufficient to impair Eμ-MYC driven oncogenesis (42). Given our findings, we hypothesized that early-life B lymphopoiesis might be more sensitive to minor constraints in protein synthesis. Indeed, our analysis show stunted B cell development in Bst neonates (Figure 4D) caused by a collapse in cell numbers at the large Pre-B cell stage and beyond. This phenotype was most severe in three-day-old neonates but became fully resolved by postnatal day 19. Taken together, these findings demonstrate that neonatal B lymphopoiesis is more sensitive to constraints in protein synthesis compared to the adult and therefore exhibits signs of ribosome addiction.

Elevated protein synthesis is required for the output of CD5+ B-1 cells early in life

The early-life expression of *Lin28b* allows for the developmental progression of self-reactive B cell precursors into the mature B cell pool, including the output of CD5+ B-1 cells (19, 20). This relaxation in the central tolerance threshold is evident in recently selected Imm-B cells where LIN28B expression increases surface levels of CD5, a marker for self-reactivity (19). Interestingly, further analysis of Imm-B cells revealed that CD5 expression induced by LIN28B positively correlated with protein synthesis levels (Figure 5A-B). Thus, enhanced protein synthesis may be important for the developmental progression of self-reactive B cells destined for the CD5+ B-1 fate. To test this, we treated LIN28B induced mice with a single dose of intraperitoneally administered Rapamycin to acutely dampen protein synthesis (Figure 5C) (43). As expected, the treatment yielded a moderate reduction in protein synthesis levels in both control and tet-LIN28B Imm-B cells (Figure 5D). This treatment selectively diminished the CD5 levels in tet-LIN28B induced Imm-B cells (Figure 5E-F) demonstrating that LIN28B dependent output of self-reactive B cells depends on increased protein synthesis.

Finally, to investigate the effect of constrained protein synthesis on early life B cell output, we measured the abundance of CD5+ B-1 cells in 19-day-old Bst mice. This timepoint was selected to evaluate B-1 cell output with minimal influence by peripheral homeostatic expansion. FACS analysis identified a significant decrease in the absolute number of peritoneal cavity (PerC) CD5+ B-1 cells of Bst mice as compared to littermate controls (Figure 5G). Further analysis of their reactivity against the predominant self-membrane component phosphatidyl choline (PtC) demonstrated a three-fold decrease of PtC reactive CD5+ B-1 B cells in Bst mice (Figure 5H-I). We therefore conclude that elevated protein synthesis is a central distinguishing parameter in neonatal B lymphopoiesis that potentiates the output of CD5+ B-1 cells with self-reactive BCR specificities.

Discussion

Our findings identify elevated protein synthesis as a hallmark of early life B cell development that is required for the time restricted output of self-reactive B-1 cells during ontogeny. This is mediated in part by the endogenous expression of the heterochronic RNA-binding protein LIN28B. Its expression naturally wanes during the first weeks of life and ectopic LIN28B expression is sufficient to recapitulate elevated protein synthesis in adult mice. Consistent with previous reports, we show that LIN28B directly binds to ribosomal protein transcripts and correlates with enhanced ribosomal protein expression in primary B cell precursors. Interestingly, this phenotype is shared across all stages of B cell development and precedes the increase in ribosomal biogenesis and protein synthesis first observed in small Pre-B cells. The delayed effect on ribosomal function coincides with the appearance of LIN28B induced c-MYC expression. Thus, increased ribosomal protein components was not sufficient to promote ribosome biogenesis in the absence of elevated c-MYC expression. We propose that LIN28B mediated post-transcriptional regulation of ribosomal proteins cooperates with the transcriptional program of c-MYC to elevate protein synthesis during B cell development. The precise extent and mechanism by which LIN28B superimposes post-transcriptional regulation on the backdrop of a c-MYC induced transcriptome have not been disentangled in this study. Nevertheless, our findings address a gap in the knowledge of neonatal B cell development with important implications in neonatal adaptive immune responses.

Protein synthesis is the most energy consuming biosynthetic process in the cell and subject to the constraints imposed by hundreds of components required for ribosome assembly (44). Indeed, protein synthesis rates give an accurate account of the overall cellular metabolism (45). And yet, its ability to control the development and function of immune cells that require rapid expansion and contraction of cellular growth has been an under-explored subject of

investigation. Our results using AHA metabolic labelling to measure protein synthesis

confirmed dynamic changes in cellular metabolism that underly proliferative bursts following

successful BCR checkpoints (1). Importantly, we demonstrate the ability of LIN28B to cushion

the biggest drop in protein synthesis in the small Pre-B cell stage. Interestingly, the restricted

metabolism of small Pre-B cells has been implicated as a mechanism to safeguard the central

tolerance checkpoint through enforcing nutrient stress and negative selection upon strong BCR

engagement (5). This notion is highly compatible with our observation that LIN28B promotes

protein synthesis to augment positive selection and the output of self-reactive CD5+ B-1 cells

early in life. Thus, we propose that the relatively mild increase in protein synthesis by LIN28B

provides sufficient relaxation of the metabolic constraints during B cell maturation, to

qualitatively shift the neonatal BCR repertoire.

In Pro-B cells, IL-7 signaling and downstream mTORC1 activation is the principal driver of c-

MYC expression and acts to promote biomass accumulation and proliferation (Zeng et al

2018). The attenuation of IL-7 signaling during B cell development marks a dramatic decrease

in overall bioenergetic state whereby preBCR/BCR mediated signals become dominant in cell

fate instruction. Interestingly, this metabolic gear shift correlates with distinct outcomes of

LIN28B mediated effects. While de-repression of Let-7 targets is most prominent during the

IL-7 dependent stages, the opposite is true of its effect on protein synthesis which appear later.

This could be due to changes in the abundance of LIN28B/Let-7 RNA substrates in

proliferating versus quiescent cells respectively. Therefore, our findings implicate metabolic

status as a powerful rheostat of LIN28B action.

In neonatal B cell development, protein synthesis is higher during all stages compared to the

adult. However, LIN28B is neither necessary nor sufficient to account for the observed effect

in Pro-B cells under IL-7 sufficient conditions, suggesting that other developmentally restricted

neonatal factors are at play. To this end, LIN28B has previously been shown to act in an

autoregulatory loop with Igf2bp3, another RNA-binding protein which binds a substantially

overlapping transcript pool to promote a fetal-like lymphopoiesis (26). It is therefore possible

that partial redundancy with Igf2bp3 could account for certain LIN28B independent effects of

early life B cell development. In the context of IL-7 deprivation however, Pro-B cells become

rapidly sensitized to LIN28B mediated effects. This finding is consistent with the previously

reported ability of LIN28B to circumvent the strict requirement for IL-7Ra during adult B

lymphopoiesis (16). Thus, overlapping effects by both LIN28B and IL-7 on the very same c-

MYC/ribosome axis may offer an explanation to the decreased reliance on IL-7 for early life

B lymphopoiesis (6).

In this study, we have not resolved the fates of LIN28B bound transcripts in the rapidly

alternating stages of B cell development. One obstacle preventing the examination of the

context dependent effects of LIN28B is the limited cellular material from quiescent cell types

in vivo. Methods such as polysome sequencing to profile changes in protein translation are

optimized for proliferating cells. However, our findings make clear that *in vitro* culturing or

oncogene mediated transformation will not recapitulate LIN28B mediated effects in small Pre-

B cells. Thus, the prohibitive nature of quiescent cells poses technical challenges that must be

overcome to address the context dependent actions of LIN28B *in vivo*. Nevertheless, our results

demonstrate that protein synthesis is an important differentially regulated feature of neonatal

B lymphopoiesis and open new avenues for future investigations. In the context of B cell

neoplasms, protein synthesis is a key mediator of c-MYC driven oncogenesis (42). Considering

the strong infant and childhood association of B cell precursor acute lymphoblastic leukemia

incidence, it will be important to investigate a potential role for increased ribosome addiction

in the pathogenesis of the disease. In the context of the neonatal B cell responses, we and others

have demonstrated that neonatal immunization produces unique B cell memory clones not

generated by adult mice immunized with the same antigen (46, 47). Our current findings that

enhanced protein synthesis critically shapes the output of naïve B cell specificities implicates

ribosomal control in the formation of a layered immune system. Going forward, the extent to

which this may instruct early life B cell memory formation remains an outstanding question of

great interest.

Acknowledgements

We thank Jonas Ungerbäck and Johanna Tingvall-Gustafsson for help with bioinformatics. We

thank Jenny Hansson, Leal Oburoglu, and Svetlana Soboleva for their input. Finally, we also

thank the FACS and Bioinformatics core facilities at the Lund Stem Cell Center for their

support.

Footnotes

Funding

J.Y. was supported by the Swedish Research Council, the Swedish Cancer Society, the

European Research Council (715313), Knut and Alice Wallenberg Foundation and the

Wenner-Gren Foundations.

Figure Legends

Figure 1. LIN28B binds to ribosomal protein transcripts in primary B cell precursors

(A) Schematic of experimental setup for the identification of LIN28B-mRNA targets. B cell

progenitors were expanded from whole, adult bone marrow in the presence of IL-7 (20 ng/mL)

and Dox (0.1 µg/mL) for five days prior to RNA-IP. Co-immunoprecipitated mRNA from two

tet-LIN28B and one control mouse were individually subjected to RNA-seq. **(B)** Scatter plot

showing enrichment correlation between the two tet-LIN28B samples. LIN28B IP/input

enrichment ratio of least two-fold (n=238) were used for subsequent analyses. **(C)** Average

enrichment ranking of the tet-LIN28B samples compared to the control sample. **(D)** Gene

ontology analysis of enriched transcripts. **(E)** Analysis of overlapping genes between our RIP-

targets and previously published LIN28B-mRNA interactome data (22, 31). **(F)** Plot showing

how gene ontology terms identified in D compares and overlaps with LIN28B interacting

transcripts identified in the indicated studies.

Figure 2. LIN28B enhances ribosome biogenesis and protein synthesis in small Pre-B and

Imm-B cells

(A) Representative western blot analysis for indicated ribosomal proteins on 50,000 FACS

sorted B cell progenitors from tet-LIN28B or littermate control mice following 10 days of Dox

induction. Representative of at least two biological replicates per cell type and genotype. Blots

were cropped to improve clarity. **(B)** qRT-PCR of 18S rRNA and **C)** 45S pre-rRNA from

50,000 FACS sorted B cell progenitors, expression values were normalized relative to the

littermate control Imm-B cells. Data from 4-5 biological replicates and 2 experimental

replicates. **(D)** Representative FACS histogram overlays of L-Azidohomoalanine (AHA)

uptake, corresponding to their protein synthesis rates, in the indicated progenitor subsets. **(E)**

Average of the median fluorescence intensity (MFI) for AHA, as shown in D. Data from 7

biological replicates and 3 experimental replicates. **(F)** Average AHA MFI from FACS analysis

on peritoneal cavity (PerC) B cell subsets. Data from 3-4 biological replicates and 2

experimental replicates. Wilcox test was used to calculate p-values. *p<0.05, **p<0.01,

***p<0.001. Bars show the mean and error bars show standard deviation.

Figure 3. IL-7 dependent stimulation of the c-MYC/ribosome axis masks the anabolic

effects of LIN28B in Pro-B cells

(A) Representative histograms showing intracellular FACS analysis of c-MYC protein

expression. **(B) c-**MYC median fluorescence intensity (MFI) levels in B cell precursors, from

the indicated genotypes following 10 days of doxycycline induction. Data is represented as

average values normalized per cell type to the littermate control. Data from 7-12 biological

replicates and 6 experimental replicates. **(C)** Gene set enrichment analysis (GSEA) of RNA-

seq data from FACS sorted B cell precursors for the indicated signatures (right Y axis) and cell

types. LIN28B enrichment rank was Wald's stat from DESeq2, comparing each cell type for

differentially expressed genes in tet-LIN28B to control mice. The Pro-B cells were sorted on

high FSC-A, in addition to the immunophenotype as stated under Materials and Methods. Data

from 3-6 biological replicates. **(D)** Experimental setup for measuring the impact of IL-7

withdrawal on LIN28B mediated effects on c-MYC protein expression and L-

azidohomoalanine (AHA) uptake in cKit+ Pro-B cells. **(E)** Representative FACS histograms

for the indicated IL-7 conditions, showing the measured c-MYC protein expression (left) and

AHA uptake (right). **(F)** Average MFI of c-MYC expression and AHA uptake FACS in Pro-B

cells kept under indicated IL-7 conditions. Values were normalized to littermate controls at

max IL-7 concentrations 20 ng/mL. Data from three separate experiments, with 3-4 littermate

and Lin28b mice depending on cell type. **(G)** Same as in (F), but for IgM+ Imm-B cells. **(H)**

Schematic illustration of the proposed LIN28B mechanism in B cell precursors. Wilcox test

was used to calculate p-values in B. T-test was used to calculate p-values in F and G. *p<0.05,

**p<0.01, ***p<0.001. Bars show the mean and error bars show standard deviation.

Figure 4. Neonatal B cell development is dependent on enhanced protein synthesis

(A) Average frequency of *Lin28b*-eGFP positivity of the indicated B cell precursor subsets,

from *Lin28b*-eGFP mice at the indicated age. 4-10 biological replicates, over the different days

of analysis. **(B)** Average AHA MFI normalized to adult samples. Three biological replicates

each, two separate experiments. **(C)** Average AHA MFI from 3-day old *Lin28b* WT (n=5) or

KO (n=5) neonates normalized to WT adult samples. Data from two separate experiments. **(D)**

Average frequency of B cell precursors from neonatal or adult Rpl24-Bst heterozygous mutants

or littermate control mice, at the indicated ages. Data from at least two separate experiments,

number of mice is indicated in figure legends. T-test was used to calculate p-values in B, C,

and D. *p<0.05, **p<0.01, ***p<0.001. Bars show the mean and error bars show standard

deviation.

Figure 5. Elevated protein synthesis is required for the output of CD5+ B-1 cells early in

life

(A) (Left) Representative FACS plots for CD5 expression in Imm-B cells from the indicated

genotypes treated with doxycycline for 10 days. (Right) Histogram showing the L-

azidohomoalanine (AHA) uptake of the individual CD5 slices. **(B)** Average AHA median

fluorescence intensity (MFI) within individual CD5 slices shown as relative values to control

Imm-B cells (n=4-5). Data from four separate experiments. **(C)** Schematic for Rapamycin

(Rapa, 2 mg/kg) treatment of tet-LIN28B and littermate control mice with Rapamycin, or

vehicle control treatment (EtOH). Mice were treated by a single, intra-peritoneal injection and

analyzed 24hrs later. **(D)** AHA MFI from Imm-B cells of the indicated genotype and treatment,

shown relative to EtOH treated littermate controls (n=5-6). Data from four separate

experiments. **(E)** Representative FACS histograms showing CD5 levels. **(F)** Average CD5 MFI

from Imm-B cells of the indicated genotypes and treatment, shown relative to EtOH treated

littermate controls (n=4-7). **(G)** Absolute number of peritoneal cavity (PerC) CD5+ B-1 cells

in day 19 neonates of the indicated genotypes (n=4). **(H)** Representative FACS plots showing

the frequency of CD5+ PtC reactive B cells from 19-day old neonates of the indicated

genotypes. **(I)** Average frequency of CD5+PtC+ B cells in day 19 neonates of the indicated

genotypes (n=4). Wilcox test was used to calculate p-values $*p<0.05$, $**p<0.01$. Bars show the

mean and error bars show standard deviation.

References

1. Clark, M. R., M. Mandal, K. Ochiai, and H. Singh. 2014. Orchestrating B cell lymphopoiesis through interplay of IL-7 receptor and pre-B cell receptor signalling. *Nat Rev Immunol* 14: 69-80.

2. Zeng, H., M. Yu, H. Tan, Y. Li, W. Su, H. Shi, Y. Dhungana, C. Guy, G. Neale, C. Cloer, J. Peng, D. Wang, and H. Chi. 2018. Discrete roles and bifurcation of PTEN signaling and mTORC1-mediated anabolic metabolism underlie IL-7-driven B lymphopoiesis. *Sci Adv* 4: eaar5701.

3. Mandal, M., M. K. Okoreeh, D. E. Kennedy, M. Maienschein-Cline, J. Ai, K. C. McLean, N. Kaverina, M. Veselits, I. Aifantis, F. Gounari, and M. R. Clark. 2019. CXCR4 signaling directs Igk recombination and the molecular mechanisms of late B lymphopoiesis. *Nat Immunol* 20: 1393-1403.

4. Nemazee, D. 2017. Mechanisms of central tolerance for B cells. *Nat Rev Immunol* 17: 281-294.

5. Muschen, M. 2019. Metabolic gatekeepers to safeguard against autoimmunity and oncogenic B cell transformation. *Nat Rev Immunol* 19: 337-348.

6. Kikuchi, K., and M. Kondo. 2006. Developmental switch of mouse hematopoietic stem cells from fetal to adult type occurs in bone marrow after birth. *Proc Natl Acad Sci U S A* 103: 17852-17857.

7. Hayakawa, K., M. Asano, S. A. Shinton, M. Gui, D. Allman, C. L. Stewart, J. Silver, and R. R. Hardy. 1999. Positive selection of natural autoreactive B cells. *Science* 285: 113-116.

8. Chen, J. W., T. A. Rice, J. M. Bannock, A. A. Bielecka, J. D. Strauss, J. R. Catanzaro, H. Wang, L. C. Menard, J. H. Anolik, N. W. Palm, and E. Meffre. 2020. Autoreactivity in naive human fetal B cells is associated with commensal bacteria recognition. *Science* 369: 320-325.

9. Wong, J. B., S. L. Hewitt, L. M. Heltemes-Harris, M. Mandal, K. Johnson, K. Rajewsky, S. B. Koralov, M. R. Clark, M. A. Farrar, and J. A. Skok. 2019. B-1a cells acquire their unique characteristics by bypassing the pre-BCR selection stage. *Nat Commun* 10: 4768.

10. Baumgarth, N. 2011. The double life of a B-1 cell: self-reactivity selects for protective effector functions. *Nat Rev Immunol* 11: 34-46.

11. Ambros, V., and H. R. Horvitz. 1984. Heterochronic mutants of the nematode Caenorhabditis elegans. *Science* 226: 409-416.

12. Copley, M. R., and C. J. Eaves. 2013. Developmental changes in hematopoietic stem cell properties. *Exp Mol Med* 45: e55.

13. Zhou, Y., Y. S. Li, S. R. Bandi, L. Tang, S. A. Shinton, K. Hayakawa, and R. R. Hardy. 2015. Lin28b promotes fetal B lymphopoiesis through the transcription factor Arid3a. *J Exp Med* 212: 569-580.

14. Copley, M. R., S. Babovic, C. Benz, D. J. Knapp, P. A. Beer, D. G. Kent, S. Wohrer, D. Q. Treloar, C. Day, K. Rowe, H. Mader, F. Kuchenbauer, R. K. Humphries, and C. J. Eaves. 2013. The Lin28b-let-7-Hmga2 axis determines the higher self-renewal potential of fetal haematopoietic stem cells. *Nat Cell Biol* 15: 916-925.

15. Sampson, V. B., N. H. Rong, J. Han, Q. Yang, V. Aris, P. Soteropoulos, N. J. Petrelli, S. P. Dunn, and L. J. Krueger. 2007. MicroRNA let-7a down-regulates MYC and reverts MYC-induced growth in Burkitt lymphoma cells. *Cancer Res* 67: 9762-9770.

16. Yuan, J., C. K. Nguyen, X. Liu, C. Kanellopoulou, and S. A. Muljo. 2012. Lin28b reprograms adult bone marrow hematopoietic progenitors to mediate fetal-like lymphopoiesis. *Science* 335: 1195-1200.

17. Kristiansen, T. A., E. Jaensson Gyllenback, A. Zriwil, T. Bjorklund, J. A. Daniel, E. Sitnicka, S. Soneji, D. Bryder, and J. Yuan. 2016. Cellular Barcoding Links B-1a B Cell Potential to a Fetal Hematopoietic Stem Cell State at the Single-Cell Level. *Immunity* 45: 346-357.

18. Wang, J., E. M. Wissink, N. B. Watson, N. L. Smith, A. Grimson, and B. D. Rudd. 2016. Fetal and adult progenitors give rise to unique populations of CD8+ T cells. *Blood* 128: 3073-3082.

19. Vanhee, S., H. Akerstrand, T. A. Kristiansen, S. Datta, G. Montano, S. Vergani, S. Lang, J.
Ungerback, A. Doyle, K. Olsson, G. Beneventi, C. T. Jensen, C. Bellodi, S. Soneji, M.
Sigvardsson, E. J. Gyllenback, and J. Yuan. 2019. Lin28b controls a neonatal to adult switch in
B cell positive selection. *Sci Immunol* 4.
20. Xu, X., M. Deobagkar-Lele, K. R. Bull, T. L. Crockford, A. J. Mead, A. P. Cribbs, D. Sims, C.
Anzilotti, and R. J. Cornall. 2020. An ontogenetic switch drives the positive and negative
selection of B cells. *Proc Natl Acad Sci U S A* 117: 3718-3727.
21. Benhamou, D., V. Labi, R. Novak, I. Dai, S. Shafir-Alon, A. Weiss, R. Gaujoux, R. Arnold, S. S.
Shen-Orr, K. Rajewsky, and D. Melamed. 2016. A c-Myc/miR17-92/Pten Axis Controls PI3K-
Mediated Positive and Negative Selection in B Cell Development and Reconstitutes CD19
Deficiency. *Cell Rep* 16: 419-431.
22. Graf, R., M. Munschauer, G. Mastrobuoni, F. Mayr, U. Heinemann, S. Kempa, N. Rajewsky,
and M. Landthaler. 2013. Identification of LIN28B-bound mRNAs reveals features of target
recognition and regulation. *RNA Biol* 10: 1146-1159.
23. Wilbert, M. L., S. C. Huelga, K. Kapeli, T. J. Stark, T. Y. Liang, S. X. Chen, B. Y. Yan, J. L.
Nathanson, K. R. Hutt, M. T. Lovci, H. Kazan, A. Q. Vu, K. B. Massirer, Q. Morris, S. Hoon, and
G. W. Yeo. 2012. LIN28 binds messenger RNAs at GGAGA motifs and regulates splicing factor
abundance. *Mol Cell* 48: 195-206.
24. Cho, J., H. Chang, S. C. Kwon, B. Kim, Y. Kim, J. Choe, M. Ha, Y. K. Kim, and V. N. Kim. 2012.
LIN28A is a suppressor of ER-associated translation in embryonic stem cells. *Cell* 151: 765-
777.
25. Hafner, M., K. E. Max, P. Bandaru, P. Morozov, S. Gerstberger, M. Brown, H. Molina, and T.
Tuschl. 2013. Identification of mRNAs bound and regulated by human LIN28 proteins and
molecular requirements for RNA recognition. *RNA* 19: 613-626.
26. Wang, S., B. Chim, Y. Su, P. Khil, M. Wong, X. Wang, A. Foroushani, P. T. Smith, X. Liu, R. Li, S.
Ganesan, C. Kanellopoulou, M. Hafner, and S. A. Muljo. 2019. Enhancement of LIN28B-
induced hematopoietic reprogramming by IGF2BP3. *Genes Dev* 33: 1048-1068.
27. Basak, A., M. Munschauer, C. A. Lareau, K. E. Montbleau, J. C. Ulirsch, C. R. Hartigan, M.
Schenone, J. Lian, Y. Wang, Y. Huang, X. Wu, L. Gehrke, C. M. Rice, X. An, H. A. Christou, N.
Mohandas, S. A. Carr, J. J. Chen, S. H. Orkin, E. S. Lander, and V. G. Sankaran. 2020. Control
of human hemoglobin switching by LIN28B-mediated regulation of BCL11A translation. *Nat*
Genet 52: 138-145.
28. Osborne, J. K., M. A. Kinney, A. Han, K. E. Akinnola, A. V. Yermalovich, L. T. Vo, D. S. Pearson,
P. M. Sousa, S. Ratanasirintrawoot, K. M. Tsanov, J. Barragan, T. E. North, R. J. Metzger, and
G. Q. Daley. 2021. Lin28 paralogs regulate lung branching morphogenesis. *Cell Rep* 36:
109408.
29. Madison, B. B., Q. Liu, X. Zhong, C. M. Hahn, N. Lin, M. J. Emmett, B. Z. Stanger, J. S. Lee, and
A. K. Rustgi. 2013. LIN28B promotes growth and tumorigenesis of the intestinal epithelium
via Let-7. *Genes Dev* 27: 2233-2245.
30. Tan, F. E., S. Sathe, E. C. Wheeler, and G. W. Yeo. 2021. Non-microRNA binding competitively
inhibits LIN28 regulation. *Cell Rep* 36: 109517.
31. Missios, P., E. L. da Rocha, D. S. Pearson, J. Philipp, M. M. Aleman, M. Pirouz, D. Farache, J.
W. Franses, C. Kubaczka, K. M. Tsanov, D. K. Jha, B. Pepe-Mooney, J. T. Powers, R. I. Gregory,
A. S. Lee, D. Dominguez, D. T. Ting, and G. Q. Daley. 2021. LIN28B alters ribosomal dynamics
to promote metastasis in MYCN-driven malignancy. *J Clin Invest* 131.
32. Peng, S., L. L. Chen, X. X. Lei, L. Yang, H. Lin, G. G. Carmichael, and Y. Huang. 2011. Genome-
wide studies reveal that Lin28 enhances the translation of genes important for growth and
survival of human embryonic stem cells. *Stem Cells* 29: 496-504.
33. Zhu, H., N. Shyh-Chang, A. V. Segre, G. Shinoda, S. P. Shah, W. S. Einhorn, A. Takeuchi, J. M.
Engreitz, J. P. Hagan, M. G. Kharas, A. Urbach, J. E. Thornton, R. Triboulet, R. I. Gregory, D.

Consortium, M. Investigators, D. Altshuler, and G. Q. Daley. 2011. The Lin28/let-7 axis
regulates glucose metabolism. *Cell* 147: 81-94.
34. Kristiansen, T. A., Q. Zhang, S. Vergani, E. Boldrin, N. Krausse, O. Andre, P. Nordenfelt, M.
Sigvardsson, D. Bryder, J. Ungerback, and J. Yuan. 2022. Developmental cues license
megakaryocyte priming in murine hematopoietic stem cells. *Blood Adv*.
35. Holl, T. M., B. F. Haynes, and G. Kelsoe. 2010. Stromal cell independent B cell development
in vitro: generation and recovery of autoreactive clones. *J Immunol Methods* 354: 53-67.
36. Ersching, J., A. Efeyan, L. Mesin, J. T. Jacobsen, G. Pasqual, B. C. Grabiner, D. Dominguez-
Sola, D. M. Sabatini, and G. D. Victora. 2017. Germinal Center Selection and Affinity
Maturation Require Dynamic Regulation of mTORC1 Kinase. *Immunity* 46: 1045-1058 e1046.
37. Shyh-Chang, N., H. Zhu, T. Yvanka de Soysa, G. Shinoda, M. T. Seligson, K. M. Tsanov, L.
Nguyen, J. M. Asara, L. C. Cantley, and G. Q. Daley. 2013. Lin28 enhances tissue repair by
reprogramming cellular metabolism. *Cell* 155: 778-792.
38. Signer, R. A., J. A. Magee, A. Salic, and S. J. Morrison. 2014. Haematopoietic stem cells
require a highly regulated protein synthesis rate. *Nature* 509: 49-54.
39. Ruggero, D. 2009. The role of Myc-induced protein synthesis in cancer. *Cancer Res* 69: 8839-
8843.
40. Morrow, M. A., G. Lee, S. Gillis, G. D. Yancopoulos, and F. W. Alt. 1992. Interleukin-7 induces
N-myc and c-myc expression in normal precursor B lymphocytes. *Genes Dev* 6: 61-70.
41. Oliver, E. R., T. L. Saunders, S. A. Tarle, and T. Glaser. 2004. Ribosomal protein L24 defect in
belly spot and tail (Bst), a mouse Minute. *Development* 131: 3907-3920.
42. Barna, M., A. Pusic, O. Zollo, M. Costa, N. Kondrashov, E. Rego, P. H. Rao, and D. Ruggero.
2008. Suppression of Myc oncogenic activity by ribosomal protein haploinsufficiency. *Nature*
456: 971-975.
43. Beretta, L., A. C. Gingras, Y. V. Svitkin, M. N. Hall, and N. Sonenberg. 1996. Rapamycin blocks
the phosphorylation of 4E-BP1 and inhibits cap-dependent initiation of translation. *EMBO J*
15: 658-664.
44. Buttgereit, F., and M. D. Brand. 1995. A hierarchy of ATP-consuming processes in
mammalian cells. *Biochem J* 312 (Pt 1): 163-167.
45. Arguello, R. J., A. J. Combes, R. Char, J. P. Gigan, A. I. Baaziz, E. Bousiquot, V. Camosseto, B.
Samad, J. Tsui, P. Yan, S. Boissonneau, D. Figarella-Branger, E. Gatti, E. Tabouret, M. F.
Krummel, and P. Pierre. 2020. SCENITH: A Flow Cytometry-Based Method to Functionally
Profile Energy Metabolism with Single-Cell Resolution. *Cell Metab* 32: 1063-1075 e1067.
46. Vergani, S., K. G. Muleta, C. Da Silva, A. Doyle, T. A. Kristiansen, S. Sodini, N. Krausse, G.
Montano, K. Kotarsky, J. Nakawesi, H. Akerstrand, S. Vanhee, S. L. Gupta, D. Bryder, W. W.
Agace, K. Lahl, and J. Yuan. 2022. A self-sustaining layer of early-life-origin B cells drives
steady-state IgA responses in the adult gut. *Immunity*.
47. New, J. S., B. L. P. Dizon, C. F. Fucile, A. F. Rosenberg, J. F. Kearney, and R. G. King. 2020.
Neonatal Exposure to Commensal-Bacteria-Derived Antigens Directs Polysaccharide-Specific
B-1 B Cell Repertoire Development. *Immunity* 53: 172-186 e176.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5