FAP106 is an interaction hub required for assembly of conserved and lineage-specific microtubule inner proteins at the cilium inner junction

[a]Michelle M. Shimogawa, [a]Angeline S. Wijono, [a,b,c]Hui Wang, [a,c,e]Jiayan Zhang, [d]Jihui Sha, [a]Natasha Szombathy, [a]Sabeeca Vadakkan, [a]Paula Pelayo, [a]Keya Jonnalagadda, [d]James Wohlschlegel, [a,c,e]Z. Hong Zhou*, and [a,c,e]Kent L. Hill*

[a]Department of Microbiology, Immunology and Molecular Genetics, University of California Los Angeles, Los Angeles, CA, 90095

[b]Department of Bioengineering, University of California Los Angeles, Los Angeles, CA, 90095

[c]California NanoSystems Institute, University of California Los Angeles, Los Angeles, CA, 90095

[d]Department of Biological Chemistry, University of California Los Angeles, Los Angeles, CA, 90095

[e]Molecular Biology Institute, University of California Los Angeles, Los Angeles, CA, 90095

Running title: *T. brucei* MIPs

*Corresponding authors:

Kent L. Hill

609 Charles E. Young Drive E, MSB 1602-G, Los Angeles, CA 90095

kenthill@microbio.ucla.edu

Z. Hong Zhou

CNSI 5511C, 570 Westwood Plaza, Los Angeles, CA 90095

Hong.Zhou@UCLA.edu

Author contributions

Designed research: MS, AW, HW, JZ, HZ, KH

Performed research: MS, AW, HW, JZ, JS, NS, SV, PP, KJ

Contributed new reagents or analytic tools:

Analyzed data: MS, AW, HW, JZ, JS, NS, SV, PP, KJ, JW, HZ, KH

Wrote the paper: MS, AW, HW, HZ, KH

ABSTRACT

Motility of pathogenic protozoa depends on flagella (synonymous with cilia) with axonemes containing nine doublet microtubules (DMTs) and two singlet microtubules. Microtubule inner proteins (MIPs) within DMTs influence axoneme stability and motility and provide lineage-specific adaptations, but individual MIP functions and assembly mechanisms are mostly unknown. Here, we show in the sleeping sickness parasite *Trypanosoma brucei*, that FAP106, a conserved MIP at the DMT inner junction, is required for trypanosome motility and functions as a critical interaction hub, directing assembly of several conserved and lineage-specific MIPs. We further use comparative cryogenic electron tomography (cryoET) and quantitative proteomics to identify novel MIP candidates, and RNAi knockdown plus fitting of AlphaFold models into cryoET maps to demonstrate one of these, MC8, is a trypanosome-specific MIP required for parasite motility. Our work advances understanding of MIP assembly mechanisms and identifies lineage-specific motility proteins that are attractive targets to consider for therapeutic intervention.

INTRODUCTION

Cilia (synonymous with eukaryotic flagella) are conserved, microtubule-based structures that provide motility and signaling functions at the surface of eukaryotic cells. Cilium motility is required for normal human reproduction, development, physiology, and to combat infection (1). Motile cilia also drive cell propulsion in eukaryotic pathogens responsible for tremendous human suffering worldwide (2, 3). Given this broad distribution and central importance to eukaryote cell biology and human health, understanding structural foundations of cilium assembly and motility is of broad interest.

Most motile cilia are built around a "9 + 2" cylindrical axoneme consisting of nine doublet microtubules (DMTs) surrounding a central pair of singlet microtubules. Each DMT is comprised

of a complete, 13-protofilament A-microtubule (A-tubule) and a partial, 10-protofilament B-

microtubule (B-tubule), with non-tubulin proteins forming an "inner junction" (IJ) filament that

connects the A- and B-tubules (4, 5). Recent studies have revealed a previously unappreciated,

yet universal feature of motile axonemes, showing that DMTs are not hollow but, instead, are

lined with a highly organized network of interconnected "microtubule inner proteins" (MIPs) (6-

9). Structures inside DMTs had been reported several decades ago in flagellated protozoa (10,

11), but the diversity and ubiquitous distribution of such structures was not recognized until

cryogenic electron tomography (cryoET) enabled high-resolution, 3D structural analysis (8, 12).

Studies in organisms from diverse eukaryotic lineages (13-17) indicate MIPs are present in both

the A- and B-tubules and occur in a regularly repeating pattern along the length of the DMT in

all motile axonemes. Their broad taxonomic distribution indicates MIPs are fundamentally

important for axoneme function and likely were present prior to eukaryote diversification.

Meanwhile, absence of regularly spaced MIP structures in non-motile cilia (18, 19) suggests

they are required specifically for motile axonemes. Recent work has shown MIPs can influence

axoneme stability, structure and beating (13, 14, 16, 20-23). Nonetheless, many aspects of MIP

functions and assembly mechanisms remain poorly understood.

In recent, groundbreaking work, cryogenic electron microscopy (cryoEM) was used to

reconstruct near-atomic resolution structures of the axoneme, thereby allowing *in situ*

identification and atomic modeling of MIPs in *Chlamydomonas*, *Tetrahymena,* and mammals

(13, 16, 17, 20, 24). These studies have expanded opportunities for studying MIP function and

revealed that while some MIPs are conserved across diverse taxa, others represent lineage-

specific adaptations (13, 16, 17, 20, 24). Axonemes of diverse organisms share general

architecture and beating mechanism, and lineage-specific MIPs therefore offer a potential

mechanism for meeting organism-specific motility characteristics (2, 15, 25), such as those

presented by movement of parasitic protozoa through tissues of their human and animal hosts

(26, 27). Identification and analysis of lineage-specific MIPs are therefore of great interest.

African trypanosomes, e.g. *Trypanosoma brucei* and related kinetoplastids, are

flagellated, protozoan pathogens that cause extensive human suffering and limit economic

development in some of the world's poorest regions (28, 29). Trypanosomes also represent

diverse eukaryotic lineages that have proven useful for uncovering novel features of eukaryote

biology (30, 31). *T. brucei* motility exhibits several distinctive features that support movement

through host tissues (15, 32) and is required for pathogenesis (33) and transmission (34). The

T. brucei flagellum has a large and diverse repertoire of MIP structures, including several that

are lineage-specific, particularly in the B-tubule (15). Except for a few conserved examples,

however, proteins comprising most *T. brucei* MIP structures are unknown, representing a critical

knowledge gap in *T. brucei* flagellum assembly and motility.

FAP106/ENKUR is a conserved B-tubule MIP that is required for motility of mammalian

sperm (35) and occupies a key position in the B-tubule, spanning the IJ filament and

interconnecting several other MIPs (13, 16, 20, 36). However, the structural foundation of the

FAP106/ENKUR motility defect is unknown, and the role of FAP106 in assembly of the IJ and

other MIPs has not been examined. To address these questions, we used RNAi knockdown to

investigate FAP106 function in *T. brucei* and show that FAP106 is required for parasite motility

and normal flagellum length. Combining CryoET and tandem mass tag (TMT) quantitative

proteomics, we further show that FAP106 knockdown results in loss of conserved IJ MIP

structures and their corresponding proteins, as well as several MIP structures and proteins that

are lineage-specific. One of these, MC8, is demonstrated by cryoET to be a novel lineage-

specific *T. brucei* MIP and is required for parasite motility. Together, our results demonstrate

that FAP106 is a critical interaction hub required for assembly of conserved and lineage-specific

MIPs at the inner junction. This work advances understanding of MIP assembly mechanisms

and provides insight into parasite-specific MIPs that control *T. brucei* motility. Because they are

not present in the mammalian host, such MIPs are attractive for consideration as targets for

therapeutic intervention in trypanosome infections.

RESULTS

FAP106 is required for parasite motility.

FAP106 forms a 'tether loop' structure that spans the inner junction and is interconnected with

several other MIPs (13, 16, 20, 36). We used tetracycline-inducible RNAi (37) to deplete

FAP106 in procyclic form cells, hereafter referred to as "FAP106 KD". RNAi knockdown reduced

FAP106 mRNA expression by more than 95% (Fig. 1A) without affecting parasite growth rate

(Fig. 1B). FAP106 KD parasites did not exhibit any gross morphological defects, although their

flagella were shorter than normal (Fig. 1C-E). Notably, despite having a beating flagellum,

FAP106 KD parasites rarely exhibited directional motility, even at high cell densities, which

typically result in faster directional motility (Fig. 1F-G; Supp. Fig. S1). Thus, FAP106 is critical

for normal parasite motility.

FAP106 is required for assembly of conserved and lineage-specific MIP structures.

To determine whether FAP106 is required for assembly of other MIP structures, we performed

cryoET on purified, demembranated flagella from control and FAP106 KD parasites. Most

known MIPs assemble with a periodicity of 16, or 48 nm (9, 13-16, 20, 24, 36, 38) and we

therefore performed subtomogram averaging to determine the 48-nm repeating unit of the DMT

(Fig. 2; Supp. Fig. S2; Movie 1) for comparison to the published structures of the 48-nm repeat

from other organisms (13, 16, 20). MIP densities in the A-tubule appeared largely unaffected in

the FAP106 KD and, given the position of FAP106 in the B-tubule, we focused our analysis on

the B-tubule. Cross-sectional views of the cryoET tomogram show several major B-tubule MIP

densities are missing in the FAP106 KD. Although we do not have sufficient resolution to

identify the corresponding proteins directly, referencing the position of MIPs conserved in other

organisms provides insight. Longitudinal and cross-sectional views (Fig. 2C-D; Movie 1) show

loss of densities corresponding to the 'tether loop' structure of FAP106 in algae and mammals

(13, 16, 20) (Fig. 2B-C, red; Movie 1), consistent with the idea that FAP106 structure is

conserved across species (13). A density corresponding to the expected position of FAP52 (16,

20, 21) appears largely unaffected in the FAP106 KD (Fig. 2B-D, tan; Movie 1). We did not

resolve filamentous densities corresponding to FAP45 on protofilaments B6-B9 in other

organisms (13, 16), but these protofilaments were heavily decorated in *T. brucei* control cells

and relatively devoid of MIP densities in FAP106 KD (Fig. 2D; Movie 1). Additionally, densities

corresponding to the expected position of the FAP45 N-terminus are reduced in the FAP106 KD

(Fig. 2D, orange). Thus, FAP106 appears to be required for proper assembly of FAP45 but not

FAP52. Interestingly, the FAP106 KD also shows loss of FAP106-associated densities that

resemble FAP210 in other organisms (Fig. 2B, 2D, green; Movie 1; (13, 16, 20, 36)), although *T.*

brucei lacks a clear FAP210 homolog (Table 1). Overall, our results indicate FAP106 is critical

for assembly of a subset of B-tubule MIP structures in the vicinity of the IJ.

Beyond known MIP structures defined in other organisms, additional densities reduced

in the FAP106 KD correspond to MIP structures that are absent in one or more other organisms.

These lineage-specific MIP structures include a protruding density on protofilament B8, "MIP

B8", densities attached to protofilament A12, "MIP A12", and a protruding density on

protofilament B5, "MIP B5" (Fig. 2B). Thus, FAP106 is required for assembly of universally-

conserved, as well as lineage-specific MIPs.

FAP106 is required for IJ filament assembly.

The IJ filament is a conserved non-tubulin structure that connects protofilaments A1 and B10 in

organisms with cilia (8, 19). The IJ filament is comprised of PACRG and FAP20 subunits that

alternate each 4nm along the axoneme (4, 5). Interestingly, most organisms have a single hole

in the IJ filament every 96nm, corresponding to one missing PACRG subunit per 96-nm

repeating unit (8, 12, 38). *T. brucei* is unusual in having two holes per 96-nm repeating unit (15)

(Fig. 2E). The function of the IJ hole is not known, nor is the structural basis for the presence of

two holes per 96nm in *T. brucei*. Strikingly, our cryoET data show that loss of FAP106 results in

two additional holes per 48-nm repeat, giving six holes per 96-nm repeat (Fig. 2E; Movie 1).

These holes correspond to loss of every fourth subunit of the IJ filament (light blue in Fig. 2E),

which, based on analogy with *Chlamydomonas* (16), likely corresponds to PACRG.

TMT proteomics identifies novel, FAP106-dependent, lineage-specific MIP proteins.

To identify proteins corresponding to FAP106-dependent MIP densities, we used tandem mass

tag (TMT) quantitative proteomics to compare the protein composition of purified,

demembranated flagella from control and FAP106 KD parasites. We quantified more than 2500

proteins and only eight of these were significantly reduced in flagella from FAP106 KD parasites

($\geq$2-fold reduced, adjusted p-value $\leq$0.06) (Fig. 3A; Table 2; Supp. Dataset S1). Three proteins

were observed to be increased in the knockdown (Fig. 3A; Supp. Dataset S1) and were not

studied further.

As expected, the protein most reduced in FAP106 KD parasites was FAP106 itself (Fig.

3A; Table 2), confirming efficient knockdown reported by qRT-PCR (Fig. 1A) and loss of

FAP106 protein from the axoneme (Fig. 2). Two other characterized proteins, FAP45 and

PACRG-B, were also reduced in the FAP106 KD (Fig. 3A-B, Table 2). FAP45 was almost 4-fold

reduced in FAP106 KD flagella (Fig. 3A; Table 2), supporting results from cryoET indicating

FAP106 is required for assembly of FAP45 into the axoneme (Fig. 2). FAP52 and FAP20 were

unaffected (Supp. Dataset S1), further supporting conclusions from cryoET that these proteins

assemble independently of FAP106 (Fig. 2). Notably, *T. brucei* has two PACRG homologs,

PACRG-A and PACRG-B (Table 1), which have been suggested to be functionally redundant

(39, 40). Despite a high degree of similarity, however, PACRG-A and PACRG-B are readily

distinguishable by mass spectrometry and our TMT proteomics analysis shows that only

PACRG-B is lost in the FAP106 KD (Fig. 3A; Table 2; Supp. Dataset S1). Loss of PACRG-B

and not PACRG-A indicates that the extra holes in the FAP106 KD (Fig. 2E) correspond to

missing PACRG-B subunits and demonstrates that PACRG-A and PACRG-B are not redundant.

Based on spacing of IJ subunits, the two IJ holes in control parasites most likely also

correspond to missing PACRG-B.

In addition to the known MIPs described above, five previously uncharacterized proteins

were reduced in flagella from FAP106 KD parasites (Fig. 3A; Table 2). One of these proteins

(RPC19, Tb927.11.8890) localizes to the nucleus (41), suggesting it is unlikely to be a MIP. The

remaining four proteins (Tb927.10.7120, Tb927.11.4920, Tb927.11.2770, and Tb927.3.3200),

are kinetoplastid-specific and localize to the flagellum (Supp. Fig. S3A; (41)), making them

strong candidates for lineage-specific, FAP106-dependent MIPs. Importantly, these four

proteins were separately identified as "MIP candidates", MC3, MC5, MC8, and MC15, in a

completely independent APEX2-based proximity proteomics analysis as being nearby FAP45

and FAP52 in the B-tubule (Supp. Fig. S4; Supp. Dataset S2). Identification of these four

proteins by independent approaches greatly supports their potential as novel, kinetoplastid-

specific MIPs.

To test the requirement of MC3, MC5, MC8 and MC15 for flagellum structure and

function, we used tetracycline-inducible RNAi to individually knockdown each of these proteins.

Knockdown resulted in loss of the respective proteins from the flagellum (Supp. Fig. S3A)

without substantial effects on parasite growth. Loss of MC5 did not affect parasite motility, while

MC3, MC8, and MC15 are each required for normal motility (Fig. 3C; Supp. Fig. S3B). MC8 KD

and MC15 KD parasites also showed small reductions in flagellum length (Fig. 3D; Supp. Fig.

S3C). To ask whether additional proteins depend on these MCs for assembly into the axoneme,

we performed TMT proteomics on flagella isolated from parasites in which the individual MCs

were knocked down (Supp. Fig. S3D-G; Supp. Dataset S1). These proteomic analyses confirm

significant reduction of each MC in the respective knockdown, but no additional proteins were

lost ($\geq$2-fold reduced, adjusted p-value $\leq$0.06) in any of the four knockdowns (Supp. Fig. S3D-G,

Supp. Dataset S1). Therefore, motility defects in MC3, MC8, and MC15 knockdowns reflect a

requirement for these individual, kinetoplastid-specific proteins in trypanosome motility.

MC8 is a novel, lineage-specific *T. brucei* MIP that comprises the MIP B8 density.

To identify MIPs corresponding to FAP106-dependent MIP densities, we fit the AlphaFold (42,

43) models of the above MCs into our cryoET structures. AlphaFold prediction of individual

proteins returned low-confidence predictions for MC3, MC5, and MC15, but predicted that MC8

folds into a structure with a high-confidence pyramid-like domain formed by α helices (Fig. 4A),

which fits well into the MIP B8 density (Fig. 4B). To determine whether MC8 in fact corresponds

to the MIP B8 density, we performed cryoET on purified, demembranated flagella from control

and MC8 KD parasites (Fig. 4C-E). Because TMT analysis showed that MC8 is the only protein

lost in the MC8 KD (Supp. Fig. S3F; Supp. Dataset S1), any density altered in the knockdown

must correspond to MC8 itself. The subtomographic average of the 48-nm repeating unit from

MC8 KD axonemes was nearly identical to that of controls except for the absence of the MIP B8

density (Fig. 4C-E). Thus, MC8 is a bona fide, *T. brucei*-specific B-tubule MIP that is required for

normal parasite motility (Fig. 3C; Supp. Fig. S3B). The motility defect of MC8 KD parasites, in

which the only missing protein is MC8, demonstrates importance of lineage-specific MIPs to

flagellar motility.

DISCUSSION

FAP106 knockdown provides insights into MIP assembly mechanisms.

Functional analysis lags behind identification and structural assignment of MIP proteins. Here

we combine functional, structural, and proteomic analysis to demonstrate that the conserved

MIP FAP106 is required for motility and is a key interaction hub for IJ filament and B-tubule MIP

structures. The role of FAP106 as a key interaction hub is emphasized by our finding that, while

FAP106 is required for assembly of FAP45 and lineage-specific MIP proteins MC3, 5, 8, and 15

(Fig. 2; Fig. 3, Fig. 5), loss of these individual proteins does not affect other MIP proteins or

structures (Supp. Fig. 3D-G; Fig. 4; Supp. Dataset S1; (21)). Thus, rather than interdependency,

our results suggest a hierarchical mechanism for B-tubule MIP assembly. FAP106-dependency

of FAP45 and a FAP210 structural homolog suggests that extensive interactions of FAP45 and

FAP210 with the tubulin lattice (13, 16) are relatively weak, while interactions with FAP106 (13)

are critical for stable binding and establishing their periodicity within the 48-nm repeat. This

supports the idea that an important role of MIPs is to enable complex assembly of proteins

along otherwise uniform protofilament polymers of tubulin (13, 14, 16) and suggests that loss of

FAP45 and FAP210 in RIB72 mutants (44) may reflect dependency on FAP106 rather than a

requirement for RIB72 directly. Our studies also provide a structural explanation for motility

defects of FAP106 mutants in trypanosomes (Fig. 2), and in sperm from FAP106/ENKUR

mutants in mice (35).

The IJ filament subunit PACRG exhibits functional heterogeneity in trypanosomes.

PACRG alternates with FAP20 to form the highly conserved IJ filament (4, 5). One of the more

mysterious features of DMTs is a hole in the IJ filament that occurs once per 96-nm repeat in

most organisms and corresponds to a missing PACRG (8, 12, 38). *T. brucei* is unusual in

having two IJ holes per 96-nm repeat, one at the site of nexin-dynein regulatory complex

(NDRC) attachment as observed in other organisms, and one proximal to the NDRC (15). The

proximal hole is devoid of external structures on the DMT (15), suggesting it is dictated by MIP

interactions rather than NDRC attachment, which is proposed for the NDRC hole (45). Our

findings support this idea, as loss of FAP106 results in extra holes in the IJ filament. Some

organisms have multiple PACRG homologs (24, 39, 46), but the contribution of individual

homologs is unknown. It is not known, for example, if they are distinctly incorporated within a

single axoneme or used for specialization of axoneme function in different cell types or

developmental stages. *T. brucei* has two PACRG homologs, PACRG-A and PACRG-B, which

have been suggested to be functionally redundant (39, 40). Our combined structural and

proteomics data instead support a model in which PACRG-A and PACRG-B both regularly

assemble within each IJ filament, alternating with FAP20 in turn, and with holes in the IJ

filament corresponding to missing PACRG-B, but not PACRG-A (Fig. 2E). Distinct locations and

assembly requirements for PACRG-A and PACRG-B indicate they play distinct roles in

assembly of the IJ. Although PACRG is conserved among ciliated organisms, the family of

PACRG proteins shows substantial heterogeneity in the N-terminal region that has been shown

to interact closely with the DMT lattice in *Chlamydomonas* (20, 46). Divergent N-termini of

PACRG-A versus PACRG-B may contribute to the differential dependence on FAP106 and their

distinct roles in assembly of the IJ.

MC8 is a novel, lineage-specific T. brucei MIP that is required for parasite motility.

The growing body of evidence suggests IJ subunits (PACRG and FAP20) and a core set of

conserved MIPs are found in all motile cilia, while other MIPs vary between organisms (13, 15-

17, 20, 24, 36, 38). FAP45, FAP52, and FAP106 represent core B-tubule MIPs while FAP126,

FAP276, and FAP210 are only found in some organisms (Table 1). MIP A12 densities (Fig. 2B-

C) are not present in *Chlamydomonas* or mammals (13, 16, 20) and appear more prominent in

T. brucei than MIP5a and MIP5b located at a similar position in *Tetrahymena* (9, 24, 36, 38), but

the proteins that comprise these densities remain unknown. We demonstrate that MC8 is a

bona fide lineage-specific MIP corresponding to the MIP B8 structure (Fig. 4). A density with

location and periodicity similar to MIP B8 is present in *Tetrahymena* (9, 24, 36, 38), but we were

unable to identify a *Tetrahymena* protein with sequence similarity to the MIP B8 protein, MC8.

Loss of MC8 in the FAP106 KD (Fig. 2B-2D) is likely due to loss of FAP45, which runs along

protofilaments B7-B8 (13, 16) and contacts FAP106 directly (13). Loss of MC8 compromises

motility (Fig. 3C; Supp. Fig. 3B) without affecting other MIPs (Fig. 4, Supp. Fig. 3F),

demonstrating that lineage-specific MIPs directly impact trypanosome motility. Validation of

MC8 as a bona fide *T. brucei* MIP in the current study, through a combination of mutant

analysis, structural, functional, and proximity proteomics approaches, demonstrates the power

of this combined approach for MIP identification and analysis. Our studies also strongly suggest

that MC3, 5, and 15 likewise correspond to novel, lineage-specific *T. brucei* MIPs. Trypanosome

motility is necessary for virulence in the mammalian host (33) and transmission through the

tsetse fly vector (34). Therefore, identification of novel, trypanosome-specific MIPs provides

attractive targets to consider for treatment of devastating diseases caused by these pathogens.

Model for assembly of the IJ

Our work provides an important advance toward elucidating the order of MIP assembly and

supports the following model for IJ assembly (Fig. 5). FAP20 binds every 8 nm, together with at

least a subset of PACRG, to form the IJ filament and this occurs independently from assembly

of FAP52 (20) and FAP106 (this work). FAP52 binds every 16 nm, independently of the IJ

filament (4) and FAP106 (this work), and may connect the IJ filament to the B-tubule via

FAP276 or other analogous structure (20, 36). FAP106 also binds every 16 nm, independently

from FAP52 (20), bridging the A- and B-tubules directly and serving as a critical guidepost for

assembling additional MIPs such as FAP45 and FAP210 (this work). Although FAP52 and

FAP106 bind regularly within each 48-nm repeat, their connections to other MIPs differ every 16

nm. Despite apparent contacts between FAP52 and other IJ MIPs (13, 16, 20, 36), assembly of

IJ MIPs in *Chlamydomonas* is largely unaffected by loss of FAP52 (20). Thus, while FAP52 and

FAP106 form contacts with each other and other IJ MIPs, they assemble independently from

one another, and FAP106 appears to be more critical for assembly of additional B-tubule MIPs.

Going forward, as additional MIPs are identified, particularly those that are lineage-specific,

application of combined proteomics, cryoEM, and mutant analysis, as done here, will be

necessary to completely define MIP assembly mechanisms and to yield a full understanding of

their contribution to flagellum function.

MATERIALS AND METHODS

Biological materials are available from the authors upon request.

***Trypanosoma brucei* culture**

Procyclic *T. brucei brucei* (strain 29-13) (47) were cultivated in SM medium (48) supplemented

with 10% heat-inactivated fetal bovine serum (FBS) at 28°C with 5% CO_2.

In situ tagging

mNeonGreen and APEX2-tagged cell lines were generated in the 29-13 background by C-

terminal tagging (41) with pPOTv6-puro-puro-mNG or pPOTv7-blast-blast-APEX2-3xHA,

respectively. pPOTv7-blast-blast-APEX2-3xHA contains the same APEX2-3xHA tag as in (49).

For MC3-NG, a transient CRISPR-Cas9 system (50) was used to tag both alleles. See Supp.

Table S2 for list of primers.

Tetracycline-inducible knockdown

Constructs for tetracycline-inducible knockdown were designed using RNAit (51) and cloned

into p2T7-177 (37). See Supp. Table S2 for list of primers. NotI-linearized plasmids were

transfected into 29-13 (FAP106 KD) or the corresponding MC-NG tagged cell lines (MC KDs)

using established methods (48). Clonal lines were generated by limiting dilution. Knockdown

was induced by growing cells in the presence of 1 µg/mL tetracycline (Tet) for at least 3d.

Control parasites were grown in the presence of tetracycline for comparison to the respective

KDs. For growth curves, cells were counted with a Beckman Coulter Z1 particle counter and

diluted daily to a concentration of 1e6 cells/mL, in triplicate.

qRT-PCR

Quantitative real-time PCR (qRT-PCR) was performed as previously described (52) with the

following modifications. FAP106-specific primers were designed using RNAit (51). Analyses

were performed in duplicate on three independent RNA preparations and values were

normalized to RPS23 (53). See Supp. Table S2 for list of primers.

Motility analyses

Cells were grown to ~1e6 cells/mL and loaded into polyglutamate-coated motility chambers (33)

sealed with Vaseline. 30s videos were acquired under dark-field illumination at ~40 frames per

second (fps) using a Hamamatsu ORCA-Flash 4.0 camera on a Zeiss Axio Imager A2 inverted

microscope with an EC Plan-Neofluar 10x/0.3 objective lens and Zen 2.6 Pro software. At least

five videos were collected for each independent biological replicate. Single cell tracking and

mean squared displacement were performed in MATLAB as previously described for fluorescent

parasites (33). For 'high density' motility analyses, cells were grown to ~1e7 cell/mL, then

diluted in conditioned medium to ~1e6 cells/mL and analyzed as above.

Purification of demembranated flagella (detergent-extracted flagellum skeletons)

Detergent extraction was done to demembranate flagellum skeletons, essentially as described

(15). Briefly, cells were washed in Dulbecco's PBS (DPBS), resuspended in Extraction buffer

(20 mM HEPES, pH 7.4, 1 mM $MgCl_2$, 150 mM NaCl, 0.5% NP40, 2x Sigmafast EDTA-free

protease inhibitor cocktail) for 15 min at room temperature, followed by depolymerization of

subpellicular microtubules with 1 mM $CaCl_2$ on ice for 30 min. Detergent-extracted flagellum

skeletons were pelleted at 1500 rcf for 10 min at 4°C.

Flagellum length measurements

For measurements of purified flagella, detergent-extracted flagellum skeletons were prepared

as described above and phase contrast images were acquired on a Zeiss Axioskop II

microscope with a Plan-Apochromat 63x/1.4 objective lens using Axiovision 4.8 software. For

measurements of the paraflagellar rod (PFR) in detergent-extracted cytoskeletons, cells were

washed in Dulbecco's PBS (DPBS) and allowed to adhere to glass cover slips for 10 min. Cover

slips were rinsed to remove unattached cells, then extracted with PEME buffer (100 mM PIPES,

2 mM EGTA, 1 mM $MgSO_4$, 0.1 mM EDTA, pH 6.8) + 1% IGEPAL CA-630 (NP40) + 2x

Sigmafast EDTA-free protease inhibitor cocktail for 10 min. Cover slips were fixed in methanol

at -20°C for 10 min, allowed to dry, then rehydrated in DPBS for 15 min. Cover slips were

blocked in DPBS + 8% normal donkey serum + 2% bovine serum albumin then incubated in

rabbit anti-PFR2 (54) primary antibody diluted 1:1000 in blocking solution, followed by donkey

anti-rabbit Alexa 594 secondary antibody diluted 1:1500 in blocking solution. Cover slips were

mounted in Vectashield containing DAPI and images were acquired on a Zeiss Axioskop II

fluorescence microscope with a Plan-Apochromat 100x/1.4 objective lens and Axiovision 4.8 or

Zen 3.5 Pro software. Flagellum lengths were measured in Fiji using the freehand line tool and

analyzed in GraphPad Prism 7.

CryoET

CryoET grids and detergent-extracted flagellum skeletons were prepared essentially as

described (15). Briefly, 2e8 log phase cells were harvested and washed twice in DPBS.

Flagellum skeletons were prepared as described above, resuspended in 100 ul Extraction buffer

and spun over a 30% sucrose cushion in Extraction buffer without NP40 at 800 rcf for 5 min at

4°C. 100-150 µL of flagellum skeletons were collected from the upper fraction, washed once

with 700 ul Extraction buffer, and once with 1 mL Extraction buffer without NP40. Flagellum

skeletons were resuspended in Extraction buffer without NP40 and vitrified on quantifoil grids

with 5nm fiducial gold beads.

With *SerialEM* (55), tilt series were collected in a Titan Krios instrument equipped with a

Gatan imaging filter (GIF) and a post-GIF K3 direct electron detector in electron-counting mode

with parameters listed in Table S1. Frames in each movie of the raw tilt series were aligned,

drift-corrected, and averaged with *Motioncor2* (56). The micrographs in each tilt series were

aligned and reconstructed into 3D tomograms by either weighted back projections (WBP, for

sub-tomographic averaging) or simultaneous iterative reconstruction technique (SIRT, for

visualization and particle picking) using the *IMOD* software package (57). The contrast transfer

function (CTF) was determined by *ctffind4* (58) and corrected with the *ctfphaseflip* program (59)

of *IMOD*.

Sub-tomogram averaging, visualization and modeling

4x binned SIRT reconstructed tomograms were further processed by IsoNet (60) to improve the

contrast for particle picking. Because of their distinct features, radial spokes were picked as

particles to represent the centers of 96nm axonemal repeat units. These particles for 96 nm

repeat units were subject to three rounds of rough alignments with PEET (7, 61). The particle

coordinates for 48-nm repeat units were generated from the coordinates of the aligned particles

for 96-nm repeat units. While keeping all coordinates for 96-nm repeat units, extra coordinates

for 48-nm repeat units were added unidirectionally from the 96-nm-repeat-units coordinates in a

distance of 48nm along the axonemal axis. PEET alignments were performed on the particles of

48-nm repeat units to generate the final maps. In Summary, the sub-tomographic averages of

flagellar components from four datasets were generated from the following number of particles:

1014 particles from 20 tomograms of Control dataset 1, 728 particles from 18 tomograms of

FAP106 KD dataset, 1936 particles from 52 tomograms of MC8 KD dataset, and 538 particles

from nine tomograms of Control dataset 2. The cryoET sub-tomographic averaged maps have

been deposited in the Electron Microscopy Data Bank under the accession codes EMD-28802,

EMD-28803, EMD-28804, and EMD-28805, for FAP106 KD, Control 1, MC8 KD, and Control 2,

respectively. The resolution of each sub-tomographic average was calculated by

calcUnbiasedFSC in *PEET* based on the 0.143 FSC criterion (Supp. Fig. S2).

UCSF ChimeraX (62) was used to visualize reconstructed tomograms and sub-

tomographic averages. For surface rendering with UCSF ChimeraX, maps were first low pass

filtered to either 20Å or 30Å.

Model prediction of MC8 was performed by AlphaFold Colab (42, 43). AlphaFold

predicted model of MC8 was superimposed into our averaged density with the molmap and

fitmap functions in UCSF ChimeraX.

TMT labeling and quantitative proteomics

Detergent-extracted flagellum skeletons were prepared from 2e8-4e8 log phase cells as

described above. After detergent extraction and depolymerization of subpellicular microtubules,

the pellet of flagellum skeletons was washed in Extraction buffer without NP40 and solubilized in

100 µl 8M urea, 100 mM Tris, pH 8 for 15 min at room temperature. Samples were centrifuged

at full speed for 15 min to pellet any insoluble material. The supernatant was diluted with 100

mM Tris, pH 8 to a final concentration of 2M urea and precipitated with 20% ice-cold

trichloroacetic acid (TCA) on ice overnight. The sample was centrifuged at full speed for 30 min

at 4°C and washed twice with -20°C acetone. The TCA-precipitated pellets were dried and

stored at $\leq$-20°C.

Sample Digestion: Protein pellets were resuspended with digestion buffer (8M urea, 0.1M Tris-

HCl pH 8.5). Each sample was normalized by absorbance at 280 nm and 25µg was aliquoted

for the digestion. Samples were reduced and alkylated via sequential 20-minute incubations

with 5 mM TCEP and 10 mM iodoacetamide at room temperature in the dark. Carboxylate-

modified magnetic beads (CMMB) also known as SP3 (63) and ethanol (50% final

concentration) were added to each sample to induce protein binding to CMMB beads. CMMB

beads were washed three times with 80% ethanol and then resuspended with 18ul of 100 mM

TEAB. Protein samples were digested overnight with 1ul of 0.1µg/ul LysC (Promega) and 2ul of

0.4µg/ul trypsin (Pierce) at 37 °C.

TMT Labeling and CIF Fractionation: 9ul of 100% acetonitrile were added to each sample to a

final concentration of 30% (v/v). Then the TMT labels were resuspended with 100% acetonitrile.

25ug of each sample was labeled using TMT6plex (Thermo Fisher Scientific) and the four

labeled samples of each set (two independent biological replicates each of control and

knockdown) were pooled. The pooled samples were fractionated by CMMB-based Isopropanol

Gradient Peptide Fractionation (CIF) method (64) into six fractions before mass spectrometry

(MS) analysis.

LC-MS Acquisition and Analysis: Fractionated samples were separated on a 75uM ID x 25cm

C18 column packed with 1.9µm C18 particles (Dr. Maisch GmbH) using a 140-minute gradient

of increasing acetonitrile and eluted directly into a Thermo Orbitrap Fusion Lumos mass

spectrometer where MS spectra were acquired using SPS-MS3. Protein identification was

performed using MaxQuant (65) v 1.6.17.0. against a user assembled database consisting of all

protein entries from the TriTrypDB (66) for *T. brucei* strain 927 (version 7.0). Searches were

performed using a 20 ppm precursor ion tolerance. TMT6plex was set as a static modification

on lysine and peptide N terminal. Carbamidomethylation of cysteine was set as static

modification, while oxidation of methionine residues and N-terminal protein acetylation were set

as variable modifications. LysC and Trypsin were selected as enzyme specificity with maximum

of two missed cleavages allowed. 1% false discovery rate was used as a filter at both protein

and PSM levels. Statistical analysis was conducted with the MSstatsTMT Bioconductor package

(67). Raw proteomic data have been deposited to the Mass Spectrometry Interactive Virtual

Environment (MassIVE) under accession ID MSV000090661.

ACKNOWLEDGMENTS

We thank Daniel Velez-Ramirez for technical assistance with APEX2 proximity proteomics and

Simon Imhof for helpful discussions. We also thank Astra Bryant for assistance with high speed

microscopy. Funding was provided by NIH grants AI052348 (K.L.H.), GM071940 (Z.H.Z), and

GM089778 (J.A.W). KJ was supported by the Beckman Scholars Program (Beckman

Foundation). We acknowledge the use of resources in the Electron Imaging Center for

Nanomachines supported by UCLA and grants from NIH (S10RR23057 and S10OD018111)

and NSF (DBI-1338135 and DMR-1548924).

428   DATA AVAILABILITY STATEMENT

The structure data generated during the current study have been deposited in the Electron

Microscopy Data Bank, accession codes EMD-28802, EMD-28803, EMD-28804, and EMD-

28805. The raw proteomics data generated during the current study have been deposited to the

Mass Spectrometry Interactive Virtual Environment (MassIVE) under accession IDs

MSV000090660 and MSV000090661 and the analyzed datasets are included in this published

article and its supplementary information files.

REFERENCES

1. J. Wallmeier *et al.*, Motile ciliopathies. *Nat Rev Dis Primers* **6**, 77 (2020).

2. T. Kruger, M. Engstler, Flagellar motility in eukaryotic human parasites. *Semin Cell Dev Biol* **46**, 113-127 (2015).

3. H. S. Judelson, F. A. Blanco, The spores of Phytophthora: weapons of the plant destroyer. *Nat Rev Microbiol* **3**, 47-58 (2005).

4. E. E. Dymek *et al.*, PACRG and FAP20 form the inner junction of axonemal doublet microtubules and regulate ciliary motility. *Mol Biol Cell* **30**, 1805-1816 (2019).

5. H. A. Yanagisawa *et al.*, FAP20 is an inner junction protein of doublet microtubules essential for both the planar asymmetrical waveform and stability of flagella in Chlamydomonas. *Mol Biol Cell* **25**, 1472-1483 (2014).

6. H. Sui, K. H. Downing, Molecular architecture of axonemal microtubule doublets revealed by cryo-electron tomography. *Nature* **442**, 475-478 (2006).

7. D. Nicastro *et al.*, The molecular architecture of axonemes revealed by cryoelectron tomography. *Science* **313**, 944-948 (2006).

8. D. Nicastro *et al.*, Cryo-electron tomography reveals conserved features of doublet microtubules in flagella. *Proc Natl Acad Sci U S A* **108**, E845-853 (2011).

9. M. Ichikawa *et al.*, Subnanometre-resolution structure of the doublet microtubule reveals new classes of microtubule-associated proteins. *Nat Commun* **8**, 15035 (2017).

10. W. A. Anderson, R. A. Ellis, Ultrastructure of Trypanosoma Lewisi - Flagellum Microtublules and Kinetoplast. *J Protozool* **12**, 483-& (1965).

11. K. Vickerman, On the surface coat and flagellar adhesion in trypanosomes. *J Cell Sci* **5**, 163-193. (1969).

12. G. Pigino *et al.*, Comparative structural analysis of eukaryotic flagella and cilia from Chlamydomonas, Tetrahymena, and sea urchins. *J Struct Biol* **178**, 199-206 (2012).

13. M. Gui *et al.*, De novo identification of mammalian ciliary motility proteins using cryo-EM. *Cell* **184**, 5791-5806.e5719 (2021).

14. M. Ichikawa *et al.*, Tubulin lattice in cilia is in a stressed form regulated by microtubule inner proteins. *Proc Natl Acad Sci U S A* **116**, 19930-19938 (2019).

15. S. Imhof *et al.*, Cryo electron tomography with volta phase plate reveals novel structural foundations of the 96-nm axonemal repeat in the pathogen. *Elife* **8** (2019).

16. M. Ma *et al.*, Structure of the Decorated Ciliary Doublet Microtubule. *Cell* **179**, 909-922.e912 (2019).

17. M. Gui *et al.*, SPACA9 is a lumenal protein of human ciliary singlet and doublet microtubules. *Proc Natl Acad Sci U S A* **119**, e2207605119 (2022).

18. P. Kiesel *et al.*, The molecular structure of mammalian primary cilia revealed by cryo-electron tomography. *Nat Struct Mol Biol* **27**, 1115-1124 (2020).

19. S. Sun, R. L. Fisher, S. S. Bowser, B. T. Pentecost, H. Sui, Three-dimensional architecture of epithelial primary cilia. *Proc Natl Acad Sci U S A* **116**, 9370-9379 (2019).

20. A. A. Z. Khalifa *et al.*, The inner junction complex of the cilia is an interaction hub that involves tubulin post-translational modifications. *Elife* **9** (2020).

21. M. Owa *et al.*, Inner lumen proteins stabilize doublet microtubules in cilia and flagella. *Nat Commun* **10**, 1143 (2019).

22. D. Stoddard *et al.*, Tetrahymena RIB72A and RIB72B are microtubule inner proteins in the ciliary doublet microtubules. *Mol Biol Cell* **29**, 2566-2577 (2018).

23. T. Ide *et al.*, CFAP53 regulates mammalian cilia-type motility patterns through differential localization and recruitment of axonemal dynein components. *PLoS Genet* **16**, e1009232 (2020).

24. S. Kubo *et al.*, Native doublet microtubules from Tetrahymena thermophila reveal the
importance of outer junction proteins. *bioRxiv* 10.1101/2022.09.30.510376,
2022.2009.2030.510376 (2022).

25. M. Rossi, G. Cicconofri, A. Beran, G. Noselli, A. DeSimone, Kinematics of flagellar swimming in
Euglena gracilis: Helical trajectories and flagellar shapes. *Proc Natl Acad Sci U S A* **114**, 13085-
13090 (2017).

26. J. L. Bargul *et al.*, Species-Specific Adaptations of Trypanosome Morphology and Motility to the
Mammalian Host. *PLoS Pathog* **12**, e1005448 (2016).

27. L. G. Wilson, L. M. Carter, S. E. Reece, High-speed holographic microscopy of malaria parasites
reveals ambidextrous flagellar waveforms. *Proc. Natl. Acad. Sci. USA* **110**, 18769-18774 (2013).

28. P. Buscher, G. Cecchi, V. Jamonneau, G. Priotto, Human African trypanosomiasis. *Lancet* **390**,
2397-2409 (2017).

29. K. I. Kasozi *et al.*, Prevalence of hemoprotozoan parasites in small ruminants along a human-
livestock-wildlife interface in western Uganda. *Vet Parasitol Reg Stud Reports* **17**, 100309 (2019).

30. L. Vincensini, T. Blisnick, P. Bastin, 1001 model organisms to study cilia and flagella. *Biol Cell* **103**,
109-130 (2011).

31. J. E. Donelson, M. J. Gardner, N. M. El-Sayed, More surprises from Kinetoplastida. *Proc Natl Acad*
Sci U S A **96**, 2579-2581 (1999).

32. T. Kruger, S. Schuster, M. Engstler, Beyond Blood: African Trypanosomes on the Move. *Trends*
Parasitol **34**, 1056-1067 (2018).

33. M. M. Shimogawa *et al.*, Parasite motility is critical for virulence of African trypanosomes. *Sci*
Rep **8**, 9122 (2018).

34. B. Rotureau, C. P. Ooi, D. Huet, S. Perrot, P. Bastin, Forward motility is essential for trypanosome
infection in the tsetse fly. *Cell Microbiol* **16**, 425-433 (2014).

35. M. K. Jungnickel *et al.*, The flagellar protein Enkurin is required for mouse sperm motility and for
transport through the female reproductive tract. *Biol Reprod* **99**, 789-797 (2018).

36. S. Li, J. J. Fernandez, A. S. Fabritius, D. A. Agard, M. Winey, Electron cryo-tomography structure
of axonemal doublet microtubule from. *Life Sci Alliance* **5** (2022).

37. B. Wickstead, K. Ersfeld, K. Gull, Targeting of a tetracycline-inducible expression system to the
transcriptionally silent minichromosomes of Trypanosoma brucei. *Mol Biochem Parasitol* **125**,
211-216 (2002).

38. K. Song *et al.*, In situ structure determination at nanometer resolution using TYGRESS. *Nat*
Methods **17**, 201-208 (2020).

39. H. R. Dawe, H. Farr, N. Portman, M. K. Shaw, K. Gull, The Parkin co-regulated gene product,
PACRG, is an evolutionarily conserved axonemal protein that functions in outer-doublet
microtubule morphogenesis. *J Cell Sci* **118**, 5421-5430 (2005).

40. R. Broadhead *et al.*, Flagellar motility is required for the viability of the bloodstream
trypanosome. *Nature* **440**, 224-227 (2006).

41. S. Dean, J. D. Sunter, R. J. Wheeler, TrypTag.org: A Trypanosome Genome-wide Protein
Localisation Resource. *Trends Parasitol* **33**, 80-82 (2017).

42. J. Jumper *et al.*, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-
589 (2021).

43. M. Mirdita *et al.*, ColabFold - Making protein folding accessible to all. *bioRxiv*
10.1101/2021.08.15.456425, 2021.2008.2015.456425 (2022).

44. A. S. Fabritius *et al.*, Proteomic analysis of microtubule inner proteins (MIPs) in Rib72 null. *Mol*
Biol Cell **32**, br8 (2021).

45. T. Heuser, M. Raytchev, J. Krell, M. E. Porter, D. Nicastro, The dynein regulatory complex is the
nexin link and a major regulatory node in cilia and flagella. *J. Cell Biol.* **187**, 921-933 (2009).

46.   N. Khan *et al.*, Crystal structure of human PACRG in complex with MEIG1 reveals roles in
axoneme formation and tubulin binding. *Structure* **29**, 572-586 e576 (2021).

47.   E. Wirtz, S. Leal, C. Ochatt, G. A. Cross, A tightly regulated inducible expression system for
conditional gene knock-outs and dominant-negative genetics in *Trypanosoma brucei*. *Mol.*
Biochem. Parasitol. **99**, 89-101 (1999).

48.   M. Oberholzer, M. A. Lopez, K. S. Ralston, K. L. Hill, Approaches for functional analysis of flagellar
proteins in African trypanosomes. *Methods in cell biology* **93**, 21-57 (2009).

49.   D. E. Velez-Ramirez *et al.*, APEX2 Proximity Proteomics Resolves Flagellum Subdomains and
Identifies Flagellum Tip-Specific Proteins in Trypanosoma brucei. *mSphere* **6** (2021).

50.   S. Shaw, S. Knusel, S. Hoenner, I. Roditi, A transient CRISPR/Cas9 expression system for genome
editing in Trypanosoma brucei. *BMC Res Notes* **13**, 268 (2020).

51.   S. Redmond, J. Vadivelu, M. C. Field, RNAit: an automated web-based tool for the selection of
RNAi targets in Trypanosoma brucei. *Mol Biochem Parasitol* **128**, 115-118 (2003).

52.   M. M. Shimogawa *et al.*, Cell Surface Proteomics Provides Insight into Stage-Specific Remodeling
of the Host-Parasite Interface in Trypanosoma brucei. *Molecular & cellular proteomics : MCP* **14**,
1977-1988 (2015).

53.   M. A. Lopez, E. A. Saada, K. L. Hill, Insect stage-specific adenylate cyclases regulate social motility
in african trypanosomes. *Eukaryotic cell* **14**, 104-112 (2015).

54.   E. A. Saada *et al.*, Insect stage-specific receptor adenylate cyclases are localized to distinct
subdomains of the Trypanosoma brucei Flagellar membrane. *Eukaryotic cell* **13**, 1064-1076
(2014).

55.   D. N. Mastronarde, Automated electron microscope tomography using robust prediction of
specimen movements. *J Struct Biol* **152**, 36-51 (2005).

56.   X. Li *et al.*, Electron counting and beam-induced motion correction enable near-atomic-
resolution single-particle cryo-EM. *Nat Methods* **10**, 584-590 (2013).

57.   J. R. Kremer, D. N. Mastronarde, J. R. McIntosh, Computer visualization of three-dimensional
image data using IMOD. *J Struct Biol* **116**, 71-76 (1996).

58.   A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron
micrographs. *J Struct Biol* **192**, 216-221 (2015).

59.   Q. Xiong, M. K. Morphew, C. L. Schwartz, A. H. Hoenger, D. N. Mastronarde, CTF determination
and correction for low dose tomographic tilt series. *J Struct Biol* **168**, 378-387 (2009).

60.   Y.-T. Liu *et al.*, Isotropic Reconstruction of Electron Tomograms with Deep Learning. *bioRxiv*
10.1101/2021.07.17.452128, 2021.2007.2017.452128 (2021).

61.   J. M. Heumann, A. Hoenger, D. N. Mastronarde, Clustering and variance maps for cryo-electron
tomography using wedge-masked differences. *J Struct Biol* **175**, 288-299 (2011).

62.   T. D. Goddard *et al.*, UCSF ChimeraX: Meeting modern challenges in visualization and analysis.
Protein Sci **27**, 14-25 (2018).

63.   C. S. Hughes *et al.*, Ultrasensitive proteome analysis using paramagnetic bead technology. *Mol*
Syst Biol **10**, 757 (2014).

64.   W. Deng, J. Sha, K. Plath, J. A. Wohlschlegel, Carboxylate-Modified Magnetic Bead (CMMB)-
Based Isopropanol Gradient Peptide Fractionation (CIF) Enables Rapid and Robust Off-Line
Peptide Mixture Fractionation in Bottom-Up Proteomics. *Molecular & cellular proteomics : MCP*
20, 100039 (2021).

65.   J. Cox, M. Mann, MaxQuant enables high peptide identification rates, individualized p.p.b.-range
mass accuracies and proteome-wide protein quantification. *Nat Biotechnol* **26**, 1367-1372
(2008).

66.   M. Aslett *et al.*, TriTrypDB: a functional genomic resource for the Trypanosomatidae. *Nucleic*
Acids Res. **38**, D457-462 (2010).

67.   T. Huang *et al.*, MSstatsTMT: Statistical Detection of Differentially Abundant Proteins in
580   Experiments with Isobaric Labeling and Multiple Mixtures. *Molecular & cellular proteomics :*
581   *MCP* **19**, 1706-1723 (2020).
582   68.   J. Zhang *et al.*, Structure of the trypanosome paraflagellar rod and insights into non-planar
583   motility of eukaryotic cells. *Cell Discov* **7**, 51 (2021).

FIGURES

Figure 1

Figure 1. FAP106 is required for parasite motility. 29-13 (Ctrl) and FAP106 knockdown (FAP106 KD) parasites were grown in the presence of tetracycline to induce knockdown. (*A*) qRT-PCR analysis of FAP106 mRNA levels. Graph shows the mean $\pm$ standard deviation from three independent biological replicates. **** Unpaired, two-tailed t-test p-value $\leq$ 0.0001. (*B*) Cumulative growth curve shows the mean cell density vs time $\pm$ standard deviation from three independent biological replicates. (*C*) Histogram shows the length distribution of purified, demembranated flagella. Inset shows a representative phase contrast image. (*D*) Histogram shows the distribution of flagellum lengths as measured by anti-paraflagellar rod (PFR) immunofluorescence microscopy in detergent-extracted cytoskeletons. Inset shows a representative image of detergent-extracted cytoskeletons with PFR in red. (*E*) Graph shows the mean $\pm$ standard deviation of flagellum lengths in *C* (Phase Contrast) and *D* (anti-PFR). Phase contrast: Ctrl = 22.0 $\pm$ 2.5 µm, FAP106 KD = 19.0 $\pm$ 3.1 µm; anti-PFR: Ctrl = 17.6 $\pm$ 2.7 µm, FAP106 KD = 14.2 $\pm$ 3.2 µm. N=50 flagella for each (*F*) Motility tracks of individual parasites. (*G*) Mean squared displacement of parasites tracked in *F*. Dotted lines indicate the upper and lower bounds of the standard error of the mean. Ctrl N=489; FAP106 KD N=675. Data from an independent biological experiment are shown in Supp. Fig. S1.

Figure 2

Figure 2. FAP106 is required for assembly of conserved and lineage-specific MIP structures. Demembranated flagella were purified from 29-13 (Ctrl) and FAP106 knockdown (FAP106 KD) parasites grown in tetracycline to induce knockdown. Structure of the 48-nm repeat of the DMT was obtained by cryoET, with subtomographic averaging. Densities are colored according to the legend and based on similarity to published structures (13, 16, 20). (*A*) Schematic illustrating a cross-section of the '9+2' axoneme and paraflagellar rod (PFR) viewed from flagellum tip to base. Adapted from (68). Enlarged panel shows a single DMT, with A- and B-tubules indicated (A_t, B_t) and protofilaments numbered. FAP106 (red) and the inner junction filament (IJ) comprised of alternating PACRG (blue) and FAP20 (purple) are colored for reference. Viewing angles shown in *C-E* are indicated. (*B*) Cross-sectional view of the DMT. Colored densities indicate structures reduced or missing in FAP106 KD compared to Ctrl. (*C* and *D*) Longitudinal views of the DMT. Viewing angles are indicated in *A, inset*. The 48-nm repeat is indicated. Note that *T. brucei* has a structure similar to FAP210 of *Chlamydmonas* (green) despite not having a clear FAP210 homolog (Table 1). (*E*) Longitudinal views of the DMT, showing the IJ filament. Viewing angles are indicated in *A, inset.* The 48-nm repeat is indicated. Black arrowheads indicate the position of the single hole found within each 48-nm repeat in Ctrl DMTs. FAP20 and PACRG subunits are colored according to the legend, with holes corresponding to missing PACRG-B subunits, as suggested by TMT proteomics (Figure 3).

Figure 3. TMT proteomics identifies novel, FAP106-dependent, lineage-specific MIP proteins. (*A*) Demembranated flagella were purified from 29-13 (Ctrl) or FAP106 KD parasites grown in tetracycline to induce knockdown. Volcano plot shows significance (-Log$_{10}$ Adjusted P-value) vs. relative abundance (Log$_2$ Fold-Change) for all proteins quantified by TMT proteomics. Relative abundance is calculated as FAP106 KD/Ctrl. Results are from two independent biological samples. Proteins that met the filtering criteria (2-fold decrease, adjusted P-value $\leq$0.06; Table 2; Supp. Dataset S1) are indicated by red dots. Labels indicate known proteins (red text) and putative MIP candidates (green text). Asterisks indicate proteins that were independently identified as putative B-tubule MIP candidates in a separate proximity labeling approach (Supp. Fig. S3; Supp. Dataset S2). (*B*) Cross-sectional view of the DMT as viewed from the flagellum tip, showing the positions of MIP structures lost in FAP106 KD as observed by cryoET (Fig. 2), with known MIPs colored. A- and B-tubules are indicated (A$_t$, B$_t$) and protofilaments are numbered. (*C*) Motility analysis showing mean squared displacement of 29-13 (Ctrl) and MC KD parasites from $\geq$2 independent biological replicates. Dotted lines indicate the upper and lower bounds of the standard error of the mean. Ctrl N=1761; MC3 KD N=627; MC5 KD N=753; MC8 KD N=709, MC15 KD N=751. Individual replicates are shown in Supp. Fig. S3B. (*D*) Detergent-extracted cytoskeletons were prepared from the indicated KD parasites or their respective mNeonGreen (NG)-tagged parental cell lines (Ctrl). Graph shows the mean $\pm$ standard deviation of flagellum lengths as measured by anti-PFR immunofluorescence microscopy. MC3-NG (Ctrl) = 19.9 $\pm$ 2.3 µm, MC3 KD = 19.3 $\pm$ 1.8 µm; MC5-NG (Ctrl) = 18.9 $\pm$ 2.4 µm, MC5 KD = 19.1 $\pm$ 2.3 µm; MC8-NG (Ctrl) = 19.6 $\pm$ 2.9 µm, MC8 KD = 18.0 $\pm$ 2.5 µm, MC15-NG (Ctrl) = 18.4 $\pm$ 2.1 µm, MC15 KD = 16.7 $\pm$ 2.6 µm. N=50 flagella for each, except N=45 flagella for MC15-NG. Unpaired, two-tailed t-test: ns = p $\geq$ 0.05, ** = p $\leq$ 0.01 and *** = p $\leq$ 0.001.

Figure 4. MC8 is a novel, lineage-specific *T. brucei* MIP that comprises the MIP B8 density. (*A*) AlphaFold prediction for MC8 (42, 43), colored according to confidence. Box indicates high-confidence region modeled in *B*. (*B*) MC8 AlphaFold prediction modeled into MIP B8 cryoET density. (*Left*) Cross-sectional view of the DMT, with the MIP B8 density colored aqua. (*Middle, Right*) Enlarged view of MIP B8 density with AlphaFold prediction shown in aqua. Viewing angle shown in *Right* panel is indicated by the dashed line in *Middle* panel. (*C-E*) Demembranated flagella were purified from 29-13 (Ctrl) and MC8 knockdown (MC8 KD) parasites grown in tetracycline to induce knockdown. Structure of the 48-nm repeat of the DMT was obtained by cryoET, followed by subtomographic averaging. (*C*) Cross-sectional view of the DMT with MIP B8 density colored in aqua. (*D* and *E*) Longitudinal views of the DMT. Viewing angles and colors are as indicated in Fig. 2.

Figure 5. FAP106 is required for parasite motility and assembly of conserved and lineage-specific MIPs at the inner junction. Normal parasite motility relies on FAP106 as a critical interaction hub for assembly of MIPs at the inner junction (IJ). Cross-sectional and longitudinal views of the DMT are illustrated. Conserved IJ MIPs PACRG-A, FAP20 and FAP52 assemble independently of FAP106, however PACRG-B does not stably assemble into the IJ filament without FAP106. The B-tubule remains intact and connected to the A-tubule at the IJ, despite loss of multiple IJ MIPs. FAP106 is also critical for assembly of the conserved MIP, FAP45, and a FAP210-like protein (modeled as filaments based on published structures (13, 16, 20)), as well as several novel, lineage-specific MIPs.

Movie Legends

Movie 1. Subtomographic average of the 48-nm repeat. Video showing 3D reconstruction of 48-nm repeat from 29-13 (Control) and FAP106 KD DMTs, rotating between views shown in Fig 2.

679 **Table 1. Known B-tubule MIPs**

Chlamydomonas	Mammals	*Tetrahymena*	*Trypanosoma*
†PACRG	PACRG	TTHERM_00446290; TTHERM_00499310; TTHERM_00499570	Tb927.3.2310; Tb927.9.9940
†FAP20	CFAP20	TTHERM_00418580	Tb927.10.2190
FAP45	CFAP45	TTHERM_001164064	Tb927.8.4580
FAP52	CFAP52	TTHERM_01094880	Tb927.11.7560
FAP106	ENKUR	TTHERM_00137550; TTHERM_00370750	Tb927.11.4880
FAP77	CFAP77	TTHERM_00974270; TTHERM_01190430	
FAP210	CFAP210	TTHERM_00643530	
FAP90	CFAP90		
FAP126	Flattop		
FAP144	FAM183A		
FAP276	CFAP276		
FAP112		TTHERM_00884620	
	EFCAB6		
	SPACA9		
			Tb927.11.2770 (MIP B8, this study,‡)

681 †FAP20 and PACRG form the IJ filament

682 ‡Corresponding structure was called 'MIP3c' in (15) but our current study suggests this structure
683 is distinct from MIP3c (FAP45).

685 **Table 2. Proteins significantly reduced in FAP106 KD flagellum skeletons by TMT**
686 **proteomics**

Gene	Gene ID	Fold Reduction	Adjusted P-value	Protein Localization	Function
FAP106	Tb927.11.4880	24.02	0.0011	Flagellum	B-tubule MIP
FAP45	Tb927.8.4580	3.95	0.0035	Flagellum	B-tubule MIP
PACRG-B	Tb927.9.9940	2.77	0.0576	Flagellum	IJ filament
MC5	Tb927.11.4920	13.74	0.0003	Flagellum	MIP candidate*
MC3	Tb927.10.7120	3.44	0.0044	Flagellum	MIP candidate*
MC8	Tb927.11.2770	2.81	0.0111	Flagellum	MIP candidate*
MC15	Tb927.3.3200	1.95	0.0177	Flagellum	MIP candidate*
RPC19	Tb927.11.8890	2.99	0.0055	Nucleus	RNA polymerase

688 *Identified by independent APEX2-based proximity proteomics of known B-tubule MIPs (see
689 Supp. Dataset S2)