Male mammalian meiosis and spermiogenesis is critically dependent on the shared

functions of the katanins KATNA1 and KATNAL1

Authors

Jessica EM Dunleavy[1], Maddison Graffeo[1], Kathryn Wozniak[2], Anne E O'Connor[1], D. Jo

Merriner[1], Joseph Nguyen[1], Ralf B Schittenhelm[3], Brendan J Houston[1], Moira K O'Bryan[1]

[1]School of BioSciences and Bio21 Institute, Faculty of Science, The University of Melbourne,

VIC, 3010, Australia. [2]Monash Biomedicine Discovery Institute and The Department of

Anatomy and Developmental Biology, Monash University, Melbourne VIC, 3800, Australia

[3]Monash Proteomics & Metabolomics Facility, Department of Biochemistry and Molecular

Biology, Biomedicine Discovery Institute, Monash University, Clayton VIC, 3800, Australia

*Correspondence: Jessica.dunleavy@unimelb.edu.au

Summary

Katanin microtubule severing enzymes are potent M-phase regulators in oocytes and somatic

cells. How the complex, and evolutionarily critical, male mammalian meiotic spindle is

sculpted remains unknown. Here, using multiple single and double gene knockout mice, we

reveal that the canonical katanin A-subunit, KATNA1, and its close paralogue, KATNAL1,

together execute multiple aspects of meiosis. We show KATNA1 and KATNAL1 collectively

regulate the male meiotic spindle, cytokinesis and midbody abscission, in addition to diverse

spermatid remodelling events, including Golgi organisation, and acrosome and manchette

formation. We also define KATNAL1-specific roles in sperm flagella development, manchette

regulation, and sperm-epithelial disengagement. Finally, using proteomic approaches we

define the KATNA1, KATNAL1, and KATNB1 mammalian testis interactome, which

includes a network of cytoskeletal and vesicle trafficking proteins. Collectively, we reveal the

presence of multiple katanin A-subunit paralogs in mammalian spermatogenesis allows for

'customized cutting' via neofunctionalization and protective buffering via gene redundancy.

Keywords: Meiotic spindle, spermatogenesis, microtubule severing, manchette, axoneme,

acrosome, HTCA, male fertility, vesicle transport, spermiation

Introduction

The katanins are an evolutionary conserved family of microtubule (MT) severing proteins that

were first identified as an essential meiosis regulator in *Caenorhabditis elegans* oocytes[1]. Since

then, they have been established as a nuanced regulator of MTs, able to employ the same action

(physical removal of tubulin heterodimers from the MT lattice) to amplify, disassemble or even

strengthen MTs (for reviews see[2,3]). At a cellular level these actions drive rapid MT

remodelling events required in many processes including ciliogenesis, cell division and

differentiation, and allow fine tuning of homeostatic MT structures and processes.

The canonical katanin complex consists of an AAA ATPase catalytic severing enzyme

KATNA1 (aka p60 or A-subunit) and a WD40-repeat containing regulatory subunit KATNB1

(aka p80 or B-subunit) and is conserved across most eukaryotes. In its active ATP bound state,

the A-subunit forms a spiral hexamer which binds MTs and threads the C-terminal tail (CTT)

of a tubulin subunit through its central pore. An ATP-driven conformational change in the

hexamer, causes an extractive force on the tubulin CTT removing the tubulin heterodimer from

the lattice. In higher eukaryotes, the B-subunit is not essential for A-subunit severing activity,

however it increases MT affinity and severing efficiency, and at a cellular level is often

essential for A-subunit localisation[2]. In vertebrates, *Drosophila,* and unicellular ciliates,

paralogous A-subunits, KATNAL1 and KATNAL2 are present, and an additional WD40-less

KATNB1 paralog, KATNBL1 is present in vertebrates. KATNAL1 is highly similar in

sequence and domain architecture to KATNA1[2] and has been shown to possess MT-severing

activity[4]. KATNAL2 however, outside of the conserved AAA ATPase MT severing domain,

differs in domain architecture to KATNA1 and KATNAL1[2] and in standard MT severing

assays did not exhibit MT severing activity[4,5]. This aside, loss of KATNAL2 function in mouse

spermatogenesis resulted in phenotypes consistent with loss of MT severing[5].

Across eukaryotes, the regulation of cell division is a defining feature of katanin action. In *C.*

elegans oocytes, the orthologues of KATNA1 and KATNB1, MEI-1 and MEI-2 drive

formation of the bipolar spindle in meiosis I via MT amplification, and subsequently regulate

spindle length, function and positioning[2,6] and in meiosis II, mediate spindle disassembly[7].

KATNA1 has also been shown to regulate female meiotic spindle length in *Xenopus laevis* and

Xenopus tropocalis[8], and in cultured mouse oocytes KATNAL1 depletion results in abnormally

long meiotic spindles and an increase in multipolar spindles [9]. Likewise, in mitosis, katanin

A-subunits have variously been shown to regulate mitotic spindle size/shape, polarity, density

and orientation, and drive midbody abscission during cytokinesis in species ranging from

unicellular ciliates and plants to mammalian cell lines[2,6,10]. Despite these data, the contribution

of katanin proteins to male meiosis has received little attention. The requirement, and identity,

of katanin A-subunits in male mammalian meiosis has not been directly tested, although data

from loss-of-functions in the regulatory, B-subunit, KATNB1, strongly suggest katanin-

mediated MT severing is required[11,12].

Herein we aimed to identify the function of katanin-A subunits in male mammalian meiosis.

Using a suite of single and double katanin A-subunit deletion mice, we reveal KATNA1 and

KATNAL1 collectively regulate male meiosis, wherein they refine metaphase spindle structure

and chromosome alignment and drive accurate chromosome segregation. We show their

functions extend into haploid germ cell development (spermiogenesis), a process that

necessitates the development of several specialised MT structures. We identify KATNA1 and

KATNAL1 as global regulators of spermatid MTs required to restrain MT bulk, and as well as

identifying several structure/process specific roles, including in Golgi function and acrosome

biogenesis. We further identify KATNAL1 specific roles in cytokinesis during meiosis, in the

formation of the sperm head-to-tail coupling apparatus (HTCA) and axoneme, in sperm head

sharping via manchette MT dynamics and dissolution, and in sperm release (spermiation).

During spermatogenesis, we reveal KATNAL1 can fully functionally compensate for loss of

KATNA1. In contrast, we show KATNA1 can only partially compensate for KATNAL1 loss,

and there is no functional compensation between KATNA1 and KATNAL2 in the mouse testis.

Finally, using an *in vivo* immunoprecipitation (IP)-mass spectrometry (MS) approach, we

reveal these diverse cellular functions are mediated by an equally diverse set of KATNA1,

KATNAL1 and KATNB1 testis interacting partners, including MT associated proteins,

regulators of actin dynamics and organisation, vesicle-transport proteins including multiple

Rab GTPase family members, and essential mediators of male meiosis and acrosome

biogenesis. Collectively these results reveal the complex reliance of spermatogenesis on

katanin function and the precise neofunctionalization of A-subunits required to achieve normal

sperm function and morphology.

Results

KATNA1 is not essential for male fertility in the mouse

Despite the clear necessity for the dynamic regulation of MT structures during meiosis, and

data showing that the loss of the KATNB1 leads to notable defects in meiosis[11,12], the 'effector'

A-subunit is unknown. Data from *C. elegans* reveal the KATNA1 orthologue MEI-1 is a key

regulator of female meiosis. To test the hypothesis that KATNA1 serves an analogous function

in male meiosis, we generated a *Katna1* germ cell specific knockout mouse model

(*Katna1$^{GCKO/GCKO}$*, Fig S1A-C). As shown in Fig S1D, *Katna1* mRNA expression was reduced

by 91.77% in *Katna1$^{GCKO/GCKO}$* spermatocytes compared to *Katnal1$^{Flox/Flox}$* controls. Loss of

KATNA1 at the protein level was confirmed via immunolabelling of testis sections (Fig S1E).

Katna1$^{GCKO/GCKO}$ mice had normal body weight (Fig S2A) and unexpectedly were fertile.

Katna1$^{GCKO/GCKO}$ mice exhibited normal testis weight, testis daily sperm production and

epididymal sperm content (Fig S2B,C,D). Histological assessment of the male reproductive

tract in *Katna1$^{GCKO/GCKO}$* mice revealed normal progression of germ cells through all phases of

spermatogenesis and the presence of sperm in the epididymis (Fig S2E,F,G). Sperm from

Katna1$^{GCKO/GCKO}$ mice were cytologically comparable to those from *Katnal1$^{Flox/Flox}$* controls

(Fig S2H) and were functionally competent, as exhibited by normal fertility when mated with

wild-type females (8.2 pups per copulatory plug in *Katna1*$^{GCKO/GCKO}$ versus 7.1 pups in

controls, P=0.0969). Collectively, these data reveal KATNA1 is dispensable for mammalian

spermatogenesis.

Germ cell derived KATNAL1 is required for optimal male fertility in mice

We next sought to explore the requirement in meiosis for the non-canonical katanin A-subunit,

KATNAL1. A role for KATNAL1 in the somatic Sertoli cell population of the testis has been

defined [13], but a specific role in male meiosis has not been tested. To do this we generated a

Katnal1 germ cell specific knockout mouse model (*Katnal1*$^{GCKO/GCKO}$, Fig S3A,B). Successful

ablation of *Katnal1* gene expression was confirmed by qPCR, western blotting of purified germ

cells and immunolabeling of testis sections (Fig 3C-E). There was a 99.56% reduction in

Katnal1 mRNA in *Katnal1*$^{GCKO/GCKO}$ compared to *Katnal1*$^{Flox/Flox}$ germ cells.

Katnal1$^{GCKO/GCKO}$ males exhibited normal mating frequency but were subfertile siring 84%

fewer pups per copulatory plug than *Katnal1*$^{Flox/Flox}$ controls (6.9 ± 3.1SD pups / plug sired by

Katnal1$^{Flox/Flox}$ males versus 1.4 ± 1.6SD pups / plug sired by *Katnal1*$^{GCKO/GCKO}$ males,

P=0.0151). As outlined below, this subfertility was due to a dramatic reduction in sperm

number, and abnormal sperm morphology and motility. *Katnal1*$^{GCKO/GCKO}$ mice had body and

testis weights comparable to *Katnal1*$^{Flox/Flox}$ mice (Fig 1A,B). Likewise, the number of

apoptotic germ cells per seminiferous tubules was not significantly different between

Katnal1$^{Flox/Flox}$ and *Katnal1*$^{GCKO/GCKO}$ mice (Fig 1D). Despite this, *Katnal1*$^{GCKO/GCKO}$ testis

daily sperm production was reduced by 42.3% compared to *Katnal1*$^{Flox/Flox}$ mice (Fig 1C),

indicative of germ cell loss in the later steps of spermiogenesis, when spermatids contribute

minimally to the overall weight of the testis. Finally, epididymal sperm content was reduced

by 73.7% compared to *Katnal1^Flox/Flox* mice (Fig 1D). This reveals that germ cell derived

KATNAL1 is required for optimal spermiogenesis and, as evidenced by the more severe

reduction in epididymal sperm content relative to DSPs, for their ultimate release from the

seminiferous epithelium via the process of spermiation. Assessment of *Katnal1^GCKO/GCKO*

seminiferous tubule histology confirmed the abnormal retention of elongated spermatids in

stage IX tubules (Fig 1G).

While all germ cell types were present in the *Katnal1^GCKO/GCKO* seminiferous epithelium, many

Katnal1^GCKO/GCKO tubules contained fewer germ cells, notably spermatids, compared to

Katnal1^Flox/Flox mice (Fig 1E,F). Areas of Sertoli cell-only tubules were observed and gaps

between adjacent Sertoli cells were frequently seen in *Katnal1^GCKO/GCKO* but never in

Katnal1^Flox/Flox (Fig 1E,F). Epididymal histology was consistent with the presence of fewer

sperm in *Katnal1^GCKO/GCKO*, in addition to the presence of prematurely sloughed germ cells in

Katnal1^GCKO/GCKO, but not *Katnal1^Flox/Flox*, mice (Fig 1H). Of note, immunolabelling testis

sections for the Sertoli cells specific β-tubulin isoform, TUBB3, confirmed the origins of the

reduced spermatogenic output in the *Katnal1^GCKO/GCKO* were distinct from those of the

previously characterised *Katnal1^1H/1H* hypomorphic mouse model where extensive disruption

of the Sertoli cell MT cytoskeleton was seen [13]. In contrast, in *Katnal1^GCKO/GCKO* testis sections

TUBB3 localisation was comparable to *Katnal1^Flox/Flox* (Fig S4A-B). These data reveal that

KATNAL1 plays an essential role in the regulation of MTs in both Sertoli and germ cells.

Of the sperm present in the epididymides of *Katnal1^GCKO/GCKO* mice, almost all were

morphologically abnormal (97.1% in *Katnal1^GCKO/GCKO* versus 23.9% in *Katnal1^Flox/Flox*, Fig

1I,J). Abnormalities included in head shape (97.1% in *Katnal1^GCKO/GCKO* versus 11.3% in

Katnal1^{Flox/Flox}), head position (36.4% versus 2.3%), coupling between the head and tail

(decapitation, 34.7% versus 11.5%), mid-pieces structure (coiling, 14.4% versus 0.0%) and

mitochondrial sheath structure (3.9% versus 0.0%) (Fig 1J). Moreover, of those sperm found

in *Katnal1^{GCKO/GCKO}* epididymides, there was a 20.2% reduction in the ability for motility and

a 52.9% reduction in progressive motility in the absence versus presence of KATNAL1 (Fig

2K,L). Considering the reduction in sperm number in *Katnal1^{GCKO/GCKO}* mice, this equates to

an 87.6% reduction in the total number of progressively motile sperm in *Katnal1^{GCKO/GCKO}*

compared to *Katnal1^{Flox/Flox}* mice. Collectively these phenotypes reveal germ cell derived

KATNAL1 is required for normal spermatogenesis and optimal male fertility.

KATNAL1 regulates chromosome alignment, segregation and cytokinesis during male

meiosis

The reduction in post-meiotic germ cells in *Katnal1^{GCKO/GCKO}* seminiferous tubules indicated a

variably penetrant meiotic defect. To assess *Katnal1^{GCKO/GCKO}* meiotic spindles, testis sections

were stained for α-tubulin as a component of MTs (Fig 2A,B). As seen in *Katnal1^{Flox/Flox}* stage

XII tubules, metaphase I and II meiotic spindle MTs typically project from the two opposing

spindle poles in a narrow and uniform angle to tightly align chromosomes/chromatids along

the metaphase plate (Fig 2Aa,2Ba-c). In *Katnal1^{GCKO/GCKO}* stage XII tubules, the vast majority

of metaphase spindles displayed normal spindle architecture and normal MT spindle density

(Fig 2Ae, Bf-h). Chromosomal alignment at the metaphase plate, however, was noticeably

abnormal in many *Katnal1^{GCKO/GCKO}* metaphase spermatocytes, with misaligned chromosomes

often observed (Fig 2A,Bf-g, green arrowheads). These defects did not however prevent

progression to anaphase, with no increase in metaphase arrest and apoptosis in stage XII-I

seminiferous tubules in *Katnal1^{GCKO/GCKO}* compared to *Katnal1^{Flox/Flox}* mice (Fig 2A,E-J).

Similarly, KATNAL1 loss in germ cells did not prevent spermatocyte progression through

anaphase and telophase. Specifically there was no discernible increase in $Katnal1^{GCKO/GCKO}$

spermatocytes stalling in anaphase or telophase or undergoing apoptosis (Fig 2A,E-J). Instead,

anaphase and telophase defects in $Katnal1^{GCKO/GCKO}$ meiosis led to the production of abnormal

haploid germ cells. In anaphase I and II, spindle MTs undergo poleward shortening, allowing

them to pull chromosomes toward opposing poles in a coordinated manner, as observed in

$Katnal1^{Flox/Flox}$ anaphase spermatocytes (Fig 2Ab,Bd-e). In $Katnal1^{GCKO/GCKO}$ anaphase

spermatocytes however, uneven segregation of chromosomes was often observed, with lagging

chromosomes commonly seen (Fig 2Af, Bi-j, green arrowheads). Consistent with uneven

chromosome segregation in anaphase, many round spermatids possessed abnormally large

nuclei (Fig 2Cb-c, yellow arrowheads). During telophase and cytokinesis midbodies were

present in $Katnal1^{GCKO/GCKO}$ and were comparable to those in $Katnal1^{Flox/Flox}$ spermatocytes

(Fig 2Ac,g). However, a subset of the resulting spermatids was binucleated (Fig 2Cd,e, orange

arrowheads), and as detailed below sperm/spermatids with multiple axonemes (Fig 3C) and

supernumerary centrioles (Fig 3Db, red arrowhead) were often observed. This indicates that

while midbodies do form in the absence of *Katnal1*, KATNAL1 plays a key role its remodelling

into an intracytoplasmic bridge and the maintenance of individual daughter spermatids.

Binucleated spermatids were never observed in $Katnal1^{Flox/Flox}$ tubules.

These data reveal KATNAL1 regulates MT-chromosome attachment and/or metaphase spindle

dynamics but is dispensable for MT generation and pruning during spindle biogenesis in male

meiosis. Moreover, our data show KATNAL1 greatly facilitates accurate chromosome

segregation during anaphase and midbody function and disassembly during cytokinesis in

either meiosis I and/or meiosis II.

KATNAL1 regulates flagella and manchette formation in male germ cells

As detailed above, of the epididymal sperm that were produced in $Katnal1^{GCKO/GCKO}$ mice 97.1% were morphologically abnormal revealing that KATNAL1 is required during spermatid differentiation (spermiogenesis) (Fig 1I,J). One of the first processes to commence in spermiogenesis is acrosome formation, and our recent data identified MT severing as a potential regulator of this process [11]. The identity of the involved MT severing A-subunit remains unknown. Acrosome formation was overtly normal in $Katnal1^{GCKO/GCKO}$ mice compared to $Katnal1^{Flox/Flox}$ controls, indicating KATNAL1 is not essential for this process (Fig S4C).

Multiple abnormalities were however observed in $Katnal1^{GCKO/GCKO}$ mouse sperm tails and the head-to-tail coupling apparatus (HTCA), including abnormal reticulation of the sperm head, decapitation, coiling of the sperm tail mid-piece and mitochondrial sheath defects, in addition to reduced sperm motility. Staining of $Katnal1^{GCKO/GCKO}$ testis sections for acetylated tubulin, as a marker of stable MTs, revealed axonemal MTs were present in sperm tails (Fig 3B), however analysis via transmission electron microscopy (TEM) revealed a range of axoneme and accessory structure abnormalities (Fig 3C). In $Katnal1^{GCKO/GCKO}$ mice, sperm flagella axonemes often lacked the typical 9+2 MT doublet arrangement observed in controls (Fig 3Ca,d,g). Instead MT doublets were often missing (Fig 3Ce,f,h, yellow asterisks), ectopically located (Fig 3C, red arrowheads) or in many cases supernumerary MT doublets were present (Fig 3Cc,l, white arrowheads). In the most dramatic instances, and consistent with the failures in cytokinesis, multiple axonemes, often each with associated accessory structures, were observed within one cell (Fig 3Cb,j-l). The abnormalities in MT doublet number and location were typically accompanied by defects in outer dense fiber (ODF) number and location (Fig 3C). In wild-type sperm, the mid-piece contains 9 ODFs closely associated with the 9 outer

MT doublets of the axoneme (Fig 3Ca). In the principal piece ODFs 3 and 8 are replaced by

the longitudinal columns of the fibrous sheath (Fig 3Cd). In *Katnal1*$^{GCKO/GCKO}$ mice however,

sperm tail ODFs 3 and/or 9 frequently persisted into the principal piece (Fig 3Cb,c, green

arrowheads). In *Katnal1*$^{GCKO/GCKO}$ mouse sperm tails wherein some or all axoneme MTs were

missing, ODFs were often also missing (Fig 3Ce,f). Consistent with the coiled mid-pieces and

mitochondrial sheath defects observed in some sperm from *Katnal1*$^{GCKO/GCKO}$ mice, TEM

revealed that in many sperm the mitochondria were not properly organised into a helical sheath

(Fig 3Cj). Fibrous sheath development in *Katnal1*$^{GCKO/GCKO}$ sperm however, appeared normal

in most cells (Fig 3C). Finally, in some sperm, only disorganised fragments of sperm tail

components were observed indicative of degenerating cells (Fig 3Cm). Collectively, these data

reveal KATNAL1-mediated MT severing regulates the assembly and potentially stability of

axoneme MT doublets, and that KATNAL1 contributes to the development of the

mitochondrial sheath and ODFs, but not the fibrous sheath. These data are consistent with

defects in cargo/organelle delivery to and along the growing sperm tail. Indeed, there is an

increasing of body of evidence (reviewed in[14]) that mitochondria and ODFs rely on MT-based

transport mechanisms for their delivery into the sperm tail.

The frequent decapitation or hyper-extension of sperm heads, in sperm from *Katnal1*$^{GCKO/GCKO}$

males, indicates the HTCA was weak and/or improperly formed in the absence of KATNAL1.

Moreover, TEM of round and elongated spermatids revealed a range of HTCA abnormalities

in *Katnal1*$^{GCKO/GCKO}$ mice compared to controls (Fig 3D). In *Katnal1*$^{Flox/Flox}$ mice, spermatid

HTCAs were easy to find in testis sections via TEM. As expected, they contained a proximal

centriole coupled to the nucleus via the basal plate and capitulum, a distal centriole coupled to

the plasma membrane and continuous with the axoneme, and the surrounding segmented

columns which are continuous with the ODFs (Fig 3Da,f). In *Katnal1*$^{GCKO/GCKO}$ mice, however

normal HTCAs were rare (Fig 3Db,g). In most spermatids, HTCAs were abnormal (Fig 3Dc-

e, h-j). In some cells, the HTCA and associated axoneme was present, but not attached to the

nucleus (Fig 3Dc,h, yellow arrowheads), while in others the basal plate could be seen

associated with the nucleus, but the rest of the HTCA was absent (Fig 3Dg). Consistent with

this data, immunolabelling of sperm for SUN5, which is an essential component of the LINC

complex that physically docks the sperm centrosome to the nucleus, revealed a 57.2%

reduction in sperm from *Katnal1*^{GCKO/GCKO} males containing SUN5 at the HTCA compared to

sperm from *Katnal1*^{Flox/Flox} mice (Fig S4D,E). These data suggest that KATNAL1 regulated-

MTs are either required to fortify the connection between the basal plate and the nucleus or are

required for the delivery and deposition of essential components of the HTCA. Our data also

suggest that SUN5 is delivered to the HTCA via KATNAL1-regulated MTs.

Abnormal head shape was the most frequent morphological abnormality present in the sperm

of *Katnal1*^{GCKO/GCKO} mice. Instead of the normal hook shape exhibited in 88.75% of

Katnal1^{Flox/Flox} mouse sperm, 34% of *Katnal1*^{GCKO/GCKO} mouse sperm heads exhibited a

'hammerhead' shape and 62% exhibited a 'knobby' head phenotype (Fig 1I, S4F,G). Abnormal

head shaping became apparent during nuclear elongation (Fig 3E) and were consistent with

defects in manchette function, where the manchette is a transient MT-based structure which

sculpts the distal half of the spermatid head. As shown in Fig 3F in wild-type mice the

manchette forms in step 8-9 spermatids (stage VIII-IX seminiferous tubules), before migrating

distally while simultaneously constricting to shape the spermatids head during steps 10-12

(stages X-XII; Fig 3F). In steps 13-14 (stage I-III) the manchette is dissembled (Fig 4F). In

spermatids from *Katnal1*^{GCKO/GCKO} mice, manchettes formed at step 8-9 but were abnormally

elongated. Elongation became increasingly pronounced as spermatid development continued

and was associated with a failure of manchette perinuclear ring migration down the sperm head,

normal HTCAs were rare (Fig 3Db,g). In most spermatids, HTCAs were abnormal (Fig 3Dc-

e, h-j). In some cells, the HTCA and associated axoneme was present, but not attached to the

nucleus (Fig 3Dc,h, yellow arrowheads), while in others the basal plate could be seen

associated with the nucleus, but the rest of the HTCA was absent (Fig 3Dg). Consistent with

this data, immunolabelling of sperm for SUN5, which is an essential component of the LINC

complex that physically docks the sperm centrosome to the nucleus, revealed a 57.2%

reduction in sperm from *Katnal1*$^{GCKO/GCKO}$ males containing SUN5 at the HTCA compared to

sperm from *Katnal1*$^{Flox/Flox}$ mice (Fig S4D,E). These data suggest that KATNAL1 regulated-

MTs are either required to fortify the connection between the basal plate and the nucleus or are

required for the delivery and deposition of essential components of the HTCA. Our data also

suggest that SUN5 is delivered to the HTCA via KATNAL1-regulated MTs.

Abnormal head shape was the most frequent morphological abnormality present in the sperm

of *Katnal1*$^{GCKO/GCKO}$ mice. Instead of the normal hook shape exhibited in 88.75% of

Katnal1$^{Flox/Flox}$ mouse sperm, 34% of *Katnal1*$^{GCKO/GCKO}$ mouse sperm heads exhibited a

'hammerhead' shape and 62% exhibited a 'knobby' head phenotype (Fig 1I, S4F,G). Abnormal

head shaping became apparent during nuclear elongation (Fig 3E) and were consistent with

defects in manchette function, where the manchette is a transient MT-based structure which

sculpts the distal half of the spermatid head. As shown in Fig 3F in wild-type mice the

manchette forms in step 8-9 spermatids (stage VIII-IX seminiferous tubules), before migrating

distally while simultaneously constricting to shape the spermatids head during steps 10-12

(stages X-XII; Fig 3F). In steps 13-14 (stage I-III) the manchette is dissembled (Fig 4F). In

spermatids from *Katnal1*$^{GCKO/GCKO}$ mice, manchettes formed at step 8-9 but were abnormally

elongated. Elongation became increasingly pronounced as spermatid development continued

and was associated with a failure of manchette perinuclear ring migration down the sperm head,

but continued constriction, and resulting in a 'knobby head' phenotype (Fig 4G). Finally,

manchette disassembly in steps 13-14 was delayed in the absence of KATNAL1 (Fig 4F).

Collectively, these results reveal KATNAL1-mediated MT severing is required to regulate

manchette MT length, migration and disassembly.

KATNA1 and KATNAL1 compensate for each other in male meiosis and spermiogenesis

Given the apparently normal fertility in $Katna1^{GCKO/GCKO}$ and residual fertility in

$Katnal1^{GCKO/GCKO}$ mice, we sought to explore the possibility of functional compensation,

between KATNA1 and its paralogs KATNAL1 and KATNAL2 in the testis. This is a

particularly critical question for meiosis wherein the identity of the critical A-subunit(s)

remains unknown. As reported previously, $Katnal2$ loss has no apparent effect on male meiosis

but does result in male infertility due to severe sperm developmental defects[5]. To test for

redundancy, we generated (1) $Katna1$ and $Katnal1$ double germ cell knockout mice

($Katna1/al1^{GCKO/GCKO}$) and (2) double $Katna1$ germ cell and $Katnal2$ knockout mice

($Katna1^{GCKO/GCKO}/al2^{KO/KO}$). The $Katna1^{GCKO/GCKO}/al2^{KO/KO}$ mouse model (Fig S5)

phenocopied our previously characterised $Katnal2$ mutant models[5], including having a

comparable spermiation failure phenotype, sperm head shaping defects (knobby heads), and

an absence of normal sperm tail development (Fig S5). No aspect of the infertility phenotype

appeared worse that the loss of KATNAL2 in isolation. These data reveal that KATNA1 and

KATNAL2 do not have compensatory function in male germ cell development.

By contrast the removal of KATNA1 and KATNAL1 from germ cells resulted in a significantly

worse male fertility phenotype than seen in either individual knockout, especially in meiosis.

Successful ablation of both $Katna1$ and $Katnal1$ from male germ cells was confirmed by qPCR

(Fig 4A,B). Test mating of the $Katna1/al1^{GCKO/GCKO}$ male mice with wild-type females,

revealed loss of both KATNA1 and KATNAL1 from male germ cells results in male sterility

(7.1±3.2 SD pups / plug sired by *Katna1/al1^Flox/Flox* males versus 0± 0 SD pups / plug sired by

Katna1/al1^GCKO/GCKO males (P=0.0175)). Moreover, assessment of the reproductive tract

revealed a 48.5% reduction in testis weight, an 89% reduction in daily sperm output and a

96.2% reduction in epididymal sperm content compared to controls (Fig 4D-F). Examination

of *Katna1/al1^GCKO/GCKO* seminiferous tubule histology instantly revealed large scale germ cell

loss during both meiosis and spermiogenesis that was significantly worse than in either single

gene deletion (Fig 4G-H). Numerous pyknotic spermatocytes were observed in all stage XII

and I *Katna1/al1^GCKO/GCKO* tubules indicative of cells stalling in meiosis followed by apoptosis

but were rare in *Katna1/al1^Flox/Flox* (Fig 4G-H, magenta arrowheads). Of the germ cells that

progressed through meiosis in *Katna1/al1^GCKO/GCKO*, the vast majority became pyknotic by

steps 12-13 (stages XII-I) of elongating spermatid development (Fig 4H, green arrowheads)

and as a result elongated spermatids were absent past stage II-VIII *Katna1/al1^GCKO/GCKO* tubules

(Fig 4H). Moreover, analysis of cleaved caspase 3/7 immunolabelling revealed a 6.4 fold

increase in the number of apoptotic germ cells in *Katna1/al1^GCKO/GCKO* versus

Katna1/al1^Flox/Flox seminiferous tubules (Fig 4O). The majority of apoptotic cells were

spermatocytes that had stalled in anaphase I and meiosis II in stage I tubules (Fig 5J,N), and to

a lesser degree those undergoing metaphase I in stage XII (Fig 5J,N). In line with these defects,

analysis of epididymal sections confirmed almost no sperm reached the *Katna1/al1^GCKO/GCKO*

epididymis. The bulk of the cells in the lumen of *Katna1/al1^GCKO/GCKO* epididymides were

instead prematurely sloughed round germ cells (Fig 4P, arrowheads).

Assessment of *Katna1/al1^GCKO/GCKO* meiotic spindles, via the immunolabelling of testis

sections for α-tubulin, revealed that while meiotic spindles formed in *Katna1/al1^GCKO/GCKO*

spermatocytes, most were morphologically abnormal compared to *Katna1/al1^Flox/Flox* controls

(Fig 5A,B). At metaphase, in contrast to the tightly packed morphology of *Katna1/al1$^{Flox/Flox}$*

controls (Fig 5Aa,Ba,e), chromosomes were loosely dispersed along the metaphase plate in

Katna1/al1$^{GCKO/GCKO}$ mice (4Bb,f). Misaligned and/or lagging chromosomes were frequently

observed in *Katna1/al1$^{GCKO/GCKO}$* metaphase and anaphase spermatocytes (Fig 5A,B). As

detailed above, these defects ultimately lead to a large proportion of *Katna1/al1$^{GCKO/GCKO}$*

metaphase and anaphase spermatocytes stalling and undergoing apoptosis in stage XII-I

seminiferous tubules (Fig 4H, I-O). Of note, examination of *Katna1/al1$^{GCKO/GCKO}$* stage I

tubules, revealed that these stalled metaphase and anaphase spermatocytes where characterised

by the ectopic accumulation of α-tubulin throughout their cytoplasm (Fig 5Ah,i) indicating

that KATNA1/KATNAL1-mediated severing restrains MT bulk in spermatocytes. Of the

subset of *Katna1/al1$^{GCKO/GCKO}$* spermatocytes that progressed through meiosis, overtly normal

midbody formation was observed in telophase (Fig 5Ag,k). However, indicative of failure of

midbody to intracellular bridge conversion and similar to *Katna1$^{GCKO/GCKO}$* mice, binucleated

spermatids were frequently observed in *Katna1/al1$^{GCKO/GCKO}$* mice but never in

Katna1/al1$^{Flox/Flox}$ controls (Fig 5Ag,k). Consistent with compromised anaphase chromosome

segregation, abnormally large round spermatids were frequent in *Katna1/al1$^{GCKO/GCKO}$* mice

but never in *Katna1/al1$^{Flox/Flox}$* controls (Fig 5Ag,h). Collectively these data indicate that

KATNAL1 and KATNA1 can at least partially compensate for each other in metaphase and

anaphase of male meiosis. The frequency of binucleated spermatids in *Katna1/al1$^{GCKO/GCKO}$*

mice however, was similar to that of *Katnal1$^{GCKO/GCKO}$* suggesting KATNA1 does not

compensate for KATNAL1 in cytokinesis.

Assessment of spermiogenesis in *Katna1/al1$^{GCKO/GCKO}$* mice revealed that unlike the single

GCKO mice, acrosome biogenesis was overtly abnormal in the absence of both KATNA1 and

KATNAL1 indicating functional compensation between these two A-subunits (Fig 6A,B). In

PAS-stained *Katna1/al1*^{Flox/Flox} control testis sections, a single PAS-positive acrosomal vesicle

was observed to adhere to step 2-3 round spermatid nuclei (stage II-III) in the Golgi phase of

acrosome development, and progressively spread to cover the apical half of the spermatid

nuclei as spermiogenesis progressed (Fig 6A). In *Katna1/al1*^{GCKO/GCKO} mice however, a large

subset of round spermatids with supernumerary ectopic acrosomal vesicles were observed (Fig

6A orange arrowheads), in addition to a smaller subset of round spermatids wherein pro-

acrosomal vesicles had failed to attach to the nucleus at the appropriate time (Fig 6A, yellow

arrowheads). As spermiogenesis progressed, elongating spermatids with fragmented and

partially detached acrosomes were common in *Katna1/al1*^{GCKO/GCKO}, but never seen in

Katna1/al1^{Flox/Flox} controls (Fig 6A, blue arrowheads). Ectopic PAS-positive vesicles were also

present throughout the seminiferous epithelium during multiple steps of acrosome biogenesis

in *Katna1/al1*^{GCKO/GCKO}, but not *Katna1/al1*^{Flox/Flox}, mice (Fig 6A, green arrowheads). TEM of

acrosome biogenesis confirmed these observations (Fig 6B), and revealed morphological

abnormalities in the spermatid Golgi apparatus of *Katna1/al1*^{GCKO/GCKO} mice (Fig 6Bi,k)

compared to *Katna1/al1*^{Flox/Flox} mice (Fig 6Bc,e). In normal round spermatids, and as seen in

Katna1/al1^{Flox/Flox} mice (Fig 6Bc,e), the Golgi apparatus has a flattened sometimes horseshoe-

like morphology wherein the trans-Golgi network (TGN) is closely associated with, and

oriented in parallel to, the apical nuclear membrane. In *Katna1/al1*^{GCKO/GCKO} mice however the

Golgi stacks were disorganised, often displaying a circular morphology, wherein the cis-Golgi

network almost entirely enclosed the TGN (Fig 6Bi), or a dispersed, splayed morphology (Fig

6Bk). These phenotypes, reveal KATNA1 and KATNAL1 collectively regulate the spatial

organisation of the Golgi apparatus and the transport of proacrosomal vesicles, and their fusion,

to the nuclear membrane. The acrosome abnormalities observed in *Katna1/al1*^{GCKO/GCKO} mice,

are reminiscent of those seen in KATNB1 loss of function mice[11] suggesting these KATNA1

and KATNAL1 likely function in complexes with KATNB1 during acrosome biogenesis.

Analysis of the key MT-based structures required for spermiogenesis revealed phenotypes consistent with a loss of MT severing and global dysregulation of the MT cytoskeleton. Similar to *Katna1*$^{GCKO/GCKO}$ mice, immunolabelling of testis sections for acetylated tubulin revealed sperm tails formed in *Katna1/al1*$^{GCKO/GCKO}$ mice, however reflective of sperm numbers fewer tails were seen than in *Katna1/al1*$^{Flox/Flox}$ controls (Fig 6C-F). Acetylated tubulin staining however, revealed the frequent ectopic accumulation of stable MTs in the cytoplasm of spermatids in *Katna1/al1*$^{GCKO/GCKO}$ mice (Fig 6E-F), consistent with the decreased turnover of MTs.

Assessment of manchette formation via the immunolabelling of testis sections for α-tubulin revealed a catastrophic dysregulation of manchette formation. MTs accumulated around the spermatid nucleus at stage VIII in *Katna1/al1*$^{GCKO/GCKO}$ mice (Fig 6G), however they failed to assemble into a sheath around, and with MTs aligning in parallel to, the distal half of the nucleus. Instead, the cytoplasm of the elongating spermatids became enriched in ectopic α-tubulin as spermiogenesis progressed (Fig 6G). Analysis of elongating spermatid ultrastructure via TEM confirmed these observations (Fig 6H). In *Katna1/al1*$^{Flox/Flox}$ mouse elongating spermatids, a typical manchette structure wherein MTs projected distally from a perinuclear ring was observed (Fig 6H). In *Katna1/al1*$^{GCKO/GCKO}$ mice, MTs were observed throughout the cytoplasm of elongating spermatids, however these were never associated with a perinuclear ring and were not arranged into typical manchette structure (Fig 6H). Collectively, these results reveal that KATNA1/KATNAL1-mediated severing is required to restrain MT growth/prune MT bulk during spermatid differentiation and to regulate manchette formation. Moreover, the absence of these phenotypes in the single GCKO KATNA1 and KATNAL1 models, suggest KATNA1 and KATNAL1 can compensate for each other in these aspects of spermiogenesis.

KATNA1, KATNAL1 and KATNB1 interact with a range of cytoskeletal and vesicular

transport proteins in the mouse testis

To explore the mechanism of action of KATNA1 and KATNAL1 during male meiosis and

spermiogenesis, and how they functionally compensate for each other, we determined the

respective interactome for KATNA1, KATNAL1 and KATNB1, within testis

immunoprecipitates via MS (Table S1 and Fig S6A-C). A total of 322, 108 and 213 proteins

were significantly enriched in the KATNA1, KATNAL1 and KATNB1 IP respectively

compared to their corresponding controls (Table S1 and Fig S6A-C). As expected, KATNA1,

KATNAL1 and KATNB1 were all significantly enriched in their own IPs validating the

specificity of each IP. KATNA1 and KATNAL1 were not found in the binding partner list for

each other, suggesting that they do not interact with each other to form heterohexamers (Table

S1). Similarly, neither bound to KATNAL2, however KATNAL1 and KATNB1 each pulled

the other down (Table S1). Consistent with the three proteins having similar functions, they

shared 48 candidate interacting proteins, in addition to 102 candidates that were shared by two

katanins (Fig S6D).

PANTHER protein class and gene ontology analysis[15] was conducted (Table S2 and

summarized in Fig S6E-J). Consistent with their regulation of MT-based processes in

spermatogenesis, KATNA1, KATNAL1 and KATNB1 each bound several cytoskeletal

proteins (Fig S6E-G, Table S2). Of particular relevance to the regulation of the Golgi and

acrosome formation all bound proteins involved in vesicle-mediated transport. In addition,

KATNA1 bound to known vesicle targeting proteins including ARL3 (Fig S6E-G, Table S2).

Functional   descriptions   of   selected   cytoskeletal,   membrane/vesicle-transport   and

spermatogenesis related proteins significantly enriched in the KATNA1, KATNAL1 and

KATNB1 IP experiments are summarised in Table S3.

Strikingly, across the three IP experiments the cytoskeletal proteins identified were

predominantly actin-related proteins, however tubulin and MT-related proteins were also

identified (Fig S6E-J, Table S2 and S3). KATNA1 also bound to the intermediate filament

protein vimentin (Table S2 and S3). Of the MT-related proteins identified as katanin testis-

interactors, multiple proteins involved in the regulation and activity of dynein MT motor

proteins were identified. Indeed, KATNA1 and KATNB1 both bound components of the

dynactin complex (Table 1), an essential cytoplasmic-dynein cofactor involved in MT-based

cargo transport. KATNA1 also bound DNAAF4, which is required for axonemal dynein

assembly and thus ciliary motility, and all three katanins bound NUDC, a dynein-dynactin

complex regulator (Table S3). Consistent with the phenotypes of our *Katnb1*[11,12], *Katnal1* and

Katna1/al1 mutant mice, the MT-related proteins identified included those with established

functions in the mitotic and/or meiotic spindle (e.g. NUDC, MAPRE1/EB1), intraflagellar

transport and ciliogenesis (ARL3) and axoneme assembly DNAAF4; Table S3). Moreover,

several have previously been implicated in spermatogenesis and male fertility (e.g. ARL3,

DNAAF4, MAPRE1/EB1) and, consistent with its mechanism of action, KATNA1 pulled

down both α- and β-tubulin isoforms (Table S3).

Of the actin-binding proteins identified across the three katanin IP experiments, multiple

regulators of actin polymerization/dynamics were identified including components of the

ARP2/3 and CapZ complexes (Table S2, S3). KATNA1, KATNAL1 and KATNB1 each bound

at least two components of the ARP2/3 complex. Data strongly suggests this complex plays

key roles in blood-testis-barrier integrity, spermiogenesis and spermiation[16-19]. Likewise, all

bound the CapZ subunit CAPZA1, and KATNA1 and KATNAL1 both bound CAPZA2. In

addition, several regulators of actin-network organization were identified to bind KATNA1

and KATNB1, and KATNA1 bound MYL6, an actin motor protein which localizes to the

manchette (Table S3). Collectively, these data indicate a novel connection between katanin

proteins and the actin cytoskeleton in male germ cells. This is consistent with data showing the

Arabidopsis katanin A-subunit KATANIN1 regulates the actin cytoskeleton[20] and that cortical

localisation of *Drosophila* KATNA1 orthologue kat-60 is actin-dependent[21].

KATNA1 and KATNB1 bound multiple Rab GTPases, a family which regulates multiple

aspects of vesicle transport (vesicle budding, actin/MT-mediated transport and fusion) (Tables

1, S1 and S2). Of note in spermatogenesis there is increasing evidence Rab and Rab-like

proteins mediate pro-acrosomal vesicle transport, in addition to intraflagellar transport and

sperm tail development[22-25]. Other transport proteins included those related to endoplasmic

reticulum (ER) to Cis-Golgi transport, intra-Golgi transport, clathrin-mediated vesicle

disassembly and endocytosis (Table 1). Further consistent with the acrosome defects in

Katna1/al1$^{GCKO/GCKO}$ mice, KATNAL1 bound ACRBP a protein essential for acrosome granule

formation and maintenance of acrosome integrity, and consistent the Golgi defects, a number

of proteins were identified with roles in Golgi organization (e.g. Rab2, Rab6, VCP, LMAN1;

Table S3).

Other notable proteins identified as binding the katanins included essential male meiosis

regulators TEX15 (KATNA1 IP) and HSPA2 (KATNA1/B1 IP), and FABP9 a protein required

for normal sperm head shape in mice (KATNA1/AL1 IP) (Table S3).

Discussion

Katanin MT severing enzymes are potent regulators of MT dynamics and organization across

eukaryotes. Herein, using a combination of single and double katanin A-subunit GCKO mice,

we show for the first time the essential requirement for katanin A-subunits in mammalian male

meiosis, provide the first direct evidence that different katanin A-subunits can functionally

compensate for one another and establish KATNA1 and KATNAL1 as novel regulators of

spermatid differentiation (spermiogenesis). Specifically, KATNA1 and KATNAL1

collectively regulate spermatocyte metaphase and anaphase spindle structure and function,

acrosome development via the Golgi apparatus and pro-acrosomal vesicle trafficking, sperm

head shaping via manchette formation, and collectively restrain MT bulk during male meiosis

and spermiogenesis. In addition to this, KATNAL1 regulates spermatocyte cytokinesis, sperm

tail and HTCA development, manchette dynamics and dissolution, and the release of sperm

from the seminiferous epithelium (spermiation). Through the analysis of KATNA1,

KATNAL1, and KATNB1 testis binding proteins, we reveal these activities involve

interactions with a diverse suite of proteins, including actin-, MT-, vesicle/membrane

trafficking-, and spermatogenesis-related proteins.

The finding that KATNA1 and KATNAL1, but not KATNA1 and KATNAL2, have gene

redundancy in mammalian spermatogenesis is consistent with the highly similar domain

structure and severing activity shared between KATNA1 and KATNAL1, but not KATNAL2

[2]. Interestingly, KATNAL1 fully compensates for KATNA1 germ cell loss, whereas KATNA1

can only partially compensate for germ cell KATNAL1 loss. This suggests KATNAL1 is the

more potent/efficient MT severing enzyme in the testis, and/or that it has neofunctionalized to

acquire additional functions. We predict it is a combination of the two. Indeed, KATNAL1 has

possesses higher MT severing activity in mouse Neuro-2a cells, and is less prone to degradation

in HEK293T cells compared to KATNA1[26]. However, we also identified several different

KATNA1 versus KATNAL1 testis interacting proteins indicating they have at least some

distinct modes of regulation and/or function. These findings have important implications for

the study of katanin proteins in contexts wherein multiple A-subunits exist, revealing that

discovery of full katanin function can require double KO models.

We identify KATNA1/KATNAL1 as essential regulators of metaphase and anaphase spindle

structure, metaphase chromosome alignment, anaphase chromosome segregation and,

KATNAL1 as a cytokinesis regulator during male meiosis. While a role for katanin A-subunits

in male mammalian meiosis has never previously been shown, we have previously shown the

B-subunit, KATNB1 is essential for normal spindle architecture and function, and for

cytokinesis in mouse spermatocytes[11,12]. In strong support of these overlapping functions being

due to KATNB1 working in partnership with KATNA1 and KATNAL1, KATNB1-KATNA1

and KATNB1-KATNAL1 complexes are present throughout mouse male meiosis wherein they

localize to spindle MTs[11]. That single GCKO of KATNAL1, but not KATNA1, is sufficient to

cause meiotic defects, agrees with our previous data showing KATNB1-KATNAL1 complexes

are the most abundant type of katanin complex during male meiosis[11].

The misalignment of metaphase chromosomes, followed by uneven anaphase segregation in

Katnal1$^{GCKO/GCKO}$ and *Katna1/al1*$^{GCKO/GCKO}$ meiosis is consistent with KATNA1/KATNAL1

regulating MT-chromosome attachment and/or regulating spindle dynamics via metaphase

poleward 'flux' (reviewed in[27]) and anaphase 'Pacman-flux' (reviewed in[28]). While future

studies will need to directly test this in meiosis, katanin-mediated MT severing has been shown

to directly contribute to metaphase spindle flux in HEK293 mitotic cells[29] and Pacman-flux in

Drosophila mitotic cells[30].

The role for KATNAL1 in cytokinesis identified herein aligns with evidence that midbody

disassembly in mitosis requires MT severing ([31] and reviewed in [2]) and our own data, regarding

KATNB1 function in mouse male meiosis[11,12]. That only a subset of germ cells fails cytokinesis

in the absence of KATNAL1 or KATNB1, suggests that other MT severing enzymes must also

be involved in this process. Indeed, we have recently shown the related MT severing enzyme,

spastin, also contributes to cytokinesis during male meiosis[32].

Our data establishes KATNA1 and KATNAL1 as novel regulators of Golgi organisation and

pro-acrosomal vesicle trafficking and fusion during acrosome biogenesis. This contributes to

an emerging body of evidence that MT-severing enzymes regulate vesicle-based transport in

spermiogenesis, as we have recently shown spastin and KATNB1 are also essential for pro-

acrosomal vesicle transport and fusion events[11,32]. The contribution of KATNA1/KATNAL1 to

these processes may be as simples as a role in pruning the MT tracks along which Golgi

membranes are spatially positioned and pro-acrosomal vesicles are delivered. However, the

interaction of KATNA1, KATNAL1 and KATNB1 with multiple proteins involved in Golgi

organisation and vesicle trafficking suggests a more direct role and raises the possibility MT-

severing can modulate cargo-MT interactions.

We have previously shown KATNAL2 is essential for sperm tail development via basal body

-plasma membrane docking and the initiation of spermatid axoneme extension[5]. Here, we show

KATNAL1 is also required, but in the subsequent assembly, and potential stability, of axoneme

MT doublets, and in mitochondrial sheath and ODF development. Moreover, while KATNAL1

is not required for basal body-membrane docking, it is required for HTCA formation/integrity.

The precise mechanism of KATNAL1 contribution to these processes remains an outstanding

question. One possibility is that KATNAL1-mediated MT severing directly amplifies and

regulates MTs for axoneme doublet formation and HTCA fortification. However, the

variability in phenotypes (e.g. missing MT doublets in some tails and supernumerary MT

doublets in others), is more suggestive of mis-regulated cargo trafficking. Of note, and

consistent with the co-incident manchette defects in KATNAL1 loss, sperm tail components

including the ODFs, and proteins required to strengthen the HTCA appear to be delivered via

the MTs of the manchette, and abnormal sperm tail and manchette phenotypes often occur in

parallel (reviewed in[33]). It is thus possible that the abnormal manchette phenotypes in

Katnal1$^{GCKO/GCKO}$ mice underpin at least some of the sperm tail defects.

The manchette phenotypes identified in the *Katnal1*$^{GCKO/GCKO}$ mice establish roles for

KATNAL1 in manchette migration down the sperm nucleus, which requires 'unzipping' of

rod-like manchette-nuclear linkers, and in controlling manchette MT length and in manchette

MT disassembly. These functions are consistent with a direct MT-severing mechanism.

Strikingly, the *Katnal1*$^{GCKO/GCKO}$ mouse manchette phenotypes, mirror not only those in

KATNB1 loss of function mice[11,12], but also those in KATNAL2 loss[5]. These identical yet non-

redundant KATNAL1 and KATNAL2 phenotypes suggest the two proteins are executing

similar functions in the manchette but severing different MT subpopulations. This is consistent

with our proteomics data, demonstrating KATNAL2 is not a binding partner of KATNAL1.

Similar to the established roles of Sertoli cell KATNAL1 is required to maintain round

spermatid-Sertoli cell adhesion[13]. We show germ cell KATNAL1 is involved in spermatid

retention and elongated spermatid release via spermiation. Data is consistent with KATNAL1

severing the cytoskeletal connections tethering elongating spermatids to the Sertoli cell. The

precise cytoskeletal targets of this action need to be identified, but of note, our IP revealed

katanins interact with multiple ARP2/3 complex proteins. The ARP2/3 complex appears to

facilitate spermiation, *albeit* via the disassembly of the Sertoli cell actin cytoskeleton [19]. Of

note spermiation defects in KATNAL2 mutant mice [5], and spermatid adhesion and spermiation

defects seen in the KATNB1 mutant mice[11,12] phenocopy those seen in *Katnal1*[GCKO/GCKO] mice.

This suggests that KATNAL1-KATNB1 complexes along with KATNAL2-KATNB1

complexes facilitate spermatid release.

Finally, we defined the testis interactome for KATNA1, KATNAL1 and KATNB1. While a

katanin interactome has previously been defined in HELA cells[4], this is the first *in vivo* tissue

wide interactome for each protein. A number of protein classes were identified that we predict

are related to the activation/inhibition of katanin function (e.g. metabolite interconversion

enzymes). Of proteins that we predict are involved specific testis functions of the katanin

enzymes, cytoskeletal and vesicle-transport related proteins emerged as key protein classes.

Future studies will need to define precise nature of these interactions, however they paint a

picture of katanin action involving a complex crosstalk between different cytoskeletal networks

notably between actin and MT networks, and suggest a direct role for katanins in regulating

the interaction of dynein- and vesicle-based cargo machinery with MTs.

Collectively, this study establishes KATNA1 and KATNAL1 as novel players in male germ

cell MT regulation. The data defined herein, taken together with our previous data on the roles

of KATNB1[11,12], KATNAL2[5] and Sertoli cell KATNAL1[13], allows us to define a model of the

complete katanin 'toolbox' during mammalian spermatogenesis (Fig 7).   This data

unequivocally shows the emergence of multiple katanin A-subunits in higher eukaryotes, not

only allows for functional sub-specialization and thus customized cutting of discrete MT

subpopulations, but also for the protection of katanin function in critical processes such as

meiosis through gene redundancy.

Methods

Mouse model production

All animal procedures were approved by the Monash University School of Biological Sciences

Animal Experimentation Ethics Committee and the University of Melbourne Ethics committee

and conducted in accordance with National Health and Medical Research Council (NHMRC)

Guidelines on Ethics in Animal Experimentation.

To the generate *Katna1^(Flox/Flox)* mice, *Katna1* KO-first conditional-ready mice (*Katna1^(KO/WT)*)

were generated through the Monash University Node of the Australian Phenomics Network

using a EUCOMM *Katna1* KO-first conditional-ready ES clone (EPD0586_1_B10) as

described[34]. To disrupt the *Katna1* gene, the FRT-LacZ-loxP-Neo-FRT-loxP-*Katna1-exons6-*

7-loxP cassette was inserted into intron 5 of the *Katna1* gene (Fig S1C).   To generate

conditional *Katna1* mouse (*Katna1^(Flox/Flox)*), mice heterozygous for the knockout-first allele

(*Katna1^(KO/WT)*) were then crossed with a transgenic line carrying the *Flp* recombinase gene.

This resulted in removal of the *FRT* flanked gene-trap cassette, exposing a floxed *Katnal1* exon

6-7 and reverting mice to a wild-type phenotype. *Katna1$^{GCKO/GCKO}$* were then generated using

a two-step breeding strategy. (1) *Katna1$^{Flox/Flox}$* females were mated with a *Stra-8-cre*

transgenic line to produce *Katna1$^{WT/Flox,Stra8-cre+}$* males. (2) *Katna1$^{WT/Flox,Stra8-cre+}$* males were

mated with *Katna1$^{Flox/Flox}$* females to produce *Katna1$^{Flox/Flox,Stra8-cre+}$* progeny

(*Katna1$^{GCKO/GCKO}$*). In these mice, gene disruption relied on Cre-lox recombination mediated

excision of *Katna1* exons 6 and 7 (ENSMUSE00001254109, ENSMUSE00001247569), and

the *Stra8-Cre* used is expressed from late spermatogonia onwards [35].

Katnal1$^{Flox/Flox}$ mice, wherein exon 3 (ENSMUSE00001360953) of the *Katnal1* gene is flanked

by loxP sites, were generated by the Monash Genome Modification Platform using

CRISPR/Cas9 technology. Guide RNA target sites flanking the target exon (5' end of exon -

5'   TCAATGTCTGAGGTGGATAA   3'   and   3'   end   of   exon   -   5'

CAAAGAAGTCTTTTAGACCC 3') were identified using the UCSC Genome Browser. A

single stranded DNA homology directed repair template, was generated from a *Katnal1*

conditional KO targeting construct using the Guide-it Long ssDNA Strandase Kit (Takara,

632645). The following oligonucleotides were used to generate a PCR product from which the

ssDNA   was   generated: 5'/5Biosg/GATGCTTAGCACTGGCTGG   3'   and   5'

GAACGTGTGCTATTTGGTCTTTAAG 3'. The CRISPR/Cas9 ribonucleoprotein complex

and single stranded DNA homology repair templates were microinjected into C57BL/6J

zygotes at the pronuclei stage and transferred into the uterus of pseudo-pregnant females.

Katnal1$^{GCKO/GCKO}$ mice were generated by intercrossing *Katnal1$^{Flox/Flox}$* mice with a *Stra8-cre*

transgenic line as above. *Katnal2$^{Flox/Flox}$* mice were generated as previously described[5].

Katna1/al1$^{Flox/Flox}$ and *Katna1/al2$^{Flox/Flox}$* mice were generated by breeding *Katna1$^{Flox/Flox}$* mice

with either *Katnal1$^{Flox/Flox}$* or *Katnal2$^{Flox/Flox}$* mice, respectively. Double germ cell deletion was

then achieved using by intercrossing double floxed mice with a *Stra8-cre* transgenic line as

above.

All mice were maintained on a C57BL/6 background. To control for background strain effects

on phenotype, *Katna1$^{Flox/Flox}$*, *Katnal1$^{Flox/Flox}$*, *Katna1/al1$^{Flox/Flox}$* and *Katna1/al2$^{Flox/Flox}$* mice

mice were used as controls for *Katna1$^{GCKO/GCKO}$*, *Katnal1$^{GCKO/GCKO}$*, *Katna1/al1$^{GCKO/GCKO}$* and

Katna1/al2$^{GCKO/GCKO}$ mice, respectively. Mouse genotypes were determined from tail biopsies

using real time PCR with specific probes designed for each allele (Transnetyx, Cordova, TN).

Efficiency of exon excision in *Katna1$^{GCKO/GCKO}$*, *Katnal1$^{GCKO/GCKO}$*, *Katna1/al1$^{GCKO/GCKO}$* and

Katna1/al2$^{GCKO/GCKO}$ mice, was determined by qPCR of isolated germ cells as described below.

Protein levels were determined by western blotting of germ cell and testis lysates as described

below.

Quantitative qPCR

Purified germ cells were prepared via the STAPUT method as detailed in [36]. Total RNA was

then extracted from purified germ cells using Trizol (Life Technologies), and reverse-

transcribed into cDNA using SuperScript III Reverse Transcriptase (Life Technologies).

Primers spanning *Katna1* exons 6 and 7 (5'-TCAGGAAGCAGTGGTGTTACC-3' forward

and 5'-AAGGGTCTTTCCAGTGCCAG-3' reverse), *Katnal1* exons 3 and 4 5'-

CCGTGTCTTGCCGAGATGAA-3'forward and 5'-GCGATTTGGACGCCTGATTT-3'

reverse) and *Katnal2* exons 2 and 3 (5'-AAGGCTTTGGAGGAGGAGAC-3' forward and 5'-

GGGGCCTTCTTAACCACTTT-3') were used to verify gene deletion in the corresponding

mouse models. All qPCRs were performed using SYBR Select Master Mix (Applied

Biosystems). Each reaction was performed in triplicate using the Agilent Mx3000P qPCR

system or an Applied Biosystems QuantStudio 3 Real-Time PCR System with the parameters

of 50°C for 2 minutes, 95°C for 10 minutes followed by 40 cycles of 95°C for 15 seconds and

60°C for 1 minute. *Ppia* was amplified at the same time as an internal control (using 5'-

CGTCTCCTTCGAGCTGTTT-'3 forward and 5'-CCCTGGCACATGAATCCT-'3 reverse

primers), and all results were normalized to *Ppia* expression. Differential expression was

analysed using the $2\Delta\Delta^{CT}$ method[37].

Male fertility characterisation

Male fertility of all mouse models was characterised as per[38]. Male fertility was assessed by

mating 10-12 week old males with wild-type females (≥6 weeks of age). Copulatory plugs

were monitored to confirm successful mating and the number of pups born per copulatory plug

was recorded. Each male was test mated with 2-3 females and pup numbers averaged. Testis

daily sperm production (DSP) and total epidydimal sperm content were determined using the

Triton X-100 solubilisation method[39] (n≥3/genotype) as modified in[11]. To assess sperm

motility, sperm were backflushed as per[40] from cauda epididymides and motility was assessed

using a computer assisted sperm analyser, as described previously[41].

For histology, testis and epididymal tissue were fixed in Bouin's fixative and histology

assessed using standard methods. Male reproductive tract histology was visualised using

periodic acid-Schiff's (PAS) and haematoxylin staining, and air-dried caudal epididymal sperm

smears were visualised using haematoxylin and eosin staining (n≥3/genotype). Germ cell

apoptosis was evaluated by immunostaining for cleaved Caspase 3 and 7 as per[42] and as

described below. The number of Caspase-positive cells in a minimum of 50 randomly selected

seminiferous tubules per mouse were counted and statistical analysis conducted as described

below and as per[11](n≥3 mice /genotype).

TEM

For analysis of seminiferous tubule ultrastructure, partially decapsulated testes and cauda epididymides were fixed with 5% glutaraldehyde/4% paraformaldehyde/0.02% picric acid in 0.1 M sodium cacodylate buffer, pH 7.4. as per[11]. Samples were *en bloc* stained with 2% uranyl acetate and all samples were embedded into Epon resin using standard procedures. Ultrathin sections were cut on a Leica EM UC7 Ultramicrotome and placed on Copper 100×100 square grids (ProSciTech). Sections were analysed using a FEI Talos L120CI TEM in the Ian Holmes Imaging Centre (The University of Melbourne, Parkville)

Antibody use

Primary antibodies used included those against acetylated-tubulin (T6793, Sigma, RRID: Ab_477585, ascites fluid, 1 in 10,000), α-tubulin (T5168, Sigma, ascites fluid, 1 in 50,000), cleaved-Caspase 3 (9664, Cell Signalling, 0.5 μg ml^{-1}), cleaved-Caspase 7 (9491, Cell Signalling, 0.23 μg ml^{-1}), KATNA1 (as described previously [11]), KATNAL1 (as described previously[11]), KATNB1 (as described in [11]), SUN5 (ProteinTech 17495-1-AP 0.8 μg/ml) and TUBB3 (Sigma-Aldrich, SAB3300047, 0.15 μg ml^{-1}). Secondary antibodies included Alexa Fluor 488 donkey anti-mouse (A212062 Invitrogen), Alexa Fluor 488 donkey anti-rabbit (A21206, Invitrogen), and Alex Fluor 555 donkey anti-rabbit (A31572, Invitrogen). EnVision+ Dual Link System-HRP (K4063, Dako) was used to amplify signal. The specificity of immunolabelling was determined by staining sections in parallel in the absence of primary antibody or where available KO mouse tissue.

Western blotting

Proteins were extracted from STAPUT purified germ cells [36], and immunoblotting conducted as previously described[36]. Images were captured using a ChemiDoc MP Imaging System

(BioRad). The molecular weights of detected proteins were estimated using a PageRuler Plus

Prestained Protein Ladder (ThermoFisher) and the Image Lab 6.0 Molecular Weight Analysis

Tool (BioRad).

Immunochemistry and imaging

To assess KATNA1 testis localisation, 5 μm cryostat testis sections were fixed in 1:1 methanol-

acetone and permeabilised in 0.2% Triton X-100 in PBS as per[43]. To assess testis localisation

of all other proteins, Bouin's fixed testes were processed into paraffin and 5 μm sections

prepared using standard methods. Paraffin sections were dewaxed and heat-mediated antigen

retrieval was conducted by microwaving sections 10 mM citrate buffer (pH 6) for 16 minutes

as per [44]. For sections to be used for chromogenic immunohistochemistry (α-tubulin,

acetylated-tubulin, cleaved-caspase 3 and 7), endogenous peroxidase activity was then blocked

via incubation with 5% H_2O_2 for 5 minutes. For all testis immunolabelling, non-specific protein

binding was minimised by incubation with CAS-block (Invitrogen) for 30 minutes and sections

were then incubated with primary antibodies diluted in DAKO Antibody Diluent overnight at

4°C. For immunofluorescent labelling, sections were incubated with the appropriate Alexa

Fluor secondary antibodies diluted 1in 500 for 1 hour at room temperature. DNA was then

visualised using 1 μg ml^{-1} 4',6-diamidino-2-phenylindole (DAPI, Invitrogen) or 4 μM TO-

PRO-3 Iodide (TOPRO, Thermo Scientific). Acrosomes were visualized using 0.5 μg ml^{-1}

lectin peanut agglutinin (PNA), Alexa Fluor 488 conjugate (L21409, Life Technologies). For

chromogenic immunohistochemistry, sections were incubated with DAKO EnVision+ Dual

Link System-HRP for 30 minutes at room temperature. The Liquid DAB+ Substrate

Chromogen System (DAKO, K3468) was then applied as per the manufacturer's instructions.

Nuclei were counterstained using Mayer's haematoxylin (Amber Scientific), followed by

Scott's tap water (Amber Scientific).

To assess SUN5 localisation in cauda epididymal sperm, sperm were washed in PBS and

airdried onto slides. Sperm were then fixed in 4% paraformaldehyde for 10 minutes. Sperm

were then permeabilised, blocked for non-specific protein binding and immunofluorescently

labelled as above.

Immunofluorescent images were collected using a Leica TCS SP8 confocal microscope (Leica

Microsystems) at the Monash Micro Imaging (Monash University) and the University of

Melbourne Biological Optical Microscopy Unit (University of Melbourne, Parkville). A

63x/1.40 HC PL APO CS2 oil immersion objective was used to take all immunofluorescent

images. Bright field images were collected cellSens Imaging Software (Olympus) on an

Olympus BX53 light microscope equipped with an Olympus DP80 camera.

IP-MS

For IP assays, wild-type mouse testes (n=3 per IP assay) were lysed in 1% NP-40 (PBS)

supplemented with protease inhibitor cocktail (Calbiochem, 539134). KATNA1, KATNAL1

and KATNB1 and their binding partners were then co-immunoprecipitated using the Pierce

Co-immunoprecipitation Kit (Thermo Scientific, 26149) as per the manufacturer's instructions,

in combination with either our KATNA1 (10μg per IP column), KATNAL1 (5μg) or KATNB1

(5μg) inhouse antibody. Briefly, to reduce non-specific binding, testis protein lysates were first

pre-cleared by incubation with 400μl of Pierce control agarose resin slurry for a 0.5-1 hour at

4°C with gently rocking. For each pre-cleared testis lysate, 5 mg was then loaded into two IP

columns, one which contained the target antibody covalently coupled to the amine reactive

Pierce AminoLink Plus Coupling Resin and one which contained the target antibody and a

non-amine reactive Pierce Control Agarose Resin (negative control), and incubated overnight

at 4°C. Lysate flow through was removed, columns were washed and bound protein eluted and

collected as per the manufacturer's instructions. The eluted proteins were neutralised with

1.5M Tris-HCL (pH 8.8), reduced with TCEP, then digested by incubation with trypsin

overnight at 37°C. Peptide samples were freeze dried and resuspended in 2% acetonitrile

(ACN)/0.1% formic acid (FA) (diluted in MilliQ water) before concentrated FA was added to

adjust the pH to <3. Samples were purified using reverse-phase chromatography C18 stage

tips, which aim to desalt and fractionate in-solution peptides at an acidic pH. The tips were

initially activated with 100% methanol and equilibrated with 0.1% FA, before samples were

loaded into them and desalted with 0.1% FA. Peptides were eluted with 80% ACN/0.1% FA,

freeze dried, resuspended in 2% ACN/0.1% FA and sonicated for 15 minutes in a water bath.

The purified peptide samples were analysed via nano liquid chromatography coupled to tandem

mass spectrometry (LC-MS/MS). Briefly, the KATNA1 IP and KATNB1 IP samples were

analysed at the Monash Proteomics and Metabolomics Facility (Monash University) using an

Orbitrap Fusion Tribrid mass spectrometer (ThermoFisher, USA) and a Q-Exactive Hybrid

Quadrupole - Orbitrap mass spectrometer (ThermoFisher, USA), respectively. Both

instruments were coupled to an UltiMate 3000 RSLCnano system (ThermoFisher, USA)

equipped with an Acclaim PepMap RSLC C18 nanoViper Analytical Column (100Å, 75 μm

X 50 cm, 2 μm; ThermoFisher, USA) and an Acclaim PepMap 100 C18 HPLC nanoViper Trap

Column (100Å, 100 μm X 2 cm, 5 μm; ThermoFisher, USA). The tryptic peptides were

separated by increasing concentrations of 80% acetonitrile (ACN) / 0.1% formic acid at a flow

rate of 250 nl/min for 128 min and 158 min, respectively. Both mass spectrometers were

operated in data-dependent acquisition (DDA) mode to automatically switch between full scan

MS and MS/MS acquisition. Each survey full scan (m/z 375–1800) in the Orbitrap Fusion

Tribrid acquisitions was acquired in the Orbitrap with 120,000 resolution (at m/z 200) after

accumulating ions to a Normalised AGC Target of 250% with a maximum injection time of 54

ms. Dynamic exclusion was set to 15 seconds. Adhering to a 2 sec cycle time, the most intense

multiply charged ions (z ≥ 2) were sequentially isolated and fragmented in the collision cell by

higher-energy collisional dissociation (HCD) and ms2 scans were acquired with a fixed

injection time of 54 ms, 30,000 resolution and a Normalised AGC Target of 400%. Very similar

parameters were used for the Q-Exactive Hybrid Quadrupole – Orbitrap acquisitions.

The KATNAL1 IP purified peptide samples were analysed at the University of Melbourne

Mass Spectrometry and Proteomics Facility, using an Orbitrap Eclipse Tribrid Mass

Spectrometer (ThermoFisher, USA) equipped with a nano ESI interface coupled to an Ultimate

3000 nano HPLC (Thermo Fisher, USA). The LC system was equipped with an Acclaim

Pepmap 100 C18 HPLC nanoViper Trap Column (100 Å, 75 μm X 2 cm, 3 μm; ThermoFisher,

USA) and an analytical column as above. The enrichment column was injected with the tryptic

peptides (1 μL, 100 ng) at an isocratic flow of 5 μL/min of 2% *v/v* CH_3CN containing 0.05%

v/v trifluoroacetic acid for 6 min, applied before the enrichment column was switched in-line

with the analytical column. The eluents were 0.1% v/v formic acid (solvent A) in H_2O and

100% *v/v* CH_3CN in 0.1% *v/v* formic acid (solvent B). The gradient was at 300 nl min^{-1} from

(i) 0–6 min, 3 % B; (ii) 6–35 min, 3–23 % B; (iii) 35-45 min, 23-40 % B; (iv) 45-50 min, 40-

80 % B; (v) 50-55 min, 80–80 % B; (vi) 55-55.1 min, 80-3 % B; (vii) 55.1-65 min, 3-3 % B.

The Eclipse Orbitrap mass spectrometer was operated in the data-dependent mode, wherein

full MS^1 spectra were acquired in a positive mode over the range of *m/z* 375-1500, with spray

voltage at 1.9kV, source temperature at 275 °C, MS^1 at 120,000 resolution, normalized AGC

target of 100 % and maximum IT time of 22 ms. The top 3 second method was used and

selecting peptide ions with charge states of ≥ 2-7 and intensity thresholds of ≥ 5E4 were

isolated for MS/MS. The isolation window was set at 1.6 *m/z*, and precursors were fragmented

using higher energy C-trap dissociation (HCD) at a normalised collision energy of 30, a

resolution of 15,000, a normalized AGC target of 100% and automated IT time.

The raw MS data were analysed with the MaxQuant software suite for the identification and

quantification of peptides/proteins from the Mouse SwissProt database. Fixed modifications of

carbamidomethylation of cysteine as well as variable oxidation of methionine and protein N-

terminal acetylation were considered. Trypsin/P was set as the protease with a maximum of 2

missed cleavages. Statistical analysis was conducted with the Perseus software considering

protein and PSM false discovery rates (FDR) both set at < 0.01. For each IP assay, the majority

UniProt ID(s) for each significantly enriched protein group were then analysed using

PANTHER Gene Ontology and protein classification[15].

Statistics

Statistical analysis of germ cell apoptosis data was conducted in R version 3.5.1[45]. Generalised

linear mixed (GLM) models were used to compare the number of caspase-positive cells per

tubule across genotypes, with animal ID included as a random effect to account for repeated

measures per individual. For each model, Akaike information criteria (AIC) estimates were

used to select the most appropriate error distribution and link function (i.e. poisson, negative

binomial, zero-inflated poisson, zero-inflated negative binomial) using the glmer function

(lme4 package[46]) and glmmTMB function (glmmTMB, package[47]). For all models, a zero-

inflated negative binomial distribution (fitted with glmmTMB, using the ziformula argument)

was selected as the most appropriate error distribution and link function (i.e. had the lowest

AIC score).

All other statistical analysis, other than for the proteomics (described above), was conducted

in Graphpad Prism 9.0. The statistical significance of differences between two groups was

determined with an unpaired Student t-test. Significance was defined as a P value of <0.05.

Acknowledgements

We thank Dr Liza O'Donnell and Dr Duangporn Jamsai for advice on the initial stages of the project. We thank Ms G. Gemma Stathatos, Ms Lauren Rogers and Ms Georgia Shering for technical assistance. We thank Associate Professor Nicholas Williamson, Dr Ching-Seng Ang and Dr Michael Leeming for technical assistance and advice regarding the MS analysis. We thank the following platforms for the use of their facilities and services: at The University of Melbourne (Parkville), the Biological Optical Imaging Platform, the Ian Holmes Imaging Centre, the Mass Spectrometry and Proteomics Facility, the Melbourne Bioresources Platform and the Melbourne Histology Platform; and at Monash University (Clayton), the Monash Animal Research Platform, the Monash Gene Modification Platform, the Monash Histology Platform, Monash Micro Imaging and the Monash Proteomic and Metabolomic Facility.

Author Contributions

Conceptualisation, J.E.M.D, M.K.O; Methodology, J.E.M.D, A.E.O, D.J.M, R.B.S, B.J.H and M.K.O; Formal Analysis, J.E.M.D, M.G, K.W, A.E.O, D.J.M, J.N, R.B.S, B.J.H, M.K.O; Investigation, J.E.M.D, M.G, K.W, A.E.O, D.J.M, J.N, R.B.S, B.J.H; Writing – Original Draft, J.E.M.D, M.G, M.K.O; Writing – Review & Editing, J.E.M.D, R.B.S, M.K.O; Visualisation, J.E.M.D, M.G, K.W; Supervision, J.E.M.D, M.K.O; Project Administration, J.E.M.D, M.K.O; Funding Acquisition, M.K.O

Declaration of interests

The authors declare no competing or financial interests.

Funding

878 J.E.M.D is supported by a National Health and Medical Research Council Ideas Grant to

879 M.K.O and J.E.M.D (APP1180929) and this project was supported in part by funding from the

880 National Health and Medical Research Council of Australia (APP1138014) and the Australian

881 Research Council (DP160100647). B.J.H is supported by funding from the Male Contraceptive

882 Initiative Fellowship.

References

1.   Mains, P.E., Kemphues, K., Sprunger, S.A., Sulston, I., and Wood, W. (1990).

Mutations affecting the meiotic and mitotic divisions of the early Caenorhabditis

elegans embryo. Genetics *126*, 593-605.

2.   McNally, F.J., and Roll-Mecak, A. (2018). Microtubule-severing enzymes: From

cellular functions to molecular mechanism. Journal of Cell Biology *217*, 4057-4069.

3.   Sarbanes, S.L., Zehr, E.A., and Roll-Mecak, A. (2022). Microtubule-severing enzymes.

Current Biology *32*, R992-R997.

4.   Cheung, K., Senese, S., Kuang, J., Bui, N., Ongpipattanakul, C., Gholkar, A., Cohn,

W., Capri, J., Whitelegge, J.P., and Torres, J.Z. (2016). Proteomic analysis of the

mammalian katanin family of microtubule-severing enzymes defines katanin p80

subunit B-like 1 (KATNBL1) as a regulator of mammalian katanin microtubule-

severing. Molecular & Cellular Proteomics *15*, 1658-1669.

5.   Dunleavy, J.E., Okuda, H., O'Connor, A.E., Merriner, D.J., O'Donnell, L., Jamsai, D.,

Bergmann, M., and O'Bryan, M.K. (2017). Katanin-like 2 (KATNAL2) functions in

multiple aspects of haploid male germ cell development in the mouse. PLOS Genetics

13, e1007078.

6.   Sharp, D.J., and Ross, J.L. (2012). Microtubule-severing enzymes at the cutting edge.

Journal of Cell Science *125*, 2561-2569. 10.1242/jcs.101139.

7.   Gomes, J.-E., Tavernier, N., Richaudeau, B., Formstecher, E., Boulin, T., Mains, P.E.,
Dumont, J., and Pintard, L. (2013). Microtubule severing by the katanin complex is
activated by PPFR-1–dependent MEI-1 dephosphorylation. Journal of Cell Biology
202, 431-439. 10.1083/jcb.201304174.

8.   Loughlin, R., Wilbur, J.D., McNally, F.J., Nédélec, F.J., and Heald, R. (2011). Katanin
contributes to interspecies spindle length scaling in Xenopus. Cell *147*, 1397-1407.

9.   Gao, L.L., Xu, F., Jin, Z., Ying, X.Y., and Liu, J.W. (2019). Microtubule-severing
protein Katanin p60 ATPase-containing subunit A-like 1 is involved in pole-based
spindle organization during mouse oocyte meiosis. Molecular Medicine Reports *20*,
3573-3582.

10. Lynn, N.A., Martinez, E., Nguyen, H., and Torres, J.Z. (2021). The mammalian family
of katanin microtubule-severing enzymes. Frontiers in Cell and Developmental
Biology *9*.

11. Dunleavy, J.E.M., O'Connor, A.E., Okuda, H., Merriner, D.J., and O'Bryan, M.K.
(2021). KATNB1 is a master regulator of multiple katanin enzymes in male meiosis
and haploid germ cell development. Development *148*, dev199922.

12. O'Donnell, L., Rhodes, D., Smith, S.J., Merriner, D.J., Clark, B.J., Borg, C., Whittle,
B., O'Connor, A.E., Smith, L.B., and McNally, F.J. (2012). An essential role for katanin
p80 and microtubule severing in male gamete production. PLOS Genetics *8*, e1002698.

13. Smith, L.B., Milne, L., Nelson, N., Eddie, S., Brown, P., Atanassova, N., O'Bryan,
M.K., O'Donnell, L., Rhodes, D., and Wells, S. (2012). KATNAL1 regulation of Sertoli
cell microtubule dynamics is essential for spermiogenesis and male fertility. PLOS
Genetics *8*, e1002697.

14.   Pleuger, C., Lehti, M.S., Dunleavy, J.E., Fietz, D., and O'Bryan, M.K. (2020). Haploid

927   male germ cells—the Grand Central Station of protein transport. Human Reproduction

928   Update *26*, 474-500. 10.1093/humupd/dmaa004.

929   15.   Thomas, P.D., Ebert, D., Muruganujan, A., Mushayahama, T., Albou, L.-P., and Mi,

930   H. (2022). PANTHER: Making genome-scale phylogenetics accessible to all. Protein

931   Science *31*, 8-22. https://doi.org/10.1002/pro.4218.

932   16.   Liu, L., Zhang, Y., Chang, X., Li, R., Wu, C., Tang, L., and Zhou, Z. (2018).

933   Fluorochloridone perturbs blood-testis barrier/Sertoli cell barrier function through

934   Arp3-mediated   F-actin   disruption.   Toxicology   Letters   *295*,   277-287.

935   https://doi.org/10.1016/j.toxlet.2018.07.001.

936   17.   Lie, P.P., Chan, A.Y., Mruk, D.D., Lee, W.M., and Cheng, C.Y. (2010). Restricted

937   Arp3 expression in the testis prevents blood–testis barrier disruption during junction

938   restructuring at spermatogenesis. Proceedings of the National Academy of Sciences

939   *107*, 11411-11416.

940   18.   Li, N., Mruk, D.D., Tang, E.I., Wong, C.K.C., Lee, W.M., Silvestrini, B., and Cheng,

941   C.Y. (2015). Formins: Actin nucleators that regulate cytoskeletal dynamics during

942   spermatogenesis. Spermatogenesis *5*, e1066476. 10.1080/21565562.2015.1066476.

943   19.   O'Donnell, L., Nicholls, P.K., O'Bryan, M.K., McLachlan, R.I., and Stanton, P.G.

944   (2011). Spermiation: the process of sperm release. Spermatogenesis *1*, 14-35.

945   20.   Takáč, T., Šamajová, O., Pechan, T., Luptovčiak, I., and Šamaj, J. (2017). Feedback

946   microtubule control and microtubule-actin cross-talk in Arabidopsis revealed by

947   integrative proteomic and cell biology analysis of KATANIN 1 mutants. Molecular &

948   Cellular Proteomics *16*, 1591-1609.

949   21.   Zhang, D., Grode, K.D., Stewman, S.F., Diaz-Valencia, J.D., Liebling, E., Rath, U.,

950   Riera, T., Currie, J.D., Buster, D.W., Asenjo, A.B., et al. (2011). Drosophila katanin is

a microtubule depolymerase that regulates cortical-microtubule plus-end interactions

and cell migration. Nature Cell Biology *13*, 361-369. 10.1038/ncb2206.

22. Ramalho-Santos, J., Moreno, R.D., Wessel, G.M., Chan, E.K.L., and Schatten, G.

(2001). Membrane Trafficking Machinery Components Associated with the

Mammalian Acrosome during Spermiogenesis. Experimental Cell Research *267*, 45-

60. https://doi.org/10.1006/excr.2000.5119.

23. Shan, M.-M., and Sun, S.-C. (2021). The multiple roles of RAB GTPases in female and

male   meiosis.   Human   Reproduction   Update   *27*,   1013-1029.

10.1093/humupd/dmab019.

24. Blacque, O.E., Scheidel, N., and Kuhns, S. (2018). Rab GTPases in cilium formation

and function. Small GTPases *9*, 76-94.

25. Lo, J.C., Jamsai, D., O'Connor, A.E., Borg, C., Clark, B.J., Whisstock, J.C., Field,

M.C., Adams, V., Ishikawa, T., and Aitken, R.J. (2012). RAB-like 2 has an essential

role in male fertility, sperm intra-flagellar transport, and tail assembly.

26. Hatakeyama, E., and Hayashi, K. (2018). KATNAL1 is a more active and stable

isoform of katanin, and is expressed dominantly in neurons. Biochemical and

biophysical research communications *507*, 389-394.

27. Rogers, G.C., Rogers, S.L., and Sharp, D.J. (2005). Spindle microtubules in flux.

Journal of Cell Science *118*, 1105-1116.

28. Rath, U., and Sharp, D.J. (2011). The molecular basis of anaphase A in animal cells.

Chromosome Research *19*, 423-432. 10.1007/s10577-011-9199-2.

29. Jiang, K., Rezabkova, L., Hua, S., Liu, Q., Capitani, G., Altelaar, A., Heck, A.J.,

Kammerer, R.A., Steinmetz, M.O., and Akhmanova, A. (2017). Microtubule minus-

end regulation at spindle poles by an ASPM–katanin complex. Nature cell biology *19*,

480-492.

30.   Zhang, D., Rogers, G.C., Buster, D.W., and Sharp, D.J. (2007). Three microtubule

977   severing enzymes contribute to the "Pacman-flux" machinery that moves

978   chromosomes. The Journal of cell biology *177*, 231-242.

979   31.   Elad, N., Abramovitch, S., Sabanay, H., and Medalia, O. (2011). Microtubule

980   organization in the final stages of cytokinesis as revealed by cryo-electron tomography.

981   Journal of Cell Science *124*, 207-215. 10.1242/jcs.073486.

982   32.   Cheers, S.R., O'Connor, A.E., Johnson, T.K., Merriner, D.J., O'Bryan, M.K., and

983   Dunleavy, J.E. (2022). Spastin is an essential regulator of male meiosis, acrosome

984   formation, manchette structure and nuclear integrity. bioRxiv.

985   33.   Lehti, M., and Sironen, A. (2016). Formation and function of the manchette and

986   flagellum during spermatogenesis. Reproduction *151*, R43-54.

987   34.   Cotton, L.M., Meilak, M.L., Templeton, T., Gonzales, J.G., Nenci, A., Cooney, M.,

988   Truman, D., Rodda, F., Lynas, A., and Viney, E. (2015). Utilising the resources of the

989   International Knockout Mouse Consortium: the Australian experience. Mammalian

990   Genome *26*, 142-153.

991   35.   Sadate-Ngatchou, P.I., Payne, C.J., Dearth, A.T., and Braun, R.E. (2008). Cre

992   recombinase activity specific to postnatal, premeiotic male germ cells in transgenic

993   mice. Genesis *46*, 738-742.

994   36.   Dunleavy, J.E.M., O'Connor, A.E., and O'Bryan, M.K. (2019). An optimised STAPUT

995   method for the purification of mouse spermatocyte and spermatid populations.

996   Molecular Human Reproduction *25*, 675-683. 10.1093/molehr/gaz056.

997   37.   Livak, K.J., and Schmittgen, T.D. (2001). Analysis of Relative Gene Expression Data

998   Using Real-Time Quantitative PCR and the 2− ΔΔCT Method. Methods *25*, 402-408.

999 38. Houston, B.J., Conrad, D.F., and O'Bryan, M.K. (2020). A framework for high-resolution phenotyping of candidate male infertility mutants: from human to mouse. Human Genetics, 1-28.

1002 39. Cotton, L., Gibbs, G.M., Sanchez-Partida, L.G., Morrison, J.R., de Kretser, D.M., and O'Bryan, M.K. (2006). FGFR-1 signaling is involved in spermiogenesis and sperm capacitation. Journal of Cell Science *119*, 75-84.

1005 40. Baker, M.A., Hetherington, L., Weinberg, A., and Velkov, T. (2014). Phosphopeptide analysis of rodent epididymal spermatozoa. JoVE (Journal of Visualized Experiments), e51546.

1008 41. Gibbs, G.M., Orta, G., Reddy, T., Koppers, A.J., Martínez-López, P., Luis de la Vega-Beltran, J., Lo, J.C., Veldhuis, N., Jamsai, D., and McIntyre, P. (2011). Cysteine-rich secretory protein 4 is an inhibitor of transient receptor potential M8 with a role in establishing sperm function. Proceedings of the National Academy of Sciences *108*, 7034-7039.

1013 42. O'Bryan, M.K., Clark, B.J., McLaughlin, E.A., D'Sylva, R.J., O'Donnell, L., Wilce, J.A., Sutherland, J., O'Connor, A.E., Whittle, B., and Goodnow, C.C. (2013). RBM5 is a male germ cell splicing factor and is required for spermatid differentiation and male fertility. PLoS genetics *9*, e1003628.

1017 43. Gibbs, G.M., Lo, J.C.Y., Nixon, B., Jamsai, D., O'Connor, A.E., Rijal, S., Sanchez-Partida, L.G., Hearn, M.T.W., Bianco, D.M., and O'Bryan, M.K. (2010). Glioma Pathogenesis-Related 1-Like 1 Is Testis Enriched, Dynamically Modified, and Redistributed during Male Germ Cell Maturation and Has a Potential Role in Sperm-Oocyte Binding. Endocrinology *151*, 2331-2342. 10.1210/en.2009-1255.

44. Jamsai, D., Bianco, D.M., Smith, S.J., Merriner, D.J., Ly-Huynh, J.D., Herlihy, A., Niranjan, B., Gibbs, G.M., and O'Bryan, M.K. (2008). Characterization of

gametogenetin 1 (GGN1) and its potential role in male fertility through the interaction

with the ion channel regulator, cysteine-rich secretory protein 2 (CRISP2) in the sperm

tail. Reproduction *135*, 751-760.

45.   R Core Development Team (2015). R: A language and environment for statistical

computing. R Foundation for Statistical Computing.

46.   Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting Linear Mixed-Effects

Models Using lme4. Journal of Statistical Software *67*. 10.18637/jss.v067.i01.

47.   Brooks, M.E., Kristensen, K., van Benthem, K.J., Magnusson, A., Berg, C.W., Nielsen,

A., Skaug, H.J., Machler, M., and Bolker, B.M. (2017). glmmTMB balances speed and

flexibility among packages for zero-inflated generalized linear mixed modeling. The R

Journal *9*, 378-400.

48.   Lin, Y.-H., Huang, C.-Y., Ke, C.-C., Wang, Y.-Y., Lai, T.-H., Liu, H.-C., Ku, W.-C.,

Chan, C.-C., and Lin, Y.-H. (2020). ACTN4 mediates SEPT14 mutation-induced sperm

head defects. Biomedicines *8*, 518.

49.   Hsu, W.H., Wang, W.J., Lin, W.Y., Huang, Y.M., Lai, C.C., Liao, J.C., and Chen, H.C.

(2018). Adducin-1 is essential for spindle pole integrity through its interaction with

TPX2. EMBO Rep *19*. 10.15252/embr.201745607.

50.   Chan, P.C., Hsu, R.Y., Liu, C.W., Lai, C.C., and Chen, H.C. (2014). Adducin-1 is

essential for mitotic spindle assembly through its interaction with myosin-X. J Cell Biol

204, 19-28. 10.1083/jcb.201306083.

51.   Adams, M., Simms, R.J., Abdelhamed, Z., Dawe, H.R., Szymanska, K., Logan, C.V.,

Wheway, G., Pitt, E., Gull, K., Knowles, M.A., et al. (2012). A meckelin-filamin A

interaction   mediates   ciliogenesis.   Hum   Mol   Genet   *21*,   1272-1286.

10.1093/hmg/ddr557.

52.   Yuan, S., Stratton, C.J., Bao, J., Zheng, H., Bhetwal, B.P., Yanagimachi, R., and Yan,

1049   W. (2015). Spata6 is required for normal assembly of the sperm connecting piece and

1050   tight head-tail conjunction. Proc Natl Acad Sci U S A *112*, E430-439.

1051   10.1073/pnas.1424648112.

1052   53.   Sangokoya, C., and Blelloch, R. (2020). MicroRNA-dependent inhibition of PFN2

1053   orchestrates ERK activation and pluripotent state transitions by regulating endocytosis.

1054   Proc Natl Acad Sci U S A *117*, 20625-20635. 10.1073/pnas.2002750117.

1055   54.   Lee, N.P.Y., Mruk, D.D., Conway, A.M., and Cheng, C.Y. (2004). Zyxin, Axin, and

1056   Wiskott-Aldrich Syndrome Protein Are Adaptors That Link the Cadherin/Catenin

1057   Protein Complex to the Cytoskeleton at Adherens Junctions in the Seminiferous

1058   Epithelium of the Rat Testis. Journal of Andrology *25*, 200-215.

1059   https://doi.org/10.1002/j.1939-4640.2004.tb02780.x.

1060   55.   Qi, Y., Jiang, M., Yuan, Y., Bi, Y., Zheng, B., Guo, X., Huang, X., Zhou, Z., and Sha,

1061   J. (2013). ADP-ribosylation factor-like 3, a manchette-associated protein, is essential

1062   for mouse spermiogenesis. Molecular Human Reproduction *19*, 327-335.

1063   10.1093/molehr/gat001.

1064   56.   Barbosa, D.J., Teixeira, V., Duro, J., Carvalho, A.X., and Gassmann, R. (2021).

1065   Dynein-dynactin segregate meiotic chromosomes in C. elegans spermatocytes.

1066   Development *148*. 10.1242/dev.197780.

1067   57.   Wu, C.-H., Zong, Q., Du, A.-L., Zhang, W., Yao, H.-C., Yu, X.-Q., and Wang, Y.-F.

1068   (2016). Knockdown of Dynamitin in testes significantly decreased male fertility in

1069   Drosophila melanogaster. Developmental biology *420*, 79-89.

1070   58.   Fouquet, J., Kann, M., Soues, S., and Melki, R. (2000). ARP1 in Golgi organisation

1071   and attachment of manchette microtubules to the nucleus during mammalian

1072   spermatogenesis. Journal of Cell Science *113*, 877-886. 10.1242/jcs.113.5.877.

59.   Tarkar, A., Loges, N.T., Slagle, C.E., Francis, R., Dougherty, G.W., Tamayo, J.V.,

Shook, B., Cantino, M., Schwartz, D., Jahnke, C., et al. (2013). DYX1C1 is required

for axonemal dynein assembly and ciliary motility. Nature Genetics *45*, 995-1003.

10.1038/ng.2707.

60.   Tang, E.I., Mok, K.W., Lee, W.M., and Cheng, C.Y. (2015). EB1 regulates tubulin and

actin cytoskeletal networks at the sertoli cell blood-testis barrier in male rats: an in vitro

study. Endocrinology *156*, 680-693. 10.1210/en.2014-1720.

61.   Wang, F., Zhang, Q., Cao, J., Huang, Q., and Zhu, X. (2008). The microtubule plus

end-binding protein EB1 is involved in Sertoli cell plasticity in testicular seminiferous

tubules. Experimental Cell Research *314*, 213-226.

https://doi.org/10.1016/j.yexcr.2007.09.022.

62.   Kanemori, Y., Koga, Y., Sudo, M., Kang, W., Kashiwabara, S.-i., Ikawa, M., Hasuwa,

H., Nagashima, K., Ishikawa, Y., and Ogonuki, N. (2016). Biogenesis of sperm

acrosome is regulated by pre-mRNA alternative splicing of Acrbp in the mouse.

Proceedings of the National Academy of Sciences *113*, E3696-E3705.

63.   Selvaraj, V., Asano, A., Page, J.L., Nelson, J.L., Kothapalli, K.S.D., Foster, J.A.,

Brenna, J.T., Weiss, R.S., and Travis, A.J. (2010). Mice lacking FABP9/PERF15

develop sperm head abnormalities but are fertile. Developmental Biology *348*, 177-

189. https://doi.org/10.1016/j.ydbio.2010.09.019.

64.   Dix, D.J., Allen, J.W., Collins, B.W., Mori, C., Nakamura, N., Poorman-Allen, P.,

Goulding, E.H., and Eddy, E.M. (1996). Targeted gene disruption of Hsp70-2 results

in failed meiosis, germ cell apoptosis, and male infertility. Proceedings of the National

Academy of Sciences *93*, 3264-3268. doi:10.1073/pnas.93.8.3264.

65.   Dix, D.J., Allen, J.W., Collins, B.W., Poorman-Allen, P., Mori, C., Blizard, D.R.,

Brown, P.R., Goulding, E.H., Strong, B.D., and Eddy, E.M. (1997). HSP70-2 is

required for desynapsis of synaptonemal complexes during meiotic prophase in

juvenile and adult mouse spermatocytes. Development *124*, 4595-4603.

10.1242/dev.124.22.4595.

66.   Govin, J., Caron, C., Escoffier, E., Ferro, M., Kuhn, L., Rousseaux, S., Eddy, E.M.,

Garin, J., and Khochbin, S. (2006). Post-meiotic shifts in HSPA2/HSP70.2 chaperone

activity during mouse spermatogenesis. J Biol Chem *281*, 37888-37892.

10.1074/jbc.M608147200.

67.   Yang , F., Eckardt , S., Leu , N.A., McLaughlin , K.J., and Wang , P.J. (2008). Mouse

TEX15 is essential for DNA double-strand break repair and chromosomal synapsis

during male meiosis. Journal of Cell Biology *180*, 673-679. 10.1083/jcb.200709057.

68.   Schöpp, T., Zoch, A., Berrens, R.V., Auchynnikava, T., Kabayama, Y., Vasiliauskaitė,

L., Rappsilber, J., Allshire, R.C., and O'Carroll, D. (2020). TEX15 is an essential

executor of MIWI2-directed transposon DNA methylation and silencing. Nature

Communications *11*, 3739. 10.1038/s41467-020-17372-5.

69.   Mountjoy, J.R., Xu, W., McLeod, D., Hyndman, D., and Oko, R. (2008). RAB2A: a

major subacrosomal protein of bovine spermatozoa implicated in acrosomal biogenesis.

Biol Reprod *79*, 223-232. 10.1095/biolreprod.107.065060.

70.   Morohoshi, A., Miyata, H., Oyama, Y., Oura, S., Noda, T., and Ikawa, M. (2021).

FAM71F1 binds to RAB2A and RAB2B and is essential for acrosome formation and

male fertility in mice. Development *148*. 10.1242/dev.199644.

71.   Da Costa, R., Bordessoules, M., Guilleman, M., Carmignac, V., Lhussiez, V., Courot,

H., Bataille, A., Chlémaire, A., Bruno, C., Fauque, P., et al. (2020). Vps13b is required

for acrosome biogenesis through functions in Golgi dynamic and membrane trafficking.

Cellular and Molecular Life Sciences *77*, 511-529. 10.1007/s00018-019-03192-4.

1122

<u>Figures</u>

Figure 1: Germ cell derived KATNAL1 is required for optimal male fertility

Body weight (**A**), testis weight (**B**), DSP (**C**), and epididymal sperm content (**D**) in

Katnal1$^{GCKO/GCKO}$ mice (black triangles) and *Katnal1$^{Flox/Flox}$* controls (white triangles)

(n≥3/genotype). Lines represent mean ± SD. PAS-stained testis (**E-G**) and epididymis (**H**)

sections. Asterisks indicate areas of seminiferous epithelium with vacuoles or devoid of germ

cells. Black arrowheads indicate retained spermatids. (**I**) Hematoxylin and eosin-stained cauda

epididymal sperm. Abnormalities were observed in the vast majority of sperm from

Katnal1$^{GCKO/GCKO}$ mice. Defects included abnormal sperm head shapes - "knobby" (b) and

"hammer" head (c, f), abnormal head sperm head position (d), decapitated sperm (e,), coiled

mid-pieces (g) and occasionally, mitochondrial sheath defects (h). (**J**) Quantification of

abnormal cauda epididymal sperm phenotypes in *Katnal1$^{Flox/Flox}$* (white triangles) versus

Katnal1$^{GCKO/GCKO}$ (black triangles). (n≥3/genotype). Percentage of motile sperm (**K**) and

percentage of progressively motile sperm (**L**) from *Katnal1$^{Flox/Flox}$* (white triangles) versus

Katnal1$^{GCKO/GCKO}$ (black triangles) males. Scale bars in **E-F** = 50 μm, **G** = 10 μm, **H** = 40 μm,

I = 10 μm. *$P<0.05$, **$P<0.01$, ***$P<0.01$, ****$P<0.001$.

Figure 2: KATNAL1 regulates chromosome alignment, segregation and cytokinesis

during male meiosis

(**A-B**) Seminiferous tubule sections immunolabelled for α-tubulin as a component of MTs.

Nuclei are counterstained with hematoxylin. Green arrowheads indicate misaligned/lagging

chromosomes. (**C**) PAS-stained testis sections showing abnormally large round spermatid

nuclei (b,c; yellow arrowhead) and binucleated round spermatids (d-e; orange arrowheads) in

1147   *Katnal1*^{GCKO/GCKO} mice. (**D-J**) Germ cell apoptosis in *Katnal1*^{GCKO/GCKO} mice was assessed by

1148   immunolabelling of seminiferous tubules for cleaved-caspase 3 and 7. The average number of

1149   cleaved-caspase 3 and/or 7 positive germ cells per seminiferous tubule is graphed in **D** and

1150   representative images are shown in **E-G**. In **D** lines represent mean ±SD. A minimum of 80

1151   randomly selected tubules per mouse were counted (n≥3/genotype). Scale bars in **A-B,G-J** =

1152   10 μm, **C**=5 μm. **E-F**= 50 μm.

1153

1154

Figure 3: KATNAL1 is a regulator of flagella formation and the MT manchette in male germ cells

Testis sections immunolabelled for the axoneme marker acetylated tubulin in *Katnal1^Flox/Flox^* (**A**) and *Katnal1^GCKO/GCKO^* (**B**). (**C**) TEM of flagella ultrastructure in cauda epididymal sperm of *Katnal1^Flox/Flox^* and *Katnal1^GCKO/GCKO^* mice. MS= mitochondrial sheath, ODFs = outer dense fibers, Ax=axoneme FS = fibrous sheath. White arrows indicate supernumerary MT doublets.

Red arrows indicate ectopic MT doublets. Green arrows indicate ectopic ODFs. Yellow

asterisks indicate missing axoneme MT doublets/or central pair. (**D**) TEM of HTCA

ultrastructure in round (**a-e**) and elongated spermatids (**f-j**) in *Katnal1*$^{Flox/Flox}$ and

Katnal1$^{GCKO/GCKO}$ mice. Red arrowhead indicated supernumerary centriole. Blue asterisks

indicate absent HTCA. Yellow arrowheads indicate HTCAs are abnormally detached from the

nucleus. (**E**) PAS-stained testis sections showing *Katnal1*$^{Flox/Flox}$ and *Katnal1*$^{GCKO/GCKO}$

elongating spermatids. (**F**) Testis sections immunolabelled for α-tubulin as a maker of the

manchette in *Katnal1*$^{Flox/Flox}$ and *Katnal1*$^{GCKO/GCKO}$ mice. Progressive steps of manchette

formation, migration and dissolution are shown left to right. (**G-H**) TEM showing manchette

structure in *Katnal1*$^{Flox/Flox}$ and *Katnal1*$^{GCKO/GCKO}$ elongating spermatids. Red arrowheads

indicate the perinuclear ring. Scale bars in **A-B** = 20 μm**, Ca-i** = 200 nm, **Cj-m** = 500 nm, **D** =

1 μm, **E-F**=10 μm, **G-H** = 2 μm**.**

1173

Figure 4: Double germ cell deletion of KATNA1 and KATNAL1 results in a severe

oligozoospermia-like phenotype

qPCR analysis of *Katna1* (**A**) and *Katnal1* (**B**) transcript levels in *Katna1/al1$^{GCKO/GCKO}$* (black

triangles, *dGCKO*) isolated spermatocytes relative to *Katna1/al1$^{Flox/Flox}$* (white triangles,

dFlox) isolated spermatocytes (n≥3/genotype). Body weight (**C**), testis weight (**D**), testis DSP

(**E**), and epididymal sperm content (**F**) in *Katna1/al1$^{GCKO/GCKO}$* (black triangles) compared to

Katna1/al1$^{Flox/Flox}$ controls (white triangles) (n≥3/genotype). Lines represent mean ± SD. PAS-

stained testis (**G-H**) sections from *Katna1/al1$^{GCKO/GCKO}$* and *Katna1/al1$^{Flox/Flox}$* mice. Magenta

arrowheads indicate pyknotic spermatocytes and green arrowheads indicate pyknotic

spermatids. Asterisks indicate areas of seminiferous epithelium with vacuoles or devoid of

germ cells. Analysis of germ cell apoptosis in *Katnal1$^{GCKO/GCKO}$* mice relative to

Katna1/al1$^{Flox/Flox}$ mice (**I-O**). Representative images of testis immunolabelling of cleaved-

caspase 3 and 7 are shown in (**I-N**). Roman numerals denote seminiferous tubule stage. The

average number of cleaved-caspase 3 and/or 7 positive germ cells per seminiferous tubule per

mouse is graphed in (**O**). In **O** lines represent mean ±SD. A minimum of 50 randomly selected

tubules per mouse were counted (n≥3/genotype). PAS-stained epididymis sections (**P**) from

Katna1/al1$^{GCKO/GCKO}$ and *Katna1/al1$^{Flox/Flox}$* mice. Scale bars in **G-J** = 50 μm, in **K-N** = 10 μm,

P = 40 μm. *$P<0.05$, **$P<0.01$, ***$P<0.01$, ****$P<0.001$.

Figure 5: KATNA1 and KATNAL1 compensate for each other in male meiosis

(**A-B**) *Katnal1/al1^(Flox/Flox)* (*dFlox*) and *Katna1/al1^(GCKO/GCKO)* (*dGCKO*) seminiferous tubule sections immunolabelled for MTs (α-tubulin). Nuclei are counterstained with hematoxylin. Green arrowheads indicate misaligned/lagging chromosomes. (**C**) PAS-stained *Katnal1/al1^(Flox/Flox)* (*dFlox*) and *Katna1/al1^(GCKO/GCKO)* (*dGCKO*) testis sections showing abnormally large round spermatid nuclei (b; yellow arrowhead) and binucleated round spermatids (c; orange arrowheads) in *Katna1/al1^(GCKO/GCKO)* mice. Scale bars in **A**= 10 μm, in **B-C**= 5 μm.

1201

Figure 6: **KATNA1 and KATNAL1 compensation in spermatid acrosome formation and**

MT regulation

(**A**) PAS-stained testis sections showing acrosome formation in *Katna1/al1$^{Flox/Flox}$* and

Katna1/al1$^{GCKO/GCKO}$ mice. Progressive steps of acrosome formation are shown left to right.

Roman numerals denote seminiferous tubule stage. Orange arrowheads indicate nuclei with

more than one proacrosomal vesicle attached. Yellow arrowheads indicate acrosomal vesicles

not attached to the nucleus. Green arrowheads indicate ectopic PAS positive vesicles

abnormally dispersed in round spermatid cytoplasm. Blue arrowheads indicate elongating

spermatids with abnormal gaps/fragmentation of the acrosomes. (**B**) Electron micrographs

showing acrosome and Golgi apparatus ultrastructure in *Katna1/al1$^{Flox/Flox}$* and

Katna1/al1$^{GCKO/GCKO}$ mice. Orange arrowheads indicate supernumerary site of acrosome

formation. Blue arrowheads indicate numerous small unfused acrosomal vesicles docked at the

nuclear membrane/fragmentation of the acrosome. (**C-F**) *Katna1/al1$^{Flox/Flox}$* (*dFlox*) and

Katna1/al1$^{GCKO/GCKO}$ (*dGCKO*) mouse testis sections immunolabelled for the sperm tails

marker acetylated tubulin. (**G**) *Katna1/al1$^{Flox/Flox}$* and *Katna1/al1$^{GCKO/GCKO}$* testis sections

immunolabelled for α-tubulin as a maker of the manchette. Progressive steps of manchette

formation, migration and dissolution are shown left to right. TEM showing nuclear and

manchette morphology in *Katna1/al1$^{Flox/Flox}$* (*dFlox*) and *Katnal1$^{GCKO/GCKO}$* mouse elongating

spermatids. Red arrowheads indicate the perinuclear ring. Blue arrowheads indicate MTs. Scale

bars in **A, C-D** = 10 μm, **B,H** = 1 μm, **E-F** = 50 μm, **G**=10 μm. Nu = Nucleus, Cy =cytoplasm

Gg=Golgi apparatus, Ac=acrosome.

1223

Figure 7: Proposed model of katanin regulation of spermatogenesis

KATNB1-independent and -dependent functions of each katanin A-subunit during male germ cell development are depicted based on the data defined herein and defined for KATNAL1 Sertoli cell function in [13], for KATNAL2 in [5] and KATNB1 defined in [11,12]. A shared line indicates two complexes have redundant/compensatory function in the process indicated. HTCA = head to tail coupling apparatus, MTs = microtubule, ODF = outer dense fibers.