1

Behavioral dissection of hunger states in *Drosophila*

KJ Weaver[1], S Raju[2], RA Rucker[3], TS Chakraborty[1], RA Holt[2], SD Pletcher[1]

1. Department of Molecular and Integrative Physiology and Geriatrics Center, Biomedical

Sciences and Research Building, University of Michigan, Ann Arbor, MI 48109, U.S.A.

2. College of Literature, Science, and the Arts, Biomedical Sciences and Research Building,

University of Michigan, Ann Arbor, MI 48109, U.S.A.

3. Neuroscience Graduate Program, University of Michigan, Ann Arbor, MI 48109, U.S.A.

Abstract

Hunger is a motivational drive that promotes feeding, and it can be generated by the

physiological need to consume nutrients as well as the hedonic properties of food. Brain circuits

and mechanisms that regulate feeding have been described, but which of these contribute to

the generation of motive forces that drive feeding is unclear. Here, we describe our first efforts

at behaviorally and neuronally distinguishing hedonic from homeostatic hunger states in

Drosophila melanogaster and propose that this system can be used as a model to dissect the

molecular mechanisms that underlie feeding motivation. We visually identify and quantify

behaviors exhibited by hungry flies and find that increased feeding duration is a behavioral

signature of hedonic feeding motivation. Using a genetically-encoded marker of neuronal

activity, we find that the mushroom body (MB) lobes are activated by hedonic food

environments, and we use optogenetic inhibition to implicate a dopaminergic neuron cluster

(PAM) to α'/β' MB circuit in hedonic feeding motivation. The identification of discrete hunger

states in flies and the development of behavioral assays to measure them offers a framework to

begin dissecting the molecular and circuit mechanisms that generate motivational states in the

brain.

Introduction

Hedonist philosophers believed that pleasure is the supreme goal of life and that behavior is determined by the pursuit of it (Aristotle et al., 2004). Hunger for tasty delicacies, for example, is a particularly potent motivator of behavior, as it can be promoted not only by a physiological need to consume nutrients, but also by the hedonic properties of food. Due to its dichotomous nature, defining behavioral and neural correlates of hunger is a challenge that has perplexed scientists for over a century. Early physiologists wondered how to explain hunger and where it might originate, and they eventually described hunger as a dull ache or gnawing sensation arising from the stomach or abdominal region (Cannon & Washburn, 1993). This historical description has aided the discovery of hormones like ghrelin, insulin, and others that communicate information from the periphery to the brain and have roles in controlling feeding in response to energetic or nutritional needs, a state we now refer to as homeostatic hunger. In the decades since these discoveries, it has become clear that peripheral signals and others are integrated in the central nervous system to promote changes in behavior and reward signaling, which we interpret as indicators of hunger, but the nature of the motivational brain systems that drive these outputs remains less straight forward to identify and quantify.

Homeostatic feeding, or "need-based" intake, is defined as food intake that is necessary to ensure caloric and nutritional needs are met, and it has been quantified by measuring bulk food intake. Such measures may be of total caloric intake or consumption of specific nutrients, but it is unclear whether these measures are representative of hunger that is driven by energetic need, nutrient deficiency, or some other motivation, which has led to variation in the way measures of homeostatic hunger are reported and interpreted (for review of mammalian brain circuits that modulate feeding, see (Alcantara et al., 2022; Andermann & Lowell, 2017; Saper et al., 2002)). However, the observation that most animals, including humans, regulate food intake around a protein target and the identification of discrete neuronal populations that regulate

protein-specific feeding in model systems suggest that levels of homeostatic hunger may be

primarily determined by the amount of protein an animal requires and thus, in recent years,

protein intake has become a common method for measuring need-based hunger (Gosby et al.,

2011; Liu et al., 2017; Ro et al., 2016; Simpson et al., 2003; Weaver et al., 2022).

Our understanding of hunger has continued to evolve, due in part to the observation that the

pleasurable effects of food also stimulate feeding independently of physiological need, a state

referred to as hedonic hunger (Lowe & Butryn, 2007). Hedonic, or "need-free", intake is thought

to be evoked by the incentive salience or tastiness of food, and thus, the sweetness of the food

environment is predicted to play a larger role (Lowe & Butryn, 2007). Indeed, hedonic hunger is

likely the predominant driver of feeding in modern human environments where food is readily

available and energy deprivation is seldom experienced. Hedonic responses to various food or

environmental manipulations have been described as either "liking" responses (relating to the

pleasure of a rewarding experience) or "wanting" responses (relating to the motivation to seek a

reward), which arise from different neuronal substrates (reviewed in (Berridge, 2009)).

Accordingly, "liking" is measured in rats by tracking facial responses in response to sweet

tastants, while "wanting" is measured by tracking latency to search for and consume a tasty

food treat or by using the novelty suppressed feeding assay that attempts to measure an

animal's ability to overcome an aversive environment to consume a food reward (Arcego et al.,

2020; Pecina et al., 2006). While observations like these have greatly aided our ability to identify

hedonic hunger states in mammals, much less is known about their neuronal and mechanistic

origins relative to homeostatic hunger.

Although we now understand that hunger stratifies into two distinct states, one driven by need

and the other by pleasure, we have a limited understanding of their behavioral and neuronal

distinctions and connections. In mammals, brain circuits that modulate homeostatic or hedonic

appetites have been identified. The well-described AGRP neurons in the arcuate nucleus of the

hypothalamus, for example, are critical components of a circuit for homeostatic feeding (for

review, see (Sternson & Eiselt, 2017)). They express receptors for peripherally released

hormones that signal energetic state, are more active upon food deprivation, and transmit a

negative valence signal that drives animals to find and consume food when the need arises

(Betley et al., 2015; Denis et al., 2015; Willesen et al., 1999; Zigman et al., 2006). In addition to

systems that seem to motivate homeostatic feeding, distinct circuits have also been associated

with hedonic feeding including the monoaminergic populations that express serotonin and/or

dopamine, and the mesolimbic dopamine pathway (Ahn et al., 2022; Altherr et al., 2021; Denis

et al., 2015; Ghiglieri et al., 1997; Rossi & Stuber, 2018). While there is evidence to suggest that

both branches of hunger are modulated by homeostatic state and hedonic feedback (e.g.,

(Denis et al., 2015; Krashes et al., 2011; Zigman et al., 2006)), the mechanistic origins that

determine how various hunger drives work together and independently to generate feeding

responses have yet to be fully described.

Simpler model systems, such as the fruit fly, *Drosophila melanogaster*, have proven valuable for

insights into the neural circuitry that modulates feeding, but the nature of their hunger drives is

unknown. Whether flies experience hedonic hunger, for example, is unclear, although it seems

likely that they do; hungry flies learn to associate odors with a sucrose reward and when fed a

high-sugar diet they overeat and become obese (Huetteroth et al., 2015; May et al., 2019).

These behaviors are all thought to occur through mechanisms involving dopaminergic neurons

in the fly brain which, when artificially activated, promote foraging and food intake (Tsao et al.,

2018). Honeybees have also exhibited signatures of food wanting, a component of hedonic

hunger, which manifests through activation of dopaminergic populations in the brain that are

reflective of reward expectation (Huang et al., 2022).

To better understand the different motive forces that drive feeding, it seems clear that there is a

need to move beyond measures of food intake or foraging behavior, which have been useful for

identifying external and internal factors that change the way an animal eats, in order to dissect

when and to what degree animals may be experiencing different types of hunger. Here we

describe our first efforts to address this need by generating clear behavioral definitions of

hunger states in *Drosophila* with the hope of providing an attractive system for defining the

molecular features of their independent roles and the relationships between them. We

characterize a repertoire of feeding micro-behaviors in the fly and use them to argue that

Drosophila experience both homeostatic and hedonic hunger. We identify neural correlates of

hedonic hunger in specific lobes of the *Drosophila* mushroom body and show that they are

required for hedonic feeding. Finally, we develop a framework to examine the relationships

between hedonic and homeostatic hunger states and present evidence for the existence of a

system that acts as a potentiator of the relationship between them. Translating findings about

the nature of feeding drives and the relationships among them could have far reaching

implications for obesity and energy balance research and other behaviors influenced by

motivational drives, including drug addiction and eating disorders.

Results

We set out to complement and enrich traditional measures of feeding by identifying new

behavioral metrics that might be useful for classifying different hunger states. Fly feeding

behavior is typically described as changes in volumetric food intake, which is difficult to

accurately quantify at the individual level due to the small amount of food ingested (Deshpande

et al., 2014; Shell et al., 2021; Shell et al., 2018), or by using automated systems that detect

physical interactions with food sources (Itskov et al., 2014; Ro et al., 2014). These automated

devices have revealed rhythmic patterns of feeding behavior in flies that mimic feeding

strategies in other animals, suggesting that the behaviors are highly organized and centrally

regulated. They are, however, unable to distinguish different stereotypical actions that a freely-

moving fly might exhibit when interacting with a food source. Detailed behavioral

characterizations of this sort have proven to be useful read-outs of neural states in other

contexts and have the potential to inform our understanding of the motivational states that drive

feeding (Chen et al., 2002; Duistermars et al., 2018; Ning et al., 2022). Much like human infants,

for example, facial expressions and tongue movements in rats in response to appetitive tastants

have been used to identify hedonic reactions and the brain systems that mediate them

(Berridge, 1996).

To identify nuances in individual fly feeding, we developed video systems for manual and high-

throughput annotation of what flies do when they eat. In all cases, we recorded freely-moving

flies placed in feeding chambers of the Fly Liquid-Food Interaction Counter (FLIC) system,

which is designed to continuously record individual flies' interactions with food sources (Ro et

al., 2014). We began by simply recording flies in the FLIC and manually curating each video in

its entirety (Fig. 1A). With the hope of observing the greatest variety of behaviors, we collected

observations from flies ranging from 5-20 days old, from mated female and male sated and

starved flies, from flies housed singly or in groups of up to three, and from flies interacting with

5% or 10% sucrose food. We observed 30 flies and nearly 300 "events", which were

demarcated by intervals greater than one second during which the fly was not in physical

contact with the food. Events were easy to identify, and different behavioral characteristics

associated with events were easy to distinguish from each other (Slooten, 1994). Rare events

(observed in fewer than 5% of flies) were not considered further, and events during which sight

of the fly was out of focus or obscured (typically 1-2 events per fly) were ignored. Based on the

remaining events, we created a library of user-defined behaviors that captured common,

distinct, and definable interactions of flies with their food (Table 1).

We classified the behaviors we observed into five main categories: two non-feeding behaviors

("other" and "interaction" events) and three feeding behaviors ("fast" feeding events, "long"

feeding events, and "long/quick" feeding events) (Table 1). The first feeding event type

comprised fast feeding events (termed "F" or "FI") during which we observed contact of the

proboscis with the food for less than four seconds. These types of events were common

(occurring in ~95% of flies, Fig. 1B, 1C) and were sometimes simultaneous with the interaction

of a fly's legs with the food. Longer feeding events (termed "L", "LI", or "LQ") were also

identified, during which time we observed contact of the proboscis with the food for more than

four seconds. Long events were less common than F-type events and could also include

simultaneous interaction of legs with food. Long feeding events were either a continuous and

sustained interaction of the proboscis with the food ("L" and "LI", occurring in ~85% of flies, Fig.

1B, 1C) or, less commonly, a continuous interaction during which the proboscis moved quickly

in and out of the food ("LQ", occurring in ~60% of flies, Fig. 1B, 1C). The first type of non-

feeding behavior was termed an interaction event ("IF" or "IH") during which the fly interacted

with the food only using its legs; no proboscis contact was observed. These putative tasting

interactions often occurred in a "pitter-patter" motion, as if the fly were "playing" with their food,

and they occurred in greater than 90% of flies (Fig. 1B, 1C). Lastly, we occasionally observed

other non-feeding events ("G" or "UC") that, nevertheless, often occurred just before or after

feeding (occurring in ~60% of flies, Fig. 1B, 1C). These tended to be various grooming

behaviors, including using the forelegs to rub the head after interacting with food.

We next asked whether our behavioral classification could be used to distinguish hungry from

sated flies. Recognizing a need to significantly increase the number of observed feeding events

to provide sufficient statistical power to detect differences in behavioral tendencies, we designed

a direct communication link between our custom DTrack video software and the FLIC system to

trigger video capture only when there was physical contact between a fly and the food source,

together with 5s windows before and after the contact. This allowed us to create video

compilations of feeding behaviors for fully fed or starved (24hr) flies over several hours. Over

500 distinct events in these compilations were then manually scored using our classification of

feeding behaviors, and comparisons were made between fully-fed and starved flies for each

behavioral category in Table 1. We observed that starved flies exhibited a greater frequency of

feeding behaviors overall, as expected (sated event frequency = 0.606 events/minute compared

to starved event frequency = 0.788 events/minute, $p = 0.061$, data not shown), and that the

relative frequency with which L-type feeding events occurred was increased (Fig. 1D, quantified

at right). The observation that starved flies exhibit more frequent long-duration events is

consistent with recent reports that starved flies increase "sips per burst" – a measure that is

similar to our definition of L and LQ events (Itskov et al., 2014).

Qualitative aspects of the relationships between individual behaviors - that is, the architecture

that results from sequences of behaviors - can reveal complexities that may also distinguish

between different neural states (Casarrubea et al., 2019; Casarrubea et al., 2015; Flavell et al.,

2022). Thus, we asked whether certain sequential patterns in fly feeding behavior appear

repeatedly and whether they are affected by starvation. To do this, we used classic sequence

analysis to map the probability that one event follows another in a sequence more often than

what is expected to occur by chance alone - an approach that has been useful for identifying

stereotyped patterns or structures of behaviors that tend to occur together (Ning et al., 2022;

Slooten, 1994). We observed that hungry flies, whom we had starved for 24 hours, but not sated

flies, followed F feeding events with additional F feeding events (transitional probability = 0.6 in

starved flies with p-value <0.001, compared to a non-significant transitional probability = 0.55 in

fully-fed flies) and L-type feeding events with additional L-type feeding events (transitional

probability = 0.439 in starved flies with p-value <0.001 compared to non-significant transitional

probability = 0.1538 in fully-fed flies) more often than would be expected by chance (Fig. 1E). In

contrast, fully-fed flies were likely to engage in F-type events after grooming or interaction

events and were more likely to groom following most other events, which may be reflective of a

less goal-oriented behavioral state compared to that of starved flies (Fig. 1E).

Having established that long-duration events occur more frequently and in a predictable

sequence in starved flies, we sought to replicate this finding in flies during their normal circadian

"meal-time", when they are presumably physiologically hungry but not yet starved. When

monitored over several days on the FLIC, flies exhibited circadian meal patterns that

corresponded to the light cycle, with one peak in the morning when lights turn on and a second

peak in the evening when lights turn off (e.g. Fig. 1 – Fig. Supp. 1). We obtained 24hr video

compilations of nearly 700 feeding events in the FLIC and used the flies' endogenous feeding

rhythm to designate "meal" and "non-meal" times of day. We manually scored the videos during

each time for feeding events (fast, long, and long/quick events). We found that during mealtimes

flies exhibited a significantly greater frequency of LQ events, while the frequencies of F and L

events were statistically indistinguishable between times of day (Fig 1F). Thus, both starved

flies and flies during their circadian meal-time exhibit an increase in the total number of feeding

events together with an increase in the relative frequency of long feeding events.

The FLIC generates several quantitative signal measures when flies interact with food, and we

asked whether any of these numerical metrics might substitute for direct visual observation and

distinguish among fast feeding events, long feeding events, and interactions. To determine how

signal measures change in accordance with a fly's behavior, we scored 15 video compilations

(resulting in 332 independent events) of flies interacting on the FLIC for F-type, L-type, or I-type

events, and we matched each visually identified behavior to its corresponding FLIC signal data

(e.g., Figs. 2A-2C). We evaluated differences in the characteristics of signal intensity for each

event (i.e., its variance, minimum, maximum, average, and total) as well as the duration of the

signal from each event. We found that only signal duration effectively distinguished long events

from both fast events and interactions (Fig. 2D-2I). Thus, we established that longer signal

durations from the FLIC are representative of L-type feeding events. We also noted that

maximum signal intensity and the variance of the signal can distinguish interactions from both

types of feeding events, which may be broadly useful to distinguish non-feeding interactions

(i.e., tasting events) from feeding events (Ro et al., 2014).

Development of an assay to identify homeostatic and hedonic feeding

Having established that hungry flies increase the duration of their feeding events as well as the

total number of events (i.e., Figure 3A, top panel), we next asked if we might be able to identify

distinct hunger drives in flies and, if so, whether our behavioral and quantitative metrics would

distinguish between them. Recognizing that the original assays used for behavioral observation,

in which flies were exposed to a single, homogenous food source, were likely insufficient to

distinguish different types of feeding drives, we designed two types of food choice environments

that we might expect would distinguish homeostatic from hedonic feeding. Homeostatic feeding

drive in flies is thought to be measured by the extent to which flies choose to consume a yeast-

containing food over one consisting of sucrose only, which we will refer to hereafter as a yeast

preference (Liu et al., 2017; Ro et al., 2016). Therefore, for measures of homeostatic drive, we

exposed flies to a choice of 2% sucrose vs. [2% yeast + 1% sucrose], and we focused our

analysis on the flies' relative interactions with the yeast containing food. To measure hedonic

drive, on the other hand, we incorporated theorized requirements to experimentally measure

hedonic hunger in mammals: animals must be energy replete and they must be tested on highly

palatable food (Lowe & Butryn, 2007). We therefore exposed flies to a "hedonic choice"

environment which mimicked the "homeostatic choice" environment except that the 2% sucrose

food was replaced with sweeter 20% sucrose, and we focused our analysis on comparing

interactions with the sucrose food between the food environments (e.g., Fig. 3A bottom panel).

This design ensured flies remained sated by access to yeast while also allowing feeding on the

highly palatable 20% sucrose solution.

We observed that the 2% sucrose environment elicited an increased yeast preference in starved

compared to fully-fed female flies (Fig. 3B), reinforcing the notion that the total number of events

with yeast vs. sucrose can be used as a relative measure of homeostatic hunger. We also

observed an increase in yeast preference in fully-fed female compared to male flies (Fig. 3C),

which may be reflective of a known difference in protein requirements, and therefore

homeostatic hunger levels, between male and female flies. We next evaluated female and male

fully-fed flies in the 20% sucrose food environment compared to the 2% sucrose environment,

which elicited increased event durations with 20% sucrose compared to the 2% sucrose food

(Fig. 3D) and also elicited decreased event durations with yeast food (Fig. 3E). The total

number of events on sucrose or yeast wells was statistically indistinguishable between food

environments (Fig. 3F, 3G), suggesting that modulation of event duration, but not total number

of events, may be a feature of hedonic food environments. We also measured sucrose intake in

the control vs. hedonic environments using a method that determines how much volume a

group of flies consume in 24 hours (termed "Con-Ex" for Consumption-Excretion) (Shell et al.,

2021; Shell et al., 2018), and found that flies consume twice as much sucrose volume in the

hedonic environment (Fig. 3 – Fig. Supp. 1A). We concluded that the measurable increase in

sucrose event durations in the hedonic food environment may reflect increases in a hedonic

hunger drive while total number of events on yeast vs. sucrose food may reflect homeostatic

state.

To challenge our measure of hedonic feeding, we reasoned that if increased event durations

were indeed a result of hedonic hunger, then sweetness alone should be sufficient to increase

event durations without modulating measures of homeostatic feeding. We tested two additional

food choice environments where we altered either the sweetness or caloric content of the sugar

choice. Sweetness was manipulated by using the non-metabolizable sweetener arabinose ([2%

sucrose + 18% arabinose], which mimics the caloric and nutritive content of the 2% sucrose but

contains the sweetness of the 20% sucrose choice. Caloric content was manipulated using the

nutritive, but not sweet, sugar alcohol, sorbitol ([2% sucrose + 18% sorbitol], which mimics the

caloric content of the 20% sucrose but contains the sweetness of the 2% sucrose (Burke &

Waddell, 2011; Hassett, 1948). We observed that the sweet arabinose food environment elicited

increased event durations that were comparable to flies in the 20% sucrose environment, with

no statistical difference in total event number (Figs. 3H, 3I, Fig. 3 – Fig. Supp. 1B) and that the

caloric but not sweet sorbitol food environment was not sufficient to increase event durations or

total event number (Figs. 3J, 3K). Finally, we tested several additional food choice environments

containing a range of sucrose concentrations and observed that event durations were shortest

when the sucrose choice contained only water or 1% sucrose and increased linearly as the

sucrose concentration increased, while event durations with yeast food and total number of

sucrose events were not statistically different (Fig. 3L, 3M, Fig. 3 – Fig. Supp. 1C). Thus, sweet

taste was sufficient to evoke longer feeding event durations, while calories alone were not,

indicating that this metric obtained from our hedonic food choice paradigm may reliably quantify

hedonic feeding.

Whether and how homeostatic and hedonic feeding mechanisms interact or overlap is

controversial, and thus, we asked whether our metrics of homeostatic and hedonic hunger are

correlated among individual flies. To obtain flies with a range of homeostatic needs, we used

fully-fed and starved flies to obtain feeding measures in our control or hedonic food choice

environments. Starvation elicited an increase in yeast interactions compared to 2% sucrose

(Fig. 4A) that was more pronounced in flies starved for 48 vs. 24 hours (Fig. 4B), which is

consistent with our expectation for increased homeostatic drive. We found that starvation

seemed to marginally increase differences in sucrose event durations in hedonic vs. control

environments, indicating that homeostatic hunger may enhance hedonic feeding (Fig. 4C). For

each fly, we next calculated metrics for homeostatic drive (# interactions with yeast food) as well

as hedonic drive (sucrose event durations) and plotted their relationship along the x- and y-

axes, respectively. We reasoned that if the circuits or programs that drive hedonic and

homeostatic feeding were completely independent, then we should observe a slope close to

zero. A significantly positive or negative slope, however, would suggest that increasing

physiological need might also modulate hedonic hunger, or vice versa. We observed a

significantly positive correlation between our measures of homeostatic and hedonic feeding

(Fig. 4D), suggesting that increasing physiological need might also potentiate hedonic hunger,

or vice versa, which is consistent with the observation in mammals that hunger increases the

reward value of food (Fulton, 2010). Interestingly, we also noted that separating regression lines

from flies in the control vs. hedonic food environments revealed parallel lines whose slopes

were not significantly different but were separated by a shift along the y-axis, indicating again

that a sweeter food environment promotes longer event durations but may not modulate the

relationship between hunger drives. These findings indicate that homeostatic state is a

potentiator of hedonic feeding and establish that inferences of hunger states and measures of

feeding should include considerations of each component independently as well as their

interaction.

Identification of neural correlates of hedonic hunger

A good deal is known about neural circuits associated with homeostatic hunger in *Drosophila*,

and we wondered whether our behavioral and quantitative classification metrics could aid in

identification of representative neural correlates of a hedonic hunger state. We began by

conducting an unbiased screen for neurons that respond to hedonic food environments using

the genetically-encoded calcium indicator CaMPARI driven by a pan-neuronal GAL4 driver

(Nsyb-GAL4). CaMPARI is a green fluorescent protein that irreversibly photoconverts to red

when flies are simultaneously exposed to increases in intracellular calcium and UV light

provided by the experimenter (Edwards et al., 2020; Fosque et al., 2015). We placed groups of

3 naïve, fully-fed flies in the hedonic food choice environment or the control food environment

and allowed them to habituate for 1 hour. Flies in each chamber were then exposed to cycles of

UV light (10s on/10s off) for a total of 1 hour to label neuronal populations that were differentially

active when flies are in the hedonic vs. the control food environment. Brains were dissected and

imaged immediately, and the ratio of red:green fluorescent protein was analyzed, with a higher

ratio indicating increased neuronal activity. We observed that the mushroom body had a higher

ratio of red:green fluorescence when flies were in the hedonic food environment, indicating that

this population was more active (Fig. 5A).

To validate the findings of our unbiased screen and as a direct test of mushroom body activity in

hedonic food environments, we restricted the expression of CaMPARI to the mushroom body

using the *238Y* enhancer-trap line (Yang et al., 1995). We observed that the hedonic food

environment increased red:green fluorescence in the horizontal mushroom body lobes

compared to the control food environment (Fig. 5B). The *Drosophila* mushroom body is a well-

described structure known to be involved in reinforcement and motivational control and is

innervated by dopaminergic populations that are thought to encode reinforcing properties of

food, including relaying information about sweet taste (Aso et al., 2009; Aso et al., 2014; Das et

al., 2016). Thus, our observation that mushroom body neurons are activated in hedonic food

environments is consistent with prior knowledge of their roles in motivation and reward

(Landayan et al., 2018).

To determine whether mushroom body neurons are involved in hedonic hunger beyond

correlative increases in activity, we tested for their requirement in hedonic and homeostatic

feeding. We used the genetically-encoded optogenetic tool GtACR to selectively inhibit

mushroom body neurons in our food choice assay. GtACR is a channelrhodopsin that is

activated upon green light exposure and increases chloride conductance, resulting in

hyperpolarization and silencing of neuronal populations (Mohammad et al., 2017). We

expressed GtACR in either the γ, α/β, or α'/β' lobes of the mushroom body (Aso et al., 2014)

and found that communication from the γ and α/β lobes is dispensable for both hedonic and

homeostatic feeding (Fig. 5C, 5D, Fig.5 – Fig. Supp. 1A, 1B). In contrast, signaling specifically

from the α'/β' lobes was required for hedonic, but not homeostatic, feeding (Fig. 5E, Fig. 5 – Fig.

Supp. 1C). The mushroom body horizonal lobes are innervated by dopaminergic neurons from

the protocerebral anterior medial (PAM) cluster, which were also required for hedonic feeding

(Fig. 5F) (Mao & Davis, 2009). These results indicate that a distinct PAM to α'/β' mushroom

body circuit regulates hedonic feeding, which is consistent with known descriptions of

anatomical and functional segregation of mushroom body lobes (Aso et al., 2014; Landayan et

al., 2018).

We next sought to determine whether communication from the α'/β' lobe was required for the

potentiating effects of homeostatic hunger on hedonic feeding. We reasoned that if the α'/β' lobe

also functions as a potentiator between the two hunger states, then inhibition of these neurons

would be expected to prevent increases in hedonic feeding when flies are starved. Instead, we

observed that starvation restored hedonic feeding in flies whose α'/β' lobes were inhibited (Fig.

5G), suggesting that potentiation of hedonic feeding by homeostatic state involves the

recruitment of additional and independent circuits for hedonic feeding.

Discussion

Animals exquisitely regulate feeding according to internal needs, but they are also subject to

their desires. Thus, food intake only sometimes occurs in response to homeostatic hunger and

can also occur in response to emotional stress, social environment, illness, and pursuit of

reward (Siemian et al., 2021; Xu et al., 2019). Because animals eat for many reasons other than

hunger, food intake alone may be a poor indicator of hunger states and may obscure

investigations into the motivational forces that drive behavior. Here, we observed the feeding

micro-behaviors of flies and developed a behavioral classification system to assist in the

identification of feeding drives. We established that flies exhibit behavioral signatures of two

hunger states, homeostatic and hedonic, and we used them to identify neurocircuits that are

required for hedonic, but not homeostatic, feeding. We also showed that homeostatic and

hedonic feeding are both independent and overlapping depending on environmental context,

and we developed a simple interaction framework that researchers can use to investigate these

relationships in detail. The development of this paradigm presents entry-points into the

discovery of new homeostatic and hedonic hunger circuits in flies, better classification of feeding

circuits which have already been described, and dissection of mechanistic relationships among

them.

The *Drosophila* MB supports a wide variety of behaviors and functions, in part due to its

complexity and heterogeneity in anatomical structure (Aso et al., 2009). The lobes of the MB are

formed by the interconnected axons of ~2000 Kenyon cells and 34 MB output neurons, and they

receive modulatory input from ~20 dopaminergic cell types (Aso et al., 2014). The MB lobes

have been anatomically and functionally divided into three classes: γ, α/β, and α'/β', which have

been further divided into functionally distinct compartments (Aso et al., 2014; Crittenden et al.,

1998; Tanaka et al., 2008). Our data suggest that compartments of the α'/β' MB lobes, but not

the γ or α/β lobes, are required for hedonic feeding. Although our experiments isolated a

discrete location within the functionally heterogenous mushroom body, the α'/β' lobe still

contains more than 350 Kenyon cells that are divided into middle (m) and anterior-posterior (ap)

compartments (Aso et al., 2014). Future investigations that utilize existing and emerging

intersectional technologies will be useful for discerning the specific cells within the α'/β' MB

lobes that promote hedonic feeding, as well as their modulatory inputs and outputs, and for

evaluation of the cell-intrinsic mechanisms that determine how hedonic hunger is encoded.

Connections from dopaminergic populations in the PAM cluster with MB-intrinsic Kenyon cells

have been implicated in appetitive learning, signaling reward in response to sugar, motivation to

overcome environmental stress, and promoting wakefulness (Burke et al., 2012; Haynes et al.,

2015; Hermanns et al., 2022; Liu et al., 2012). PAM projections to Kenyon cell axons in the β'

compartment of the MB respond to sugar by increasing calcium transients, which has led to the

idea that these synapses transmit information about sweet taste (Liu et al., 2012; May et al.,

2020). Interestingly, these responses are increased 2-fold by starvation, suggesting that they

scale in response to environmental context or degree of need (Liu et al., 2012), and artificial

activation of PAM neurons also increases foraging behavior (Tsao et al., 2018). Our finding that

the PAM>α'/β' circuit is required for promoting hedonic hunger builds upon a rich body of

literature, and together these observations offer a model that may unite the many described

functions of this node. One possibility is that this circuit acts as a permissive or motive force that

gates or fuels downstream mechanisms involved in behavioral responses to environmental

stimuli. Indeed, learning, reward, and wakefulness are all processes that might benefit from or

require a force that sustains them. This view offers an attractive entry into understanding the

seat of motivation in the fly brain.

Decades of research on the neural control of feeding have resulted in the identification of

mammalian neurocircuits involved in the control of feeding, and technological advances have

allowed researchers to manipulate them with an increasing degree of specificity. We are,

however, approaching the limits of what we can learn from manipulations and technologies of

these sorts and thus, complimenting them with discoveries from model systems in which more

specific manipulations can be tested may accelerate our ability to understand how the brain

generates motivated states. The hedonic circuits we have identified here in flies, for example,

can be immediately investigated at multiple levels to assess the neuronal and molecular

mechanisms that determine their role in feeding, including spatiotemporal manipulation of cell-

signaling pathways, electrophysiological recordings in live animals in response to various food

environments or energetic deficits, and optogenetic manipulation paired with close behavioral

monitoring. These types of experiments would fulfill several main goals in the study of the neural

control of appetite regulation (Ahn et al., 2022) and, despite differences in the precise identity of

feeding neurocircuits between flies and mammals, discoveries about foundational principles and

conserved signaling pathways may inform and direct mammalian studies.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
genetic reagent (*Drosophila melanogaster*)	*Canton-S*	Bloomington *Drosophila* Stock Center		

genetic reagent (*Drosophila melanogaster*)	*UAS-CaMPARI*	Bloomington *Drosophila* Stock Center	BDSC #68762 RRID:BDSC _68762	
genetic reagent (*Drosophila melanogaster*)	*MB238Y-GAL4*	Bloomington *Drosophila* Stock Center	BDSC #81009 RRID:BDSC _81009	
genetic reagent (*Drosophila melanogaster*)	*MB131B-GAL4*	Bloomington *Drosophila* Stock Center	BDSC #68265 RRID:BDSC _68265	
genetic reagent (*Drosophila melanogaster*)	*MB008B-GAL4*	Bloomington *Drosophila* Stock Center	BDSC #68291 RRID:BDSC _68291	
genetic reagent (*Drosophila melanogaster*)	*MB461B-GAL4*	Bloomington *Drosophila* Stock Center	BDSC #68327 RRID:BDSC _68327	
genetic reagent (*Drosophila melanogaster*)	*GMR58E02-GAL4*	Bloomington *Drosophila* Stock Center	BDSC #41347 RRID:BDSC _41347	
genetic reagent (*Drosophila melanogaster*)	*LexAop2-CsChrimson;UAS-CaMPARI2;GMR57C10-GAL4*	Bloomington *Drosophila* Stock Center	BDSC #81089 RRID:BDSC _81089	

genetic reagent (*Drosophila melanogaster*)	*UAS-GtACR*	Other		M. Dus (University of Michigan)
software, algorithm	RStudio	RStudio	*RRID:SCR_000432*	
software, algorithm	Fiji	ImageJ	RRID:SCR_002285	
software, algorithm	FLIC analysis R code	Flidea	RRID:SCR_018386	
software, algorithm	DTrack	Other		S. Pletcher (University of Michigan)
Other	FLIC *Drosophila Feeding Monitors*	Sable Systems	Model DFMV3	https://www.sablesys.com/products/classic-line/flic_drosophila_behavior_system/
Other	FLIR grasshopper camera	FLIR	GS3-U3-28S5C-C	#88-052 - GS3-U3-28S5C-C 2/3" FLIR Grasshopper®3 High Performance USB 3.0 Color Camera
Other	Fujinon varifocal lens	Fujinon	MFR #DV3.4x3.8SA-1	Fujinon 3 MP Varifocal Lens (3.8-13mm, 3.4x Zoom)

Fly stocks and husbandry

Fly stocks were maintained on a standard cornmeal-based larval growth medium and in a

controlled environment (21C, 60% humidity) with a 12:12hr light:dark cycle. We controlled the

developmental larval density by manually aliquoting 32 ul of collected eggs into individual

bottles containing 25-50mL of food at 25C. For FLIC experiments, following eclosion mixed sex

flies were kept on cornmeal medium at 18C until they were 7-14 days old. One day prior to the

experiments, flies were sorted by sex and transferred onto either SY10 medium (10% [w/v]

sucrose and 10% [w/v] yeast) or wet starved (1 Kimwipe with 2mL of H20) at 25C. Experiments

that used flies starved for 48 hours were sorted to starvation vials and kept at 25C two days

prior to the experiments. We used mated female and male flies that were between 7-14 days old

for all video characterization experiments and mated female flies for all subsequent

experiments, except where noted. The following stocks were used for experiments: *Canton-S,*

UAS-CaMPARI (BDSC #68762)*, MB238Y-GAL4* (BDSC #81009)*, MB131B-GAL4* (BDSC

#68265)*, MB008B-GAL4* (BDSC #68291)*, MB461B-GAL4* (BDSC #68327)*, GMR58E02-GAL4*

(BDSC #41347)*,* and *LexAop2-CsChrimson;UAS-CaMPARI2;GMR57C10-GAL4* (BDSC

#81089) were obtained from the Bloomington *Drosophila* Stock Center. *UAS-GtACR* was

provided by M. Dus (University of Michigan, MI).

Video Recording and Scoring

The video recording setup for initial behavioral characterizations (Fig. 1B, 1C, Supp. Table 1)

consisted of the Fly Liquid-Food Counter (FLIC), a custom single-well enclosure, and a light to

illuminate the recording area. A FLIR Grasshopper camera (GS3-U3-28S5C-C) fitted with a

Fujinon varifocal lens (MFR #DV3.4x3.8SA-1) was positioned against the lid to record the FLIC

well and the surrounding enclosed area. Video recording and FLIC data collection were started

simultaneously, and video length was dependent on the maximum number of frames that could

be stored. This setup recorded continuous video of the fly and saved this footage unaltered. For

experiments in Figures 1D-1F, we developed a direct communication link between our custom

DTrack video software and the FLIC system to trigger video capture only when there was

physical contact between a fly and the food source, together with 5s windows before and after

the contact (see Fig. 1A).

Videos of sated or 24-hour wet-starved male and female Canton-S flies were recorded. A single

fly was aspirated into the FLIC, which contained 5% or 10% sucrose food solution in 4 mg/l

MgCl$_2$. Videos were recorded at various times of day between 10AM and 7PM and had an

average length of about 51 minutes, except for videos collected for circadian analysis, which

were recorded over a 24-hour period. "Meal-times" were defined as 7AM-10AM and 7PM-10PM.

All videos were recorded on the lab bench. Manual, frame-by-frame annotation was used to

classify user-defined behaviors. Experimenters were blinded to the treatment for video scoring.

Data was analyzed using the open-source JWatcher software and R studio.

Fly Liquid-Food Interaction Counter (FLIC) assays

Flies were tested on the Fly Liquid-Food Interaction Counter (FLIC) system as previously

described (Ro et al., 2014) (Sable Systems International). FLIC *Drosophila* Feeding Monitors

(DFMs, Sable Systems International, model DFMV3) were used in the two-choice configuration

and each chamber was loaded with food solutions containing a sugar choice (1%-20% sucrose,

[18% arabinose + 2% sucrose], or [18% sorbitol + 2% sucrose]) and a yeast choice ([2% bacto

yeast extract + 1% sucrose]) in 4 mg/l MgCl$_2$. Flies were briefly anesthetized on ice and

aspirated into the DFM chambers, ensuring that each DFM received flies from all treatments.

We began recording immediately after loading flies (generally, loading all DFMs requires <10

minutes) and measured FLIC interactions for 6 hours during the "non-meal" time of day. Each

FLIC experiment contains pooled data from two independent replicates. FLIC data were

analyzed using custom R code, which is available at

https://github.com/PletcherLab/FLIC_R_Code. Default thresholds were used for analysis except

for the following: minimum feeding threshold = 10, tasting threshold = (0,10). Animals that did

not participate (i.e. returned zero values), whose DFM returned an unstable baseline signal, or

who produced extreme outliers (criteria for outliers were predetermined as exceeding twice the

mean of the population) were excluded from analysis.

Consumption-excretion feeding assays

Con-Ex experiments were carried out as previously described (Shell et al., 2021; Shell et al.,

2018). Experimental female flies were sorted to appropriate control (choice between SY10 food

or 2% sucrose) or hedonic (choice between SY10 or 20% sucrose) food environments (10 flies

per vial) when they were 4 days old and maintained on this food for 3 weeks. Food was

changed every 2-3 days. After the dietary pre-treatment period, blue test food was prepared in

removable choice caps by adding 1% (w/v) FD&C Blue No. 1 to only the side of the cap

containing the sucrose food choice. Flies were moved to fresh vials with the removable blue-

food caps on the top of the vials and were allowed to feed and excrete for 24 hours. Caps and

flies were removed after the 24-hour test period and flies were counted. Excreted dye was

collected by vortexing each vial with 3mL of water. Concentration of the dye was determined by

absorbance at 630nm and compared to a standard curve of known concentrations.

Optogenetic FLIC Assays

Flies expressing UAS-GtACR in desired neuronal populations were reared as described above.

When 7-14 days old, flies were sorted to cornmeal medium containing 200 μm all-*trans*-retinal

(from a stock solution of 100 mM ATR in 100% ethanol) and kept in the dark for 1-4 days at

18C. Flies were shifted to 25C for 24 hours before the experiment and flipped to either SY10

media containing 800 μm all-*trans*-retinal or starvation vials. Custom FLIC lids were fitted with

530 nm LEDs (Luxeon). Custom hardware and firmware were designed to allow the

experimenter to control LED intensity and a range of light stimulus parameters. We used a

stimulus frequency of 40 Hz and a pulse width of 800 ms for the entire duration of the 6 hour

FLIC experiment.

CaMPARI

CaMPARI photoconversion was carried out according to previously established protocols with some modifications (Edwards et al., 2020). A 405 nM LED (Thorlabs M405L3) was connected to a 1.2A LED driver (ThorLabs LEDD1B). Custom hardware and firmware were designed to allow the experimenter to control LED intensity and a range of light stimulus parameters. Female flies expressing *UAS-CaMPARI* were reared as described above. Custom choice caps were filled with fresh SY10% on one side and either 2% or 20% sucrose in 2% bacto agar on the other side. Groups of 5 flies were briefly anesthetized on ice and aspirated into the appropriate food environments. A custom clear cap was placed on top of the mocap to enclose the flies. The 405 nM LED was placed on top (~1/4 inch above the food surface) and the flies were allowed to habituate to the environment for 1 hour. After habituation, flies received UV light stimulation for 1 hour (cycling 10s on and 10s off, 1.0A). Immediately following light stimulation, flies were protected from light and brains were dissected as previously described (Wu & Luo, 2006). Brains were mounted between a glass microscope slide and a #1.5 cover glass separated by a custom bridge in VECTASHIELD Antifade Mounting Medium (Vector Laboratories). Samples were imaged on a Nikon A1 Confocal Microscope with a 20X air objective using 488 nm and 561 nm laser lines. Z-stacks were collected at a 1-2 μm step. All treatments were mounted under the same cover slip. Image processing was performed using ImageJ (NIH), and the experimenter was blinded to treatment. Slices containing the ROI were identified and collapsed into an average intensity projection. ROIs were measured in the red and green channel after background subtraction, and a ratio of red:green fluorescence is reported.

Statistics

For comparisons involving only one level, we used Student's t-test to detect significant differences between two treatments or one-way ANOVA followed by Tukey's post-hoc test after verifying normality and equality of variances. T-tests were two-tailed during initial

characterization experiments, or one-tailed in future experiments where the predicted direction

of change was known. For comparisons involving more than one level, we used two-way

ANOVA to detect significant interactions between the levels and followed up with Tukey's post-

hoc when significance was detected ($p<0.05$). In cases where experimental replicates were

pooled, a two-way ANOVA with blocking for experiment was performed to ensure non-

significant experimental effects. For all dot and bar plots, error bars represent the SEM. All

statistical tests and graphing were performed using R. Specific details of statistical analyses are

presented in the figure legends.

Acknowledgments: We wish to thank past and present members of the Pletcher laboratory for

their support and comments about the experimental design of these studies, and express

gratitude to David Paris for his help engineering and creating some tools used in this study. We

acknowledge Binyamin Jacobovitz in the Michigan Medicine Microscopy Core for training and

advice on confocal imaging. Figure 1A was created with BioRender.com. **Funding:** This

research was supported by The National Science Foundation Graduate Research Fellowship

Program (No. DGE 1256260) and the Howard Hughes Medical Institute through the James H.

Gilliam Fellowships for Advanced Study program to K.J.W (#GT11426) and the US National

Institute of Health, National Institute on Aging (R01 AG051649, R01 AG030593, and R01

AG063371) and the Glenn Medical Foundation to S.D.P.

Competing interests: The senior author (S.D.P) is a share holder in the company, Flidea,

which has developed technology related to the FLIC feeding system. **Data and materials**

availability: All data generated during this study are provided with the manuscript.

FLIC data were analyzed using custom R code, which is available at

https://github.com/PletcherLab/FLIC_R_Code.

Supplementary Files

Figure 5 – video 1 NsybCaMPARI stacks control environment green

Figure 5 – video 2 NsybCaMPARI stacks control environment red

Figure 5 – video 3 NsybCaMPARI stacks control environment merge

Figure 5 – video 4 NsybCaMPARI stacks hedonic environment green

Figure 5 – video 5 NsybCaMPARI stacks hedonic environment red

Figure 5 – video 6 NsybCaMPARI stacks hedonic environment merge

Figure 5 – video 7 MBCaMPARI stacks control environment green

Figure 5 – video 8 MBCaMPARI stacks control environment red

Figure 5 – video 9 MBCaMPARI stacks control environment merge

Figure 5 – video 10 MBCaMPARI stacks hedonic environment green

Figure 5 – video 11 MBCaMPARI stacks hedonic environment red

Figure 5 – video 12 MBCaMPARI stacks hedonic environment merge

References

Ahn, B. H., Kim, M., & Kim, S. Y. (2022). Brain circuits for promoting homeostatic and non-homeostatic appetites. *Exp Mol Med*, *54*(4), 349-357. https://doi.org/10.1038/s12276-022-00758-4

Alcantara, I. C., Tapia, A. P. M., Aponte, Y., & Krashes, M. J. (2022). Acts of appetite: neural circuits governing the appetitive, consummatory, and terminating phases of feeding. *Nat Metab*, *4*(7), 836-847. https://doi.org/10.1038/s42255-022-00611-y

Altherr, E., Rainwater, A., Kaviani, D., Tang, Q., & Guler, A. D. (2021). Long-term high fat diet consumption reversibly alters feeding behavior via a dopamine-associated mechanism in mice. *Behav Brain Res*, *414*, 113470. https://doi.org/10.1016/j.bbr.2021.113470

Andermann, M. L., & Lowell, B. B. (2017). Toward a Wiring Diagram Understanding of Appetite Control. *Neuron*, *95*(4), 757-778. https://doi.org/10.1016/j.neuron.2017.06.014

Arcego, D. M., Krolow, R., Lampert, C., Toniazzo, A. P., Garcia, E. D. S., Lazzaretti, C., Costa, G., Scorza, C., & Dalmaz, C. (2020). Chronic high-fat diet affects food-motivated behavior and hedonic systems in the nucleus accumbens of male rats. *Appetite*, *153*, 104739. https://doi.org/10.1016/j.appet.2020.104739

Aristotle, Thomson, J. A. K., Tredennick, H., & Aristotle. (2004). *The Nicomachean ethics* (Further rev. ed.). Penguin Books. Publisher description http://www.loc.gov/catdir/enhancements/fy0716/2004271883-d.html

Contributor biographical information
http://www.loc.gov/catdir/enhancements/fy0716/2004271883-b.html
Aso, Y., Grubel, K., Busch, S., Friedrich, A. B., Siwanowicz, I., & Tanimoto, H. (2009). The
mushroom body of adult Drosophila characterized by GAL4 drivers. *J Neurogenet, 23*(1-
2), 156-172. https://doi.org/10.1080/01677060802471718
Aso, Y., Hattori, D., Yu, Y., Johnston, R. M., Iyer, N. A., Ngo, T. T., Dionne, H., Abbott, L. F., Axel,
R., Tanimoto, H., & Rubin, G. M. (2014). The neuronal architecture of the mushroom
body provides a logic for associative learning. *Elife, 3*, e04577.
https://doi.org/10.7554/eLife.04577
Berridge, K. C. (1996). Food reward: brain substrates of wanting and liking. *Neurosci Biobehav*
Rev, 20(1), 1-25. https://doi.org/10.1016/0149-7634(95)00033-b
Berridge, K. C. (2009). 'Liking' and 'wanting' food rewards: brain substrates and roles in eating
disorders. *Physiol Behav, 97*(5), 537-550.
https://doi.org/10.1016/j.physbeh.2009.02.044
Betley, J. N., Xu, S., Cao, Z. F. H., Gong, R., Magnus, C. J., Yu, Y., & Sternson, S. M. (2015).
Neurons for hunger and thirst transmit a negative-valence teaching signal. *Nature,*
521(7551), 180-185. https://doi.org/10.1038/nature14416
Blumstein, D. T., & Daniel, J. C. (2007). *Quantifying behavior the JWatcher way*. Sinauer
Associates. Table of contents only
http://www.loc.gov/catdir/toc/ecip0713/2007011510.html
Burke, C. J., Huetteroth, W., Owald, D., Perisse, E., Krashes, M. J., Das, G., Gohl, D., Silies, M.,
Certel, S., & Waddell, S. (2012). Layered reward signalling through octopamine and
dopamine in Drosophila. *Nature, 492*(7429), 433-437.
https://doi.org/10.1038/nature11614
Burke, C. J., & Waddell, S. (2011). Remembering nutrient quality of sugar in Drosophila. *Curr*
Biol, 21(9), 746-750. https://doi.org/10.1016/j.cub.2011.03.032
Cannon, W. B., & Washburn, A. L. (1993). An explanation of hunger. 1911. *Obes Res, 1*(6), 494-
500. https://doi.org/10.1002/j.1550-8528.1993.tb00033.x
Casarrubea, M., Aiello, S., Di Giovanni, G., Santangelo, A., Palacino, M., & Crescimanno, G.
(2019). Combining Quantitative and Qualitative Data in the Study of Feeding Behavior in
Male Wistar Rats. *Front Psychol, 10*, 881. https://doi.org/10.3389/fpsyg.2019.00881
Casarrubea, M., Jonsson, G. K., Faulisi, F., Sorbera, F., Di Giovanni, G., Benigno, A., Crescimanno,
G., & Magnusson, M. S. (2015). T-pattern analysis for the study of temporal structure of
animal and human behavior: a comprehensive review. *J Neurosci Methods, 239*, 34-46.
https://doi.org/10.1016/j.jneumeth.2014.09.024
Chen, S., Lee, A. Y., Bowens, N. M., Huber, R., & Kravitz, E. A. (2002). Fighting fruit flies: a model
system for the study of aggression. *Proc Natl Acad Sci U S A, 99*(8), 5664-5668.
https://doi.org/10.1073/pnas.082102599
Crittenden, J. R., Skoulakis, E. M., Han, K. A., Kalderon, D., & Davis, R. L. (1998). Tripartite
mushroom body architecture revealed by antigenic markers. *Learn Mem, 5*(1-2), 38-51.
https://www.ncbi.nlm.nih.gov/pubmed/10454371
Das, G., Lin, S., & Waddell, S. (2016). Remembering Components of Food in Drosophila. *Front*
Integr Neurosci, 10, 4. https://doi.org/10.3389/fnint.2016.00004

Denis, R. G., Joly-Amado, A., Webber, E., Langlet, F., Schaeffer, M., Padilla, S. L., Cansell, C.,
Dehouck, B., Castel, J., Delbes, A. S., Martinez, S., Lacombe, A., Rouch, C., Kassis, N.,
Fehrentz, J. A., Martinez, J., Verdie, P., Hnasko, T. S., Palmiter, R. D., . . . Luquet, S.
(2015). Palatability Can Drive Feeding Independent of AgRP Neurons. *Cell Metab, 22*(4),
646-657. https://doi.org/10.1016/j.cmet.2015.07.011
Deshpande, S. A., Carvalho, G. B., Amador, A., Phillips, A. M., Hoxha, S., Lizotte, K. J., & Ja, W. W.
(2014). Quantifying Drosophila food intake: comparative analysis of current
methodology. *Nat Methods, 11*(5), 535-540. https://doi.org/10.1038/nmeth.2899
Duistermars, B. J., Pfeiffer, B. D., Hoopfer, E. D., & Anderson, D. J. (2018). A Brain Module for
Scalable Control of Complex, Multi-motor Threat Displays. *Neuron, 100*(6), 1474-1490
e1474. https://doi.org/10.1016/j.neuron.2018.10.027
Edwards, K. A., Hoppa, M. B., & Bosco, G. (2020). The Photoconvertible Fluorescent Probe,
CaMPARI, Labels Active Neurons in Freely-Moving Intact Adult Fruit Flies. *Front Neural*
Circuits, 14, 22. https://doi.org/10.3389/fncir.2020.00022
Flavell, S. W., Gogolla, N., Lovett-Barron, M., & Zelikowsky, M. (2022). The emergence and
influence of internal states. *Neuron, 110*(16), 2545-2570.
https://doi.org/10.1016/j.neuron.2022.04.030
Fosque, B. F., Sun, Y., Dana, H., Yang, C. T., Ohyama, T., Tadross, M. R., Patel, R., Zlatic, M., Kim,
D. S., Ahrens, M. B., Jayaraman, V., Looger, L. L., & Schreiter, E. R. (2015). Neural circuits.
Labeling of active neural circuits in vivo with designed calcium integrators. *Science,*
347(6223), 755-760. https://doi.org/10.1126/science.1260922
Fulton, S. (2010). Appetite and reward. *Front Neuroendocrinol, 31*(1), 85-103.
https://doi.org/10.1016/j.yfrne.2009.10.003
Ghiglieri, O., Gambarana, C., Scheggi, S., Tagliamonte, A., Willner, P., & De Montis, M. G. (1997).
Palatable food induces an appetitive behaviour in satiated rats which can be inhibited by
chronic stress. *Behav Pharmacol, 8*(6-7), 619-628. https://doi.org/10.1097/00008877-
199711000-00018
Gosby, A. K., Conigrave, A. D., Lau, N. S., Iglesias, M. A., Hall, R. M., Jebb, S. A., Brand-Miller, J.,
Caterson, I. D., Raubenheimer, D., & Simpson, S. J. (2011). Testing protein leverage in
lean humans: a randomised controlled experimental study. *PLoS One, 6*(10), e25929.
https://doi.org/10.1371/journal.pone.0025929
Hassett, C. C. (1948). The utilization of sugars and other substances by Drosophila. *Biol Bull,*
95(1), 114-123. https://www.ncbi.nlm.nih.gov/pubmed/18874957
Haynes, P. R., Christmann, B. L., & Griffith, L. C. (2015). A single pair of neurons links sleep to
memory consolidation in Drosophila melanogaster. *Elife, 4*.
https://doi.org/10.7554/eLife.03868
Hermanns, T., Graf-Boxhorn, S., Poeck, B., & Strauss, R. (2022). Octopamine mediates sugar
relief from a chronic-stress-induced depression-like state in Drosophila. *Curr Biol.*
https://doi.org/10.1016/j.cub.2022.07.016
Huang, J., Zhang, Z., Feng, W., Zhao, Y., Aldanondo, A., de Brito Sanchez, M. G., Paoli, M.,
Rolland, A., Li, Z., Nie, H., Lin, Y., Zhang, S., Giurfa, M., & Su, S. (2022). Food wanting is
mediated by transient activation of dopaminergic signaling in the honey bee brain.
Science, 376(6592), 508-512. https://doi.org/10.1126/science.abn9920

Huetteroth, W., Perisse, E., Lin, S., Klappenbach, M., Burke, C., & Waddell, S. (2015). Sweet
taste and nutrient value subdivide rewarding dopaminergic neurons in Drosophila. *Curr*
Biol, 25(6), 751-758. https://doi.org/10.1016/j.cub.2015.01.036
Itskov, P. M., Moreira, J. M., Vinnik, E., Lopes, G., Safarik, S., Dickinson, M. H., & Ribeiro, C.
(2014). Automated monitoring and quantitative analysis of feeding behaviour in
Drosophila. *Nat Commun, 5*, 4560. https://doi.org/10.1038/ncomms5560
Krashes, M. J., Koda, S., Ye, C., Rogan, S. C., Adams, A. C., Cusher, D. S., Maratos-Flier, E., Roth,
B. L., & Lowell, B. B. (2011). Rapid, reversible activation of AgRP neurons drives feeding
behavior in mice. *J Clin Invest, 121*(4), 1424-1428. https://doi.org/10.1172/JCI46229
Landayan, D., Feldman, D. S., & Wolf, F. W. (2018). Satiation state-dependent dopaminergic
control of foraging in Drosophila. *Sci Rep, 8*(1), 5777. https://doi.org/10.1038/s41598-
018-24217-1
Liu, C., Placais, P. Y., Yamagata, N., Pfeiffer, B. D., Aso, Y., Friedrich, A. B., Siwanowicz, I., Rubin,
G. M., Preat, T., & Tanimoto, H. (2012). A subset of dopamine neurons signals reward for
odour memory in Drosophila. *Nature, 488*(7412), 512-516.
https://doi.org/10.1038/nature11304
Liu, Q., Tabuchi, M., Liu, S., Kodama, L., Horiuchi, W., Daniels, J., Chiu, L., Baldoni, D., & Wu, M.
N. (2017). Branch-specific plasticity of a bifunctional dopamine circuit encodes protein
hunger. *Science, 356*(6337), 534-539. https://doi.org/10.1126/science.aal3245
Lowe, M. R., & Butryn, M. L. (2007). Hedonic hunger: a new dimension of appetite? *Physiol*
Behav, 91(4), 432-439. https://doi.org/10.1016/j.physbeh.2007.04.006
Mao, Z., & Davis, R. L. (2009). Eight different types of dopaminergic neurons innervate the
Drosophila mushroom body neuropil: anatomical and physiological heterogeneity. *Front*
Neural Circuits, 3, 5. https://doi.org/10.3389/neuro.04.005.2009
May, C. E., Rosander, J., Gottfried, J., Dennis, E., & Dus, M. (2020). Dietary sugar inhibits
satiation by decreasing the central processing of sweet taste. *Elife, 9*.
https://doi.org/10.7554/eLife.54530
May, C. E., Vaziri, A., Lin, Y. Q., Grushko, O., Khabiri, M., Wang, Q. P., Holme, K. J., Pletcher, S.
D., Freddolino, P. L., Neely, G. G., & Dus, M. (2019). High Dietary Sugar Reshapes Sweet
Taste to Promote Feeding Behavior in Drosophila melanogaster. *Cell Rep, 27*(6), 1675-
1685 e1677. https://doi.org/10.1016/j.celrep.2019.04.027
Mohammad, F., Stewart, J. C., Ott, S., Chlebikova, K., Chua, J. Y., Koh, T. W., Ho, J., & Claridge-
Chang, A. (2017). Optogenetic inhibition of behavior with anion channelrhodopsins. *Nat*
Methods, 14(3), 271-274. https://doi.org/10.1038/nmeth.4148
Ning, J., Li, Z., Zhang, X., Wang, J., Chen, D., Liu, Q., & Sun, Y. (2022). Behavioral signatures of
structured feature detection during courtship in Drosophila. *Curr Biol, 32*(6), 1211-1231
e1217. https://doi.org/10.1016/j.cub.2022.01.024
Pecina, S., Smith, K. S., & Berridge, K. C. (2006). Hedonic hot spots in the brain. *Neuroscientist,*
12(6), 500-511. https://doi.org/10.1177/1073858406293154
Ro, J., Harvanek, Z. M., & Pletcher, S. D. (2014). FLIC: high-throughput, continuous analysis of
feeding behaviors in Drosophila. *PLoS One, 9*(6), e101107.
https://doi.org/10.1371/journal.pone.0101107

Ro, J., Pak, G., Malec, P. A., Lyu, Y., Allison, D. B., Kennedy, R. T., & Pletcher, S. D. (2016).
Serotonin signaling mediates protein valuation and aging. *Elife*, *5*.
https://doi.org/10.7554/eLife.16843
Rossi, M. A., & Stuber, G. D. (2018). Overlapping Brain Circuits for Homeostatic and Hedonic
Feeding. *Cell Metab*, *27*(1), 42-56. https://doi.org/10.1016/j.cmet.2017.09.021
Saper, C. B., Chou, T. C., & Elmquist, J. K. (2002). The need to feed: homeostatic and hedonic
control of eating. *Neuron*, *36*(2), 199-211. https://doi.org/10.1016/s0896-
6273(02)00969-8
Shell, B. C., Luo, Y., Pletcher, S., & Grotewiel, M. (2021). Expansion and application of dye
tracers for measuring solid food intake and food preference in Drosophila. *Sci Rep*,
11(1), 20044. https://doi.org/10.1038/s41598-021-99483-7
Shell, B. C., Schmitt, R. E., Lee, K. M., Johnson, J. C., Chung, B. Y., Pletcher, S. D., & Grotewiel, M.
(2018). Measurement of solid food intake in Drosophila via consumption-excretion of a
dye tracer. *Sci Rep*, *8*(1), 11536. https://doi.org/10.1038/s41598-018-29813-9
Siemian, J. N., Arenivar, M. A., Sarsfield, S., & Aponte, Y. (2021). Hypothalamic control of
interoceptive hunger. *Curr Biol*, *31*(17), 3797-3809 e3795.
https://doi.org/10.1016/j.cub.2021.06.048
Simpson, S. J., Batley, R., & Raubenheimer, D. (2003). Geometric analysis of macronutrient
intake in humans: the power of protein? *Appetite*, *41*(2), 123-140.
https://doi.org/10.1016/s0195-6663(03)00049-7
Slooten, E. (1994). Behavior of Hector's Dolphin: Classifying Behavior by Sequence Analysis.
Journal of Mammalogy, *75*(4), 956-964. https://doi.org/10.2307/1382477
Sternson, S. M., & Eiselt, A. K. (2017). Three Pillars for the Neural Control of Appetite. *Annu Rev
Physiol*, *79*, 401-423. https://doi.org/10.1146/annurev-physiol-021115-104948
Tanaka, N. K., Tanimoto, H., & Ito, K. (2008). Neuronal assemblies of the Drosophila mushroom
body. *J Comp Neurol*, *508*(5), 711-755. https://doi.org/10.1002/cne.21692
Tsao, C. H., Chen, C. C., Lin, C. H., Yang, H. Y., & Lin, S. (2018). Drosophila mushroom bodies
integrate hunger and satiety signals to control innate food-seeking behavior. *Elife*, *7*.
https://doi.org/10.7554/eLife.35264
Weaver, K. J., Holt, R. A., Henry, E., & Pletcher, S. D. (2022). Encoding of hunger by the neuronal
epigenome slows aging in Drosophila [preprint]. *Biorxiv*. https://doi.org/doi:
https://doi.org/10.1101/2022.07.21.501022
Willesen, M. G., Kristensen, P., & Romer, J. (1999). Co-localization of growth hormone
secretagogue receptor and NPY mRNA in the arcuate nucleus of the rat.
Neuroendocrinology, *70*(5), 306-316. https://doi.org/10.1159/000054491
Wu, J. S., & Luo, L. (2006). A protocol for dissecting Drosophila melanogaster brains for live
imaging or immunostaining. *Nat Protoc*, *1*(4), 2110-2115.
https://doi.org/10.1038/nprot.2006.336
Xu, Y., Lu, Y., Cassidy, R. M., Mangieri, L. R., Zhu, C., Huang, X., Jiang, Z., Justice, N. J., Xu, Y.,
Arenkiel, B. R., & Tong, Q. (2019). Identification of a neurocircuit underlying regulation
of feeding by stress-related emotional responses. *Nat Commun*, *10*(1), 3446.
https://doi.org/10.1038/s41467-019-11399-z

Yang, M. Y., Armstrong, J. D., Vilinsky, I., Strausfeld, N. J., & Kaiser, K. (1995). Subdivision of the
820   Drosophila mushroom bodies by enhancer-trap expression patterns. *Neuron*, *15*(1), 45-
821   54. https://doi.org/10.1016/0896-6273(95)90063-2
822   Zigman, J. M., Jones, J. E., Lee, C. E., Saper, C. B., & Elmquist, J. K. (2006). Expression of ghrelin
823   receptor mRNA in the rat and the mouse brain. *J Comp Neurol*, *494*(3), 528-548.
824   https://doi.org/10.1002/cne.20823
825

827
Figure 1. Characterization of feeding micro-behaviors in flies. (A) Cartoon schematic of setup
for filming flies on the FLIC. Frame capture is triggered by interactions of flies with the food source.
(B) Frequency of all five main categories of observed feeding micro-behaviors: other (O), interaction
(I), fast (F), long (L), long/quick (LQ). Each data point represents the total number of times an event
occurred normalized to all observed events for each fly. N=8 flies, 528 events. **(C)** Percentage of
flies from (B) that engaged in each feeding micro-behavior. (**D**) Representative temporal plots of
sated (N=4 flies, 171 events) and starved (N=4 flies, 357 events) feeding micro-behaviors (left panel)
and frequency of each behavior (right panel, two-tailed t-tests). Each row in left panel represents one
fly, and frequencies are relative to all behaviors (two-tailed t-tests). **(E)** Heat maps of starved (left)
and sated (right) transitional probabilities of observed behaviors from (D) (transitional probabilities
are generated by dividing the observed count for each event pair by the total occurrences of the
given event. P-values are determined based on Z-scores, as described in (Blumstein & Daniel,
2007)). Long events contain both L and LQ events. **(F)** Frequency of each feeding micro-behavior
during meal (7AM/PM-10AM/PM) vs. non-meal times of day (one-tailed t-tests). N=9 flies, 693
events, frequency is relative to total interactions + feeding events.
843
844

845
846 **Figure 1 – Figure Supplement 1. Flies exhibit distinct morning and evening feeding**
847 **peaks. (A)** Representative temporal plot of food interactions on 2% sucrose FLIC from female
848 *Canton-S* flies monitored over 24 hrs. Each data point represents the binned average number of
849 events from a population of 38 flies in the previous 30-minute interval. Shaded regions indicate
850 the times of day designated as "meal-times".
851

852
Figure 2. FLIC signal characteristics distinguish between event-types. (A-C)
Representative FLIC signal generated during a visually identified interaction (I), fast feeding (F),
or long feeding (L/LQ) event. **(D-I)** Signal characteristics generated by the FLIC during visually
identified I, F, or L/LQ events. Each data point represents one event from one fly (one-way
ANOVA with Tukey's post-hoc).
858

859
Figure 3. Feeding event durations are increased by hedonic food environments. (A) Cartoon
schematic of event number and event duration (top). Cartoon schematic of hedonic and control food
choice environments, and comparisons used to define metrics for homeostatic and hedonic feeding
(bottom). **(B)** Total number of events with sucrose or yeast in the control food choice environment
from sated or 24-hr starved female *Canton-S* flies (two-way ANOVA with Tukey's post-hoc). **(C)**
Total number of events with sucrose or yeast in the control food choice environment from sated
female or male *Canton-S* flies (two-way ANOVA with Tukey's post-hoc). **(D-E)** Median event
durations with sucrose (D) or yeast (E) in the control vs. hedonic food choice environments from
sated female and male *Canton-S* flies (two-way ANOVA). **(F-G)** Total number of events with sucrose
(F) or yeast (G) in the control vs. hedonic food choice environments from sated female and male
Canton-S flies (two-way ANOVA). **(H-I)** Median event durations (H) and total number of events (I)
with sugar in the indicated food choice environments from sated female *Canton-S* flies (one-way
ANOVA with Tukey's post-hoc). **(J-K)** Median event durations (J) and total number of events (K) with
sugar in the indicated food choice environment from sated female *Canton-S* flies (one-way ANOVA
with Tukey's post-hoc). **(L-M)** Median event durations (L) and total number of events (M) with
sucrose in the indicated food choice environments (one-way ANOVA).
876

877
Figure 3 – Figure Supplement 1. Hedonic food environments do not elicit increases in
total number of events or event durations with yeast but do promote increased
volumetric intake. (A) Con-Ex measurement of volumetric sucrose consumptions (left panel)
and total mass consumed (right panel) in the control vs. hedonic food environments from female
Canton-S flies. (one-way ANOVAs). **(B)** Total number of events with sugar or yeast in indicated
food choice environments from sated female *Canton-S* flies (two-way ANOVA). **(C)** Median
event durations with yeast in the indicated food choice environments from sated female *Canton-*
S flies (one-way ANOVA).
886

Figure 4. Hedonic feeding is modulated by homeostatic hunger. (A) Total number of events with sucrose or yeast in the control food choice environment from sated or 24-hr starved female *Canton-S* flies (two-way ANOVA with Tukey's post-hoc). **(B)** Total number of events with sucrose or yeast in the control food choice environment from 24-hr vs. 48-hr starved female *Canton-S* flies (two-way ANOVA with Tukey's post-hoc). **(C)** Median event durations with sucrose in the control vs. hedonic food environments from sated vs. 24-hr starved female *Canton-S* flies (two-way ANOVA). **(D)** Linear regression of interactions with yeast (licks) versus mean event durations with indicated sugar. Each data point represents one fly (yeast licks are expressed as a transformation to the 0.25 power and event durations are expressed as a log transformation to achieve normality, call = lm(SucroseDurations~YeastLicks*FoodEnvironment), two-way ANOVA, adjusted R^2=0.1835).

Figure 5. Hedonic hunger requires distinct mushroom body lobes. (A) Representative maximum intensity projections from brains of flies expressing *Nsyb-GAL4>UAS-CaMPARI* exposed to control or hedonic food environments, quantified at right (scale bar = 100 uM, two-tailed t-test). **(B)** Representative maximum intensity projections from brains of flies expressing *MB238Y-GAL4>UAS-CaMPARI*, quantified at right (scale bar = 10 uM, one-tailed t-test). For both (A) and (B), green represents unconverted CaMPARI and magenta represents converted CaMPARI. **(C-E)** Median event durations with sucrose in control vs. hedonic food environments from sated female flies expressing *UAS-GtACR* driven by either (C) *MB131B-GAL4* (γ(d);γ(m))*,* (D) *MB008B-GAL4* (α/β(c);α/β(p);α/β(s)), or (E) *MB461B-GAL4* (α'/β'(ap);α'β'(m)) (two-way ANOVAs with Tukey's post-hoc). **(F)** Median event durations with sucrose in control vs. hedonic food environments from sated female flies expressing *PAM-GAL4>UAS-GtACR* (two-way ANOVA with Tukey's post-hoc). **(G)** Median event durations with sucrose in control vs. hedonic food environments from 24-hr starved female flies expressing *MB461B-GAL4>UAS-GtACR* (two-way ANOVA).

915
916 **Figure 5 – Figure Supplement 1. Mushroom body lobes are not required for homeostatic**
917 **hunger. (A-C)** Total number of events with sucrose or yeast in the control food choice
918 environment from 24-hr starved female flies expressing *UAS-GtACR* driven by either (A)
919 *MB131B-GAL4* (γ(d);γ(m))*, (B) *MB008B-GAL4* (α/β(c);α/β(p);α/β(s)), or (C) *MB461B-GAL4*
920 (α'/β'(ap);α'β'(m)) (two-way ANOVAs).
921

922   **Figure 5 – Videos 1-12. Movies of confocal stacks, related to Figure 5A-B.** Movies of
923   confocal z-stacks through whole brains of flies carrying *Nsyb-Gal4>UAS-CaMPARI* (videos 1-6)
924   or *MB238Y-GAL4>UAS-CaMPARI* (videos 7-12) in control or hedonic food environments.
925   Green channels represent unconverted CaMPARI and magenta channels represent
926   photoconverted CaMPARI.
927

928 **Table 1. Library of feeding micro-behaviors**
929

Main Event Type	Code	Behavior Description
Other (O)	G	Groom
	UC	Unknown contact
Interaction (I)	IF	Front legs touch food
	IH	Hind legs touch food
Fast feeding (F)	F	Single movement of proboscis into food to feed for 1-3 seconds
	FI	Meets requirements for F and I
Long feeding (L)	L	Proboscis continuously in food while feeding for > 4 seconds
	LI	Meets requirements for L and I
Long/quick feeding (L)	LQ	Proboscis moves in and out quickly while feeding for > 4 seconds