Defining transcription factor nucleosome binding with Pioneer-seq

Maria Tsompana[1], Patrick Wilson[1], Vijaya Murugaiyan[1], Christopher R. Handelmann[1,2], Michael J. Buck[1,2]

[1]Department of Biochemistry, Jacobs School of Medicine and Biomedical Sciences, State University of New York at Buffalo, Buffalo, NY, United States

[2]Department of Biomedical Informatics, Jacobs School of Medicine and Biomedical Sciences, State University of New York at Buffalo, Buffalo, NY, United States

ABSTRACT

Gene expression requires the targeting of transcription factors (TFs) to regulatory sequences often occluded within nucleosomes. To comprehensively examine TF nucleosome binding we developed Pioneer-Seq. In Pioneer-seq a library of nucleosomes containing thousands of DNA sequences with TF binding sites in all possible nucleosome orientations is bound to a TF and the protein-nucleosome complex is isolated and quantified. To demonstrate Pioneer-seq we examined nucleosome binding by OCT4, SOX2, KLF4, and c-MYC. Our results demonstrate that KLF4 and SOX2 can bind close to the nucleosome dyad with nucleosome sequence being the major factor regulating TF binding across all studied TFs.

Introduction

The interactions between proteins and chromosomal DNA underlie basic nuclear processes, such as transcriptional regulation, DNA replication, repair, and recombination, chromosome segregation, and epigenetic inheritance, as well as many fundamental biological responses, including cell growth, division, and differentiation, embryonic development, environmental stress responses, apoptosis, and disease state development. According to the human protein atlas, 8,887 proteins localize to the nucleus, with an estimated 1,500 DNA-binding transcription factors (TFs)[1]. Most TFs preferentially bind nucleosome-free DNA, which appears to be a conundrum during development because many gene regulatory regions are in nucleosomal DNA. However, there are a few TFs that belong to a specific class, known as pioneer factors, that can bind to closed chromatin and open nucleosomal domains [2-6].

Pioneer factors are not able to bind all their targets throughout the genome, indicating that there are some constraints. There is evidence that the location of the TF binding site (TFBS) within a nucleosome (known as the translational setting) is one determinant[6]. For example, binding of the glucocorticoid receptor at its TFBS near the nucleosome edge is 4-fold greater than at an identical site positioned 20 bp from the dyad[7]. The translational setting can inhibit TF binding 2- to 100-fold[8-11]. TF binding is also influenced by the orientation of a TFBS on a nucleosome (known as the rotational setting) that results from the twist of the helical DNA structure. FoxA binds to its well-defined target site in the *Alb* enhancer (5–15 bp from the nucleosome dyad in liver cells)[12] only at specific rotational settings[13]. The way in which TFs recognize their binding sites (e.g., partial motif recognition) also likely influences their ability to bind nucleosomal DNA[14].

Until recently, the only way to determine if and how a TF binds its target within nucleosomal DNA was by a low-throughput molecular biology assay, in which *in vitro*-formed nucleosomes with a single targeted DNA sequence were bound by a regulatory protein at

increasing concentrations[15]. To examine various TFBSs or to define their optimal translational or rotational settings within the nucleosome, this assay had to be repeated for each template DNA. An approach called NCAP-SELEX (nucleosome consecutive affinity purification–systematic evolution of ligands by exponential enrichment) was recently developed that uses randomized nucleosomal DNA libraries with TF-nucleosome enrichment followed by next-generation sequencing[16]. However, NCAP-SELEX is limited to short nucleosome fragments and cannot be used to directly compare binding to different locations in the nucleosome or to linker DNA. To compare TFBSs at various nucleosomal positions, competitive nucleosome binding assays and Senu-seq combine electrophoretic mobility shift assays (EMSAs) with a nucleosome library and next-generation sequencing[4, 5, 17], but these approaches are limited in the number of sequences that can be examined and are restricted to nucleosome positioning sequences (NPSs). To comprehensively examine TF-nucleosome binding across various NPSs and TFBSs and in *in vivo*-targeted nucleosomes (ITNs), we developed Pioneer-seq (transcription factor nucleosome binding protocol), which can be used to examine 7,500 nucleosomes in a single assay.

Results

Pioneer-seq workflow

Pioneer-seq expands on competitive nucleosome binding assays by including thousands of sequences and includes additional NPSs with ITNs. Pioneer-seq uses 5S rDNA [18], MMTV LTR [19], and the artificial synthetic sequence Widom 601 DNA as NPSs [20]. Use of these highly characterized NPSs is advantageous due to the availability of their defined structures and dynamics [15]. For each TF to be studied, a TFBS is added to the NPS at a 1-bp resolution across each NPS with additional sites outside the core 147-bp nucleosome in the linker regions

(multiple TFBSs can be studied in a single assay). Each NPS with the inserted TFBS is flanked by PCR primers (19–20 bp) for a total sequence length of 230 bp.

In addition to highly characterized NPSs, Pioneer-seq can also include genomic locations targeted *in vivo* by a TF of interest. These ITNs are defined by examining nucleosome positioning data from MNase-seq or NOMe-seq with TF binding data from ChIP-seq, ChIP-nexus, or Cut & Tag. The 191-bp sequences centered at the ITN are then filtered for predicted nucleosome formation ability[21] and the presence of the specific TFBS. In total, 7,500 sequences are designed for each library such that most sequences are not specific for any particular TF. All DNA sequences are synthesized as an Agilent 230-bp oligonucleotide library. Nucleosomes are then assembled with salt gradient dialysis using all nucleosome sequences simultaneously; free DNA is removed from the nucleotide library by a sucrose gradient.

The purified nucleosome library is used in binding assays in which the TF of interest is added at increasing concentrations. TF-nucleosome complexes are detected after a short incubation by separating the reaction mixtures on a native polyacrylamide gel (the first lane of which contains only nucleosomes to measure the background and input levels for each experimental replicate). Nucleosomes that are bound by the TF are identified by sequencing the DNA that is extracted and purified from shifted bands in the gel. The sequencing results are then analyzed and mapped back to the original 7,500-nucleosome library. Figure 1 illustrates the Pioneer-seq workflow.

TF-nucleosome binding is then determined by calculating the "relative supershift," which compares a nucleosome sequence to a collection of non-specific sequences in the same binding experiment. By comparing each nucleosome sequence to the non-specific control sequences from the same lane, loading, PCR amplification, next-generation sequencing, and binding to additional unknown motifs are all internally controlled.

Nucleosome formation and positioning for the Pioneer-seq library is then determined by MNase-seq [22, 23] . For these experiments, MNase-digested DNA from the nucleosome library

Fig 1. Overview of Pioneer-seq. With this method, 230-bp DNA sequences of interest are designed in batches of 7,500 sequences, including well-characterized nucleosome positioning sequences (NPS; Widom 601 or 5S rDNA) and sequences for *in vivo* targeted nucleosomes (ITNs) (left). The method enables testing of binding site variation and positioning, flanking sequence content, and combinatorial binding events. Nucleosomes are formed and purified on all 7,500 sequences to generate a nucleosome library (middle). The entire nucleosome library is incubated with increasing amounts of a transcription factor (TF) of interest (bottom right). The TF-nucleosome complexes are separated by electrophoretic mobility shift assay (EMSA), and the bound and unbound DNA are recovered, quantified, and sequenced. Nucleosome positioning and accessibility for every DNA sequence in the nucleosome library are determined by digestion with micrococcal nuclease (MNase) (top right). The resulting DNA fragments are sequenced and mapped back to the initial library. TFBS, TF binding site.

is sequenced and mapped back to the 7,500-nucleosome library to determine which regions are protected; results from well-defined NPS are consistent with known nucleosome structures (Extended Data Fig. 1).

Defining TF-nucleosome binding

To evaluate Pioneer-seq, we examined nucleosome binding for the Yamanaka factors OCT4, SOX2, KLF4, and c-MYC [24]. OCT4, SOX2 and KLF4 have been shown to act as pioneer factors that directly bind to chromatin regions inaccessible to other TFs and subsequently trigger transcriptional competency by directing chromatin remodeling [25, 26]. Despite their pioneering capabilities OCT4, SOX2 and KLF4 cannot bind to all their TFBS within a genome [27].

The strongest binding occurs for all factors at TFBS located outside of the nucleosome region within the linkers (Extended data Fig. 2). Within nucleosomes there is extensive binding variability depending on the NPS and TFBS. The Widom 601 NPS is the most extensively studied NPS and 601-nucleosomes has been a model for studying nucleosome structures and dynamics [20] and in our experiments, 601 nucleosomes are the most efficiently formed nucleosomes (Extended Data Fig. 3). The binding of OCT4, SOX2, KLF4, and MYC to the 601 nucleosomes was inhibited when their TFBSs were near the nucleosome dyad (Fig. 2), similarly to that observed for TP53 and TP63 in previous studies[4, 5]. For KLF4 there were two TFBSs (with corresponding reverse complement sequences): Klf4-1 (CCCCACCC) is derived from the motif MA0039.4 from Jaspar[28] and Klf4-2 (GCCCCGCCCCGCCCC) is derived from the KLF4 long motif discovered in mouse embryonic stem cells[29]. KLF4 bound strongly to Klf4-1 only when the site was positioned >55 bp from the 601 dyad. Results for binding to Klf4-2 were noisier at some internal nucleosome positions because Klf4-2 appears to disrupts nucleosome formation. The results for the reverse complement sequences (Klf4-1RC and Klf4-2RC) are similar to those for the direct sequences. For OCT4, there is an OCT4 TFBS (TATGCAAAT) and a joint Oct4-Sox2 TFBS (CTTTGTTATGCAAAT). Binding of OCT4 to both TFBS is very similar, as the OCT4 target sequence is the same. However, there were small differences in their relative supershifts that resulted from differences in the center positioning of the TFBSs. For SOX2, there is a joint Oct4-Sox2 TFBS and a SOX2 TFBS (ACAATGG). SOX2 binds both sequences in the linkers and can bind the Oct4-Sox2 TFBS at the nucleosome edge. The non-pioneer factor MYC was also examined with a single palindromic TFBS (ACCACGTGGT) derived from MA0059.1. MYC can bind its TFBS in the linker and when its TFBS is located > 55bp from the dyad.

With Pioneer-seq, we can compare the binding of TFs across different NPS in a single assay. We compared binding to nucleosomes of 601 NPS, 5S rDNA, and the MMTV long

Fig 2. TF binding to the Widom 601 nucleosome. The TFBS is positioned across all possible locations along the 601 nucleosome with TFBSs in the left and right linkers to generate a total of 149 unique nucleosomes per TFBS. The relative supershift for each nucleosome is determined by counting the frequency of each sequence within the shifted band in the electrophoretic mobility shift assay and comparing it to that for nonspecific binding (i.e., binding to a sequence without the TFBS for that particular TF). This value is then normalized to the input ratio of nucleosomes (see Eq. 1). Shading around each line is SEM. Binding for KLF4 (**a**), SOX2 (**b**), OCT4 (**c**), and MYC (**d**) is shown for nucleosomes with their specific TFBSs along with binding to a nonspecific TFBS nucleosome sequence (shown in gray). Breaks in the trace for Klf4-2 TFBS indicate missing data as a result of inefficient nucleosome formation. RC, reverse complement.

terminal repeat. The binding of KLF4, OCT4, SOX2, and MYC was closer to the dyad for 5S nucleosomes than for 601 nucleosomes, with binding inhibited ±30 bp from the dyad. In addition, binding was asymmetric, with increased binding on the left side of the 5S nucleosome. Binding to the MMTV nucleosome was similar to that for the 5S nucleosome, with limited binding within 30 bp of the dyad. However, asymmetric binding occurred on the right face of the MMTV nucleosome. All TFs tested had greater binding to TFBSs in 5S and MMTV nucleosomes than to those in the 601 nucleosomes.

Rotational and translational settings of nucleosome sequences drive binding differences

Because all 7,500 nucleosome sequences are exposed simultaneously to TFs in a single experiment, the binding to different NPSs or TFBSs can be directly compared. We found that in general, binding to TFBSs within the 601 nucleosome was strongly inhibited, whereas binding increased as TFBSs approached the edges of MMTV and 5S nucleosomes. As shown in Fig.

Fig 3. TF binding to 5S and MMTV nucleosomes. The TFBSs were positioned across all possible locations along the 5S (**a**–**d**) and MMTV (**e**–**h**) nucleosomes, with TFBSs in the left and right linkers. Relative supershifts for KLF4 (**a**, **e**), SOX2 (**b**, **f**), OCT4 (**c**, **g**), and MYC (**d**, **h**) are shown for their specific TFBSs along with binding to a nonspecific TFBS nucleosome sequence (shown in gray). Breaks in the trace for Klf4-2 TFBS in panels a and e indicate missing data as a result of inefficient nucleosome formation.

4a, SOX2 binding to the OCT4-SOX2 TFBS differed depending on the NPS. SOX2 bound strongly to the left face of the 5S nucleosome, with periodic binding at -30 and -40 sites. SOX2 has a high-mobility group (HMG) box domain, which binds in the minor groove of DNA and induces helical bending[30]. The pre-bent conformation of nucleosomal DNA may thus facilitate the binding of SOX2 and other HMG domain-containing proteins[31].

In addition to the KLF4 TFBSs in the library, KLF4 can bind to the TFBSs for p53. At these sequences, KLF4 can bind at the linkers and at specific rotational and translational settings near the nucleosome dyad. Its preferred TFBS is unbound when positioned at the same internal locations. The p53-1 and p53-2 TFBS are both bound by KLF4 in the linker regions but not as strong as the KLF4 TFBS (Extended Data Fig. 2). On the other hand, when

positioned near Super-Helix Location (SHL) -2 (-20 bp from dyad) within 601 both of the p53

motifs are bound (Fig 4b, Extended Data Fig 4). To confirm KLF4 binding to the p53-1

(GGGCATGTCCGGGCATGTCC) site on 601, we performed validation assays with DNaseI

footprinting. In this experiment, the single nucleosome containing the p53-1 TFBS at SHL-2 is

treated with DNaseI before and after KLF4 binding (Fig 4c). Changes in protection are

distinguishable at the p53-1 TFBS. KLF4 appears to be binding at two GGGC sequences

spaced 10bp apart in the p53-1 TFBS. The two GGGC sequences are partially exposed in

neighboring major grooves (Fig 4d). The DNaseI footprinting also showed an additional

footprint located at SHL 5.5. This region shown in red in Fig 5d is directly next to the p53-1

TFBS located on the other DNA gyre, suggesting that KLF4 binding to p53-1 at SHL -2 disrupts

the region near the entry/exit on the opposite side.

Examining ITNs

In addition to well-characterized NPSs, ITNs at gene regulatory regions were included in the

nucleosome library. These nucleosomes were identified using ChIP-seq binding data from

induction of OCT4, SOX2, KLF4, and MYC in fibroblast during IPSC generation [27].

Nucleosome positions were defined by NOMe-seq from the fibroblast cell line (IMR90) [32].

After filtering nucleosomes by predicted nucleosome formation efficiency, 372 nucleosomes

were designed. In total, ~20% of the ITNs failed to make stable nucleosomes for our assay.

The remaining ITNs had a formation efficiency similar to 5S and MMTV nucleosomes (Extended

Data Fig. 3).

The Lin28B nucleosome is a ITN that has been examined by multiple groups [14]. We

defined the nucleosome-protected region for Lin28B with MNase-seq and show a protected

region centered at 150 bp with an OCT4 TFBS at position 38-46 bp. Pioneer-seq results show

Fig. 4. Nucleosome binding depends on nucleosome positioning sequence and binding site. a, Relative supershifts for SOX2 binding to the Oct4-Sox2 TFBS (CTTTGTTATGCAAAT) within the 5S, MMTV, and 601 nucleosomes. **B,** Relative supershifts for KLF4 binding to Klf4-1 (CCCCACCC), Klf4-1RC (GGGTGGGG), and p53-1 (GGGCATGTCCGGGCATGTCC) within the 601 nucleosome; "*" indicates the nucleosome validated in DNase I footprinting experiments. RC, reverse complement. **c,** DNase I footprinting of nucleosomes containing the p53-1 TFBS at SHL -2; an additional footprint observed on the neighboring gyre is indicated with the dashed-line box. Nuc, DNase I digestion. **d,** Model of the 601 nucleosome with the p53-1 motif highlighted in yellow, the two GGGC in blue, and the additional footprinted region in red.

that only OCT4 can bind specifically to the Lin28B nucleosome. These results are consistent

with DNase-footprinting and binding specificity assays for KLF4, MYC, OCT4, and SOX2 [14].

The previously characterized ITNs located at ALBN1, NRCAM, CX3CR1, and ESRRB were also

included within this library (Extended Data Fig. 5). Other ITNs that were bound by OCT4, SOX2,

KLF4, and MYC are shown in Fig. 5b–e.

By examining the Oct4-Sox2 sites we can directly compare OCT4 and SOX2 binding at

their TFBS when they are at the same nucleosome positions. At most Oct4-Sox2 TFBSs SOX2

can bind while OCT4 only binds when the TFBS is outside the protected nucleosomal region

(Fig 5 b,d,e; Extended Data Fig 6). To compare binding between factors at ITNs we have

Fig 5. Binding to *in vivo*-targeted nucleosomes. The locations of TFBSs with MNase protection for *in vivo*-targeted nucleosomes (ITNs) are shown (red color scale at bottom). MNase protection was measured as the percentage of nucleosome bases that were protected from MNase digestion and calculated for each base pair as the ratio of base pair coverage to the total reads for that specific nucleosome: **a**, Lin28B nucleosome sequence bound with O; **b**, ITN bound with O, S, K, and M. **c–d**, ITN bound with O, S, and K; **e,** ITN bound with O and S. O, OCT4; S, SOX2; K, KLF4; M, MYC. The relative supershifts for each nucleosome are shown for KLF4, MYC, OCT4, and SOX2 binding on the right.

plotted there relative supershift for all ITNs containing a binding site. KLF4 and SOX2 bind to a

larger percentage of their ITNs then MYC and OCT4 (Fig. 6a).

To understand the role of nucleosome positioning for the ITNs we mapped the TFBS in

relationship to the center of MNase protection (Fig. 6b-e). Binding of KLF4, OCT4, SOX2, and

MYC are significantly correlated with the distance of the TFBS from MNase protection center.

These results are very similar to the results for 5S and MMTV, suggesting that distance to dyad

is a major factor regulating TF nucleosome binding. To examine binding near the nucleosome

center we selected the region 40 bp from the MNase center and determined the percentage of

bound sites to the total number of sites within 40 bp. KLF4 bound 67%, SOX2 32%, MYC 13%,

and OCT4 0%. This suggests that KLF4 and SOX2 have a special ability to target some of their

Fig 6. Binding to *in vivo*-targeted nucleosomes. a, Violin plot showing relative supershifts for KLF4, MYC, OCT4, and SOX2 binding to *in vivo*-targeted nucleosomes (ITNs) containing their transcription factor binding sites (TFBSs). **b–e,** Relative supershifts for binding to ITNs compared to the distance of the TFBS distance from the center of MNase protection.

TFBSs within a nucleosome and support the results with NPSs showing that KLF4 bound at

SHL -2 and -3 and SOX2 bound at SHL -3 and -4.

DISCUSSION

Pioneer factors are a proposed class of TFs that can bind inaccessible genomic regions and

facilitate the binding of other TFs. This ability is attributed to specific DNA binding domains that

are able to target their binding sites within a nucleosome[31] and bind to partial motifs that are

accessible on the nucleosome surface[14]. However, recent studies with the archetypal pioneer

factor FOXA1 and its non-pioneer cofactor HNF4A have shown that binding is defined by the

TFBS, specifically its density and affinity, rather than differences in TFs themselves[33]. Our

results suggest a more complex model.

The TFs we tested bound at the nucleosome edge, but this differed according to the NPS. These results are consistent with a dynamic partial unwrapping of DNA from histones at sites where the DNA enters or exits the nucleosome, exposing the DNA to TFs[34, 35]. Edge binding would be heavily dependent on TFBS affinity and the number of sites. We also found that two of the TFs (KLF4 and SOX2) targeted sites close to the nucleosome dyad at specific rotational settings. By using ITNs, we were able to show that KLF4 and SOX2 behave differently than OCT4 and can target their binding sites near the nucleosome center.

KLF4 appears to be binding to two partial motifs (CCCG) located in consecutive partially exposed major grooves, similarly to the nucleosome binding of another zinc finger TF, GATA3. GATA3 binds to split 5′-GAT-3′ motifs in solvent-exposed major grooves[36]. KLF4 will only bind the non-typical motif at these internal locations, which suggests that the partial motif or neighboring bases provide a structure binding context that is missing for other TFBSs. The binding of KLF4 to the internal p53 sites in the 601 NPS but not the other NPS suggests that nucleosome structure also facilitates certain binding events. Although the overall nucleosome structure is consistent across different NPS, certain patterns of dinucleotides cause differences in the groove width and helical deformation of nucleosomal DNA[37, 38]. Indeed, groove width and the extent of helical deformation vary substantially among different sites with the same rotational settings[39, 40]. Single-molecule DNA unzipping experiments have shown that position-specific histone-DNA interactions also vary across the nucleosome[41].

There are still many questions about how TFs can bind nucleosomal DNA that can be addressed with Pioneer-seq. The vast majority of studies on nucleosomes have used the Widom 601 NPS. This sequence is extremely well-studied and is the sequence used for 114 structural studies. Widom 601 forms nucleosomes *in vitro* very efficiently in a single predominate position allowing reproducible and well-defined structures [15]. 5S and MMTV nucleosomes are biologically derived sequences that have been used for various studies [37,

42]. 5S and MMTV nucleosomes are not as stable as 601 and form with efficiencies similar to ITNs (Extended Data Fig. 3). Binding at nucleosomes with different NPSs also varied among all TFs in this study. In general, 601 nucleosomes were strongly inhibitory for TF binding, whereas 5S and MMTV nucleosomes were only inhibitory when the TBFSs were close to the nucleosome dyad. These results suggest that model nucleosomes should not be the sole type of nucleosomes relied on for future studies. TF binding may be best understood by examining both ITNs and model nucleosomes. Use of ITNs does have limitations, because exact positioning of the dyad is unknown and weak nucleosome formation could limit specific sequences from being tested. For Pioneer-seq, we propose the use of ITNs along with a model nucleosome in which the TFBS can be tiled in all possible locations.

The ability for some TFs to bind a specific nucleosome site appears to depend on the binding sequence. This suggests that certain sites that require a pioneer factor may also have an alternative motif. The bases flanking the motif could impact nucleosome binding by affecting the structural presentation of the TFBS. TF binding could also be affected by events that influence the shape of the DNA around the nucleosome, such as DNA methylation, histone variants, and histone modifications. Notably, Pioneer-seq can be used to investigate these possibilities.

Lastly, transcription is regulated by a complex of multiple TFs that bind proximal regulatory regions. These multi-TF binding events can be directly cooperative, as seen when TFs physically interact, or indirectly cooperative to displace the nucleosome[43]. Pioneer-seq is ideally suited to test various models of cooperativity between TFs and enable the mechanistic dissection of these crucial regulatory events.

Conclusion

In summary, Pioneer-seq is a powerful method for investigating the essential first step in gene regulation, binding of TF to inaccessible DNA located within nucleosomes. Due to its nature of comparing binding to non-specific binding across a whole nucleosome library, Pioneer-seq allows the direct comparison of sites located in various nucleosome positions, with differing NPSs, and across TFBS variants. Pioneer-seq can be applied to directly address various mechanistic models for TF-nucleosome binding and can be used to uncover inherent TF interaction differences.

Methods

Pioneer-seq library design

Three nucleosome positioning sequences (NPS), namely, Widom 601, 5S rDNA, and mouse mammary tumor virus (MMTV)-A, were used to form stable nucleosomes. Each sequence has been characterized by multiple biochemical assays[44] and was scanned for the presence of binding sites for the transcription factors (TFs) of interest using FIMO; sequences were modified to remove the binding sites[45]. Sequences were generated with the TF binding site (TFBS) of interest placed at every base pair positing in the nucleosome, with sites in both linker regions. In total, 149 sequences were designed for every TFBS with each NPS.

In vivo-targeted nucleosomes (ITNs) were determined by integrating datasets from chromatin immunoprecipitation with sequencing (ChIP-seq) and nucleosome positioning datasets. Locations bound by OCT4, SOX2, KLF4, or MYC were determined from ChIP-seq datasets[27]. Bound sites were checked for the specific TFBS, and locations lacking an identifiable TFBS were removed. Nucleosome positions were determined from NOMe-seq (nucleosome occupancy and methylome sequencing) from IMR90 cells (GSM543311) using the DANPOS algorithm[32, 46]. The position serving as the nucleosome center was then expanded to 191 bp and evaluated with a nucleosome scoring function[21]. The probability that the center base is part of a nucleosome was used as the probability score for each nucleosome. Nucleosomes with a score of <0.7 were removed from the design.

Nucleosome library assembly

All nucleosome sequences were flanked by primer sequences to generate 230-bp sequences. The nucleosome library containing a total of 7,500 unique sequences was acquired from Agilent as a custom oligonucleotide library, which was amplified using Herculase II Fusion DNA

polymerase in 100-µl reaction mixtures (1x Herculase II reaction buffer, 1 mM dNTPs, 200 pM

Agilent library, 250 nM forward and reverse primers) with 15 PCR cycles. For a typical

experiment, the DNA obtained from 11 reactions was purified with a QIAquick PCR purification

kit (cat. no. 28104; Qiagen) and quantified with a NanoDrop spectrometer; fragment size was

confirmed with a 2% agarose gel. Nucleosomes were then generated from H2A/H2B dimers and

H3.1/H4 tetramers (NEB) by incubating the DNA sequences and the histones at an

octamer/DNA molar ratio of 1:1.2 (in a solution containing 10 mM dithiothreitol [DTT] and 1.8 M

NaCl) for 30 min at room temperature. The reaction mixture was transferred to a Slide-A-Lyzer

MINI dialysis unit (10,000 MWCO, cat. no. 69750; Thermo Scientific). Dialysis was performed

with 1.2 ml of the dialysis buffers at 4°C in 1.0 M NaCl for 2 h, 0.8 M NaCl for 2 h, 0.6 NaCl for 2

h, and TE buffer (pH 8.0) overnight at 4°C. Nucleosomes were then transferred to a clean 1.5-

ml tube pretreated with 0.3 mg/ml bovine serum albumin (BSA). Nucleosome formation was

then confirmed by 4% native polyacrylamide gel electrophoresis. Free DNA was removed from

nucleosomes by using a 7%–20% sucrose gradient, and nucleosomes were concentrated and

quantified via qPCR[4, 5]. Nucleosomes were then stored at 4 °C for up to 1 month.

DNA binding assay followed by EMSA

The protein-nucleosome binding assays were carried out by incubating the purified nucleosome

libraries described above and human full-length KLF4 (Origene TP306691), OCT4 (Origene

TP311998), SOX2 (Origene TP300757), and MYC (Origene TP307611) with MAX (Origene

TP306812) (in 7 µl DNA binding buffer (10 mM Tris-Cl [pH7.5], 50 mM NaCl, 1 mM DTT, 0.25

mg/ml BSA, 2 mM $MgCl_2$, 0.025% Nonidet P-40, and 5% glycerol) for 10 min on ice and then 30

min at room temperature. Increasing concentrations of TF (0 to 10 pmol) were added to 0.2 pmol

purified nucleosomes. Protein binding was detected by electrophoretic mobility shift assays

(EMSAs) on 4% (w/v) native polyacrylamide gels (acrylamide/bisacrylamide, 29:1 [w/w], 7 × 10 cm) in 0.5× Tris-borate-EDTA buffer at 100 V at 4 °C.

DNA isolation and purification

After electrophoresis, DNA was imaged by staining with SYBR green (LONZA). All visual bands as well as the bands at the same locations in the other lanes were excised from the gel. The chopped gel slices were soaked in diffusion buffer (0.5 M ammonium acetate, 10 mM magnesium acetate, 1 mM EDTA [pH 8.0], 0.1% SDS) and incubated at 50 °C overnight. The supernatant was collected, residual polyacrylamide was removed with glass wool, and the DNA was purified with QIAquick spin columns (Qiagen). The DNA concentration for each sample was determined by qPCR using a standard curve generated from a control sequence.

Library construction and sequencing

Illumina sequencing libraries were generated using a two-step PCR method, with 8 to 12 cycles of amplification for the first step including four sets of primers designed to offset sequence reads and dephase the libraries during Illumina sequencing (Extended Data Table 1). The number of cycles for the first-round PCR was determined using the sample concentration determined by qPCR. Each sample was then indexed using Nextera dual indices (Nextera XT index primer 1 [N7xx] and Nextera XT index primer 2 [S5xx]). After each PCR, reaction mixtures were cleaned up with AMPure XP beads (Beckman Coulter). The concentration of each sample was determined using the Invitrogen Quant-iT dsDNA assay kit, and equal amounts of each sample DNA were pooled and sequenced on an Illumina NextSeq 2x150. Sequencing and quality control were performed at the University at Buffalo Genomics and Bioinformatics Core.

Pioneer-seq analysis

Illumina sequence reads were processed with an automated Snakemake pipeline of applications to refine and identify the sequences present in the sample pool[47]. The 3′ ends of Illumina FASTQ reads with low-quality scores were removed with Cutadapt using a quality cutoff of 30 (-q 30)[48]. Forward and reverse FASTQC reads were merged with Vsearch (--fastq_mergepairs) only if they shared at least 20 overlapping nucleotides (--fastq_minovlen 20) and had no more than two mismatched nucleotides between them (--fastq_maxdiffs 2)[49]. Primer sequences present at the ends of FASTQ reads were removed with Cutadapt. FASTQ reads of >220 nucleotides (nt) or <174 nt were filtered out with Cutadapt (--maximum-length 220 --minimum-length 174)[48]. FASTQ reads were converted to FASTA format using the FASTX-Toolkit FASTQ-to-FASTA command[50]. FASTA reads were mapped to a sequence in the reference library of 7,500 nucleosome sequences with Vsearch (--dbmatched) only if they had alignment lengths of at least 150 nt (--mincols 150), had at least 98.5% similarity (--id 0.985), and were the query and database sequence pairing with the highest percentage of identity (--top_hits_only)[49]. The results were then analyzed relative to control/nonspecific binding (relative supershift).

Relative supershift was determined from the supershift bands and controls for technical variability introduced by gel excision, PCR, library construction, or sequencing. In this method, each specific nucleosome sequence is measured relative to that for nonspecific binding (nucleosomes lacking a TFBS):

$$\text{Relative supershift} = \log_2\left(\frac{\frac{\text{reads supershift}_N}{(\text{reads supershift}_{NS})}}{\frac{\text{reads nucleosome band TF null}_N}{(\text{reads nucleosome band TF null}_{NS})}}\right), \qquad \text{(Eq. 1)}$$

where N is one of the 7,500 nucleosome sequences, NS is the control nucleosome sequences, "reads supershift" is the supershift band, and "reads nucleosome band TF null" is the nucleosome band in the TF-null lane.

Pioneer-seq was performed with multiple concentrations of TFs because of differences in inherent binding affinities and protein purity. An initial analysis of linker binding events was used to (i) confirm specific binding of the TF of interest and (ii) define the TF concentration with the largest binding signal compared to that for nonspecific binding. The relative supershift for a single TF concentration is presented throughout this manuscript: 0.4 pmol for KLF4, 0.2 pmol for OCT4 and SOX2, and 0.8 pmol for MYC/MAX.

MNase-seq on nucleosome library

The nucleosome library (0.2 pmol/μl) was digested by MNase (0.05 U/μl) in nuclease digestion buffer (10 mM Tris-HCL [pH 8.0], 2 mM CaCl$_2$) for 25 min at 37 $^\circ$C; digestion was stopped with 2% SDS and 40 mM EDTA). Each sample was then incubated with proteinase K (16 μg) for 1 h at 55 $^\circ$C. The DNA was purified from the reaction and concentrated with the QIAquick PCR purification kit. The concentrations of DNA in each sample were determined with the Invitrogen Quant-iT dsDNA assay kit and equalized. Illumina sequencing libraries were generated using an NEBNext Ultra II DNA library prep kit. Individual samples were multiplexed and sequenced via Illumina MiSeq 2x150.

MNase-seq results were quality filtered ($q > 30$) and adapter trimmed using Cutadapt[51]. The quality reads were merged and mapped to the 7,500 nucleosome library sequences using Vsearch[52]. The read counts and end positions were used to measure MNase protection, which was calculated for each base pair as the ratio of base pair coverage to total reads for that specific nucleosome.

DNase I footprinting

Nucleosomal DNA 5′ labeled with FAM (6-carboxyfluorescein) was formed into nucleosomes and purified as described above. Nucleosomes (50 ng) were bound with 0.4 pmol of KLF4 in DNA binding buffer (10 mM Tris-HCL [pH 7.5], 1 mM $MgCl_2$, 10 µM $ZnCl_2$, 1 mM DTT, 10 mM KCl, 0.5 mg BSA, 5% glycerol) at room temp for 1 h. Each sample was then incubated with 0.06 U DNase I in 50 µl of digestion buffer (10 mM $MgCl_2$, 5 mM $ZnCl_2$) at 25 °C for 1 min; digestion was stopped with 90 µl stop solution ($NaCl_2$, 30 mM EDTA, 1% SDS). Digested DNA was then purified using phenol/chloroform/isoamyl alcohol and submitted to Roswell Park Genomic Facility for capillary electrophoresis fragment analysis on an ABI PRISM 3130xl Genetic Analyzer. The resulting data were analyzed using the Microsatellite analysis app from Thermo Fisher Scientific.

Modeling of a KLF4-bound nucleosome

The structure for the Widom 601 nucleosome[53] was retrieved from the Protein Data Bank (PDB)[54] (PDB identifier 5OXV). The location of the KLF4-bound motif was determined from the Pioneer-seq results, and the structural residues were colored using ChimeraX software [55].

Data availability

All Pioneer-seq results are stored on NCBI SRA at accession PRJNA892950.

References

1. Uhlen M, Oksvold P, Fagerberg L, Lundberg E, Jonasson K, Forsberg M, Zwahlen M, Kampf C, Wester K, Hober S, et al: **Towards a knowledge-based Human Protein Atlas.** *Nat Biotechnol* 2010, **28:**1248-1250.
2. Cirillo LA, Lin FR, Cuesta I, Friedman D, Jarnik M, Zaret KS: **Opening of compacted chromatin by early developmental transcription factors HNF3 (FoxA) and GATA-4.** *Mol Cell* 2002, **9:**279-289.
3. Iwafuchi-Doi M, Zaret KS: **Pioneer transcription factors in cell reprogramming.** *Genes Dev* 2014, **28:**2679-2692.
4. Yu X, Singh PK, Tabrejee S, Sinha S, Buck MJ: **DeltaNp63 is a pioneer factor that binds inaccessible chromatin and elicits chromatin remodeling.** *Epigenetics Chromatin* 2021, **14:**20.
5. Yu X, Buck MJ: **Defining TP53 pioneering capabilities with competitive nucleosome binding assays.** *Genome Res* 2019, **29:**107-115.
6. Michael AK, Thoma NH: **Reading the chromatinized genome.** *Cell* 2021, **184:**3599-3611.
7. Li Q, Wrange O: **Translational positioning of a nucleosomal glucocorticoid response element modulates glucocorticoid receptor affinity.** *Genes Dev* 1993, **7:**2471-2482.
8. Rossetti L, Cacchione S, De Menna A, Chapman L, Rhodes D, Savino M: **Specific interactions of the telomeric protein Rap1p with nucleosomal binding sites.** *J Mol Biol* 2001, **306:**903-913.
9. Angelov D, Lenouvel F, Hans F, Muller CW, Bouvet P, Bednar J, Moudrianakis EN, Cadet J, Dimitrov S: **The histone octamer is invisible when NF-kappaB binds to the nucleosome.** *J Biol Chem* 2004, **279:**42374-42382.
10. Blomquist P, Li Q, Wrange O: **The affinity of nuclear factor 1 for its DNA site is drastically reduced by nucleosome organization irrespective of its rotational or translational position.** *J Biol Chem* 1996, **271:**153-159.
11. Vettese-Dadey M, Walter P, Chen H, Juan LJ, Workman JL: **Role of the histone amino termini in facilitated binding of a transcription factor, GAL4-AH, to nucleosome cores.** *Mol Cell Biol* 1994, **14:**970-981.
12. Chaya D, Hayamizu T, Bustin M, Zaret KS: **Transcription factor FoxA (HNF3) on a nucleosome at an enhancer complex in liver chromatin.** *J Biol Chem* 2001, **276:**44385-44389.
13. Sekiya T, Muthurajan UM, Luger K, Tulin AV, Zaret KS: **Nucleosome-binding affinity as a primary determinant of the nuclear mobility of the pioneer transcription factor FoxA.** *Genes Dev* 2009, **23:**804-809.
14. Soufi A, Garcia MF, Jaroszewicz A, Osman N, Pellegrini M, Zaret KS: **Pioneer transcription factors target partial DNA motifs on nucleosomes to initiate reprogramming.** *Cell* 2015, **161:**555-568.
15. Yu X, Buck MJ: **Pioneer factors and their in vitro identification methods.** *Mol Genet Genomics* 2020.
16. Zhu F, Farnung L, Kaasinen E, Sahu B, Yin Y, Wei B, Dodonova SO, Nitta KR, Morgunova E, Taipale M, et al: **The interaction landscape between transcription factors and the nucleosome.** *Nature* 2018, **562:**76-81.
17. Michael AK, Grand RS, Isbel L, Cavadini S, Kozicka Z, Kempf G, Bunker RD, Schenk AD, Graff-Meyer A, Pathare GR, et al: **Mechanisms of OCT4-SOX2 motif readout on nucleosomes.** *Science* 2020, **368:**1460-1465.
18. Simpson RT, Stafford DW: **Structural features of a phased nucleosome core particle.** *Proc Natl Acad Sci U S A* 1983, **80:**51-55.
19. Richard-Foy H, Hager GL: **Sequence-specific positioning of nucleosomes over the steroid-inducible MMTV promoter.** *EMBO J* 1987, **6:**2321-2328.

20.   Lowary PT, Widom J: **New DNA sequence rules for high affinity binding to histone octamer and sequence-directed nucleosome positioning.** *J Mol Biol* 1998, **276:**19-42.

21.   Kaplan N, Moore IK, Fondufe-Mittendorf Y, Gossett AJ, Tillo D, Field Y, LeProust EM, Hughes TR, Lieb JD, Widom J, Segal E: **The DNA-encoded nucleosome organization of a eukaryotic genome.** *Nature* 2009, **458:**362-366.

22.   Tsompana M, Buck MJ: **Chromatin accessibility: a window into the genome.** *Epigenetics Chromatin* 2014, **7:**33.

23.   Rizzo JM, Bard JE, Buck MJ: **Standardized collection of MNase-seq experiments enables unbiased dataset comparisons.** *BMC Mol Biol* 2012, **13:**15.

24.   Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S: **Induction of pluripotent stem cells from adult human fibroblasts by defined factors.** *Cell* 2007, **131:**861-872.

25.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK: **Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities.** *Mol Cell* 2010, **38:**576-589.

26.   Serandour AA, Avner S, Percevault F, Demay F, Bizot M, Lucchetti-Miganeh C, Barloy-Hubler F, Brown M, Lupien M, Metivier R, et al: **Epigenetic switch involved in activation of pioneer factor FOXA1-dependent enhancers.** *Genome Res* 2011, **21:**555-565.

27.   Soufi A, Donahue G, Zaret KS: **Facilitators and impediments of the pluripotency reprogramming factors' initial engagement with the genome.** *Cell* 2012, **151:**994-1004.

28.   Fornes O, Castro-Mondragon JA, Khan A, van der Lee R, Zhang X, Richmond PA, Modi BP, Correard S, Gheorghe M, Baranasic D, et al: **JASPAR 2020: update of the open-access database of transcription factor binding profiles.** *Nucleic Acids Res* 2020, **48:**D87-D92.

29.   Avsec Z, Weilert M, Shrikumar A, Krueger S, Alexandari A, Dalal K, Fropf R, McAnany C, Gagneur J, Kundaje A, Zeitlinger J: **Base-resolution models of transcription-factor binding reveal soft motif syntax.** *Nat Genet* 2021, **53:**354-366.

30.   Hou L, Srivastava Y, Jauch R: **Molecular basis for the genome engagement by Sox proteins.** *Semin Cell Dev Biol* 2017, **63:**2-12.

31.   Fernandez Garcia M, Moore CD, Schulz KN, Alberto O, Donague G, Harrison MM, Zhu H, Zaret KS: **Structural Features of Transcription Factors Associating with Nucleosome Binding.** *Mol Cell* 2019, **75:**921-932 e926.

32.   Kelly TK, Liu Y, Lay FD, Liang G, Berman BP, Jones PA: **Genome-wide mapping of nucleosome positioning and DNA methylation within individual DNA molecules.** *Genome Res* 2012, **22:**2497-2506.

33.   Hansen JL, Loell KJ, Cohen BA: **A test of the pioneer factor hypothesis using ectopic liver gene activation.** *Elife* 2022, **11**.

34.   Li G, Widom J: **Nucleosomes facilitate their own invasion.** *Nat Struct Mol Biol* 2004, **11:**763-769.

35.   Polach KJ, Widom J: **Mechanism of protein access to specific DNA sequences in chromatin: a dynamic equilibrium model for gene regulation.** *J Mol Biol* 1995, **254:**130-149.

36.   Tanaka H, Takizawa Y, Takaku M, Kato D, Kumagawa Y, Grimm SA, Wade PA, Kurumizaka H: **Interaction of the pioneer transcription factor GATA3 with nucleosomes.** *Nat Commun* 2020, **11:**4136.

37.   Frouws TD, Duda SC, Richmond TJ: **X-ray structure of the MMTV-A nucleosome core.** *Proc Natl Acad Sci U S A* 2016, **113:**1214-1219.

38.   Koyama M, Kurumizaka H: **Structural diversity of the nucleosome.** *J Biochem* 2018, **163:**85-95.

39.   Xu F, Olson WK: **DNA architecture, deformability, and nucleosome positioning.** *J Biomol Struct Dyn* 2010, **27:**725-739.

40. West SM, Rohs R, Mann RS, Honig B: **Electrostatic interactions between arginines and the minor groove in the nucleosome.** *J Biomol Struct Dyn* 2010, **27:**861-866.
41. Ettig R, Kepper N, Stehr R, Wedemann G, Rippe K: **Dissecting DNA-histone interactions in the nucleosome by molecular dynamics simulations of DNA unwrapping.** *Biophys J* 2011, **101:**1999-2008.
42. Thiriet C, Hayes JJ: **Functionally relevant histone-DNA interactions extend beyond the classically defined nucleosome core region.** *J Biol Chem* 1998, **273:**21352-21358.
43. Mirny LA: **Nucleosome-mediated cooperativity between transcription factors.** *Proc Natl Acad Sci U S A* 2010, **107:**22534-22539.
44. Flaus A: **Principles and practice of nucleosome positioning in vitro.** *Frontiers in Life Science* 2011, **5:**5-27.
45. Grant CE, Bailey TL, Noble WS: **FIMO: scanning for occurrences of a given motif.** *Bioinformatics* 2011, **27:**1017-1018.
46. Chen K, Xi Y, Pan X, Li Z, Kaestner K, Tyler J, Dent S, He X, Li W: **DANPOS: dynamic analysis of nucleosome position and occupancy by sequencing.** *Genome Res* 2013, **23:**341-351.
47. Koster J, Rahmann S: **Snakemake-a scalable bioinformatics workflow engine.** *Bioinformatics* 2018, **34:**3600.
48. Martin M: **Cutadapt removes adapter sequences from high-throughput sequencing reads.** *2011* 2011, **17:**3.
49. Rognes T, Flouri T, Nichols B, Quince C, Mahe F: **VSEARCH: a versatile open source tool for metagenomics.** *PeerJ* 2016, **4:**e2584.
50. Gordon A. HG: **FASTQ/A short-reads preprocessing tools (unpublished).** 2010.
51. Martin M: **Cutadapt removes adapter sequences from high-throughput sequencing reads.** *EMBnetjournal* 2011, **17:**10.
52. **FASTX-Toolkit** [http://hannonlab.cshl.edu/fastx_toolkit/index.html]
53. Ekundayo B, Richmond TJ, Schalch T: **Capturing Structural Heterogeneity in Chromatin Fibers.** *J Mol Biol* 2017, **429:**3031-3042.
54. Berman HM, Westbrook J, Feng Z, Gilliland G, Bhat TN, Weissig H, Shindyalov IN, Bourne PE: **The Protein Data Bank.** *Nucleic Acids Research* 2000, **28:**235-242.
55. Pettersen EF, Goddard TD, Huang CC, Meng EC, Couch GS, Croll TI, Morris JH, Ferrin TE: **UCSF ChimeraX: Structure visualization for researchers, educators, and developers.** *Protein Sci* 2021, **30:**70-82.