Auditory cortex encodes lipreading information through spatially distributed activity

Ganesan Karthik[1], Cody Zhewei Cao[1], Michael I. Demidenko[1], Andrew Jahn[1], William C. Stacey[2], Vibhangini S. Wasade[3,4], David Brang*[1]

[1] Department of Psychology, University of Michigan, Ann Arbor, MI 48109
[2] Department of Neurology, University of Michigan, Ann Arbor, MI 48109
[3] Henry Ford Hospital, Detroit, MI 48202
[4] Department of Neurology, Wayne State University School of Medicine, MI 48201

Corresponding Author: djbrang@umich.edu

Data Availability Statement

The data that support the findings of this study will be made openly available through the

University of Michigan Deep Blue Repository.

Conflict of Interest Statement

The authors declare no competing financial interests.

Acknowledgements

This study was supported by NIH Grants R00DC013828, 1R01DC020717, and R01NS094399.

Keywords

Multisensory; Audiovisual; Speech; ECoG; iEEG; sEEG

Abstract

Watching a speaker's face improves speech perception accuracy. These benefits are owed, in part,

to implicit lipreading abilities present in the general population. While it is established that

lipreading can alter the perception of a heard word, it is unknown how information that is extracted

from lipread words is transformed into a neural code that the auditory system can use. One

influential, but untested, hypothesis is that visual speech modulates the population coded

representations of phonetic and phonemic features in the auditory system. This model is largely

supported by data showing that silent lipreading evokes activity in auditory cortex, but these

activations could alternatively reflect general effects of arousal or attention, or the encoding of

non-linguistic features such as visual timing information. This gap limits our understanding of how

vision supports speech perception processes. To test the hypothesis that the auditory system

encodes visual speech information, we acquired fMRI data from healthy adults and intracranial

recordings from electrodes implanted in patients with epilepsy during auditory and visual speech

perception tasks. Across both methods, linear classifiers successfully decoded the identity of

silently lipread words using the spatial pattern of auditory cortex responses. Examining the time-

course of classification using intracranial recordings, lipread words were classified at significantly

earlier time-points relative to heard words, suggesting a predictive mechanism for facilitating

speech. These results support a model in which the auditory system combines the joint neural

distributions evoked by heard and lipread words to generate a more precise estimate of what was

said.

Significance Statement

When we listen to someone speak in a noisy environment, watching their face can help us

understand them better, largely due to automatic lipreading abilities. However, it unknown how

lipreading information is transformed into a neural code that the auditory system can use. We used

fMRI and intracranial recordings in patients to study how the brain processes silently lipread words

and found that the auditory system encodes the identity of lipread words through spatially

distributed activity. These results suggest that the auditory system combines information from both

lipreading and hearing to generate more precise estimates of what is said, potentially by both

activating the corresponding representation of the heard word and suppressing incorrect phonemic

representations.

Introduction

Visual speech improves auditory speech perception during face-to-face conversations (1, 2). These benefits are strongest in noisy situations (3) and in individuals with hearing loss due to healthy aging (4), intrinsic brain tumor (5), stroke (6, 7), concussion (8, 9), or cochlear implants (10). However, there is limited understanding of how the brain enables vision to facilitate hearing processes.

The ability to extract useful information from visual speech signals (e.g., lipreading) is an implicit behavior that is rooted in the statistical relationship between auditory and visual cues in the natural environment (11). Lip dynamics are strongly correlated with different features of speech including temporal information (onset of words, rate of speech, and the boundaries between words) and relative spectral pitch based on the acoustics of the oral cavity (1). Most recognizably, the shape of the lips during speech is reliably associated with corresponding speech sounds (12); these simple lip shapes are described as visemes and are analogous to phonemes in the auditory domain (basic units of speech sounds).

Research has demonstrated that silent visual speech (e.g., lipreading) evokes activity within the auditory system (13, 14). Indeed, intracranial electroencephalography (iEEG) recordings indicate that visual speech influences processing in auditory regions through multiple temporal, spectral, and spatial configurations (15). While these findings highlight the broad influence of visual information on auditory speech processing, differences in activity do not provide a mechanistic account for how visual speech signals integrate with auditory neuronal populations. Among potential mechanisms, it is best understood that visual timing information during continuous speech biases auditory timing through phase-resetting mechanisms (16, 17). However,

it remains unclear how lipreading information (visemes) is transformed into a signal used by the

auditory system.

Within the auditory domain, phonetic and phonemic features are encoded by local and

distributed populations of neurons, respectively (18, 19). Mesgarani and colleagues (18) used

human iEEG recorded from high-density electrodes to demonstrate that phonemes are represented

by distributed populations of neurons in the STG. Combined with past research, these data support

a model in which the STG contains a patchy distribution of neurons that are tuned to specific

phonetic features via their spectro-temporal profiles (18, 20). For example, data show spatially

distinct responses in these regions to spectrally similar phonemes such as /ba/ and /da/ (20-22),

and clustered responses across a large phoneme-space (e.g., the distributed pattern of activity to

/ma/ is more similar to /na/ than it is to the spectro-temporally distinct phoneme /ba/ (18)). Indeed,

the identities of different *heard* phonemes can be decoded by the distribution of activity in the

auditory cortex (23), even when the physical auditory stimulus remains the same.

Building on this framework of auditory perception, we and others proposed that activity

from lipread visemes is relayed from visual regions to auditory cortex, preferentially modulating

the same populations of neurons that encode matching phoneme responses (15, 24). In this

hypothesis, heard and lipread activations in auditory cortex are combined through a winner-take-

all mechanism, in which the phoneme population with highest activation profile leads to the

phoneme that is perceived (24).

Here we test the hypothesis that the identities of individual visemes are represented in the

auditory system through distributed patterns of activation, and these spatial distributions match

corresponding phoneme representations. Auditory cortex activation magnitude and informational

content were examined using functional magnetic resonance imaging (fMRI) in healthy

individuals and iEEG recordings in patients with epilepsy during word perception tasks, in which

patients either saw the lip movements or heard the speech sounds for the same groups of words.

The identities of the different words were classified from fMRI and iEEG signals in auditory cortex

using support vector machine classifiers (SVMs). Results demonstrate that the auditory system

reliably encodes the identity of visemes using spatially distributed activity in a similar manner to

heard words. Moreover, visemes evoked spatially similar activity to matching phonemes,

consistent with the hypothesis that visual speech targets corresponding phoneme representations.

Figure 1. fMRI task schematic. Schematic of auditory and visual trials. Auditory trials began with a fixation cross followed by a
CV stimulus (either /mama/, /fafa/, or /kaka/). Visual trials presented the visual components of these same recordings without the
corresponding audio track. After stimulus offset, subjects were cued to identify which of the three phonemes (or visemes) they saw
(or heard) via button press.

Results

fMRI Experiment

In Experiment 1 we presented subjects ($n = 64$) with consonant vowel (CV) syllables while

fMRI activity was acquired. Each trial included the silent video or auditory stimulus taken from a

speaker producing the CVs /mama/, /fafa/, or /kaka/ (Fig. 1). Stimuli were presented using an

optimized event-related design and data were analyzed using univariate and decoding approaches

to examine the activation and information present in heard and lipread signals. Simultaneous

auditory-visual stimuli were not included in this design to ensure adequate signal-to-noise ratios

for classifying silent visual-only speech.

Mean behavioral accuracy was high in both conditions: 95.67% (SD = 3.01%) in the

auditory-only condition and 92.31% (SD = 3.72%) in the visual-only condition. As expected, the

mean accuracy in the auditory-only condition was significantly higher than the visual-only

condition; $t(63) = 6.57$, $p < 0.001$, $d = 0.96$. None of the 64 subjects performed below the pre-

registered exclusion threshold (accuracy in either condition below 75%).

Previous research demonstrated that silent lipreading increased fMRI BOLD responses

within auditory regions (13). First, we replicated this finding using univariate contrasts in the

auditory-only and visual-only conditions (phonemes vs fixation and visemes vs fixation). Fig. 2a-

c shows the results of the whole-brain analyses, corrected for multiple comparisons using cluster

statistics (vertex-wise threshold of $P < 0.001$, cluster-corrected to $P < 0.05$, frontal regions

excluded as described in the Methods but see Supp. Fig. 1 for unmasked data). Complete statistics

for each analysis are reported in Supp. Tables 1-3 and beta estimates extracted from auditory and

visual regions of interest (ROIs) are shown in Supp. Fig. 2. Phonemes elicited significantly

increased BOLD activity within the STG bilaterally and decreased BOLD within visual regions.

Visemes showed a similar pattern in the auditory system with increased BOLD in the STG and

pSTS bilaterally, along with increased BOLD in visual regions, including hMT+, the fusiform

gyrus, and visual cortex. Importantly, phonemes elicited stronger activity relative to phonemes

throughout the STG.

Fig. 2d shows the normalized BOLD time-series from four regions of interest. Notably,

BOLD response timing in the STG was similar across phoneme and visemes, consistent with a

model in which visual speech biases early auditory perceptual processes. Conversely, visemes

elicited activity with the pSTS significantly before phonemes, suggesting a potential pathway for

the transformation of visemes to phonemes (consistent with previous studies; e.g. (25)). For

example, at 3.2 seconds following stimulus onset, visemes evoked significantly greater activity in

the pSTS, $t(63) = 5.68$, $p < .001$ (FDR-corrected), $d = 0.710$. These plots show the time-series

scaled to have the same amplitude across conditions to better see similar onset times in the STG

and different onset times elsewhere. Unnormalized data are shown in Supp. Fig. 3.

Figure 2. Univariate activations. (a) phonemes vs fixation (b) visemes vs fixation, and (c) phonemes vs visemes. Phonemes
evoked maximally increased activity in the STG bilaterally. Visemes evoked increased activity within bilateral visual cortex, left
pSTS, right MT, and the STG bilaterally, though significantly weaker within the STG compared to phonemes. Colored regions
reflect significant increases (red and yellow) or decreases (blues) in task-related activation (thresholded at p<.001 and corrected for
multiple comparisons using cluster-statistics). (d) BOLD time-series extracted from four regions (averaged across left and right
hemispheres). Data were normalized based on the maximum average value between 3.2 (TR=5) and 9.6 (TR=13) seconds to better

visualize relative shapes of the time-series. Notably, phonemes and visemes showed a similar temporal response in the STG but
visemes elicited activity in the pSTS more quickly than phonemes. Stimulus onset occurred at TR=1.

Univariate contrasts reveal activation magnitude but not informational content or

representational structure. Previous decoding-based approaches using fMRI (26, 27) and iEEG

(28, 29) demonstrated that auditory speech identities could be reconstructed from spatially

distributed activity in auditory cortex. To examine whether viseme identity is represented in the

auditory system by distributed patterns of activity, we used multivariate pattern analysis (MVPA)

(30) to classify individual phoneme and viseme labels.

Whole-brain searchlight-based MVPA was applied at the individual-subject level (31)

conducted separately for each of the two conditions of interest (auditory-only and visual-only

trials). Results of the whole-brain analysis, corrected for multiple comparisons using cluster

statistics are shown in Fig. 3 (vertex-wise threshold of $P < 0.001$, cluster-corrected to $P < 0.05$);

full statistics for each analysis is reported in Supp. Tables 4 and 5. In the auditory-only condition,

peak decoding accuracy was observed bilaterally in the STG and pSTS. This is consistent with

previous studies demonstrating phonetic representations in the STG using MVPA (32). In the

visual-only condition, peak decoding accuracy was observed within the STG bilaterally, the left

pSTS, visual cortex bilaterally, and right hMT+. To understand the spatial overlap of phoneme and

viseme representations in the auditory system we compared the spatial distribution of the

classification maps. Within the STG, results showed that a majority of vertices contained either

only phoneme information or both phoneme and viseme information, with very few vertices

representing viseme information alone (Fig. 3c). Across the left and right STG, phonemes (but not

visemes) were significantly classified at 27.2% of vertices, phonemes and visemes were jointly

classified from 25.2% of vertices , and visemes (but not phonemes) were significantly classified

at 0.16% of vertices. In total, phonemes were classified at twice as many vertices compared to

visemes within the STG (52.4% vs 25.4%).

The univariate analysis showed that visual speech increased BOLD activity broadly across

auditory regions and viseme decoding analyses identified above-chance classification accuracy in

a more restricted set of vertices throughout the STG. Comparing the two results, the univariate

visual-only analysis showed two times as many significant vertices in the STG (bilaterally)

compared to the area with significant viseme classification in the MVPA analysis (56.7% vs 25.4%

of STG vertices). This is consistent with the prediction that only a restricted proportion of the STG

encodes visemic information, while other regions reflect domain general responses to the visual

signals or the presentation of other visual information (e.g., temporal or spectral information (1)).

Conversely, the auditory-only analysis showed significant univariate responses in 79.2% of the

STG, and phonemes were significantly classified from 52.4% of vertices within the STG.

To further quantify the relative information across target regions, we performed individual-

subject SVM classification in five regions of interest (ROIs) in each hemisphere (dimension of

vertices within the ROI; Fig. 3d) and corrected p-values using FDR. As shown in Fig. 3e and Supp.

Fig. 4, phoneme classification accuracy was strongly above chance (33.3%) in the left and right

STG and pSTS (all $p < .001$). Viseme classification accuracy was similarly above chance in the

left STG ($p < .001$) and left pSTS ($p < .05$). Visemes were additionally able to be classified from

multiple visual regions, most notably in right hMT+ ($p < .001$), right fusiform gyrus ($p < .01$), and

right V1/V2 ($p < .01$). To confirm that our viseme decoding results were generated from auditory

selective regions as opposed to non-selective regions of the STG, we repeated viseme classification

only at voxels within the STG that showed significant increase in BOLD activity to phonemes

relative to fixation. These auditory selective results showed a similar pattern of effects with

significant viseme classification in both the left STG (*p* < .001) and right STG (*p* < .05).

Figure 3. fMRI decoding of phoneme and viseme information in an event-related design. (a-c) Searchlight-based MVPA
classification in *n* = 64 subjects. Classifiers were trained to identify (a) the phoneme heard (/fafa/, /mama/, or /kaka/) in the auditory-
only condition, (b) the viseme seen in the visual-only condition, or (c) condition differences between auditory-only and visual alone
trials. Decoding was conducted at the individual subject level and only group-level differences greater than chance (thresholded at
p<.001 and corrected for multiple comparisons using cluster-statistics) are shown. (a) Peak phoneme decoding was observed in the
bilateral STG. (b) Significant viseme decoding was observed in the bilateral STG, left pSTS, and visual regions. (c) Vertices with
significant classification of phonemes but not visemes (red), visemes but not phonemes (blue), or with significant classification of
both phonemes and visemes (purple). There is a large overlap in the vertices at which visemes and phonemes could be classified.
Restricted to the just the STG, vertices at which viseme classification was significant covered roughly half of the area in the STG
that phonemes were classified successfully at (48.1% overlap) with negligible area uniquely able to classify visemes. (d) Regions
of interest (ROIs) used for hypothesis driven classification at the single-subject level. (e) Results of classification at selected ROIs.
Phonemes were significantly classified from the left STG and pSTS. Visemes were significantly classified from the left STG
(consistent with the hypothesis that information about visemes is represented within the STG), pSTS, and visual cortex. Center line
reflects the mean, colored box SE, and the tails 95% confidence intervals. *p<.05, ***p<.001. Chance accuracy is 33.3%.
218
219

iEEG Experiment

Results from the fMRI study demonstrated that viseme information is represented in

auditory areas. Moreover, because visemes were classified based on the spatial distribution of

vertices in the STG, this supports a model in which lipreading is represented through population-

coded responses in the auditory system, similar to the neural representation underlying

phonemes(18). However, the slow temporal dynamics of fMRI signals prevent a fine-grained

analysis of the time-course of lipreading activation to examine when this information is available

to the auditory system. Additionally, the use of only three dissimilar CV stimuli prevented a more

graded analysis of these population-coded responses, such as whether more perceptually similar

phonemes (e.g., /ga/ and /da/) elicit more similar population-coded responses relative to

perceptually distinct phonemes (e.g., /fa/ and /ba/). To answer both of these questions we collected

data from a similar auditory-visual speech paradigm from $n = 6$ patients with epilepsy who had

electrodes implanted within auditory areas of the brain (Fig. 4a). Patients were presented with 240

auditory-only (listening) and 240 visual-only (lipreading) trials containing a single 1-2 syllable

word. Each word began with one of four consonants ('B', 'F', 'G', or 'D') to enable the classification

of distinct phonemic patterns. 40 distinct words were used (10 containing each of the 4 initial

consonants; Supp. Table 6) and each word was repeated 6 times within each condition. On each

trial subjects selected the initial consonant heard or seen from four options (4-alternative forced

choice paradigm). Subjects' mean behavioral accuracy across auditory-only and visual-only trials

was significantly above chance (25%) at the group level: auditory-only (M = 91.9%, SD = 8.1%,

$t(5) = 20.2$, $p < .001$, $d = 8.23$), visual-only (M = 67.4%, SD = 17.9%, $t(5) = 5.78$, $p = .002$, $d =$

2.36). As expected, auditory-only trials were correctly identified significantly more often than

visual-only trials, $t(5) = 5.93$, $p = .002$, $d = 2.42$.

Words in both the auditory-only and visual-only conditions evoked activity broadly

throughout the STG and MTG consistent with prior work (15). Examining the spectral breakdown

of these responses (Fig. 4b-d), phonemes evoked increased theta and high gamma power (HGp)

and suppressed beta power following word onset; Supp. Fig. 5 shows the spectral responses for

different regions of the STG. In natural speech, visual articulations typically occur before the onset

of speech-related sounds (typically within 40 - 200 ms of speech onset (33)). Because of this pre-

articulatory visual information, visemes evoked increased beta suppression beginning before the

expected phoneme onset time (-250 to 0 ms) $[t(5) = 2.78, p = .039, d = 1.13]$. This is consistent

with past research (15) and indicative of possible lateral(34) or feedback inputs (35-37) into the

auditory system.

Additionally, visemes elicited moderate changes in HGp following sound onset, even

though no sound was present. Viseme-related HGp increases were maximal at the posterior STG,

consistent with past research (15, 38, 39). Fig. 4e shows this pattern in single electrode HGp

responses from two patients (both electrodes within the left posterior STG), with HGp changes

occurring in response to visemes before phonemes. This pattern was distinct from responses in the

fusiform gyrus, at which visemes evoked early HGp increases following the onset of the visual

stimulus and no reliable response at any point during auditory-only trials (Fig. 4f).

Figure 4. iEEG results during an auditory-only (listening) and visual-only (lipreading) speech perception paradigm. (a) Distribution of all recorded electrodes (those beneath the pial surface not shown) (*n* = 6 patients). (b-d) Event-related spectral perturbations (ERSP) plots from all STG electrodes, averaged across subjects. (b) Phoneme responses peaked after sound onset with theta and high-gamma power (HGp) increases, as well as beta suppression. (c) Viseme responses evoked maximal changes in beta power, with increased beta suppression starting before the expected time of sound onset. (d) Difference between phoneme and viseme ERSP plots. (e) HGp responses from two STG electrodes in response to auditory-only trials (phonemes; red lines) and visual-only trials (visemes; blue lines). Posterior STG electrodes showed increased HGp responses to visemes before the time when speech sounds would be expected to begin. (f) HGp responses from two fusiform gyrus electrodes. Visemes evoked increased HGp shortly after onset of the face, with elevated HGp persisted throughout the visual movement period. Phonemes failed to evoke reliable changes in activity within the fusiform gyrus. Shaded regions reflect single condition 95% confidence intervals. Light gray boxes show significant between condition differences (multiple comparisons corrected using FDR).

To replicate the results of our fMRI experiment and examine whether viseme information is represented within auditory regions, we decoded word information using spatial and temporal signals from iEEG electrodes. Fig. 5a shows group-level average classification accuracy for decoding the initial word consonant for auditory-only and visual-only trials. iEEG signals used for decoding analysis were recorded from electrodes from superiortemporal, middletemporal and supramarginal areas during 0-500 ms during each trial and downsampled to 10Hz. Electrodes were only included if they were deemed functionally significant: i.e., regardless of condition they had significant activity using a one-sample t-test (p<0.01, FDR-corrected). Classification was conducted separately in each subject using SVM classifiers on single-trial event-related potential

(ERP) responses (60 per consonant-initial auditory and visual words) using time points and

electrodes as dimensions. We observed significant classification (evaluated using binomial

statistics) in all six patients for both auditory-only (all $p < .05$) and visual-only conditions (all $p <$

.05) (single-subject statistics shown in Supplemental Table 7). Similarly, at the group-level we

observed classification accuracy reliably above chance for both auditory-only ($t(5) = 5.39$, $p =$

.0030, $d = 2.20$), and visual-only trials ($t(5) = 5.57$, $p = .0026$, $d = 2.27$). We additionally observed

a trend towards greater classification in auditory-only trials relative to visual-only trials ($t(5) =$

2.08, $p = .0916$, $d = 0.851$).

The successful classification of phonemes and visemes indicated that auditory areas

represent information about the consonant initial words for both auditory-only and visual-only

speech stimuli. The diagonal of the confusion matrices (Fig. 5b-c) shows that this classification

was robust for each of the four auditory-only and visual-only stimuli considered ($p < .05$ for 3 out

of 4 phonemes and 3 out of 4 visemes) (statistics shown in Supp. Table 8). Previous research has

shown that local auditory responses spatially cluster according to phonetic features (18); for the

stimuli used here, B, G, and D form one cluster and F another. Consistent with these clusters,

classification was higher for auditory words with the consonant F compared to words with the

consonants B, G, or D ($t(5) = 2.95$, $p = .0319$, $d = 1.20$); a similar trend was observed for visual-

only trials ($t(5) = 1.80$, $p = .1318$, $d = 0.736$), consistent with perceptual ambiguity of these items

in phoneme-space.

Following the analyses in the fMRI experiment, we additionally performed a follow-up

analysis examining visual-speech information only in electrodes that demonstrated significant

auditory activity. Electrodes were identified for inclusion to those within the STG that showed

significant auditory responses to phonemes. Even within this more restricted range of electrodes

we observed significant viseme classification in four of the six patients at the individual subject

level (all p-value less than .01) and significantly greater classification of visemes at the group level

relative to chance ($t(5) = 2.62$, $p = .047$, $d = 1.07$).

Figure 5. iEEG classification of phoneme and viseme identities from auditory ($n = 6$ patients) and visual ($n = 3$ patients)
regions. (a) Accuracy of an SVM classifier in identifying the correct initial consonant ('B', 'F', 'G', or 'D') from either auditory-only
or visual-only words classified at the individual-subject level from spatial and temporal iEEG information. Both visemes and
phonemes were reliably classified from auditory electrodes. Chance accuracy is 25% and plots show group-level boxplots. (b-c)
Group-averaged confusion matrices taken from 4-class auditory-only and visual-only SVM classifiers. Cells denote the frequency
at which each consonant-initial word was predicted (x-axis) relative to the true labels (y-axis). 'F' initial words were best classified
across both auditory-only and visual-only conditions, whereas 'G' and 'D' initial words were more readily confused. (d) Group-
averaged classification at individual time-points from auditory electrodes (phoneme-onset at 0 sec) showing significant
classification accuracy for both auditory-only and visual-only trials shortly after phoneme onset; in the visual-only condition, this
time-point reflected the associated speech onset time even though no auditory stimulus was presented. Shaded region reflects SEM.
(e) Spatial distribution of electrodes at which auditory-only (red) or visual-only (blue) trials were reliably classified (p<.05 based
on binomial statistics); purple electrodes reflect significant classification in both conditions (with 11 out of 14 of these electrodes
present in the superior temporal gyrus) and gray electrodes reflect non-significant classification in either condition. Electrodes

beneath the pial surface were projected out to the lateral surface for visualization. (f) Scatterplot quantifying the similarity of
classification frequency for auditory-only trials and visual-only trials from auditory electrodes (taken from 8-class classifier). Data
reflect pairwise classification values, with the first letter reflecting the real consonant label and the second letter the predicted
consonant label. For example, 'F' trials were predicted correctly at high frequency for both auditory and visual trials, whereas 'D'
trials were incorrectly labeled as 'F' trials infrequently across both auditory and visual trials. The high correlation ($r^2 = .8003$, p<.001
permutation test) is consistent with the hypothesis that visual speech evokes responses targeting similar distributions of neurons to
corresponding phoneme responses in the STG. (g) Group-level classification accuracy showing that responses in the fusiform gyrus
can distinguish between different visemes but not phonemes. (h-i) Group-level confusion matrices for auditory-only and visual-
only trials from fusiform gyrus electrodes.
330

To examine the time-course of auditory and visual speech representations within the

auditory system, we classified the identity of stimuli independently at 1 ms intervals (from -1000

ms to +1000 ms relative to phoneme onset time and smoothed the resultant time-series using a 50

ms moving average). Classification was applied separately for each subject and group-level

statistics were calculated across subjects (multiple comparisons were corrected using FDR applied

from 0 to 1000 ms) (Fig. 5d). Visemes were reliably classified (all $p < .05$ corrected) at the start

of this range from 0 - 23 ms (relative to sound onset) as well as multiple later periods in the time-

range. In contrast, phonemes were not classified until 96 ms after sound onset but demonstrated

continuous significant classification until 657 ms. Importantly, the early classification of visual-

only trials demonstrates that visemic information is available to the auditory system at the same

early perceptual stage as is phonemic information, instead of reflecting later categorical or

decision-level processes. It is additionally possible that viseme information is available to the

auditory system before phoneme information (see ref (40)) but the evidence for this in the current

data is weak and requires future confirmation: while classification of visemes was higher than

phonemes at 0 ms ($p < .05$ uncorrected), no multiple comparison corrected significant differences

were observed between the conditions at any time-point, likely due to the small sample size.

In a parallel set of analyses, we classified the identity of stimuli independently at each

electrode within an auditory region (including the STG, MTG, SMG) to understand the spatial

distribution of phoneme and viseme classification and their overlap. Phonemes were significantly

($p < .05$ using binomial statistics) classified from 65 out of 260 electrodes (25.0%) while visemes

were significantly classified from only 38 electrodes (14.6%). This pattern is similar to the overall

classification rate observed in our fMRI data, where phonemes were classified at three times as

many vertices compared to visemes within the STG. Restricted to only the STG, 14 electrodes

significantly classified visemic information in total, with 11 of these also significantly classifying

phonemic information, highlighting the spatial overlap of these processes. Again, this is consistent

with the pattern observed in our fMRI data, in which few vertices were sensitive to only visemic

information.

Because phonemes are represented through population coded responses, misclassification

can reveal information about related neural processes. For example, if the rate at which the

consonant /d/ is misclassified as /g/ in both auditory-only and visual-only trials is similar, it

suggests similar underlying representations. To test whether auditory cortex shows similar

representations for phonemic and visemic information, we calculated a correlation between each

of the phoneme-pairs across the phoneme and viseme group-averaged confusion matrices. Fig. 5f

shows the scatter plot reflecting classification rate for each consonant pair. Across auditory-only

and visual-only trials, classification (and misclassification) rates were highly correlated ($r^2 =$

$.8003, p < .001$ permutation test). Significance was calculated by randomly permuting the stimulus

labels of each trial and repeating the full classification analysis $n = 1000$ times. This is consistent

with our hypothesis that the spatiotemporal neural representation of viseme identities in the

auditory areas is similar to that of phonemes.

Three of the six subjects had electrodes along the ventral temporal lobe (including the

fusiform gyrus). To examine phoneme and viseme representations in this visual region, we

repeated classification on this restricted set of electrodes. Within visual electrodes, group-level

classification was significantly above chance for visual-only trials ($t(2) = 8.74, p = .0128, d = 5.05$)

but not auditory-only trials ($t(2) = 1.15$, $p = .3701$, $d = 0.662$). We additionally observed greater

classification in visual-only trials relative to auditory-only trials ($t(2) = 5.29$, $p = .0339$, $d = 3.05$).

This pattern was seen at the individual-subject level in all three subjects using binomial statistics

(all $p < .01$ for visual-only trials and all $p > .24$ for auditory-only trials).

Classification of ERPs revealed robust encoding of phoneme and viseme information in

the auditory system, driven by low-frequency oscillatory information (power and phase) that

reflects the excitatory/inhibitory balance of local neuronal populations (41). Higher frequency

activity (high-gamma power; HGp), in contrast, is associated with the average rate of action

potentials generated by a local population of neurons (42). Across HGp from all auditory

electrodes, we observed significant classification (evaluated using binomial statistics; $p < .05$) in

five out of six patients for auditory-only trials and three out of six patients for visual-only trials.

Similarly, at the group-level we observed classification accuracy reliably above chance for both

auditory-only ($t(5) = 3.74$, $p = .013$, $d = 1.53$), and visual-only trials ($t(5) = 3.56$, $p = .0162$, $d =$

1.45). We additionally observed a trend towards greater classification in auditory-only trials

relative to visual-only trials ($t(5) = 2.40$, $p = .0614$, $d = 0.981$). More reliable classification for

low-frequency signals evoked by visemes is consistent with the fMRI finding that classification

can occur in auditory regions that do not show increased firing rates in response to visual speech.

Discussion

Extensive research has shown that silent visual speech can modulate activity within

primary auditory regions in humans (13-15, 24, 43, 44). However, multiple sources of information

could be contained in these visual-driven auditory responses including visual motion and timing

information (45), visual parsing of speech rate (11), visual-derived spectral information (1),

general effects on attention or arousal (34), or viseme-to-phoneme transformations (15). Here we

tested the hypothesis that the identities of individual visemes are represented in the auditory system

through distributed patterns of activation, and these spatial distributions match corresponding

phoneme representations. Using fMRI and intracranial electrodes implanted in auditory regions

we found that the auditory system reliably encodes the identity of visemes using spatially

distributed activity in a similar manner to phonemes. Moreover, visemes evoked similar patterns

of confusability as phonemes, consistent with the hypothesis that visual speech activates

corresponding phoneme representations.

Data from both fMRI and iEEG showed reliable classification of visemes from auditory

regions, maximal in the left pSTS and STG bilaterally. The spatial distribution of significant

viseme classification suggest multiple routes from visual cortex (46) to auditory cortex via hMT+

and the pSTS. Consistent with this view, visemes increased BOLD activity in hMT+ and the pSTS

before these regions responded to phonemes. Future functional connectivity analyses can be used

to examine the paths of transmission of lipreading information from visual to auditory regions and

computational analyses to examine how viseme information is transformed into phoneme or

phonetically tuned features. For example, dynamic causal modelling (DCM) has previously shown

that visual speech modulates auditory processing through ventral and dorsal routes (47).

Classification of iEEG data enables inferences about *when* phoneme and viseme

information is available to the auditory system. This temporal resolution is necessary to understand

whether visemes are used by the auditory system during the same time window that auditory

phonemes are processed in order to support the population-coded perceptual representations of

phonemes. Alternatively, if viseme representations follow the perceptual encoding of phonemes

(e.g., >500 ms) visemes may instead support categorical decisions about what was heard. The

present data showed significant classification accuracy for both auditory-only and visual-only

trials shortly after phoneme onset, indicating that visemic information is available to the auditory

system at the same perceptual stage as is phonemic information. Nonetheless, it remains possible

that silent visual speech can provide visemic information to the auditory system before phoneme

onset in cases which visual speech precedes auditory onset (40).

While we observed increased BOLD activity in response to lipreading broadly throughout

the STG, this does not necessarily indicate that lipreading increased the spiking rate of excitatory

neurons within all auditory regions. Specifically, research indicates that BOLD increases can

reflect changes in either local excitation or inhibition as both can increase metabolic activity (48,

49) and recent work indicates that even positive BOLD can reflect decreased metabolic activity in

some contexts (50). iEEG responses provide greater clarity on these mechanisms as lipreading has

been shown to elicit increased HGp in the posterior STG (15, 40), associated with increased

neuronal firing rates (51), but beta suppression in the mid to anterior STG (15), associated with

inhibitory processes (37). The notion that silent visual speech may suppress neural activity within

portions of the auditory system has been theorized to reflect the optimized tuning of neurons

specialized for auditory speech (52). Despite these differences in activation, above-chance

classification was observed throughout the STG and pSTS suggesting two distinct mechanisms

through which visual information is used to modulate phoneme populations. In posterior

activations, we hypothesize that silent visual speech selectively activates matching phonetically or

phonemically-tuned neurons in a categorical manner. Conversely, visemes may suppress activity

in the STG in a targeted manner to inhibit incorrect representations in phonetically tuned neuronal

populations (40) to indirectly refine the representation of correct phonetic features. While

speculative, one possible explanation for why visemes avoid directly activating matching

phonetically tuned neurons is to limit the potential for crossmodal hallucinations (53); i.e., hearing

speech during silent lipreading.

Our current study demonstrates that visual speech information can be decoded in the

auditory region, aligning with a growing body of research, including the findings of Brohl et al.

(54), which underscores the multisensory nature of speech processing. While both studies

underscore auditory cortex's role in integrating visual speech, our approach, utilizing a

combination of fMRI and intracranial EEG, provides unique spatial precision. This level of detail

complements the temporal resolution offered by MEG, as used by Brohl et al., enabling a more

comprehensive understanding of how visual and auditory speech information is integrated at

different neural levels. Our controlled study design, focusing on specific parameters like stimulus

length and voice onset time (VOT), offers distinct advantages over naturalistic designs such as

those used by Brohl et al. (54). By minimizing confounds due to natural correlations between

auditory and visual speech elements, we can more accurately isolate and test the encoding of visual

speech information in the auditory cortex. This approach is particularly relevant when considering

the findings of Brohl et al., who highlighted the auditory cortex's role in reflecting unheard acoustic

features during lip reading and established a behavioral relevance to these neural representations.

A limitation of the present work is that the small set of phonemes and visemes presented

provide a limited account of the full distribution of phonemes and visemes present in English. This

limitation was necessary to ensure adequate signal-to-noise ratios to enable classification of the

individual phonemes and visemes, but future research can examine the full distribution of phoneme

and viseme representations using more natural speech stimuli (55) in auditory-visual contexts

using high-density intracranial electrodes. Data from such experiments would be predicted to show

that phoneme tuning functions (the spatial selectivity of responses to a specific phoneme) will be

more precise (narrower and more distinct from other phonemes) during auditory-visual speech

compared to auditory-only speech. Moreover, we predict that phoneme and viseme spatial maps

will imperfectly overlap (as the same viseme could denote 'pet' or 'bet') and that the dissimilarity

in phoneme and viseme maps explain categorical shifts in perception during the McGurk effect (a

perceptual illusion in which visual speech alters which phoneme is heard (56)).

In sum, the present studies provide the first direct evidence that the visemic features from

lipread words are represented in the auditory system at a perceptual stage through spatially

distributed neural activity. The data are consistent with the model that visemes share auditory

representations with phonemes, which may serve to refine phonetic and phonemic population

responses, to in turn support speech perception fluency. Importantly, while viseme identities could

be classified from a majority of vertices in the STG that responded in general to visual speech,

auditory regions likely encode a variety of other visual features (e.g., visual motion timing, visual

parsing of speech rate, visual-derived spectral information) useful for facilitating speech

perception processes in the natural environment.

Methods

fMRI Experiment

Planned analyses and sample size stopping justification for the fMRI study were pre-

registered at OSF (https://osf.io/6fzwd/?view_only=60484583a2bb4dcdb8e27788c7c4c373).

Minor deviations from the pre-registered protocol are noted throughout the methods section. The

study was approved by the Institutional Review Board (IRB) of the University of Michigan.

Subjects

FMRI data was acquired from n = 64 subjects (F = 47, M = 17) recruited from the

University of Michigan's Psychology paid-subject pool (individuals who had previously expressed

interest in research studies) and through word of mouth. Subjects' ages ranged from 18-32 (Mean:

22.87, SD = 3.29) and included 56 right-handed, 7 left-handed, and 1 ambidextrous individual.

Written consent was obtained from each subject. subjects were paid USD $20 per hour for their

time. Data was collected from each subject in a single session lasting approximately 1 hour and 15

minutes. Because power analyses using multivariate pattern analyses (MVPA) remain a challenge,

we determined our sample size based on univariate power analysis (on the assumption that this

would yield a minimum acceptable sample size). Sample size to detect visual-only effects was

determined using data from the auditory-only condition in a preliminary sample using NeuroPower

(using random field theory, cluster threshold p=.05, alpha=.05, n=27). Estimated sample sizes

ranged from n=62 to 64 across the pairwise phoneme comparisons (/fafa/ vs /mama/, /fafa/ vs

/kaka/, and /kaka/ vs /mama/) and n=64 was selected to ensure adequate power. No data from the

visual-only condition was analyzed prior to submission of the pre-registration.

Tasks, Stimuli and Experimental Design

We used an auditory and visual speech paradigm optimized for an event-related fMRI

design. On each trial, subjects were presented with a three-alternative forced-choice task that

consisted of either an auditory-only stimulus or a visual-only stimulus. Three types of phonemes;

/fafa/, /kaka/ and /mama/ and three types of visemes; /fafa/, /kaka/ and /mama/ were used for this

task and randomly mixed within blocks. These specific phonemes were chosen to maximize the

differentiability between the individual phonemic representations in the neuronal populations of

the STG (28, 44).

512   Fig. 1 shows the timing and structure of the task. Each trial for both the auditory-only and

513 visual-only conditions lasted for 2 seconds. The auditory-only trials began with a fixation cross

514 against a black screen, with the phonemes presented 250 ms after the appearance of the fixation

515 cross. The visual-only trial began with the appearance of a female actor's face on the screen, with

516 lip movements beginning 250 ms after face onset. After the presentation of each auditory-only or

517 visual-only trial, subjects were presented with 3 options (/fa/, /ka/, and /ma/) and were instructed

518 to press one of three associated buttons on an MRI-safe button response box.

519   The first 24 subjects were shown response choices that always appeared in the same order

520 (/fa/, /ka/, or /ma/) with a stable mapping between response choice and button (the index finger

521 was always used to make the response for /fa/, the middle finger for /ka/ and the ring finger for

522 /ma/). While performing the sample size estimates for our power analysis, we saw that the stable

523 mapping between response choices and button presses resulted in response type differentiability

524 in the motor cortex consistent with prior evidence for motor regions encoding information about

525 finger movements (57). Hence, to counteract this effect and to negate the confounds of motor

526 region responses during speech perception (58), we altered the pre-registered protocol for the

527 remaining 40 subjects, who were shown response choices that were randomized after each trial.

528 Group-level comparisons of MVPA data between these two sub-groups revealed significant

529 differences only in the hand area of left M1/S1 for both the auditory-only and visual-only

530 conditions.

531   Subjects had 1.25 seconds to respond to the answer choices. If the subject failed to register

532 a response within 1.25 seconds, the trial was recorded as a missed response. Every trial was

533 followed by a 5-6 second jitter period (sampled from a uniform random distribution) which acted

534 as the intertrial interval (ITI) (59). In each run, subjects completed 60 trials that were split between

30 auditory-only and 30 visual-only trials, with 10 trials each for every phoneme and viseme; trial

types and stimuli were randomly intermixed in each run.

In total, subjects completed five runs, resulting in 300 trials in total (150 phonemes, 150

visemes) during the task, with each run lasting 8 minutes and 30 seconds. Psychtoolbox was used

for stimulus delivery and recording timing information and subject responses. Auditory stimuli

were presented using fMRI compatible Avotec headphones that had integrated earmuffs in order

to achieve maximum reduction of scanner noise. The sound level of stimuli was held constant for

all subjects. While presenting auditory speech stimuli in an MRI scanner can be challenging, the

undegraded nature of the auditory stimuli enabled near perfect accuracy throughout the task. A

mirror system reflected the visual stimuli from an LCD projector onto a mirror (width of the mirror:

12cm, approximate viewing distance between eye and mirror: 15cm; width and height of the face

on screen: 9cm x 12cm) located inside the magnet bore of the scanner.

Data Exclusion Criteria

To ensure that subjects included in analyses demonstrated persistent attention throughout

the task, we pre-registered exclusion criteria to remove subjects with behavioral accuracy rates

less than 75% for either auditory-only or visual-only conditions: no subjects were excluded based

on this cutoff.

fMRI Data Collection

Subjects were scanned in a GE Discovery MR750 3.0 Tesla scanner with a Nova 32

channel standard adult-sized coil (Milwaukee, WI). One high-resolution T1-weighted structural

image was obtained for each subject that was used in preprocessing, flip angle = 8, FOV = 25.6

mm, slice thickness = 1 mm, 256 slices. Then, for each of the five runs, functional T2*-weighted

BOLD images were obtained using a multiband gradient-echo, echo planar imaging sequence with

a resolution of 2.4 x 2.4 x 2.4 mm^3, TR of 800 ms and, TE of 30 ms, Flip Angle of 52, for a total

of 644 3D volumes of the whole brain with a FOV of 216 mm. To account for signal saturation,

the task did not start until the first 10 TRs were acquired and discarded by the scanner in each run.

Data Processing

fMRI data was reconstructed with realignment and fieldmap correction applied using

SPM12 to each of the five T2* runs for inhomogeneity recovery of signal in the B0 field.

Physiological noise was removed using RETROICOR (60). For both the univariate and

multivariate analysis, preprocessing steps were completed using SPM12 (Wellcome Department

of Cognitive Neurology, London, UK; https://www.fil.ion.ucl.ac.uk/spm/software/spm12/). We

utilized The Decoding Toolbox (https://sites.google.com/site/tdtdecodingtoolbox/; version 3.997)

for the whole-brain multivariate analyses.

Preprocessing

Before preprocessing the functional images, SPM's display tool was used to set the origin

of the anatomical volumes for each subject manually by picking the location of the anterior

commissure. After this, functional volumes were reconstructed and realigned, physiological noise

was removed, and field map correction was applied. This was followed by slice time correction to

account for acquisition time differences between slices for each of the whole brain functional

volumes. This data was then co-registered to the subject's anatomical space using a 4th degree B-

spline, followed by segmentation of the tissues from the anatomical image with a forward

deformation field. Information generated during the segmentation process was then used to

transform the co-registered functional volumes into the standard MNI anatomical space with

isotropic voxel volume dimensions of 2mm. The normalized data was then spatially smoothed

using a full-width half maximum (FWHM) kernel of 5mm.

Univariate Analyses

We performed a univariate, contrast-based analysis of auditory-only phonemes (averaged

across the 3 phonemes) and visual-only visemes (averaged across the 3 visemes) in order to

identify the regions that demonstrate significantly different activation patterns across stimulus

types. We utilized a canonical hemodynamic response function with event duration set to 2 seconds

for each of the phonemes (AuditoryFA + AuditoryKA + AuditoryMA) and visemes VisualFA +

VisualKA + VisualMA) and 5.5 seconds for the fixation periods (Fixation). Event onsets times

were defined as the moment when the fixation cross (for auditory trials) or face (visual trials)

appeared on the screen.

In the first level analysis, whole brain beta maps were generated individually for all seven

conditions for each of the 64 subjects. These maps also included information from regressors for

motion correction (six head movement parameters). Freesurfer's group-analysis pipeline was used

for second level analyses (61). Specifically, each subject's data was projected onto the cortical

surface of the fsaverage subject (using the command mris_preproc) and smoothed using a FWHM

of 10mm (using the command mri_surf2surf). General linear models were estimated with the

command mri_glmfit separately for each hemisphere and condition, excluding motor and frontal

areas due to the initial $n = 24$ subjects with consistent phoneme-motor mappings (analyses

inclusive of all regions are shown in Supp. Fig. 1). Significant vertices were identified at the group

level using the command mri_glmfit-sim using a vertex level threshold of $p < .001$ and cluster-

level threshold of $p < .05$ (estimated with 10000 permutations) to control for multiple comparisons;

p-values were adjusted for separate tests of the two hemispheres. All tests were two-tailed.

Univariate time-series were calculated from the individual subject volumes following

fMRI realignment, unwarping with field map correction applied, and slice-timing correction.

Volume data were epoched relative to phoneme or viseme onset for each individual trial, and the

fourth volume (2.4 seconds post-stimulus) was used as a baseline for each trial. To quantify the

relative differences in timing between phonemes and visemes, data were additionally normalized

so that data in each condition for each subject ranged from 0-1.

Multivariate Analyses

To identify regions that reliably differentiated classes of phonemes and classes of visemes,

we performed searchlight based MVPA analyses. Preprocessing steps for univariate and

multivariate analyses were matched except for the normalization and smoothing, such that for the

multivariate analysis, these two steps were performed after the first level analysis was completed.

For the decoding analysis, we utilized The Decoding Toolbox (62) with a LIBSVM (63) based

support vector machine (SVM) implementation. For each of the individual subjects, we built a

SVM classifier with a cross-validation scheme for the five runs. We used these classifiers to build

two separate models: one to classify between the three phonemes and the other to classify between

the three visemes. The phoneme models were constructed to identify voxels that reliably decoded

the identity of each of the three phonemes while the viseme models were built to identify voxels

that reliably decoded the identity for each of the three visemes. These models were implemented

as independent whole-brain searchlight analysis in the first level of the MVPA model. For each of

the models, beta estimates were calculated and extracted from a 3-voxel radius sphere. 4 fMRI

runs were used for training and 1 run for testing in an iterative manner. The searchlight center was

shifted through voxel-wise patterns throughout the brain to extract whole-brain accuracy maps for

auditory-only and visual-only conditions. Chance-level accuracy (33.3%) was subtracted from

individual subjects and conditions so that null-hypothesis values could be set to zero. Group-level

analyses and multiple comparison corrections were performed using Freesurfer and matched those

in the Univariate Analyses.

ROI-Based Decoding Analyses

Following the whole-brain searchlight analysis, we selected five regions of interests from

each hemisphere (ROI) based on results from literature (24, 43, 44). Four ROIs (STG, pSTS,

fusiform, and hMT+) were pre-registered. The fifth ROI (V1/V2) was included in the classification

analyses given the strong univariate response in the visual-only condition. ROIs were identified at

the individual subject level based on Freesurfer aparc-aseg labeling (64). Selected labels included

'superiortemporal', 'bankssts','MT_exvivo.thresh', the combined labels 'FG1.mpm.vpnl' to

'FG4.mpm.vpnl', and the combined labels 'V1_exvivo.thresh' and 'V2_exvivo.thresh'. Contrast

beta estimates (condition vs fixation) were extracted for each subject, stimulus (6 phonemes and

visemes), block, and ROI. SVM analyses were performed at the individual subject level with

models trained on n-1 blocks (leave-one-out classification) using the 'fitcecoc' function in

MATLAB.

IEEG Experiment

The study was approved by the Institutional Review Boards (IRB) at the University of

Michigan and Henry Ford Hospitals.

Subjects and Recordings

$N = 6$ patients (2 female, 4 male) undergoing clinical evaluation using iEEG for intractable

epilepsy consented to participate in this study under an institutional review board (IRB) approved

protocol at the University of Michigan or Henry Ford hospital. Patients' ages ranged from 12-39

years of age (mean = 29.7, std = 9.8) and 5 were right-handed (one patient self-reported to be

ambidextrous). All patients were native English speakers. Clinically implanted depth electrodes (5

mm center-to-center spacing) and/or subdural electrodes (10 mm center-to-center spacing) were

used to acquire iEEG data from subjects. IEEG data from a total of 459 electrodes were recorded

from the six subjects. The type and location of electrodes implanted were based on the clinical

needs of the patients. Electrodes were implanted within left auditory areas for 2 patients and right

auditory areas for 4 patients. IEEG recordings were acquired at either 4096 Hz ($n = 4$ patients) or

1000 Hz ($n = 2$ patients) due to differences in clinical amplifiers.

MRI and CT Acquisition and Processing

Preoperative T1-weighted magnetic resonance imaging (MRI) and postoperative computer

tomography (CT) scans were acquired for all subjects. The preoperative T1 MRI was registered to

the postoperative CT using SPM12 using the 'mutual information' method (65). The CT was not

resliced or resampled. The localization of each electrode was performed using custom software

(66). The algorithm works by identifying and segmenting electrodes from the CT image based on

gray scale intensity, and projects subdural electrodes to the dura surface using the shape of the

electrode disk to counteract post-operative compression. For all subsequent analyses including

reconstruction of cortical surfaces, volume segmentation and anatomical labeling, the Freesurfer

image analysis suite was utilized (http://surfer.nmr.mgh.harvard.edu/ (67, 68)).

Task and Stimuli

Subjects were tested at their bedside in an Epilepsy Monitoring Unit using a laptop running

Psychtoolbox (69). The task paradigm was adapted from a prior study (70) which was designed to

behaviorally study multiple aspects of auditory-visual speech integration. The stimuli consisted of

a female speaker who produced 40 commonly used 1-2 syllable words that each started with one

of the four consonants: *'b', 'f', 'g', 'd'* (10 of each). The phoneme in the second position of each

of these words was generally balanced across each of the four groups. Each stimulus was recorded

at a frame rate of 29.97 frames per second, and trimmed to 1100 ms in length. Further adjustments

were made such that the first consonantal burst of each word occurred at 500 ms during the video

playback by removing leading video frames.

Each subject underwent two task variants using the same stimuli and task design to increase

trial numbers, and to reduce classifier overfitting. Supp. Fig. 6 shows the task schematic for both

variants of the task. In variant one, subjects were presented with words one at a time, in one of two

main conditions: auditory-only or visual-only. Subjects then identified the initial speech sound of

the presented stimulus using a button press to select one of four options shown on the computer

screen. For example, on a trial with the word *"bag"*, the options presented to the subject were *'b',*

'g', 'd', 'th'. The paradigm included 40 trials per consonant in each main condition, such that each

of the 40 words were presented 4 times in the visual-only condition and another 4 times in the

auditory-only condition. This resulted in a total of 320 trials for each subject using task variant 1.

The words used in our task are presented in Supplemental Table 6.

In task variant 2, subjects were presented with trials in one of four main conditions:

auditory-only, visual-only, congruent audiovisual, or incongruent audiovisual. Task stimuli and

instructions were the same as in variant 1. Variant 2 included 20 trials per consonant in each main

condition. A second factor that was manipulated in this variant was the background noise level of

the stimuli such that half of the words used in each condition were presented in either a low noise

or a high noise context. In the low noise context, the auditory stimuli were presented as they were

recorded (SNR = 32.2 dB SPL). In the high noise context, pink noise was added to reduce the

signal-to-noise (SNR) ratio of the signals to -6 db SPL. Data were combined across the low and

high noise contexts to increase the number of trials available for classification analyses. Only data

from the auditory-only and visual-only conditions were included in analyses because they matched

the main conditions obtained from Task variant 1 and as the auditory-visual conditions contained

too few trials to be used in the classification analyses.

A total of 480 trials (Task variant 1: 320 trials, task variant 2: 160 trials) with 60 trials for

each consonant (*'b', 'g', 'd', 'f'*) per condition was obtained from the combined data of both task

variants. Each subject received a randomized trial order. For the auditory-only condition, a gray

rectangle was presented 500 ms before sound onset. Stimuli offset occurred 600 ms after sound

onset time. In the visual-only condition, face onset occurred 500 ms before the time when phoneme

onset would naturally occur. A wait time of 1.25 seconds was provided for the subjects to respond

to each of the stimuli.

IEEG Data Preprocessing

Data were preprocessed using bipolar referencing, such that signals from adjacent

electrodes were subtracted in a pairwise manner. This ensured that the final signals of interest were

obtained from neuronal populations that provided maximal localized responses (71). Analyses in

auditory regions were restricted to electrodes (registered in MNI space) that were within 10 mm

of the Freesurfer anatomical labels 'superiotemporal', 'middletemporal' or 'supramarginal'.

Excessively noisy electrodes were removed either manually or statistically by identifying

electrodes with raw signals that were 5 SD greater in comparison to all other electrodes. For

complementary analyses in visual regions, electrode locations were anatomically restricted to the

'inferiortemporal' and 'fusiform' labels.

Drift was removed from each channel (using residuals from fits to a 3rd order polynomial

and high-pass filtering at 0.1 Hz). Power-line interference was removed by notch-filtering at 60

Hz and its harmonics. ERPs were extracted from this minimally processed signal. HGp activity

was extracted from the continuous time-series after wavelet convolution and power transformation

(70-150 Hz in 5 Hz intervals, wavelet cycles = 20 at 70 Hz, and increased linearly to maintain the

same wavelet duration across frequencies). ERP and HGp data were segmented into 2 second

epochs centered around speech onset time for a specific stimulus: trial onset was defined as the

point when the initial consonant burst occurred. All data were then resampled to 1000 Hz.

Electrodes from both the left and right hemispheres were projected into the left hemisphere

for analyses and visualization. This projection was performed by registering each subject's skull-

stripped brain to the Freesurfer cvs_avg35_inMN152 template image through affine registration

using the Freesurfer function 'mri_robust_register' (72). Right hemisphere electrode coordinates

were then reflected onto the left hemisphere across the sagittal axis.

Classifiers for Calculating Decoding Accuracy

A support vector machine (73) classifier was utilized for calculating decoding accuracy.

Classifiers for stimulus trials were built for individual subjects and group-level analyses were

performed by combining results from individual subjects (subject as a random effect).

Classification was performed on downsampled data (10 Hz except where stated otherwise) to

reduce dimensional complexity. Phonemes and visemes were classified using a 4-fold multiclass

classifier from 0 to 500 ms following sound onset time (or the corresponding point in the visual

movie); electrodes and time-points were treated as dimensions in the classification of individual

trials. Time-series analyses were performed independently at each time-point (1000 Hz) and

accuracies were smoothed at the individual subject-level across 50 time points using the Matlab

function 'movmean'. Electrode-level analyses were performed on individual electrodes located in

auditory regions.

Similarity Analysis

To test whether auditory cortex showed a similar representation for phonemic and visemic

information, we examined the similarity of the phoneme and viseme confusion matrices.

Specifically, we paired each of the 16 cells in the two confusion matrices and used Pearson

correlation to examine their relationship. Significance was calculated by randomly permuting the

stimulus labels of each trial and repeating the full classification analysis $n = 1000$ times.

Calculating Individual Subject Classification Significance

The four classes tested within each condition yielded chance levels of classification at 25%.

To calculate significance above this chance level, we used binomial statistics for within-subject

significance testing (74, 75). We used the 'binocdf' function in MATLAB for this, by considering

two parameters: the number of trials, and probability of success at each instance (25%). This gives

rise to a binomial chance-level probability that varies depending on the number of data points used

for classification in each of the models that were built. This resulted in a chance probability of

29.58% ($p = 0.05$) for a 4-class classifier with 240 trials.

References

1. J. Plass, D. Brang, S. Suzuki, M. Grabowecky, Vision perceptually restores auditory spectral dynamics in speech. *Proc Natl Acad Sci U S A* **117**, 16920-16927 (2020).
2. C. Micheli *et al.*, Electrocorticography reveals continuous auditory and visual speech tracking in temporal and occipital cortex. *Eur J Neurosci* **51**, 1364-1376 (2020).
3. L. A. Ross, D. Saint-Amour, V. M. Leavitt, D. C. Javitt, J. J. Foxe, Do you see what I am saying? Exploring visual enhancement of speech comprehension in noisy environments. *Cerebral cortex* **17**, 1147-1153 (2007).
4. S. Rosemann, C. M. Thiel, Audio-visual speech processing in age-related hearing loss: Stronger integration and increased frontal lobe recruitment. *Neuroimage* **175**, 425-437 (2018).
5. A. A. Aabedi *et al.*, Convergence of heteromodal lexical retrieval in the lateral prefrontal cortex. *Sci Rep* **11**, 6305 (2021).
6. D. Caplan, D. Gow, N. Makris, Analysis of lesions by MRI in stroke patients with acoustic-phonetic processing deficits. *Neurology* **45**, 293-298 (1995).
7. J. T. Crinion, E. A. Warburton, M. A. Lambon-Ralph, D. Howard, R. J. Wise, Listening to narrative speech after aphasic stroke: the role of the left anterior temporal lobe. *Cereb Cortex* **16**, 1116-1125 (2006).
8. N. Kraus *et al.*, The neural legacy of a single concussion. *Neurosci Lett* **646**, 21-23 (2017).
9. E. C. Thompson *et al.*, Difficulty hearing in noise: a sequela of concussion in children. *Brain Inj* **32**, 763-769 (2018).
10. C. A. Anderson, I. M. Wiggins, P. T. Kitterick, D. E. Hartley, Adaptive benefit of cross-modal plasticity following cochlear implantation in deaf adults. *Proceedings of the National Academy of Sciences* **114**, 10256-10261 (2017).
11. C. Chandrasekaran, A. Trubanova, S. Stillittano, A. Caplier, A. A. Ghazanfar, The natural statistics of audiovisual speech. *PLoS computational biology* **5**, e1000436 (2009).
12. C. G. Fisher, Confusions among visually perceived consonants. *Journal of speech and hearing research* **11**, 796-804 (1968).
13. G. A. Calvert *et al.*, Activation of auditory cortex during silent lipreading. *science* **276**, 593-596 (1997).
14. J. Pekkola *et al.*, Primary auditory cortex activation by visual speech: an fMRI study at 3 T. *Neuroreport* **16**, 125 (2005).
15. G. Karthik *et al.*, Visual speech differentially modulates beta, theta, and high gamma bands in auditory cortex. *Eur J Neurosci* **54**, 7301-7317 (2021).
16. R. Thézé, A.-L. Giraud, P. Mégevand, The phase of cortical oscillations determines the perceptual fate of visual cues in naturalistic audiovisual speech. *Science advances* **6**, eabc6348 (2020).
17. P. Mégevand *et al.*, Crossmodal phase reset and evoked responses provide complementary mechanisms for the influence of visual speech in auditory cortex. *Journal of Neuroscience* **40**, 8530-8542 (2020).
18. N. Mesgarani, C. Cheung, K. Johnson, E. F. Chang, Phonetic feature encoding in human superior temporal gyrus. *Science* **343**, 1006-1010 (2014).
19. M. K. Leonard *et al.*, Large-scale single-neuron speech sound encoding across the depth of human cortex. *Nature*, 1-10 (2023).

20. E. Formisano, F. De Martino, M. Bonte, R. Goebel, " Who" is saying" what"? Brain-based decoding of human voice and speech. *Science* **322**, 970-973 (2008).

21. E. F. Chang *et al.*, Categorical speech representation in human superior temporal gyrus. *Nat Neurosci* **13**, 1428-1432 (2010).

22. R. D. Raizada, F. M. Tsao, H. M. Liu, P. K. Kuhl, Quantifying the adequacy of neural representations for a cross-language phonetic discrimination task: prediction of individual differences. *Cereb Cortex* **20**, 1-12 (2010).

23. M. K. Leonard, M. O. Baud, M. J. Sjerps, E. F. Chang, Perceptual restoration of masked speech in human cortex. *Nat Commun* **7**, 13619 (2016).

24. M. S. Beauchamp, A. R. Nath, S. Pasalar, fMRI-guided transcranial magnetic stimulation reveals that the superior temporal sulcus is a cortical locus of the McGurk effect. *The Journal of Neuroscience* **30**, 2414-2417 (2010).

25. A. R. Nath, M. S. Beauchamp, Dynamic changes in superior temporal sulcus connectivity during perception of noisy audiovisual speech. *J Neurosci* **31**, 1704-1714 (2011).

26. N. Kilian-Hütten, G. Valente, J. Vroomen, E. Formisano, Auditory cortex encodes the perceptual interpretation of ambiguous sound. *Journal of Neuroscience* **31**, 1715-1720 (2011).

27. M. Bonte, L. Hausfeld, W. Scharke, G. Valente, E. Formisano, Task-dependent decoding of speaker and vowel identity from auditory cortical response patterns. *Journal of Neuroscience* **34**, 4548-4557 (2014).

28. N. Mesgarani, E. F. Chang, Selective cortical representation of attended speaker in multi-talker speech perception. *Nature* **485**, 233-236 (2012).

29. J. G. Makin, D. A. Moses, E. F. Chang, Machine translation of cortical activity to text with an encoder–decoder framework. *Nature neuroscience* **23**, 575-582 (2020).

30. J. V. Haxby, A. C. Connolly, J. S. Guntupalli, Decoding neural representational spaces using multivariate pattern analysis. *Annual review of neuroscience* **37**, 435-456 (2014).

31. N. Kriegeskorte, R. Goebel, P. Bandettini, Information-based functional brain mapping. *Proc Natl Acad Sci U S A* **103**, 3863-3868 (2006).

32. J. S. Arsenault, B. R. Buchsbaum, Distributed neural representations of phonological features during speech perception. *Journal of Neuroscience* **35**, 634-642 (2015).

33. J.-L. Schwartz, C. Savariaux, No, there is no 150 ms lead of visual speech on auditory speech, but a range of audiovisual asynchronies varying from small audio lead to large audio lag. *PLoS Computational Biology* **10**, e1003743 (2014).

34. C. E. Schroeder, P. Lakatos, Y. Kajikawa, S. Partan, A. Puce, Neuronal oscillations and visual amplification of speech. *Trends Cogn Sci* **12**, 106-113 (2008).

35. L. H. Arnal, B. Morillon, C. A. Kell, A. L. Giraud, Dual neural routing of visual facilitation in speech processing. *J Neurosci* **29**, 13445-13453 (2009).

36. J. Besle, A. Fort, C. Delpuech, M. H. Giard, Bimodal speech: early suppressive visual effects in human auditory cortex. *European Journal of Neuroscience* **20**, 2225-2234 (2004).

37. A. M. Bastos *et al.*, Visual areas exert feedforward and feedback influences through distinct frequency channels. *Neuron* **85**, 390-401 (2015).

38. L. E. Bernstein, J. Jiang, D. Pantazis, Z.-L. Lu, A. Joshi, Visual phonetic processing localized using speech and nonspeech face gestures in video and point-light displays. *Human Brain Mapping* **32**, 1660-1676 (2011).

860   39.   M. Ozker, I. M. Schepers, J. F. Magnotti, D. Yoshor, M. S. Beauchamp, A double
861   dissociation between anterior and posterior superior temporal gyrus for processing
862   audiovisual speech demonstrated by electrocorticography. *Journal of cognitive*
863   *neuroscience* **29**, 1044-1060 (2017).

864   40.   P. J. Karas *et al.*, The visual speech head start improves perception and reduces superior
865   temporal cortex responses to auditory speech. *Elife* **8**, e48116 (2019).

866   41.   R. Gao, E. J. Peterson, B. Voytek, Inferring synaptic excitation/inhibition balance from
867   field potentials. *Neuroimage* **158**, 70-78 (2017).

868   42.   S. Ray, N. E. Crone, E. Niebur, P. J. Franaszczuk, S. S. Hsiao, Neural correlates of high-
869   gamma oscillations (60–200 Hz) in macaque local field potentials and their potential
870   implications in electrocorticography. *Journal of Neuroscience* **28**, 11526-11536 (2008).

871   43.   M. S. Beauchamp, B. D. Argall, J. Bodurka, J. H. Duyn, A. Martin, Unraveling
872   multisensory integration: patchy organization within human STS multisensory cortex.
873   *Nature neuroscience* **7**, 1190-1192 (2004).

874   44.   H. G. Yi, M. K. Leonard, E. F. Chang, The encoding of speech sounds in the superior
875   temporal gyrus. *Neuron* **102**, 1096-1110 (2019).

876   45.   M. McGrath, Q. Summerfield, Intermodal timing relations and audio-visual speech
877   recognition by normal-hearing adults. *The Journal of the Acoustical Society of America* **77**,
878   678-685 (1985).

879   46.   A. R. Nidiffer, C. Z. Cao, A. O'Sullivan, E. C. Lalor, A linguistic representation in the
880   visual system underlies successful lipreading. *bioRxiv* (2021).

881   47.   L. Zhang, Y. Du, Lip movements enhance speech representations and effective
882   connectivity in auditory dorsal stream. *NeuroImage*, 119311 (2022).

883   48.   N. K. Logothetis, What we can do and what we cannot do with fMRI. *Nature* **453**, 869-
884   878 (2008).

885   49.   M. Jueptner, C. Weiller, does measurement of regional cerebral blood flow reflect synaptic
886   activity? Implications for PET and fMRI. *Neuroimage* **2**, 148-156 (1995).

887   50.   S. M. Epp, G. Castrillon, J. Andrews-Hanna, C. Preibisch, V. Riedl, Two distinct modes
888   of hemodynamic responses in the human brain. *bioRxiv*, 2023.2012. 2008.570806 (2023).

889   51.   S. Ray, E. Niebur, S. S. Hsiao, A. Sinai, N. E. Crone, High-frequency gamma activity (80–
890   150 Hz) is increased in human cortex during selective attention. *Clinical Neurophysiology*
891   **119**, 116-133 (2008).

892   52.   H. Blank, M. H. Davis, Prediction errors but not sharpened signals simulate multivoxel
893   fMRI patterns during speech perception. *PLoS biology* **14**, e1002577 (2016).

894   53.   A. Nair, D. Brang, Inducing synesthesia in non-synesthetes: Short-term visual deprivation
895   facilitates auditory-evoked visual percepts. *Consciousness and cognition* **70**, 70-79 (2019).

896   54.   F. Bröhl, A. Keitel, C. Kayser, MEG activity in visual and auditory cortices represents
897   acoustic speech-related information during silent lip reading. *eneuro* **9** (2022).

898   55.   N. Mesgarani, C. Cheung, K. Johnson, E. F. Chang, Phonetic feature encoding in human
899   superior temporal gyrus. *Science* **343**, 1006-1010 (2014).

900   56.   H. McGurk, J. MacDonald, Hearing lips and seeing voices. *Nature* **264**, 746-748 (1976).

901   57.   G. Shen *et al.*, Decoding the individual finger movements from single-trial functional
902   magnetic resonance imaging recordings of human brain activity. *European Journal of*
903   *Neuroscience* **39**, 2071-2082 (2014).

904   58.   S. M. Wilson, A. P. Saygin, M. I. Sereno, M. Iacoboni, Listening to speech activates motor
905   areas involved in speech production. *Nature neuroscience* **7**, 701-702 (2004).

59. D. Zeithamova, M.-A. de Araujo Sanchez, A. Adke, Trial timing and pattern-information
analyses of fMRI data. *Neuroimage* **153**, 221-231 (2017).
60. G. H. Glover, T. Q. Li, D. Ress, Image-based method for retrospective correction of
physiological motion effects in fMRI: RETROICOR. *Magnetic Resonance in Medicine:*
An Official Journal of the International Society for Magnetic Resonance in Medicine **44**,
162-167 (2000).
61. D. J. Hagler Jr, A. P. Saygin, M. I. Sereno, Smoothing and cluster thresholding for cortical
surface-based group analysis of fMRI data. *Neuroimage* **33**, 1093-1103 (2006).
62. M. N. Hebart, K. Görgen, J.-D. Haynes, The Decoding Toolbox (TDT): a versatile software
package for multivariate analyses of functional imaging data. *Frontiers in*
neuroinformatics **8**, 88 (2015).
63. C.-C. Chang, C.-J. Lin, LIBSVM: a library for support vector machines. *ACM transactions*
on intelligent systems and technology (TIST) **2**, 1-27 (2011).
64. R. S. Desikan *et al.*, An automated labeling system for subdividing the human cerebral
cortex on MRI scans into gyral based regions of interest. *Neuroimage* **31**, 968-980 (2006).
65. P. Viola, W. M. Wells III, Alignment by maximization of mutual information.
International journal of computer vision **24**, 137-154 (1997).
66. D. Brang, Z. Dai, W. Zheng, V. L. Towle, Registering imaged ECoG electrodes to human
cortex: A geometry-based technique. *J Neurosci Methods* **273**, 64-73 (2016).
67. A. M. Dale, B. Fischl, M. I. Sereno, Cortical surface-based analysis: I. Segmentation and
surface reconstruction. *NeuroImage* **9**, 179-194 (1999).
68. B. Fischl, M. I. Sereno, A. M. Dale, Cortical surface-based analysis: II: Inflation, flattening,
and a surface-based coordinate system. *NeuroImage* **9**, 195-207 (1999).
69. M. Kleiner, D. Brainard, D. Pelli, What's new in Psychtoolbox-3? (2007).
70. L. A. Ross *et al.*, Impaired multisensory processing in schizophrenia: deficits in the visual
enhancement of speech comprehension under noisy environmental conditions. *Schizophr*
Res **97**, 173-183 (2007).
71. D. Yao *et al.*, Which reference should we use for EEG and ERP practice? *Brain topography*
32, 530-549 (2019).
72. M. Reuter, H. D. Rosas, B. Fischl, Highly accurate inverse consistent registration: a robust
approach. *Neuroimage* **53**, 1181-1196 (2010).
73. C. Cortes, V. Vapnik, Support-vector networks. *Machine learning* **20**, 273-297 (1995).
74. E. Demandt *et al.*, Reaching movement onset-and end-related characteristics of EEG
spectral power modulations. *Frontiers in neuroscience* **6**, 65 (2012).
75. E. Combrisson, K. Jerbi, Exceeding chance level by chance: The caveat of theoretical
chance levels in brain signal classification and statistical assessment of decoding accuracy.
Journal of neuroscience methods **250**, 126-136 (2015).
943