# Combinatorial interpretation of BMP and WNT allows BMP to act as a morphogen in time but not in concentration.

Elena Camacho-Aguilar[1], Sumin Yoon[1], Miguel A. Ortiz-Salazar[1], Aryeh Warmflash[1,2]*

[1]Department of Biosciences and [2]Department of Bioengineering, Rice University, Houston TX 77005
*author for correspondence: aryeh.warmflash@rice.edu

## Abstract

Secreted morphogen signals play a key role in the determination of cell fates during embryonic development. BMP signaling is essential for vertebrate gastrulation, as it initiates a cascade of signals that controls the self-organized patterning of the three germ layers. Although morphogen signals are typically thought to induce cell fates in a concentration-dependent manner, development is a highly dynamic process, so it is crucial to understand how time-dependent signaling affects cellular differentiation. Here we show that varying the duration of BMP signaling leads to either pluripotent, mesodermal, and extraembryonic states, while varying the concentration does not cause efficient mesodermal differentiation at any dose. Thus, there is a morphogen effect in time but not in concentration, and an appropriately timed pulse of BMP induces hPSCs to a mesodermal fate more efficiently that sustained signaling at any concentration. Using live cell imaging of signaling and cell fate reporters together with a simple mathematical model, we show that this effect is due to a combinatorial interpretation of the applied BMP signal and induced endogenous WNT signaling. Our findings have implications for how signaling pathways control the landscape of early human development.

## Introduction

In mammalian development, gastrulation is the first differentiation event of the embryo proper (Arnold and Robertson, 2009; Bardot and Hadjantonakis, 2020) where the pluripotent epiblast differentiates into the three germ layers of the embryo: ectoderm, mesoderm, and endoderm. This process is orchestrated by a cascade of signals initiated by BMP signaling from the extraembryonic tissues which triggers Wnt and Nodal signaling in the epiblast (Ben-Haim et al., 2006; Brennan et al., 2001; Tortelote et al., 2013). The ligands for all three pathways find their highest expression in the proximal posterior portion of the embryo, where the primitive streak emerges, defining the anterior-posterior axis of the embryo (Conlon et al., 1994; Liu et al., 1999; Winnier et al., 1995). However, how cell differentiation is combinatorially controlled by this cascade of signals at this stage is still not completely understood.

Technical and ethical issues prevent the study of these questions *in vivo*, especially in the human embryo. To overcome this problem, several *in vitro* models of mammalian gastrulation have been developed that can serve as proxies for investigating these stages of development (Camacho-Aguilar and Warmflash, 2020; Fu et al., 2021; Heemskerk, 2020; Shahbazi et al., 2019). In particular, we and others have shown that human pluripotent stem cells (hPSCs) grown in colonies of precise size and shape using micropatterning technology and treated with the upstream gastrulation-inducing signal BMP4 pattern into all three germ layers plus a fourth one transcriptionally similar to extraembryonic tissues which most likely represents the amnion (Chhabra and Warmflash, 2021; Chhabra et al., 2019; Etoc et al., 2016; Minn et al., 2020; Warmflash et al., 2014). While some studies have suggested that these fates may emerge from differential concentrations of BMP generated by diffusion and interactions with the inhibitor Noggin (Tewary et al., 2017), several studies contradict this idea.

First, in colonies of sizes of 10 cells or smaller, BMP acts as a switch controlling a cell state transition from pluripotency to extraembryonic as BMP concentration increases without inducing mesendodermal fates at any dose (Nemashkalo et al., 2017). In fact, primitive steak induction only takes place when the cell density is above a threshold, and it is blocked if either the Wnt or Nodal pathways are inhibited (Chhabra et al., 2019; Nemashkalo et al., 2017). Thus, secondary signals are necessary for BMP to induce differentiation into the three germ layers. However, much remains unknown about the underlying mechanisms.

Here we studied how cell fate decisions are controlled by BMP signaling during human pluripotent stem cell differentiation *in vitro*. We investigated how hPSCs interpret duration and concentration of the applied BMP4 signal. Our results unveiled a *morphogen effect* in time, where short, intermediate, and long pulses of BMP4 signaling result in cells adopting pluripotent, mesodermal, and extraembryonic states, respectively. In contrast, there was no comparable morphogen effect in concentration: varying the BMP concentration does not cause comparably efficient mesodermal differentiation at any dose. Using live cell imaging of signaling reporters we discovered that the temporal

morphogen effect is controlled by combinatorial interpretation of the exogenously supplied BMP4 and the endogenously induced WNT signaling. Taking advantage of mathematical modelling, we uncovered the minimal requirements for the logic that controls these cell fate decisions, providing a fate map that explains the mechanism that allows BMP to act as a morphogen in time but not in concentration. Our findings have implications for how signaling pathways control the landscape of early human development and highlight the importance of time in *in vitro* differentiation protocols.

## Results

**BMP signaling produces a morphogen effect in time but not in concentration**
As BMP induces extraembryonic fates directly but other fates through intermediates (Ben-Haim et al., 2006; Martyn et al., 2018; Nemashkalo et al., 2017), we speculated that the timescales for these processes may differ, and the duration of BMP treatment may be an important variable. Previous differentiation protocols in hPSCs also indicated that the timing of BMP treatment may influence the outcome (Zhang et al., 2008). We cultured hPSCs for two days, exposing them to varying durations of BMP4 treatment of a fixed concentration, 10 ng/mL, at the outset (Fig. 1A). We observed that cells that were exposed to a BMP pulse of 8 hours or less returned to the pluripotent or undifferentiated state, as marked by high SOX2, OCT4, and NANOG expression (Fig. 1B-D, S1A,B). On the other hand, BMP pulses longer than 32 hours were necessary to differentiate cells uniformly into an extraembryonic state, marked by high CDX2, ISL1, HAND1 and GATA3 expression (Fig. 1B-D, S1C-F). Strikingly, pulses of 16 hours showed a high level of mesoderm or primitive streak differentiation, marked by high BRA expression (Fig. 1B-D). These results unveiled a *morphogen effect* in the duration of BMP signaling, with short, intermediate, and long pulses resulting in pluripotent, mesodermal, and extraembryonic fates, respectively.

Several studies have suggested that for other pathways, including Sonic Hedgehog, duration and amplitude of signaling may be interchangeable, with shorter treatment with higher doses being equivalent to longer exposure to low doses (Dessaud et al., 2007; Sagner and Briscoe, 2017). If this was the case for BMP signaling in hPSCs, the high proportion of mesodermal cells obtained with a pulse of intermediate duration should be reproduced by treating hPSCs with a constant 2-day pulse of a lower BMP concentration. To test this, we compared the results of treating cells with 10 ng/mL BMP4 pulses of intermediate durations (ranging from 14 to 24 hours) (Fig. 2A,B), to the results obtained by inducing hPSCs with several constant concentrations (ranging from 1 to 4 ng/ml) for 48 hours (Fig. 2C,D). We observed that constant induction by any lower BMP4 concentration could not reproduce the high percentage of mesoderm cells obtained with 10ng/mL BMP4 for intermediate durations (Fig. 2A-D).

To further investigate the interplay between concentration and time of exposure to control hPSCs fate specification, we induced hPSCs with different pulses (increasing from 0 to 48 hours in 6 hours increments) of varying concentrations of BMP4 (ranging from 2 to 50

ng/mL) and observed the cell fates after 48 hours (Fig. S2A,B). Interestingly, when BMP4
containing media was withdrawn and replaced by fresh media, the data suggested that
for higher concentrations, shorter pulses were needed to induce mesoderm fates. While

**Figure 1. BMP4 signaling produces a morphogen effect in time.** (A) Schematic of the induction protocol. hPSCs are treated with a pulse of 10ng/mL BMP4 of varying durations (t*) after which the BMP-containing media is removed and replaced by the pluripotency maintainence media mTeSR1. hPSCs are always fixed and immunostained 48h after the onset of induction. (B) Example images of immunofluorescence for DAPI, SOX2, BRA, and CDX2 after the indicated BMP4 treatments. (C) Quantifications of cell fate proportions for the experimental conditions in (B). (D) From left to right, scatter plots of the quantifications of SOX2, BRA and CDX2 for the conditions mTeSR 0-48h, BMP 10ng/mL 0-16h followed by mTeSR 16-48h, and BMP 10ng/mL 0-48h, respectively. Each dot corresponds to a single cell, and its color marks the cell fate assigned to the cell as shown in (C). Scale bars: 100um.

this result might suggest a tradeoff between time and concentration, an alternative
possibility is that it could be due to a partial removal of the BMP4 ligand, with higher doses
being more difficult to completely remove compared to lower ones. To distinguish these
possibilities, we compared removing BMP ligand with or without adding the BMP inhibitor
Noggin to extinguish any remaining signal. Consistent with the idea that added BMP
cannot be removed by washing alone, even 3 mTeSR washes were insufficient to lower
the levels of differentiation to those observed by adding Noggin (Fig. S2C).
127
We therefore repeated the above experiment using a variety of different BMP durations
and concentrations with a modified protocol in which BMP containing media was replaced
by Noggin containing media.  Mesoderm induction peaked with a BMP4 pulse of around
30h for nearly all concentrations except for the lowest ones (Fig. 2E,F). The later peak at
lower concentrations is due to a slower induction of secondary signals necessary for
differentiation, as we show below (see Fig. S3E). Notably, even though the most efficient
duration of the BMP4 pulse was common for all concentrations above 3 ng/ml,
quantification showed that higher concentrations yielded higher proportions of BRA
positive cells. Taken together, the results suggest that concentration and time of exposure
are not interchangeable, and that efficient mesoderm induction after 48 hours of treatment
required a pulse of a high BMP4 concentration.
139
**BMP signal induces WNT in a bistable fashion**
In order to understand how signaling dynamics underlie our results on cell fate induction,
we took advantage of live cell reporters of signaling activity to measure the signaling
response in time under different treatment conditions. In particular, we first tracked the
BMP4 response by performing live imaging of hPSCs with a GFP::SMAD4 fusion in the
endogenous locus (Nemashkalo et al., 2017), and quantified the strength of the signaling
response as the ratio of nuclear and cytoplasmic SMAD4 intensity (Fig. 3A-D, S3A,D).
Consistent with our previous work (Heemskerk et al., 2019; Nemashkalo et al., 2017), a
sudden increase in BMP4 leads to a rapid translocation of SMAD4 into the nucleus in less
than 30 minutes. This response is sustained with a gradual decrease in activity over time
for the remaining 48 hours if BMP4 was kept in the media (Fig. 3D, S3A). Addition of
Noggin resulted in a rapid decrease in nuclear SMAD4 to baseline (Fig. 3D, S3A).
Consistent with our previous observations, removal of BMP4 without Noggin addition

resulted in a rapid decrease in nuclear SMAD4 but to a higher level than baseline,
indicating that some BMP4 remains in the media (Fig. 3D, S3A,D).
155
156 We then explored WNT/β-catenin response, the next member of the signaling cascade
157 that plays a role in gastrulation and mesoderm (Arnold and Robertson, 2009). We tracked
158 WNT response by performing live imaging of hPSCs with GFP fused to the N terminus of
159 endogenous β-catenin (Massey et al., 2019), and quantified the strength of the signaling
160 response as the mean nuclear β-catenin intensity (Fig. 3E-H). Strikingly, measurement of

**Figure 2. Duration and concentration of BMP4 signaling are not interchangeable. (A)** Example images of immunofluorescence for SOX2, BRA, and CDX2 in magenta, yellow, and cyan, respectively, after the indicated pulses of 10ng/mL BMP4 treatments followed by mTeSR. **(B)** Quantifications of the proportions of cells classified as BRA+ in the treatments shown in (A). **(C)** Example images of immunofluorescence for SOX2, BRA, and CDX2 in magenta, yellow, and cyan, respectively, after the treatments of the indicated constant concentrations of BMP4. **(D)** Quantifications of the proportions of cells classified as BRA+ in the treatments shown in (C). **(E)** Example images of immunofluorescence for SOX2, BRA, and CDX2 in magenta, yellow, and cyan, respectively, after the 48h treatments with pulses of the indicated times (columns) followed by Noggin, of the indicated concentrations (rows) of BMP. **(F)** Quantifications of the proportions of cells classified as BRA+ in the treatments shown in (E). Scale bars: 100um. Error bars in (B), (D), and (F) show the SEM.

161 Wnt dynamics revealed a bistable WNT/β-catenin response (Fig. 3H, S3B-C). If BMP was
162 withdrawn early, β-catenin levels converged to a low level in time. However, if BMP was
163 presented for sufficiently long, WNT activity became self-sustaining and remained stable
164 even after BMP withdrawal. This likely results from positive feedback in which WNT

**Figure 3. BMP signaling induces a bistable endogenous WNT. (A)** BMP signal is transduced through SMAD4 translocation to the nucleus. Noggin inhibits BMP signal by preventing it from binding to the receptors. **(B,C)** hPSCs expressing GFP::SMAD4 before BMP4 treatment (B), and 5 hours after treatment with BMP4 (C). **(D)** GFP::SMAD4 average nuclear:cytoplasmic intensity ratio after the treatments indicated at the bottom of the figure. Additional experimental conditions are shown in Fig. S3A. **(E)** Simplified canonical WNT/β-catenin pathway. IWP2 prevents canonical WNT secretion. **(F,G)** hPSCs expressing GFP::β-catenin before BMP treatment (F), and 40 hours after treatment with BMP4 (G). **(H)** GFP::β-catenin average nuclear intensity after the treatments indicated at the bottom of the figure. Additional experimental conditions are shown in Fig. S3B-C. Scale bars: 50um.

165 signaling activates its own ligands, as has been shown in other contexts (Wang et al.,
166 2012).
167
Comparing the signaling trajectories with the resulting fates reveals that only in cases
where BMP is repressed after having induced WNT to a high level are cells able to
differentiate to mesoderm (Fig. S3B,C). This suggests that a combinatorial effect between
BMP and BMP-induced, bistable Wnt signals underlies the observed morphogen effect in
time. Short BMP durations of high BMP concentration are not enough to activate WNT
and therefore cells remain pluripotent. On the other hand, if BMP is maintained for longer
than 32 hours, Wnt signaling is induced but is insufficient to override the effects of BMP
signaling, and cells become extraembryonic. It is in a window of middle durations, where
BMP is maintained for long enough to induce Wnt autoactivation but is then suppressed,
that cells are exposed to a low BMP, high Wnt signaling profile, and mesoderm
differentiation is obtained.
179
We also investigated how WNT/$\beta$-catenin response depended on cell density, as previous
studies have shown that cell density is critical for BMP signal to induce mesodermal fates
(Nemashkalo et al., 2017). Consistent with these studies, we observed that if we reduced
the number of cells seeded from 40K cells/cm$^2$ to 15K cells/cm$^2$, pulses of BMP of high
concentration could no longer induce mesoderm (Fig. S4A-D). To understand whether
the lack of mesoderm correlated with reduced WNT signaling, we compared the WNT
response of cells seeded in a density of 40K cells/cm$^2$ (high density) versus cells seeded
in a density of 15K cells/cm$^2$ (low density) by performing live imaging of the $\beta$-catenin
reporter cells (Fig. S4E). Indeed, we observed that the WNT response under BMP
treatment was lower at low densities. While BMP induced a high WNT response in cells
seeded at high density, the WNT response of cells seeded at low density under the same
BMP treatment was comparable to non-treated cells at high density (Fig. S4E). Moreover,
treating cells with IWP2 at the time of BMP treatment to inhibit the secretion of
endogenous Wnt ligands suppressed mesoderm induction (Fig. S5). Finally, we
investigated how the WNT response depended on BMP concentration, and observed that
the rate of increase of nuclear $\beta$-catenin increased as the BMP dose increased (Fig. S3E).
Taken together, these results suggest that BMP induces a bistable endogenous WNT
signal in a concentration dependent manner, that is necessary for mesoderm induction.
The mechanism behind this process will be discussed further below.
199
**A simple mathematical model reproduces the observed WNT bistability**
To better understand the observed WNT bistability, we developed a minimal
mathematical model that recapitulated the signaling dynamics described above. This
model is comprised of two sub-systems of differential equations that model the BMP and
Wnt responses, respectively, and which are connected via BMP4 activation of Wnt, which
reflects the transcriptional activation of *WNT3* by BMP signaling (Fig. 4A-B) (Kurek et al.,
2015).
207

**Figure 4. BMP-induced WNT bistability is reproduced by a simple mathematical model. (A)** Schematic of the network regulating BMP and WNT response. **(B)** System of ODEs that model the network dynamics in (A). **(C)** Simulations of BMP and WNT dynamics under the indicated BMP4 treatments. Media change is also simulated at 24 hours for all conditions. BMP = BMP4 ligand. BMPR = BMP4 receptors. SMAD4 = Nuclear SMAD4. Nog = Noggin. WNT = WNT ligands. βCat = Nuclear β-catenin. IWP2 = IWP2.

The equations in (B) are:

$$\frac{d\text{BMP}}{dt} = \frac{1}{\tau}(-(1-\text{Nog})\,b_{aff}\text{BMP} \times \text{BMPR} + \gamma\text{SMAD4})$$

$$\frac{d\text{BMPR}}{dt} = \frac{1}{\tau}(\beta - (\delta_I + (1-\text{Nog})\,b_{aff}\text{BMP}) \times \text{BMPR} + \gamma\text{SMAD4})$$

$$\frac{d\text{SMAD4}}{dt} = \frac{1}{\tau}((1-\text{Nog})\,b_{aff}\text{BMP} \times \text{BMPR} - (\gamma+\delta)\text{SMAD4})$$

$$\frac{d\text{WNT}}{dt} = (1-\text{IWP2})\left(a_{SW}\frac{\text{SMAD4}^{nsw}}{K_{SW}^{nsw} + \text{SMAD4}^{nsw}} + a_{WW}\frac{\beta\text{Cat}^{nww}}{K_{WW}^{nww} + \beta\text{Cat}^{nww}}\right) - \delta_W\text{WNT}$$

$$\frac{d\beta\text{Cat}}{dt} = a_{W\beta}\frac{(\text{WNT} + a_{C\beta}\text{CHIR})^{nw\beta}}{K_{W\beta}^{nw\beta} + (\text{WNT} + a_{C\beta}\text{CHIR})^{nw\beta}} - \delta_\beta\beta\text{Cat}$$

We simulated the BMP response using the model proposed in (Heemskerk et al., 2019), where binding of BMP ligands to the receptor complex activates SMAD4 but enhances the degradation of the BMP receptor (Fig. 4A-B). The effect of the BMP inhibitor Noggin is modelled by a direct inhibition of BMP binding to free receptors (Fig. 4A-B). BMP withdrawal without Noggin addition was modelled by multiplying the original BMP concentration by a constant that represented the percentage of BMP that remained in the culture, as observed in Fig. 3 and S3. With this model, features of SMAD4 dynamics observed in the data such as fast SMAD4 upregulation upon BMP4 treatment, rapid SMAD4 downregulation by Noggin, and slow decay were reproduced (Fig. 4C, S6A,B).

The second sub-system models the Wnt response. Wnt expression is activated by SMAD4 and activates nuclear β-catenin. β-catenin in turn further activates Wnt signaling. The need to accumulate sufficient Wnt protein before β-catenin becomes active explains the delay in the rise in β-catenin activity observed experimentally (see Fig. 3H). Details of this model can be found in the Supplemental Material.

The model comprised by the two-subsystems could reproduce the observed BMP-induced WNT bistability (Fig. 4C, S6). Short BMP pulses allowed the β-catenin variable to remain in a low steady state, while longer BMP pulses resulted in the β-catenin variable converging to a high steady state where it remained even in the absence of BMP (Fig. 4C). The model also reproduced the signaling dynamics obtained under BMP withdrawal without Noggin and the BMP concentration-dependent increase of WNT signaling (Fig. S6). Interestingly, as in the experimental data, while a 16h pulse of BMP followed by

Noggin inhibition resulted in β-catenin returning to a low value, a 16h pulse of BMP
followed by withdrawal without Noggin resulted in β-catenin auto-activating to a high level
by 48 hours (Fig. S3A-C, S6A,C). Thus, modeling supports the idea that residual BMP in
the media explains why a 16h pulse of BMP 10 ng/mL is sufficient to induce mesoderm if
the BMP containing media is replaced by media alone, while longer pulses are needed if
Noggin is used.
238
**Mesoderm induction correlates with a slow loss of pluripotency**
Our next goal was to understand the dynamics of the cell fate transitions to understand
how signaling dynamics determined the observed cell fates. We treated hPSCs with
pulses of 10 ng/mL BMP of varying durations and examined marker gene expression at
16h, 24h, 32h and 48h after treatment (Fig. 5A-C, S7). When cells were not exposed to
BMP, cells remained pluripotent as marked by high SOX2 expression at every time point

**Figure 5. Cell fate transitions under BMP treatment. (A-C)** Scatter plots of the quantifications of SOX2, BRA and CDX2 under the treatments mTeSR 0-48h (A), BMP 10ng/mL 0-48h (B), and BMP 10ng/mL 0-16h followed by mTeSR 16-48h (C), at 16h, 24h, 32h, and 48h. Each dot corresponds to a single cell, and its color marks the cell fate assigned to the cell as shown in the legend at the bottom. **(D)** Schematic of resulting mRNA transcribed from the labelled SOX2 allele after CRISPR-Cas-mediated SOX2-mCitrine-T2A-BSD knockin. The blasticidin resistance protein (BSD) facilitates selection of labelled cells. T2A is a self-cleaving peptide that enables separation of BSD from SOX2::mCitrine. **(E)** Representative images of hPSCs with mCitrine-labelled SOX2 and the nuclear marker CFP::H2B. (Left) Under no treatment. (Right) SOX2 expression is lost after 48h of 10ng/mL BMP4 treatment. **(F)** Confocal microscopy images of live SOX2::mCitrine hPSCs at 0h, 24h, and 48h after treatment with 10ng/mL BMP4. **(G)** SOX2::mCitrine average nuclear intensity after the indicated treatments. Scale bars: 100um.

observed (Fig. 5A). On the other hand, if hPSCs were exposed to BMP for the whole 48h, a pluripotent-to-extraembryonic transition was observed between 24 and 32 hours, when cells downregulated SOX2 and then upregulated CDX2 (Fig. 5B). Importantly, no BRA expression was observed in this condition at any time point, indicating that cells transition directly to extraembryonic without going through an intermediate mesoderm state. In the experimental conditions where mesoderm differentiation was obtained, such as 16 hours of 10 ng/mL BMP4 followed by mTeSR, a slow SOX2 downregulation was observed followed by a late BRA upregulation, starting after 32h of induction (Fig. 5C). This late mesoderm induction is consistent with recent studies that propose that SOX2 levels need to be low for Wnt to induce mesoderm differentiation (Blassberg et al., 2022, 2).

In order to observe gene expression dynamics with a higher time resolution, we used CRISPR-Cas9 genome engineering to insert mCitrine at the endogenous locus to form a C-terminal fusion with SOX2 (SOX2::mCitrine) (Fig. 5D, S8, Materials and Methods). To facilitate nuclear identification and analysis, the cells also express CFP::H2B, which was incorporated into the genome using the ePiggyBac transposable element system (Lacoste et al., 2009). Treatment of these cells with BMP4 showed a rapid downregulation of SOX2, with undetectable expression after two days of treatment (Fig. 5E,F, S8A,B), and antibody staining for SOX2 and other pluripotency markers showed that these cells shared similar dynamics with WT hPSCs (Fig. S8).

We imaged SOX2::mCitrine cells under different BMP4 treatments and observed the SOX2 dynamics for two days. Consistent with results above, SOX2 expression was maintained in cells that were not exposed to BMP4 (Fig. 5G). On the other hand, SOX2 expression rapidly began to decay under BMP4 treatment, with a rate that was initially proportional to BMP4 concentration (Fig. 5G, S8D). If BMP4 was withdrawn from the media early enough, SOX2 expression returned to high levels, while if cells were induced with long pulses of BMP4, they lost SOX2 expression. Interestingly, with a 16-hour pulse of 10ng/mL BMP4, which leads to peak mesoderm differentiation, SOX2 downregulation slowed after 16 hours. A second phase of SOX2 decay led to hPSCs eventually loosing SOX2 expression but later than cells induced under longer BMP4 pulses (Fig. 5G). These results show that mesoderm induction correlates with a slow SOX2 decay after BMP withdrawal.

The initial decay of SOX2 expression is strongly correlated with the integral of BMP4 in time (Fig. S9A-E) suggesting that there is a tradeoff between the magnitude and the duration of signaling in the dissolution of the pluripotent state. This correlation no longer holds in later times (Fig. S9E), and the total integral of BMP4 signal does not directly translate to cell fate (Fig. S9F). This supports the hypothesis that BMP initially controls the exit from pluripotency, and the combinatorial interpretation of BMP and endogenous WNT signaling controls the decision between ExE and mesoderm fates.

Taken together, these results suggest that BMP4 concentration initially regulates SOX2 expression, which rapidly starts to decay after BMP4 treatment with a rate that is

**Figure 6. A minimal cell fate network can recapitulate the observed dynamics. (A)** Schematics of the coupling of the signalling network and the minimal cell fate network. **(B)** Simulated SOX2, BRA and CDX2 dynamics from the model in (A) for the indicated experimental conditions. **(C)** Experimental (top) and simulated (bottom) cell fate proportions under constant low concentrations (left) and pulses of BMP 10ng/mL (right).

proportional to SMAD4 activity. If BMP4 is removed early, SOX2 expression recovers to baseline and cells remain pluripotent. Long BMP4 treatments produce a direct pluripotency-to-ExE transition, without going through an intermediate mesoderm state. On the other hand, intermediate BMP4 treatments produce a two-phased SOX2 downregulation, with an initial fast partial decay followed by a slow decay to baseline which correlates with the onset of mesoderm induction.

**A simple network model can recapitulate the observed dynamics**
To understand how signal dynamics controlled the observed cell fate transitions, we created a minimal cell fate network (CFN) model, where the nodes of the network correspond to the cell fates observed, i.e. pluripotent, mesodermal, and extra-embryonic, characterized by high SOX2, BRA, or CDX2 expression, respectively. We coupled this CFN model with the signaling model described above (Fig. 6A). For simplicity, we modelled SOX2, BRA and CDX2 expression in time as a proxy for the corresponding cell fates. To generate mutually exclusive cell states, we included mutually repressive interactions between SOX2, BRA and CDX2 (Fig. 6A).

Secondly, as previous studies have shown and our data supports (Chhabra and
Warmflash, 2021; Li et al., 2013; Minn et al., 2020; Xu et al., 2002; Yang et al., 2021),
BMP activation of SMAD complexes leads to transcription of ExE markers, such as CDX2,
and therefore we included activation of CDX2 expression by nuclear SMAD4 through a
Hill equation (Fig. 6A, Supplemental Information). In this regard, we also observed that
CDX2 and ISL1 expression levels, once active, remained high even if BMP was
withdrawn from the media, suggesting a possible auto-regulation of ExE markers, which
we modelled by an additive auto-activation (Fig. 6A, Supplemental Information).

As WNT signals are necessary for BMP-induced mesoderm formation (Chhabra et al.,
2019; Martyn et al., 2018; Nemashkalo et al., 2017), we considered that mesoderm
markers are upregulated through $\beta$-catenin activation by WNT, and modelled the
activation of BRA expression through a Hill function of $\beta$-catenin levels (Fig. 6A,
Supplemental Information).

Moreover, the SOX2 expression dynamics observed in Fig. 5G suggested a
downregulation of SOX2 by BMP, through a SMAD intermediate, as SOX2 expression
begins to decay rapidly upon BMP treatment, at which stage neither Wnt signaling, as
reflected in nuclear $\beta$-catenin, nor mesoderm or ExE markers are yet upregulated (Fig.
3H, 5A-C). In fact, as mentioned above, the initial rate of SOX2 decay showed a strong
correlation with SMAD4 levels for different concentrations and durations (Fig. S9). Hence,
a direct inhibition of SOX2 expression levels by SMAD4 was included in the CFN model
(Fig. 6A, Supplemental Information).

This minimal deterministic CFN model was fitted to simplified expression levels obtained
in a subset of experiments by using a Monte Carlo optimization algorithm, the details of
which can be found in the Supplemental Information. The model was compatible with the
dynamics observed in the data such as direct pluripotency-to-ExE transition without
expressing BRA under constant 2-day BMP4 induction. The model was also able to mimic
the slow SOX2 decay under an intermediate pulse of BMP4 which also resulted in a late
BRA upregulation (Fig. 6B). Importantly, the model was able to reproduce features and
conditions that were not used in the fitting, such as the observed SOX2 dynamics for
different BMP4 concentrations, and that a longer pulse of 5ng/ml BMP is necessary for
cells to differentiate to the ExE than under 10ng/ml BMP in followed by mTeSR withdrawal
(Fig. S10).

Having fitted the deterministic CFN model to the available data, we generated a stochastic
version by considering an additive white noise of strength $\xi$, to test whether the model
could reproduce the cell fate proportions observed in the data. We fixed all the parameters
from the deterministic model and fitted only the single noise parameter and the initial
distribution of the SOX2 expression to the cell fate proportions obtained in a subset of
experimental data (Supplemental Information). We found that without modification to any
of the parameters from the deterministic model, the stochastic model reproduced the
observed BMP morphogen effect in time and the low mesoderm induction at any low

**Figure 7. Cell fate is a combinatorial response to BMP and WNT. (A)** Fate map defined by BMP (SMAD4) and WNT (bCat) levels obtained from the model in Figure 6. P = Pluripotency. M = Mesoderm. E = Extra-embryonic. **(B)** Signaling trajectories on the fate map for the indicated BMP10ng/mL pulses. **(C)** Simulated SMAD4, b-Catenin, SOX2 and BRA dynamics for the conditions indicated on the legend. **(D)** Simulated prediction (right) SOX2 and BRA intensities for the same conditions as in (D), and corresponding experimental confirmation (left).

350    constant concentration (Fig. 6C). Taken together the simple cell fate network model
351    recapitulated the dynamics and cell proportions observed in the data including in a large
352    amount of data not used to fit the model parameters.
353
**Cell fate is a combinatorial response to BMP and WNT**
With the obtained model, we then aimed to understand how the signals BMP and WNT
controlled cell fate determination. We created a phase diagram, or fate map, that shows,
for a given value of SMAD4 and β-catenin, which cell fates are stable, or, in other words,
accessible, in the deterministic model under those signaling levels (Fig. 7A). For example,
for low SMAD4 and β-catenin levels, only the pluripotent (P; SOX2+) state is stable and,

therefore, cells remain pluripotent under low BMP and WNT levels. On the other hand, for high SMAD4 values and low β-catenin levels, the only stable state is the ExE (E; CDX2+) fate, therefore the model suggests that cells would eventually adopt the ExE fate if induced with high BMP while inhibiting WNT signaling. For high SMAD4 and high β-catenin levels, a bistable region is defined, where both the mesodermal (M; BRA+) and ExE fates are accessible, and cells will become one or the other depending on the initial state of the cell.

However, cells are normally not exposed to constant SMAD4 and β-catenin levels, but they experience the signaling dynamics discussed previously, some of which are measured in Fig. 3. Combining these signaling trajectories and the fate map, the model explains the experimental observations by showing that cell fate is a combinatorial response to BMP and WNT. Under short pulses of BMP, endogenous WNT is not activated, so the signaling trajectory moves to a high BMP region for a short period of time after which it goes back to a low BMP, low WNT profile and cells stay pluripotent (Fig. 7B, pink curve, and S11). Under constant high BMP, endogenous WNT is activated, and the signal trajectory ends up in a region of high BMP and high WNT, where both ExE and mesoderm differentiation are possible, however, the trajectory lays in the basin of attraction of the ExE fate and therefore cells adopt this fate (Fig. 7B, blue curve). Under medium durations where WNT becomes self-sustained, even though the signal trajectory initially moves in a high BMP region for some time, decreasing the SOX2 levels in the cell, the withdrawal of the BMP signal situates the signal trajectory in a region of tristability (Fig. 7A-B, gray region), which slows down the decay of pluripotent markers and prevents cells from increasing the expression of ExE markers (Fig. 7B, yellow curve, and S11). This allows cells to become mesoderm once WNT increases and the signaling trajectory moves to the bistable ExE and mesoderm region (Fig. 7B, yellow curve, and S11). On the other hand, constant low concentrations result in a slow WNT upregulation, and therefore the signal trajectory stays in a pluripotency or ExE-promoting region for too long, and are inefficient at promoting mesoderm differentiation (Fig. 7C, S11). Taken together, the model unveils a strategy for efficiently inducing mesoderm with BMP: induction with a high BMP dose that rapidly increases endogenous WNT signaling, after which BMP must be withdrawn for cells to become mesoderm over ExE.

Here we have shown that by pulsing BMP signal, we can obtain a relatively high population of cells that exclusively expresses the mesoderm marker BRA after 2 days. However, this population is heterogeneous and we speculated that this was due to the limitations of the dynamics of endogenous WNT in response to BMP. Inducing hPSCs by CHIR results in a strong, rapid and sustained WNT response (Massey et al., 2019) and yields a very high fraction of mesoderm cells (Fig. S11E). However, this induction is slow, most of the cells still expressing SOX2 after 2 days of induction (Fig. 7E, S11), and taking 3 days to obtain a homogeneous mesodermal population (Fig. S11E). Modelling CHIR through β-catenin activation (Fig. 4) we could reproduce this effect (Fig. 7D,E, yellow condition). We wondered whether we could take advantage of the insights above to speed up mesoderm induction by CHIR. Our goal was to find a 2-day protocol that yielded the

highest BRA expression by varying a sequence of inductions with BMP and CHIR. Our model suggested that induction of mesoderm by CHIR was slow because of a strong SOX2 inhibition of the mesoderm state. We reasoned that a pulse of BMP4 signal could be used to destabilize the pluripotency state which, if then followed by CHIR induction, could yield a higher proportion of mesoderm cells by 48 hours. Our model predicted that a 4-hour pulse of BMP was sufficient for this purpose, and if followed by 44 hours of CHIR, would result in a very high BRA expression by day 2 (Fig. 7D,E, red condition). Indeed, experimentally, by day 2 we obtained higher BRA expression and much more complete SOX2 downregulation than with CHIR alone (Fig. 7E, yellow condition). The model explains the combinatorial effect between BMP and Wnt signals in mediating decisions between the pluripotent, mesoderm and ExE fates and offers a platform to rationally design more optimal protocols. Taken together, our study highlights how understanding signaling dynamics can be exploited for developing efficient differentiation protocols.

## Discussion

In this study, we have unveiled a combinatorial mechanism by which BMP and downstream endogenous WNT signaling combinatorially control the cell state transitions observed in mammalian gastrulation and previous *in vitro* studies. We showed a BMP-induced morphogen effect in the duration but not the concentration of signaling, indicating that duration and concentration of BMP signal are not interchangeable in this context. In particular, a specific pulse of high concentration of BMP4 signal is much more efficient at inducing a mesoderm-like state than any constant concentration. Taking advantage of signaling and fate reporter cell lines, we showed that these results depend on an endogenous BMP-induced WNT bistability. A simple minimal cell fate network model explains how the observed cell state transitions are regulated by combinatorial interpretation of these two signals. Induction with a short BMP pulse is not enough to induce endogenous WNT signaling, resulting in hPSCs remaining in the initial pluripotent state. Long pulses of high BMP4 concentration, on the contrary, induce endogenous WNT signals but a high BMP signal overrides the endogenous WNT resulting in extraembryonic differentiation. A pulse of BMP4 of intermediate length that activates WNT autoregulation results in efficient mesoderm differentiation. At lower constant concentrations, WNT activation is slow and cells are therefore pushed towards an ExE fate before experiencing this signal, explaining why low doses of BMP are not interchangeable with a pulse of a high BMP dose. Taken together, our study reveals an underlying logic where, in order to induce efficient mesodermal induction, BMP signaling needs to be sufficiently strong to rapidly induce endogenous WNT signaling upregulation, but sufficiently short for cells to differentiate in a low BMP, high WNT environment.

Our work adds to the growing body of knowledge on how the dynamics of signals are interpreted by downstream regulatory networks. In the case of Shh signaling in the murine neural tube, there is an effective tradeoff between duration and concentration so that shorter durations at higher concentration are sufficient to induce the same fates as longer exposure to lower concentrations (Dessaud et al., 2007). In the case of the three-way

decision between pluripotent, primitive streak mesoderm, and extraembryonic amnion,
the situation is more complex and relies on the interplay between BMP and Wnt signaling,
and cannot be predicted by a parameter of any one of these pathways. Interestingly, we
observed that the initial decrease in SOX2 correlated very well with the cumulative
integral of BMP signaling as measured by SMAD4 across several different time courses
of applied BMP signal (Fig. S9E). This suggests that the initial dissolution of the
pluripotent state by BMP signaling may indeed by controlled by the integral of signaling,
but that the decision between different potential fates relies on the interpretation of
multiple pathways. Indeed, when Wnt signaling is inhibited so that cells can only switch
from the pluripotent to the extraembryonic fate, the fraction of cells adopting this fate is
well predicted by the integral of the BMP signal (Teague and Heemskerk, submitted).

It is also important to note that measuring the intensity of signaling with a reporter is
essential as the external concentration doesn't translate directly into signaling activity. In
the case of BMP signaling, the activation of the SMAD proteins is switch-like and
regulated over a narrow range of concentrations, so that it is difficult to regulate the
effective activity simply by changing the concentration of ligand in the media (Heemskerk
et al., 2019). Further, the loss of the BMP from the media over time causes more
prolonged signaling at higher doses (Heemskerk et al., 2019), making it difficult to
decouple duration from concentration. Together these features mean that even though
the dissolution of pluripotency is responsive to the integral of signaling, in practice, the
duration of signal is more easily controlled, and is more likely to be the determining factor
in vivo.

It has recently been proposed that BMP4 differentiation rapidly becomes irreversible due
to positive feedback through GATA3 (Gunne-braden et al., 2020). In that study, a one-
hour pulse of BMP4 was sufficient to induce irreversible differentiation in hPSC colonies,
which is in clear contradiction with our results (Fig. 1, 2, S1, S2). Irreversible differentiation
following short exposure to BMP is contradictory to several studies from our lab and
others (Chhabra et al., 2019; Etoc et al., 2016; Jo et al., 2022; Nemashkalo et al., 2017)
that showed that BMP signaling is rapidly downregulated by small molecule or
extracellular protein inhibitors and that pulses of longer than 10 or 24 hours are needed
to induce differentiation in both in micropatterned colonies and regular culture,
respectively. Apparently irreversible differentiation is likely due to incomplete removal of
the BMP in the media (Fig. S2C-D), and indeed we found using a live cell reporter for
BMP signaling that signaling activity remained substantially above baseline when media
containing high doses of BMP was replaced by media alone, however, treatment with
Noggin quickly abrogated this continued signaling. Further experiments with washing
confirmed that BMP ligands cannot be removed from the media by washing alone (see
Fig. S2C).

Our results showed that under BMP induction of hPSCs there is a relatively narrow
window of durations of BMP exposure which are able to specify mesoderm, and this was
related to the dynamics of the induced endogenous WNT signaling. This highlights the

need to consider dynamics when developing in vitro differentiation protocols. In particular,
while here we investigated the importance of the duration of a single pulse of BMP, more
complex time courses of BMP stimulation are possible, and it would be interesting to
determine the optimal time courses for achieving different fates. In vivo, while the same
cascade of signals controls gastrulation in other mammals, there is a wide range of
developmental time scales, and it is unclear by what mechanism the window of BMP
duration needed might be matched to the developmental time scale.  Interestingly, recent
studies have uncovered cell movement as a mechanism that controls the temporal
exposure to morphogens (Fulton et al., 2022). Further work could elucidate whether the
differences might be due to differential protein stability and cell cycle duration as has been
recently proposed in other contexts (Matsuda et al., 2020; Rayon et al., 2020).
503
We show that there is a form of memory in the WNT signaling so that transient exposure
to BMP can be sufficient to induce WNT in a sustained fashioned. This is reminiscent of
the interplay between WNT and NODAL signaling in which cells remember prior exposure
to WNT, which alters their subsequent response to NODAL (Yoney et al., 2018; Yoney et
al., 2022). The mechanisms are different as in the case described here BMP leads to
ongoing WNT signaling, likely through sustained expression of WNT ligands, while WNT
affects the interpretation of NODAL by inducing EOMES without a subsequent
requirement for the continuation of WNT activity.
512
A popular recent approach based in the Waddington landscape metaphor has been
shown to be a powerful yet simple way to study cell fate transitions in different contexts
(Camacho-Aguilar et al., 2021; Coomer et al., 2022; Corson and Siggia, 2012; Corson
and Siggia, 2017; Huang, 2012; Mojtahedi et al., 2016; Sáez et al., 2022; Valcourt et al.,
2021). In these models, the differentiation of a cell is depicted as ball rolling down a
landscape of hills and valleys, that represent the different cell types the stem cell can
differentiate into. In particular, we and others have shown that using dynamical systems
theory one can enumerate the nature of the possible bifurcations, and parametrize these
through the signals to build the landscape model, which then can be quantitatively fitted
to the data (Camacho-Aguilar et al., 2021; Sáez et al., 2022). These studies, however,
lacked signaling dynamics data and the underlying bifurcations had to be guessed and
then validated through fitting. Also, these models have not been compared yet to any
more mechanistic models. Here using a cell fate network model, which although simple
contains some mechanistic knowledge, we have deciphered the underlying fate map
without any prior knowledge about the candidate bifurcations. Interestingly, our three
mutually repulsive state network contains the bifurcations present in the elliptic-umbilic
catastrophe, as had been proposed before (Rand et al., 2021). To our knowledge, this is
the first experimental system that confirms this idea. We believe that our results provide
a foundation to quantitatively compare mechanistic and landscape models using
quantitative data on signaling dynamics.
533
Much work has been done to decipher the dynamics of signals that control the patterning
of micropatterned hPSC colonies treated with BMP4 (Chhabra et al., 2019; Etoc et al.,

2016; Heemskerk et al., 2019; Warmflash et al., 2014). These studies have unveiled that,
initially, there is a homogeneous response to BMP4 across the whole colony, which is
restricted to the edge via receptor localization and accumulation of Noggin at the colony
center between 10 and 20 hours after treatment (Etoc et al., 2016; Heemskerk et al.,
2019). Subsequently, around 30h, waves of WNT and Nodal signaling start near the edge
of the colony and move inwards, spatially correlating with a ring of BRA-positive
mesodermal cells (Chhabra et al., 2019; Heemskerk et al., 2019). Although our results
have been obtained in a culture with lower cell density where self-organized patterning
does not occur, they are consistent with the observations of a pulse of BMP throughout
the colony which induces endogenous WNT signal. The cells that adopt a mesodermal
fate are those that are displaced from the edge and therefore only experience transient
BMP signaling followed by upregulation of Wnt signaling to high levels. Thus, the model
developed here could be a good starting point to build a spatial model to understand how
patterns arise from the interplay of dynamic signaling and combinatorial interpretation.
550
551
552
553

**Materials and Methods**

**Cell lines.**
The cell lines used were ESI017 (NIHhPSC-11-0093), ESI017 GFP::$\beta$-Catenin RFP::H2B
(Massey et al., 2019), RUES2 GFP::Smad4 RFP::H2B (Nemashkalo et al., 2017), and
ESI017 SOX2::mCitrine CFP::H2B (this study). ESI017 cells were obtained directly from
ESIBIO while RUES2 were a gift of Ali Brivanlou (Rockefeller University). The identity of
these cells as pluripotent cells was confirmed via triple staining for pluripotency markers
OCT4, SOX2, and NANOG. All cells were routinely tested for mycoplasma contamination
and found negative.

**Cell Culture, Treatments, and Differentiation.**
All cell lines were maintained in pluripotency maintenance culture as described in
(Nemashkalo et al., 2017). ESI017 SOX2::mCitrine CFP::H2B were maintained in mTeSR
Plus medium (STEMCELL Technologies; 100-0276) and Blasticidin (5$\mu$g/ml; A.G.
Scientific; B-1247-SOL) for selection, which was removed before experiments.
Ibidi $\mu$-Slide 8 Well plates (Ibidi; 80826) were used for experiments, which were coated
with Matrigel (5$\mu$l/ml; Corning; 354277) diluted in DMEM/F12 (VWR;45000-344). For all
experiments, cells were seeded into mTeSR1 medium (STEMCELL Technologies;
85857) containing rock inhibitor Y27672 (10$\mu$M; STEMCELL Technologies; 05875) at a
density of 4 x 10$^4$/cm$^2$ (except when noted otherwise). Treatment started 21 hours after
seeding, and media was always changed every 24 hours, and when performing the
indicated specific treatments.
The following recombinant proteins and small molecules were used: BMP4 (R&D
Systems; 314BP050), Noggin (500ng/ml; R&D Systems; 6057-NG-100), IWP2
(Stemgent; 040034; 3$\mu$M), CHIR99021 (24$\mu$M; MedChem Express; HY-10182).

For micropatterning experiments, these were done in a 96-well plate (CYTOO) of 700 $\mu m$
circular micropatterns. Coating and seeding were done as previously described by
(Warmflash et al., 2014). Briefly, wells were coated with Human Recombinant Laminin
511 (Fisher Scientific) in a 1:20 dilution in PBS with calcium and magnesium for 3 hours
at 37 °C. hESCs were then seeded as single cells and maintained in mTeSR medium.
After overnight incubation, cells were treated with 50 ng/ml BMP4 (R&D systems) for 48
hours and with 1 hour pulse of 50 ng/ml BMP4 (R&D systems), as previously described
in (Gunne-braden et al., 2020), with or without 100 ng/ml of Noggin (Fisher Scientific).
The following recombinant proteins and small molecules were used: BMP4 (R&D
Systems; 314BP050), Noggin (250ng/ml; R&D Systems; 6057-NG-100), IWP2
(Stemgent; 040034; 3$\mu$M), CHIR99021 (24$\mu$M; MedChem Express; HY-10182).
592

**Plasmids and Generation of SOX2-mCitrine-Labeled hPSC Cell Line.**
We used CRISPR-Cas9 technology for gene editing, where the ESI017 hPSC line was
used as the parental line. We used previously published constructs to fuse mCitrine
directly with SOX2 at the C-terminus of the SOX2 coding sequence (Martyn et al., 2018).
The SOX2 homology donor consists of a 1-kb homology arm, an
mCitrine::T2A::blasticidin cassette, and a 1-kb right homology arm. Cas9 expression
plasmid, homology donor DNA (Plasmid AW-P46), and guide RNA (Plasmid AW-P45;
GTGCCCGGCACGGCCATTAA) were nucleofected in hPSCs using the P3 Primary Cell
4D-Nucleofector X Kit (Lonza; V4Xp-3012), and positive transformants were selected with
blasticidin (10$\mu$g/ml; A.G. Scientific; B-1247-SOL) and CloneR (STEMCELL
Technologies; 05889) for two days, after which cells were passaged and single clones
were handpicked and amplified. Sanger sequencing was performed to screen and confirm
a successful clone (Primer sequences are listed in Table 1). After establishment, the
stable line was checked for pluripotency markers, i.e. OCT4, SOX2 and NANOG
expression, as well as BMP differentiation both in regular culture and micropatterning,
and was found indistinguishable from WT ESI017 hPSCs.
An ePiggyBac (ePB) master vector based on the pBSSK backbone (Lacoste et al., 2009),
harboring transposon-specific inverted terminal repeat sequences (ITR) was modified to
deliver a nuclear marker CFP::H2B (Plasmid AW-P68). ePB master vector and helper
(Plasmid AW-27) were nucleofected into the established SOX2::mCitrine cell line using
the P3 Primary Cell 4D-Nucleofector X Kit (Lonza; V4Xp-3012). G-418 (40ng/mL;
ThermoFisher; 10131035) started two days after nucleofection and lasted for at least 7
days.
616

617 **Table 1**

| Region | Forward primer | Reverse primer |
|---|---|---|
| SOX2::mcitrine | ccagctcgcagacctacatgaa | TGGCGGATCTTGAAGTTCACCT |
| mcitrine::T2A::BSD::SOX2 | AGGTGAACTTCAAGATCCGCCA | GTCATTTGCTGTGGGTGATGGG |

618
619

620 **Immunofluorescence Antibody Staining.**

621 Cells were fixed with 4% PFA and stained as described in (Nemashkalo et al., 2017).
622 Antibodies and dilutions used are listed in Table 2.
623
624 **Table 2**

| Protein | Species | Dilution | Vendor | Catalog no. |
|---------|---------|----------|--------|-------------|
| Sox2 | Rabbit | 1:200 | Cell Signalling Technologies | 23064s |
| Brachyury | Goat | 1:300 | R&D Systems | AF2085 |
| Brachyury | Rabbit | 1:400 | R&D Systems | MAB20851 |
| Cdx2 | Mouse | 1:100 | Biogenex | MU392A-5UC |
| Isl1 | Mouse | 1:50 | DSHB | 39.4D-5 |
| Oct4 | Mouse | 1:400 | BD Biosciences | 611203 |
| Nanog | Mouse | 1:400 | BD Biosciences | 560482 |
| Nanog | Goat | 1:200 | R&D Systems | AF1997 |
| Hand1 | Goat | 1:200 | R&D Systems | AF3168 |
| Gata3 | Rabbit | 1:100 | ThermoFisher | PA1-101 |
| Tbx6 | Goat | 1:200 | R&D Systems | AF4744 |

625
626
**Imaging and Analysis.**
Imaging acquisition was done on an Olympus/Andor spinning disk confocal microscope
with a 20x, 0.75NA air objective.
For live imaging experiments, reporter cell lines were maintained in antibiotic selection
for the H2B fluorescent marker for at least three days and stopping 2 days before seeding,
at the latest, to maximize the number of fluorescent cells in the culture. Time-lapse
imaging intervals were 15 minutes, and Z-stacks were acquired in three planes spaced
2.5μM apart. During imaging, temperature (37 $^\circ$C), humidity (~50%), and $CO_2$ (5%) were
controlled, and media change was performed without moving the plate from the stage. 8
positions of each condition were selected for imaging.
Image analysis was performed using Ilastik (Berg et al., 2019; Sommer et al., 2011) for
initial segmentation and custom-written MATLAB code available at
https://github.com/warmflashlab/Camacho-Aguilar2022_BMPWNT for further analysis.
Smad4 dynamics was quantified as the nuclear to cytoplasmic Smad4 ratio, (Heemskerk
et al., 2019). Nuclear protein expression was measured by mean nuclear intensity and
normalized by mean nuclear DAPI to correct for intensity variations due to optics.
For the analysis of the micropatterning experiment, stitching of the colonies was
performed in Fiji using the algorithm in (Preibisch et al., 2009). Segmentation and mean
intensity quantification were done on Ilastik (Berg et al., 2019; Sommer et al., 2011) and
custom software written in MATLAB (MathWorks), previously described in (Warmflash et
al., 2014).
648
649

650
651 **Cell Fate Classification.**
652 (see Appendix)
653
654
655 **Mathematical Models.**
656 (see Appendix)
657
658
# Acknowledgements

We thank Idse Heemskerk, Seth Teague, Francisco-Jesús Castro-Jiménez, Nathan Lord,
and Eric Siggia for helpful discussions on the project. Sapna Chhabra for performing the
experiment in Fig S8C and for helpful feedback on the project. Joseph Massey for sharing
helpful pieces of code for analyzing live-cell imaging data. Cecilia Guerra and Lizhong Liu
for technical assistance with experiments; and all the members of the Warmflash lab for
helpful feedback. The work was supported by grants to AW from the National Science
Foundation (MCB-1553228 and MCB-2135296) and the Simons Foundation (511079).

667
668
669

**Arnold, S. J. and Robertson, E. J.** (2009). Making a commitment: cell lineage allocation and axis
patterning in the early mouse embryo. *Nature reviews Molecular cell biology* **10**, 91–
103.

**Bardot, E. S. and Hadjantonakis, A.-K.** (2020). Mouse gastrulation: Coordination of tissue
patterning, specification and diversification of cell fate. *Mechanisms of Development*
**163**, 103617.

**Ben-Haim, N., Lu, C., Guzman-Ayala, M., Pescatore, L., Mesnard, D., Bischofberger, M., Naef,**
**F., Robertson, E. J. and Constam, D. B.** (2006). The nodal precursor acting via activin
receptors induces mesoderm by maintaining a source of its convertases and BMP4.
*Developmental cell* **11**, 313–323.

**Berg, S., Kutra, D., Kroeger, T., Straehle, C. N., Kausler, B. X., Haubold, C., Schiegg, M., Ales, J.,**
**Beier, T., Rudy, M., et al.** (2019). ilastik: interactive machine learning for (bio)image
analysis. *Nat Methods* **16**, 1226–1232.

**Blassberg, R., Patel, H., Watson, T., Gouti, M., Metzis, V., Delás, M. J. and Briscoe, J.** (2022).
Sox2 levels regulate the chromatin occupancy of WNT mediators in epiblast progenitors
responsible for vertebrate body formation. *Nat Cell Biol* **24**, 633–644.

**Brennan, J., Lu, C. C., Norris, D. P., Rodriguez, T. A., Beddington, R. S. P. and Robertson, E. J.**
(2001). Nodal signalling in the epiblast patterns the early mouse embryo. *Nature* **411**,
965–969.

**Camacho-Aguilar, E. and Warmflash, A.** (2020). Insights into mammalian morphogen dynamics
from embryonic stem cell systems. *Current Topics in Developmental Biology* **137**, 279–
305.

**Camacho-Aguilar, E., Warmflash, A. and Rand, D. A.** (2021). Quantifying cell transitions in C.
elegans with data-fitted landscape models. *PLoS Computational Biology* **17**, 1–28.

**Chhabra, S. and Warmflash, A.** (2021). BMP-treated human embryonic stem cells
transcriptionally resemble amnion cells in the monkey embryo. *Biology Open* **10**,
bio058617.

**Chhabra, S., Liu, L., Goh Id, R., Kong, X. and Warmflash Id, A.** (2019). Dissecting the dynamics
of signaling events in the BMP, WNT, and NODAL cascade during self-organized fate
patterning in human gastruloids. *PLoS biology* **17**,.

**Conlon, F. L., Lyons, K. M., Takaesu, N., Barth, K. S., Kispert, A., Herrmann, B. and Robertson,**
**E. J.** (1994). A primary requirement for nodal in the formation and maintenance of the
primitive streak in the mouse. *Development (Cambridge, England)* **120**, 1919–1928.

**Coomer, M. A., Ham, L. and Stumpf, M. P. H.** (2022). Noise distorts the epigenetic landscape
and shapes cell-fate decisions. *Cell Systems* **13**, 83-102.e6.

**Corson, F. and Siggia, E. D.** (2012). Geometry, epistasis, and developmental patterning.
*Proceedings of the National Academy of Sciences of the United States of America* **109**,
5568–5575.

**Corson, F. and Siggia, E. D.** (2017). Gene-free methodology for cell fate dynamics during
development. *eLife* **6**, e30743.

**Dessaud, E., Yang, L. L., Hill, K., Cox, B., Ulloa, F., Ribeiro, A., Mynett, A., Novitch, B. G. and**
**Briscoe, J.** (2007). Interpretation of the sonic hedgehog morphogen gradient by a
temporal adaptation mechanism. *Nature* **450**, 717–720.

**Etoc, F., Metzger, J., Ruzo, A., Kirst, C., Yoney, A., Ozair, M. Z., Brivanlou, A. H. and Siggia, E. D.**
(2016). A Balance between Secreted Inhibitors and Edge Sensing Controls Gastruloid
Self-Organization. *Developmental Cell* **39**, 302–315.

**Fu, J., Warmflash, A. and Lutolf, M. P.** (2021). Stem-cell-based embryo models for fundamental
research and translation. *Nature Materials* **20**, 132–144.

**Fulton, T., Spiess, K., Thomson, L., Wang, Y., Clark, B., Hwang, S., Paige, B., Verd, B. and**
**Steventon, B.** (2022). Cell Rearrangement Generates Pattern Emergence as a Function
of Temporal Morphogen Exposure. 2021.02.05.429898.

**Gunne-braden, A., Sullivan, A., Gharibi, B., Wollman, R., East, P., Santos, S. D. M., Gunne-**
**braden, A., Sullivan, A., Gharibi, B., Sheriff, R. S. M., et al.** (2020). GATA3 Mediates a

Fast , Irreversible Commitment to BMP4-Driven Differentiation in Human Embryonic
Stem Cells. *Stem Cell* 1–14.

**Heemskerk, I.** (2020). Full of potential: Pluripotent stem cells for the systems biology of
embryonic patterning. *Developmental Biology* **460**, 86–98.

**Heemskerk, I., Burt, K., Miller, M., Chhabra, S., Guerra, M. C., Liu, L. and Warmflash, A.** (2019).
Rapid changes in morphogen concentration control self-organized patterning in human
embryonic stem cells. *eLife* **8**,.

**Huang, S.** (2012). The molecular and mathematical basis of Waddington's epigenetic landscape:
A framework for post-Darwinian biology? *BioEssays* **34**, 149–157.

**Jo, K., Teague, S., Chen, B., Khan, H. A., Freeburne, E., Li, H., Li, B., Ran, R., Spence, J. R. and**
**Heemskerk, I.** (2022). Efficient differentiation of human primordial germ cells through
geometric control reveals a key role for Nodal signaling. *eLife* **11**, e72811.

**Kurek, D., Neagu, A., Tastemel, M., Tüysüz, N., Lehmann, J., van de Werken, H. J. G., Philipsen,**
**S., van der Linden, R., Maas, A., van IJcken, W. F. J., et al.** (2015). Endogenous WNT
Signals Mediate BMP-Induced and Spontaneous Differentiation of Epiblast Stem Cells
and Human Embryonic Stem Cells. *Stem Cell Reports* **4**, 114–128.

**Lacoste, A., Berenshteyn, F. and Brivanlou, A. H.** (2009). An Efficient and Reversible
Transposable System for Gene Delivery and Lineage-Specific Differentiation in Human
Embryonic Stem Cells. *Cell Stem Cell* **5**, 332–342.

**Li, Y., Moretto-Zita, M., Soncin, F., Wakeland, A., Wolfe, L., Leon-Garcia, S., Pandian, R., Pizzo,**
**D., Cui, L., Nazor, K., et al.** (2013). BMP4-directed trophoblast differentiation of human
embryonic stem cells is mediated through a ΔNp63+ cytotrophoblast stem cell state.
*Development (Cambridge, England)* **140**, 3965–3976.

**Liu, P., Wakamiya, M., Shea, M. J., Albrecht, U., Behringer, R. R. and Bradley, A.** (1999).
Requirement for Wnt3 in vertebrate axis formation. *Nature Genetics* **22**, 361–365.

**Martyn, I., Kanno, T. Y., Ruzo, A., Siggia, E. D. and Brivanlou, A. H.** (2018). Self-organization of
a human organizer by combined Wnt and Nodal signalling. *Nature* **558**, 132–135.

**Massey, J., Liu, Y., Alvarenga, O., Saez, T., Schmerer, M. and Warmflash, A.** (2019). Synergy
with TGFβ ligands switches WNT pathway dynamics from transient to sustained during
human pluripotent cell differentiation. *Proceedings of the National Academy of Sciences*
**116**, 4989.

**Matsuda, M., Hayashi, H., Garcia-Ojalvo, J., Yoshioka-Kobayashi, K., Kageyama, R., Yamanaka,**
**Y., Ikeya, M., Toguchida, J., Alev, C. and Ebisuya, M.** (2020). Species-specific
segmentation clock periods are due to differential biochemical reaction speeds. *Science*
**369**, 1450–1455.

**Minn, K. T., Fu, Y. C., He, S., Dietmann, S., George, S. C., Anastasio, M. A., Morris, S. A. and**
**Solnica-Krezel, L.** (2020). High-resolution transcriptional and morphogenetic profiling of
cells from micropatterned human ESC gastruloid cultures. *eLife* **9**, e59445.

**Mojtahedi, M., Skupin, A., Zhou, J., Castaño, I. G., Leong-Quong, R. Y. Y., Chang, H., Trachana,**
**K., Giuliani, A. and Huang, S.** (2016). Cell Fate Decision as High-Dimensional Critical
State Transition. *PLOS Biology* **14**, e2000640.

**Nemashkalo, A., Ruzo, A., Heemskerk, I. and Warmflash, A.** (2017). Morphogen and
community effects determine cell fates in response to BMP4 signaling in human
embryonic stem cells. *Development (Cambridge)* **144**,.

**Preibisch, S., Saalfeld, S. and Tomancak, P.** (2009). Globally optimal stitching of tiled 3D
microscopic image acquisitions. *Bioinformatics* **25**, 1463–1465.

**Rand, D. A., Raju, A., Sáez, M., Corson, F. and Siggia, E. D.** (2021). Geometry of gene regulatory
dynamics. *Proceedings of the National Academy of Sciences* **118**, e2109729118.

**Rayon, T., Stamataki, D., Perez-Carrasco, R., Garcia-Perez, L., Barrington, C., Melchionda, M.,**
**Exelby, K., Lazaro, J., Tybulewicz, V. L. J., Fisher, E. M. C., et al.** (2020). Species-specific
pace of development is associated with differences in protein stability. *Science* **369**,
eaba7667.

**Sáez, M., Blassberg, R., Camacho-Aguilar, E., Siggia, E. D., Rand, D. A. and Briscoe, J.** (2022).
Statistically derived geometrical landscapes capture principles of decision-making
dynamics during cell fate transitions. *Cell Systems* **13**, 12-28.e3.

**Sagner, A. and Briscoe, J.** (2017). Morphogen interpretation: concentration, time, competence,
and signaling dynamics. *Wiley Interdisciplinary Reviews: Developmental Biology* e271.

**Shahbazi, M. N., Siggia, E. D. and Zernicka-Goetz, M.** (2019). Self-organization of stem cells into
embryos: A window on early mammalian development. *Science* **364**, 948–951.

**Sommer, C., Straehle, C., Köthe, U. and Hamprecht, F. A.** (2011). Ilastik: Interactive learning
and segmentation toolkit. In *2011 IEEE International Symposium on Biomedical Imaging:*
*From Nano to Macro*, pp. 230–233.

**Tewary, M., Ostblom, J., Prochazka, L., Zulueta-Coarasa, T., Shakiba, N., Fernandez-Gonzalez,**
**R. and Zandstra, P. W.** (2017). A stepwise model of reaction-diffusion and positional
information governs self-organized human peri-gastrulation-like patterning.
*Development* **144**, 4298–4312.

**Tortelote, G. G., Hernández-Hernández, J. M., Quaresma, A. J. C., Nickerson, J. A., Imbalzano,**
**A. N. and Rivera-Pérez, J. A.** (2013). Wnt3 function in the epiblast is required for the
maintenance but not the initiation of gastrulation in mice. *Developmental biology* **374**,
164–173.

**Valcourt, J. R., Huang, R. E., Kundu, S., Venkatasubramanian, D., Kingston, R. E. and**
**Ramanathan, S.** (2021). Modulating mesendoderm competence during human germ
layer differentiation. *Cell Reports* **37**, 109990.

**Wang, J., Sinha, T. and Wynshaw-Boris, A.** (2012). Wnt Signaling in Mammalian Development:
Lessons from Mouse Genetics. *Cold Spring Harb Perspect Biol* **4**, a007963.

**Warmflash, A., Sorre, B., Etoc, F., Siggia, E. D. and Brivanlou, A. H.** (2014). A method to
recapitulate early embryonic spatial patterning in human embryonic stem cells. *Nature*
*methods* **11**, 847–54.

**Winnier, G., Blessing, M., Labosky, P. A. and Hogan, B. L.** (1995). Bone morphogenetic protein-
4 is required for mesoderm formation and patterning in the mouse. *Genes &*
*development* **9**, 2105–2116.

**Xu, R.-H., Chen, X., Li, D. S., Li, R., Addicks, G. C., Glennon, C., Zwaka, T. P. and Thomson, J. A.**
(2002). BMP4 initiates human embryonic stem cell differentiation to trophoblast. *Nature*
*biotechnology* **20**, 1261–1264.

**Yang, R., Goedel, A., Kang, Y., Si, C., Chu, C., Zheng, Y., Chen, Z., Gruber, P. J., Xiao, Y., Zhou,**
**C., et al.** (2021). Amnion signals are essential for mesoderm formation in primates. *Nat*
*Commun* **12**, 5126.

**Yoney, A., Etoc, F., Ruzo, A., Carroll, T., Metzger, J. J., Martyn, I., Li, S., Kirst, C., Siggia, E. D.**
**and Brivanlou, A. H.** (2018). WNT signaling memory is required for ACTIVIN to function
as a morphogen in human gastruloids. *eLife* **7**, 1–28.

**Yoney, A., Bai, L., Brivanlou, A. H. and Siggia, E. D.** (2022). Mechanisms underlying WNT-
mediated priming of human embryonic stem cells. *Development* **149**, dev200335.

**Zhang, P., Li, J., Tan, Z., Wang, C., Liu, T., Chen, L., Yong, J., Jiang, W., Sun, X., Du, L., et al.**
(2008). Short-term BMP-4 treatment initiates mesoderm induction in human embryonic
stem cells. *Blood* **111**, 1933–1941.