Title: Systematic characterization of brain cellular crosstalk signaling networks in Alzheimer's disease reveals a novel role for SEMA6D in TREM2-dependent microglial activation

Authors: Ricardo D'Oliveira Albanus[1,2,3]*, Gina M. Finan[4,5]*, Logan Brase[1,2,3], Shuo Chen[6], Qi Guo[7], Abhirami Kannan[1,2,3], Mariana Acquarone[1,2,3], Shih-Feng You[1,2,3], Brenna C Novotny[1,2,3], Patricia M Ribeiro Pereira[8], John C Morris[2,9,10,11], Eric McDade[1,11], Carlos Cruchaga[1,2,3,11], Jasmeer P Chhatwal[11,12], Dominantly Inherited Alzheimer Network (DIAN)[11], Emily E Mace[13], Bruno A Benitez[14], Laura Piccio[2,15,16], Greg T Sutherland[17], Qin Ma[7], Hongjun Fu[6], Celeste M Karch[1,2,3,11], Oscar Harari[1,2,3]†, Tae-Wan Kim[4,5]†

[1]Department of Psychiatry, Washington University School of Medicine, St. Louis, MO, USA
[2]Hope Center for Neurological Disorders, Washington University School of Medicine, St. Louis, MO, USA
[3]NeuroGenomics and Informatics, Department of Psychiatry Washington University in St. Louis
[4]Department of Pathology and Cell Biology, Columbia University Irving Medical Center, New York, NY, USA
[5]Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University Irving Medical Center, New York, NY, USA
[6]Department of Neuroscience, The Ohio State University Wexner Medical Center, Columbus, OH, USA
[7]Department of Biomedical Informatics, Ohio State Comprehensive Cancer Center, Columbus, OH, USA
[8]Department of Radiology, Division of Radiological Sciences, Washington University School of Medicine, St. Louis, MO, USA
[9]Knight Alzheimer Disease Research Center, Washington University School of Medicine, St. Louis, MO, USA
[10]Department of Neurology, Washington University School of Medicine, St. Louis, MO, USA
[11]Dominantly Inherited Alzheimer Network (DIAN)
[12]Department of Neurology, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA
[13]Department of Pediatrics, Columbia University, New York, NY, USA
[14]Department of Neurology, Beth Israel Deaconess, Harvard Medical School, Boston, MA, USA
[15]Department of Pathology and Immunology, Washington University School of Medicine, St. Louis, MO, USA
[16]Brain and Mind Centre, School of Medical Sciences, University of Sydney, Camperdown, NSW, Australia
[17]School of Medical Sciences and Charles Perkins Centre, Faculty of Medicine and Health, The University of Sydney, Sydney, NSW, Australia

* Contributed equally

† Corresponding authors

Abstract

Disrupted cross-cellular communication signaling (cellular crosstalk) has been implicated in neurodegenerative diseases, including Alzheimer's disease (AD). However, there is currently no systematic characterization of brain crosstalk networks in health and disease. We systematically characterized brain cellular crosstalk networks using single-nucleus transcriptomics data from a large cohort of control and AD brain donors (n=67). We found that crosstalk interactions between microglia and neurons were highly enriched to directly involve reported AD risk genes as ligands or receptors. Computational reconstruction of the co-expression networks associated with neuron-microglia crosstalk revealed they perturb additional known AD risk genes in microglia. We identified the interaction of neuronal SEMA6D (a PLXNA1 ligand) with a highly connected microglial regulatory sub-network involving *TREM2*, *APOE*, and *HLA* genes, which we predict is disrupted in late AD stages. Using CRISPR-modified human induced pluripotent stem cell (iPSC)-derived microglia and treatment with recombinant SEMA6D, we experimentally demonstrated that SEMA6D promotes microglial phagocytosis and cytokine (TNFα and IL-6) release in a TREM2-dependent manner. The novel discovery that the SEMA6D-PLXNA1/TREM2 signaling axis is involved in the regulation of microglia function demonstrates that our systematic characterization of cellular crosstalk networks is an important strategy for discovering specific mediators of significant cross-cellular interactions important to AD pathogenesis, gaining wider insights into the biology of this disease, and uncovering novel therapeutics.

Introduction

Cross-cellular signaling (cellular crosstalk) is integral to brain function. By establishing cellular networks mediated by membrane receptors and their corresponding ligands, cells can gather information from their immediate environment and respond accordingly. Cellular crosstalk is crucial to brain homeostasis and crucial processes of neurodevelopment, particularly for synaptic pruning and axon guidance (*1*, *2*). However, increasing experimental and genetic evidence implicates aberrant crosstalk in neurodegenerative diseases (*3–5*). For instance, a recent study systematically characterized crosstalk interactions involving astrocytes and microglia in a mouse model of multiple sclerosis and identified a role for the plexin, semaphorin, and ephrin protein families in neuroinflammation (*5*). Similarly, another study identified a role for astrocyte-microglia crosstalk in modulating amyloid pathology (*6*), and other studies highlighted the role of plexin-semaphorin signaling in mediating immune responses in microglia and other myeloid cells (*7*, *8*). These findings indicate that aberrant cellular crosstalk interactions likely contribute to neurodegeneration. From a translational view, cellular crosstalk is an attractive molecular target for drug development, as s membrane receptors have been widely studied as better therapeutic targets (*9–11*).

Genome-wide association studies have successfully identified genetic risk factors for Alzheimer's disease (AD) and the gene products likely to mediate these signals (*12–15*). Functional genomics leveraging human tissue (*16–19*) and experimental data from human induced pluripotent stem cell (iPSC)-derived cells (*20*) identified a significant fraction of AD risk genes expressed by microglial cells. However, how are the AD risk genes modulated is still unknown. Microglia are the resident immune cells in the brain and are highly attuned to their

surrounding environment sensing signals from neighboring cells (*21*). Therefore, we
hypothesized that microglial AD risk genes modulate cellular crosstalk between microglia and
other brain cell types.

In this study, we used single-nucleus transcriptomic profiles from neuropath-free controls and
AD human brains across a broad spectrum of neuropathological states and genetic
backgrounds to systematically reconstruct the cellular crosstalk networks across seven major
brain cell types: microglia, astrocytes, oligodendrocytes, oligodendrocytes precursors (OPC),
excitatory (exc.) and inhibitory (inh.) neurons, and endothelial cells. We found that neuron-
microglia crosstalk interactions were highly enriched compared to other cross-cellular
interactions to directly involve known AD risk genes. In addition, we identified a sub-network of
microglial genes centered around *TREM2*. We predict this subnetwork is disrupted in late-stage
AD and modulated by the crosstalk interaction between neuronal semaphorin 6D (SEMA6D)
and microglial TREM2. Interestingly, SEMA6D is a known ligand of Plexin-A1, which interacts
with TREM2 and TYROBP (DAP12) to form a receptor complex(*7*). Semaphorins and their
receptors regulate immune cell function and are genetically and functionally implicated in AD (*6,*
22–24). Using wild-type (WT) and *TREM2* knockout (KO) iPSC-derived microglia (iMGL), we
showed that SEMA6D promotes microglia functions, including phagocytosis and cytokine
release in a TREM2-dependent manner. Our findings demonstrate that systematic
characterization of cellular crosstalk networks in human brains is a viable strategy to elucidate
aberrant regulatory biology in AD and other neurodegenerative diseases.

Results

Analyses of human brain transcriptomic profiles identify a role for cellular crosstalk in AD

To systematically characterize cellular crosstalk interactions in neuropath-free controls and AD
brains, we analyzed single-nucleus transcriptomic profiles (snRNA-seq) of superior parietal
cortex tissue samples from donors of the Knight Alzheimer Disease Research Center (Knight
ADRC) and the Dominantly Inherited Alzheimer Network (DIAN) (*16*). This dataset
encompasses different AD subtypes, including sporadic AD and autosomal dominant AD, with
donors distributed in a broad spectrum of neuropathological states and genetic backgrounds,
including carriers of *TREM2* risk variants **(Figure 1a, Supp. Table 1)**. We identified patterns of
ligand-receptor gene expression across cell type pairs and their corresponding transcriptional
states using CellPhoneDB (*25*), which has been successfully used to predict patterns of cellular
crosstalk. We performed the analyses to detect condition-specific crosstalk interactions by
stratifying brains by disease status and genetic group. In total, we identified between 961 and
1,600 (median 1,521) significant (Bonferroni-corrected $p < 0.05$) crosstalk interactions between
cellular state pairs across all donor categories **(Figure 1b; Supp. Figure 1a; Supp. Table 2)**.
Furthermore, the observed crosstalk patterns were highly robust to dataset downsampling,
including removing a subset of donors and using 100 barcodes or less per transcriptional state
(fraction of replication 0.81 to 0.86 across donor categories, median = 0.84; **Supp. Figure 1b-**
c). These results indicate that the crosstalk interaction patterns identified using CellPhoneDB
were not driven by technical factors such as the number of donors, nuclei, and barcode
sequencing depth.

We compared the crosstalk patterns across donor categories to identify AD-related global
changes. Overall, crosstalk interactions involving microglia were significantly increased in all AD
donors compared to controls (OR=1.21, adj. p=5.22e-4), and we detected a significant decrease
in crosstalk involving excitatory neurons in sporadic AD donors (OR = 0.82, adj. p=0.03; **Supp.**
Figure 2). These results are consistent with neuronal loss and the critical role of microglia in AD
onset. We next performed a functional enrichment of the genes associated with the up- and
down-regulated crosstalk interactions in each cell type to determine the biological pathways
perturbed in AD. Consistent with significant changes in cellular homeostasis in AD brains, we
observed changes in the cellular crosstalk patterns between neuropath-free control and AD
donors. The crosstalk interactions up-regulated in AD were significantly enriched for pathways
indicative of immune activation and cellular stress (**Figure 1c**). Interestingly, we observed a
significant increase in ephrin signaling in autosomal dominant AD donors (p = 1.48e-12). This
finding agrees with the previously described role of this signaling pathway in neuroinflammation
and neurodegeneration (*5*). Aiming to understand how each cell type likely contributes to AD
crosstalk, we also performed a similar functional enrichment analysis focusing on cell types to
identify crosstalk interactions changed in AD. We observed a potential role for multiple cell types
contributing to immune activation and neuronal stress (**Supp. Figure 3**). Together, these results
indicate that AD involves widespread dysregulation of homeostatic cross-cellular signaling
pathways.

Our initial analysis identified thousands of significant crosstalk interactions across all cell types
and donor categories. We next sought to focus on interactions involving genes linked to AD by
genetic and functional studies, which we hypothesized were more likely to be early drivers of
AD. (**Supp. Table 3**). We identified 90 crosstalk interactions directly involving an AD gene as
either the ligand or receptor, of which 34 were considered significant by CellPhoneDB in at least
one cell type pair (**Figure 1d**). Microglia had the highest enrichment for crosstalk interactions
involving AD genes ($\log_2$ OR range: 0.85 to 1.76, median = 1.23, *p* range: 4.69e-3 to 3.08e-8,
median = 2.68e-4; **Figure 1e**), and most of these interactions (64.9%) involved microglia as the
likely receptor cell type. These enrichment patterns were robust to the observed increase in the
number of crosstalk interactions involving microglia in AD vs. controls and the overall higher
representation of AD genes in microglia (**Supp. Figure 4**, Methods).

We next analyzed which cell type(s) were most likely to interact with microglia crosstalk by
conditioning the previous analyses on microglia. We observed that excitatory neurons had the
highest enrichment point estimates to partner with microglia for crosstalk interactions involving
AD genes mainly due to pre-symptomatics ($\log_2$ OR = 1.98, p = 0.035; **Figure 1f**). We replicated
these analyses in three case-control snRNA-seq datasets of the prefrontal cortex region from
the South West Dementia Brain Bank (SWDBB), the Rush ADRC, and the University of
California Irvine Institute for Memory Impairments and Neurological Disorders (UCI MIND)
ADRC (**Supp. Table 4**) (*18*, *26*, *19*). We observed the same strong enrichment of microglia for
crosstalk interactions involving AD genes in all three studies (**Supp. Figure 5**). We observed
differences across studies regarding which neuronal cell type was most enriched for crosstalk
interactions with microglia involving AD genes. Nonetheless, the enrichment direction of effect
for excitatory neurons agreed with our findings in all studies. These results suggest that a

subset of genes previously implicated in AD mediate cell signaling pathways between neurons
and microglia.

To further validate if the previous results were not driven by an overrepresentation bias of
microglia interactions in the CellPhoneDB database, we used the control donors to calculate the
enrichment of crosstalk interactions in genes found by GWAS for other neurological traits
(**Supp. Table 5**). If the crosstalk interactions were biased towards one or a subset of cell types
due to database overrepresentation, our expectation would be that we would observe similar
enrichment patterns across traits. On the contrary, we observed distinct crosstalk enrichment
patterns across neurological traits (**Figure 1g**). Crosstalk interactions involving oligodendrocyte
precursors were significantly enriched for genes from one schizophrenia GWAS (OR = 2.64,
adj. p = 3.12e-5). We also observed nominally significant enrichment for astrocyte crosstalk
interactions for genes from a Parkinson's disease GWAS (OR = 2.45, p = 0.029). Importantly,
we replicated the enrichment for microglia crosstalk interactions in genes from two independent
AD GWAS (ORs = 3.39 and 3.07, adj. p = 3.16e-4 and 0.0023). We next used a strategy to
nominate AD genes based on snATAC-seq co-accessibility between the gene promoter regions
with a fine-mapped (*14*) AD GWAS variant (see Methods). This rigorous approach nominates
candidate AD genes and their corresponding cell types based on direct evidence from functional
genomics. We observed an even higher enrichment for microglia crosstalk interactions in AD
(OR = 9.07, p = 1.24e-15; **Figure 1h**). In combination, these results indicate that the observed
crosstalk enrichment patterns were not artificially biased toward any cell type. Furthermore,
these analyses highlight that our crosstalk framework can be used to understand biological
processes associated with GWAS variants for multiple neurological diseases.

Crosstalk interactions between microglia and excitatory neurons are predicted to
modulate genes implicated in AD

We next sought to investigate how the crosstalk signals between neurons and microglia
modulate downstream microglia gene regulatory networks. Using a system biology approach
based on extending the functionality of the CytoTalk software (*27*) (see methods), we inferred
the gene co-expression networks upstream of the ligands and downstream of the receptors.
CytoTalk is complementary to CellPhoneDB, which does not inform the biological processes
likely downstream of crosstalk interactions. Using CytoTalk, we reconstructed the gene
regulatory network associated with crosstalk interactions between excitatory neurons and
microglia (**Figure 2a-b**). Strikingly, the microglia co-expression network downstream of the
neuron-microglia crosstalk interactions was enriched for genes previously associated with AD
(cases OR range = 1.99 to 3.52, median = 3.06, adj. p range = 0.017 to 2.86e-8, median =
1.06e-4; **Figure 2c**). In addition, the microglia crosstalk network identified by CytoTalk was
overall enriched for immune processes, including phagocytosis and cytokine production, among
others (**Figure 2d**), consistent with neuron-microglia crosstalk interactions modulating microglia
activation states (*28*). These results indicate that neuron-microglia crosstalk interactions
propagate signals that modulate genes previously implicated in AD, including those involved in
regulating microglial activation.

The SEMA6D-TREM2 crosstalk axis is predicted to modulate microglia activation and to
be disrupted in late-stage AD

Among the seven crosstalk interactions prioritized by CytoTalk based on the co-expression
network topology, we identified the crosstalk interaction between neuronal ligand semaphorin
6D (SEMA6D) and TREM2/TYROBP (**Figure 2b**). In the brain, semaphorin signaling was
initially described as a mediator of axon guidance via the plexin family of receptors (*29*).
However, an increasing body of evidence indicates that these molecules are also involved in
immune responses. Notably, a study in bone marrow-derived macrophages discovered that the
SEMA6D-PLXNA1 interaction induces macrophage activation in a TREM2-dependent manner
via activation of DAP12 (*7*). Importantly, this study found that TREM2, PLXNA1, and DAP12 co-
immunoprecipitate in a TREM2-dependent manner, consistent with the formation of a complex
(*30, 31*). Furthermore, additional studies also linked plexin-semaphorin signaling to immune
activation and neurodegeneration (*5, 22–24, 32*). Given this role for SEMA6D in the immune
activation of myeloid cells, we aimed to test the potential role of the SEMA6D-TREM2/PLXNA1
signaling axis in AD.

Using a clustering algorithm to divide the microglia crosstalk network into sub-networks (see
methods), we identified a microglia sub-network comprised of a large clique of genes highly
connected to *TREM2* and *TYROBP*. This TREM2 sub-network was predicted to be modulated
by neuronal SEMA6D (**Figure 2e**). Furthermore, the TREM2 sub-network was enriched for
microglia activation genes, indicating that we recapitulated the well-established link between
TREM2 and microglia activation (*33*) through an unsupervised approach (**Figure 2d**).
Interestingly, in addition to genes linked to microglia activation, the TREM2 crosstalk sub-
network included *APOE* and *HLA* genes previously reported as AD risk genes. The connection
between *TREM2* and *APOE* is consistent with studies showing that APOE is a TREM2 ligand
(*34, 35*) and suggests that the PLXNA1-SEMA6D crosstalk interaction modulates AD risk
genes.

To validate these results, we repeated the CytoTalk analyses in the snRNA-seq studies from the
SWDBB, Rush ADRC, and UCI MIND ADRC cohorts (*18, 19, 26*). Consistent with our original
results, CytoTalk prioritized the SEMA6D-TREM2 signaling axis mediating the crosstalk
interactions between exc. neurons and microglia in all three cohorts (**Supp. Figure 6a**). Finally,
we determined that the TREM2 sub-network and its predicted modulation by SEMA6D were
robust to the choice of donors and number of nuclei (**Supp. Figure 6b**). Together, these results
reinforce that the unsupervised methodological approach in this study identified core elements
of microglia gene regulation, which are predicted to be modulated by neuron-microglia cellular
crosstalk interactions and are robust to brain donor and region.

We next sought to understand how the TREM2 sub-network related to AD progression. We
leveraged the range of neuropathological states in our dataset to develop a statistical framework
to test the association of this sub-network with the Braak stages, controlling for genetic and
other confounding factors (see methods). Using this approach, we determined that the
expression of the TREM2 sub-network was negatively associated with the Braak stages (beta =
-0.31, adj. p = 4.32e-57). To contextualize this result, we calculated the association of all
neuron-microglia crosstalk sub-networks with the Braak stages. Strikingly, the majority (11 of
14) of the microglia crosstalk sub-networks were negatively associated with the Braak stages,
and the TREM2 sub-network was among the most negatively associated with Braak stages
(**Figure 2f**). These results suggest that neuron-microglia crosstalk interactions and their

downstream targets in microglia are impaired in the later stages of AD. Next, we sought to
determine if SEMA6D is a potential modulator of the TREM2 microglia crosstalk sub-network.
We reasoned that if this were the case, the neuronal SEMA6D crosstalk sub-network direction
of association with the Braak stages would agree with the TREM2 sub-network. Indeed, the
neuronal SEMA6D sub-network was negatively associated with the Braak stages (beta = -0.09,
adj. p = 1.63e-04; **Figure 2h**). Therefore, our findings indicate that the biological processes
associated with the SEMA6D-PLXNA1 neuron-microglia crosstalk interactions are disrupted in
AD and likely confer AD resilience by modulating TREM2-dependent microglia activation.

Multiple microglia co-expression sub-networks are disrupted during AD progression

We next adapted our network analysis framework to analyze the microglia co-expression
network aiming to gain insights into the role of microglia in AD. This expanded approach allowed
us to test the association of sub-networks not predicted to be directly involved in neuron-
microglia crosstalk with the Braak stages (see methods). Using the entire reconstructed
microglia co-expression network, we identified 360 sub-networks and recapitulated several
crosstalk sub-networks identified previously with CytoTalk, including the TREM2 sub-network
(**Supp. Figure 7**). These microglia sub-networks were divided between positive and negative
associations with the Braak stages (**Figure 2g**). Consistent with the well-established roles of
PSEN1 and *APP* in AD onset (*36*), the *PSEN1* and *APP* co-expression sub-networks were
among the most positively correlated with the Braak stages (betas = 0.32 and 0.16, adj. p =
8.25e-57 and 2.17e-13, respectively). In contrast, the sub-network of *SORL1*, a gene associated
with protective roles in AD (*37*, *38*), was negatively associated with the Braak stages (beta = -
0.15, adj. p = 3.52e-11). Interestingly, our unsupervised approach identified two separate sub-
networks with opposing directions of effect for the genes in the MS4A locus, which genetically
controls soluble TREM2 levels (*39*) (*MS4A4A* and *MS4A6A* betas = -0.11 and 0.19, adj. p =
2.87e-06 and 2.93e-20, respectively). This result suggests that the MS4A genetic signal
regulates at least two independent biological processes, as we reported previously (*39*). Within
the context of all microglia genes, the TREM2 sub-network was among the most negatively
associated with the Braak stages, consistent with our previous analysis using the prioritized
crosstalk network from CytoTalk. These results indicate that multiple biological pathways
downstream of microglia-neuronal crosstalk are disrupted in AD. Furthermore, our unsupervised
computational framework identifies impaired TREM2-dependent microglia activation as a
contributing factor to AD progression.

**SEMA6D treatment potentiates immune activation in iPSC-derived microglia in a TREM2-
dependent manner**

Microglia, among other functions, regulate brain homeostasis through phagocytic activity and
modulate neuroinflammation by releasing inflammatory cytokines, such as TNF-α and IL-6 (*40*).
We generated iMGL to examine if SEMA6D modulates iMGL functions (**Supp. Figure 8a**, see
methods). (*41*) iMGL exhibited established microglia markers, including TREM2, IBA1, and
TMEM119 (**Supp. Figure 8b**). We also generated *TREM2* knockout (KO) human iPSCs using
CRISPR/Cas9 to examine the role of the SEMA6D-TREM2 signaling axis on microglia function
(**Supp. Figure 8c**). We verified the loss of TREM2 expression at the protein level in the KO cell
line by Western blot analysis (**Supp. Figure 8d**).

To determine if SEMA6D can regulate iMGL phagocytic activity and if this is TREM2 dependent,
we treated WT and KO iMGL with SEMA6D protein and measured the degree of phagocytic
activity on pHrodo-labeled human synaptosomes (**Supp. Figure 9**). Live-cell imaging over 24
hours showed increased phagocytosis of WT iMGL treated with SEMA6D starting after 6 hours.
TREM2 KO iMGL treated with SEMA6D did not demonstrate increased phagocytosis compared
to untreated TREM2 KO cells (**Figure 3a-b**). These results indicate that SEMA6D can modulate
iMGL phagocytosis in a TREM2-dependent manner. We analyzed conditioned media using a
multiplex immunoassay to determine if SEMA6D can regulate iMGL cytokine release mediated
by TREM2from control or SEMA6D-treated iMGL. We found that SEMA6D increased the
secretion of TNF-α and IL-6 in WT iMGL. SEMA6D did not increase cytokine release in TREM2
KO iMGL (**Figure 3c**). SEMA6D regulation of iMGL cytokine release also appears to be TREM2
dependent.

TREM2 mediates signaling through the adaptor protein TYROBP (DAP12). Activation of TREM2
results in tyrosine phosphorylation within the ITAM motif and subsequent SYK phosphorylation
(**Figure 3f**) (*42*). To determine if SEMA6D activates TREM2 downstream signaling, we analyzed
WT and KO iMGL lysates for phosphorylated SYK expression (p-SYK) normalized to total SYK
expression (SYK). Treatment of WT iMGL with SEMA6D induced SYK phosphorylation, while
KO iMGL did not. (**Figure 3d**)-**e**.). This demonstrates that SEMA6D can activate TREM2
signaling and suggests that SEMA6D signals through the PLXNA1/TREM2/DAP12 complex.

In parallel, to understand the regulatory changes induced by SEMA6D treatment in iMGL, we
generated bulk RNA-seq data for the SEMA6D-treated and control WT and KO iMGL (**Supp.**
Figure 10). We observed significant changes in gene expression associated with TREM2 KO
(**Supp. Figure 10a-b**). As expected, *TREM2* was among the most downregulated genes in the
KO iMGL (**Supp. Figure 10a**). Strikingly, we observed robust transcriptional changes in WT
SEMA6D-treated iMGL but not *TREM2* KO iMGL (**Figure 3g; Supp. Figure 10a-b**), supporting
a pivotal role for TREM2 in mediating SEMA6D-PLXNA1 crosstalk signaling. To further
understand how TREM2 modulates this crosstalk, we focused on the *TREM2* co-expression
sub-network we predicted from the snRNA-seq data using CytoTalk. The *TREM2* sub-network
had significantly lower expression in the untreated KO iMGL compared to WT, consistent with
TREM2 being a key regulator of this sub-network. In line with this interpretation, the TREM2
sub-network was activated by SEMA6D treatment in the WT iMGL but significantly less so in the
KO iMGL (**Figure 3h-i**).

To gain further insights into how SEMA6D treatment regulates microglial transcriptional
programs, we checked biologically relevant transcriptional signatures previously described in
microglia. These included genes up-regulated in phagocytosing microglia (*43*) and those up-
regulated in response to LPS treatment (*20*). As control signatures, we included genes down-
regulated in TREM2 KO iMGL (*44*) and a set of randomly selected genes for which we would
not expect any concerted transcriptional changes. We observed the strongest effects (Log2FC
>0.4) of SEMA6D treatment in the phagocytosing microglia gene signature (**Figure 3i**),
indicating that SEMA6D activates genes crucial to phagocytosis in the WT but not TREM2 KO
iMGL. Notably, TREM2, APOE, and RPS19 are among the most up-regulated genes by
SEMA6D treatment in the WT iMGL, which were either present in the phagocytosing microglia
gene signature or correspond to genes previously linked to microglia activation (*45, 17*) (**Figure**

3j). Our results indicate that SEMA6D-TREM2 crosstalk signaling induces a TREM2-mediated
cascade of transcriptional events resulting in microglia activation.

Discussion

In this study, we leveraged single-nucleus gene expression profiles from a diverse cohort of
brain donors to systematically dissect the contribution of cross-cellular signaling (cellular
crosstalk) networks to AD progression. Our data-driven approach to identifying active crosstalk
interactions and reconstructing their corresponding downstream pathways provides additional
evidence that disrupted cellular crosstalk networks contribute to neurodegeneration. One
remarkable finding from our study is that a significant portion of AD risk genes are either directly
involved in crosstalk interactions or immediately downstream of microglia interactions. These
results highlight the difficulty of characterizing the prominent role of microglia in AD progression,
as these cells integrate complex signals originating in other brain cell types. Specifically, our
results support that dysregulation in the intricate signaling between neurons and microglia is
linked to AD progression. Therefore, focusing on cellular crosstalk networks can provide further
functional context to understand the biology of genes associated with AD risk in a cell non-
autonomous and autonomous manner.

Among the interactions we detected between neurons and microglia, we identified a novel
functional link between neuronal SEMA6D and microglial TREM2. Furthermore, by leveraging
iMGL, we demonstrate that SEMA6D signaling induces a TREM2-dependent microglia
activation phenotype transcriptionally similar to phagocytosing microglia (*43*). These results,
combined with our observation that the transcriptional networks upstream and downstream of
the SEMA6D-PLXNA1 interaction are downregulated in AD, suggest that loss of this crosstalk
interaction exacerbates the deleterious processes occurring in the later stages of this disease.

We used iMGL to demonstrate that SEMA6D promotes microglial phagocytosis and
inflammatory cytokine release. SEMA6D-induced microglial and downstream SYK activation
were not observed in the TREM2 KO iMGL. Previous reports showed that SEMA6D promotes
peripheral dendritic cell activation and osteoclast differentiation via the receptor complex
harboring PLXNA1 and TREM2 (*46*). Thus, it is conceivable that SEMA6D functions as a natural
ligand for the PLXNA1/TREM2 co-receptor and enhances TREM2 signaling in human microglia.
Therefore, neuronal SEMA6D could influence functional properties and fate via the stimulation
of TREM2-dependent intracellular signaling and induction of TREM2-network gene expression.
Nevertheless, it remains to be determined if SEMA6D induces an anti-inflammatory state of
microglia that might be beneficial in clearing neuropathological changes associated with AD.

SEMA6D also regulates lipid metabolism and polarization of macrophages via the interaction
with another class A plexin family member, PlexinA4 (*47, 48*). Several coding variants in
PlexinA4 are associated with AD risk (*48, 49*) and modulate both amyloid and tau pathology
(*50*). Thus, semaphorin-plexin signaling may play a fundamental role in regulating the functional
interactions between glutaminergic neurons and a subset of microglia and may be perturbed in
AD.

While this study focused on a restricted subset of crosstalk interactions involving microglia and
neurons, our systematic characterization of cross-cellular signaling patterns identified thousands

of candidate interactions involving all brain cell types represented in our snRNA-seq data. Some of these interactions warrant further investigation. For example, the interleukin receptor IL1RAP has been previously implicated in genetic studies of AD endophenotypes (*3*, *51–53*), and the contribution of IL-1 signaling to neurodegenerative diseases has been well-established (*54–56*). In line with these previous findings, we identified the *IL1RAP* sub-network in microglia as the most negatively associated with the Braak stages. These results suggest that disruption of the IL-1 signaling pathway is likely to be involved in AD progression. More broadly, the IL1RAP case highlights that the continued exploration of brain crosstalk networks identified in this study will yield valuable biological insights into AD biology. Finally, we identified distinct crosstalk enrichment patterns for genes identified by genetic studies of other neurological traits or diseases. These results emphasize that cellular crosstalk is an integral part of normal brain physiology, and accounting for this regulatory layer will help contextualize how candidate disease risk genes fit into biological pathways. Together, our findings offer a strong rationale that the systematic characterization of cellular crosstalk networks is a viable strategy for gaining insight into the biology of neurodegenerative diseases.

Acknowledgments

Data collection and sharing for this project was supported by The Dominantly Inherited Alzheimer Network (DIAN, U19AG032438), funded by the National Institute on Aging (NIA), the Alzheimer's Association (SG-20-690363-DIAN), the German Center for Neurodegenerative Diseases (DZNE), Raul Carrea Institute for Neurological Research (FLENI), Partial support by the Research and Development Grants for Dementia from Japan Agency for Medical Research and Development, AMED, and the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI), Spanish Institute of Health Carlos III (ISCIII), Canadian Institutes of Health Research (CIHR), Canadian Consortium of Neurodegeneration and Aging, Brain Canada Foundation, and Fonds de Recherche du Québec – Santé. This manuscript has been reviewed by DIAN Study investigatorsfor scientific content and consistency of data interpretation with previous DIAN Study publications. We acknowledge the altruism of the participants and their families and the contributions of the DIAN research and support staff at each of the participating sites for their contributions to this study. This research was supported by NIH grants R01 AG067606 (to T.-W.K), AG072464 (CMK, OH), NINDS R01NS118146 and R21NS127211 (to BAB), NIH AG062734 (CMK), and R56AG067764 (GS). Additional funding from the Chan Zuckerberg Initiative (CMK, OH, CC). We thank Barbara Corneo for advice on iPSC protocols, and Barbara Corneo, Wei Wang; Caisheng (Luke) Lu, MD, Ph.D. for assisting FACS sorting. IMGL sequencing and preliminary data processing was conducted by Novogene.

Author contributions

OOO

Competing interests

T.-W.K. is a cofounder of BL Melanis Co. Ltd.

416
417
Figure 1: Overview of predicted cellular crosstalk patterns in human brains during health and AD. a) Uniform Manifold Approximation and Projection (UMAP) representation of the snRNA-seq dataset. **b)** Number of unique significant CellPhoneDB interactions detected involving each cell type as either the ligand or receptor across donor categories. **c)** Gene ontology enrichments of crosstalk interactions up- or down-regulated between each donor category versus controls. **d)** Cellular crosstalk interactions involving one AD gene as either ligand or receptor across all cellular state pairs. **e)** Enrichment of AD crosstalk interactions involving each cell type in each donor category. **f)** Enrichment of AD crosstalk interactions between microglia and other cell types (labeled in the y-axis). **g)** Crosstalk enrichments for each cell type (control donors only) in genes nominated by GWAS in other neuropsychiatric traits. **h)** Similar to (**g**) but using only AD genes supported by snATAC-seq co-accessibility (Methods). Odds ratios in (**e-h**) were calculated using Fisher exact tests (Methods).

Figure 2: Crosstalk interactions between neurons and microglia are predicted to modulate AD risk genes. a) Microglia-excitatory neuron crosstalk network inferred by CytoTalk. **b)** Prioritized crosstalk interactions using CytoTalk. **c)** Enrichment of AD genes in the microglia crosstalk network across donor categories. **d**) GO enrichments for the genes participating in microglia and excitatory neurons crosstalk networks. **e)** Predicted TREM2 crosstalk subnetwork **f)** Association of microglia crosstalk subnetworks with Braak stages. **g)** Association of all microglia subnetworks with Braak stages. **h)** Association excitatory neuron crosstalk subnetworks with the Braak stages.

Figure 3. SEMA6D treatment induces microglia activation in a TREM2-dependent manner.
a) Representative graph of phagocytosis of synaptosomes by WT or TREM2 KO iMGL treated
with Sema6D (5 µg 5ug/ml). **b)** Quantification of phagocytosis of synaptosomes by WT or
TREM2 KO iMGL treated with Sema6D at 24 hr. Data represent three independent experiments
performed in triplicate; mean ± SEM unpaired two-tailed t-test, $^{**}p < 0.01$ at 24 hrs. **c)**
Quantification of media from WT or KO iMGL treated with Sema6D (10 µg/ml) for cytokine
release of TNF-α and IL-6, using the Mesoscale V-plex neuroinflammation panel. **d)**
Representative western blot analysis of p-SYK and total SYK in WT and KO iMGL treated with
Sema6D (10 µg/ml). β-actin as loading control **e)** Quantification of p-SYK/total SYK western blot
analysis, n=3 **(d)**. **f)** Schematic of the proposed PlexinA-TREM2/DAP12 signaling complex. **g)**
RNA-seq effect size distribution on SEMA6D-treated iMGL (WT and *TREM2* KO). **h)**
Transcriptional changes associated with SEMA6D treatment (WT or *TREM2* KO vs. control) or
TREM2 knockout (KO vs. WT, no treatment) in the TREM2 crosstalk sub-network. Only TREM2
sub-network genes differentially expressed in at least one comparison were included. **i)**
Transcriptional effects of the same conditions as **(b)** across biologically relevant gene
signatures. Background corresponds to 500 randomly selected genes. Dots correspond to a 1%
FDR significance threshold for comparing the effect size distribution to the corresponding
background in each comparison (Wilcoxon rank-sum test). **j)** Transcriptional effects of the
conditions across highly differentially expressed genes from a subset of signatures in **(i)**.

References

References

1. D. P. Schafer, E. K. Lehrman, A. G. Kautzman, R. Koyama, A. R. Mardinly, R. Yamasaki,
R. M. Ransohoff, M. E. Greenberg, B. A. Barres, B. Stevens, Microglia sculpt postnatal
neural circuits in an activity and complement-dependent manner. *Neuron*. **74**, 691–705
(2012).

2. D. Fard, L. Tamagnone, Semaphorins in health and disease. *Cytokine Growth Factor Rev*.
57, 55–63 (2021).

3. V. K. Ramanan, S. L. Risacher, K. Nho, S. Kim, L. Shen, B. C. McDonald, K. K. Yoder, G.
D. Hutchins, J. D. West, E. F. Tallman, S. Gao, T. M. Foroud, M. R. Farlow, P. L. De Jager,
D. A. Bennett, P. S. Aisen, R. C. Petersen, C. R. Jack, A. W. Toga, R. C. Green, W. J.
Jagust, M. W. Weiner, A. J. Saykin, Alzheimer's Disease Neuroimaging Initiative (ADNI),
GWAS of longitudinal amyloid accumulation on 18F-florbetapir PET in Alzheimer's disease
implicates microglial activation gene IL1RAP. *Brain*. **138**, 3076–3088 (2015).

4. C. S. McAlpine, J. Park, A. Griciuc, E. Kim, S. H. Choi, Y. Iwamoto, M. G. Kiss, K. A.
Christie, C. Vinegoni, W. C. Poller, J. E. Mindur, C. T. Chan, S. He, H. Janssen, L. P.
Wong, J. Downey, S. Singh, A. Anzai, F. Kahles, M. Jorfi, P. F. Feruglio, R. I. Sadreyev, R.
Weissleder, B. P. Kleinstiver, M. Nahrendorf, R. E. Tanzi, F. K. Swirski, Astrocytic
interleukin-3 programs microglia and limits Alzheimer's disease. *Nature* (2021),
doi:10.1038/s41586-021-03734-6.

5. I. C. Clark, C. Gutiérrez-Vázquez, M. A. Wheeler, Z. Li, V. Rothhammer, M. Linnerbauer, L.
M. Sanmarco, L. Guo, M. Blain, S. E. J. Zandee, C.-C. Chao, K. V. Batterman, M.
Schwabenland, P. Lotfy, A. Tejeda-Velarde, P. Hewson, C. Manganeli Polonio, M. W.
Shultis, Y. Salem, E. C. Tjon, P. H. Fonseca-Castro, D. M. Borucki, K. Alves de Lima, A.
Plasencia, A. R. Abate, D. L. Rosene, K. J. Hodgetts, M. Prinz, J. P. Antel, A. Prat, F. J.
Quintana, Barcoded viral tracing of single-cell interactions in central nervous system
inflammation. *Science*. **372**, eabf1230 (2021).

6. H. Lian, A. Litvinchuk, A. C.-A. Chiang, N. Aithmitti, J. L. Jankowsky, H. Zheng, Astrocyte-
Microglia Cross Talk through Complement Activation Modulates Amyloid Pathology in
Mouse Models of Alzheimer's Disease. *J Neurosci*. **36**, 577–589 (2016).

7. N. Takegahara, H. Takamatsu, T. Toyofuku, T. Tsujimura, T. Okuno, K. Yukawa, M. Mizui,
M. Yamamoto, D. V. R. Prasad, K. Suzuki, M. Ishii, K. Terai, M. Moriya, Y. Nakatsuji, S.
Sakoda, S. Sato, S. Akira, K. Takeda, M. Inui, T. Takai, M. Ikawa, M. Okabe, A.
Kumanogoh, H. Kikutani, Plexin-A1 and its interaction with DAP12 in immune responses
and bone homeostasis. *Nat Cell Biol*. **8**, 615–622 (2006).

8. T. Ito, K. Yoshida, T. Negishi, M. Miyajima, H. Takamatsu, H. Kikutani, A. Kumanogoh, K.
Yukawa, Plexin-A1 is required for Toll-like receptor-mediated microglial activation in the
development of lipopolysaccharide-induced encephalopathy. *Int J Mol Med*. **33**, 1122–
1130 (2014).

9. H. Yin, A. D. Flynn, Drugging Membrane Protein Interactions. *Annu Rev Biomed Eng*. **18**,
51–76 (2016).

10.   J. P. Overington, B. Al-Lazikani, A. L. Hopkins, How many drug targets are there? *Nat Rev*
502   *Drug Discov.* **5**, 993–996 (2006).

503   11.   T. Worzfeld, S. Offermanns, Semaphorins and plexins as therapeutic targets. *Nat Rev*
504   *Drug Discov.* **13**, 603–621 (2014).

505   12.   J. C. Lambert, C. A. Ibrahim-Verbaas, D. Harold, A. C. Naj, R. Sims, C. Bellenguez, A. L.
506   DeStafano, J. C. Bis, G. W. Beecham, B. Grenier-Boley, G. Russo, T. A. Thorton-Wells, N.
507   Jones, A. V. Smith, V. Chouraki, C. Thomas, M. A. Ikram, D. Zelenika, B. N. Vardarajan, Y.
508   Kamatani, C. F. Lin, A. Gerrish, H. Schmidt, B. Kunkle, M. L. Dunstan, A. Ruiz, M. T.
509   Bihoreau, S. H. Choi, C. Reitz, F. Pasquier, C. Cruchaga, D. Craig, N. Amin, C. Berr, O. L.
510   Lopez, P. L. De Jager, V. Deramecourt, J. A. Johnston, D. Evans, S. Lovestone, L.
511   Letenneur, F. J. Morón, D. C. Rubinsztein, G. Eiriksdottir, K. Sleegers, A. M. Goate, N.
512   Fiévet, M. W. Huentelman, M. Gill, K. Brown, M. I. Kamboh, L. Keller, P. Barberger-
513   Gateau, B. McGuiness, E. B. Larson, R. Green, A. J. Myers, C. Dufouil, S. Todd, D.
514   Wallon, S. Love, E. Rogaeva, J. Gallacher, P. St George-Hyslop, J. Clarimon, A. Lleo, A.
515   Bayer, D. W. Tsuang, L. Yu, M. Tsolaki, P. Bossù, G. Spalletta, P. Proitsi, J. Collinge, S.
516   Sorbi, F. Sanchez-Garcia, N. C. Fox, J. Hardy, M. C. Deniz Naranjo, P. Bosco, R. Clarke,
517   C. Brayne, D. Galimberti, M. Mancuso, F. Matthews, European Alzheimer's Disease
518   Initiative (EADI), Genetic and Environmental Risk in Alzheimer's Disease, Alzheimer's
519   Disease Genetic Consortium, Cohorts for Heart and Aging Research in Genomic
520   Epidemiology, S. Moebus, P. Mecocci, M. Del Zompo, W. Maier, H. Hampel, A. Pilotto, M.
521   Bullido, F. Panza, P. Caffarra, B. Nacmias, J. R. Gilbert, M. Mayhaus, L. Lannefelt, H.
522   Hakonarson, S. Pichler, M. M. Carrasquillo, M. Ingelsson, D. Beekly, V. Alvarez, F. Zou, O.
523   Valladares, S. G. Younkin, E. Coto, K. L. Hamilton-Nelson, W. Gu, C. Razquin, P. Pastor,
524   I. Mateo, M. J. Owen, K. M. Faber, P. V. Jonsson, O. Combarros, M. C. O'Donovan, L. B.
525   Cantwell, H. Soininen, D. Blacker, S. Mead, T. H. Mosley, D. A. Bennett, T. B. Harris, L.
526   Fratiglioni, C. Holmes, R. F. de Bruijn, P. Passmore, T. J. Montine, K. Bettens, J. I. Rotter,
527   A. Brice, K. Morgan, T. M. Foroud, W. A. Kukull, D. Hannequin, J. F. Powell, M. A. Nalls, K.
528   Ritchie, K. L. Lunetta, J. S. Kauwe, E. Boerwinkle, M. Riemenschneider, M. Boada, M.
529   Hiltuenen, E. R. Martin, R. Schmidt, D. Rujescu, L. S. Wang, J. F. Dartigues, R. Mayeux,
530   C. Tzourio, A. Hofman, M. M. Nöthen, C. Graff, B. M. Psaty, L. Jones, J. L. Haines, P. A.
531   Holmans, M. Lathrop, M. A. Pericak-Vance, L. J. Launer, L. A. Farrer, C. M. van Duijn, C.
532   Van Broeckhoven, V. Moskvina, S. Seshadri, J. Williams, G. D. Schellenberg, P. Amouyel,
533   Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's
534   disease. *Nat Genet.* **45**, 1452–1458 (2013).

535   13.   B. W. Kunkle, B. Grenier-Boley, R. Sims, J. C. Bis, V. Damotte, A. C. Naj, A. Boland, M.
536   Vronskaya, S. J. van der Lee, A. Amlie-Wolf, C. Bellenguez, A. Frizatti, V. Chouraki, E. R.
537   Martin, K. Sleegers, N. Badarinarayan, J. Jakobsdottir, K. L. Hamilton-Nelson, S. Moreno-
538   Grau, R. Olaso, R. Raybould, Y. Chen, A. B. Kuzma, M. Hiltunen, T. Morgan, S. Ahmad, B.
539   N. Vardarajan, J. Epelbaum, P. Hoffmann, M. Boada, G. W. Beecham, J.-G. Garnier, D.
540   Harold, A. L. Fitzpatrick, O. Valladares, M.-L. Moutet, A. Gerrish, A. V. Smith, L. Qu, D.
541   Bacq, N. Denning, X. Jian, Y. Zhao, M. Del Zompo, N. C. Fox, S.-H. Choi, I. Mateo, J. T.
542   Hughes, H. H. Adams, J. Malamon, F. Sanchez-Garcia, Y. Patel, J. A. Brody, B. A.
543   Dombroski, M. C. D. Naranjo, M. Daniilidou, G. Eiriksdottir, S. Mukherjee, D. Wallon, J.
544   Uphill, T. Aspelund, L. B. Cantwell, F. Garzia, D. Galimberti, E. Hofer, M. Butkiewicz, B.
545   Fin, E. Scarpini, C. Sarnowski, W. S. Bush, S. Meslage, J. Kornhuber, C. C. White, Y.
546   Song, R. C. Barber, S. Engelborghs, S. Sordon, D. Voijnovic, P. M. Adams, R.
547   Vandenberghe, M. Mayhaus, L. A. Cupples, M. S. Albert, P. P. De Deyn, W. Gu, J. J.
548   Himali, D. Beekly, A. Squassina, A. M. Hartmann, A. Orellana, D. Blacker, E. Rodriguez-

Rodriguez, S. Lovestone, M. E. Garcia, R. S. Doody, C. Munoz-Fernadez, R. Sussams, H.
Lin, T. J. Fairchild, Y. A. Benito, C. Holmes, H. Karamujić-Čomić, M. P. Frosch, H.
Thonberg, W. Maier, G. Roshchupkin, B. Ghetti, V. Giedraitis, A. Kawalia, S. Li, R. M.
Huebinger, L. Kilander, S. Moebus, I. Hernández, M. I. Kamboh, R. Brundin, J. Turton, Q.
Yang, M. J. Katz, L. Concari, J. Lord, A. S. Beiser, C. D. Keene, S. Helisalmi, I.
Kloszewska, W. A. Kukull, A. M. Koivisto, A. Lynch, L. Tarraga, E. B. Larson, A.
Haapasalo, B. Lawlor, T. H. Mosley, R. B. Lipton, V. Solfrizzi, M. Gill, W. T. Longstreth, T.
J. Montine, V. Frisardi, M. Diez-Fairen, F. Rivadeneira, R. C. Petersen, V. Deramecourt, I.
Alvarez, F. Salani, A. Ciaramella, E. Boerwinkle, E. M. Reiman, N. Fievet, J. I. Rotter, J. S.
Reisch, O. Hanon, C. Cupidi, A. G. Andre Uitterlinden, D. R. Royall, C. Dufouil, R. G.
Maletta, I. de Rojas, M. Sano, A. Brice, R. Cecchetti, P. S. George-Hyslop, K. Ritchie, M.
Tsolaki, D. W. Tsuang, B. Dubois, D. Craig, C.-K. Wu, H. Soininen, D. Avramidou, R. L.
Albin, L. Fratiglioni, A. Germanou, L. G. Apostolova, L. Keller, M. Koutroumani, S. E.
Arnold, F. Panza, O. Gkatzima, S. Asthana, D. Hannequin, P. Whitehead, C. S. Atwood, P.
Caffarra, H. Hampel, I. Quintela, Á. Carracedo, L. Lannfelt, D. C. Rubinsztein, L. L. Barnes,
F. Pasquier, L. Frölich, S. Barral, B. McGuinness, T. G. Beach, J. A. Johnston, J. T.
Becker, P. Passmore, E. H. Bigio, J. M. Schott, T. D. Bird, J. D. Warren, B. F. Boeve, M. K.
Lupton, J. D. Bowen, P. Proitsi, A. Boxer, J. F. Powell, J. R. Burke, J. S. K. Kauwe, J. M.
Burns, M. Mancuso, J. D. Buxbaum, U. Bonuccelli, N. J. Cairns, A. McQuillin, C. Cao, G.
Livingston, C. S. Carlson, N. J. Bass, C. M. Carlsson, J. Hardy, R. M. Carney, J. Bras, M.
M. Carrasquillo, R. Guerreiro, M. Allen, H. C. Chui, E. Fisher, C. Masullo, E. A. Crocco, C.
DeCarli, G. Bisceglio, M. Dick, L. Ma, R. Duara, N. R. Graff-Radford, D. A. Evans, A.
Hodges, K. M. Faber, M. Scherer, K. B. Fallon, M. Riemenschneider, D. W. Fardo, R.
Heun, M. R. Farlow, H. Kölsch, S. Ferris, M. Leber, T. M. Foroud, I. Heuser, D. R. Galasko,
I. Giegling, M. Gearing, M. Hüll, D. H. Geschwind, J. R. Gilbert, J. Morris, R. C. Green, K.
Mayo, J. H. Growdon, T. Feulner, R. L. Hamilton, L. E. Harrell, D. Drichel, L. S. Honig, T.
D. Cushion, M. J. Huentelman, P. Hollingworth, C. M. Hulette, B. T. Hyman, R. Marshall,
G. P. Jarvik, A. Meggy, E. Abner, G. E. Menzies, L.-W. Jin, G. Leonenko, L. M. Real, G. R.
Jun, C. T. Baldwin, D. Grozeva, A. Karydas, G. Russo, J. A. Kaye, R. Kim, F. Jessen, N.
W. Kowall, B. Vellas, J. H. Kramer, E. Vardy, F. M. LaFerla, K.-H. Jöckel, J. J. Lah, M.
Dichgans, J. B. Leverenz, D. Mann, A. I. Levey, S. Pickering-Brown, A. P. Lieberman, N.
Klopp, K. L. Lunetta, H.-E. Wichmann, C. G. Lyketsos, K. Morgan, D. C. Marson, K. Brown,
F. Martiniuk, C. Medway, D. C. Mash, M. M. Nöthen, E. Masliah, N. M. Hooper, W. C.
McCormick, A. Daniele, S. M. McCurry, A. Bayer, A. N. McDavid, J. Gallacher, A. C.
McKee, H. van den Bussche, M. Mesulam, C. Brayne, B. L. Miller, S. Riedel-Heller, C. A.
Miller, J. W. Miller, A. Al-Chalabi, J. C. Morris, C. E. Shaw, A. J. Myers, J. Wiltfang, S.
O'Bryant, J. M. Olichney, V. Alvarez, J. E. Parisi, A. B. Singleton, H. L. Paulson, J.
Collinge, W. R. Perry, S. Mead, E. Peskind, D. H. Cribbs, M. Rossor, A. Pierce, N. S.
Ryan, W. W. Poon, B. Nacmias, H. Potter, S. Sorbi, J. F. Quinn, E. Sacchinelli, A. Raj, G.
Spalletta, M. Raskind, C. Caltagirone, P. Bossù, M. D. Orfei, B. Reisberg, R. Clarke, C.
Reitz, A. D. Smith, J. M. Ringman, D. Warden, E. D. Roberson, G. Wilcock, E. Rogaeva, A.
C. Bruni, H. J. Rosen, M. Gallo, R. N. Rosenberg, Y. Ben-Shlomo, M. A. Sager, P.
Mecocci, A. J. Saykin, P. Pastor, M. L. Cuccaro, J. M. Vance, J. A. Schneider, L. S.
Schneider, S. Slifer, W. W. Seeley, A. G. Smith, J. A. Sonnen, S. Spina, R. A. Stern, R. H.
Swerdlow, M. Tang, R. E. Tanzi, J. Q. Trojanowski, J. C. Troncoso, V. M. Van Deerlin, L. J.
Van Eldik, H. V. Vinters, J. P. Vonsattel, S. Weintraub, K. A. Welsh-Bohmer, K. C.
Wilhelmsen, J. Williamson, T. S. Wingo, R. L. Woltjer, C. B. Wright, C.-E. Yu, L. Yu, Y.
Saba, A. Pilotto, M. J. Bullido, O. Peters, P. K. Crane, D. Bennett, P. Bosco, E. Coto, V.
Boccardi, P. L. De Jager, A. Lleo, N. Warner, O. L. Lopez, M. Ingelsson, P. Deloukas, C.
Cruchaga, C. Graff, R. Gwilliam, M. Fornage, A. M. Goate, P. Sanchez-Juan, P. G. Kehoe,
N. Amin, N. Ertekin-Taner, C. Berr, S. Debette, S. Love, L. J. Launer, S. G. Younkin, J.-F.

Dartigues, C. Corcoran, M. A. Ikram, D. W. Dickson, G. Nicolas, D. Campion, J. Tschanz, H. Schmidt, H. Hakonarson, J. Clarimon, R. Munger, R. Schmidt, L. A. Farrer, C. Van Broeckhoven, M. C. O'Donovan, A. L. DeStefano, L. Jones, J. L. Haines, J.-F. Deleuze, M. J. Owen, V. Gudnason, R. Mayeux, V. Escott-Price, B. M. Psaty, A. Ramirez, L.-S. Wang, A. Ruiz, C. M. van Duijn, P. A. Holmans, S. Seshadri, J. Williams, P. Amouyel, G. D. Schellenberg, J.-C. Lambert, M. A. Pericak-Vance, Genetic meta-analysis of diagnosed Alzheimer's disease identifies new risk loci and implicates Aβ, tau, immunity and lipid processing. *Nat Genet.* **51**, 414–430 (2019).

14. J. Schwartzentruber, S. Cooper, J. Z. Liu, I. Barrio-Hernandez, E. Bello, N. Kumasaka, A. M. H. Young, R. J. M. Franklin, T. Johnson, K. Estrada, D. J. Gaffney, P. Beltrao, A. Bassett, Genome-wide meta-analysis, fine-mapping and integrative prioritization implicate new Alzheimer's disease risk genes. *Nat Genet.* **53**, 392–402 (2021).

15. C. Bellenguez, F. Küçükali, I. E. Jansen, L. Kleineidam, S. Moreno-Grau, N. Amin, A. C. Naj, R. Campos-Martin, B. Grenier-Boley, V. Andrade, P. A. Holmans, A. Boland, V. Damotte, S. J. van der Lee, M. R. Costa, T. Kuulasmaa, Q. Yang, I. de Rojas, J. C. Bis, A. Yaqub, I. Prokic, J. Chapuis, S. Ahmad, V. Giedraitis, D. Aarsland, P. Garcia-Gonzalez, C. Abdelnour, E. Alarcón-Martín, D. Alcolea, M. Alegret, I. Alvarez, V. Álvarez, N. J. Armstrong, A. Tsolaki, C. Antúnez, I. Appollonio, M. Arcaro, S. Archetti, A. A. Pastor, B. Arosio, L. Athanasiu, H. Bailly, N. Banaj, M. Baquero, S. Barral, A. Beiser, A. B. Pastor, J. E. Below, P. Benchek, L. Benussi, C. Berr, C. Besse, V. Bessi, G. Binetti, A. Bizarro, R. Blesa, M. Boada, E. Boerwinkle, B. Borroni, S. Boschi, P. Bossù, G. Bråthen, J. Bressler, C. Bresner, H. Brodaty, K. J. Brookes, L. I. Brusco, D. Buiza-Rueda, K. Bürger, V. Burholt, W. S. Bush, M. Calero, L. B. Cantwell, G. Chene, J. Chung, M. L. Cuccaro, Á. Carracedo, R. Cecchetti, L. Cervera-Carles, C. Charbonnier, H.-H. Chen, C. Chillotti, S. Ciccone, J. A. H. R. Claassen, C. Clark, E. Conti, A. Corma-Gómez, E. Costantini, C. Custodero, D. Daian, M. C. Dalmasso, A. Daniele, E. Dardiotis, J.-F. Dartigues, P. P. de Deyn, K. de Paiva Lopes, L. D. de Witte, S. Debette, J. Deckert, T. Del Ser, N. Denning, A. DeStefano, M. Dichgans, J. Diehl-Schmid, M. Diez-Fairen, P. D. Rossi, S. Djurovic, E. Duron, E. Düzel, C. Dufouil, G. Eiriksdottir, S. Engelborghs, V. Escott-Price, A. Espinosa, M. Ewers, K. M. Faber, T. Fabrizio, S. F. Nielsen, D. W. Fardo, L. Farotti, C. Fenoglio, M. Fernández-Fuertes, R. Ferrari, C. B. Ferreira, E. Ferri, B. Fin, P. Fischer, T. Fladby, K. Fließbach, B. Fongang, M. Fornage, J. Fortea, T. M. Foroud, S. Fostinelli, N. C. Fox, E. Franco-Macías, M. J. Bullido, A. Frank-García, L. Froelich, B. Fulton-Howard, D. Galimberti, J. M. García-Alberca, P. García-González, S. Garcia-Madrona, G. Garcia-Ribas, R. Ghidoni, I. Giegling, G. Giorgio, A. M. Goate, O. Goldhardt, D. Gomez-Fonseca, A. González-Pérez, C. Graff, G. Grande, E. Green, T. Grimmer, E. Grünblatt, M. Grunin, V. Gudnason, T. Guetta-Baranes, A. Haapasalo, G. Hadjigeorgiou, J. L. Haines, K. L. Hamilton-Nelson, H. Hampel, O. Hanon, J. Hardy, A. M. Hartmann, L. Hausner, J. Harwood, S. Heilmann-Heimbach, S. Helisalmi, M. T. Heneka, I. Hernández, M. J. Herrmann, P. Hoffmann, C. Holmes, H. Holstege, R. H. Vilas, M. Hulsman, J. Humphrey, G. J. Biessels, X. Jian, C. Johansson, G. R. Jun, Y. Kastumata, J. Kauwe, P. G. Kehoe, L. Kilander, A. K. Ståhlbom, M. Kivipelto, A. Koivisto, J. Kornhuber, M. H. Kosmidis, W. A. Kukull, P. P. Kuksa, B. W. Kunkle, A. B. Kuzma, C. Lage, E. J. Laukka, L. Launer, A. Lauria, C.-Y. Lee, J. Lehtisalo, O. Lerch, A. Lleó, W. Longstreth, O. Lopez, A. L. de Munain, S. Love, M. Löwemark, L. Luckcuck, K. L. Lunetta, Y. Ma, J. Macías, C. A. MacLeod, W. Maier, F. Mangialasche, M. Spallazzi, M. Marquié, R. Marshall, E. R. Martin, A. M. Montes, C. M. Rodríguez, C. Masullo, R. Mayeux, S. Mead, P. Mecocci, M. Medina, A. Meggy, S. Mehrabian, S. Mendoza, M. Menéndez-González, P. Mir, S. Moebus, M. Mol, L. Molina-Porcel, L. Montrreal, L. Morelli, F. Moreno, K. Morgan, T. Mosley, M. M. Nöthen, C. Muchnik, S. Mukherjee, B. Nacmias, T. Ngandu,

G. Nicolas, B. G. Nordestgaard, R. Olaso, A. Orellana, M. Orsini, G. Ortega, A. Padovani, C. Paolo, G. Papenberg, L. Parnetti, F. Pasquier, P. Pastor, G. Peloso, A. Pérez-Cordón, J. Pérez-Tur, P. Pericard, O. Peters, Y. A. L. Pijnenburg, J. A. Pineda, G. Piñol-Ripoll, C. Pisanu, T. Polak, J. Popp, D. Posthuma, J. Priller, R. Puerta, O. Quenez, I. Quintela, J. Q. Thomassen, A. Rábano, I. Rainero, F. Rajabli, I. Ramakers, L. M. Real, M. J. T. Reinders, C. Reitz, D. Reyes-Dumeyer, P. Ridge, S. Riedel-Heller, P. Riederer, N. Roberto, E. Rodriguez-Rodriguez, A. Rongve, I. R. Allende, M. Rosende-Roca, J. L. Royo, E. Rubino, D. Rujescu, M. E. Sáez, P. Sakka, I. Saltvedt, Á. Sanabria, M. B. Sánchez-Arjona, F. Sanchez-Garcia, P. S. Juan, R. Sánchez-Valle, S. B. Sando, C. Sarnowski, C. L. Satizabal, M. Scamosci, N. Scarmeas, E. Scarpini, P. Scheltens, N. Scherbaum, M. Scherer, M. Schmid, A. Schneider, J. M. Schott, G. Selbæk, D. Seripa, M. Serrano, J. Sha, A. A. Shadrin, O. Skrobot, S. Slifer, G. J. L. Snijders, H. Soininen, V. Solfrizzi, A. Solomon, Y. Song, S. Sorbi, O. Sotolongo-Grau, G. Spalletta, A. Spottke, A. Squassina, E. Stordal, J. P. Tartan, L. Tárraga, N. Tesí, A. Thalamuthu, T. Thomas, G. Tosto, L. Traykov, L. Tremolizzo, A. Tybjærg-Hansen, A. Uitterlinden, A. Ullgren, I. Ulstein, S. Valero, O. Valladares, C. V. Broeckhoven, J. Vance, B. N. Vardarajan, A. van der Lugt, J. V. Dongen, J. van Rooij, J. van Swieten, R. Vandenberghe, F. Verhey, J.-S. Vidal, J. Vogelgsang, M. Vyhnalek, M. Wagner, D. Wallon, L.-S. Wang, R. Wang, L. Weinhold, J. Wiltfang, G. Windle, B. Woods, M. Yannakoulia, H. Zare, Y. Zhao, X. Zhang, C. Zhu, M. Zulaica, EADB, GR@ACE, DEGESCO, EADI, GERAD, Demgene, FinnGen, ADGC, CHARGE, L. A. Farrer, B. M. Psaty, M. Ghanbari, T. Raj, P. Sachdev, K. Mather, F. Jessen, M. A. Ikram, A. de Mendonça, J. Hort, M. Tsolaki, M. A. Pericak-Vance, P. Amouyel, J. Williams, R. Frikke-Schmidt, J. Clarimon, J.-F. Deleuze, G. Rossi, S. Seshadri, O. A. Andreassen, M. Ingelsson, M. Hiltunen, K. Sleegers, G. D. Schellenberg, C. M. van Duijn, R. Sims, W. M. van der Flier, A. Ruiz, A. Ramirez, J.-C. Lambert, New insights into the genetic etiology of Alzheimer's disease and related dementias. *Nat Genet* (2022), doi:10.1038/s41588-022-01024-z.

16. L. Brase, S.-F. You, R. D. Albanus, J. L. Del-Aguila, Y. Dai, B. C. Novotny, C. Soriano-Tarraga, T. Dykstra, M. V. Fernandez, J. P. Budde, K. Bergmann, J. C. Morris, R. J. Bateman, R. J. Perrin, E. McDade, C. Xiong, A. Goate, M. Farlow, J. P. Chhatwal, P. Schofield, H. Chui, Dominantly Inherited Alzheimer Network (DIAN), G. T. Sutherland, J. Kipnis, C. M. Karch, B. A. Benitez, C. Cruchaga, O. Harari, "A landscape of the genetic and cellular heterogeneity in Alzheimer disease" (preprint, Neurology, 2021), , doi:10.1101/2021.11.30.21267072.

17. M. Olah, V. Menon, N. Habib, M. F. Taga, Y. Ma, C. J. Yung, M. Cimpean, A. Khairallah, G. Coronas-Samano, R. Sankowski, D. Grün, A. A. Kroshilina, D. Dionne, R. A. Sarkis, G. R. Cosgrove, J. Helgager, J. A. Golden, P. B. Pennell, M. Prinz, J. P. G. Vonsattel, A. F. Teich, J. A. Schneider, D. A. Bennett, A. Regev, W. Elyaman, E. M. Bradshaw, P. L. De Jager, Single cell RNA sequencing of human microglia uncovers a subset associated with Alzheimer's disease. *Nat Commun*. **11**, 6129 (2020).

18. S.-F. Lau, H. Cao, A. K. Y. Fu, N. Y. Ip, Single-nucleus transcriptome analysis reveals dysregulation of angiogenic endothelial cells and neuroprotective glia in Alzheimer's disease. *Proc Natl Acad Sci U S A*. **117**, 25800–25809 (2020).

19. S. Morabito, E. Miyoshi, N. Michael, S. Shahin, A. C. Martini, E. Head, J. Silva, K. Leavy, M. Perez-Rosendahl, V. Swarup, Single-nucleus chromatin accessibility and transcriptomic characterization of Alzheimer's disease. *Nat Genet* (2021), doi:10.1038/s41588-021-00894-z.

20.   N. M. Dräger, S. M. Sattler, C. T.-L. Huang, O. M. Teter, K. Leng, S. H. Hashemi, J. Hong,
C. D. Clelland, L. Zhan, L. Kodama, A. B. Singleton, M. A. Nalls, J. Ichida, M. E. Ward, F.
Faghri, L. Gan, M. Kampmann, "A CRISPRi/a platform in iPSC-derived microglia uncovers
regulators of disease states" (preprint, Neuroscience, 2021), ,
doi:10.1101/2021.06.16.448639.

21.   Q. Li, B. A. Barres, Microglia and macrophages in brain homeostasis and disease. *Nat Rev*
Immunol. **18**, 225–242 (2018).

22.   A. Kumanogoh, H. Kikutani, Immunological functions of the neuropilins and plexins as
receptors for semaphorins. *Nat Rev Immunol*. **13**, 802–814 (2013).

23.   G. Jun, H. Asai, E. Zeldich, E. Drapeau, C. Chen, J. Chung, J.-H. Park, S. Kim, V.
Haroutunian, T. Foroud, R. Kuwano, J. L. Haines, M. A. Pericak-Vance, G. D.
Schellenberg, K. L. Lunetta, J.-W. Kim, J. D. Buxbaum, R. Mayeux, T. Ikezu, C. R.
Abraham, L. A. Farrer, PLXNA4 is associated with Alzheimer disease and modulates tau
phosphorylation. *Ann Neurol*. **76**, 379–392 (2014).

24.   P. F. Good, D. Alapat, A. Hsu, C. Chu, D. Perl, X. Wen, D. E. Burstein, D. S. Kohtz, A role
for semaphorin 3A signaling in the degeneration of hippocampal neurons during
Alzheimer's disease. *J Neurochem*. **91**, 716–736 (2004).

25.   M. Efremova, M. Vento-Tormo, S. A. Teichmann, R. Vento-Tormo, CellPhoneDB: inferring
cell–cell communication from combined expression of multi-subunit ligand–receptor
complexes. *Nat Protoc*. **15**, 1484–1506 (2020).

26.   Y. Zhou, W. M. Song, P. S. Andhey, A. Swain, T. Levy, K. R. Miller, P. L. Poliani, M.
Cominelli, S. Grover, S. Gilfillan, M. Cella, T. K. Ulland, K. Zaitsev, A. Miyashita, T. Ikeuchi,
M. Sainouchi, A. Kakita, D. A. Bennett, J. A. Schneider, M. R. Nichols, S. A. Beausoleil, J.
D. Ulrich, D. M. Holtzman, M. N. Artyomov, M. Colonna, Human and mouse single-nucleus
transcriptomics reveal TREM2-dependent and TREM2-independent cellular responses in
Alzheimer's disease. *Nat Med*. **26**, 131–142 (2020).

27.   Y. Hu, T. Peng, L. Gao, K. Tan, CytoTalk: De novo construction of signal transduction
networks using single-cell transcriptomic data. *Sci. Adv.* **7**, eabf1356 (2021).

28.   Z. Szepesi, O. Manouchehrian, S. Bachiller, T. Deierborg, Bidirectional Microglia-Neuron
Communication in Health and Disease. *Front Cell Neurosci*. **12**, 323 (2018).

29.   F. Nakamura, R. G. Kalb, S. M. Strittmatter, Molecular basis of semaphorin-mediated axon
guidance. *J Neurobiol*. **44**, 219–229 (2000).

30.   J. Klesney-Tait, I. R. Turnbull, M. Colonna, The TREM receptor family and signal
integration. *Nat Immunol*. **7**, 1266–1273 (2006).

31.   D. L. Kober, T. J. Brett, TREM2-Ligand Interactions in Health and Disease. *J Mol Biol*. **429**,
1607–1629 (2017).

32.   E. E. Evans, V. Mishra, C. Mallow, E. M. Gersz, L. Balch, A. Howell, C. Reilly, E. S. Smith,
T. L. Fisher, M. Zauderer, Semaphorin 4D is up-regulated in neurons of diseased brains
and triggers astrocyte reactivity. *J Neuroinflammation*. **19**, 200 (2022).

33. T. K. Ulland, M. Colonna, TREM2 — a key player in microglial biology and Alzheimer
disease. *Nat Rev Neurol*. **14**, 667–675 (2018).

34. Y. Atagi, C.-C. Liu, M. M. Painter, X.-F. Chen, C. Verbeeck, H. Zheng, X. Li, R.
Rademakers, S. S. Kang, H. Xu, S. Younkin, P. Das, J. D. Fryer, G. Bu, Apolipoprotein E
Is a Ligand for Triggering Receptor Expressed on Myeloid Cells 2 (TREM2). *J Biol Chem*.
290, 26043–26050 (2015).

35. F. L. Yeh, Y. Wang, I. Tom, L. C. Gonzalez, M. Sheng, TREM2 Binds to Apolipoproteins,
Including APOE and CLU/APOJ, and Thereby Facilitates Uptake of Amyloid-Beta by
Microglia. *Neuron*. **91**, 328–340 (2016).

36. S. M. Neuner, J. Tcw, A. M. Goate, Genetic architecture of Alzheimer's disease. *Neurobiol
Dis*. **143**, 104976 (2020).

37. E. Rogaeva, Y. Meng, J. H. Lee, Y. Gu, T. Kawarai, F. Zou, T. Katayama, C. T. Baldwin, R.
Cheng, H. Hasegawa, F. Chen, N. Shibata, K. L. Lunetta, R. Pardossi-Piquard, C. Bohm,
Y. Wakutani, L. A. Cupples, K. T. Cuenco, R. C. Green, L. Pinessi, I. Rainero, S. Sorbi, A.
Bruni, R. Duara, R. P. Friedland, R. Inzelberg, W. Hampe, H. Bujo, Y.-Q. Song, O. M.
Andersen, T. E. Willnow, N. Graff-Radford, R. C. Petersen, D. Dickson, S. D. Der, P. E.
Fraser, G. Schmitt-Ulms, S. Younkin, R. Mayeux, L. A. Farrer, P. St George-Hyslop, The
neuronal sortilin-related receptor SORL1 is genetically associated with Alzheimer disease.
Nat Genet. **39**, 168–177 (2007).

38. R.-H. Yin, J.-T. Yu, L. Tan, The Role of SORL1 in Alzheimer's Disease. *Mol Neurobiol*. **51**,
909–918 (2015).

39. Y. Deming, F. Filipello, F. Cignarella, C. Cantoni, S. Hsu, R. Mikesell, Z. Li, J. L. Del-
Aguila, U. Dube, F. G. Farias, J. Bradley, J. Budde, L. Ibanez, M. V. Fernandez, K.
Blennow, H. Zetterberg, A. Heslegrave, P. M. Johansson, J. Svensson, B. Nellgård, A.
Lleo, D. Alcolea, J. Clarimon, L. Rami, J. L. Molinuevo, M. Suárez-Calvet, E. Morenas-
Rodríguez, G. Kleinberger, M. Ewers, O. Harari, C. Haass, T. J. Brett, B. A. Benitez, C. M.
Karch, L. Piccio, C. Cruchaga, The MS4A gene cluster is a key modulator of soluble
TREM2 and Alzheimer's disease risk. *Sci Transl Med*. **11**, eaau2291 (2019).

40. W.-Y. Wang, M.-S. Tan, J.-T. Yu, L. Tan, Role of pro-inflammatory cytokines released from
microglia in Alzheimer's disease. *Ann Transl Med*. **3**, 136 (2015).

41. A. McQuade, M. Coburn, C. H. Tu, J. Hasselmann, H. Davtyan, M. Blurton-Jones,
Development and validation of a simplified method to generate human microglia from
pluripotent stem cells. *Mol Neurodegener*. **13**, 67 (2018).

42. T. R. Jay, V. E. von Saucken, G. E. Landreth, TREM2 in Neurodegenerative Diseases. *Mol
Neurodegener*. **12**, 56 (2017).

43. A. Grubman, X. Y. Choo, G. Chew, J. F. Ouyang, G. Sun, N. P. Croft, F. J. Rossello, R.
Simmons, S. Buckberry, D. V. Landin, J. Pflueger, T. H. Vandekolk, Z. Abay, Y. Zhou, X.
Liu, J. Chen, M. Larcombe, J. M. Haynes, C. McLean, S. Williams, S. Y. Chai, T. Wilson,
R. Lister, C. W. Pouton, A. W. Purcell, O. J. L. Rackham, E. Petretto, J. M. Polo,
Transcriptional signature in microglia associated with Aβ plaque phagocytosis. *Nat
Commun*. **12**, 3015 (2021).

44. A. McQuade, Y. J. Kang, J. Hasselmann, A. Jairaman, A. Sotelo, M. Coburn, S. K.
Shabestari, J. P. Chadarevian, G. Fote, C. H. Tu, E. Danhash, J. Silva, E. Martinez, C.
Cotman, G. A. Prieto, L. M. Thompson, J. S. Steffan, I. Smith, H. Davtyan, M. Cahalan, H.
Cho, M. Blurton-Jones, Gene expression and functional deficits underlie TREM2-knockout
microglia responses in human models of Alzheimer's disease. *Nat Commun.* **11**, 5370
(2020).

45. H. Keren-Shaul, A. Spinrad, A. Weiner, O. Matcovitch-Natan, R. Dvir-Szternfeld, T. K.
Ulland, E. David, K. Baruch, D. Lara-Astaiso, B. Toth, S. Itzkovitz, M. Colonna, M.
Schwartz, I. Amit, A Unique Microglia Type Associated with Restricting Development of
Alzheimer's Disease. *Cell.* **169**, 1276-1290.e17 (2017).

46. N. Takegahara, H. Takamatsu, T. Toyofuku, T. Tsujimura, T. Okuno, K. Yukawa, M. Mizui,
M. Yamamoto, D. V. R. Prasad, K. Suzuki, M. Ishii, K. Terai, M. Moriya, Y. Nakatsuji, S.
Sakoda, Plexin-A1 and its interaction with DAP12 in immune responses and bone
homeostasis. *Nature Cell Biology.* **8**, 615–22 (2006).

47. A. Kumanogoh, H. Kikutani, Immunological functions of the neuropilins and plexins as
receptors for semaphorins. *Nature Reviews. Immunology.* **13**, 802–14 (2013).

48. S. S. Kang, A. Kurti, A. Wojtas, K. E. Baker, C.-C. Liu, T. Kanekiyo, Y. Deming, C.
Cruchaga, S. Estus, G. Bu, J. D. Fryer, Identification of plexin A4 as a novel clusterin
receptor links two Alzheimer's disease risk genes. *Hum Mol Genet.* **25**, 3467–3475 (2016).

49. Q. Han, Y.-A. Sun, Y. Zong, C. Chen, H.-F. Wang, L. Tan, Alzheimer's Disease
Neuroimaging Initiative, Common Variants in PLXNA4 and Correlation to CSF-related
Phenotypes in Alzheimer's Disease. *Frontiers in Neuroscience.* **12** (2018) (available at
https://www.frontiersin.org/articles/10.3389/fnins.2018.00946).

50. S. Chung, J. Yang, H. J. Kim, E. M. Hwang, W. Lee, K. Suh, H. Choi, I. Mook-Jung, Plexin-
A4 mediates amyloid-β-induced tau pathology in Alzheimer's disease animal model. *Prog
Neurobiol.* **203**, 102075 (2021).

51. A. Macedo, C. Gómez, M. Â. Rebelo, J. Poza, I. Gomes, S. Martins, A. Maturana-
Candelas, V. G. Pablo, L. Durães, P. Sousa, M. Figueruelo, M. Rodríguez, C. Pita, M.
Arenas, L. Álvarez, R. Hornero, A. M. Lopes, N. Pinto, Risk Variants in Three Alzheimer's
Disease Genes Show Association with EEG Endophenotypes. *J Alzheimers Dis.* **80**, 209–
223 (2021).

52. Y. Deming, Z. Li, M. Kapoor, O. Harari, J. L. Del-Aguila, K. Black, D. Carrell, Y. Cai, M. V.
Fernandez, J. Budde, S. Ma, B. Saef, B. Howells, K.-L. Huang, S. Bertelsen, A. M. Fagan,
D. M. Holtzman, J. C. Morris, S. Kim, A. J. Saykin, P. L. De Jager, M. Albert, A. Moghekar,
R. O'Brien, M. Riemenschneider, R. C. Petersen, K. Blennow, H. Zetterberg, L. Minthon, V.
M. Van Deerlin, V. M.-Y. Lee, L. M. Shaw, J. Q. Trojanowski, G. Schellenberg, J. L.
Haines, R. Mayeux, M. A. Pericak-Vance, L. A. Farrer, E. R. Peskind, G. Li, A. F. Di Narzo,
Alzheimer's Disease Neuroimaging Initiative (ADNI), Alzheimer Disease Genetic
Consortium (ADGC), J. S. K. Kauwe, A. M. Goate, C. Cruchaga, Genome-wide association
study identifies four novel loci associated with Alzheimer's endophenotypes and disease
modifiers. *Acta Neuropathol.* **133**, 839–856 (2017).

53.   A. Zettergren, K. Höglund, S. Kern, V. Thorvaldsson, M. Johan Skoog, O. Hansson, N.
Andreasen, N. Bogdanovic, K. Blennow, I. Skoog, H. Zetterberg, Association of IL1RAP-
related genetic variation with cerebrospinal fluid concentration of Alzheimer-associated tau
protein. *Sci Rep.* **9**, 2460 (2019).

54.   F. Su, F. Bai, Z. Zhang, Inflammatory Cytokines and Alzheimer's Disease: A Review from
the Perspective of Genetic Polymorphisms. *Neurosci Bull.* **32**, 469–480 (2016).

55.   G. Ogunmokun, S. Dewanjee, P. Chakraborty, C. Valupadas, A. Chaudhary, V. Kolli, U.
Anand, J. Vallamkondu, P. Goel, H. P. R. Paluru, K. D. Gill, P. H. Reddy, V. De Feo, R.
Kandimalla, The Potential Role of Cytokines and Growth Factors in the Pathogenesis of
Alzheimer's Disease. *Cells.* **10**, 2790 (2021).

56.   A. S. Mendiola, A. E. Cardona, The IL-1β phenomena in neuroinflammatory diseases. *J
Neural Transm (Vienna).* **125**, 781–795 (2018).

57.   T. Stuart, A. Butler, P. Hoffman, C. Hafemeister, E. Papalexi, W. M. Mauck, Y. Hao, M.
Stoeckius, P. Smibert, R. Satija, Comprehensive Integration of Single-Cell Data. *Cell.* **177**,
1888-1902.e21 (2019).

58.   C. Hafemeister, R. Satija, Normalization and variance stabilization of single-cell RNA-seq
data using regularized negative binomial regression. *Genome Biol.* **20**, 296 (2019).

59.   Y. Zhang, T. Liu, C. A. Meyer, J. Eeckhoute, D. S. Johnson, B. E. Bernstein, C. Nussbaum,
R. M. Myers, M. Brown, W. Li, X. S. Liu, Model-based Analysis of ChIP-Seq (MACS).
Genome Biol. **9**, R137 (2008).

60.   H. M. Amemiya, A. Kundaje, A. P. Boyle, The ENCODE Blacklist: Identification of
Problematic Regions of the Genome. *Sci Rep.* **9**, 9354 (2019).

61.   H. A. Pliner, J. S. Packer, J. L. McFaline-Figueroa, D. A. Cusanovich, R. M. Daza, D.
Aghamirzaie, S. Srivatsan, X. Qiu, D. Jackson, A. Minkina, A. C. Adey, F. J. Steemers, J.
Shendure, C. Trapnell, Cicero Predicts cis-Regulatory DNA Interactions from Single-Cell
Chromatin Accessibility Data. *Mol. Cell.* **71**, 858-871.e8 (2018).

62.   A. Patel, A. Garcia Diaz, J. C. Moore, D. Sirabella, B. Corneo, Establishment and
characterization of two iPSC lines derived from healthy controls. *Stem Cell Res.* **47**,
101926 (2020).

63.   W. B. Stine, K. N. Dahlgren, G. A. Krafft, M. J. LaDu, In vitro characterization of conditions
for amyloid-beta peptide oligomerization and fibrillogenesis. *J Biol Chem.* **278**, 11612–
11622 (2003).

64.   D. Kim, J. M. Paggi, C. Park, C. Bennett, S. L. Salzberg, Graph-based genome alignment
and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol.* **37**, 907–915 (2019).

65.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

66.   Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate: A Practical and Powerful
854   Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B*
855   *(Methodological)*. **57**, 289–300 (1995).

856   67.   D. Szklarczyk, A. L. Gable, K. C. Nastou, D. Lyon, R. Kirsch, S. Pyysalo, N. T. Doncheva,
857   M. Legeay, T. Fang, P. Bork, L. J. Jensen, C. von Mering, The STRING database in 2021:
858   customizable protein-protein networks, and functional characterization of user-uploaded
859   gene/measurement sets. *Nucleic Acids Res.* **49**, D605–D612 (2021).

860   68.   P. Shannon, A. Markiel, O. Ozier, N. S. Baliga, J. T. Wang, D. Ramage, N. Amin, B.
861   Schwikowski, T. Ideker, Cytoscape: a software environment for integrated models of
862   biomolecular interaction networks. *Genome Res.* **13**, 2498–2504 (2003).

863   69.   J. H. Morris, L. Apeltsin, A. M. Newman, J. Baumbach, T. Wittkop, G. Su, G. D. Bader, T.
864   E. Ferrin, clusterMaker: a multi-algorithm clustering plugin for Cytoscape. *BMC*
865   *Bioinformatics.* **12**, 436 (2011).

866   70.   C. E. Shannon, A Mathematical Theory of Communication. *Bell System Technical Journal.*
867   **27**, 379–423 (1948).

868   71.   A. A. Margolin, I. Nemenman, K. Basso, C. Wiggins, G. Stolovitzky, R. Dalla Favera, A.
869   Califano, ARACNE: an algorithm for the reconstruction of gene regulatory networks in a
870   mammalian cellular context. *BMC Bioinformatics.* **7 Suppl 1**, S7 (2006).

871   72.   P. Langfelder, S. Horvath, WGCNA: an R package for weighted correlation network
872   analysis. *BMC Bioinformatics.* **9**, 559 (2008).

Methods

snRNA-seq data processing

Our single-nucleus RNA-seq dataset from 67 donors was processed as described previously (*16*). Briefly, we used the 10X Chromium single-cell Reagent Kit v3 aiming for 10,000 nuclei per individual and 50,000 reads per nucleus. We mapped the sequenced reads to the GRCh38 human reference genome using CellRanger (v. 6.1.1) and filtered nuclei based on sequencing depth and percent mitochondrial reads. In total, we obtained snRNA-seq data from 294,114 nuclei after stringent QC. We used Seurat (*57*) (v. 3.2.3) to identify cell types and transcriptional states.

We obtained the count matrices and barcode metadata from other public snRNA-seq datasets from human brains (*18*, *19*, *26*) as made available by the authors. We used the barcode metadata to annotate the nuclei into cell types and donor categories. We considered any nuclei not in the final metadata table for the respective study as failing QC. These nuclei were discarded from downstream analyses. In order to run CellPhoneDB and CytoTalk in these datasets, we normalized each study separately using the SCT normalization (*58*) from Seurat.

Reanalysis of brain snATAC-seq data

We downloaded the raw snATAC-seq fastq files from the Morabito *et al.* study (*19*) and re-processed this dataset using CellRanger-ATAC (v. 2.0.0) count function. Firstly, we filtered the resulting GRCh38 BAM files for each sample using flags *-f 3 -F 4 -F 8 -F 256 -F 1024 -F 2048 -q 30* to retain high-quality, properly mapped alignments. Second, we split each sample BAM file into separate files corresponding to each cell type using the barcode annotations reported by the study authors. For each cell type, we merged all BAM files across donors and called narrow peaks using MACS2 (v. 2.2.7.1) (*59*) using flags *--nomodel --shift -100 --extsize 200 --keep-dup all --call-summits -B*. Third, we filtered all peaks that overlapped the GRCh38 ENCODE exclusion list regions (ENCODE accession ENCFF356LFX) (*60*). Finally, we used CICERO (*61*), with default parameters, to calculate co-accessibility across ATAC-seq narrow peaks for each cell type separately.

Cell Culture

iPSCs were maintained feeder-free on matrigel-coated plates in StemFlex medium (Gibco) in a humidified incubator (5% CO_2, 37 □°C).

Differentiation of iPSC to iPSC-derived microglia (iMGL)

iPSC-microglia were generated as described in McQuade et al., 2018 (*41*). Briefly, iPSCs were directed down a hematopoietic lineage using the STEMdiff Hematopoietic kit (STEMCELL Technologies). After 12 days in culture, hematopoietic progenitor cells (HPCs) were transferred into a microglia differentiation medium containing DMEM/F12, 2× insulin-transferrin-selenite, 2× B27, 0.5× N2, 1× GlutaMAX, 1× nonessential amino acids, 400 □µM monothioglycerol, and 5 □µg/mL human insulin. Media was added to cultures every other day and supplemented with 100 □ng/mL IL-34, 50 □ng/mL TGF-β1, and 25 ng/mL M-CSF (PeproTech) for 28 days. In the final three days of differentiation, 100 ng/mL CD200 (Novoprotein) and 100 □ng/mL CX3CL1 (PeproTech) were added to the culture.

Generation of TREM2 KO iPSCs

The WT iPSC line (APOE 3/3) was a generous gift from Dr. Barbara Corneo (*62*). TREM2 KO iiPSCs were generated using the Human Stem Cell Nucleofector™ Kit 2 (Lonza) according to the manufacturer's protocol. Briefly, iiPSCs were resuspended in 100 µL nucleofection buffer and 5 µg of vector lentiCRISPRv2GFP expressing sgRNA against TREM2 (5'ATCACAGACGATACCCTGGG 3'). LentiCRISPRv2GFP was a generous gift from David Feldser (Addgene plasmid# 82416; http://n2t.net/addgene:82416; RRID:Addgene_82416). The suspension was transferred to the Amaxa Nucleofector cuvette and transfected using program CA-137. GFP-expressing cells were single-cell sorted into 96-well plates using the BD Influx cell sorter (Columbia University CCTI/HICCC Flow Core). Cell colonies were maintained and expanded. TREM2 KO cell lines were verified by Sequencing (Genewiz).

Western Blotting Analysis and Antibodies

Proteins were resolved via SDS-PAGE using NuPAGE™ 4-12% Bis-Tris gels (Invitrogen) and probed for expression. Proteins were detected using the Odyssey XF Imager (LI-COR Biosciences). Antibodies used included: anti-Phospho-SYK (Cell Signaling), anti-SYK (Cell Signaling), anti-TREM2 (Cell Signaling), and anti-β-actin (Sigma).).

Immunocytochemistry

iMGL were fixed in 4% paraformaldehyde, permeabilized, and blocked for 1 hr in 10% normal goat serum or 10% fetal bovine serum. Primary antibodies were diluted in 10% normal goat serum containing 0.1% Triton X-100 and incubated overnight at 4°C. Primary antibodies included anti-TREM2 (R&D Systems), anti-IBA1 (Fujifilm), and anti-TMEM119 (Novus)). Following incubation of the primary antibody, cells were washed and incubated for 1 hr at room temperature in Alexa Fluor 568- or 488-conjugated secondary antibody (Invitrogen). Immunostained cells were mounted using VECTASHIELD Mounting Medium (Vector Laboratories) and imaged using the Nikon C1 Digital Confocal System.

Synaptosome Preparation

The human temporal lobe was homogenized using a Dounce tissue grinder in homogenization buffer (0.032 M Sucrose, 0.5 mM $CaCl_2$, 1 mM $MgCl_2$, 1 mM $NaHCO_3$ supplemented with protease/phosphatase inhibitor) and centrifuged at 1400×g. The supernatant was saved (S1), and the resulting pellet was re-homogenized in homogenization buffer and centrifuged at 700×g, resulting in supernatant, S1'. Supernatants S1 and S1' were mixed and centrifuged at 700×g resulting in supernatant, S2. S2 was centrifuged at 14000×g. The resulting pellet containing synaptosomes was resuspended in buffer (0.32 M Sucrose, 1 mM $NaHCO_3$).

Oligomer and fibril formation

Aβ42 oligomers and fibrils were prepared according to the established protocol by Stine et al. (*63*). Briefly, Aβ42 (Anaspec) was resuspended to 1 mM solution in HFIP. Aliquots were evaporated in a fume hood for 2 hrs, and the resulting peptide film was dried under vacuum in a SpeedVac and stored at -20°C. Immediately prior to use, the films were solubilized to 1 mM in anhydrous DMSO, sonicated in a bath sonicator for 10 min, and diluted to 100 µM in oligomer-

954 forming conditions (cold PBS; incubation at 4°C for 24 h) or fibril-forming conditions (10mM HCl;
955 incubation at 37°C for 24 h).

pHrodo Labeling

pHrodo Red (Invitrogen) labeling protocol was adapted from the manufacturer's instructions.
Briefly, Aβ42 oligomer or fibril solutions (100 µM) were centrifuged at 16000×g for 2 min to
collect the aggregates. Aggregates were washed in 1 mL HBSS and centrifuged for 2 min. The
supernatants were aspirated, and 200 µl of pellet aggregates were resuspended in by adding
200 µl 0.1M NaHCO$_3$ to the pellets. The pHrodo Red dye (10 mg/ml) was then added as per the
manufacturer's instructions. pHrodo labeling was performed and incubated for 1 hr in the dark,
followed by centrifugation at 16000×g for 2 min. Aggregates were washed 3x with HBSS and
resuspended to 100 µM in HBSS.

Phagocytosis assay

iMGL were plated at a density of 3x10^5 cells per well in 96-well plates and incubated for 24 hrs.
iMGL were treated with 10 µM SEMA6D in media supplemented with either 30 µg/ml pHrodo
Red tagged synaptosomes, 1 mM pHrodo Red tagged Aβ fibril or 400 nM Aβ oligomer and
incubated for 24 hrs. Cells were imaged using an IncuCyte live cell imaging system. Analysis
was performed by Incucyte Zoom Software. The phagocytic activity of iMGL was quantified
using the Incucyte® SX5 Live-Cell Analysis instrument. Total cell counts were performed using
the CellQuanti-Blue™ Cell Viability Assay kit (BioAssay Systems).

Cytokine assay

974 iMGL were plated at a density of 3x10^5 cells per well in 96-well plates and incubated for 24 hrs.
975 iMGL were treated with 10 µM SEMA6D for 24 hrs. Cell media was harvested, and cytokines
976 were quantified using the V-PLEX Human Proinflammatory Panel II (4-Plex) (MSD) according to
977 the manufacturer's instructions. Total cell counts were performed using the CellQuanti-Blue™
978 Cell Viability Assay kit.

iMGL RNAseq

WT or TREM2 KO iMGL were treated with 10 µM SEMA6D for 24 hrs. mRNA was extracted
from iMGL using RNeasy Mini kit (Qiagen) following the manufacturer's instructions. Human
mRNA sequencing and bioinformatics analysis were performed by Novogene. Raw fastq files
were mapped to the GRCh38 reference genome using HISAT2 (v. 2.0.5) (*64*) and the
corresponding GRCh38 index files (http://daehwankimlab.github.io/hisat2/download). We used
DESeq2 (*65*) for differential analyses. Briefly, we retained genes with ≥ 10 counts in at least
three samples and processed all samples together to increase the statistical power to calculate
dispersion. We then retrieved the fold changes and significances for the comparisons of interest
individually (*e.g.*, SEMA6D-treated WT vs. untreated WT). Finally, we adjusted for multiple
testing per comparison using the Benjamini-Hochberg correction (*66*) and used a 10% false
discovery rate (FDR) threshold to identify differentially expressed genes.

CellPhoneDB analyses

We used CellPhoneDB (*25*) (v. 2.1.7) to estimate crosstalk interactions based on the expression
of known ligand-receptor pairs. We used the *cellphonedb method statistical_analysis* function

with default parameters to calculate crosstalk interactions at the cluster level (transcriptional
state) using the sctransform-normalized snRNA-seq count matrices as input. We performed
CellPhoneDB analyses for each subset of donors separately (*e.g.*, controls, sporadic AD). We
considered significant all those interactions that passed multiple testing correction (Bonferroni *p*
< 0.05) for the entire set of analyses (all transcriptional states and donor categories).

To address issues of potential bias from differences in cell type representation, we performed
an additional CellPhoneDB analysis, downsampling each cluster in our snRNA-seq data to 100
nuclei barcodes. Similarly, we re-ran CellPhoneDB in multiple combinations of two to three cell
types in our data (e.g., astrocytes and OPC only, astrocytes, microglia, and OPC only) to
confirm that the resulting interactions were stable across subsets of cell types. Finally, we
repeated the CellPhoneDB analyses individually for a subset of brains to confirm that the
crosstalk patterns were identified at a single brain level. For other public snRNA-seq datasets
analyzed in this study, we used the cluster and donor labels provided by the authors of each
study.

To calculate the significance of CellPhoneDB interactions, we used Fisher exact tests (FETs) to
compare the number of interactions at the cell type level. We used the union of all significant
interactions across transcriptional states for each cell type. We generated one 2x2 contingency
table per cell type and donor category combination for the comparisons between controls versus
other donor categories. One dimension of the contingency table encoded the number of unique
significant CellPhoneDB interactions involving the cell type of interest versus all other cell types
combined. The other dimension encoded the number of interactions identified in the controls
versus the donor category of interest. To calculate the enrichment of crosstalk interactions
involving AD-related genes per cell type, we generated, for each donor category, 2x2
contingency tables encoding the number of interactions involving the cell type of interest (yes
versus no) and involving AD genes (yes versus no). Finally, we calculated the enrichment of
AD-related crosstalk interactions pairwise across all cell-type pairs by further subsetting the
previous contingency tables to separate between AD-related interactions involving cell type of
interest A and cell type of interest B versus cell type of interest A without cell type of interest B
(*e.g.*, microglia and excitatory neurons versus to microglia and the remaining cell types). All FET
p-values were corrected for multiple testing using the Bonferroni correction based on the total
number of tests performed for each analysis.

Crosstalk enrichments for other neuropsychiatric traits

We downloaded the GWAS associations of neuropsychiatric traits from the European
Bioinformatics Institute GWAS catalog (https://www.ebi.ac.uk/gwas) for the studies listed in
Supp. Table 5. We used the mapped genes for each locus as the input gene list to calculate the
crosstalk enrichments in each study. We used the same FET approach described above for the
AD-related crosstalk interactions to calculate the enrichment of genes nominated by each
GWAS study (Figure 2D). To calculate the enrichment of crosstalk interactions involving genes
associated with AD GWAS based on co-accessibility (Figure 2E), we used as the input gene list
all genes with a transcription start site (TSS) region either 1) overlapping an ATAC-seq peak co-
accessible with another ATAC-seq peak harboring a fine-mapped AD GWAS variant (PPA >
0.01) from the Schwartzentruber *et al.* study (*14*) or 2) overlapping an ATAC-seq peak in any
cell type harboring a fine-mapped variant.

Downstream crosstalk networks reconstruction using CytoTalk

1037

To identify the signaling networks downstream of crosstalk interactions, we used CytoTalk (v.
4.0.3) with minor modifications. CytoTalk first reconstructs the co-expression network of two cell
types using information theory, then prioritizes biologically relevant crosstalk interactions based
on their connection to central genes in each network using graph theory. This analysis results in
a prioritized network of genes predicted to mediate the crosstalk signals between the two cell
types. Briefly, CytoTalk reconstructs the co-expression network for each cell type pair and
connects them based on a database of ligand-receptor interactions. It then uses a prize-
collecting Steiner forest algorithm to identify the crosstalk signaling network between the two
cell types (*27*). We generated input csv files per cell type for each donor category containing the
sctransform-normalized snRNA-seq log-transformed counts per barcode. Additionally, we
increased the stringency of the CytoTalk networks by manually inputting only interactions that
were also present in CellPhoneDB. This step was done because CytoTalk uses protein-protein
interactions and text mining from STRING-DB (*67*) to build its crosstalk interactions database,
which we consider overly permissive. For the network visualization in **Figure 2a**, we used the
union of the crosstalk signaling network identified for each donor category in microglia versus
excitatory neurons. We generated network visualizations using Cytoscape (v. 3.9.0) (*68*) and
identified crosstalk sub-networks using the community cluster (GLay) function of clusterMaker
(v. 2.2) (*69*). The modified CytoTalk version used in this study
(https://github.com/rdalbanus/CytoTalk) fixed minor bugs and enabled better parallelization
control for a cluster environment. These issues have since been fixed in newer CytoTalk
releases.

To calculate the enrichment of AD-related genes downstream of the crosstalk interactions
(**Figure 2c**), we used FETs to compare the number of AD-related genes in each side of the exc.
neuron-microglia crosstalk network versus all AD-related genes expressed in the corresponding
cell type. All FET p-values were corrected for multiple testing using the Bonferroni correction
based on the total number of tests performed for each analysis.

Network analyses

1064

We used the *compute_mutual_information_single* function from CytoTalk to reconstruct the
excitatory neurons and microglia co-expression networks for each donor category. Briefly, this
function calculates the mutual information (MI) (*70*) across all pairs of genes per cell type and
then applies the ARACNE algorithm (*71*) to eliminate most indirect interactions between genes.
We then merged the ARACNE-filtered MI matrices for each cell type across donor categories by
keeping the highest MI value for each interaction. In order to identify co-expression sub-
networks, we applied the WGCNA pipeline (*72*) on the merged MI matrix. To test the
association of each sub-network (n = 360) with the Braak stage, we first calculated its
eigengene (the first principal component of gene expression) for each sub-network. We then
tested the association of each eigengene with a high Braak stage using the binomial regression
model:

$$Braak\ high_i \sim eigengene_{ij} + sex_i + APOE4_i + TREM2_i$$

In this model, i represents a barcode and j, a co-expression sub-network. $Braak\ high_i$, sex_i,
$APOE4_i$, and $TREM2_i$ are binary variables encoding the donor Braak stage (Braak > 3 versus

Braak ≤ 3), sex, *APOE* genotype (one or two copies of *APOE4* versus zero), and *TREM2*
genotype (common variant versus R47H, R136W, or R62H). We ran this model for each sub-
network and obtained the corresponding coefficient and p-value of the eigengene term. Then,
we adjusted the significance for multiple testing using the Bonferroni correction. Similarly, we
performed this same analysis in the sub-networks obtained from the CytoTalk microglia-exc.
neurons crosstalk network.

Functional enrichments

All gene ontology (GO) functional enrichments in this study were calculated with the
WebGestaltR R package (v. 0.4.4), using the *ORA* method, the
geneontology_Biological_Process_noRedundant annotation, and the *genome_protein-coding*
reference set. For all analyses, we used an FDR threshold of 1 to recover all enrichments and
manually adjusted p-values for multiple testing using the Bonferroni correction (significance
threshold used: $p < 0.05$). To further collapse redundancy in GO terms, we used the R package
rrvgo (v. 1.9.1). For each WebGestaltR analysis, we generated a similarity matrix across all GO
terms significant in at least one comparison using the *calculateSimMatrix* function. We then
reduced the significant terms for each analysis using the *reduceSimMatrix* function. The
threshold values for the *reduceSimMatrix* function were selected based on visual inspection of
the resulting terms (0.8 for the analyses in Figures 1F and 3C and 0.9 for Figure 3F). Finally, we
reported the most significant p-value among the grouped terms under each parent term.

Data availability

1098 iMGL RNA-seq data will be publicly available in GEO upon publication. The snRNA-seq data
1099 from the Knight ADRC is publicly available by request from the National Institute on Aging
1100 Genetics of Alzheimer's Disease Data Storage Site (NIAGADS) under accession number
1101 NG00108 (https://www.niagads.org/datasets/ng00108). DIAN brain bank snRNA-seq data
1102 access requires a request through https://dian.wustl.edu/our-research/for-investigators.

Code availability

1104 All scripts necessary to reproduce the figures from this manuscript will be available at
1105 https://github.com/albanus-research/2021_trem2_adad_snRNAseq. The modified CytoTalk
1106 version used for this manuscript is available at https://github.com/rdalbanus/CytoTalk.

1107

a

Full data

Downsampled data

b

c

Single brain

d

Subset of cell types

1108

1109 **Supplementary Figure 1. a)** Overview of CellPhoneDB results on full dataset versus a heavily
1110 downsampled dataset (maximum 100 nuclei per snRNA-seq cluster). **b)** Fraction of replicated
1111 CPDB interactions in downsampled dataset compared to the full dataset.

1112

1113 **Supplementary Figure 2.** Global changes in crosstalk frequencies per cell type across donor
1114 categories versus controls.

1115

1116 **Supplementary Figure 3.** Gene ontology enrichments of the genes associated with crosstalk
1117 interactions only detected in cases or controls, grouped by cell type.

1118

1119

1120 **Supplementary Figure 4. a)** Bias of expressed genes in each cell type belonging to a CPDB
1121 interaction. b) Fraction of AD-related genes expressed in each cell type that participate in any
1122 CPDB interaction.

1123

1124

1125

1126 **Supplementary Figure 5.** Enrichment of AD crosstalk interactions involving each cell type in
1127 each donor category for different snRNA-seq datasets (left plots) and their corresponding
1128 enrichment of AD crosstalk interactions between microglia and other cell types (blowout panels).

1129

1130

1131 **Supplementary Figure 6. a)** Reconstructed exc. neurons and microglia crosstalk networks
1132 using CytoTalk in different snRNA-seq datasets. **b)** Reconstructed TREM2 crosstalk network
1133 with CytoTalk by downsampling the number of nuclei to the donor category with the smallest
1134 number of nuclei.

1135

1136

1137 **Supplementary Figure 7.** Comparison of the TREM2 subnetwork prioritized by CytoTalk **(a)**
1138 and identified directly from the microglia co-expression network **(b)**.

1139

1140

Supplementary Figure 8. a) Schematic of the differentiation procedure and brightfield imaging

of iPSC-derived microglia (iMGL). **b)** mature iMGL stained with microglial markers, TREM2,

Iba1, and TMEM119 and co-stained with DAPI. **c)** Schematic of DNA (red) targeted by gRNA

against TREM2 for CRISPR/Cas9 modification and showing an 8 base pair homozygous

deletion within exon 2 of the TREM2 gene. **d)** Western blot analysis of wild-type (WT) and

TREM2 KO iPS cell lysates immunostained for TREM2 with β-actin as the loading control.

1147

1148

Supplementary Figure 9. a) Phagocytosis quantification using Incucyte live-cell analysis
system on iMGL treated with pHrodo-labeled synaptosomes (syn). **b)** Fluorescent and
brightfield microscopy of iMGL treated with pHrodo-labeled synaptosomes (syn) with/out
cytochalasin D. **c)** Phagocytosis quantification using Incucyte live-cell analysis system on iMGL
treated with pHrodo-labeled Aβ oligomer (Abo) and Aβ fibril (Abf). **d)** Fluorescent microscopy of
iMGL treated with pHrodo-labeled Aβo with/out cytochalasin D and co-stained with 5-CFDA
membrane marker.

1156

Supplementary Figure 10. a) Volcano plots for the differentially expressed genes in different
iMGL RNA-seq experiments. Red genes correspond to differentially expressed at 10% FDR. **b)**
Effect size distribution of the three iMGL experiments. **c)** Effect size distribution of
downregulated genes in TREM2 KO iMGL from the McQuade et al. 2018 study in the three
iMGL RNA-seq experiments. **d)** Comparison of reported effect sizes from the McQuade study
and the TREM2 KO vs. WT iMGL RNA-seq experiment from this study.

1163