NhaR, LeuO and H-NS are part of an expanded regulatory

network for ectoine biosynthesis expression

Katherine E. Boas Lichty, Gwendolyn J. Gregory, E. Fidelma Boyd*

Department of Biological Sciences, University of Delaware, Newark, DE, 19716

Corresponding author*

E. Fidelma Boyd

Department of Biological Sciences

341 Wolf Hall, University of Delaware
Newark, DE 19716

Phone: (302) 831-1088. Fax: (302) 831-2281 Email: <u>fboyd@udel.edu</u>

Keywords: Ectoine biosynthesis regulation, *Vibrio*, LeuO, NhaR, H-NS, ToxR

Abstract

Bacteria accumulate compatible solutes, to maintain cellular turgor pressure when

exposed to high salinity. In the marine halophile *Vibrio parahaemolyticus*, the compatible

solute ectoine is biosynthesized *de novo*, which is energetically more costly than uptake;

therefore, tight regulation is required. To uncover novel regulators of the ectoine

biosynthesis *ectABC-asp_ect* operon, a DNA affinity pulldown of proteins interacting

with the *ectABC-asp_ect* regulatory region was performed. Mass spectrometry analysis

identified, amongst others, three regulators: LeuO, NhaR, and the nucleoid associated

protein H-NS. In-frame non-polar deletions were made for each gene and P_{ectA}-*gfp*

promoter reporter assays were performed in exponential and stationary phase cells.

P_{ectA}-*gfp* expression was significantly repressed in the Δ*leuO* mutant and significantly

induced in the Δ*nhaR* mutant compared to wild type, suggesting positive and negative

regulation, respectively. In the Δ*hns* mutant, P_{ectA}-*gfp* showed increased expression in

exponential phase cells, but no change compared to wild type in stationary phase cells.

To examine whether H-NS interacts with LeuO or NhaR at the ectoine regulatory

region, double deletion mutants were created. In a Δ*leuO*/Δ*hns* mutant, P_{ectA}-*gfp* showed

reduced expression, but significantly more than Δ*leuO* suggesting H-NS and LeuO

interact to regulate ectoine expression. Whereas Δ*nhaR*/Δ*hns* had no additional effect as

compared to Δ*nhaR* suggesting NhaR regulation is independent of H-NS. To examine

leuO regulation further, a P_{leuO}-*gfp* reporter analysis was examined that showed

significantly increased expression in the Δ*leuO*, Δ*hns* and Δ*leuO*/Δ*hns* mutants as

compared to wild type, indicating both are repressors. Growth pattern analysis of the

mutants in M9G 6%NaCl, showed growth defects compared to wild type, indicating

that these regulators play an important physiological role in salinity stress tolerance.

Importance: Ectoine is a commercially used compatible solute that acts as a biomolecule

stabilizer because of its additional role as a chemical chaperone. A better understanding

of how the ectoine biosynthetic pathway is regulated in natural bacterial producers can

be used to increase efficient industrial production. The *de novo* biosynthesis of ectoine is

essential for bacteria to survive osmotic stress when exogenous compatible solutes are

absent. This study identified LeuO as a positive regulator and NhaR as a negative

regulator of ectoine biosynthesis and also showed that similar to enteric species, LeuO

is an anti-silencer of H-NS. In addition, defects in growth in high salinity among all the

mutants suggest that these regulators play a broader role in the osmotic stress response

beyond ectoine biosynthesis regulation.

Introduction

When bacteria are exposed to high salinity conditions, an efflux of water across the osmotic gradient occurs lowering cellular turgor pressure (1-5). The initial short-term response to osmotic up shock is the uptake of potassium (K+), which is accompanied by glutamate accumulation in Gram-negative bacteria to maintain electro neutrality. This is the short-term strategy because high concentrations of K+ have deleterious effects on cellular processes (1, 2, 6, 7). The long term response to increased osmotic stress is characterized by the uptake or biosynthesis of compatible solutes, which are small organic molecules that provide osmotic protection by helping to maintain the turgor pressure of the cell and can be accumulated to extremely high intracellular concentrations (3, 8, 9).

The compatible solute ectoine was initially discovered to provide osmotic protection in anoxygenic phototrophs (10), but subsequent analysis showed that it functions as a compatible solute in numerous Gram-negative and Gram-positive bacteria (reviewed in (3, 8, 11-14), which included many members of the family *Vibrionaceae* (15-19). Accumulation to very high intracellular concentrations without interfering with molecular processes is a hallmark of compatible solutes. Additionally, ectoine acts as a chemical chaperone through the stabilization of proteins, thus it is an important compound for commercial uses in medicine and cosmetics to stabilize biomolecules against factors such as heating, freezing, desiccation, and UV radiation (20-23).

Therefore, understanding the regulation of the ectoine biosynthetic pathway is

important to increase efficient industrial production using natural bacterial producers

(24).

Biosynthesis of ectoine (1,4,5,6-tetrahydro-2-methyl-4-pyrimidinecarboxylic acid) is *de*

novo from L-aspartic acid, and performed by the evolutionarily conserved operon

ectABC, which encodes the EctA, EctB, and EctC proteins (25-27) (**Fig. S1**). In many

species an aspartokinase (*asp_ect*) is clustered with the ectoine biosynthesis operon,

which converts aspartic acid to β-aspartyl phosphate, which is then converted to L-

aspartate-β-semialdehyde by aspartate semialdehyde dehydrogenase (Asd). This

intermediate from the aspartic acid pathway is incorporated into the ectoine

biosynthesis pathway when it is converted to L-2,4-diaminobutyrate by EctB (L-2,4-

diaminobutyrate transaminase). EctA (L-2,4-diaminobutyrate N^γ-acetyltransferase) then

acetylates this product to form N^γ-acetyldiaminobutyrate. The final step to produce

ectoine is a cyclic condensation reaction performed by EctC (L-ectoine synthase) (13,

27).

In some species, clustered with the ectoine biosynthesis operon is *ectR*, a MarR-type

regulator shown to repress transcription of the ectoine operon (24, 28-30). Among

Vibrionaceae, an *ectR* homolog is present only in *Aliivibrio* species (31, 32). Previous

studies in *V. parahaemolyticus* and *V. cholerae* showed that *ectABC-asp_ect* transcription

was repressed in low salinity by another MarR-type regulator named CosR (17, 31, 33).

CosR from *V. parahaemolyticus* showed 31% amino acid identity to *Aliivibrio fischeri* EctR,

with less than 60% query coverage and does not cluster with *ectABC-asp_ect*, but is

located elsewhere in the genome (31). The quorum sensing master regulators AphA and

OpaR activate and repress expression of *ectABC-asp_ect*, respectively in *V.*

parahaemolyticus (17). AphA and OpaR also regulate expression of CosR, creating a

tightly controlled feed-forward loop for ectoine expression across the growth curve (17).

In this study, we set out to identify novel regulators of the ectoine biosynthesis genes by

performing a DNA affinity chromatography pulldown using the regulatory region of

the *ectABC-asp_ect* operon. We identified numerous candidates and selected the

regulators LeuO, NhaR, OmpR, TorR, and the nucleoid associated protein (NAP) H-NS

to examine further. We constructed in-frame nonpolar deletions in each gene. These

regulators play a role in the regulation of the ectoine biosynthesis genes based upon

P_{ectA}-*gfp* expression reporter assays. In addition, analysis of double mutants Δ*leuO*/Δ*hns*

and Δ*nhaR*/Δ*hns* showed that H-NS interacted with LeuO but not NhaR to control

ectoine gene expression. Importantly, when these deletion mutants were grown under

high salinity conditions a growth defect was observed, suggesting that the regulators

NhaR, LeuO, and H-NS are physiologically important for growth in high salinity

conditions.

Results

Ectoine is biosynthesized in cells grown in M9G 3%NaCl, but absent in M9G 1%NaCl. To confirm *V. parahaemolyticus de novo* ectoine biosynthesis is a response to increased salinity, ^{1}H nuclear magnetic resonance spectroscopy (^{1}H-NMR) at 600MHz in the presence of deuterium oxide as the solvent was performed. Ethanol extracts of *V. parahaemolyticus* wild type cells grown overnight at 37°C in minimal medium supplemented with glucose (M9G) with either 1%NaCl or 3%NaCl were used to obtain ^{1}H-NMR spectra (**Fig. S2A**). Peaks corresponding to the hydrogen atoms of ectoine are labeled and were present in M9G 3%NaCl extracts but were not detectable in M9G 1%NaCl extracts as previously shown (15) (**Fig. S2A**). In addition to intracellular accumulation of ectoine, we also wanted to demonstrate that *ectABC-asp_ect* expression is increased in M9G 3%NaCl compared to 1% NaCl. A transcriptional reporter assay was performed, where *gfp* was placed under the control of the ectoine regulatory region (P$_{ectA}$-*gfp*). The expression was examined in wild type cells grown in either M9G 1%NaCl or M9G 3%NaCl by measuring the specific fluorescence of P$_{ectA}$-*gfp* as a cumulative readout of *ectA* transcription (**Fig. S2B**). The observed level of specific fluorescence of P$_{ectA}$-*gfp* was significantly increased in M9G 3%NaCl (4.0-fold) relative to that of cells grown in M9G 1%NaCl (**Fig. S2B**).

Identification of putative regulators of the ectoine biosynthesis operon. With the knowledge that ectoine is present in cells grown in M9G 3%NaCl, we performed a

promoter pulldown under this growth condition. A promoter pulldown was performed

using a 5′-biotin tagged 323-bp DNA fragment of the *ectABC-asp_ect* regulatory region

bound to streptavidin beads. As a negative control DNA probe, the coding region of the

ectB gene was used. To identify novel regulators of ectoine biosynthesis, we grew cells

to mid-exponential phase in M9G 3%NaCl and subsequent cell lysate was incubated

with DNA probe-coated beads. Bound proteins were then eluted using a stepwise NaCl

gradient and separated with SDS-PAGE gel (**Fig. S3A**). Bands that were present in the

target lanes, but not present in the *ectB* negative control lanes were selected for analysis

via mass spectrometry (**Fig. S3B**). The proteins identified with mass spectrometry

analysis were aligned to the *V. parahaemolyticus* genome and sorted by gene class and

ranked by score (**Table S1**). The mass spectrometry score is the combined sum of all

scores for an observed spectra that can be matched to an individual peptide. The higher

the score the more confident the match to the specified peptide. Candidate proteins of

interest that belonged to the transcriptional regulator gene class were selected. These

candidate proteins are described in Table S1 and include homologs of regulators NhaR

(VP0527), OmpR (VP0154), and TorR (VP1032). NhaR and OmpR were shown in

Escherichia coli, which does not produce ectoine, to be involved in the osmotic stress

response (34-39). *V. parahaemolyticus* NhaR shows 62%, OmpR 84%, and TorR 57%

amino acid identities to homologs in *E. coli*. NhaR, OmpR, and TorR have not been

examined in *Vibrio* species. An additional regulators of interest, LeuO (VP0350), a LysR-

type regulator and H-NS (histone-like nucleoid structuring) protein were also examined

further. In *V. parahaemolyticus*, LeuO was previously shown to be positively regulated

by ToxR and is essential for acid stress tolerance (40, 41). H-NS in *E. coli* and *Salmonella*

enterica, is a universal silencer of gene expression (42-47). In addition, an *E. coli* study by

Shimada and colleagues identified 140 LeuO-binding sites with over 90% of these sites

also showing H-NS binding. Further studies have shown that LeuO plays a significant

role as an antagonist of H-NS gene silencing (42, 48, 49). While not chosen for

examination in this study, two additional regulators did stand out for possible future

analysis, VPA0036 (accession number Q87K62) and VP0489 (accession number Q87SD5)

(**Table S1**). VPA0036 gave a high score, but is annotated as a hypothetical protein with

no conserved domains. HHPred analysis uncovered a putative N-terminal winged-helix

protein domain with a probability of 96.53% and E-value 0.016. VP0489 is annotated as

an aerobic respiration control protein FecA and ArcA a two-component system

response regulator.

As a proof of concept for the pulldown, LeuO and NhaR proteins were purified and

visualized on SDS-PAGE gels showing an expected 36.2 kDa LeuO protein and a 33.9

kDa NhaR protein (**Fig. S4**). Electrophoretic mobility shift assays (EMSAs) were

performed first using the 323-bp regulatory region of *ectABC-asp_ect* as a DNA probe

(**Fig. S5A**). For the LeuO EMSA, we used the regulatory region of the *leuO* gene as a

positive control and show binding in a concentration dependent manner for this and

also for the P*ectA* probe 1 (**Fig. S5B**). Binding of NhaR to P*ectA* was also confirmed in

the EMSA (**Fig. S5C**). To examine binding of LeuO and NhaR further, probe 1 was

divided into three similarly sized probes P*ectA* 1A, 1B and 1C. Purified protein LeuO

showed band shifts for probe 1A and a complete shift for probe 1C (**Fig. S5D**). For

NhaR all three probes showed shifts in a concentration dependent manner (**Fig. S5E**).

P*ectA*-*gfp* reporter assays shows differential expression in Δ*leuO*, Δ*nhaR*, Δ*ompR* and

Δ*torR* mutants. To examine the role of *leuO, nhaR, ompR,* and *torR* candidate regulators,

we performed GFP reporter assays using a reporter plasmid where *gfp* was under the

control of the ectoine regulatory region (P*ectA*-*gfp*). This reporter plasmid was

transformed into wild type or in-frame non-polar mutants of Δ*leuO,* Δ*nhaR,* Δ*ompR,* and

Δ*torR*. Cultures were grown overnight in M9G 3% NaCl and relative fluorescence

intensity (RFU) was measured. Specific fluorescence was calculated by dividing RFU by

OD. From this analysis, the relative fluorescence (RFU) measured in the Δ*leuO* mutant

background was 1.7-fold reduced as compared to wild type (**Fig. 1**). This data suggests

that LeuO is an activator of *ectABC-asp_ect*. The relative fluorescence measured in a

Δ*nhaR* background was significantly increased 1.8-fold as compared to wild type,

suggesting that NhaR is a repressor of *ectABC-asp_ect* (**Fig. 1**). P*ectA*-*gfp* expression in the

Δ*ompR* (1.3-fold) and Δ*torR* (1.4-fold) mutants were also increased as compared to wild

type, but not to the same level as NhaR.

LeuO and H-NS interplay to control P$_{ectA}$-*gfp*. In *E. coli* and *V. cholerae*, H-NS has been shown to interact with LeuO in both cooperative and antagonistic relationships at different gene loci (42, 50-53). Therefore, we wanted to examine whether H-NS and LeuO interacted at the *ectABC-asp_ect* regulatory region. We constructed an in-frame non-polar Δ*hns* mutant and a double Δ*leuO*/Δ*hns* mutant. In exponential phase cells (OD ~ 0.4), P$_{ectA}$-*gfp* in the Δ*leuO* mutant showed significant reduced expression, and in the Δ*hns* mutant expression was significantly increased (**Fig. 2A**). However, in the Δ*leuO*/Δ*hns* double mutant P$_{ectA}$-*gfp* expression was reduced, but not to the same level as the Δ*leuO* single mutant (**Fig. 2A**). In stationary phase cells (OD ~ 1.0), P$_{ectA}$-*gfp* relative fluorescence was similar to exponential cells in all strains except the Δ*hns* mutant, which was similar to wild-type (**Fig. 2B**). These data indicate interplay between both LeuO and H-NS to control ectoine gene expression with LeuO playing an anti-silencer role.

To further investigate the relationship between LeuO and H-NS in *V. parahaemolyticus*, a *gfp*-expressing reporter plasmid under the control of the LeuO regulatory region (P$_{leuO}$-*gfp*) was constructed. Relative fluorescence was measured in cells grown to OD ~1.0 in M9G 3%NaCl in wild type, Δ*leuO*, Δ*hns*, Δ*leuO*/Δ*hns*, and Δ*toxR*. When compared to wild type, P$_{leuO}$-*gfp* expression in the Δ*hns* mutant showed a 4.0-fold increased relative fluorescence indicating that H-NS is a repressor of *leuO* expression (**Fig. 3**). P$_{leuO}$-*gfp* expression was 2.4-fold higher in Δ*leuO* and 2.8-fold higher in Δ*leuO*/Δ*hns* when compared to wild type. In *V. parahaemolyticus*, ToxR is a positive regulator of *leuO*

expression (40, 54, 55) and this was confirmed in this reporter assay, with Δ*toxR*

showing reduced P$_{leuO}$-*gfp* (1.8-fold) expression when compared to wild type (**Fig. 3**).

These data indicate that both H-NS and LeuO are repressors of *leuO* expression and in

the absence of both negative regulators other proteins such as ToxR are likely able to

access the regulatory region.

H-NS and LeuO are important for growth in high salinity. To assess whether the

deletions of *leuO* and *hns* have a physiological effect, we examined the ability of *V.*

parahaemolyticus wild type, Δ*leuO*, Δ*hns*, and Δ*leuO*/Δ*hns* mutants to grow in M9G 1%

NaCl, 3%NaCl, or 6%NaCl at 37°C. In M9G 1%NaCl and 3%NaCl, there were no

significant growth defects observed among most strains, with the exception of Δ*leuO*,

which reached a lower final OD (0.44) as compared to wild type (0.57) (**Fig. S6A, 6B**).

Growth in M9G 6%NaCl showed the wild type and Δ*hns* had a 5 h lag phase, but the

Δ*hns* mutant reached a lower final OD (0.37) as compared to wild type (0.54) (**Fig. 4**).

The Δ*leuO* mutant had a lag phase of 6 h and Δ*leuO*/Δ*hns* had a 13 h lag phase. In

addition, the double deletion mutant had a final OD$_{595}$ of 0.2, which was significantly

less than wild type with final OD of 0.54 (**Fig. 4**). Complementation of Δ*leuO* with

pBA*leuO* and Δ*hns* with pBA*hns*, restored growth to wild type levels (**Fig. 4**). However,

we were unable to complement the double mutant with either pBA*leuO* or pBA*hns*.

Overall, these data demonstrate that both LeuO and H-NS play a significant role in the

osmotic stress response and are required for optimal growth in high salinity.

NhaR is a negative regulator of the ectoine biosynthesis operon. Next, we considered

whether there was interplay between NhaR and H-NS in the regulation of *ectABC-ask*.

To accomplish this, we examined P$_{ectA}$-*gfp* in Δ*nhaR*, Δ*hns* and a Δ*nhaR*/Δ*hns* double

mutant in M9G 3%NaCl in exponential and stationary phase cells (**Fig. 5**). In

exponential cells (~OD 0.4). P$_{ectA}$-*gfp* expression was higher in Δ*nhaR* (1.8-fold) as

compared to wild type confirming NhaR as a negative regulator (**Fig. 5A**). P$_{ectA}$-*gfp*

expression in the Δ*hns* and Δ*hns*/Δ*nhaR* mutants were both similarly increased

compared to wild type (**Fig. 5A**). Examination of stationary phase cells (OD ~ 1.0) gave

similar results to expression in exponential cells with the exception of Δ*hns*, which

showed P$_{ectA}$-*gfp* expression levels similar to wild type (**Fig. 5B**). Overall, the data

suggest that there is no interplay between H-NS and NhaR in ectoine expression

because the double mutant was not significantly different than the single mutants.

Next, we examined the effect of deleting *nhaR*, *hns*, and *hns*/*nhaR* on growth of *V.*

parahaemolyticus in high salinity. Growth of the deletion mutants in M9G 1%NaCl and

3% NaCl was similar to wild type indicating no overall defect (**Fig. S6C, D**). However,

Δ*nhaR*, Δ*hns*, and Δ*hns*/Δ*nhaR* mutants grown in M9G 6%NaCl, reached a final OD$_{595}$ of

0.37, 0.36, and 0.38 respectively as compared to wild type OD$_{595}$ of 0.5 (**Fig. 6**).

Complementation of Δ*nhaR* with pBA*nhaR* restored growth to wild type levels (**Fig. 6**).

These data indicates that there is a growth defect in high salinity for Δ*nhaR*, but deletion

of *hns* does not contribute further to this defect. This supports the expression data that

NhaR and H-NS do not interact to control ectoine expression. The data does indicate

that NhaR is important for high salinity tolerance.

Discussion

The accumulation of compatible solutes occurs either via uptake or biosynthesis and in

V. parahaemolyticus there are six transporters and two biosynthesis operons for

compatible solutes present (19, 56). The biosynthesis of the compatible solute ectoine

and not glycine betaine was previously shown to be essential for *V. parahaemolyticus*

survival in high salinity (15). However, ectoine biosynthesis is costly as it drains

intracellular pools of metabolites such as aspartic acid and multiple regulators that

provide tight control are needed (57).

In this study, we identified several new regulators of ectoine biosynthesis (**Fig. 7**). We

found that NhaR acted as a repressor of ectoine biosynthesis. However, deletion of this

repressor, showed a defect when grown under high salinity conditions (6% NaCl). This

could be explained by the fact that NhaR is known to regulate genes related to the

osmotic stress response outside of ectoine biosynthesis (58-61). In *E. coli*, *nhaR* is present

in an operon with *nhaA*, which encodes a Na+/H+ antiporter. NhaR was shown to

activate transcription of *nhaA*, which is induced in high salt and alkaline pH and is

essential for survival under these conditions. It was also suggested that NhaR is a

sensor and transducer of Na+ because the presence of Na+ alters the contact points

between NhaR and the *nhaA* regulatory region (62). A homolog of *nhaA* is present in *V.*

parahaemolyticus, and thus may also be activated by NhaR, which could explain the

growth defect in high salinity for the *V. parahaemolyticus ΔnhaR* mutant.

Our data also established that LeuO is a positive regulator of ectoine biosynthesis gene

expression and H-NS acts as a gene silencer. The analysis of a double *ΔleuO/Δhns*

mutant suggested that LeuO also likely plays a role as an anti-silencer of H-NS at this

locus. In addition, we found that LeuO and H-NS both act as repressors of *leuO* in *V.*

parahaemolyticus, further supporting the interaction between these two regulators.

LeuO was first identified as a LysR-type regulator and shown to positively regulate the

leuABCD leucine synthesis operon in *Salmonella enterica* (63, 64). Subsequently,

chromatin-immunoprecipitation-on-chip analysis showed 178 LeuO binding sites in *S.*

enterica, indicating its role as a global regulator (65). H-NS was also shown to be a global

regulator in this species and in *E. coli*, silencing the transcription of a large number of

horizontally acquired genes (46, 47, 66). In *V. parahaemolyticus*, LeuO was first named

CalR as it was negatively regulated by calcium (41). LeuO was shown to positively

regulate *ompU*, which encodes an outer membrane porin important for acid stress (**Fig.**

7) (40). Additionally, the global regulator ToxR was shown to be a positive regulator of

leuO expression and deletion of either *leuO* or *toxR* resulted in a growth defect in acid

conditions (40). In our study, *leuO* showed significantly increased expression in the

ΔleuO mutant whereas in the *ΔtoxR* mutant *leuO* expression was reduced showing it is

an activator (**Fig. 7**). In *V. vulnificus*, H-NS was shown to repress *leuO* by binding to sites

that overlap with ToxR and LeuO binding sites (67, 68). Our data also showed that *leuO*

is an auto-repressor and that H-NS represses *leuO* in *V. parahaemolyticus*, which is likely

using a similar regulatory mechanism to that in *V. vulnificus* (**Fig. 7**). Our growth data

showed that deletion of either *leuO* or *hns* resulted in defects when grown at high

salinity and a double Δ*leuO*/Δ*hns* mutant showed a severe growth defect that was more

pronounced than either of the single mutants. The observed defects in high salinity

suggests a larger role for both regulators in the osmotic stress response. The more

pronounced growth defect observed for Δ*leuO*/Δ*hns* reinforces the hypothesis that these

proteins interact at other loci important for increased salinity toleranace.

Methods

Bacterial strains, media and culture conditions. All strains and plasmids used in this

study are listed in Table S2. *V. parahaemolyticus* RIMD2210633, a streptomycin-resistant

clinical isolate, was used in this study as the wild-type (WT) strain (69, 70). Unless

stated otherwise, *V. parahaemolyticus* was grown in either lysogeny broth (LB; Fisher

Scientific, Fair Lawn, NJ) with 3% (wt./vole) NaCl (LB 3%) or M9 minimal media (47.8

mM Na_2HPO_4, 22 mM KH_2PO_4, 18.7 mM NH_4Cl, 8.6 mM NaCl; Sigma Aldrich)

supplemented with 2 mM $MgSO_4$, 0.1 mM $CaCl_2$, 20 mM glucose as the sole carbon

source (M9G) and 3% (wt/vol) NaCl at 37°C. *E. coli* strains were grown in either LB

supplemented with 1% (wt/vol) NaCl (LB1%) or M9G supplemented with 1% (wt/vol)

NaCl. The strain *E. coli* β2155 λ*pir* strain is a diaminopimelic acid (DAP) auxotrophic

strain and was grown with 0.3 mM DAP (71). All strains were grown at 37°C with

aeration. Antibiotics were added to growth media at the following concentrations as

necessary: ampicillin (Amp), 100 µg/mL; chloramphenicol (Cm), 12.5 µg/mL;

tetracycline (Tet), 1 µg/mL; kanamycin (Km), 50 µg/mL.

Preparation of cellular extracts and proton nuclear magnetic resonance (^{1}H-NMR).

Wild-type *V. parahaemolyticus* was grown overnight at 37°C in either M9G 1% NaCl or

M9G 3%NaCl. Stationary phase cells were then pelleted and washed twice with 1XPBS.

Three freeze-thaw cycles were performed with the cell pellets to increase lysis, and the

cells were then suspended in 750 µL ethanol. Debris was pelleted by centrifugation, and

the ethanol solution was transferred to a clean tube and evaporated under vacuum.

Then deuterium oxide (D_2O) was used to resuspend the pellet, and insoluble material

was removed by centrifugation. The solution was then transferred to a 5-mm NMR tube

for analysis on a Bruker AVANCE 600NMR spectrometer at a proton frequency of

600.13 MHz with a sweep of 12,376 Hz and a relaxation delay of 5 s. Sixteen scans were

co-added for each spectrum.

DNA-Affinity Pulldown. DNA-affinity chromatography pulldown was performed as

previously described (72-74). The bait DNA primers were designed to amplify the 323-

bp regulatory region of *ectA* (VP1722) with a biotin moiety added to the 5′ end. A 327-

bp negative-control bait DNA probe was amplified from a coding region of *ectB*

(VP1721), which was also labeled with a biotin moiety at the 5′ end. Both probes were

amplified using Phusion HF polymerase (New England Biolabs) PCR. Ten reactions

were completed for each probe and then pooled and purified using an ethanol

extraction technique (75). The pulldown was completed twice, once after growth in non-

inducing conditions (LB3%) and once after growth in inducing conditions (M9G3%) in

an effort to obtain both positive and negative regulators of the ectoine biosynthesis

operon. The wild-type strain was grown overnight in LB3%, cells were pelleted and

washed two times with 1XPBS and diluted 1:50 into 500 mL of LB3% or M9G3%. The

cultures were then grown to an OD_{595} of 0.5, pelleted and stored overnight at -80°C. The

pellet was then suspended in 1.5 mL Fast-Break lysis buffer (Promega, Madison, WI)

with 1 mM PMSF, 0.5 mM benzamidine, 0.1 µg/mL lysozyme and incubated for 30 min.

at room temperature. The cells were then sonicated on ice at 30% pulse amplitude for 15

seconds with 1 min. rest, and this was repeated 5 times. The cell lysate was then

clarified by centrifugation for 30 min. at 4°C. The cell lysate was precleared with

streptavidin DynaBeads (Thermo Scientific, Waltham, MA) to remove nonspecific

protein-bead interactions.

Streptavidin M280 DynaBeads (ThermoFisher) were then incubated two times with 10

µg of biotinylated probe for 20 min. The precleared cell lysate was then incubated for 30

min. with the bait-coated beads at room temperature with constant rotation in the

presence of 100 µg sheared salmon sperm DNA as a non-specific competitor. This was

completed twice with washes in between. Protein candidates were then eluted from the

bait DNA-bead complex with a stepwise NaCl gradient (100 mM, 200 mM, 300 mM, 500

mM, 170 mM, and 1 M). Next, 6X SDS and 1 mM β-mercaptoethanol (BME) was added

to elution samples which were then boiled at 95°C for 5 min. Samples were then

separated using a 12% SDS-PAGE and visualized using the Pierce Silver Stain Kit

(ThermoFisher). The P$_{ectA}$ bait and *ectB* negative control were loaded next to each other

in order of increasing NaCl concentrations. Bands that were present in the P$_{ectA}$ bait but

not the negative control lanes were excised, de-stained, and digested with trypsin

following the standard procedure for mass spectrophotometry preparation with C18

ZipTips (Fisher Scientific, Fair Lawn, NJ). The samples were then individually analyzed

with a Q-exactive Orbitrap mass spectrometer with nano-flow electrospray (Thermo).

Proteins were subsequently identified using Proteome Discover 2.1 software.

Protein Purification. Protein were purified as previously described (76). The full-length

genes for *leuO* (VP0350) and *nhaR* (VP0527) were cloned into a pET-28a(+) vector with a

C-terminal histidine tag, transformed into *E. coli* Dh5α, purified, and sequenced. These

vectors pET*leuO* and pET*nhaR* were then transformed into *E. coli* BL21 (DE3) cells, and

expression was induced with 0.5mM IPTG at 0.4 OD$_{595}$ and then cells were grown

overnight at 25°C. Cells were then pelleted and resuspended with lysis buffer (50 mM

sodium phosphate, 200 mM NaCl, 20 mM imidazole, pH 7.4, 1.0 mM

phenylmethanesulfonyl fluoride, 0.5 mM benzamidine) and lysed via sonication. Debris

was pelleted, and supernatant applied to an IMAC column packed with HisPur Ni-

NTA resin (ThermoFisher) that was equilibrated with column buffer (50 mM sodium

phosphate, 200 mM NaCl, 20 mM imidazole, pH 7.4). The column was washed with

buffer containing increasing concentrations of imidazole (20mM – 100mM) to remove

any contaminants. His-tagged proteins were eluted using 500mM imidazole and then

dialyzed overnight at 4°C in sodium phosphate buffer to remove any excess salts.

Samples of supernatant and each flow through, wash, and elution were analyzed via

SDS-PAGE gel to assess protein purity and verify purification.

Electrophoretic mobility shift assay. DNA fragments were designed as outlined in

Table S3, to encompass the entire region (322-bp) or sections of the ectoine biosynthesis

regulatory region (125, 137, 106-bp). The concentration of purified protein (LeuO-His or

NhaR-His) was determined using the Bradford Assay, and LeuO (0 to 2.175 uM) or

NhaR (0 to 2.175 uM) was incubated for 20 minutes with 30 ng of each DNA fragment

in a binding buffer (10 mM Tris [pH 7.4 at 4°C], 150 mM KCl, 0.1 mM dithiothreitol, 0.1

mM EDTA [pH 8.0], 5% polyethylene glycol). The reactions were then loaded (10 μL) on

a 6% native acrylamide gel (pre-run 200V for 2 hours at 4°C) and run for 2 hours at 4°C

(200V) in 0.5X TBE running buffer. To visualize, gels were stained in an ethidium

bromide bath for 15 minutes.

Mutant strain construction. In-frame deletion mutants of *nhaR* (VP0527), *torR*

(VP1032), and *hns* (VP1133), were designed as previously described via allelic exchange

(17). A truncated region of the *nhaR* (15-bp of the 891-bp), *torR* (54-bp of the 714-bp),

and *hns* (33-bp of the 408-bp) genes was generated using the primers in Table S3. The

truncated products were then ligated with the suicide vector pDS132 using the Gibson

assembly protocol and transformed into *E. coli* Dh5α (77, 78). The resulting plasmids

pDSΔ*nhaR*, pDSΔ*torR*, and pDSΔ*hns* were purified and transformed into *E. coli* β2155

λ*pir* (diaminopimelic acid auxotroph), followed by conjugation and homologous

recombination into the *V. parahaemolyticus* RIMD2210633 genome. Single crossover of

the plasmids into the genome were selected for by plating onto chloramphenicol (CM)

and screened via PCR for a truncated allele. To induce a double-crossover event, the

single-cross strain was grown overnight in the absence of CM, leaving behind either the

truncated allele or the wild-type allele. The cultures were spread on 10% sucrose plates,

and healthy colonies were screened for double-crossover and gene truncation; colonies

that contain the plasmid will appear soupy on the plate due to the presence of the *sacB*

selectable marker. In-frame deletions were confirmed with sequencing.

An in-frame deletion of *ompR* (VP0154) was created in *V. parahaemolyticus* RIMD2210633

through splicing by overlap extension (SOE) PCR and homologous recombination (79).

Primers were designed to create a 692-bp truncated allele of *ompR* using the primer pair

listed in Table S3. This truncated product was ligated into the cloning vector pJET1.2

with T4 DNA ligase and transformed into *E. coli* Dh5α λ*pir*. The truncated *ompR* allele

was then excised from the pJETΔ*ompR* vector using restriction enzymes (SacI, XbaI) and

ligated into the suicide vector pDS132. The pDSΔ*ompR* vector was then transformed

into *E. coli* β2155 λ*pir* (DAP auxotroph), and the same procedure as described above

was completed. Double mutants Δ*leuO*/Δ*hns* and Δ*hns*/Δ*nhaR* were generated by

conjugating *E. coli* β2155 λ*pir* + pDS*hns* with *V. parahaemolyticus* Δ*leuO* and β2155 λ*pir* +

pDS*nhaR* with *V. parahaemolyticus* Δ*hns.* The same procedure as described above was

carried out to achieve these double deletions. To confirm that the phenotypes observed

in the Δ*leuO*, Δ*hns* and Δ*nhaR* mutants were not due to secondary mutations within the

genome, the strains were complemented with functional copies of the gene. The coding

regions were amplified from *V. parahaemolyticus* RIMD2210633 genome and cloned into

the expression vector pBAD33. Then transformation into *E. coli* β2155 was completed

for both vectors. The expression vectors, pBAD*leuO*, pBAD*hns*, pBAD*nhaR* were then

conjugated into the Δ*leuO*, Δ*hns* and Δ*nhaR* strains.

Transcriptional GFP-reporter assay. The reporter construct pRUP$_{ectA}$-*gfp* was created

previously using the pRU1064 vector, which contains a promoterless *gfp* cassette, as

well as Tet and Amp resistance genes (17, 80). Using *E. coli* β2155 λ*pir* containing

pRUP$_{ectA}$-*gfp,* the reporter plasmid was conjugated into each of the *V. parahaemolyticus*

mutant strains Δ*leuO*, Δ*nhaR*, Δ*torR*, Δ*ompR*, Δ*hns*, Δ*leuO*/Δ*hns*, and Δ*hns*/Δ*nhaR.* Strains

were grown overnight in LB3% with tetracycline. The cultures were then washed two

times with 1XPBS, diluted 1:100 into M9G3% with tetracycline, and grown to OD ~ 0.4

or OD ~ 1.0 as indicated. The reporter expression was determined by measuring the

relative fluorescence with excitation at 385 nm and emission at 509 nm in black, clear

bottomed 96-well microplates on a Tecan Spark microplate reader with Magellan

software (Tecan Systems, Inc., San Jose, CA). Specific fluorescence was calculated by

dividing the relative fluorescence units (RFU) by the optical density of each well. At

least two biological replicates were performed for each experiment. Statistics were

completed either with a Student's T-test or ANOVA followed by Tukey-Kramer Post

Hoc test, as designated.

ACKNOWLEDGEMENTS

This research was supported by a National Science Foundation grant (award IOS-

1656688) to E.F.B. KBL and GJG were funded in part by a University of Delaware

graduate fellowship award and KBL in part by a Chemistry-Biology Interface

predoctoral training program grant: 5T32GM008550.

Figure Legends

Fig. 1. Expression of a P_{ectA}-*gfp* transcriptional fusion in wild-type (WT) *V.*

parahaemolyticus Δ*leuO*, Δ*nhaR*, Δ*ompR*, and Δ*torR* mutants. Cultures were grown

overnight in M9G 3% NaCl and relative fluorescence intensity (RFU) was measured.

Specific fluorescence was calculated by dividing RFU by OD. Mean and standard

deviation of three biological replicates are shown. Statistics were calculated using a

Student's t-test; (*, $P < 0.05$; ***, $P < 0.001$).

Fig. 2. Expression of a P_{ectA}-*gfp* transcriptional fusion in wild-type (WT) *V.*

parahaemolyticus Δ*leuO*, Δ*hns*, and Δ*leuO*/Δ*hns* mutants. Cultures were grown **A**. to OD

~0.4 or **B**. OD ~1.0 in M9G 3%NaCl and relative fluorescence intensity (RFU) was

measured. Specific fluorescence was calculated by dividing RFU by OD. Mean and

standard deviation of three biological replicates are shown. Statistics were calculated

using an ANOVA followed by Tukey-Kramer post hoc test; (***, $P < 0.001$).

Fig. 3. Expression reporter assays of P_{leuO}-*gfp* transcriptional fusion in *V.*

parahaemolyticus wild type (WT), Δ*leuO*, Δ*hns*, Δ*leuO*/Δ*hns, and* Δ*toxRS* mutants.

Cultures were grown to OD ~1.0 in M9G 3%NaCL and relative fluorescence intensity

(RFU) was measured. Specific fluorescence was calculated by dividing RFU by OD.

Mean and standard deviation of two biological replicates are shown. Statistics were

calculated using a Student's t-test; (*, $P < 0.05$; **, $P < 0.005$).

Fig. 4. Growth analysis of Δ*leuO*, Δ*hns* and Δ*leuO*/Δ*hns* mutants in M9G 6%NaCl. Cells

were grown to OD 0.5 in M9G 1%NaCl and then inoculated into M9G 6%NaCl and

growth was measured every hour for 24h at 37°C. Mean and standard deviation of two

biological replicates are shown. Also shown, complementation of Δ*leuO* with pBA*leuO*

and Δ*hns* with pBA*hns*. For all growth curves the area under the curve (AUC) was

calculated for each strain and a Student's t-test was performed (*, $P < 0.05$; **, $P < 0.005$;

***, $P < 0.001$).

Fig. 5. Expression of a P_{ectA}-*gfp* transcriptional fusion in wild type (WT) *V.*

parahaemolyticus, ΔnhaR, Δhns, and Δhns/ΔnhaR mutants. Cultures were grown **A**. to OD

~0.4 or **B**. OD ~1.0 in M9G 3%NaCl and relative fluorescence intensity (RFU) was

measured. Specific fluorescence was calculated by dividing RFU by OD. Mean and

standard deviation of three biological replicates are shown. Statistics were calculated

using an ANOVA followed by Tukey-Kramer post hoc test; (***, $P < 0.001$).

Fig. 6. Growth analysis of Δ*nhaR*, Δ*hns* and Δ*nhaR*/Δ*hns* mutants in M9G 6%NaCl. Cells

were grown to OD 0.5 in M9G 1%NaCl and then inoculated into M9G 6%NaCl and

growth was measured every hour for 24 h at 37°C. Mean and standard deviation of two

biological replicates are shown. Also shown, complementation of Δ*nhaR* with pBA*nhaR*.

For all growth curves the area under the curve (AUC) was calculated for each strain and

a Student's t-test was performed (**, $P < 0.005$).

Fig. 7. Model of ectoine biosynthesis gene expression in *V. parahaemolyticus*. Solid

arrows represent positive regulation while hammer lines represent negative regulation.

Previously it was shown that LeuO is required for OmpU expression, an important

component of the acid stress response. H-NS was also shown to negatively regulated

ToxR. In *E. coli*, NhaR is an activator of NhaA a symporter required for NaCl tolerance.

The ? indicates that this has not been shown in *V. parahaemolyticus*. Our data show that

the NhaR and LeuO regulate *ectABC-asp_ect* and that H-NS is also part of this

regulatory circuit.

References

1.   Wood JM. 1999. Osmosensing by bacteria: signals and membrane-based sensors.

Microbiol Mol Biol Rev MMBR 63:239-262.

2.   Poolman B, Glaasker E. 1998. Regulation of compatible solute accumulation in

bacteria. Mol Microbiol 29:397-407.

3.   Galinski EA. 1995. Osmoadaptation in bacteria. Adv Microb Physiol 37:272-328.

4.   Wood JM. 2011. Bacterial osmoregulation: a paradigm for the study of cellular

homeostasis. Annu Rev Microbiol 65:215-38.

5.   Wood J. 2015. Bacterial responses to osmotic challenges. J Gen Physiol 145: 381-

388.

6.   Csonka L. 1989. Physiological and genetic responses of bacteria to osmotic stress.

Microbiol Rev 53: 121-47.

7.   McLaggan D, Naprstek J, Buurman ET, Epstein W. 1994. Interdependence of K+

and glutamate accumulation during osmotic adaptation of *Escherichia coli*. J Biol

Chem 269:1911-7.

8.   da Costa M, Santos H, Galinski E. 1998. An overview of the role and diversity of

compatible solutes in Bacteria and Archaea. Ad Biochem Eng/Biotech 61: 117-53.

9.   Galinski EA, Oren A. 1991. Isolation and structure determination of a novel

compatible solute from the moderately halophilic purple sulfur bacterium

Ectothiorhodospira marismortui. Eur J Biochem 198:593-8.

10.   Galinski EA, Pfeiffer HP, Trüper HG. 1985. 1,4,5,6-Tetrahydro-2-methyl-4-

pyrimidinecarboxylic acid. A novel cyclic amino acid from halophilic

phototrophic bacteria of the genus Ectothiorhodospira. Eur J Biochem 149:135-9.

11.   Kempf B, Bremer E. 1998. Uptake and synthesis of compatible solutes as

microbial stress responses to high-osmolality environments. Arch Microbiol 170:

319-30.

12.   Severin J, Wohlfarth A, Galinski EA 1992. The predominant role of recently

discovered tetrahydropyrimidines for the osmoadaptation of halophilic

eubacteria. Microbiology 138:1629-1638.

13.   Czech L, Hermann L, Stöveken N, Richter AA, Höppner A, Smits SHJ, Heider J,

Bremer E. 2018. Role of the extremolytes ectoine and hydroxyectoine as stress

protectants and nutrients: Genetics, phylogenomics, biochemistry, and structural

Analysis. Genes (Basel) 9: 177.

14.   Hermann L, Mais CN, Czech L, Smits SHJ, Bange G, Bremer E. 2020. The ups and

downs of ectoine: structural enzymology of a major microbial stress protectant

and versatile nutrient. Biol Chem 401:1443-1468.

15.   Ongagna-Yhombi SY, Boyd EF. 2013. Biosynthesis of the osmoprotectant ectoine,

but not glycine betaine, is critical for survival of osmotically stressed *Vibrio*

parahaemolyticus cells. Appl Environ Microbiol, 79: 5038-49.

16.   Pflughoeft KJ, Kierek K, Watnick PI. 2003. Role of ectoine in *Vibrio cholerae*

osmoadaptation. Appl Environ Microbiol 69:5919-27.

17.   Gregory GJ, Morreale DP, Carpenter MR, Kalburge SS, Boyd EF. 2019. Quorum

Sensing Regulators AphA and OpaR Control Expression of the Operon

Responsible for Biosynthesis of the Compatible Solute Ectoine. Appl Environ

Microbiol 85: e01543-19.

18.   Gregory GJ, Dutta A, Parashar V, Boyd EF. 2020. Investigations of

Dimethylglycine, Glycine Betaine, and Ectoine Uptake by a Betaine-Carnitine-

Choline Transporter Family Transporter with Diverse Substrate Specificity in

Vibrio Species. J Bacteriol 202:e00314-20.

19.   Naughton LM, Blumerman SL, Carlberg M, Boyd EF. 2009. Osmoadaptation

among Vibrio species and unique genomic features and physiological responses

of *Vibrio parahaemolyticus*. Appl Environ Microbiol 75:2802-10.

20.   Graf R, Anzali S, Buenger J, Pfluecker F, Driller H. 2008. The multifunctional role

of ectoine as a natural cell protectant. Clin Dermatol 26:326-33.

21.   Buenger J, Driller H. 2004. Ectoin: an effective natural substance to prevent UVA-

induced premature photoaging. Skin Pharmacol Physiol 17:232-7.

22.   Jebbar M, Talibart R, Gloux K, Bernard T, Blanco C. 1992. Osmoprotection of

Escherichia coli by ectoine: uptake and accumulation characteristics. J Bact 174:

5210-7.

23.   Sauer T, Galinski EA. 1998. Bacterial milking: A novel bioprocess for production

559   of compatible solutes. Biotechnol Bioeng 57:306-13.

560   24.   Czech L, Poehl S, Hub P, Stöveken N, Bremer E. 2018. Tinkering with osmotically

561   controlled transcription allows enhanced production and excretion of ectoine

562   and hydroxyectoine from a microbial cell factory. Appl Environ Microbiol 84:

563   e01772-17.

564   25.   Kuhlmann AU, Bremer E. 2002. Osmotically regulated synthesis of the

565   compatible solute ectoine in *Bacillus pasteurii* and related *Bacillus* spp. Appl

566   Environ Microbiol 68:772-83.

567   26.   Reshetnikov AS, Khmelenina VN, Trotsenko YA. 2006. Characterization of the

568   ectoine biosynthesis genes of haloalkalotolerant obligate methanotroph

569   "Methylomicrobium alcaliphilum 20Z". Arch Microbiol 184:286-97.

570   27.   Louis P, Galinski EA. 1997. Characterization of genes for the biosynthesis of the

571   compatible solute ectoine from Marinococcus halophilus and osmoregulated

572   expression in *Escherichia coli*. Microbiol (Reading) 143:1141-1149.

573   28.   Mustakhimov, II, Reshetnikov AS, Fedorov DN, Khmelenina VN, Trotsenko YA.

574   2012. Role of EctR as transcriptional regulator of ectoine biosynthesis genes in

575   Methylophaga thalassica. Biochem (Mosc) 77:857-63.

576   29.   Mustakhimov, II, Reshetnikov AS, Khmelenina VN, Trotsenko YA. 2009. EctR--a

577   novel transcriptional regulator of ectoine biosynthesis genes in the

578   haloalcaliphilic methylotrophic bacterium Methylophaga alcalica. Dokl Biochem

579   Biophys 429:305-8.

580   30.   Mustakhimov, II, Reshetnikov AS, Glukhov AS, Khmelenina VN, Kalyuzhnaya

581   MG, Trotsenko YA. 2010. Identification and characterization of EctR1, a new

582   transcriptional regulator of the ectoine biosynthesis genes in the halotolerant

583   methanotroph Methylomicrobium alcaliphilum 20Z. J Bacteriol 192:410-7.

584   31.   Gregory GJ, Morreale DP, Boyd EF. 2020. CosR is a global regulator of the

585   osmotic stress response with widespread distribution among Bacteria. Appl

586   Environ Microbiol 86:e00120-20.

587   32.   Gregory GJ, Boyd EF. 2021. Stressed out: Bacterial response to high salinity using

588   compatible solute biosynthesis and uptake systems, lessons from Vibrionaceae.

589   Comput Struct Biotechnol J 19:1014-1027.

590   33.   Shikuma NJ, Davis KR, Fong JN, Yildiz FH. 2013. The transcriptional regulator,

591   CosR, controls compatible solute biosynthesis and transport, motility and biofilm

592   formation in *Vibrio cholerae*. Environ Microbiol 15:1387-99.

593   34.   Williams SG, Carmel-Harel O, Manning PA. 1998. A functional homolog of

594   *Escherichia coli* NhaR in *Vibrio cholerae*. J Bacteriol 180:762-5.

595   35.   Rahav-Manor O, Carmel O, Karpel R, Taglicht D, Glaser G, Schuldiner S, Padan

596   E. 1992. NhaR, a protein homologous to a family of bacterial regulatory proteins

(LysR), regulates nhaA, the sodium proton antiporter gene in *Escherichia coli*. J

Biol Chem 267:10433-8.

36. Król JE. 2018. Regulatory loop between the CsrA system and NhaR, a high

salt/high pH regulator. PLoS One 13:e0209554.

37. Pratt LA, Hsing W, Gibson KE, Silhavy TJ. 1996. From acids to osmZ: multiple

factors influence synthesis of the OmpF and OmpC porins in *Escherichia coli*. Mol

Microbiol 20:911-7.

38. Kenney LJ, Bauer MD, Silhavy TJ. 1995. Phosphorylation-dependent

conformational changes in OmpR, an osmoregulatory DNA-binding protein of

Escherichia coli. Proc Natl Acad Sci U S A 92:8866-70.

39. Chakraborty S, Kenney LJ. 2018. A New Role of OmpR in Acid and Osmotic

Stress in Salmonella and E. coli. Front Microbiol 9:2656.

40. Whitaker WB, Parent MA, Boyd A, Richards GP, Boyd EF. 2012. The *Vibrio*

parahaemolyticus ToxRS regulator is required for stress tolerance and colonization

in a novel orogastric streptomycin-induced adult murine model. Infect Immun

80:1834-45.

41. Gode-Potratz CJ, Chodur DM, McCarter LL. 2010. Calcium and iron regulate

swarming and type III secretion in *Vibrio parahaemolyticus*. J Bacteriol 192:6025-38.

42. Shimada T, Bridier A, Briandet R, Ishihama A. 2011. Novel roles of LeuO in

transcription regulation of E. coli genome: antagonistic interplay with the

universal silencer H-NS. Mol Microbiol 82:378-97.

43. Navarre WW, McClelland M, Libby SJ, Fang FC. 2007. Silencing of xenogeneic

DNA by H-NS-facilitation of lateral gene transfer in bacteria by a defense system

that recognizes foreign DNA. Genes Dev 21:1456-71.

44. Dorman CJ. 2004. H-NS: a universal regulator for a dynamic genome. Nat Rev

Microbiol 2:391-400.

45. Dorman CJ. 2007. H-NS, the genome sentinel. Nat Rev Microbiol 5:157-61.

46. Lucchini S, Rowley G, Goldberg MD, Hurd D, Harrison M, Hinton JC. 2006. H-

NS mediates the silencing of laterally acquired genes in bacteria. PLoS Pathog

2:e81.

47. Oshima T, Ishikawa S, Kurokawa K, Aiba H, Ogasawara N. 2006. *Escherichia coli*

histone-like protein H-NS preferentially binds to horizontally acquired DNA in

association with RNA polymerase. DNA Res 13:141-53.

48. Hernández-Lucas I, Gallego-Hernández AL, Encarnación S, Fernández-Mora M,

Martínez-Batallar AG, Salgado H, Oropeza R, Calva E. 2008. The LysR-type

transcriptional regulator LeuO controls expression of several genes in *Salmonella*

enterica serovar Typhi. J Bacteriol 190:1658-70.

49.   Shimada T, Yamamoto K, Ishihama A. 2009. Involvement of the leucine response

635   transcription factor LeuO in regulation of the genes for sulfa drug efflux. J

636   Bacteriol 191:4562-71.

637   50.   De la Cruz MA, Fernández-Mora M, Guadarrama C, Flores-Valdez MA,

638   Bustamante VH, Vázquez A, Calva E. 2007. LeuO antagonizes H-NS and StpA-

639   dependent repression in Salmonella enterica ompS1. Mol Microbiol 66:727-43.

640   51.   Hernández-Lucas I, Calva E. 2012. The coming of age of the LeuO regulator. Mol

641   Microbiol 85:1026-8.

642   52.   Ayala JC, Wang H, Benitez JA, Silva AJ. 2018. Molecular basis for the differential

643   expression of the global regulator VieA in *Vibrio cholerae* biotypes directed by H-

644   NS, LeuO and quorum sensing. Mol Microbiol 107:330-343.

645   53.   Medina-Aparicio L, Rebollar-Flores JE, Gallego-Hernández AL, Vázquez A,

646   Olvera L, Gutiérrez-Ríos RM, Calva E, Hernández-Lucas I. 2011. The

647   CRISPR/Cas immune system is an operon regulated by LeuO, H-NS, and

648   leucine-responsive regulatory protein in *Salmonella enterica* serovar Typhi. J

649   Bacteriol 193:2396-407.

650   54.   Ante VM, Bina XR, Howard MF, Sayeed S, Taylor DL, Bina JE. 2015. *Vibrio*

651   *cholerae leuO* transcription is positively regulated by ToxR and contributes to Bile

652   resistance. J Bacteriol 197:3499-510.

55.   Bina XR, Howard MF, Ante VM, Bina JE. 2016. *Vibrio cholerae* LeuO links the

ToxR regulon to expression of Lipid A remodeling genes. Infect Immun 84:3161-

3171.

56.   Ongagna-Yhombi SY, McDonald ND, Boyd EF. 2015. Deciphering the role of

multiple betaine-carnitine-choline transporters in the Halophile *Vibrio*

parahaemolyticus. Appl Environ Microbiol 81:351-63.

57.   Oren A. 1999. Bioenergetic aspects of halophilism. Microbiol Mol Biol Rev.

63:334-48

58.   Dover N, Higgins CF, Carmel O, Rimon A, Pinner E, Padan E. 1996. Na+-induced

transcription of *nhaA*, which encodes an Na+/H+ antiporter in *Escherichia coli*, is

positively regulated by nhaR and affected by hns. J Bacteriol 178:6508-17.

59.   Dover N, Padan E. 2001. Transcription of nhaA, the main Na(+)/H(+) antiporter

of *Escherichia coli*, is regulated by Na(+) and growth phase. J Bacteriol 183:644-53.

60.   Padan E, Gerchman Y, Rimon A, Rothman A, Dover N, Carmel-Harel O. 1999.

The molecular mechanism of regulation of the NhaA Na+/H+ antiporter of

Escherichia coli, a key transporter in the adaptation to Na+ and H+. Novartis

Found Symp 221:183-96.

61.   Toesca I, Perard C, Bouvier J, Gutierrez C, Conter A. 2001. The transcriptional

activator NhaR is responsible for the osmotic induction of osmC(p1), a promoter

672 of the stress-inducible gene osmC in *Escherichia coli*. Microbiology (Reading)

673 147:2795-2803.

674 62. Carmel O, Rahav-Manor O, Dover N, Shaanan B, Padan E. 1997. The Na+-specific

675 interaction between the LysR-type regulator, NhaR, and the *nhaA* gene encoding

676 the Na+/H+ antiporter of *Escherichia coli*. Embo j 16:5922-9.

677 63. Fang M, Majumder A, Tsai KJ, Wu HY. 2000. ppGpp-dependent *leuO* expression

678 in bacteria under stress. Biochem Biophys Res Commun 276:64-70.

679 64. Hertzberg KM, Gemmill R, Jones J, Calvo JM. 1980. Cloning of an EcoRI-

680 generated fragment of the leucine operon of *Salmonella typhimurium*. Gene 8:135-

681 52.

682 65. Dillon SC, Espinosa E, Hokamp K, Ussery DW, Casadesús J, Dorman CJ. 2012.

683 LeuO is a global regulator of gene expression in *Salmonella enterica* serovar

684 Typhimurium. Mol Microbiol 85:1072-89.

685 66. Navarre WW, Porwollik S, Wang Y, McClelland M, Rosen H, Libby SJ, Fang FC.

686 2006. Selective silencing of foreign DNA with low GC content by the H-NS

687 protein in *Salmonella*. Science 313:236-8.

688 67. Park NY, Kim IH, Wen Y, Lee KW, Lee S, Kim JA, Jung KH, Lee KH, Kim KS.

689 2019. Multi-Factor regulation of the master modulator LeuO for the Cyclic-(Phe-

690 Pro) signaling pathway in *Vibrio vulnificus*. Sci Rep 9:20135.

68.   Park NY, Lee KW, Kim KS. 2020. H-NS Silences Gene Expression of LeuO, the

Master Regulator of the Cyclic(Phe-Pro)-dependent Signal Pathway, in *Vibrio*

vulnificus. J Microbiol Biotechnol 30:830-838.

69.   Makino K, Oshima K, Kurokawa K, Yokoyama K, Uda T, Tagomori K, Iijima Y,

Najima M, Nakano M, Yamashita A, Kubota Y, Kimura S, Yasunaga T, Honda T,

Shinagawa H, Hattori M, Iida T. 2003. Genome sequence of *Vibrio*

parahaemolyticus: a pathogenic mechanism distinct from that of V cholerae. Lancet

361:743-9.

70.   Whitaker WB, Parent MA, Naughton LM, Richards GP, Blumerman SL, Boyd EF.

2010. Modulation of Responses of *Vibrio parahaemolyticus* O3:K6 to pH and

Temperature Stresses by Growth at Different Salt Concentrations. Appl Environ

Microbiol, 76: 4720-9.

71.   Dehio C, Meyer M. 1997. Maintenance of broad-host-range incompatibility group

P and group Q plasmids and transposition of Tn5 in Bartonella henselae

following conjugal plasmid transfer from *Escherichia coli*. J Bacteriol 179:538-40.

72.   Jutras BL, Verma A, Stevenson B. 2012. Identification of novel DNA-binding

proteins using DNA-affinity chromatography/pull down. Curr Protoc Microbiol

Chapt 1:F.1.

73.   Chaparian RR, Olney SG, Hustmyer CM, Rowe-Magnus DA, van Kessel JC. 2016.

Integration host factor and LuxR synergistically bind DNA to coactivate quorum-

sensing genes in Vibrio harveyi. Mol Microbiol 101:823-40.

74.   Tague JG, Hong J, Kalburge SS, Boyd EF. 2022. Regulatory Small RNA Qrr2 Is

Expressed Independently of Sigma Factor-54 and Can Function as the Sole Qrr

Small RNA To Control Quorum Sensing in *Vibrio parahaemolyticus*. J Bacteriol

204:e0035021.

75.   Jutras BL, Liu Z, Brissette CA. 2010. Simultaneous isolation of Ixodidae and

bacterial (*Borrelia* spp.) genomic DNA. Curr Protoc Microbiol Chapt 1:E.2.

76.   Tague JG, Regmi A, Gregory GJ, Boyd EF. 2021. Fis connects two sensory

pathways, quorum sensing and surface sensing, to control motility in *Vibrio*

parahaemolyticus. Front Microbiol 12:669447.

77.   Gibson DG. 2011. Enzymatic assembly of overlapping DNA fragments. Methods

Enzymol 498:349-61.

78.   Philippe N, Alcaraz JP, Coursange E, Geiselmann J, Schneider D. 2004.

Improvement of pCVD442, a suicide plasmid for gene allele exchange in bacteria.

Plasmid 51:246-55.

79.   Horton RM, Hunt HD, Ho SN, Pullen JK, Pease LR. 1989. Engineering hybrid

genes without the use of restriction enzymes: gene splicing by overlap extension.

Gene 77:61-8.

80.   Karunakaran R, Mauchline TH, Hosie AHF, Poole PS. 2005. A family of promoter

730   probe vectors incorporating autofluorescent and chromogenic reporter proteins

731   for studying gene expression in Gram-negative bacteria. Microbiol (Reading)

732   151:3249-3256.

P*ectA*-*gfp* Reporter Assay

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Growth curves in M9G 6% NaCl

Fig. 6

Fig. 7