Trait salivary alpha-amylase activity levels define the conditions for

facilitation by L-DOPA of extinction consolidation

Andres, E.[1+*], Chuan-Peng, H.[1+], Gerlicher, A.M.V.[2,3,4], Meyer, B.[1,2], Tüscher, O.[1,5], Kalisch, R.[1,2]

[1]Leibniz Institute for Resilience Research (LIR), 55122 Mainz, Germany

[2]Neuroimaging Center (NIC), Focus Program Translational Neuroscience (FTN), Johannes Gutenberg University Medical Center Mainz, 55131 Mainz, Germany

[3]Department of Experimental Psychology, Helmholtz Institute, Utrecht University, 3584 CS Utrecht, The Netherlands

[4]Department of Clinical Psychology, University of Amsterdam, 1018 WS Amsterdam, The Netherlands

[5]Department of Psychiatry and Psychotherapy, Johannes Gutenberg University Medical Center, 55131 Mainz, Germany

[+]These authors contributed equally.

Even after successful extinction, conditioned fear can return. Strengthening the consolidation of the fear-inhibitory safety memory formed during extinction is one way to counteract return of fear. In this preregistered direct replication study in male participants, we confirm that spontaneous post-extinction reactivations of a neural activation pattern evoked in the ventromedial prefrontal cortex (vmPFC) during extinction predict extinction memory retrieval 24 h later. We do not confirm that L-DOPA administration after extinction enhances retrieval and that this is mediated by enhancement of the number of vmPFC reactivations. However, additional non-preregistered analyses reveal a beneficial effect of L-DOPA on extinction retrieval when controlling for the trait-like stable baseline levels of salivary alpha-amylase enzymatic activity (trait sAA) levels that participants show on the three experimental days. Further, trait sAA negatively predicts retrieval, and this effect is rescued by L-DOPA treatment. Our results suggest that individuals with high basal levels of sympathetic nervous system (SNS) activity may have poor extinction and that L-DOPA may be selectively beneficial for these individuals, which holds potential for clinical applications.

Introduction

Recognizing an event or situation as dangerous and reacting defensively when it re-occurs is a learning mechanism (fear or threat conditioning) that is fundamental for survival. However, learning when a previous threat no longer signals danger and ceasing costly defense behavior (fear or threat extinction) also has great adaptive value and has been related to a reduced risk for threat-related mental disorders such as post-traumatic stress disorder (PTSD) or anxiety disorders[1]. Also, extinction learning is the probable mechanism that underlies the exposure-based treatment of threat-related disorders[2,3].

To investigate whether fear extinction lastingly reduces CRs, a three-phase paradigm is commonly used, consisting of fear conditioning, fear extinction, and a memory test. During fear conditioning, an innocuous conditioned stimulus (CS) is repeatedly paired with an unconditioned stimulus (US). Participants start to exhibit CRs to the formerly neutral CS, and a CS-US association, or 'fear memory', is formed[4]. During extinction, participants are re-exposed to the CS in the absence of the US several times, and CRs decrease. During the test phase, participants are once more exposed to the CS, again in the absence of the US, and CRs are measured. Based on this paradigm, studies have reported that even after a complete reduction of CRs over the course of extinction, CRs often return during the test phase ("return of fear"[5]). Thus, extinction learning is not an unlearning or erasure of the original fear memory, but formation of a new CS-noUS association or 'extinction memory'[5], and the test phase effectively examines the retrieval and/or expression of the fear in competition with the extinction memory.

In exposure-based treatment, relapse after successful exposure is not uncommon, and return of fear is presumably a precursor for relapse[3]. Therefore, it would be highly desirable to develop methods to prevent the return of fear. Given that consolidation processes are crucial for long-term memory expression[6–8], reinforcing the consolidation of the extinction memory may be one promising avenue[9].

The dopaminergic system plays a crucial role in memory consolidation[10]. In the case of

extinction, memory retrieval has been shown to be worse when dopaminergic activity is

decreased after extinction training. After microinfusion of a D1 receptor antagonist into the

infra-limbic part of the medial prefrontal cortex (IL) following extinction training in rats, later

extinction memory retrieval was reduced relatively to a vehicle condition, suggesting

impaired consolidation[11]. Conversely, increasing dopaminergic activity after extinction

training results in better long-term extinction memory retrieval. So, the systemic

administration in mice of methylphenidate, a dopamine and noradrenaline reuptake inhibitor,

after extinction learning led to relatively decreased fear responses at test 24 h later[12]. Of

particular relevance for potential clinical applications, post-extinction systemic administration

of the anti-Parkinson drug L-DOPA, a precursor of dopamine that preferentially enhances

dopaminergic turnover in the frontal cortex[13], improved extinction memory retrieval both in

mice and healthy normal humans in altogether six experiments[14–16]. A negative result in one

human study was accompanied by reduced neural activity at test in brain areas related to

conditioned fear[17]. In two human experiments conducted outside the magnetic resonance

imaging (MRI) environment[18], post-extinction L-DOPA reduced CRs at test only in

participants who showed successful extinction learning; in one other non-MRI study, the L-

DOPA effect was absent[19]. Hence, L-DOPA is a likely pro-consolidation agent, although the

boundary conditions for its effectiveness still have to be established.

A strong body of research supports a causal role for the IL, the rodent homolog of the

vmPFC, in the consolidation of extinction memories[20]. In humans using functional MRI

(fMRI), Gerlicher et al. (2018)[16] reported that a neural activation pattern in the vmPFC, which

they initially observed during extinction when the US was unexpectedly omitted from CS

presentations, spontaneously re-occurred during post-extinction rest. They further observed

that the number of reactivations of this multi-voxel pattern (MVP) in the resting state

predicted extinction memory retrieval as well as vmPFC activation at the memory test 24 h

later. Importantly, the IL/vmPFC is a target of dopaminergic projections from the ventral

tegmental area (VTA), and extinction has been shown to evoke lasting dopamine release in

this brain area[21]. Extending the rodent microinfusion data[11], Gerlicher et al. (2018)[16] also

reported that post-extinction L-DOPA administration enhanced the number of spontaneous

vmPFC MVP reactivations and that this mediated the beneficial effect of L-DOPA on

extinction retrieval. No other brain area showed this relationship, suggesting a pro-

consolidation action of dopamine that depends on consolidation-related vmPFC activity.

A further intriguing aspect of this study was that all findings were specific to vmPFC MVPs

from early extinction trials. Early in extinction, the CS still elicits a high US expectation, and

the surprise (or "prediction error", PE) generated by US omission is highest[22]. It is generally

accepted that the formation of the new CS-noUS association in extinction is driven by this

PE[23–25], and extinction PEs in turn have been shown to be encoded by phasic dopamine

release in the ventral striatum[26]. It is currently unclear whether the vmPFC also receives

dopaminergic extinction PE signals, but extinction PE-correlated fMRI activity has been

observed in humans also outside the ventral striatum[27]. Taken together, the VTA-originating

dopaminergic system may tie together extinction learning, memory formation, and memory

consolidation via influences on the prefrontal cortex.

We here aimed to directly replicate the findings of Gerlicher et al. (2018)[16], as preregistered

in Chuan-Peng et al. (2018)[28]. For this, we employed the same fMRI paradigm (Fig. 1a) with

differential fear conditioning in context A (background picture) on day 1. Two geometric

symbols served as CS+ (reinforced at its offset in 50% of conditioning trials with a painful

US) and CS- (non-reinforced CS), respectively. Extinction learning in context B on day 2 was

immediately followed by oral placebo or L-DOPA administration (randomized between-

subject design) and resting-state fMRI scans 10, 45, and 90 min after the end of extinction,

to cover the early consolidation window. On day 3, we tested extinction memory retrieval in

context B. During extinction and test, no US was administered. Skin conductance responses

(SCRs) were used as CRs (Fig. 1b).

We tested three main replication hypotheses[28]: First, we expected that post-extinction L-

DOPA as compared to placebo administration would improve extinction memory retrieval at

test on day 3. That is, we expected smaller differential (CS+>CS-) SCRs for L-DOPA-

compared to placebo-treated participants (hypothesis 1). Second, we expected that the

number of spontaneous post-extinction reactivations during the post-extinction resting-state

scans of a vmPFC MVP linked with CS+ offsets in the first five extinction trials (first third of

extinction) would predict extinction memory retrieval during test on day 3 in both placebo-

and L-DOPA-treated participants (hypothesis 2). We were particularly interested in the

resting-state scan conducted 45 min after extinction, as this is where Gerlicher et al. (2018)[16]

had observed a predictive relationship to retrieval, but also allowed ourselves to test the

scans conducted 10 and 90 min after extinction, as a precise timing of, or very restricted

time window for, the effect was considered unlikely. Third, we expected that L-DOPA would

significantly increase the number of vmPFC reactivations (hypothesis 3).

To determine the optimal replication sample size, we combined different approaches[28]. We

conducted a power analysis based on the critical effect size from Gerlicher et al. (2018)[16] for

the most important hypothesis 1 (see Methods for details). This led to a required sample size

of N=22 (i.e., eleven participants per treatment group). However, considering that effects in

discovery samples may overestimate the true effect size, Simonsohn (2015)[29] suggested

that replication studies should have a sample size that is at least 2.5 times greater than that

of the discovery study (N=40 in Gerlicher et al., 2018[16]), leading to a required sample size of

N=100 (50 per group). This approach would mean that we would need a sample size 4.5

times bigger than the sample size estimated based on the critical effect size in the discovery

study, which we considered exaggerated. To balance the feasibility of a three-day

pharmacological fMRI paradigm and the requirement to limit the number of participants

exposed to study-associated burden[30] with the requirement of sufficient statistical power, we

settled on a sample size of N=70 (35 per group). With an estimated drop-out rate of 10%,

this means we could expect to achieve a final sample size of N=63, which is 1.5 times

greater than in the discovery study and nearly 3 times greater than based on power

calculation. The responsible medical ethics committee approved this approach.

We furthermore extended the paradigm to clarify two remaining questions (see

preregistration[28]). First, to assess whether L-DOPA not only protects against the

spontaneous recovery of extinguished fear after a mere passage of time but also against the

renewal of fear gated by a context change[5], we added a test in a new context C (non-

extinction context, a new background picture) on day 3 (secondary research question 1; Fig.

1a). Protection against renewal would suggest that L-DOPA makes the extinction memory

context independent, as indicated by the results of Haaker et al. (2013)[14]. Second, we

collected saliva samples throughout all experimental days to measure enzymatic activity

levels of salivary alpha-amylase (sAA) and the concentration of salivary cortisol (sCORT) as

markers of the activity of the sympathetic nervous system (SNS) and of the hypothalamus-

pituitary-adrenal gland (HPA) axis, respectively[31,32] (Fig. 1c). The activity of these systems,

linked to arousal and stress, have been suggested to affect consolidation processes in

general and extinction consolidation in particular[33–35]. We had previously observed L-DOPA

main effects in two out of three MRI experiments[14,16,17], while two purely behavioral

experiments showed only conditional L-DOPA effects, moderated by prior extinction

success[18], and one other behavioral study showed no effect[19]. This might indicate a potential

influence of the MRI environment on the effect of L-DOPA[19]. Previous studies have reported

relatively increased sAA[36,37] and sCORT levels[36,38] in the scanner environment. It is

conceivable that higher arousal or stress might impair extinction[33,39], leaving more room for

an augmenting effect of L-DOPA; alternatively, dopamine might positively interact with

arousal- or stress-related neural activity in its effect on extinction consolidation[13,33]. We

therefore wondered if peak sAA or sCORT levels on the extinction day 2 would affect

extinction retrieval on day 3 and interact with treatment (placebo vs. L-DOPA) (secondary

research question 2).

sAA and sCORT have been observed to increase in response to arousing and stressful

stimuli, such as being scanned[36,38,40] or receiving painful stimulation[41–43], but they have also

been reported to exhibit pronounced stability across time when repeatedly measured in

baseline states of rest[44–46]. For sAA in particular, variance across a day or across days is

better explained by between- than within-subject differences[44,47]. Out et al. (2011) have also

demonstrated that baseline sAA is substantially heritable[48]. Hence, repeated baseline sAA

assessments (as well as repeated sCORT measures) may also be employed as trait-like

individual-differences markers (trait sAA, trait sCORT). These individual differences may

partly determine individual differences in extinction and/or in the effect of L-DOPA on

extinction memory consolidation. At the request of one reviewer, we therefore also tested

whether sAA and sCORT baseline levels on the three experimental days interact with

extinction learning, extinction retrieval, and drug treatment (non-preregistered secondary

research question 3).

In this work, we show that the number of spontaneous reactivations in a resting-state scan

90 min after extinction of an MVP elicited specifically in the vmPFC during early extinction

learning when the US is omitted at CS+ offsets positively predicts extinction memory

retrieval 24 h later. We also find that trait sAA as well as peak sAA at the time of extinction

negatively predict retrieval. Controlling for trait and state sAA reveals a beneficial L-DOPA

main effect and a trait sAA by group interaction, such that the negative effect of trait sAA is

rescued by L-DOPA treatment. Our findings suggest that boosting dopaminergic activity

promotes the consolidation of extinction in individuals with elevated basal SNS activity.

Results

Effect of L-DOPA administration post extinction on extinction memory retrieval

First, we tested whether post-extinction L-DOPA compared to placebo administration on day

2 enhances extinction memory retrieval on day 3 (preregistered hypothesis 1). We compared

the extent of spontaneous recovery of CRs in the extinction context B between the two

treatment groups (see Fig. 1b). In addition, we asked whether L-DOPA administration also

reduces contextual renewal of fear in a new context C on the test day (preregistered

secondary research question 1).

Fear acquisition on day 1 was equally successful in both groups, as indicated by a significant

effect of stimulus (CS+>CS- SCRs averaged across last 20% trials: $F_{1,64}=70.52$, p<.001,

generalized $\eta^2=.23$) in the absence of group (placebo/L-DOPA) and stimulus by group

effects (ps>.33, n=66 participants with sufficient SCR data quality; Fig. 1b; see Methods and

Supplementary Table 2 for details on exclusions and resulting final sample sizes per

analysis). Fear was retrieved at the beginning of extinction on day 2 in both groups (start-

fear: averaged CRs across first 20% trials; stimulus: $F_{1,49}=36.23$, p<.001, generalized

$\eta^2=.16$; group and stimulus*group: ps>.14; n=51). By the end of extinction on day 2 (end-

fear: SCRs across last 20% trials), an unforeseen group difference emerged (stimulus:

$F_{1,49}=8.81$, p=.005, generalized $\eta^2=.04$; group: $F_{1,49}=2.07$, p=.16; stimulus*group: $F_{1,49}=4.92$,

p=.02, generalized $\eta^2=.02$, n=51), characterized by higher CRs towards the CS+ in the

placebo group (two-sample t-test: $t_{49}=2.09$, p=.04, CI 95% [0.006 0.31]).

Contrary to our predictions, administration of L-DOPA on day 2 did not result in a significant

reduction of spontaneous recovery on day 3 (averaged CRs across all trials; stimulus:

$F_{1,53}=55.30$, p<.001, generalized $\eta^2=.20$; group: $F_{1,53}=.91$, p=.34; stimulus*group: $F_{1,53}=0.02$,

p=.89; n=55). Also, contextual renewal of fear was not reduced (stimulus: $F_{1,53}=23.66$,

p<.001, generalized $\eta^2=.07$; group: $F_{1,53}=0.91$, p=.35, stimulus*group: $F_{1,53}=0.17$, p=.68).

Adjusting for the observed group differences at the end of extinction did not change the

result (multiple regression including end-fear at extinction as well as fear acquisition and

start-fear at extinction as covariates, non-preregistered analysis: spontaneous recovery:

$\beta_{group}=.02$, SE=.05, $t_{40}=0.51$, p=.61; renewal: $\beta_{group}=.02$, SE=.04, $t_{40}=0.55$, p=.58; n=45),

although it showed in both cases that end-fear at extinction significantly and positively

predicted CRs at test (spontaneous recovery: $\beta_{end-fear}$=.25, SE=.10, t_{40}=2.52, p=.02; renewal:

$\beta_{end-fear}$=.27, SE=.09, t_{40}=3.06, p=.004; Fig. 2), as also observed in Gerlicher et al. (2018)[16].

Since preregistration, three studies have reported that better extinction (lesser end-fear) is

associated with better extinction retrieval in L-DOPA-treated, but not placebo-treated,

participants, suggesting the L-DOPA effect may under some circumstances be restricted to

successful extinguishers[18,19]. However, in an additional non-preregistered interaction

analysis, group did not interact with end-fear of extinction in predicting CRs at spontaneous

recovery or renewal (all ps>.40). Instead, comparable relationships between end-fear at

extinction and CRs at the spontaneous recovery and renewal tests were found in both

groups (simple slope analysis: spontaneous recovery: $\beta_{end-fear/placebo}$=.24, SE=.13, t_{39}=1.86,

p=.07; $\beta_{end-fear/L-DOPA}$=.26, SE=.16, t_{39}=1.65, p=.11; renewal: $\beta_{end-fear/placebo}$=.32, SE=.11,

t_{39}=2.82, p=.008; $\beta_{endf-ear/L-DOPA}$=.19, SE=.14, t_{39}=1.35, p=.19).

Together, these findings do not confirm an effect of L-DOPA on extinction memory

consolidation but show a relationship between extinction success and extinction memory

retrieval.

Relationship between vmPFC pattern reactivations post extinction and extinction

memory retrieval

Next, we tested whether spontaneous post-extinction reactivations of a multi-voxel activity

pattern (MVP) in the vmPFC, elicited by the offset of the first five CS+ trials early in

extinction, on day 2 predict extinction memory retrieval at test on day 3 (preregistered

hypothesis 2). We observed the predicted negative correlation between the number of

vmPFC reactivations and CRs in the spontaneous recovery test for the resting-state scan

conducted 90 min after extinction (β=-.10, SE=.03, t_{41}=3.17, p=.003, Bonferroni threshold for

testing four time points: .013, n=46; Fig. 3), but not for the scans conducted before as well

as 10 and 45 min after extinction (ps>.27). Like in the discovery study, where the effect was

observed at 45 min, the relationship did not differ between groups (interaction number of

reactivations at 90 min*group: p=.66, simple slope analysis: $\beta_{placebo}$=−.09, p=.04, β_{L-}

$_{DOPA}$=−.12, p=.02). The relationship did not extend to renewal (non-preregistered analysis,

ps>.77 for all time points).

Additional analyses confirmed the anatomical specificity of the relationship at 90 min to the

vmPFC (Supplementary Fig. 2a-h). Also at 45 min, no region emerged as showing a

relationship (Supplementary Fig. 3 a-i). The relationship at 90 min in the vmPFC was also

specific to early CS+ offsets, as no effect could be detected for early CS- offsets at 90 min in

this region (non-preregistered analysis, ps>.55).

Together, these results establish an important role for spontaneous post-extinction

reactivations of an extinction-related activity pattern in the vmPFC in the consolidation of

long-term extinction memories.

The number of vmPFC reactivations at any of the post-extinction time points did not

detectably predict differential BOLD responses (contrast CS+ > CS-) on day 3. Accordingly,

the preregistered analysis of the effects of vmPFC reactivations on task-modulated

functional connectivity (contrast CS+ > CS-) on this day could not be conducted.

Effect of L-DOPA administration post extinction on vmPFC reactivations

Next, we tested whether L-DOPA compared to placebo increases the number of

spontaneous post-extinction vmPFC reactivations 10, 45, and 90 min after extinction

(preregistered hypothesis 3). In congruence with an absent effect of L-DOPA on extinction

retrieval, there was no observable effect of L-DOPA on the number of vmPFC reactivations

during either of the resting-state scans (repeated-measures ANOVA: group: p=.36,

time*group: p=.70, n=46; Fig. 4).

Together, these findings do not confirm an effect of L-DOPA on post-extinction vmPFC

reactivations of an extinction-related activity pattern.

Relationship between salivary alpha-amylase and cortisol with extinction memory

retrieval, and interactions with L-DOPA administration post extinction

Our preregistered secondary research question 2 concerned whether individual differences in sAA and sCORT responses to the experimental manipulation on the extinction day 2, involving scanning and an extinction learning session, had an influence on extinction memory retrieval and might moderate the L-DOPA effect. To this aim we collected sAA and sCORT before participants entered the scanner and received extinction training ('baseline', time 0 min in Fig. 1c) and at several time points thereafter on day 2. Time courses of both markers on day 2 exhibited linear decreases (effects of time: sAA: $F_{6,324}=9.50$, p<.001, n=56; sCORT: $F_{6,346}=27.45$, p<.001, n=60) in the absence of group influences (group and group*time: ps>.12), but did not show the expected peaks at a post-extinction time point relative to baseline (Fig. 5). Maximum levels of sAA and sCORT were observed at baseline in 71.7% and 60.3% of participants, respectively, and at the first post-extinction time point (approx. 25 min in Fig. 1c) in the remaining participants. This suggests the manipulation did not robustly induce arousal and/or stress.

One reviewer pointed out that trait-like sAA activity levels, as apparent from the baseline measurements of sAA on the three experimental days, may also potentially explain individual differences in extinction retrieval and in L-DOPA effects (non-preregistered secondary research question 3). Supporting the notion of trait-like stability in sAA, we observed highly significant medium- and large-sized correlations in baseline sAA between the three days (day 1-2: R=.46, day 1-3: R=.61, day 2-3: R=.60; all ps<.001; Supplementary Fig. 4a-c). Test-retest reliability, assessed in a two-way mixed-effects model with participants as random and rater/time as fixed factors, was good (intraclass correlation [ICC]=.701, p<.0001). This allowed us to use average baseline sAA values from the three days as signifying an individual's 'trait sAA'. To not neglect potential influences of a current state of the SNS at the time of extinction, we also defined a 'state sAA' level as the deviation of an individual's maximum value on day 2 from the trait value (thus avoiding collinearity), and

entered both into a prediction model for extinction memory retrieval along with factors group

and the trait by group and state by group interactions. Note that trait and state values did not

differ between groups (two-sample t-tests: trait sAA: t_{57}=0.21, p=.84; state sAA: t_{57}=0.88,

p=.38; n=59).

Both trait and state sAA positively predicted CRs at spontaneous recovery (trait sAA:

β_{sAA}=.0008, SE=.0002, t_{37}=2.60, p=.013; state sAA: β_{sAA}=.0008, SE=.0003, t_{37}=2.80, p=.008;

n=43; Fig. 6). That is, higher trait-like or state levels of SNS activity appeared to be

associated with relatively impaired extinction learning or consolidation (rather than with

improved consolidation, as predicted for peak sAA responses on day 2 by us[28]). More

intriguingly, when controlling for trait and state sAA, the analysis also showed a positive

effect of L-DOPA (β_{group}=.17, SE=.08, t_{37}=2.04, p=.048) and a trait sAA by group interaction

($\beta_{sAA*group}$=-.0009, SE=.0004, t_{37}=2.13, p=.04). There was a non-significant trend-wise state

sAA by group interaction ($\beta_{sAA*group}$=-.0008, SE=.0004, t_{37}=1.75, p=.089). See Fig. 6. This

means that controlling for individual levels of sAA activity revealed a beneficial main effect of

L-DOPA on extinction retrieval, and this L-DOPA effect was particularly pronounced at high

levels of trait sAA. Whereas trait sAA positively predicted CRs at spontaneous recovery in

the placebo group (Fig. 6a), a simple slope analysis showed an inversion of the sAA-to-

spontaneous recovery relationship in the L-DOPA group ($\beta_{placebo}$=.0006, p=.012; β_{L-DOPA}

=−.0003, p=.38). Hence, the analysis of sAA effects in our study suggests that trait SNS

activity is an important moderator of dopaminergic extinction effects, as previously

proposed[18,19]. High basal SNS activity may thus be a necessary condition for extinction

consolidation augmentation by L-DOPA. The moderating influence of state sAA remains

open.

A further non-preregistered exploratory analysis revealed that trait (but not state) sAA was

significantly associated with high extinction end-fear, that is, poor extinction success (trait

sAA: β_{sAA}=.002, SE=.0008, t_{33}=2.53, p=.02; state sAA: β_{sAA}=-.001, SE=.0007, t_{33}=1.63,

p=.11; group: β_{group}=-.04, SE=.15, t_{33}=0.24, p=1.00; n=39; Supplementary Fig. 5). These

results may support generally impairing effects of trait-like SNS activity on extinction, as

suggested by animal studies[39], which in turn might be carried over to spontaneous recovery

(see results above). There were no effects of trait or state sAA on extinction retrieval in the

renewal test (non-preregistered exploratory analysis, not shown).

For baseline sCORT, we did not observe temporal stability comparable to baseline sAA (day

1-2: R=.53, day 1-3: R=.32, day 2-3: R=.29; all ps>.11; ICC=.03, p=.43; Supplementary Fig.

4d-f). We nevertheless for completeness also tested the effects of trait and state sCORT on

extinction retrieval and L-DOPA action as well on extinction success. Trait and state sCORT

did not differ between groups (trait sCORT: t_{61}=0.87, p=.39; state sCORT: t_{61}=0.06, p=.96;

n=63). There were no significant effects.

Relationship between salivary alpha-amylase with vmPFC reactivations during memory

consolidation, and interaction with L-DOPA administration post extinction

As controlling for trait and state sAA in the present study had revealed the hypothesized

beneficial effect of L-DOPA on extinction memory retrieval (which had not been found when

testing for an L-DOPA effect alone), we generated the new hypothesis that controlling for

sAA might also reveal a beneficial effect of L-DOPA on vmPFC reactivations 90 min after

extinction. This non-preregistered exploratory analysis found significant state sAA ($β_{sAA}$=-

.003, SE=.002, t_{43}=2.04, p=.047, n=49) and state sAA by group effects ($β_{sAA*group}$=.005,

SE=.003, t_{43}=2.04, p=.048). See Fig. 7. There were non-significant trend-wise effects of trait

sAA ($β_{sAA}$=-.002, SE=.001, t_{43}=1.86, p=.07), group ($β_{group}$=.83, SE=.002, t_{43}=1.88, p=.07), and

trait sAA by group ($β_{sAA*group}$=.004, SE=.002, t_{43}=1.79, p=.08). This suggests that, unlike in

the case of extinction memory retrieval, the effect of state sAA may be relatively more

important when investigating consolidation-related vmPFC activity.

Simple slope analysis showed a tendency for a negative influence of state sAA on vmPFC

reactivations in the placebo group ($β_{placebo}$=-.003, p=.05), in line with the above-reported

negative association between state sAA and extinction memory retrieval and with the

negative main effect of state sAA found in this analysis. This negative state sAA-to-vmPFC

reactivations relationship was absent, and tended to be inversed, in the L-DOPA group ($\beta_{L\text{-}}$

$_{DOPA}$ =.002, p=.34; Fig. 7b). That is, higher apparent current SNS activity was associated

with fewer reactivations in the placebo, but not in the L-DOPA group. This latter result

resembles in its pattern the result shown in Fig. 6a of an interaction between trait sAA and L-

DOPA in their influence on spontaneous recovery.

There were no significant effects when analyzing reactivations at 45 min after extinction.

Discussion

The present study aimed to replicate the main study results of Gerlicher et al. (2018)[16]. We

tested whether post-extinction L-DOPA as compared to placebo administration would

improve extinction memory retrieval in healthy normal male participants, as visible from

reduced spontaneous recovery during an extinction memory test on day 3 of our

experimental paradigm (hypothesis 1); whether the number of spontaneous reactivations of

an extinction learning-related multi-voxel pattern (MVP) observed in the vmPFC in post-

extinction resting-state scans would be predictive for extinction memory retrieval across

treatment groups (hypothesis 2); and whether L-DOPA administration would relatively

increase post-extinction vmPFC reactivations (hypothesis 3). The latter was done with an

eye on potentially finding the mediation of the L-DOPA effect on retrieval by the increased

number of vmPFC reactivations that was reported by Gerlicher et al. (2018)[16]. Two

preregistered extensions further aimed at clarifying whether L-DOPA administration protects

against the renewal of fear, tested on day 3 by presenting the CSs in a non-extinction

context C (secondary research question 1), and whether peak sAA or sCORT responses to

the experimental manipulation (scanning, extinction) on the extinction day 2 predict

extinction memory retrieval on day 3 and interact with drug treatment in doing so (secondary

research question 2). Upon request of a reviewer, we also tested whether trait-like baseline

activity levels of sAA (and for completeness, trait-like baseline concentrations of sCORT)

showed predictive effects (non-preregistered secondary research question 3).

We confirmed one of our three main hypotheses, namely that the number of spontaneous

post-extinction reactivations of a vmPFC activity pattern observed during CS+ offsets in

early extinction is predictive for extinction memory retrieval (hypothesis 2). Specifically, this

was observed for the resting-state scan conducted 90 min after extinction, though not at 45

min as in the discovery study. We could not replicate the facilitating effect of L-DOPA on

extinction retrieval (hypothesis 1) and on the number of vmPFC reactivations (hypothesis 3).

Examining whether L-DOPA could make extinction memories context independent, we also

did not find an advantage of L-DOPA administration for extinction memory retrieval in a

renewal test (secondary research question 1). Answering secondary research question 2

was complicated by absence of a clear response in sAA and sCORT measures to the

experimental manipulation. Contrary to our expectations, being placed in a scanner,

experiencing scanner noise, and being subjected to an extinction training session did not

lead to detectable increases at the group level in these state markers of SNS and HPA

activity, which would have been indicative of an acute arousal and stress response. We

therefore focused our analysis on secondary research question 3, centered around potential

predictive effects of trait-like sAA and sCORT levels. Trait sAA and sCORT were

operationalized as average baseline values from all three experimental days, while potential

influences of current activity levels were assessed by entering the deviation of individuals'

maximum sAA and sCORT levels on day 2 from their respective trait levels into a common

prediction model with treatment group and the trait by group and state by group interaction

terms. These analyses found that trait and state sAA were negatively associated with

extinction memory retrieval. Further, trait sAA (but not state sAA) levels interacted with drug

treatment in that the generally negative association of trait sAA with extinction retrieval

(higher spontaneous recovery) was abolished in the L-DOPA group. Unexpectedly, the

analysis also showed a beneficial main effect of L-DOPA on extinction retrieval. Hence,

controlling for trait and state sAA levels revealed an L-DOPA effect that was not observable

in the uncontrolled analysis conducted to test hypothesis 1.

We complemented these main analyses with an exploratory analysis of trait and state sAA

on extinction learning success on day 2, finding that extinction learning is poor in individuals

with high trait sAA, and with an exploratory analysis of L-DOPA effects on vmPFC

reactivations 90 min after extinction that took into consideration trait and state sAA. Here,

state sAA negatively predicted reactivations and state sAA and L-DOPA treatment interacted

in the sense that state sAA was negatively related with the number of vmPFC reactivations

in the placebo group, but tended to be positively related in the L-DOPA group.

Our main replication finding is that spontaneous post-extinction reactivations of a CS+ offset-

related vmPFC activity pattern positively associate with later extinction memory retrieval.

This result is in line with previous animal research, showing that spontaneous IL activity after

extinction learning is crucial for consolidation and predictive for memory retrieval[49].

Additionally, stimulus-specific fMRI multi-voxel reactivation patterns in other learning

domains and brain regions also have been reported to be associated with later memory

performance outcomes[50]. The reactivation pattern in our work specifically recapitulates CS+

offset-related activity in the vmPFC during *early* extinction. The CS+ offset is the time point

in a trial at which participants have learned to expect a US during prior conditioning, and the

omission of the US that occurs during the extinction phase is most unexpected at the

beginning of extinction, while less of a surprise at later extinction trials. This suggests that

the vmPFC may reactivate a prediction error (PE) signal, which learning theory considers to

be the key teaching signal in associative learning[22,51]. In extinction specifically, the PE may

not so much lead to an update (reduction) of the aversive value of the CS, but rather to the

build-up of a new, appetitive safety memory (a CS–noUS association, or extinction memory)

that antagonistically inhibits the CS-US association (or fear memory)[52,53]. On this basis, one

can postulate that vmPFC reactivations of important extinction learning events after

extinction provide the link between extinction learning and extinction memory retrieval, by

facilitating the storage and consolidation of a safety association.

One open question is why we find a predictive effect of vmPFC reactivations for memory

retrieval at 90 min after extinction and not at 45 min, as in the discovery study. Animal

studies indicate that the time window for consolidation processes can be broad, spanning

hours and even extending into sleep, depending on which molecular or network mechanisms

are involved[54]. Further, while molecular consolidation processes are bound to specific

timelines[55], they may still show substantial interindividual differences, for instance, due to

differences in the speed or success of learning, in the individual make-up of the molecular

systems, or in the current state of the system in which a consolidation process occurs (e.g.,

in stress, vigilance, arousal etc.). Such differences have not been explored in human work

but may lead to substantial heterogeneity in consolidation dynamics across studies. It is also

worth noting that oral administration of a drug can lead to substantial pharmacokinetic

variability.

Previous studies from our group using a cue conditioning and extinction paradigm have

reported pronounced L-DOPA main effects when experiments were conducted in MRI[14,16],

but only conditional L-DOPA effects (in participants with successful extinction learning) when

experiments were purely behavioral (two experiments[18]). Because there is evidence that the

scanner environment is frequently perceived by participants as arousing and stressful[36–38],

this pattern of results prompted us to postulate that L-DOPA might be more effective in

promoting extinction memory consolidation under conditions of high arousal and/or stress,

as should be apparent from high sAA or sCORT responses on the extinction day[28].

Our analysis of sAA and sCORT time courses found no evidence of such responses at the

group level. The same was observed on the other experimental days (conditioning, retrieval

test; see Fig. 6). The absence of a clear cortisol response is not very surprising, given that

cortisol is typically not increased by conditioning experiments in humans[56]. The absence of

sAA responses could potentially be attributed to a habituation effect, whereby the initial peak

of sAA activity might not have been captured, since the time participants spent in the

scanner each day before the second saliva sampling was at least 20 minutes. Another part

of the explanation may lie in the limited reliability of sAA as a reactivity marker[57].

Unexpectedly, however, we observed a pronounced predictive effect of trait sAA on

extinction memory retrieval and on the influence of L-DOPA treatment on this outcome. sAA

activity levels repeatedly measured in states of rest show high temporal stability, higher

between- than within-subject differences, and substantial heritability[44,45,47,48]. To establish

trait sAA levels, it is recommended to assess sAA on two or three days, whereby the number

of days is more important than the number of measurements per day[44]. This here allowed us

to use average baseline sAA values from three experimental days to obtain a trait-like

individual-differences marker.

Baseline, or resting, sAA is lowered by systemic administration of the beta-adrenergic

receptor blocker propranolol[58,59] and raised by the alpha2-adrenergic antagonist yohimbine,

which also increases plasma noradrenaline levels[60], indicating a direct link between baseline

sAA and noradrenergic activity and, by extension, the basal activity of the sympathetic

nervous system (SNS). The suitability of baseline sAA as a marker for basal SNS activity is

further supported by a systematic positive link between baseline sAA and age[61-63], which is

in line with the established finding that older adults have higher chronic SNS activation[64],

and by the observation that morning sAA levels are associated with hypertension[63].

The same study showed a positive relationship between morning sAA and Mild Cognitive

Impairment, while another study showed a relationship between morning sAA and cognitive

impairments in younger adults[65]. Baseline sAA has also been linked with poor pattern

separation[66]. These data indicate a negative relationship between baseline sAA and

cognition and suggest that this measure may be usable as a marker for neural function

beyond basal SNS activity. The apparent link between baseline sAA and cognition may be

explained via their common relationship with general noradrenergic activity, based on the

finding that higher noradrenaline levels in cerebrospinal fluid are associated with worse

cognition[67].

Taken together, the literature on baseline sAA indicates that trait-like sAA values from

baseline states reflect basal SNS activity and index poorer cognitive performance. This in

turn makes it a plausible extension of the current knowledge that the trait-like individual

differences in baseline sAA activity observed in our study are a predictor of individual

differences in extinction memory performance.

Trait sAA in our data predicted poorer extinction memory retrieval, poorer extinction success,

and (at trend level) fewer post-extinction vmPFC reactivations.

The negative association with extinction retrieval was abolished by L-DOPA administration

after extinction. In a similar vein, there was a trend-level interaction between trait sAA and

treatment group in the analysis predicting vmPFC reactivations. These findings suggest that

L-DOPA is more effective in participants with high basal SNS activity. From a clinical point of

view, this points to an important boundary condition for a potential application of L-DOPA as

a pharmacological augmentation of exposure-based therapy[9]. L-DOPA-based augmentation

may be specifically indicated in patients in which a prior determination of trait sAA levels has

revealed high basal arousal. From a scientific point of view, we may have identified an

important individual-differences factor that should be considered in future studies testing

methods to optimize extinction or exposure therapy – via L-DOPA and potentially also via

other routes.

The present data are not conclusive as to whether current sAA levels (state sAA) may be of

similar importance as trait sAA. Although we did observe a negative predictive effect of state

sAA for extinction retrieval as well as vmPFC reactivations, there was only a trend-level

interaction with L-DOPA treatment on retrieval. Because state sAA in our analysis was

defined in relation to individuals' trait sAA levels, an independent influence of current sAA

activity remains elusive. Future studies will in any case have to test the combined influence

of both variables.

Mechanistically, a simple explanation for an apparently better efficacy of L-DOPA with high

basal SNS activity may be that L-DOPA has more room to improve extinction consolidation

when it is compromised by (nor)adrenergic activity. Alternatively, L-DOPA may also directly

interfere with the impairing effects of central noradrenaline on extinction consolidation.

Previous research has suggested that, at high arousal levels, noradrenaline released from

locus coeruleus (LC) projections in the basolateral amygdala (BLA) acts via α1- and β-

adrenoceptors to increase BLA activity, while noradrenaline release in the mPFC reduces

activity there. Further, the increased BLA activity also attenuates mPFC output via inhibitory

BLA-mPFC projections. In sum, there is an increase in fear and impeded learning and/or

consolidation[33,39]. Dopamine may counteract these impairing effects of noradrenergic

activity, first, by decreasing prefrontal extracellular noradrenaline levels via increased

neuronal reuptake and activation of inhibitory α2-adrenoceptors[13], and second, by directly

reducing BLA-mediated inhibition of the mPFC[68]. Together, these findings point to a possible

mechanism by which systemic L-DOPA administration abolishes the detrimental effects of

high arousal on extinction learning and consolidation.

It is important to point out that our findings of trait and, to a lesser extent, state sAA

influences on extinction and L-DOPA-based extinction augmentation result from non-

preregistered analyses and that further investigation is needed to replicate them.

Importantly, future work should also experimentally manipulate the SNS. A further limitation

is that our method to collect salivary samples for sAA analysis could be improved, in

particular by utilizing collection devices with synthetic swabs or other appropriate materials

instead of cotton swabs. Beyond the role of salivary measures, it should be noted that all

reported effects on extinction retrieval in this study were observed in skin conductance

responses as our main index of conditioned responding. There were no predictive

relationships of vmPFC reactivations, sAA, or drug treatment on US expectancy ratings.

SCRs are implicit and objectively measurable, but expectancy or other ratings have the

practical advantage that they can be easily collected, in particular also in clinical settings.

One explanation for the absence of effects on ratings may lie in fear responses being carried

by dissociable systems (e.g.[69]), not all of which may be affected by our manipulation. Finally,

a clear gap in the current extinction and L-DOPA literature is that findings have not yet been

extended to women.

In this replication study, we aimed at reproducing the main findings from Gerlicher et al.

(2018)[16] with minor extensions. Our results establish the predictive role of a potentially PE-

related reactivation pattern in the vmPFC for extinction memory retrieval, tested in

spontaneous recovery. We discover that high basal SNS activity in male humans is

associated with impaired extinction consolidation and that L-DOPA promotes extinction

consolidation specifically in individuals with high basal SNS activity. This result opens up

new avenues for the investigation of L-DOPA as an enhancer of exposure therapy. For fear

extinction research, our results lead to new mechanistic hypotheses and suggest a new

possibility to explain the vast individual differences frequently observed in human extinction

research.

Methods

Design. The study design is as in the discovery study, with the exception of the added sAA

and sCORT measurements on days 1 and 2 and the added renewal test on day 3.

Technical parameters. All technical parameters, including the MRI scanner and equipment,

were kept identical to those used in the initial study.

Sample size. Our major effect of interest in the discovery study[16] was the interaction effect

of stimuli (CS+/CS-) and group (placebo/L-DOPA) on average SCRs during the test in the

extinction context on day 3, corresponding to our replication hypothesis 1. This effect had a

size of $\eta_p2=.166$, calculated by IBM SPSS Statistics (version 23, Chicago IL). Using this

effect size, we estimated using G*power 3.1.9.2[70] that with a power of 80% and an alpha of

0.05, the required sample size would be N=22, i.e., n=11 participants per group. As stated in

the Introduction, further statistical, logistical, and ethical considerations led us to aim at a

sample size of N=63 after drop-outs, requiring us to recruit 70 participants. Supplementary

Table 1 gives demographic information and further characteristics of the full recruited

sample.

Technical problems in data acquisition, described below for each data modality in the

corresponding Methods sections, reduced sample sizes for the analysis of the data

modalities. Supplementary Table 2 lists the missing data per participant, data modality, and

day. The final n for each analysis is indicated in the corresponding sections of the Results.

For the testing of our main hypothesis 1, using SCR data, we achieved a final n of 55, which

is 1.4 times greater than the discovery sample and 2.5 times greater than the critical sample

size estimated based on power calculation. This suggests that the null finding in the analysis

testing hypothesis 1 is not due to insufficient power.

Participants. As in the discovery study, we restricted recruitment to individuals of male sex.

The estrous cycle has been shown to interact with extinction memory consolidation[71,72], and

dopamine has been shown to have opposing effects on extinction depending on estrous

cycle phase[73]. Thus, by limiting ourselves to male participants, we could expect higher

sample homogeneity and correspondingly higher chances for detecting a neurobiological

mechanism. After safely demonstrating an effect with this strategy, a necessary next step is

to test transfer to non-male populations. A board-certified physician screened participants for

contraindications of L-DOPA intake, current physiological, neurological, or psychiatric

disorders, excessive consumption of nicotine (>10 cigarettes/day), alcohol (>15 glasses of

beer/wine per week), or cannabis (>1 joint/month), participation in other pharmacological

studies, and skin conductance non-responding (assessed by eSense Skin response,

Mindfield® Biosystems Ltd., Berlin, Germany). Drug abuse was assessed via a urine test

(M-10/3DT; Diagnostik Nord, Schwerin, Germany). The experiment was approved by the

local ethics committee (Ethikkommission der Landesaⵧrztekammer, Rhineland-Palatinate,

Germany) and was conducted in accordance with the Declaration of Helsinki. All participants

gave informed consent.

Experimental Design. We used a three-day fear conditioning and extinction paradigm

consisting of conditioning (day 1) in context A, extinction (day 2) in context B, and tests for

the effect of L-DOPA on extinction memory retrieval in the original extinction context (B) and

in a new context (C) (day 3). See Fig. 1a. Our study involves a 2x2 mixed factorial design

with stimulus as within- (CS+ vs. CS-) and drug group as between-subject (placebo vs. L-

DOPA) factor. As in our previous studies[14,16,18,19], the main outcome measures were average

CS evoked SCRs during each of the test contexts on day 3.

Stimuli. Two black geometric symbols (a square and a rhombus) presented in the center of

the screen served as CSs. The symbols were super-imposed on background pictures of one

of three different gray photos (living room, kitchen, and sleeping room), which served as

contexts A, B, and C. The assignment of symbols to the CS+ and CS- and backgrounds to

the conditioning, extinction or renewal context were randomized between participants and

groups. Diminishing the risk of low-visual feature differences between the CS+ and the CS-,

contrast and luminance of stimuli were adjusted using SHINE toolbox[74]. Stimuli were

presented using Presentation Software (Presentation®, Neurobehavioral Systems, Inc.,

Berkeley, CA, USA). A painful electrical stimulation consisting of three square-wave pulses

of 2 ms (50 ms inter-stimulus interval) was employed as US. Pain stimuli were generated by

using a DS7A electrical stimulator (Digitimer, Weybridge) and delivered to the skin through a

surface electrode with a platinum pin (Specialty Developments, Bexley, UK). Due to

observed incidences of high-voltage MRI artefacts in previous studies, we moved the

stimulation further away from the magnet's bore from the dorsal hand (pre-registered

location of stimulation) to the ankle.

Drug treatment. Participants were randomly assigned to the L-DOPA or the placebo group

using a randomization list generated before the start of the study, with the restriction that

groups had to be matched on self-reported trait anxiety based on the State-Trait Anxiety

Inventory questionnaire (STAI-T[75]). STAI-T scores did not differ between groups after

acquisition of n=45 participants, and therefore the predefined treatment group randomization

order was kept. After full data acquisition, STAI-T values did not differ between groups

(Supplementary Table 1). Other than in Gerlicher et al. (2018)[16] and than preregistered[28],

anxiety sensitivity index (ASI) scores, originally intended to also be matched between

groups, were not acquired due to an initially undetected technical failure. Drug preparation

was done by a person not involved in the experiments or analyses. Participants were

administered either 150/37.5 mg L- DOPA-benserazide (Levodopa-Benserazid-ratiopharm®,

Germany; for dosage see[14,17]) or a visually identical capsule filled with mannitol and aerosil

(i.e. placebo). Drugs were prepared and provided by the pharmacy of the University Medical

Center Mainz and administered in a double-blind fashion. Participants were asked to refrain

from eating, consuming caffeinated drinks, and smoking two hours prior to drug intake.

Fasting L-DOPA absorption is rapid, with L-DOPA peak plasma concentration reaching

within 15 to 60 min after oral intake[76]. Furthermore, L-DOPA half-time is also short (90

minutes), excluding that drug effects on extinction retrieval can be explained by direct drug

action on the test day.

Experimental procedures

Day 1 - fear learning. Participants filled out the state version of the state-trait anxiety

inventory (STAI-S), the Acceptance and Action questionnaire (AAQ[77]) the General Health

Questionnaire (GHQ[78]) (no group differences in all questionnaires; for analyses, see

Supplementary Table 1) and answered a list of questions assessing behaviors potentially

influencing on sCORT measurements (no group differences in all measures; for analyses,

see Supplementary Table 3). Subsequently, a saliva sample was collected. Participants

were placed in the MRI scanner, and SCR and pain stimulation electrodes were attached.

An 8-min resting-state scan was conducted. Participants then were familiarized with the

experiment through two CS presentations in each of the three contexts and a training of US-

expectancy ratings (for analyses, see Supplementary Fig 1). US-expectancy ratings were

taken on all three days as additional check for the success of conditioning, extinction, and

extinction retrieval, respectively. Subsequently, US-intensity was calibrated to a level rated

as "maximally painful, but still tolerable" (for analyses, see Supplementary Table 1).

Participants were then instructed that one symbol would never be followed by a pain

stimulus and that their task was to find out what rule applied to the other symbol. After

scanning onset, the paradigm started with US-expectancy ratings for each CS. After the

initial rating, a background picture representing context A appeared on the screen. The

context picture remained on the screen continuously throughout conditioning. Participants

were presented with 10 CS+ and 10 CS- trials. Notably, 5 out of 10 CS+ presentations (i.e.,

50%) were reinforced. CSs were presented for 4.5 seconds. In case of reinforced CS+

presentations, the US was delivered such that it co-terminated with the CS presentation.

Inter-trial intervals (ITIs) lasted 17, 18, or 19 seconds. Trial order was randomized in such a

way that not more than two trials of the same type (i.e., CS+ with US, CS+ without US, CS-)

succeed each other. Conditioning lasted approximately 12 min and scanning ended after

picture offset and US-expectancy ratings. After conditioning, another 8-min resting-state

scan was conducted, followed by anatomical scans (T1, T2, DTI, see Acquisition of MRI

data). Subsequently, electrodes were detached, participants gave another saliva sample and

filled out a list of questions assessing contingency knowledge (for details, see

Supplementary Methods). The whole procedure lasted approximately 90 min.

Day 2 - Extinction learning and consolidation. After 24 h (±2 h), participants came back

to the laboratory to fill out the STAI-S questionnaire and answer questions on behaviors

potentially influencing sCORT measures. Participants provided a saliva sample and were

placed in the scanner. Electrodes were attached, and participants were instructed that the

experiment would continue, and that their individual US strength from day 1 would be

applied. An 8-min resting-state scan was conducted. Before and after extinction, US-

expectancy ratings were taken. During extinction, a background picture representing context

B was continuously shown on the screen and participants were presented 15 CS+ and CS-

trials, using the same timings and pseudo-randomization algorithm as in conditioning.

Extinction lasted approximately 15 min. Subsequently, participants were taken out of the

scanner for saliva sample collection, detaching of electrodes and receiving either a placebo

or L-DOPA pill. Participants stayed under observation for 90 minutes. During this period,

further 8-min resting-state scans were performed ~10, 45, and 90 minutes after extinction

end, and saliva samples were collected each 20 min starting from the saliva collection time

point at extinction end (see Fig. 1c). Before leaving the laboratory, participants filled out the

STAI-S, a list of questions assessing contingency knowledge and a questionnaire assessing

possible side-effects of L-DOPA intake (for details, see Supplementary Table 4). The whole

procedure lasted approximately 150 min.

Day 3 – Extinction memory retrieval test (spontaneous recovery and renewal). After 24

h (±2 h), participants came back to the laboratory and filled out the STAI-S and side-effects

questionnaires and answered questions regarding potential confounding factors on sCORT

measurements. Participants provided a saliva sample and were placed in the scanner,

electrodes were attached, and participants were instructed that their US strength from day 1

would be applied and that the experiment would continue. An 8-min resting-state scan was

conducted. Before and after the test, US-expectancy ratings were taken. During the test,

participants were then presented 10 CS+ and 10 CS- trials in context B first (spontaneous

recovery test) and subsequently the same number of trials in context C (renewal test),

applying timings and randomization as on day 1. Finally, another saliva sample was taken,

and participants filled out a questionnaire assessing contingency knowledge. The test lasted

about 20 min; the whole procedure lasted approximately 60 min.

Skin conductance responses (SCRs). CS elicited SCRs were employed as conditioned

fear responses (CRs). Electrodermal activity was recorded from the thenar and hypo-thenar

of the non-dominant hand using self-adhesive Ag/AgACl electrodes prefilled with an isotonic

electrolyte medium and the Biopac MP150 with EDA100C device (EL-507, BIOPAC®

Systems Inc., Goleta, California, USA). The raw signal was amplified and low-pass filtered

with a cut-off frequency of 1 Hz. The onset of SCRs was visually scored offline in a time

window from 900 to 4000 ms after CS onset. The phasic amplitude of SCRs was calculated

by subtracting the onset background tonic skin conductance level (SCL) from the

subsequent peak, using a custom-made analysis script. Technical problems in data

acquisition led to missing data in n=0/3/3 participants for days 1/2/3, respectively. SCRs with

amplitudes smaller than 0.02 μs were scored as zero and remained in the analysis. If more

than 75% of trials during an experimental session were scored as zero, data of this

participant during that session was regarded as invalid and excluded from SCR analysis

(n=4/16/12 on days 1/2/3). Hence, in total, data from n=4/19/15 participants on days 1/2/3

were not available for analysis. To normalize distributions, data was log-transformed (+1 and

log) and range-corrected for each participant and experimental session (i.e., (SCR -

SCRmin) / SCRmax[79]).

Heart rate. Heart rate was recorded with an MRI-compatible fibre-optic pulse oximeter

during both resting-state scans and experimental sessions. We assessed heart rate during

the resting-state phases preceding each experimental session as an indicator of autonomic

stress response. Further analyses were not implemented, as data was largely lost due to a

storage error.

Pupil dilation. Pupil dilation was assessed monocularly using an MRI-compatible camera

(MR Cam Model 12M; MRC systems, Germany). The iViewX 2.8 software (Sensormotoric

Instruments, Germany) was used for recordings. Contrary to our pre-registration, no

analyses were conducted, as data was largely incomplete due to technical camera

problems.

US-expectancy ratings. Participants were asked to indicate the expectancy of receiving an

electric pain stimulation for each CS with a cursor on a visual analogue scale from 0 = "no

expectancy" to 100 = "high expectancy". The start position of the cursor on the scale was

determined randomly. Ratings were not available for n=0/3/3 participants on days 1/2/3.

According to the ratings, participants showed successful fear acquisition, extinction, and

extinction memory retrieval. For results, see Supplementary Fig 1.

Salivary alpha-amylase and cortisol measurement. To account for potential fluctuations

in salivary cortisol levels, all experimental procedures were scheduled to commence no

earlier than 4pm. This time allocation was implemented to ensure a balanced representation

of cortisol levels across participants and minimize any confounding effects that may arise

from diurnal variations. Saliva samples were collected from the participants outside the

scanner using cotton swabs. Samples were taken before and after the fMRI scan on days 1

and 3. On day 2, the saliva samples were collected before and after the first fMRI scan. Five

additional saliva samples were collected approximately every 20 minutes after extinction

learning (total n=11, Fig 1c for timeline). Saliva sample collection followed a standardized

procedure using Salivettes with cotton swab and white cap from the Sarstedt manufacturer.

Participants were instructed to place the swab in their cheek without chewing on it. The swab

remained in the cheek timed for one minute to allow for sufficient saliva collection. Samples

were collected throughout each day's experiment, and they were stored in the fridge at -20

°C approximately 10 minutes after the experiment's completion. The elapsed time from the

first sample collection (baseline) to freezing was approximately 100 minutes for day 1, 160

minutes for day 2, and 70 minutes for day 3. All samples remained frozen throughout the

data collection period of three years, and they were thawed one day prior to sending them to

the analyzing laboratory. All samples were analyzed within two weeks after being sent to the

laboratory. sCORT and sAA were determined using a commercial enzyme immunoassay,

see Supplementary Methods. Data was missing due to lack of saliva, or incomplete due to

insufficient saliva for more than one measure from one swab, that is, sCORT was measured

first, and no more saliva was left for the sAA measurement (missing values: sCORT: n=4 on

day 1, n=5 on day 2 pre extinction, n=11 of six swab times post extinction, n=10 on day 3;

sAA: n=9 on day 1, n=6 on day 2 pre extinction, n=14 of six swab times post extinction, n=14

on day 3).

Acquisition of MRI data. MRI data was acquired on a Siemens MAGNETOM Trio 3 Tesla

MRI System using a 32-channel head coil. Resting-state and task data were recorded using

gradient echo, echo planar imaging (EPI) with a multiband sequence covering the whole

brain (TR: 1000 ms, TE: 29 ms, multi-band acceleration factor: 4, voxel-size: 2.5 mm

isotropic, flip angle 56°, field of view: 210 mm;[80]). A high-resolution T1 weighted image was

acquired after the experiment on day 1 for anatomical visualization and normalization of the

EPI data (TR: 1900 ms, TE: 2540 ms, voxel size: 0.8 mm isotropic, flip angle 9°, field of

view: 260 mm, MPRAGE sequence). T2 weighted images were collected for preventative

neuro-radiological diagnostics for all participants (45 slices, TR: 6100 ms, TE: 79 ms, voxel

size: 3 mm isotropic, flip angle: 120°, Turbo Spin Echo (TSE) sequence). Lastly, we

collected multidimensional diffusion-weighted tensor images (DTI) from each participant (72

slices, voxel-size: 2 mm isotropic, TR: 9100 ms, TE: 85 ms, number of directions: 64,

diffusion weights: 2, b-value 1: 0 s/mm^2, b-value 2: 1000 s/mm^2, Multi-Directional Diffusion

Weighted (MDDW) sequence)).

For the acquisition of the resting-state scans, participants were instructed to remain awake,

keep their eyes open, fixate a black cross presented in the center of a grey screen, and let

their mind wander freely. Compliance with the instruction to remain awake was monitored

online using video recordings of the right eye. After each resting-state scan, participants

rated their tiredness on a scale from 0 = "not tired at all" to 100 = "almost fell asleep" (for

analyses, see Supplementary Table 1). No resting-state data had to be excluded due to

sleep.

MRI data were not available due to technical problems from n=0/3/3 participants on days

1/2/3.

Preprocessing of fMRI data. Only task and resting-state fMRI data from day 2, relevant to

address our hypotheses, were analyzed and are reported here. fMRI data was preprocessed

and analyzed using statistical parametric mapping (SPM12, Wellcome Trust Centre for

Neuroimaging, London, UK, http://www.fil.ion.ucl.ac.uk/) running on Matlab 2015b

(MathWorks®, Natick, Massachusetts, USA). The first 5 volumes of each scan were

discarded due to equilibrium effects. Preprocessing includes realignment and co-registration

of the mean functional image to the T1 weighted anatomical image. Subsequently, the T1

weighted anatomical image was segmented and normalized to add Montreal Neurologal

Institute (MNI) space based on SPM's tissue probability maps. Normalization of the

functional images was achieved by applying the resulting deformation fields to the realigned

and co-registered functional images. Lastly, functional data was smoothed using a 6mm full-

width-at-half-maximum Gaussian smoothing kernel. Data of participants was excluded when

movement peaks exceeded more than 3 mm or 2° (task data, n=7, resting-state data: n=6

further participants).

Data analysis

Statistical analysis of behavioral and psychophysiological data. For statistical analysis

of SCR and US-expectancy ratings, we conducted repeated measures ANOVA with stimulus

(CS+/CS-) as within- and drug (placebo/L-DOPA) as between-subject factor. Following our

previous studies[16,18,19] and preregistration, we tested whether fear was successfully acquired

based on CRs averaged across the last 20% of trials during fear conditioning (i.e., last 2

trials). Start-fear at extinction on day 2 was assessed using the average first 20% of trials

during extinction (i.e., first 3 trials). End-fear of extinction was determined across the last

20% of trials (i.e., last 3 trials) of the extinction session on day 2. The effect of L-DOPA on

retrieval was tested using CRs averaged across all trials of either the spontaneous recovery

or the renewal test on day 3. For all models that included trait and state sAA, we examined

the model assumptions such as normality of residuals, homoscedasticity, and linearity to

ensure their validity. All statistical analyses were implemented in R version 4.1.2 (2021-11-

01).

Multivariate fMRI analysis. Investigating potential reactivations of extinction specific MVPs

in the vmPFC, we analyzed the extinction task data (day 2) using a model including one

regressor for CS+ and CS- onsets, respectively, US-expectancy ratings, and context on/-

offset. Furthermore, the model included one regressor for the first five CS+ offsets, where

omission of the US was unexpected, and one for the first five CS- offsets, where US

omission was expected, as well as one regressor each for the remaining ten CS+ and CS-

offsets. All regressors were delta-functions convolved with the hemodynamic response

function (HRF). The MVP evoked by the first five US omissions at CS+ offset in the vmPFC

was extracted from the resulting beta-map in the vmPFC region of interest (ROI; see below).

Resting-state data was analyzed in accordance with a previous study examining memory

reactivation[81] (see also[16]), i.e., general linear models (GLMs) for each day 2 resting-state

scan (pre-, ~10, 45 and 90 minutes post-extinction) included delta-function regressors for

each volume (TR: 1 sec), thereby accounting for potential reactivations which may have

occurred during any point of the resting-state scan. No high-pass filtering was applied in the

resting-state models and AR(1) auto-correlation correction was employed. MVPs in the pre-

defined ROI during the resting state were extracted from the resulting beta-image series (TR

= 1 second, i.e., 480 – 5 = 475 beta images).

Subsequently, we correlated (Pearson correlation coefficient) the pattern evoked by the first

five US omissions at CS+ offset during extinction with the resulting 475 patterns of all four

resting-state scans and Fisher Z- transformed the correlation coefficients. The 475

correlations of the US omission pattern with the resting-state pattern recorded before

extinction learning was employed to create a baseline distribution. The mean and standard

deviation of this baseline distribution was used to transform each correlation between the

template and the resting-state patterns into a Z- score ($Zi = (ri - \mu i)/\sigma$). Correlations with a Z-

score exceeding a value of 2 ($Z > 2 \approx p < .05$) were counted as potential reactivations of the

CS+ offset-related vmPFC pattern. Reactivations were summed per participant and resting-

state scan. Subsequently, multiple linear regression analyses with number of CS+ offset-

related vmPFC reactivations at baseline, ~10, 45, and 90 min after extinction as predictors

and average differential (CS+>CS-) SCR during either the spontaneous recovery or the

renewal test as dependent variables were performed separately.

Regions of interest (ROIs). We focused our analysis on the vmPFC based on previous

work[16]. Control regions involved in fear and extinction learning were included comprising of:

anterior cingulate cortex (ACC), superior frontal gyrus (SFG), left and right insula, left and

right amygdala, and left and right hippocampus (see Supplementary Fig. 2 and 3 for results).

All ROIs were extracted from the Harvard-Oxford Atlas and thresholded at 50%-tissue

probability by previous work[16].

Data availability. The data that support the findings of this study are available upon request

from the corresponding author E.A. (e-mail: elena.andres@lir-mainz.de).

References

1.　Zuj, D. V., Palmer, M. A., Lommen, M. J. J. & Felmingham, K. L. The centrality of fear

extinction in linking risk factors to PTSD: A narrative review. *Neurosci. Biobehav. Rev.*

69, 15–35 (2016).

2.　Craske, M. G., Hermans, D. & Vervliet, B. State-of-the-art and future directions for

extinction as a translational model for fear and anxiety. *Philos. Trans. R. Soc. B Biol.*

Sci. **373**, (2018).

3.　Vervliet, B., Craske, M. G. & Hermans, D. Fear Extinction and Relapse: State of the

Art. *Annu. Rev. Clin. Psychol.* **9**, 215–248 (2013).

4.   Pavlov, I. P. Conditioned Reflexes. An Investigation of the Physiological Activity of the

Cerebral Cortex. *Oxford Univ. Press* (1927) doi:10.2307/1134737.

5.   Bouton, M. E. Context and behavioral processes in extinction. *Learn. Mem.* **11**, 485–

494 (2004).

6.   Müller, G. E. & Pilzecker, A. Experimentelle Beiträge zur Lehre vom Gedächtnis. *Z.*

Psychol. Ergänzungsband 1–300 (1900).

7.   Nadel, L. & Moscovitch, M. Memory consolidation, retrograde amnesia and the

hippocampal complex. *Curr. Opin. Neurobiol.* **7**, 217–227 (1997).

8.   Squire, L. R. Mechanisms of Memory. *Science (80-.).* **232**, 1612–1619 (1986).

9.   Singewald, N., Schmuckermair, C., Whittle, N., Holmes, A. & Ressler, K. J.

Pharmacology of cognitive enhancers for exposure-based therapy of fear, anxiety and

trauma-related disorders. *Pharmacol. Ther.* **149**, 150–190 (2015).

10.   Iemolo, A., De Risi, M. & De Leonibus, E. *Role of Dopamine in memory consolidation.*

Sakakibara, M and Ito E eds. Memory Consolidation. Nova Science Publisher (2015).

11.   Hikind, N. & Maroun, M. Microinfusion of the D1 receptor antagonist, SCH23390 into

the IL but not the BLA impairs consolidation of extinction of auditory fear conditioning.

Neurobiol. Learn. Mem. **90**, 217–222 (2008).

12.   Abraham, A. D., Cunningham, C. L. & Lattal, K. M. Methylphenidate enhances

extinction of contextual fear. *Learn. Mem.* **19**, 67–72 (2012).

13.   Dayan, L. & Finberg, J. P. M. L-DOPA increases noradrenaline turnover in central and

peripheral nervous systems. *Neuropharmacology* **45**, 524–533 (2003).

14.   Haaker, J. *et al.* Single dose of <scp>l</scp> -dopa makes extinction memories

context-independent and prevents the return of fear. *Proc. Natl. Acad. Sci.* **110**,

E2428–E2436 (2013).

15.   Whittle, N. *et al.* Enhancing dopaminergic signaling and histone acetylation promotes
long-term rescue of deficient fear extinction. *Transl. Psychiatry* **6**, (2016).

16.   Gerlicher, A. M. V., Tüscher, O. & Kalisch, R. Dopamine-dependent prefrontal
reactivations explain long-term benefit of fear extinction. *Nat. Commun.* **9**, 4294
(2018).

17.   Haaker, J., Lonsdorf, T. B. & Kalisch, R. Effects of post-extinction l-DOPA
administration on the spontaneous recovery and reinstatement of fear in a human
fMRI study. *Eur. Neuropsychopharmacol.* **25**, 1544–1555 (2015).

18.   Gerlicher, A. M. V., Tüscher, O. & Kalisch, R. L-DOPA improves extinction memory
retrieval after successful fear extinction. *Psychopharmacology (Berl).* (2019)
doi:10.1007/s00213-019-05301-4.

19.   Gerlicher, A. M. V., Ilhan-Bayrakci, M., Tüscher, O. & Kalisch, R. Post-extinction L-
DOPA administration to reduce the return of fear in humans. *PsyArXiv Prepr.* (2022)
doi:10.31234/osf.io/rhnsa.

20.   Milad, M. R. & Quirk, G. J. Fear Extinction as a Model for Translational Neuroscience:
Ten Years of Progress. *Annu. Rev. Psychol.* **63**, 129–151 (2012).

21.   Hugues, S., Garcia, R. & Léna, I. Time course of extracellular catecholamine and
glutamate levels in the rat medial prefrontal cortex during and after extinction of
conditioned fear. *Synapse* **61**, 933–937 (2007).

22.   Rescorla, R. A. & Wagner, A. R. A Theory of Pavlovian Conditioning: Variations in the
Effectiveness of Reinforcement and Nonreinforcement. *Class. Cond. II Curr. Res.*
theory 64–99 (1972) doi:10.1101/gr.110528.110.

23.   Pearce, J. M. & Hall, G. A model for Pavlovian learning: Variations in the

882   effectiveness of conditioned but not of unconditioned stimuli. *Psychol. Rev.* **87**, 532–

883   552 (1980).

884   24.   Dunsmoor, J. E., Niv, Y., Daw, N. & Phelps, E. A. Rethinking Extinction. *Neuron* **88**,

885   47–63 (2015).

886   25.   Papalini, S., Beckers, T. & Vervliet, B. Dopamine: from prediction error to

887   psychotherapy. *Transl. Psychiatry* **10**, 164 (2020).

888   26.   Kalisch, R., Gerlicher, A. M. V. & Duvarci, S. A Dopaminergic Basis for Fear

889   Extinction. *Trends Cogn. Sci.* **23**, 274–277 (2019).

890   27.   Thiele, M., Yuen, K. S. L., Gerlicher, A. V. M. & Kalisch, R. A ventral striatal prediction

891   error signal in human fear extinction learning. *Neuroimage* **229**, 117709 (2021).

892   28.   Chuan-Peng, H. *et al.* Dopamine-dependent prefrontal reactivations explain long-term

893   benefit of fear extinction: A direct replication attempt. *osf.io/x64cn* (2018).

894   29.   Simonsohn, U. Small Telescopes: Detectability and the Evaluation of Replication

895   Results. *Psychol. Sci.* **26**, 559–569 (2015).

896   30.   Bacchetti, P., Wolf, L. E., Segal, M. R. & McCulloch, C. E. Ethics and Sample Size.

897   *Am. J. Epidemiol.* **161**, 105–110 (2005).

898   31.   Kirschbaum, C. & Hellhammer, D. H. Salivary Cortisol in Psychobiological Research:

899   An Overview. *Neuropsychobiology* **22**, 150–169 (1989).

900   32.   Nater, U. M. & Rohleder, N. Salivary alpha-amylase as a non-invasive biomarker for

901   the sympathetic nervous system: Current state of research.

902   *Psychoneuroendocrinology* **34**, 486–496 (2009).

33.   Giustino, T. F. & Maren, S. Noradrenergic Modulation of Fear Conditioning and

Extinction. *Front. Behav. Neurosci.* **12**, 1–20 (2018).

34.   Joëls, M., Pu, Z., Wiegert, O., Oitzl, M. S. & Krugers, H. J. Learning under stress: how

does it work? *Trends Cogn. Sci.* **10**, 152–158 (2006).

35.   Meir Drexler, S., Merz, C. J., Jentsch, V. L. & Wolf, O. T. Stress modulation of fear

and extinction in psychopathology and treatment. *Neuroforum* **26**, 133–141 (2020).

36.   Muehlhan, M., Lueken, U., Wittchen, H. U. & Kirschbaum, C. The scanner as a

stressor: Evidence from subjective and neuroendocrine stress parameters in the time

course of a functional magnetic resonance imaging session. *Int. J. Psychophysiol.* **79**,

118–126 (2011).

37.   Visser, R. M., Kunze, A. E., Westhoff, B., Scholte, H. S. & Kindt, M. Representational

similarity analysis offers a preview of the noradrenergic modulation of long-term fear

memory at the time of encoding. *Psychoneuroendocrinology* **55**, 8–20 (2015).

38.   Eatough, E. M., Shirtcliff, E. A., Hanson, J. L. & Pollak, S. D. Hormonal reactivity to

MRI scanning in adolescents. *Psychoneuroendocrinology* **34**, 1242–1246 (2009).

39.   Maren, S. Unrelenting Fear Under Stress: Neural Circuits and Mechanisms for the

Immediate Extinction Deficit. *Front. Syst. Neurosci.* **16**, 1–13 (2022).

40.   Nater, U. M. *et al.* Stress-induced changes in human salivary alpha-amylase activity -

Associations with adrenergic activity. *Psychoneuroendocrinology* **31**, 49–58 (2006).

41.   Bali, A. & Jaggi, A. S. Clinical experimental stress studies: Methods and assessment.

Rev. Neurosci. **26**, 555–579 (2015).

42.   Koh, D., Ng, V. & Naing, L. Alpha amylase as a salivary biomarker of acute stress of

venepuncture from periodic medical examinations. *Front. Public Heal.* **2**, 1–5 (2014).

43.   Timmers, I. *et al.* Fear of pain and cortisol reactivity predict the strength of stress-

induced hypoalgesia. *Eur. J. Pain (United Kingdom)* **22**, 1291–1303 (2018).

44. Out, D., Granger, D. A., Sephton, S. E. & Segerstrom, S. C. Disentangling sources of
individual differences in diurnal salivary α-amylase: Reliability, stability and sensitivity
to context. *Psychoneuroendocrinology* **38**, 367–375 (2013).

45. Granger, D. A. *et al.* Integrating the measurement of salivary α-amylase into studies of
child health, development, and social relationships. *J. Soc. Pers. Relat.* **23**, 267–290
(2006).

46. Wang, X. *et al.* Stability and predictors of change in salivary cortisol measures over
six years: MESA. *Psychoneuroendocrinology* **49**, 310–320 (2014).

47. Nater, U. M., Rohleder, N., Schlotz, W., Ehlert, U. & Kirschbaum, C. Determinants of
the diurnal course of salivary alpha-amylase. *Psychoneuroendocrinology* **32**, 392–401
(2007).

48. Out, D., Bakermans-Kranenburg, M. J., Granger, D. A., Cobbaert, C. M. & van
Ijzendoorn, M. H. State and trait variance in salivary α-amylase: A behavioral genetic
study. *Biol. Psychol.* **88**, 147–154 (2011).

49. Burgos-Robles, A., Vidal-Gonzalez, I., Santini, E. & Quirk, G. J. Consolidation of Fear
Extinction Requires NMDA Receptor-Dependent Bursting in the Ventromedial
Prefrontal Cortex. *Neuron* **53**, 871–880 (2007).

50. Tambini, A. & Davachi, L. Awake Reactivation of Prior Experiences Consolidates
Memories and Biases Cognition. *Trends Cogn. Sci.* **23**, 876–890 (2019).

51. Schultz, W., Dayan, P. & Montague, P. R. A neural substrate of prediction and
reward. *Science (80-.).* **275**, 1593–1599 (1997).

52. Bouton, M. E. Context, ambiguity, and unlearning: sources of relapse after behavioral

extinction. *Biol. Psychiatry* **52**, 976–986 (2002).

53. Nasser, H. M. & McNally, G. P. Appetitive-aversive interactions in Pavlovian fear

conditioning. *Behav. Neurosci.* **126**, 404–422 (2012).

54. Kandel, E. R., Dudai, Y. & Mayford, M. R. The molecular and systems biology of

memory. *Cell* **157**, 163–186 (2014).

55. Izquierdo, I. *et al.* Different molecular cascades in different sites of the brain control

memory consolidation. *Trends Neurosci.* **29**, 496–505 (2006).

56. Zorawski, M., Cook, C. A., Kuhn, C. M. & Labar, K. S. Sex, stress, and fear: Individual

differences in conditioned learning. *Cogn. Affect. Behav. Neurosci.* **5**, 191–201

(2005).

57. Ditzen, B., Ehlert, U. & Nater, U. M. Associations between salivary alpha-amylase and

catecholamines - A multilevel modeling approach. *Biol. Psychol.* **103**, 15–18 (2014).

58. Van Stegeren, A., Rohleder, N., Everaerd, W. & Wolf, O. T. Salivary alpha amylase as

marker for adrenergic activity during stress: Effect of betablockade.

Psychoneuroendocrinology **31**, 137–141 (2006).

59. Hermans, E. J. *et al.* Stress-related noradrenergic activity prompts large-scale neural

network reconfiguration. *Science (80-.).* **334**, 1151–1153 (2011).

60. Ehlert, U., Erni, K., Hebisch, G. & Nater, U. Salivary α-amylase levels after yohimbine

challenge in healthy men. *J. Clin. Endocrinol. Metab.* **91**, 5130–5133 (2006).

61. Hidalgo, V., Almela, M., Villada, C. & Salvador, A. Acute stress impairs recall after

interference in older people, but not in young people. *Horm. Behav.* **65**, 264–272

(2014).

62. Schwarz, J. *et al.* Does sleep deprivation increase the vulnerability to acute

psychosocial stress in young and older adults? *Psychoneuroendocrinology* **96**, 155–
165 (2018).

63. Yamane, N. *et al.* Salivary Alpha-Amylase Activity and Mild Cognitive Impairment
    among Japanese Older Adults: The Toon Health Study. *J. Prev. Alzheimer's Dis.* **4**,
    752–757 (2022).

64. Seals, D. R. & Dinenno, F. A. Collateral damage: cardiovascular consequences of
    chronic sympathetic activation with human aging. *Am. J. Physiol. Circ. Physiol.* **287**,
    H1895–H1905 (2004).

65. Almstrup, F. B. *et al.* Associations between saliva alpha-amylase, heart rate
    variability, saliva cortisol and cognitive performance in individuals at ultra high-risk for
    psychosis. *Schizophr. Res.* **255**, 165–172 (2023).

66. Pagen, L. H. G. *et al.* Elevated Activity of the Sympathetic Nervous System Is Related
    to Diminished Practice Effects in Memory: A Pilot Study. *J. Alzheimer's Dis.* **80**, 1675–
    1685 (2021).

67. Wang, L. Y. *et al.* Cerebrospinal fluid norepinephrine and cognition in subjects across
    the adult age span. *Neurobiol. Aging* **34**, 2287–2292 (2013).

68. Floresco, S. B. & Tse, M. T. Dopaminergic regulation of inhibitory and excitatory
    transmission in the basolateral amygdala-prefrontal cortical pathway. *J. Neurosci.* **27**,
    2045–2057 (2007).

69. Soeter, M. & Kindt, M. Erasing fear for an imagined threat event.
    Psychoneuroendocrinology **37**, 1769–1779 (2012).

70. Faul, F., Erdfelder, E., Lang, A.-G. & Buchner, A. G*Power 3: A flexible statistical
    power analysis program for the social, behavioral, and biomedical sciences. *Behav.
    Res. Methods* **39**, 175–191 (2007).

71.   Cover, K. K., Maeng, L. Y., Lebrón-Milad, K. & Milad, M. R. Mechanisms of estradiol

in fear circuitry: Implications for sex differences in psychopathology. *Transl.*

Psychiatry **4**, (2014).

72.   Stark, R. *et al.* Influence of the stress hormone cortisol on fear conditioning in

humans: Evidence for sex differences in the response of the prefrontal cortex.

Neuroimage **32**, 1290–1298 (2006).

73.   Rey, C. D., Lipps, J. & Shansky, R. M. Dopamine D1 receptor activation rescues

extinction impairments in low-estrogen female rats and induces cortical layer-specific

activation changes in prefrontal-amygdala circuits. *Neuropsychopharmacology* **39**,

1282–1289 (2014).

74.   Willenbockel, V. *et al.* Controlling low-level image properties: The SHINE toolbox.

Behav. Res. Methods **42**, 671–684 (2010).

75.   Spielberger, C. D., Gorsuch, R. L. & Lushene, R. E. Manual for the state-trait anxiety

inventory. *Consult. Psychol. Press* (1970).

76.   Contin, M. & Martinelli, P. Pharmacokinetics of levodopa. *J. Neurol.* **257**, (2010).

77.   Hayes, S. C. *et al.* Measuring experiential avoidance: A preliminary test of a working

model. *Psychol. Rec.* **54**, 553–578 (2004).

78.   Goldberg, D. P. & Hillier, V. F. A scaled version of the General Health Questionnaire.

Psychol. Med. **9**, 139–145 (1979).

79.   Dawson, M. E., Schell, A. M. & Filion, D. L. The electrodermal system. *Handb.*

Psychophysiology, Fourth Ed. 217–243 (2007) doi:10.1017/9781107415782.010.

80.   Feinberg, D. A. *et al.* Multiplexed echo planar imaging for sub-second whole brain fmri

and fast diffusion imaging. *PLoS One* **5**, (2010).

81.   Staresina, B. P., Alink, A., Kriegeskorte, N. & Henson, R. N. Awake reactivation

1021   predicts memory in humans. *Proc. Natl. Acad. Sci.* **110**, 21159–21164 (2013).

Acknowledgments

This work was funded by the Deutsche Forschungsgemeinschaft (DFG), SFB1193,

subproject C01 to R.K.

Author contributions

E.A., C.-P.H., A.M.V., O.T. and R.K. designed the experiment; E.A. and C.-P.H. collected the

data; E.A., C.-P.H., and B.M. analyzed the data; and E.A. and R.K. wrote the paper.

Competing interests

R.K. has received advisory honoraria from JoyVentures, Herzlia, Israel.

Fig. 1 Experimental design and skin conductance responses. a A three-day fMRI paradigm was employed with fear conditioning on day 1 in context A, extinction on day 2 in context B, and test on day 3 first in context B (the context in which the extinction learning took place), and then in a new context C. Immediately after extinction learning, participants were administered either a placebo or a L-DOPA pill. Drug administration was randomized and double-blinded (placebo: n=35, L-DOPA n= 35, all male, for group characteristics, see Supplementary Table 1) . All three phases occurred in the MRI scanner. Further, resting-state scans (R) were collected before and after fear learning, before the start of extinction, ~10, 45 and 90 minutes after extinction, and before test. **b** Conditioned responses (CRs) were assessed as SC responses (SCRs) to the CS+ and CS- throughout all experimental phases. The upper panel depicts the mean SCRs of the placebo group (n=33/27/30 on days 1/2/3), the lower panel shows results of the L-DOPA-treated group (n=33/24/25 on days 1/2/3). Data is presented as mean ± standard error of the mean (sem). **c** Saliva samples were collected on day 2 before the start of the experimental paradigm (time 0 min) followed by a resting-state scan (approx. 8 min) and extinction (approx. 15 min). Subsequent saliva samples were taken right after extinction before pill intake and then about every 20 minutes (+/- 2 minutes), with the last sample taken after the last resting-state measurement at 90 min post-extinction. Samples on day 1 and 3 were collected before and after the paradigm in the scanner. Tubes represent saliva sample collections. Scanner bore symbols represent scanning times. Cond = Conditioning, Ext = Extinction.

Fig. 2 Relationship between CRs at the end of extinction (end -fear) on day 2 and CRs at test in the extinction context B and a new context C on day 3. Extinction success (assessed by differential CRs (CS+>CS-) at the end of extinction) positively predicted spontaneous recovery (differential CRs (CS+>CS-) at test in extinction context B; **a**) and renewal (differential CRs (CS+>CS-) at test in a new context; **b**) in both the placebo- and the L-DOPA-treated groups.

Fig. 3 Number of spontaneous reactivations of early CS+ offset -related vmPFC patterns 90 min post extinction on day 2 predicts CRs at test on day 3 in both groups. Extinction memory retrieval prediction by number of vmPFC reactivations was observed across both L-DOPA- and placebo-treated participants 90 min after extinction. Regression line applies to both groups.

Fig. 4 Effect of post-extinction L-DOPA administration on the number of spontaneous reactivations of CS+ offset-related vmPFC patterns during resting-state scans on day 2. There was no effect of L-DOPA on numbers of vmPFC reactivation patterns during resting -state fRMI 10, 45, and 90 min post extinction learning (n=46). Data is presented as median ± 1.5*IQR. Dots represent outliers lying outside this range.

Fig. 5 sAA and sCORT measurements on days 1-3. a sAA and **b** sCORT levels exhibited linear decreases on day 2 with peak levels in the group observed before extinction. sAA and sCORT did not differ on day 1 between time points and groups, that is, before and after conditioning (sAA: time: $F_{1,59}=2.28$, p=.14, group: $F_{1,59}=.72$, p=.40, time*group: $F_{1,59}=.88$, p=.35, n=61; sCORT: time: $F_{1,64}=2.95$, p=.09, group: $F_{1,64}=2.78$, p=.10, time*group: $F_{1,64}=.28$, p=.60, n=66). On day 3, only sCORT was significantly decreased after the experimental procedure in both groups (sAA: time: $F_{1,54}=.26$, p=.62, group: $F_{1,54}=.29$, p=.60, time*group: $F_{1,54}=2.97$, p=.09, n=57; sCORT: time: $F_{1,57}=10.67$, p=.002, group: $F_{1,57}=.07$, p=.80, time*group: $F_{1,57}=.18$, p=.67, n=60). Saliva samples were collected as shown in Fig. 1c. Data is presented as median ± 1.5*IQR. Dots represent outliers lying outside this range.

Fig. 6 Relationship between trait and state sAA and CRs at test in context B on day 3. a Whereas high trait sAA was predictive for impaired extinction memory retrieval at test in the extinction context B (spontaneous recovery) in the placebo group, this effect was rescued by L-DOPA administration. **b** A similar, though non-significant pattern emerged also in participants with high state sAA on day 2, defined as deviation of the individual's peak sAA level on that day from their trait sAA value. Regression lines shown represent the predicted relationship between trait or state sAA and CRs at test based on the implemented model.

Fig. 7 Relationship between trait and state sAA and number of vmPFC reactivations 90 min after extinction learning. a High trait sAA was non-significantly associated with vmPFC reactivations 90 min after extinction. **b** High state sAA showed a negative association with vmPFC reactivations 90 min after extinction in the placebo group, whereas this effect was abolished in the L-DOPA group. Regression lines shown represent the predicted relationship between trait or state sAA and CRs at test based on the implemented model.