Title: Degenerative changes are associated with severity of anterior cruciate

ligament injury within the skeletally immature joint

Authors: Danielle Howe[1,2], Jacob D. Thompson[1,2], Stephanie D. Teeter[1], Margaret Easson[1],

Olivia Barlow[3], Emily H. Griffith[4], Lauren V. Schnabel[2,5], Jeffrey T. Spang[6],

Matthew B. Fisher[1,2,6*]

Affiliations:

[1]Joint Department of Biomedical Engineering, North Carolina State University and the

University of North Carolina at Chapel Hill; Raleigh, NC, 27695, USA

[2]Comparative Medicine Institute, North Carolina State University; Raleigh, NC, 27607, USA

[3]Department of Mechanical Engineering, North Carolina State University; Raleigh, NC, 27607,

USA

[4]Department of Statistics, North Carolina State University; Raleigh, NC 27695

[5]Department of Clinical Sciences, North Carolina State University; Raleigh, NC, 27695, USA

[6]Department of Orthopaedics, University of North Carolina at Chapel Hill; Chapel Hill, NC,

27599, USA

*Corresponding author. Email: mbfisher@ncsu.edu

One Sentence Summary: Joint degeneration was associated with degree of joint instability after

partial and complete ACL injury in a skeletally immature porcine model.

Abstract

Anterior cruciate ligament (ACL) injuries are a major problem in the pediatric and adolescent

populations. Some of these injuries extend only partially through the tissue cross-section; yet,

there is limited data to inform clinical treatment of such partial tears. In particular, it is unknown

how injury severity impacts long-term degenerative changes in the joint. Here, we leverage a

skeletally immature preclinical porcine model to evaluate joint biomechanics and degeneration

after partial (isolated anteromedial (AM) or posterolateral (PL) bundle) or complete ACL injury.

Six months after injury, joint laxity increases were minimal after PL bundle injury, minor after

AM bundle injury, and major after ACL injury. Joint degeneration (evaluated in the cartilage and

meniscus) was minimal after PL bundle injury, moderate after AM bundle injury, and substantial

after ACL injury. With subjects grouped by clinical Lachman grade (indicating the extent of

joint destabilization), degeneration was associated with increasing grade, irrespective of injury

type. These findings point to the importance of considering joint laxity as a factor when treating

young patients, particularly those with partial ACL injuries.

Main Text

Injuries to the anterior cruciate ligament (ACL) of the knee, particularly those occurring in children and adolescents, are a significant clinical and societal concern (*1*). The total number of ACL injuries across all age groups is substantial (200,000-250,000 yearly in the United States alone) (*2, 3*). A specific concern is the 2-4 fold increase in the rate of injury and subsequent surgical treatment in the pediatric population compared to adults (*4-6*).

Preventing joint degeneration is a major concern. The natural occurrence of joint degeneration after ACL rupture is well documented in adults (*7*). In the pediatric population, full-thickness ACL tears result in two-fold increases in knee laxity (*8*) and 3-4 times greater risk of developing meniscal and cartilage lesions with delayed surgical intervention (*9*). It is commonly assumed that joint instability after ACL injury is a major factor in subsequent joint degeneration (*10-12*), though the degree of joint instability required to cause degeneration is unknown for skeletally immature patients. Nevertheless, surgical reconstruction is the standard practice to restore joint stability for young, active patients with complete ACL tears (*13*), despite concerns about growth plate disruption.

For partial-thickness ACL tears, which account for 10-27% of diagnosed ACL injuries (*14*) and result in a more minor (~30% on average) increase in knee laxity (*15*), the superiority of operative or non-operative treatment to limit degenerative changes is less evidence-based. On one hand, subsequent meniscal and chondral pathologies after non-surgical treatment of partial tears are reported in young patients (*16, 17*). On the other hand, the invasive nature of reconstructive surgery can exacerbate degenerative changes (*18*), and approximately 50% of young patients who undergo ACL reconstruction still develop osteoarthritis after 10 years (*19*).

A major complicating factor in determining the best clinical treatment is that not all

partial tears are the same. The ACL has two major subregions, the anteromedial (AM) bundle

and the posterolateral (PL) bundle, and partial tears can occur in either. Tear location may impact

functional outcomes after partial injury, and the results may differ between children and adults.

Specifically, a study in adult patients reported similar joint stability after isolated AM and PL

bundle injuries (*15*), while a study in pediatric patients reported better outcomes for AM bundle

injuries (*20*). Furthermore, in a skeletally immature pig model, the biomechanical function of the

AM and PL bundles of the ACL depends on age and sex, such that the AM bundle plays a larger

functional role in older versus younger subjects and males versus females (*21, 22*). Since

evidence regarding treatment of bundle-specific partial injuries in children and adolescents is

sparse, there is a need for further investigation of long-term outcomes and degenerative changes

in skeletally immature subjects. Large exploratory studies in humans are challenging to perform

in the pediatric patient population. However, the pig is a valuable preclinical model with similar

ACL anatomy and function to humans (*23-25*), which can be used to control injury severity and

assess functional outcomes and degenerative changes after injury.

In the present study, we investigated biomechanics, remodeling, and degeneration in the

knee joint after isolated AM bundle, PL bundle, and complete ACL tears in a juvenile porcine

model. We hypothesized firstly that partial (AM and PL bundle) tears would result in small

levels of instability and degeneration relative to complete ACL tears. Secondly, we hypothesized

that remaining soft tissues in the joint would remodel in response to changes in biomechanical

loading after ACL injury, such that the degree of remodeling would associate with injury

severity. At 6 months after injury, joint laxity showed no change after PL bundle injury, minor

increases after AM bundle injury and major increases after ACL injury. Similarly, PL bundle

injury resulted in minimal degeneration, AM bundle injury resulted in moderate degeneration,

and complete ACL injury led to substantial degeneration, such that the degree of degenerative

changes was associated with joint instability.

Results

Joint kinematics after partial and complete ACL injury

To evaluate the impact of AM and PL bundle tears on knee function and degeneration in

skeletally immature subjects, we performed arthroscopic transection and removal of the AM

bundle (AM-), PL bundle (PL-) or ACL (ACL-) in juvenile Yorkshire crossbreed pigs and

assessed biomechanics and joint health after 24 weeks (Fig. 1a). Joint kinematics were evaluated

using a robotic testing system first under applied anterior-posterior (AP) drawer to simulate

clinical exams (Fig. 1b-e). AP tibial translation (APTT) was greater in AM- joints (46%,

$P<0.0001$) and ACL- joints (195%, $P<0.0001$) compared to contralateral controls, while PL-

joints showed no significant difference (P=0.22) (Fig. 1c-d). Between injury groups, the inter-

limb difference in APTT was significantly different for all comparisons ($P\leq0.0061$) (Fig. 1e).

Under applied compression (Fig. 1f-i) (to simulate weight bearing) and varus-valgus moments

(Fig. 1j-m) (to simulate rotational loads), anterior tibial translation (ATT) was only greater in

ACL- joints compared to contralateral controls (327%, P=0.0005) (Fig. 1h), and the interlimb

difference in ATT was greater for the ACL- group compared to AM- and PL- groups ($P<0.0001$

for each) (Fig. 1i). Likewise, varus-valgus (VV) rotation was greater in ACL- joints compared to

contralateral controls (33%, P=0.012) (Fig. 1l), with a larger interlimb difference compared to

AM- and PL- groups ($P\leq0.0037$) (Fig. 1m). Sex did not significantly affect kinematics under AP

or VV loads, but under compression, ATT was significantly greater in males than females within

the AM- group (P=0.003) (Supplementary Fig. 1). Overall, PL bundle injury induced minimal

joint instability, AM bundle injury induced moderate joint instability, and complete ACL injury

induced major joint instability.

Fig. 1. Biomechanical assessment of porcine joints 6 months after partial and complete

ACL transection. a, Schematic of arthroscopic transection of the AM bundle, PL bundle, or

entire ACL, followed by evaluation after 24 weeks. b, Diagram of anterior-posterior (AP) loads

applied to the tibia. c, Resulting mean force-translation curves. AP tibial translation (APTT)

shown as d, absolute values and e, interlimb differences. f, Diagram of compressive load applied

to the tibia. g, Resulting mean force-translation curves. Anterior tibial translation (ATT) shown

as h, absolute values and i, interlimb differences. j, Diagram of varus-valgus (VV) torques

applied to the tibia. k, Resulting mean torque-rotation curves. VV rotation shown as l, absolute

values and m, interlimb differences. Red arrows indicate the direction of the applied load, and

black arrows indicate the direction of measured translation (b, f, j). Points and shaded area

represent mean ± 95% CI (c, g, k). Lines connect paired joints (d, h, l). Individual data points

(n=12 for PL- and AM- groups, n=8 for ACL- group) presented with mean ± 95% CI (e, i, m). *

P<0.05 between injured and contralateral, from paired t-test with Holm-Sidak correction.

#P<0.05 between injury type, from one-way ANOVA with Tukey post hoc test.

Subhead 2: Remodeling of the uninjured bundle of the ACL

In partial injury (AM- and PL-) joints, the in-situ force (Fig. 2a) carried by the remaining

ACL bundle was greater compared to the corresponding bundle force in contralateral controls

under applied anterior drawer (P≤0.00019 for both injury types) (Fig. 2b). However, under joint

compression, the remaining bundle force was elevated only for AM- joints (P=0.0016) (Fig. 2c).

Remaining bundle in-situ force did not differ on average under varus or valgus moments,

although considerable specimen-to-specimen variability existed (Supplementary Fig. 2). To

assess remodeling of the ACL that may drive functional changes, the cross-sectional area (CSA)

of the remaining bundle in injured joints was measured from MRI and compared to the

corresponding bundle in contralateral joints (Fig. 2d). The remaining bundle was larger in CSA

compared to contralateral controls for PL- joints (AM bundle) by 13% and AM- joints (PL

bundle) by 60% (P≤0.022, for both) (Fig. 2e). In addition, bundle collagen organization was

assessed through T2* mapping using MRI (Fig. 2f). No differences were found in median T2*

value of the remaining bundle in injured joints compared to contralateral controls for either

injury group (Fig. 2g).

Fig. 2. Remodeling of the remaining bundle after transection of the AM or PL bundle of the

ACL. a, Schematic of in-situ bundle forces measured under applied loads. Resultant force in the

uninjured bundle under applied b, anterior drawer and c, compression. d, Schematic

demonstrating methods used to measure bundle cross-sectional area (CSA) from segmented MRI

scans. e, CSA of the uninjured bundle. f, Schematic showing representative sagittal MR image

overlaid with the T2* map of the ACL. Plot shows signal intensity at the six measured echo

times (blue circles) and fitted monoexponential decay curve (yellow) for a representative voxel.

Calculated T2* relaxation time and goodness-of-fit (r^2) shown in graph. g, Median T2* value of

the uninjured bundle. Individual data points (n=12 per injury group) presented with

mean/median. Lines connect paired joints. *P<0.05 between injured and contralateral, from

paired t-test with Holm-Sidak correction.

Subhead 3: Cartilage and synovium degeneration

To assess if the type of injury impacted degeneration of articular cartilage in the joint,

T1ρ mapping was performed via MRI, and the median T1ρ value was calculated for the medial

and lateral femoral and tibial articular cartilage (Fig. 3a). Higher T1ρ values are linked to lower

proteoglycan content (*26, 27*). T1ρ values relative to the contralateral control were not

significantly elevated in any compartment for the PL- group, but were elevated in the medial

femoral and lateral tibial compartments for the AM- group and the lateral femoral and lateral

tibial compartments for the ACL- group (P≤0.022 for all, Fig. 3b-e). Furthermore, the interlimb

difference in T1ρ values were significantly greater in the AM- group relative to the PL- group for

the medial femoral compartment and in the ACL- group relative to the PL- group for the lateral

femoral and lateral tibial compartments (P≤0.012 for all).

Macroscopic articular cartilage damage was scored according to Osteoarthritis Research

Society International (OARSI) guidelines (*28*) for the four compartments (Fig. 3f,

Supplementary Fig. 3) and summed for a composite joint score. The composite joint scores were

significantly elevated only the ACL- group compared to contralateral controls (P=0.016 Fig. 3g).

Synovium pathology was similarly evaluated through macroscopic scoring, though no significant

differences were found between groups (Fig. 3h). A small subset of cartilage was evaluated

histologically, but there was high sample-to-sample variability and little discernable differences

between injury groups. (Supplementary Fig. 4). Overall, minimal cartilage degradation was

observed in PL- joints, and similar increases in cartilage T1ρ values (an early indicator of

cartilage degradation) were observed in AM- and ACL- joints; however, macroscopic cartilage

degeneration was elevated only in ACL- joints.

Fig. 3. Cartilage degeneration and synovium inflammation. a, Representative sagittal MR

images overlaid with T1ρ maps of articular cartilage. Interlimb differences in T1ρ value for b,

lateral femoral, c, medial femoral, d, lateral tibial, and e, medial tibial articular cartilage. #P<0.05

compared to 0 from one sample t-test; bars indicate P<0.05 between injury type, from one-way

ANOVA with Tukey post hoc test or Welch ANOVA with Dunnett post hoc test for data sets

with unequal variance. f, Representative images of articular cartilage and synovium, with

macroscopic scores shown in the corner of each image. White arrows indicate cartilage erosions

reaching the subchondral bone. g, Composite macroscopic OARSI scores summed from the four

articular cartilage compartments. *P<0.05 compared to contralateral controls from Wilcoxon

matched-pairs test. h, Histogram of macroscopic OARSI scores for synovium, presented as a

percent of samples for each group. Individual data points (n=12 for PL- and AM- groups, n=8 for

ACL- group) presented with mean ± 95% CI (b-e) or median ± IQR (g).

Subhead 4: Meniscus degeneration

The biomechanical effect of injury on the menisci of the knee joint was assessed by

measuring the in-situ load carried by each tissue under the applied loading conditions. The load

carried in the medial meniscus under applied anterior drawer was 33% greater for AM- joints

and 3.7-fold greater for ACL- joints compared to contralateral joints (P≤0.029 for each) (Fig.

4a), while no significant differences were measured under applied compression (Fig. 4b). The

load carried in the lateral meniscus was greater compared to contralateral joints for AM- joints

under applied anterior drawer (by 2.7-fold, P=0.008, Fig. 4c) and for ACL- joints under applied

compression (by 2.1-fold, P=0.045, Fig. 4d). No differences in meniscal loads were measured in

PL- joints relative to contralateral controls. Next, the volume of the menisci were measured from

MRI (Fig. 4d) and compared between injured and control joints. The medial and lateral menisci

were 11% and 7% larger in AM- joints and 28% and 30% larger in ACL- joints, respectively

(P≤0.048 for all, Fig. 4f-g). No changes in meniscal size were measured in PL- joints. For ACL-

joints, in addition to the measurable increases in size, the menisci qualitatively developed an

oblique shape, and medial menisci developed an unorganized fibrous posterior outgrowth,

evident macroscopically and on MRI (Supplementary Fig. 5). Overall, both in-situ loading and

volume of the menisci showed no change in PL- joints, moderate increases in AM- joints, and

substantial increases in ACL- joints.

Fig. 4. Changes in meniscal loading and size after partial and complete ACL transection.

In-situ forces in the medial meniscus under applied a, anterior drawer and b, axial compression.

In-situ forces in the lateral meniscus under applied c, anterior drawer and d, axial compression. e,

Representative 3D models of menisci from each injury group, generated from MRI scans. Medial

menisci shown in blue, lateral menisci in red. Volume of the f, medial meniscus and g, lateral

meniscus. Individual data points (n=12 for PL- and AM- groups, n=8 for ACL- group) presented

with mean. Lines connect paired joints. *P<0.05 between injured and contralateral, from paired

t-test with Holm-Sidak correction.

Subhead 5: Clinical implications and associations

To investigate whether findings were similar for partial injuries at different ages during

skeletal growth, AM and PL bundle injuries were created in adolescent (6-month-old) pigs, and

similar results were found (See Supplemental Results and Supplemental Figures 6-7). Using

pooled data from both age groups, changes in joint laxity following injury were associated with

degenerative changes in the cartilage. Linear regressions were run between interlimb differences

in APTT and degenerative outcomes (Fig. 5a-h, Supplementary Fig. 8). Across injury types,

significant positive associations existed between interlimb differences in APTT and mean side-

to-side differences in T1ρ of the lateral femoral cartilage (r^2=0.16, P=0.0089), lateral meniscus

volume (r^2=0.73, P<0.0001), and medial meniscus volume (r^2=0.47, P<0.0001) (Fig. 5a,e,

Supplementary Fig. 8). Within injury type, however, there were generally weaker associations

for lateral femoral cartilage T1ρ and the interlimb APTT difference (r^2=0.001-0.24) (Fig. 5b-d).

For medial meniscus volume, poor associations were also found between the changes in medial

meniscus volume and interlimb APTT increases within PL-injured (P=0.58) and ACL-injured

(P=0.42) joints. The strongest positive association was found between the medial meniscus

volume change and APTT in AM-injured joints, though again with an r^2 value of only 0.26

(P=0.046, Fig. 5f-h, Supplementary Fig. 8m-p). Similarly, no significant associations were

observed for changes in medial femoral cartilage T1ρ values or lateral meniscus volume and

changes in APTT for any injury group (Supplementary Fig. 8e-l).

To place the observed degenerative outcomes in a clinically relevant framework,

specimens were grouped by equivalent Lachman test grade (Fig. 5i-l, Supplementary Fig. 9),

which is a common clinical exam to assess joint stability after ACL injury. Generally, mean side-

side differences in lateral femoral cartilage T1ρ values (P=0.048) and medial (P=0.0002)

meniscus volume increased progressively for increasing Lachman test grades. Within grade,

significant interlimb differences were measured for lateral femoral cartilage T1ρ values only for

specimens with Lachman test grades C and D (P$\leq$0.0086), but significant increases in medial

meniscus volume were measured for specimens with grade B, C, and D (P$\leq$0.023 for each).

Linear regressions were performed within Lachman grades to further investigate associations

between joint stability and degeneration (Supplementary Fig. 9), but these were generally weak

(r²<=0.18). Pooling all injury types, these findings indicate that joint destabilization is positively

associated with the extent of joint degradation, but strong associations were not readily apparent

within an individual injury type or clinical test grade level.

Fig. 5. Degenerative outcomes as a function of changes in laxity. a-h, Linear regressions of

interlimb difference in APTT versus (a-d) lateral femoral joint cartilage T1ρ and (e-h) medial

meniscus volume, across injury groups and separated by PL-, AM-, and ACL-injured joints for

pooled juvenile and adolescent pigs. i,j, Lachman test grade from applied anterior-posterior

drawer in biomechanical testing used to separate interlimb difference in k, lateral femoral joint

cartilage T1ρ and l, medial meniscus volume. *P<0.05 compared to 0 from one sample t-test;

bars indicate P<0.05 between injury type, from one-way ANOVA with Tukey post hoc test.

Statistical results from linear regression analyses shown in graph. Individual data points (n=17

for PL- and AM- groups, n=8 for ACL- group) presented with mean ± 95% CI (k-l).

Discussion

There is presently a lack of data to inform clinical treatment of partial ACL injury in young patients, and large exploratory clinical studies are difficult to perform in this patient population. The present study provides a comprehensive analysis of functional and degenerative changes in the knee joint after partial (isolated AM or PL bundle) and complete ACL injury in skeletally immature subjects using a preclinical porcine model and clinically relevant outcome measures. Our results indicate that ACL injury severity (partial or complete) impacts outcomes, and equally importantly, that partial injury location (AM or PL bundle) also impacts outcomes. Across all injury types, the degree of degenerative changes was strongly associated with the extent of joint destabilization. Joint laxity increases were minimal for PL bundle injury, minor for AM bundle injury, and major for ACL injury. In both partial injury cases, the remaining bundle remodeled through an increase in CSA to attempt to restore ACL function and stabilize the knee joint. Cartilage degeneration and meniscal hypertrophy were observed to a minor degree after AM bundle injury and to a major degree after ACL injury, but not after PL bundle injury. Generally, sex did not impact degenerative outcomes in this model.

The joint laxity findings in this study are consistent with studies in humans, which report minor (0-6 mm) increases in laxity after partial ACL tears (*8, 15, 18, 20*) compared to major (6-12 mm) increases in laxity after complete ACL tears (*8, 15*). ACL injuries are commonly diagnosed on the field or in the office through physical examination using a Lachman test, which assesses tibial translation under anterior loads, or a pivot shift test, which evaluates tibial subluxation under compressive and rotational loads (*29*). Therefore, the small differences in laxity for partial ACL injuries point to challenges in diagnosis, particularly for PL bundle tears. This is supported by a clinical study in adults which found that 50% of partial tears confirmed

290 through arthroscopy were diagnosed as a false negative using an arthrometer designed to

291 quantify joint laxity (*30*). In the current study, no significant difference in side-to-side laxity was

292 measured under anterior or rotational loads for the PL bundle injury group. In fact, all specimens

293 from the PL bundle injury group fell within the Lachman grade A (indicative of a "normal"

294 joint), suggesting that many of these injuries may go undiagnosed and untreated. Imaging,

295 primarily MRI, can add value as a complementary diagnostic tool in cases where physical

296 examination is inconclusive. Therefore, physical exams paired with MRI appear to be the best

297 current process for diagnosing partial ACL tears, although there remains space for improvement

298 of diagnostic sensitivity and specificity (*15, 31*). Diagnostic accuracy has the potential for

299 improvement through implementing higher resolution MRI protocols or newer sequences

300 capable of better contrast for ligaments (*32*). There also remains the potential for development of

301 new clinical exams which can detect partial injuries.

302   Our functional data is consistent with clinical studies in adults which reported that a

303 greater number of PL bundle injuries were biomechanically functional than AM bundle injuries

304 (*15, 33-35*). Data is less definitive in pediatric patients, since currently only two studies evaluate

305 outcomes after partial injury in young patients (*18, 20*). Kocher et al. found patients with a PL

306 bundle injury were more likely to require a subsequent surgery (*20*), which seems contradictory

307 to findings in this study. However, survivor bias may be a source for this discrepancy. In that

308 study, patients with Lachman grade C or D underwent surgical reconstruction and were not

309 included, which would exclude more AM bundle injuries than PL bundle injury based on our

310 laxity findings. Additionally, it is possible that a greater percentage of AM bundle tears progress

311 to complete ACL tears before receiving treatment. Even without progression to a complete tear,

312 our data shows that degenerative changes can occur after partial ACL injury with joint

instability, suggesting that surgical intervention may be of value even in cases of minor joint

instability (Lachman Grade B or higher). Conversely, the lack of degenerative changes in partial

ACL injury with no joint instability could provide a rationale for non-operative treatment,

allowing more growth to occur prior to surgical intervention if needed.

Our findings are consistent with preclinical in-vivo studies of partial ACL injury as well.

A study in skeletally mature sheep found gross and histological cartilage degradation and

meniscal damage 40 weeks after AM bundle injury (*36*). Another study in skeletally mature

rabbits found histological evidence of meniscal degeneration 8 weeks after both AM and PL

bundle injuries (*37*). In adult rabbits, partial ("medial half") and complete ACL injuries led to an

increase in joint laxity and histological cartilage damage, and histological scoring was associated

with anterior joint stiffness (*38*). Although factors including skeletal maturity, injury location,

and time from injury differ between these studies and our study, the consistent findings show that

partial ACL injury alone can lead to long-term joint degeneration across different preclinical

small and large animal models. Furthermore, our study extends these prior findings to show that

degeneration occurs even in the context of skeletally immature joints, despite the potential for

higher healing/remodeling capacity.

Several animal studies have tried to associate joint instability and degeneration (*38-40*).

However, these studies often involve creating injuries to multiple ligaments (*39*) or stabilizing

the joint via surgery (*40*), which limits clinical relevance. Moreover, no study spans the full

spectrum of instability corresponding to Lachman grades A through D. This is crucial since

patients with grades A or B would be most likely to be treated non-surgically in humans.

Additionally, degenerative outcomes are often assessed via histology, which limits their clinical

translatability. In our work, we comprehensively compare different types of ACL injury that

create a full range of joint instability from Lachman grades A through D, allowing a

comprehensive association to degeneration. Additionally, we leverage clinically relevant

outcome measures (MRI) that could be used to motivate confirmatory studies in humans (though

we also performed traditional gross and histological scoring).

Along these lines, the cartilage degeneration that we measured macroscopically and from

MRI did not necessarily correspond to one another, particularly in the AM bundle injured group

which showed significantly elevated cartilage T1ρ values but little macroscopic cartilage

damage. Furthermore, histological scoring of the cartilage was highly variable and did not show

clear associations with macroscopic scoring or MRI T1ρ relaxation values. These metrics

inherently measure different outcomes, with macroscopic scoring reflecting large structural

damage and loss of tissue, T1ρ values reflecting the matrix composition of existing cartilage

tissue, and histological scoring reflecting local, 2D structural and cellular damage. While

elevated T1ρ values are often used as an early predictor of future degeneration (*41*), macroscopic

and histological scoring reflects later-stage structural degeneration. These metrics may capture

different phases of degeneration, and we would predict that macroscopic scores for the AM

bundle injury group would increase at a longer follow-up timepoint. The difference between

outcome metrics also highlights the limited 2D nature of histology (only looking at a thin slice)

and suggests that 3D metrics provide a clearer picture of overall cartilage degeneration.

In addition to cartilage, changes in the meniscus were also associated with joint

instability, suggesting meniscal hypertrophy may also be an early indicator of joint degeneration.

This is supported by data from a guineapig model, where meniscal hypertrophy was tied to

cartilage damage after ACL injury (*42*). Although little to no data exist regarding meniscal size

changes after ACL injury in humans, larger meniscus volume in middle-aged women was tied to

later development of structural OA (*43*). While it is common to evaluate meniscal tears after

ACL injury, meniscal size, independent of the presence of injury, is often overlooked. However,

meniscus size may be an important indicator of overall joint health. Furthermore, meniscus size

may be a more repeatable metric across time and would not require specialized MRI sequences,

such as T1ρ.

The current work has implications for treatment of partial ACL tears, since we showed

significant meniscal and chondral pathology after partial ACL injury at the equivalence of a

grade B or higher Lachman test. Although we saw differences in outcomes due to injury type, the

extent of destabilization may be a better metric as a basis for treatment, since we saw significant

associations between cartilage degeneration and joint laxity even within AM bundle injuries. Our

results suggest that surgical intervention may be appropriate for patients with Lachman grade B

or above, and more likely to be necessary in patients with an AM bundle tear rather than PL

bundle tear. Additionally, the increase in remaining bundle CSA with no change in T2* value

indicates that viable tissue capable of functional remodeling can remain after a partial ACL

injury. This points to advantages in surgical techniques which preserve the intact tissue. Single

bundle augmentation techniques which repair or reconstruct only the injured bundle are

relatively new but gaining in popularity (*44-48*). These techniques have the advantage of

preserving the functional remaining tissue and may cause less donor site morbidity since smaller

grafts would be required.

This study includes several limitations, including the use of an animal model. While the

pig is an established preclinical model for musculoskeletal studies (*23*) and the ACL bundles

function similarly between pigs and humans (*24*), there are some differences in knee anatomy

(*25*) and physiological loading compared to humans. Furthermore, human patients are prescribed

physical therapy and activity restrictions, while we did not attempt to restrict animal activity or

acquire metrics for physical activity. Moreover, our evaluation of joint stability does not account

for active stabilization provided by the muscles, so it is possible that the joint is more stable in-

vivo. Nevertheless, the degenerative changes we detected occurred in the presence of muscle

function in-vivo. Additionally, our surgical model of ACL injury does not account for damage to

other tissues that may occur during the high impact of a naturally occurring ACL injury.

However, our model does isolate the effects of progressive degeneration caused by the loss of

joint stability. The injured bundle or entire ACL was removed in our surgical model. This

prevents any healing, which has been observed in a limited capacity in clinical cases (*49*). An

additional limitation includes the use of contralateral limbs as a control. These may differ from

joints in healthy animals due to altered gait or limb compensation after injury of the ipsilateral

joint. Lastly, the conclusions in this study are limited to only one timepoint after injury and do

not capture the time course of degenerative changes. Future longitudinal studies to assess the

progression of the degenerative outcomes would be useful for distinguishing early predictors of

degeneration and identifying an ideal window for implementing treatment interventions to

prevent degenerative changes.

This work fills key gaps in knowledge about degenerative changes after partial and

complete ACL injury in skeletally immature subjects. We leverage a large animal model to

collect detailed data about joint biomechanics, tissue remodeling, and cartilage degeneration

which are not possible to collect in clinical studies. Our data show that the extent of

destabilization after an ACL injury may serve as a predictor of degeneration, irrespective of the

injury type (partial or complete) or partial injury location (AM or PL bundle). Associations in

this study also imply that there may not be a key threshold beyond which degenerative changes

occur, but instead, a continuous gradient where even small increases in laxity can cause minor

degeneration. Overall, this study provides a fundamental basis for informing clinical treatments

in young patients with partial ACL tears.

Methods

Subhead 1: Study design

The objective of this study was to evaluate the outcomes of partial ACL injury as a

function of tear location (AM or PL bundle), with covariates of sex and age, in a skeletally

immature surgical porcine model. All animals were bred at the North Carolina State University

Swine Educational Unit, and all experimental protocols were approved by the North Carolina

State University Institutional Animal Care and Use Committee. Sample sizes were calculated

using a power analysis described in the "Statistical analysis" section. A total of 32 juvenile

Yorkshire crossbreed pigs (age: 13 ± 1 week, weight: 34 ± 5 kg) underwent unilateral

arthroscopic transection of the AM bundle (n=6 female, n=6 male), the PL bundle (n=6 female,

n=6 male), or the complete ACL (n=4 female, n=4 male) with a sham operation performed on

the contralateral joint. Injury type and injured limb side were randomized. To confirm findings in

an older age group, a small subset of 9 adolescent pigs (age: 25 ± 1 week, weight: 129 ± 21 kg)

underwent unilateral arthroscopic transection of the AM bundle (n=3 female, n=1 male) or PL

bundle (n=4 female, n=1 male). One adolescent animal which was meant to undergo AM bundle

transection was excluded due to a pre-existing partial ACL tear to the PL bundle. The endpoint

was set at 24 weeks to experimentally evaluate joint biomechanics, cartilage health, and

remodeling of soft tissues in the knee joint. Data were collected and analyses were performed in

a blinded manner, where possible. No outliers were excluded in the presented data.

Subhead 2: Surgical procedure

Prior to surgery, all pigs were sedated and received analgesics and antimicrobials. Operations were performed arthroscopically (Arthrex Synergy System) under general anesthesia. The knee joints were accessed through medial and lateral parapatellar incisions (~0.5 cm long). The infrapatellar fat pad was partially removed using a 3.5 mm motorized shaver (SabreTooth, Arthrex) to visualize the ACL. A freer elevator was inserted between the AM and PL bundles near the tibial insertion and used to bluntly separate the bundles. A triangle knife (Ectra II Knife 4448, Smith & Nephew) was inserted into the separation and (1) turned anteriorly to transect the AM bundle, (2) turned posteriorly to transect the PL bundle, or (3) used to transect the full cross-section of the ACL. The remnant of the transected bundle or complete ligament was debrided using a shaver to ensure loss of function and prevent healing. In the contralateral sham-operated control limb, the infrapatellar pad was similarly partially removed. The ACL bundles were separated with a freer elevator but not transected. Following the operation, skin incisions were suture closed, a local analgesic was administered through subcutaneous injection at the portal sites, and the joint was covered with an antimicrobial adhesive drape. Animals were given analgesics and antimicrobials for 3 days post-operatively. Animals were allowed full weight bearing immediately following surgery and housed in pens with no restriction of movement. After 6 months (23.9 ± 0.3 weeks), all animals were euthanized, and hind limbs were collected. Skin and excess muscle were removed, and knee joints were wrapped in saline-soaked pads and stored at -20 °C.

Subhead 3: Magnetic resonance imaging

Prior to scanning, joints were thawed. Joints were imaged at full extension. Imaging was performed using a Siemens Magnetom Prisma 3T MRI system and a knee coil. Joints were

scanned using a T1ρ fast low-angle shot (FLASH) sequence (FOV: 128x128x84mm, voxel size:

0.4x0.4x1.5mm, FA: 10°, TR: 8.74ms, TE: 4.60ms, spin-lock frequency: 500 Hz, TSL: 0, 10, 20,

30, 40ms) and a T2 susceptibility weighted imaging (SWI) sequence (FOV: 160x160x128mm,

voxel size: 0.4x0.4x0.8mm, FA: 12°, TR: 31.0ms, TE: 3.3,7.1,11.4,15.7,20.0,24.4ms). Following

scanning, joints were stored at -20 °C for subsequent biomechanical testing. T1ρ and T2*

relaxation maps were created by fitting monoexponential decay function on a voxel-by-voxel

basis to the 5 spin-lock times and 6 echo times, respectively, using a custom MATLAB script.

Articular cartilage from the medial and lateral condyles of the femur and tibia were

manually segmented by a single user from T1ρ images with 0 ms spin-lock time using

commercial software (Simpleware ScanIP). Using a MATLAB script, the segmented areas were

transposed onto T1ρ relaxation maps described above, and the median T1ρ relaxation time was

calculated for each of the four segmented cartilage compartments.

Similarly, the AM bundle, PL bundle, complete ACL, and medial and lateral menisci

were manually segmented by a single user from T2* images with 15.7ms echo time. The median

T2* relaxation time was calculated for each ACL bundle. Additionally, the volume of each

segmented tissue was recorded, and the cross-sectional area (CSA) of the AM and PL bundles

and the complete ACL were calculated as previously described (*21, 22*). Briefly, each segmented

tissue was exported as a surface mesh, and analyzed using a MATLAB script. Each mesh was

rotated such that a line of best fit through the mesh vertices aligned with the z-axis. The CSA

was calculated for slices of the mesh in 1 mm increments along the z-axis, then averaged over

the middle 50% of slices.

Subhead 4: Biomechanical testing

Biomechanics of the knee (stifle) joint were evaluated using a 6 degree-of-freedom

(DOF) robotic testing system (KR300 R2500, KRC4, Kuka) equipped with a universal force-

moment sensor (Omega160 IP65, ATI Industrial Automation), and integrated through the

simVitro software knee module (Cleveland Clinic). To prepare for testing, joints were thawed at

room temperature and trimmed to expose the femoral and tibial diaphyses. The femoral head and

the distal end of the tibia were removed using a bone saw, then bones were fixed using

fiberglass-reinforced epoxy (Everglass, Evercoat) in molds designed to fit in custom clamps for

testing. Joints were again stored at -20 °C and thawed at room temperature one day prior to robot

testing.

The femoral end of the joint was fixed to a rigid testing platform, while the tibial end was

fixed to the robotic system at 40° of flexion. A 3D digitizer (G2X MicroScribe) was used to

determine an anatomic coordinate system. Joints were sprayed with saline and lightly wrapped in

a pad throughout testing to maintain hydration. A passive flexion path was determined by flexing

the joint from 40° to 90° while minimizing forces/moments in the remaining 5 DOFs. Then,

loads/torques were separately applied to the tibia in the anterior-posterior (AP) (100 N), proximal

(axial compression, 140 N), and VV (5 Nm) directions/rotations at 60° of joint flexion while

recording kinematic paths. Kinematics were then replayed while recording forces. Next, the joint

capsule was removed, and kinematics were replayed while recording forces. The difference in

forces measured between the intact and capsule-deficient condition was calculated as the force

carried by the joint capsule, by the principle of superposition (*50*). This process was repeated

after transection of the AM bundle (or any remnant), PL bundle (or any remnant), MCL, LCL,

PCL, medial meniscus, lateral meniscus, medial femoral condyle, and lateral femoral condyle.

Joint laxity was measured by anterior-posterior tibial translation (APTT) under AP drawer,

anterior tibial translation (ATT) under axial compression, and VV rotation under VV moments.

Load-transformation curves were plotted for each loading condition using data points collected at

intervals of 20% of the maximum applied load.

Subhead 5: Macroscopic scoring

During robotic testing and tissue dissection, gross images were taken of the synovium

following joint capsule dissection, as well as the femoral and tibial articular cartilage, following

removal of all other tissues, during robotic testing. Synovium and cartilage were macroscopically

scored using Osteoarthritis Research Society International (OARSI) guidelines for sheep and

goat (*28*). Although are no scoring guidelines for pigs, this scoring system has been used to

evaluate cartilage in previous porcine studies (*51, 52*). Synovial pathology was scored from 0

(normal) to 5 (severe), and articular cartilage of medial and lateral condyles of the femur and

tibia were scored from 0 (normal) to 4 (large erosions down to subchondral bone) by two blinded

reviewers. Disagreements between users were re-examined to determine a consensus. Scores

from the four cartilage compartments were added to give a total joint score.

Subhead 6: Histological analyses

Immediately following robotic testing, 5 mm coronal slabs were cut from the lateral

femoral condyle (1.5-2 cm from the femoral notch apex) and lateral tibial plateau (1.5-2 cm

anterior to the posterior boundary of the tibial plateau) using a bone saw. Samples were fixed in

4% paraformaldehyde solution at 4°C for 72 hours, then decalcified in Formical-2000 for 17

days. Samples were trimmed to 2-3mm in thickness and embedded in paraffin. 8 μm sections

were taken from each sample and stained with safranin-O/fast-green. Whole sections were

imaged at 4x magnification using an Olympus IX73 microscope with CellSens software. Two

blinded reviewers scored samples adapted from OARSI guidelines (*53*) in the following

categories: matrix staining, fibrillation, and focal cell loss.

Subhead 7: Statistical analyses

Sample sizes were determined using a power analysis based on prior work that

demonstrated the ability of transection of the AM bundle and ACL in cadaveric joints to generate

different levels of initial joint instability (*54*). Using an overall α level of 0.05 (accounting for

multiple comparisons), a power of 0.8, and an effect size of 1.5 for the AM- group, a sample size

of n=12 was selected to measure a difference between injured and sham-operated joints. For

complete loss of ACL function, much larger effect sizes are expected (>3), so a sample size of

n=8 was used for this group.

Normality of all data sets was confirmed using Shapiro-Wilk test, and equal variance

between data sets was tested using Brown-Forsythe test. Joint kinematics, tissue forces under

applied loads, tissue size metrics, and bundle T2* values were compared between injured and

sham-operated joints using paired t-tests with Holm-Sidak's correction for multiple comparisons.

Outliers were removed from lateral meniscus forces under anterior drawer using the iterative

Grubbs' method with an alpha of 0.05. Interlimb differences in joint kinematics, cartilage T1ρ,

and meniscal volumes were compared between injury types and ages of injury using a two-way

ANOVA with Holm-Sidak post hoc comparison. Interlimb differences in macroscopic and

histologic cartilage scores were compared between AM-injury and PL-injured joints in juvenile

and adolescent pigs with Mann-Whitney tests. For data sets with unequal variance, Welch

ANOVA with Dunnett post hoc comparison was used. To evaluate clinically relevant outcomes,

samples were grouped into equivalent Lachman test grades according to IKDC guidelines (A: <3

mm, B: 3-5 mm, C: 5-10 mm, D: >10 mm) (*55*), based on joint translations under AP drawer.

Interlimb differences in cartilage T1ρ, macroscopic cartilage scores, and meniscal volume were

compared between Lachman grades using one-way ANOVA and compared to zero with one

sample t-test. Associations were assessed between interlimb differences in APTT and interlimb

differences in cartilage T1ρ and meniscal volume using linear regression across all injury types

and within injury types. All statistical analyses were performed in Prism (GraphPad), with

overall significance set at α=0.05 for all analyses with corrections for multiple comparisons done

as needed. Adjusted P values are reported. Data presented as mean ± 95% CI, unless stated

otherwise, and data are reported in supplemental material.

Data availability

The main data supporting the results in this study are available within the paper and its

Supplementary Information.

References

1.   G. Kolata, If You Tear a Knee Ligament, Arthritis Is Likely to Follow in 10 YearsN. Y.
Times (2017) (available at https://www.nytimes.com/2017/11/06/health/arthritis-risk-
acl.html).

2.   T. L. Sanders, H. Maradit Kremers, A. J. Bryan, D. R. Larson, D. L. Dahm, B. A. Levy,
M. J. Stuart, A. J. Krych, Incidence of Anterior Cruciate Ligament Tears and
Reconstruction: A 21-Year Population-Based Study. *Am J Sports Med* **44**, 1502-1507
(2016).

3.   N. A. Mall, P. N. Chalmers, M. Moric, M. J. Tanaka, B. J. Cole, B. R. Bach, Jr., G. A.
Paletta, Jr., Incidence and trends of anterior cruciate ligament reconstruction in the
United States. *Am J Sports Med* **42**, 2363-2370 (2014).

4.   B. C. Werner, S. Yang, A. M. Looney, F. W. Gwathmey, Jr., Trends in Pediatric and
Adolescent Anterior Cruciate Ligament Injury and Reconstruction. *J Pediatr Orthop* **36**,
447-452 (2016).

5.   D. Zbrojkiewicz, C. Vertullo, J. E. Grayson, Increasing rates of anterior cruciate ligament
569   reconstruction in young Australians, 2000-2015. *Med J Aust* **208**, 354-358 (2018).

570   6.   E. R. Dodwell, L. E. Lamont, D. W. Green, T. J. Pan, R. G. Marx, S. Lyman, 20 years of
571   pediatric anterior cruciate ligament reconstruction in New York State. *Am J Sports Med*
572   **42**, 675-680 (2014).

573   7.   D. Dare, S. Rodeo, Mechanisms of post-traumatic osteoarthritis after ACL injury. *Curr*
574   *Rheumatol Rep* **16**, 448 (2014).

575   8.   D. M. Daniel, M. L. Stone, R. Sachs, L. Malcom, Instrumented measurement of anterior
576   knee laxity in patients with acute anterior cruciate ligament disruption. *Am J Sports Med*
577   **13**, 401-407 (1985).

578   9.   I. Kawashima, H. Hiraiwa, S. Ishizuka, R. Kawai, Y. Kusaka, K. Ohtomo, T. Tsukahara,
579   Incidence of Medial and Lateral Meniscal Tears After Delayed Anterior Cruciate
580   Ligament Reconstruction in Pediatric Patients. *Orthop J Sports Med* **8**,
581   2325967120964603 (2020).

582   10.   P. Neuman, I. Kostogiannis, T. Friden, H. Roos, L. E. Dahlberg, M. Englund, Knee laxity
583   after complete anterior cruciate ligament tear: a prospective study over 15 years. *Scand J*
584   *Med Sci Sports* **22**, 156-163 (2012).

585   11.   P. D. Fabricant, M. S. Kocher, Management of ACL Injuries in Children and
586   Adolescents. *J Bone Joint Surg Am* **99**, 600-612 (2017).

587   12.   H. Mizuta, K. Kubota, M. Shiraishi, Y. Otsuka, N. Nagamoto, K. Takagi, The
588   conservative treatment of complete tears of the anterior cruciate ligament in skeletally
589   immature patients. *J Bone Joint Surg Br* **77**, 890-894 (1995).

590   13.   C. L. Ardern, G. Ekas, H. Grindem, H. Moksnes, A. Anderson, F. Chotel, M. Cohen, M.
591   Forssblad, T. J. Ganley, J. A. Feller, J. Karlsson, M. S. Kocher, R. F. LaPrade, M.
592   McNamee, B. Mandelbaum, L. Micheli, N. Mohtadi, B. Reider, J. Roe, R. Seil, R.
593   Siebold, H. J. Silvers-Granelli, T. Soligard, E. Witvrouw, L. Engebretsen, 2018
594   International Olympic Committee consensus statement on prevention, diagnosis and
595   management of paediatric anterior cruciate ligament (ACL) injuries. *Knee Surg Sports*
596   *Traumatol Arthrosc*, (2018).

597   14.   P. Colombet, D. Dejour, J. C. Panisset, R. Siebold, S. French Arthroscopy, Current
598   concept of partial anterior cruciate ligament ruptures. *Orthop Traumatol Surg Res* **96**,
599   S109-118 (2010).

600   15.   D. Dejour, P. G. Ntagiopoulos, P. R. Saggin, J. C. Panisset, The diagnostic value of
601   clinical tests, magnetic resonance imaging, and instrumented laxity in the differentiation
602   of complete versus partial anterior cruciate ligament tears. *Arthroscopy* **29**, 491-499
603   (2013).

16.   E. R. Hunt, C. A. Jacobs, C. E. Conley, M. L. Ireland, D. L. Johnson, C. Lattermann,
Anterior cruciate ligament reconstruction reinitiates an inflammatory and
chondrodegenerative process in the knee joint. *J Orthop Res* **39**, 1281-1288 (2021).

17.   P. N. Chalmers, N. A. Mall, M. Moric, S. L. Sherman, G. P. Paletta, B. J. Cole, B. R.
Bach, Jr., Does ACL reconstruction alter natural history?: A systematic literature review
of long-term outcomes. *J Bone Joint Surg Am* **96**, 292-300 (2014).

18.   J. M. Fayard, B. Sonnery-Cottet, G. Vrgoc, P. O'Loughlin, G. D. de Mont Marin, B.
Freychet, T. D. Vieira, M. Thaunat, Incidence and Risk Factors for a Partial Anterior
Cruciate Ligament Tear Progressing to a Complete Tear After Nonoperative Treatment in
Patients Younger Than 30 Years. *Orthop J Sports Med* **7**, 2325967119856624 (2019).

19.   M. Group, J. S. Everhart, M. H. Jones, S. Yalcin, E. K. Reinke, L. J. Huston, J. T.
Andrish, C. L. Cox, D. C. Flanigan, C. C. Kaeding, R. A. Magnussen, N. Obuchowski, R.
D. Parker, A. D. Pedroza, R. A. Sanders, C. S. Winalski, K. P. Spindler, The Clinical
Radiographic Incidence of Posttraumatic Osteoarthritis 10 Years After Anterior Cruciate
Ligament Reconstruction: Data From the MOON Nested Cohort. *Am J Sports Med* **49**,
1251-1261 (2021).

20.   M. S. Kocher, L. J. Micheli, D. Zurakowski, A. Luke, Partial tears of the anterior cruciate
ligament in children and adolescents. *Am J Sports Med* **30**, 697-703 (2002).

21.   S. G. Cone, E. P. Lambeth, H. Ru, L. A. Fordham, J. A. Piedrahita, J. T. Spang, M. B.
Fisher, Biomechanical Function and Size of the Anteromedial and Posterolateral Bundles
of the ACL Change Differently with Skeletal Growth in the Pig Model. *Clin Orthop*
Relat Res **477**, 2161-2174 (2019).

22.   D. Howe, S. G. Cone, J. A. Piedrahita, B. Collins, L. A. Fordham, E. H. Griffith, J. T.
Spang, M. B. Fisher, Sex-specific biomechanics and morphology of the anterior cruciate
ligament during skeletal growth in a porcine model. *J Orthop Res*, (2021).

23.   S. G. Cone, P. B. Warren, M. B. Fisher, Rise of the Pigs: Utilization of the Porcine Model
to Study Musculoskeletal Biomechanics and Tissue Engineering During Skeletal Growth.
Tissue Eng Part C Methods **23**, 763-780 (2017).

24.   J. W. Xerogeanes, R. J. Fox, Y. Takeda, H. S. Kim, Y. Ishibashi, G. J. Carlin, S. L. Woo,
A functional comparison of animal anterior cruciate ligament models to the human
anterior cruciate ligament. *Ann Biomed Eng* **26**, 345-352 (1998).

25.   B. L. Proffen, M. McElfresh, B. C. Fleming, M. M. Murray, A comparative anatomical
study of the human knee and six animal species. *Knee* **19**, 493-499 (2012).

26.   R. R. Regatte, S. V. Akella, A. J. Wheaton, G. Lech, A. Borthakur, J. B. Kneeland, R.
Reddy, 3D-T1rho-relaxation mapping of articular cartilage: in vivo assessment of early
degenerative changes in symptomatic osteoarthritic subjects. *Acad Radiol* **11**, 741-749
(2004).

27. S. V. Akella, R. R. Regatte, A. J. Gougoutas, A. Borthakur, E. M. Shapiro, J. B. Kneeland, J. S. Leigh, R. Reddy, Proteoglycan-induced changes in T1rho-relaxation of articular cartilage at 4T. *Magn Reson Med* **46**, 419-423 (2001).

28. C. B. Little, M. M. Smith, M. A. Cake, R. A. Read, M. J. Murphy, F. P. Barry, The OARSI histopathology initiative - recommendations for histological assessments of osteoarthritis in sheep and goats. *Osteoarthritis Cartilage* **18 Suppl 3**, S80-92 (2010).

29. V. Musahl, I. D. Engler, E. M. Nazzal, J. F. Dalton, G. A. Lucidi, J. D. Hughes, S. Zaffagnini, F. Della Villa, J. J. Irrgang, F. H. Fu, J. Karlsson, Current trends in the anterior cruciate ligament part II: evaluation, surgical technique, prevention, and rehabilitation. *Knee Surg Sports Traumatol Arthrosc* **30**, 34-51 (2022).

30. J. Isberg, E. Faxen, S. Brandsson, B. I. Eriksson, J. Karrholm, J. Karlsson, KT-1000 records smaller side-to-side differences than radiostereometric analysis before and after an ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* **14**, 529-535 (2006).

31. P. Van Dyck, E. De Smet, J. Veryser, V. Lambrecht, J. L. Gielen, F. M. Vanhoenacker, L. Dossche, P. M. Parizel, Partial tear of the anterior cruciate ligament of the knee: injury patterns on MR imaging. *Knee Surg Sports Traumatol Arthrosc* **20**, 256-261 (2012).

32. N. C. Nacey, M. G. Geeslin, G. W. Miller, J. L. Pierce, Magnetic resonance imaging of the knee: An overview and update of conventional and state of the art imaging. *J Magn Reson Imaging* **45**, 1257-1275 (2017).

33. P. S. Christel, U. Akgun, T. Yasar, M. Karahan, B. Demirel, The contribution of each anterior cruciate ligament bundle to the Lachman test: a cadaver investigation. *J Bone Joint Surg Br* **94**, 68-74 (2012).

34. R. L. Hole, D. M. Lintner, E. Kamaric, J. B. Moseley, Increased tibial translation after partial sectioning of the anterior cruciate ligament. The posterolateral bundle. *Am J Sports Med* **24**, 556-560 (1996).

35. R. Siebold, F. H. Fu, Assessment and augmentation of symptomatic anteromedial or posterolateral bundle tears of the anterior cruciate ligament. *Arthroscopy* **24**, 1289-1298 (2008).

36. K. I. Barton, B. J. Heard, J. L. Sevick, C. R. Martin, S. M. M. Shekarforoush, M. Chung, Y. Achari, C. B. Frank, N. G. Shrive, D. A. Hart, Posttraumatic Osteoarthritis Development and Progression in an Ovine Model of Partial Anterior Cruciate Ligament Transection and Effect of Repeated Intra-articular Methylprednisolone Acetate Injections on Early Disease. *Am J Sports Med* **46**, 1596-1605 (2018).

37. D. Wang, Z. Wang, M. Li, S. Xu, The underlying mechanism of partial anterior cruciate ligament injuries to the meniscus degeneration of knee joint in rabbit models. *J Orthop Surg Res* **15**, 428 (2020).

38. Y. Tochigi, T. Vaseenon, A. D. Heiner, D. C. Fredericks, J. A. Martin, M. J. Rudert, S. L. Hillis, T. D. Brown, T. O. McKinley, Instability dependency of osteoarthritis development in a rabbit model of graded anterior cruciate ligament transection. *J Bone Joint Surg Am* **93**, 640-647 (2011).

39. C. B. Frank, J. E. Beveridge, K. D. Huebner, B. J. Heard, J. E. Tapper, E. J. O'Brien, N. G. Shrive, Complete ACL/MCL deficiency induces variable degrees of instability in sheep with specific kinematic abnormalities correlating with degrees of early osteoarthritis. *J Orthop Res* **30**, 384-392 (2012).

40. K. Murata, N. Kanemura, T. Kokubun, T. Fujino, Y. Morishita, K. Onitsuka, S. Fujiwara, A. Nakajima, D. Shimizu, K. Takayanagi, Controlling joint instability delays the degeneration of articular cartilage in a rat model. *Osteoarthritis Cartilage* **25**, 297-308 (2017).

41. H. F. Atkinson, T. B. Birmingham, R. F. Moyer, D. Yacoub, L. E. Kanko, D. M. Bryant, J. D. Thiessen, R. T. Thompson, MRI T2 and T1rho relaxation in patients at risk for knee osteoarthritis: a systematic review and meta-analysis. *BMC Musculoskelet Disord* **20**, 182 (2019).

42. G. Du, H. Zhan, D. Ding, S. Wang, X. Wei, F. Wei, J. Zhang, B. Bilgen, A. M. Reginato, B. C. Fleming, J. Deng, L. Wei, Abnormal Mechanical Loading Induces Cartilage Degeneration by Accelerating Meniscus Hypertrophy and Mineralization After ACL Injuries In Vivo. *Am J Sports Med* **44**, 652-663 (2016).

43. D. Xu, J. van der Voet, J. H. Waarsing, E. H. Oei, S. Klein, M. Englund, F. Zhang, S. Bierma-Zeinstra, J. Runhaar, Are changes in meniscus volume and extrusion associated to knee osteoarthritis development? A structural equation model. *Osteoarthritis Cartilage* **29**, 1426-1431 (2021).

44. B. Sonnery-Cottet, R. Zayni, J. Conteduca, P. Archbold, T. Prost, Y. Carrillon, J. Clechet, M. Thaunat, Posterolateral Bundle Reconstruction With Anteromedial Bundle Remnant Preservation in ACL Tears: Clinical and MRI Evaluation of 39 Patients With 24-Month Follow-up. *Orthop J Sports Med* **1**, 2325967113501624 (2013).

45. N. Shohat, D. Lindner, E. Tamir, Y. Beer, G. Agar, Prospective Comparison between Remnant-Preserving Augmentation and Double-Bundle Reconstruction in Anterior Cruciate Ligament Tears. *Isr Med Assoc J* **19**, 355-359 (2017).

46. A. M. Gipsman, N. Trasolini, G. F. R. Hatch, 3rd, Primary Anterior Cruciate Ligament Single-Bundle Repair With Augmentation for a Partial Anterior Cruciate Ligament Tear. *Arthrosc Tech* **7**, e367-e372 (2018).

47. G. Y. Ahn, T. H. Lee, K. J. Lee, S. Woo, Comparison of clinical outcomes and second-look arthroscopic evaluations between anterior cruciate ligament anteromedial bundle augmentation and single-bundle anterior cruciate ligament reconstruction. *Knee Surg Relat Res* **32**, 45 (2020).

715   48.   T. Tajima, N. Yamaguchi, Y. Morita, T. Yokoe, M. Nagasawa, T. Ota, S. Kawagoe, Y.
716   Nakamura, E. Chosa, Remnant-Preserved Single-Bundle Anterior Cruciate Ligament
717   Augmentation in Multi-ligamentous Knee Reconstruction. *J Knee Surg*, (2022).

718   49.   E. H. Crain, D. C. Fithian, E. W. Paxton, W. F. Luetzow, Variation in anterior cruciate
719   ligament scar pattern: does the scar pattern affect anterior laxity in anterior cruciate
720   ligament-deficient knees? *Arthroscopy* **21**, 19-24 (2005).

721   50.   G. A. Livesay, H. Fujie, S. Kashiwaguchi, D. A. Morrow, F. H. Fu, S. L. Woo,
722   Determination of the in situ forces and force distribution within the human anterior
723   cruciate ligament. *Ann Biomed Eng* **23**, 467-474 (1995).

724   51.   N. P. Karamchedu, M. M. Murray, J. T. Sieker, B. L. Proffen, G. Portilla, M. Q. Costa, J.
725   Molino, B. C. Fleming, Bridge-Enhanced Anterior Cruciate Ligament Repair Leads to
726   Greater Limb Asymmetry and Less Cartilage Damage Than Untreated ACL Transection
727   or ACL Reconstruction in the Porcine Model. *Am J Sports Med* **49**, 667-674 (2021).

728   52.   M. P. Pinette, J. Molino, B. L. Proffen, M. M. Murray, B. C. Fleming, Effects of Male
729   and Female Sex on the Development of Posttraumatic Osteoarthritis in the Porcine Knee
730   After Anterior Cruciate Ligament Surgery. *Am J Sports Med*, 3635465221102118 (2022).

731   53.   C. W. McIlwraith, D. D. Frisbie, C. E. Kawcak, C. J. Fuller, M. Hurtig, A. Cruz, The
732   OARSI histopathology initiative - recommendations for histological assessments of
733   osteoarthritis in the horse. *Osteoarthritis Cartilage* **18 Suppl 3**, S93-105 (2010).

734   54.   D. Howe, S. G. Cone, J. A. Piedrahita, J. T. Spang, M. B. Fisher, Age and Sex-Specific
735   Joint Biomechanics in Response to Partial and Complete ACL Injury in the Porcine
736   Model. *J Athl Train*, (2021).

737   55.   E. P. Mulligan, D. Q. McGuffie, K. Coyner, M. Khazzam, The reliability and diagnostic
738   accuracy of assessing the translation endpoint during the lachman test. *Int J Sports Phys*
739   *Ther* **10**, 52-61 (2015).

741 **Acknowledgments**

742 We would like to thank Laboratory Animal Resources (NC State) and the Biomedical Research

743 Imaging Center (UNC-CH) for their contributions to this work. This work was supported by the

744 National Institutes of Health (NIH) grants F31AR077997 and R01AR071985, and the National

745 Science Foundation (NSF) grant DGE-1746939.

Author contributions

D.H., L.V.S., J.T.S. and M.B.F. conceptualized the work. D.H., J.D.T., S.D.T., M.E., E.H.G., L.V.S., J.T.S. and M.B.F. designed the methodology. D.H., J.D.T., S.D.T., M.E., O.B., L.V.S., J.T.S. and M.B.F. investigated the work. Formal analysis was carried out by D.H., J.D.T., S.D.T., M.E., E.H.G. and M.B.F. Data was curated by D.H., J.D.T., S.D.T., and M.E. D.H., J.D.T., M.E. and M.B.F. created the figures for this work. Funding was acquired by D.H., J.D.T., L.V.S., J.T.S. and M.B.F., and the project administration was D.H. and M.B.F. with M.B.F. providing supervision. D.H. wrote the original draft and all authors reviewed and edited the paper.

Competing interests

Authors declare that they have no competing interests.

Supplementary Fig. 1. Sex-specific biomechanical outcomes. Interlimb differences in (A) anterior-posterior tibial

translation (APTT) under applied AP drawer, (B) anterior tibial translation (ATT) under applied axial compression,

and (C) varus-valgus (VV) rotation under applied VV moments. Individual data points presented with mean ± 95%

CI. * P<0.05 between male and female, from one-way ANOVA with Tukey post hoc test.

Supplementary Fig. 2. In-situ bundle forces under rotational loads. Resultant force in the uninjured bundle

under applied (A) valgus moments and (B) varus moments. Lines connect paired joints. Individual data points

presented with mean.

Supplementary Fig. 3. T1ρ values and macroscopic scores of cartilage compartments. T1ρ values of the (A)

lateral femoral, (B) lateral tibial, (C) medial femoral, and (D) medial tibial cartilage compartments. Macroscopic

OARSI score of the (E) lateral femoral, (F) lateral tibial, (G) medial femoral, and (H) medial tibial cartilage

compartments. Individual data points presented with mean with lines connecting paired joints (A-D) or median ±

IQR (E-H). *P<0.05 between injured and contralateral, from paired t-test with Holm-Sidak correction.

Supplementary Fig. 4. Histological scoring of lateral femoral and tibial cartilage samples stained with

safranin-O and fast green. Representative images taken from samples in sham-operated, PL-, AM-, and ACL-

joints. Summed composite histological scores shown for the (A) lateral femoral cartilage and (B) lateral tibial

cartilage. Individual data points presented with median ± IQR.

Supplementary Fig. 5. MRI and macroscopic images of medial menisci from sham-operated and ACL- joints.

(A,C) Sham-operated control shows normal anatomy of the medial meniscus, where (B,D) red dotted lines and

arrows indicate degenerative, fibrous outgrowth in ACL-injured joints. Anterior-posterior and medial-lateral

directions indicated.

Supplementary Results & Discussion: Extension to adolescent animals

To investigate whether findings were similar for partial injuries at different ages during skeletal growth, AM and PL bundle injuries were created in adolescent (6-month-old) pigs. Joint biomechanics, cartilage health, and meniscus volume were evaluated after 24 weeks. Interlimb differences in joint laxity were similarly elevated for AM- joints under AP drawer (P<0.05) in both the adolescent and juvenile injury groups, with no significant difference in laxity for the PL-group (Supplementary Fig. 6A, Fig. S6A). Laxity results were also similar between the two age groups under varus-valgus torque and compression for the PL-injured joints (Fig. S6B-C and Fig. 1). Both juvenile and adolescent age groups had similar increases of ~1-2 mm in laxity under compression for AM-injured joints, yet the adolescent group had a statistically significant increase in laxity where the juvenile group did not (Fig. S6B and Fig. 1). For both age groups, lateral and medial femoral cartilage T1ρ values and composite macroscopic cartilage scores appeared similarly elevated in AM- joints but not PL- joints, with the medial femoral cartilage T1ρ having statistically significantly higher (P<0.05) T1ρ values for AM-injured joints for both juvenile and adolescent age groups (Fig. 5B-D and Fig. S6F-J). Lastly, the interlimb differences in medial and lateral meniscus volume were elevated ~5-10% after partial injury in adolescent animals similar to juvenile animals (Fig. S6D and E). Although sample size was limited for adolescent animals (n=4-5 per injury type), these data highlight similarities across two age groups, adding confidence to our findings in younger animals with larger sample sizes.

Supplementary Fig. 6. Joint laxity and tissue remodeling after partial ACL injury created in adolescent

compared to juvenile pigs. (A) Interlimb differences in APTT under applied AP drawer. (B) Interlimb differences

in T1ρ value for lateral femoral articular cartilage and (C) medial femoral articular cartilage. (D) Interlimb

differences in macroscopic cartilage. Individual data points (n=5 for PL- and n=4 for AM- adolescent groups, n=12

for PL- and AM- juvenile groups) presented with mean ± 95% CI (A,B,D,E) or median ± IQR (C). *P<0.05 between

PL- and AM- injured joints, from two-way ANOVA with Holm-Sidak correction. Multiple comparisons were run

using Mann-Whitney test for interlimb differences in macroscopic scoring between PL- and AM- injured joints for

juvenile and adolescent animals.

Supplementary Fig. 7. Biomechanical and degenerative outcomes after partial injuries created in adolescent

(6 month) pigs. (A) APTT under applied AP drawer, (B) ATT under applied axial compression, and (C) VV

rotation under applied VV moments. Volume of the (D) medial meniscus and (E) lateral meniscus. (F) Composite

macroscopic OARSI scores summed from the four articular cartilage compartments. T1ρ value for (G) lateral

femoral, (H) medial femoral, (I) lateral tibial, and (J) medial tibial articular cartilage. Lines connect paired joints.

Supplementary Fig. 8. Interlimb difference in (A) lateral and (E) medial femoral cartilage T1rho values and (I)

lateral and (M) medial meniscus volumes versus interlimb difference in APTT across all injury groups in pooled

juvenile and adolescent pigs. Injury groups separated by PL-, AM-, and ACL- for interlimb differences in (B-D)

lateral femoral cartilage T1rho values, (F-H) medial femoral cartilage T1rho values, (J-L) lateral meniscus volume,

and (N-P) medial meniscus volume. Statistical results from linear regression analyses shown in graph.

Supplementary Fig. 9. Interlimb difference in (A) lateral and (F) medial femoral cartilage T1rho values and (K)

lateral and (P) medial meniscus volumes, grouped by the equivalent Lachman test grade in pooled juvenile and

adolescent pigs. *P<0.05 compared to 0 from one sample t-test; bars indicate P<0.05 between injury type, from one-

way ANOVA with Tukey post hoc test. Injury groups separated by Lachman grades A-D for interlimb differences in

(B-E) lateral femoral cartilage T1rho values, (G-J) medial femoral cartilage T1rho values, (L-O) lateral meniscus

volume, and (Q-T) medial meniscus volume. Statistical results from linear regression analyses shown in graph.

Individual data points presented with mean ± 95% CI (A,F,K,P).