Decoding the variant-to-function relationship for *LIPA*, a risk locus for coronary artery disease

Fang Li,[1*] Elise Flynn, [2,3] Philip Ha, [1] Mazal N. Zebak, [1] Haoxiang Cheng, [4] Chenyi Xue, [1] Jianting Shi, [1] Xun Wu, [1] Ziyi Wang, [1] Yujiao Meng, [1] Jian Cui, [1] Yizhou Zhu, [5] Annie Rozenblyum, [1] Jeana Chun, [1] Antonio Hernandez-Ono, [6] Ali Javaheri, [7,8] Babak Razani, [9] Marit Westerterp, [10] Robert C Bauer, [1] Yousin Suh, [5] Ke Hao, [4] Tuuli Lappalainen, [2,3,11] Hanrui Zhang, [1*]

1. Cardiometabolic Genomics Program, Division of Cardiology, Department of Medicine, Columbia University Irving Medical Center, New York, NY, USA

2. New York Genome Center, New York, NY, USA

3. Department of Systems Biology at Columbia University, New York, NY, USA

4. Department of Genetic and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA

5. Department of Obstetrics and Gynecology, Department of Genetics and Development, Columbia University Irving Medical Center, New York, NY, USA

6. Department of Medicine, Vagelos College of Physicians and Surgeons, Columbia University, New York, NY, USA

7. Cardiovascular Division, Washington University School of Medicine, St Louis, MO, USA

8. John Cochran Veterans Affairs Hospital, St Louis, MO, USA

9. Division of Cardiology and Vascular Medicine Institute, the Department of Medicine, and the Department of Veterans Affairs, University of Pittsburgh, Pittsburgh, PA, USA

10. Department of Pediatrics, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands

11. Department of Gene Technology, Science for Life Laboratory, Royal Institute of Technology, Stockholm, Sweden

* Corresponding authors:

Fang Li PhD

Cardiometabolic Genomics Program,

Division of Cardiology

Department of Medicine

Columbia University Irving Medical Center

630 West 168th Street, P&S10-401

New York, NY 10032

Tel: (212) 305-3508

Email: fl2532@cumc.columbia.edu

Hanrui Zhang MB PhD

Cardiometabolic Genomics Program,

Division of Cardiology

Department of Medicine

Columbia University Irving Medical Center

630 West 168th Street, P&S10-401

New York, NY 10032

Tel: (212) 304-7905

Email: hz2418@cumc.columbia.edu

Conflicts-of-interest statement

Graphical Abstract

Abstract

Translating human genomic discoveries into mechanistic insights requires linking genetic variations to candidate genes and their causal functional phenotypes. Genome-wide association studies have consistently identified *LIPA* as a risk locus for coronary artery disease (CAD), with previous expression quantitative trait loci (eQTL) analyses prioritizing *LIPA* as a candidate causal gene. However, functional studies elucidating the causal variants, regulatory mechanisms, target cell types, and their causal impact on atherosclerosis have been lacking. To address this gap, we applied functional genomics and experimental mouse models to establish the variant-to-function relationship at the *LIPA* locus. Our findings show that CAD risk alleles in the *LIPA* locus increase LIPA expression and enzyme activity specifically in monocytes/macrophages by enhancing PU.1 binding to an intronic enhancer region that interacts with the *LIPA* promoter. In myeloid *Lipa*-overexpressing mice, we observed larger atherosclerotic lesions accompanied by altered macrophage function, including increased macrophage accumulation due to enhanced monocyte recruitment, reduced neutral lipid accumulation, and upregulation of integrin and extracellular matrix pathways. Our work establishes a direct causal link between *LIPA* risk alleles and increased monocyte/macrophage LIPA that exacerbates atherosclerosis, bridging human functional genomic evidence to the mechanistic understanding of CAD.

Introduction

Coronary artery disease (CAD) remains the leading cause of death worldwide. The inheritability of CAD has been estimated to be 40% to 60%, supporting the contribution of inherited genetic variation to its pathogenesis (1). Genome-wide association studies (GWASs) determine the common genetic variants associated with diseases and traits, having collectively identified more than 300 CAD-associated risk loci (2-13). Identifying the causal genes that connect genetic variation to disease risk will inform novel biological mechanisms and therapeutic targets for CAD.

The *LIPA* locus has been reproducibly identified as a risk locus for CAD in multiple GWASs and meta-analyses across ethnic groups (**Table S1**) (2-13), but not plasma lipid traits (14), indicating a CAD-specific yet lipid metabolism-independent genetic mechanism. Colocalization analysis suggests that genetic variants associated with CAD are also associated with *LIPA* expression in the Stockholm-Tartu Atherosclerosis Reverse Networks Engineering Task study (STARNET) dataset (15), suggesting *LIPA* as the candidate causal gene at the locus. The *LIPA* gene encodes lysosomal acid lipase (also known as LAL), the only known acidic lipase hydrolyzing cholesteryl ester and triglyceride within lysosomes (16). Rare coding and splicing junction mutations in *LIPA* result in Mendelian disorders, including the infant-onset Wolman Disease due to a complete loss of LIPA and the later-onset Cholesteryl Ester Storage Disease (CESD) with up to 10% residual enzyme activity (16). CESD is characterized by hyperlipidemia, hepatosplenomegaly, and premature atherosclerosis, presumably due to severe dyslipidemia (17). Mice with whole body knockout of *Lipa* phenotypically resemble the

late-onset CESD in humans despite the complete loss of *Lipa*, and die within days when on an *Ldlr*$^{-/-}$ background and challenged with dietary cholesterol (18). However, almost all common CAD variants at the *LIPA* locus, including the lead single nucleotide polymorphism (SNP) rs1412444, are intronic, suggesting that they likely exert *cis-regulatory* effects rather than impacting protein function directly. The sole exonic SNP in this linkage disequilibrium (LD) block, i.e. rs1051338, does not alter the expression, activity, lysosomal trafficking, or secretion of LIPA, further implicating that these noncoding variants likely contribute to CAD risk by modulating gene regulation via *cis-regulatory* effects (19).

The SNP rs1412444, and other SNPs in high LD, showed robust quantitative trait locus (QTL) signals in monocytes, with risk alleles of CAD linked to higher *LIPA* mRNA (2, 3), and more importantly, higher enzyme activity (19) in human peripheral blood monocytes. These findings support that CAD risk alleles are associated with higher LIPA activity in monocytes, which may be causally involved in CAD. Pathologically, monocytes play essential roles in the development and exacerbation of atherosclerosis. In atherosclerosis, plaque macrophages are largely derived from the infiltration of circulating monocytes (20). Monocytes and macrophages may therefore represent the most relevant cell types for the regulatory effects of CAD *LIPA* variants.

Despite strong statistical association and initial QTL studies supporting the directionality and the potential causal cell types, the variant-to-function relationship and how this important locus causally impacts atherosclerosis are unclear. Herein, we sought to

delineate the causal variants, the effector gene, the cell types involved, and the regulatory mechanisms of this CAD risk locus; and to elucidate how elevated LIPA in these key causal cell types, as implicated by human functional genomic evidence, contributes to atherosclerosis pathogenesis. This work connects CAD-associated *LIPA* genetic variants to their biological functional, advancing our understanding of this locus's causal role in elevating CAD risk.

Results

Integrative genomic analyses identify *LIPA* as the candidate causal gene within the *LIPA* locus

GWASs have identified a number of common genetic variants at the *LIPA* locus that are strongly associated with risks of CAD as summarized in **Table S1**. Leveraging the CARDIoGRAMplusC4D GWAS dataset (7, 13), we identified twenty SNPs ($P < 1 \times 10^{-8}$) significantly associated with CAD risk, with rs1412444 as the lead SNP ($P = 5.15 \times 10^{-12}$) **(Figure 1A** and **Table S2**).

To identify the causal gene and causal tissue/cell types at the locus, we applied integrative genomic analysis pipeline employing three methods from two broad classes: 1) Colocalization analysis; 2) Summarized Mendelian Randomization (SMR) followed by Heterogeneity in Dependent Instruments Test (HEIDI) (21) or MetaXcan (22). Using CARDIoGRAMplusC4D GWAS and GTEx eQTL data, which shows the strongest *LIPA* eQTL in blood (**Figure S1A**), colocalization analysis by ENLOC (23) suggest that *LIPA*

is the only nearby gene showing strong regional colocalization probability (rcp) in this

region (± 1 Mb flanking gene *LIPA,* rcp = 0.657, whereas all other genes have rcp = 0)

(**Figure 1B** and **Figure S1B)**. By integrating CARDIoGRAMplusC4D and UKBB

GWASs with STARNET eQTL data of 9 tissue/cell types in this region (± 500 kb flanking

gene *LIPA*), we confirmed that *LIPA* showed significant colocalization signal (Coloc H_4 >

0.8) in CAD-relevant tissue/cell types (**Table S3**). SMR-HEIDI test examines if the same

variant is associated with gene expression and complex traits (24). SMR-HEIDI analysis

also identified *LIPA* as the only candidate causal gene with P_{SMR} < 0.001 and P_{HEIDI} >=

0.05 (**Table S4)**. MetaXcan further confirmed *LIPA* as the candidate causal gene

demonstrating the strongest MetaXcan evidence (P < 5 × 10^{-8}) (**Table S5**). Expanding

on prior work that hinted at *LIPA* as a causal gene (2, 3), our integrative genomic

analyses provided strong statistical evidence establishing *LIPA* as the causal gene at

this locus.

Functional genomic data support monocytes and macrophages as the causal cell

types

Functional regulatory variants and genomic elements often exert tissue- and cell type-

specific effects to regulate the expression of their target genes. Settling causal

tissues/cell types will enhance our understanding of disease-relevant regulatory

mechanisms. We applied three strategies leveraging large-scale transcriptomic and

epigenomic datasets to elucidate the involved cell types for the regulatory role of the

LIPA CAD variants.

First, GTEx eQTL analysis revealed the strongest *LIPA* eQTL association in whole

blood, followed by spleen and visceral adipose tissue (**Figure S1A**). Given the high

immune cell content in these tissues, we examined the BLUEPRINT dataset to identify

specific immune cell association (25). As shown in **Figure 1C**, monocytes showed the

strongest eQTL associations for CAD SNPs across all three eQTL methods, including

WASP-CHT (WASP-corrected combined haplotype test),(26) WASP-ASE (WASP-

corrected allele-specific expression) (26), and WP10 eQTL, while T-cells showing no

eQTL signals. Neutrophil exhibited a weak eQTL signal for SNP rs304412, which lacks

significant CAD association (**Table S2**).

Second, eQTL studies in diseased tissue allow the discovery of associations not

identifiable in healthy tissues, likely due to shifts in cell composition and regulatory

landscape changes. Using STARNET, with eQTL studies in cardiometabolic tissues

from patients with CAD, and CARDIoGRAMplusC4D datasets (± 500 kb flanking gene

LIPA), our analysis found more evidence for the role of monocytes and macrophages in

CAD-risk at the *LIPA* locus: 1) atherosclerotic aortic wall (AOR), blood, and human

peripheral blood monocyte-derived macrophage (HMDM) (**Figure 1D-1F**, respectively)

represent the strongest evidence as the causal tissues/cell types ($P_{eQTL} < 5 \times 10^{-8}$,

Table S6), passing colocalization, SMR-HEIDI, and metaXan tests (Coloc $H_4 > 0.8$,

$P_{SMR} < 0.001$ and $P_{HEIDI} >= 0.05$, and $P_{metaXan} < 5 \times 10^{-8}$, **Table S3-S5**); 2) foamy

macrophages (acetylated-LDL loaded HMDMs) showed strong eQTL (**Figure 1G**)

meeting SMR-HEIDI (**Table S4**); 3) *LIPA* eQTL associations in liver, visceral abdominal

adipose (VAF), and skeletal muscle (SKLM) had $P_{eQTL} < 5 \times 10^{-8}$ (**Figure S2, Table S6**),

though SMR-HEIDI suggested that different variants (in high LD) independently

regulated CAD risk and *LIPA* expression in these tissues ($P_{HEIDI} < 0.05$, **Table S4**); 4)

no *LIPA* eQTL association was found in pre/early-atherosclerotic mammary arteries

(MAM, **Figure S2**, **Table S6**), the healthy arteries mainly consist of endothelial cells and

vascular smooth muscle cells unlikely to have immune cell accumulation. These results

together support that monocytes/macrophages are likely the causal cell types for the

LIPA locus.

Finally, we harnessed single-cell datasets to provide insights into cell type-specific

regulatory mechanisms. Single-cell ATAC-seq of human atherosclerotic plaque

identifies cell type-specific regulatory elements, with recent scATAC-seq of human

atherosclerotic plaques (n = 41 subjects and 28,316 cells) revealing a macrophage-

specific regulatory element containing rs1320496 and the lead SNP (rs1412444) for

CAD (27). The regulatory element was absent in vascular smooth muscle cell-derived

cells and other cell types in the plaque, underscoring a macrophage-specific regulatory

mechanism. In addition, in the STARNET dataset, rs1320496 was the lead eQTL SNP

(P = 1.11×10^{-11}) for *LIPA* in foamy macrophages (**Figure 1G, Table S6**). Collectively,

bulk and single-cell data highlight monocytes and macrophages as the causal cell type

for *LIPA* variant regulation.

LIPA CAD risk alleles are associated with higher _LIPA_ mRNA, protein, and

enzyme activity in monocytes and macrophages

Upon establishing the causal cell types, it is important to confirm how the CAD risk alleles affect *LIPA* expression in these cells. Previous eQTL studies show that the risk alleles (T) of rs1412444 and rs2246833 (in high LD, $r^2 = 0.98$) are associated with higher *LIPA* mRNA (2, 3), and more importantly, higher LIPA enzyme activity (19) in human monocytes. In light of the strong *LIPA* eQTL signals in HMDM and foamy macrophages (**Figure 1F** and **1G**) and the vital role of macrophages in atherosclerosis (28), the cause of CAD, we examined *LIPA* expression and enzyme activity in human macrophages.

Using HMDMs differentiated from peripheral blood mononuclear cells isolated from fully genotyped healthy individuals (**Table S7**), we observed a 1.42-fold and 1.57-fold increase in *LIPA* mRNA for C/T and T/T genotypes of rs1412444, respectively, and significant increase in LIPA enzyme activity (by 1.37-fold for C/T and 1.47-fold for T/T) compared to C/C, the non-risk genotype (**Figure 1H** and **1I**). LIPA protein levels also increase in risk allele carriers, confirmed by Western blot (29) (**Figure 1J**). Thus, both previous functional genomic datasets and our experimental cohort validate that CAD risk alleles link to higher LIPA expression and enzyme activity in monocytes (19) and macrophages.

Experimental validation establishes the functional regulatory element and the regulatory mechanisms of the candidate causal variants

Identifying causal variants and linking them to the target gene remains challenging, yet

it represents a critical step to definitively establish the causality of the locus. Most

disease-associated noncoding variants reside in *cis*-regulatory elements (cREs) that

regulate gene expression in a cell type-specific manner (30). We experimentally

validated the cREs and functional variants regulating *LIPA* expression using the

following strategies: 1) prioritize cREs harboring GWAS variants using ChIP-seq data in

monocytes and macrophages (31) and luciferase assay; 2) use high-resolution Tri-HiC

to identify chromatin looping between *LIPA* promoter and putative cREs, and CRISPR

interference (CRISPRi) to validate the role of the cRE in regulating *LIPA* expression; 3)

perform luciferase assay with site-directed mutagenesis to identify functional variant(s);

and 4) analyze allele-specific binding (ASB) and motifs to understand transcription

factor (TF) interactions.

CAD GWASs reproducibly identified rs1412444 and rs2246833, both intronic in *LIPA*,

as the lead SNP or tag SNP (**Table S1**). Using epigenomic data in monocytes and

macrophages (31), we visualized the locus, including histone marks (H3K4me1 and

H3K27ac) and TF binding sites (**Figure 2A** and **Figure S3**). The rs1412444 region

aligns with stronger enhancer marks and PU.1 binding (**Figure 2A**), showing enhancer

activity by luciferase assay in THP-1 monocytes but not HEK293 cells, indicating a

monocyte/macrophage-specific regulatory mechanism (**Figure 2B**). The rs2246833

region showed no enhancer activity in either cell line (**Figure 2B**), consistent with

weaker enhancer features (**Figure 2A**).

Enhancers regulate gene expression by recruiting TFs and looping with the target gene

promoter, a key mechanism for how noncoding variants influence transcription (30).

Using high-resolution Tri-HiC data (32), we identified a strong interaction between the

rs1412444 region and the *LIPA* promoter in human monocytes, but not T-cells (**Figure**

2C), aligning with eQTL data showing lack of eQTL association in T-cells (**Figure 1C**).

CRISPRi targeting the rs1412444 region suppressed *LIPA* expression in THP-1 cells,

especially by gRNA spanning rs1320496 (**Figure 2D**).

The rs1412444 region, containing SNPs rs1412445, rs1320496 and rs1412444, was

prioritized as the causal cRE to further identify the functional variants in the region using

both experimental and statistical methods. The SNP rs1412444 is the lead SNP in CAD

GWAS (**Figure 1A**) and *LIPA* eQTL in atherosclerotic aortas (**Figure 1D**). The SNP

rs1412445 is in extremely high LD with rs1412444 (r^2 = 0.99), thereby serving as a

proxy. The SNP rs1320496 shows the strongest signal for *LIPA* eQTL in HMDM and

foamy macrophages (**Figure 1F-1G**), and is partly linked with SNP rs1412444 (r^2 = 0.45,

D' = 1). Among three major haplotypes (CCC, CTC, and TTT, defined by rs1412445,

rs1320496, and rs1412444 genotypes, **Table S8**), compared to the non-risk haplotype

CCC, the CTC and CTT haplotypes showed increased enhancer activity in monocytes

by luciferase assay (**Figure 3A**), suggesting that both rs1412445/rs1412444 and

rs10320496 may affect *LIPA* expression. Conditional analysis partially disentangles the

contribution of rs1412445/rs1412444 and rs1320496 on *LIPA* expression and CAD risk,

with both SNPs significantly linked to LIPA expression and CAD when analyzed

independently, supporting the presence of multiple causal variants within the rs1412444

enhancer region (**Figure 3B**).

We next sought to understand the mechanisms of action of the functional variants in the rs1412444 enhancer region. Many eQTL SNPs regulate target gene expression by altering TF binding. Using eQTL data from 49 tissues in GTEx, we have previously established that analyzing eQTL effect size as a function of TF expression level offers a generalizable approach to discover TF regulators of genetic variant effects (33). Using this analysis pipeline, we observed significant correlations between eQTL effects of rs1320496 and the expression levels of 9 TFs, with PU.1, encoded by the *SPI1* gene, showing the strongest effect (Spearman rho = 0.48, P = 6.0 × 10^{-4}, **Figure S4A**). Indeed, the rs1412444 enhancer region overlaps with the PU.1 binding site, an important TF required for myeloid differentiation (**Figure 2A**) (34). We, therefore, hypothesize that the functional variants in the rs1412444 region may alter PU.1 binding affinity. To test this, we used: 1) allele-specific PU.1 ChIP-seq; 2) motif analysis; 3) electrophoresis mobility shift assay (EMSA); and 4) knockdown of *SPI1* to confirm causality. PU.1 ChIP-seq (35) showed higher PU.1 binding for the risk alleles (T) of rs1320496 and rs1412444 compared to non-risk alleles (C) (**Figure 3C**). Among the SNPs, only rs1320496 (T) was predicted to directly change a key nucleotide to enhance motif-matching (**Figure 3D**). EMSA further validated that the T allele of rs1320496 created a binding site for PU.1 (**Figure 3E**). The T allele of rs1412444 showed increased PU.1 binding compared to the C allele (**Figure 3E**), while the T allele of rs1412445 showed no evidence of binding to PU.1 (**Figure S4B**). We further validated that knockdown of *SPI1* reduces *LIPA* mRNA expression (**Figure 3F**), and partially abolished the effects of the risk alleles on increasing the enhancer activity of the rs1412444 region (**Figure 3G**). These findings

confirm that risk alleles of rs1320496 and rs1412444 increase *LIPA* expression and

contribute to CAD risk by enhancing PU.1 binding. (**Figure 3H**)

We also examined TFs beyond PU.1. A modest correlation between rs1412444 eQTL

effects and *STAT1* expression was noted in GTEx data (Spearman rho = 0.26, P = 0.07,

Figure S5A) and ENCODE data (**Figure S5D**), with STAT1 ChIP-seq in CD14^{+}

monocytes showing an allelic imbalance (**Figure S5B**). However, *STAT1* knockdown did

not affect *LIPA* expression (**Figure S5C**), suggesting STAT1 unlikely regulate *LIPA* CAD

variants effects.

In summary, our functional genomic studies confirm that the risk alleles of genetic

variants associated with CAD lead to increased *LIPA* mRNA, protein, and enzyme

activity in monocytes and macrophages. These variants within the enhancer region

interact with the *LIPA* promotor and regulate *LIPA* expression by altering PU.1 binding.

Having established the causative role of these variants in elevating LIPA, we next aim to

assess whether this increase exacerbates atherosclerosis using myeloid-specific *Lipa*

overexpression in preclinical mouse models.

Preclinical mouse studies demonstrate that myeloid-specific overexpression of

***Lipa* increases atherosclerosis**

To elucidate how increased myeloid LIPA impacts atherosclerosis *in vivo*, we developed

a myeloid-specific *Lipa* overexpression mouse model (*LysMCre$^{+/-}$, Lipa$^{KI/WT}$, Ldlr$^{-/-}$,*

referred to as *M-Lipa^{KI}*, **Figure 4A** and **S6A** for illustration of the construct and breeding

strategy). Their corresponding littermates *LysMCre^{-/-}, Lipa^{KI/WT}, Ldlr^{-/-}* were used as

control (*Ctrl*). The percentage of GFP^+ cells, a reporter of *Lipa* overexpression, was

quantified across multiple tissues to demonstrate specificity and efficiency of

overexpression (**Figure S6C-S6E**). In *M-Lipa^{KI}* mice, GFP^+ cells comprised 36.2%,

22.0%, 34.8%, and 68.5% of CD45^+CD115^+ monocytes/macrophages in blood, bone

marrow (BM), spleen, and peritoneum, respectively (**Figure S6C**). GFP^+ cells

represented 46.3% and 70.2% of CD115^-Ly6G^{hi} neutrophils in blood and BM (**Figure**

S6D), with minimal GFP^+ CD3^+ lymphocytes (< 2.7%, **Figure S6E**), confirming myeloid-

specific overexpression. The efficiency of *LysMCre* recombination is comparable to

previous reports on *LysMCre*-mediated overexpression in mouse models (36).

The overexpression of *Lipa* at mRNA, protein, and enzyme activity levels was confirmed

in bone marrow-derived macrophages (BMDMs) and peritoneal macrophages (PMs)

from *M-Lipa^{KI}* mice of both sexes (**Figure S6F-S6O**). In male mice, *Lipa* mRNA and

enzyme activity increased by 1.56-fold and 1.26-fold in *M-Lipa^{KI}* BMDMs (**Figure S6F**

and S6G) and by 3.49-fold and 1.44-fold in *M-Lipa^{KI}* PMs (**Figure S6K and S6L**)

compared to *Ctrl*, respectively. In female mice, the *Lipa* mRNA and enzyme activity

increased by 1.41-fold and 1.21-fold in *M-Lipa^{KI}* BMDMs (**Figure S6H and S6I**) and

2.52-fold and 1.24-fold in *M-Lipa^{KI}* PMs (**Figure S6M and S6N**) compared to *Ctrl*,

respectively. Consistently, LIPA protein significantly increased in both BMDMs (**Figure**

S6J) and PMs (**Figure S6O**) from *M-Lipa^{KI}* mice compared to *Ctrl*. Despite the modest

overexpression, the enzyme activity increase was comparable to HMDMs of risk allele

carriers (**Figure 1I**), supporting *M-Lipa^{KI}* mice as a model for studying the causal effects

of gain-of-function of LIPA observed in human functional genomic data.

Ctrl and *M-Lipa^{KI}* mice were fed a Western diet (WD, TD88137, Envigo Teklad) for 16

weeks to induce atherosclerosis, with body weight, organ weight, and lipid profiles

assessed (**Figure S7A-S7N**). Body weight, spleen weight/body weight, and liver

weight/body weight were comparable between *Ctrl* and *M-Lipa^{KI}* mice in both males

(**Figure S7A-S7C**) and females (**Figure S7H-S7J**). No differences in total cholesterol

(**Figure S7D** and **S7K**) or cholesterol in low-density lipoprotein (LDL) and high-density

lipoprotein (HDL) fractions were observed (**Figure S7E** and **S7L,** after 4 h fasting and in

both sexes). Female *M-Lipa^{KI}* mice showed a trend toward higher plasma triglyceride (P

= 0.050, **Figure S7M**) and higher triglyceride in both LDL and HDL fractions (**Figure

S7N**), an effect not seen in males (**Figure S7F** and **S7G**). Overall, myeloid *Lipa*

overexpression did not alter body weight or plasma cholesterol, consistent with human

data linking *LIPA* variants to CAD but not metabolic traits, while the modest effects of

myeloid *Lipa* overexpression on triglyceride metabolism specifically in female mice on a

WD require further examination (37).

Atherosclerotic lesion size was significantly larger, with a 17.78% increase in male and

21.85% in female *M-Lipa^{KI}* mice compared to *Ctrl* mice (**Figure 4B**). Lesion size

positively correlated with LIPA activity in PMs (**Figure 4C**, P = 0.014, *r* = 0.4499),

supporting the association between increased macrophage LIPA activity and

atherogenesis. No differences in necrotic core area or fibrous cap thickness, features of

plaque instability (38), were observed (**Figure 4D**, **4E**).

In summary, our findings highlight a pro-atherogenic role for increased myeloid Lipa

expression without affecting body weight or lipid profile, in line with human genomic

data.

Myeloid *Lipa* overexpression promotes atherosclerosis with an increased

macrophage content in the lesion, mainly attributable to increase monocyte

recruitment

Myeloid *Lipa* overexpression significantly increased the CD68$^+$ macrophage area in

lesions for both sexes (**Figure 5A**), with no differences in macrophage proliferation or

apoptosis, as determined by Ki67 and TUNEL (terminal deoxynucleotidyl transferase

dUTP nick-end labeling) staining (**Figure 5B-5C**), suggesting other mechanisms for the

increase.

During atherogenesis, blood monocytes continuously influx into the subendothelial

space and contribute to lesion macrophage accumulation (28). Since plaque

macrophage proliferation or apoptosis remained unaffected, we hypothesize that

increased macrophage area may result from increased monocyte recruitment. To test

this, Ly6C^{hi} circulating monocytes were pulse-labeled with fluorescent beads (**Figure 5D**)

as previously described (39). Three days after labeling, bead numbers within the lesion

were quantified. *M-Lipa^{KI}* mice showed increased beads in CD68$^+$ macrophage area

within plaques (**Figure 5D**), indicating higher monocyte recruitment. We further

confirmed that *M-Lipa^{KI}* with WD feeding did not alter myelopoiesis in blood or BM

(**Figure S8A and S8B**) or affect the percentage of hematopoietic progenitor cells in BM

(**Figure S8C**). Chemokines, *e.g.,* CCL2 (40, 41), CCL8 (42), and CCL7 (43), have been

implicated in regulating monocyte infiltration in atherosclerosis progression. The levels

of plasma CCL2, CCL7, and CCL8, and 13 other chemokines and cytokines, showed no

difference between genotypes (**Figure S9**). These results suggest that myeloid *Lipa*

overexpression increases monocyte recruitment to the plaques, likely due to changes in

the local plaque microenvironment rather than monocyte numbers or systemic

inflammation.

Myeloid *Lipa* overexpression leads to reduced neutral lipid accumulation and

increased free cholesterol accumulation in macrophages

Beyond the increased macrophages in *M-Lipa^{KI}* plaques, we further validated the

functional effects of *Lipa* overexpression in macrophages. Using flow cytometry, we

characterized foamy (SSC^{hi}LipidTOX^{hi}) aortic macrophages (CD45$^+$CD11b$^+$CD64$^+$)

(**Figure 6A** for study design and **Figure S10A** for gating strategy) (44). We confirmed

that in atherosclerotic plaques, the percentage of SSC^{hi}LipidTOX^{hi} foamy aortic

macrophages was similar between *M-Lipa^{KI}* and *Ctrl* mice (**Figure 6B**), but the mean

fluorescent intensity (MFI) of LipidTOX in foamy aortic macrophages isolated from *M-*

Lipa^{KI} mice was significantly lower than that in *Ctrl* mice (**Figure 6C**), indicating reduced

neutral lipids. Within *M-Lipa^{KI}* mice, GFP^+ aortic macrophages had lower neutral lipids

than GFP^- aortic macrophages (**Figure 6D**), confirming intrinsic effects driven by *Lipa*

overexpression.

We also assessed SSC^{hi}LipidTOX^{hi} foamy PMs (CD45^+F4/80^+) from WD-fed mice

(**Figure 6E** for study design and **Figure S10B** for gating strategy). The percentage and

LipidTOX MFI of foamy PMs were significantly lower in *M-Lipa^{KI}* mice compared to *Ctrl*

mice, indicating reduced neutral lipid accumulation (**Figure 6F**, **6G**). Fluorometric

cholesterol assays in PMs (**Figure 6H** for study design) further showed higher total and

free cholesterol but lower cholesteryl ester levels in *M-Lipa^{KI}* PMs (**Figure 6I**), aligning

with LipidTOX findings. These data consistently show reduced neutral lipid

accumulation in *Lipa^{KI}* aortic macrophages and PMs.

We further explored if reduced neutral lipid accumulation in *M-Lipa^{KI}* PMs from WD-fed

mice could be replicated *in vitro* in cultured macrophages loaded with modified

lipoproteins. PMs from mice fed a normal laboratory diet (ND) were treated with

oxidized-LDL (oxLDL, **Figure S11A** for study design). We indeed observe a lower MFI

of LipidTOX in oxLDL-loaded *M-Lipa^{KI}* PMs compared to *Ctrl* PMs (**Figure S11B**), which

was not due to a significant change in oxLDL-binding or uptake (**Figure S11C** and

S11D). As expected, *Lipa* mRNA was elevated in *M-Lipa^{KI}* PMs (**Figure S11E**), while

genes for lipoprotein uptake were similar (**Figure S11E)**. *M-Lipa^{KI}* PMs had lower

expression of cholesterol biosynthesis genes (**Figure S11F**), suggesting increased free

cholesterol accumulation in the endoplasmic reticulum thus a suppression of cholesterol

biosynthesis genes. No differences were seen in cholesterol efflux gene expression (45),

including *Abca1* and *Abcg1* (**Figure S11G**). It is worth noting that oxLDL loading only

led to a slight increase in total and free cholesterol, but levels were similar between *M-*

Lipa^{KI} and *Ctrl* PMs (**Figure S11H**). Thus, oxLDL-loaded PMs replicate some, but not all,

phenotypic features of PMs from WD-fed mice, underscoring the importance of carefully

assessing the appropriate cellular model for mechanistic studies.

Myeloid *Lipa* overexpression distinctly alters the transcriptomic signature of

aortic macrophages and PMs

Building on the observed phenotypic changes, we sought to determine how myeloid

Lipa overexpression alters macrophage transcriptomes. To compare the transcriptomic

signature of aortic macrophages between *Ctrl* and *M-Lipa^{KI}* mice fed a WD for 16 weeks,

we performed bulk RNA-seq using the low-input RNA-seq method owing to the limited

materials of aortic macrophages (**Figure 7A** for study design). We identified 300

upregulated and 123 downregulated DE genes in *M-Lipa^{KI}* vs. *Ctrl* aortic macrophages

(**Figure 7B** and **Table S9**, absolute fold change > 1.5, FDR-adjusted P < 0.05). Gene

Set Enrichment Analysis (GSEA) showed enrichment of integrin signaling and cell-

matrix adhesion pathways in the upregulated DE genes (**Figure 7C** and **Table S10**).

The increased integrins (*Itga6*, *Itga9*, *Itgb3*, *Itgax*, etc.) and adhesion molecules and

receptors (*Cspg4*, *Dag1, Fn1, Kdr, Tnc, Npnt,* etc.) in *M-Lipa^{KI}* aortic macrophages may

contribute to the increased monocyte adhesion and infiltration and macrophage

adhesion in the atherosclerotic plaque microenvironment, thus promoting atherosclerosis (**Figure 7D)**.

To further examine if *Lipa* overexpression uniquely affect aortic macrophages, we performed RNA-seq on PMs from *Ctrl* and *M-Lipa^KI* mice fed a WD for 16 weeks (**Figure 7E** for study design). We identified a total of 69 upregulated and 31 downregulated DE genes (**Figure 7F** and **Table S11**, absolute fold change > 1.5, FDR-adjusted P < 0.05). GSEA indicated enrichment in interferon signaling and antigen presentation pathways in the upregulated DE genes in *M-Lipa^KI* vs. *Ctrl* PMs (**Figure 7G** and **Table S12**). This is consistent with the recent finding that lipid loading suppressed the expression of interferon-stimulated genes, e.g., *Ifit2*, *Ifit1*, *Oas3*, and *Stat2*, in mouse and human macrophages (46), as visualized in **Figure 7H**. The upregulated expression of major histocompatibility complex genes, e.g., *H2-Eb1*, *H2-Q6*, *H2-Ab1*, *H2-K1*, etc. in PMs of *M-Lipa^KI* mice implicate immune activation (**Figure 7I**). To explore whether our observation of increased free cholesterol in PMs of *M-Lipa^KI* mice may link to a proinflammatory transcriptomic signature, we further validated the mRNA of *Il18* and *Il1b* in PMs using qRT-PCR (**Figure S12**) and the plasma level of IL-18 and IL1β (**Figure S9**) in mice. However, no significant differences were observed between *Ctrl* and *M-Lipa^KI*, suggesting NLRP3 inflammasome may not be the primary proinflammatory driving force in this context, despite the known role of excessive free cholesterol accumulation in the endoplasmic reticulum in activating the NLRP3 inflammasome (47, 48).

These findings emphasize the distinct effects of *Lipa* overexpression on aortic

macrophages compared to other macrophage types, highlighting the importance of

examining context-specific cellular responses in disease mechanisms.

Single-cell RNA-seq analysis reveals that myeloid *Lipa* overexpression alters cell

heterogeneity and transcriptomic profile of atherosclerotic aortic cells

The transcriptomic signature of increased integrin and extracellular matrix (ECM) genes

in *Lipa*-overexpressing aortic macrophage prompts us to speculate whether the

changes reflect altered macrophage subpopulations and whether macrophage

transcriptomic changes can alter other plaque cell types via cell-cell interaction. To gain

insights into the effects of myeloid *Lipa* overexpression on the phenotypic and functional

heterogeneity of plaque cells, we carried out single-cell RNA-sequencing (scRNA-seq)

analysis. Aortic single-cell suspensions were obtained from *Ctrl* and *M-Lipa^{KI}* mice fed a

WD for 16 weeks. Live single cells were sequenced using the 10X Chromium platform

(**Figure 8A** for study design and **Figure S13A** for gating strategy). Data analysis was

performed using Seurat v4.3.0.1 (49). Annotation of cell types was based on top marker

genes of each cluster and by comparing with marker genes defined in previous meta-

analyses of atherosclerotic plaque scRNA-seq data (**Table S13** for the top 15 marker

genes and **Figure 8B** for the visualization of cell clusters and **Figure 8C** the top 3

marker genes) (50). Data confirmed an increased macrophage proportion in *M-Lipa^{KI}*

mice (**Figure 8D** and **Table S14**), consistent with histological data (**Figure 5A**). Results

also suggest decreased SMC proportion and increased fibroblast proportion (**Figure 8D**

and **Table S14**). Understanding the noisy nature of scRNA-seq, we performed DE

analysis for each cell type to gain exploratory insights into effects of myeloid *Lipa*

overexpression on the transcriptomic signature of different plaque cell types. While we

observed modest changes, notably, among the 6 cell types, we identified the most DE

genes in SMCs between *M-Lipa^{KI}* and *Ctrl* mice (**Table S15** for DE genes and **Table**

S16 for canonical pathway analysis using Ingenuity Pathway Analysis, IPA). Focused

analysis of SMCs reveals 6 SMC subclusters consistent with these identified in the

literature (50) (**Figure 8E** for UMAP visualization and **Figure S13B** for marker gene

visualization). In *M-Lipa^{KI}* mice, a reduction in SMC1 and an increase in Cxcl12 SMC

and fibroblast-like SMCs were observed (**Figure 8F**). IPA suggests the activation of

canonical pathways in SMCs of *M-Lipa^{KI}* mice, including the Integrin Cell Surface

Interactions and Atherosclerosis Signaling canonical pathways (**Figure 8G** and **Table**

S16), which corroborates with the observed increase in atherosclerotic lesion and the

increased integrin and ECM genes in aortic macrophages by bulk RNA-seq.

Upregulated DE genes in SMCs of *M-Lipa^{KI}* mice include *Col1a1*, *Col1a2*, *Lum*, *Thbs1*,

Vcam1, and *Mmp2* (**Table S15**), all of which have been implicated in contributing to

increased ECM remodeling,(51) with *Vcam1* also playing a role in the recruitment and

retention of immune cells within plaques (52). Among the 6 SMC subclusters, SMC1

and Cxcl12 SMCs showed the most DE genes between *Ctrl* and *M-Lipa^{KI}* mice. IPA

reveals activation of the Integrin Cell Surface Interactions canonical pathway in the

SMC1 subcluster (**Figure 8H** and **Table S16**), while inhibition of the Smooth Muscle

Contraction pathway in the Cxcl12 SMC subcluster (**Figure 8I** and **Table S16**).

Collectively, these data suggest that in *M-Lipa^{KI}* mice, plaque SMCs may undergo

increased dedifferentiation, and macrophage-SMC interactions could be part of the

mechanisms contributing to the enhanced atheroprogression. Even though the

transcriptomic signature of macrophages did not differ between *M-Lipa^{KI}* and *Ctrl* mice

in this dataset (**Table S15** and **S16**), clustering analysis identified subclusters of aortic

macrophages, including Trem2$^+$, Ccl2$^+$, Lyve1$^+$ (resident-like), Il1b$^+$ (inflammatory), and

proliferative macrophages, aligning with those consistently reported in the literature (53,

54), showing comparable subcluster proportions (**Figure S13C-S13E**).

Given the transcriptomic differences in aortic macrophages observed in bulk RNA-seq

data, we speculate that while scRNA-seq provides valuable information on cellular

heterogeneity, it may be less sensitive in detecting transcriptomic differences, especially

when sequencing all aortic cells with limited reads per cell type. To gain deeper

molecular insights into the effects of *M-Lipa^{KI}* on the phenotypic and functional

heterogeneity of aortic macrophages, we conducted scRNA-seq analysis specifically

focused on these cells. Since *LysMCre* does not result in 100% recombination, we

leverage this by separating CD45$^+$GFP$^-$ (no *Lipa* overexpression) and CD45$^+$GFP$^+$

(*Lipa*-overexpressing) cells from enzymatically dissociated aortas of *M-Lipa^{KI}* mice

(**Figure S14A** for study design and **Figure S14B** for gating strategies). This approach

allowed us to compare aortic macrophages with or without *Lipa* overexpression within

the same plaque microenvironment (**Table S17** for the top 20 marker genes and **Figure**

S14C for subcluster identification and assignment by comparing with marker genes

defined in previous meta-analyses of atherosclerotic plaque scRNA-seq data) (53, 54).

A total of 12 distinct cell clusters were identified, including 5 macrophage clusters and 4

monocyte/dendritic clusters, as well as small numbers of B cells, fibroblasts, and SMCs

that are present likely because of sorting-related contamination (**Figure S14C).** The

GFP$^+$ sample has more than 95% of the cells annotated as macrophages/monocytes,

confirming the myeloid specificity of *Lipa* overexpression in our model. We have also

validated the increased *Lipa* mRNA in GFP$^+$ aortic macrophages compared to GFP$^-$

aortic macrophages sorted from the aortas of *M-Lipa^{KI}* mice, as well as in macrophages

sorted from the aortas of *Ctrl* mice (**Figure S14D**). A total of 5 macrophage subclusters

were identified in both GFP$^-$ and GFP$^+$ populations (**Figure S14E**), including Trem2

macrophages (cluster 0), Trem2 foamy macrophages (cluster 1), Lyve1 macrophages

(cluster 2), Il1b macrophages (cluster 4), and proliferative macrophages (cluster 5).

Both cluster 0 and cluster 1 had higher *Trem2* expression (**Figure S14E**), while cluster

1 had high expression of lipid processing genes *Fabp5* and *Gpnmb*, and lysosomal

genes *Ctsl* and *Atp6v0d2* (**Figure S14E** and **Table S17**), a signature resembling

increased lipid metabolism, that we refer to as Trem2 foamy macrophages.(54) Notably,

the percentage of Trem2 foamy macrophages (cluster 1) is lower in the GFP$^+$ population

(17.32%) compared to the GFP$^-$ population (36.00%, **Figure S14F** and **Table S18**). This

lower proportion of lipid metabolism-associated aortic macrophage based on scRNA-

seq analysis is consistent with lower neutral lipids in GFP$^+$ aortic macrophages

observed by flow cytometry (**Figure 6D**). DE and IPA analyses suggest cluster 0 Trem2

macrophages had the most DE genes (**Table S19**), representing a transcriptomic

signature showing predicted activation of pathways involving oxidative stress and

inflammation, such as iNOS Signaling, TREM1 Signaling, and TNFR1 Signaling (**Figure**

S14G and **Table S20**); these pathways were also identified by IPA in the merged

datasets between GFP- and GFP+ macrophages (**Figure S14H, Table S20**).

Taken together, histological and phenotypic analyses confirm an increased macrophage

area and decreased neutral lipid accumulation in aortic macrophages of *M-Lipa^{KI}* mice;

scRNA-seq analysis further indicates that myeloid *Lipa* overexpression increases the

overall proportion of plaque macrophages while reducing the proportion of the

macrophage subcluster associated with lipid metabolism. Furthermore, the data suggest

that myeloid *Lipa* overexpression influences SMCs, leading to a transcriptomic

signature showing the activation of Integrin Cell Surface Interactions and

Atherosclerosis Signaling canonical pathways, which may contribute to plaque

progression.

Discussion

Despite the first CAD GWAS being more than a decade ago, successful translation of

genetic associations into causal genes and pathways has been confined to a small

subset of GWAS loci (30). A number of genetic risk loci for CAD act through established

CAD risk factors, such as lipid traits (13), which themselves have a significant genetic

determination. Yet, many confirmed loci appear to act through novel mechanisms that

are independent of traditional risk factors. Elucidating these mechanisms is critical for a

better understanding of CAD pathogenesis and the identification of therapeutic targets.

LIPA is one of the CAD loci that is specifically associated with CAD (2-13), but not with

lipid traits (14). It is therefore of great interest to tackle the variant-to-function

relationship for the *LIPA* locus and to determine how altered *LIPA* expression causally

impacts atherosclerosis through mechanisms beyond its effects on plasma lipids. Here,

using functional genomic approaches and experimental mouse models, we present

evidence that *LIPA* is causal in CAD. Our results support that: 1) *LIPA* expression and

enzyme activity are higher in HMDM of CAD risk allele carriers; 2) the CAD GWAS lead

SNP rs1412444 resides in an enhancer region that shows strong interaction with the

LIPA promoter in a monocyte-specific manner, with CRISPRi targeting the region

suppressing *LIPA* expression; 3) within the enhancer region, risk alleles of rs1412444

and rs1320496 show higher binding to PU.1 and lead to increased enhancer activity; 4)

Myeloid overexpression of *Lipa* leads to increased atherosclerosis lesion area and

macrophage area in mice on *Ldlr*[-/-] background fed a WD, due to increased monocyte

recruitment; 5) Myeloid overexpression of *Lipa* leads to an increase in free cholesterol

and decrease in cholesteryl ester accumulation in PMs of mice fed a WD, accompanied

by upregulation of genes in interferon signaling and antigen presentation pathways; 6)

Myeloid overexpression of *Lipa* alters plaque cell heterogeneity and transcriptomic

profile; the upregulation of genes in integrin signaling and cell-matrix adhesion

pathways may contribute to macrophage-SMC interaction and facilitate monocyte

recruitment, thus exacerbating atherosclerosis. Taken together, these data demonstrate

that increased *LIPA* in myeloid cells drives atheroprogression, a directionality consistent

with human GWAS and eQTLs finding for the *LIPA* locus (2-4).

Delineating causal variants is often challenging, partly because most GWAS variants

are noncoding and in high LD with multiple variants within the same locus. A coding

variant rs1051338, in high LD with the lead SNP rs1412444 (r^2 = 0.86), causes a

nonsynonymous threonine to proline change within the signal peptide of LIPA protein.

Because of the essential role of LIPA in lipid metabolism, the initial speculation was that

rs1051338 serves as a causal variant contributing to lower LIPA activity (55). Further

work by Evans *et al.* revealed that the nonsynonymous change within the signal peptide

of LIPA protein did not affect protein trafficking, enzyme activity, or secretion (19).

Evans *et al.* also confirmed that the risk allele (C) of rs1051338 is associated with

increased *LIPA* expression and enzyme activity in human monocytes (19). Our work

established the same directionality in human macrophages. We have also

demonstrated the cell type-specific regulatory role of the *LIPA* CAD variants, *i.e.,* the

enhancer activity of the rs1412444 region is monocyte-specific, and the chromatin

interaction between the rs1412444 enhancer region and *LIPA* promoter was observed

in monocytes, but not T-cells. We established that rs1320496 and rs1412444 are the

functional SNPs within the rs1412444 enhancer region that regulate allele-specific

interaction with PU.1, collectively exhibiting additive effects and suggesting a model of

multiple causal variants within a single locus. Protein QTL (pQTL) studies represent a

critical level of information for understanding the genetic regulation of protein expression.

Although pQTL studies were performed in STARNET plasma, LIPA was not included in

the original panel (NPX Olink platform). Future functional genomic data at the proteomic

and metabolomic levels, in a cell and tissue-specific manner, and experimental

validation by high-throughput functional genomic screening, will further enhance our

ability to connect genetic variation to the functional mechanisms at scale.

This work has provided valuable mechanistic insights, while highlighting the importance

of selecting appropriate cellular models for mechanistic studies. PMs of *M-Lipa^{KI}* mice

fed a WD showed increased free cholesterol accumulation and a transcriptomic

signature showing upregulated interferon signaling. We attempted to model this

phenotype *in vitro* using oxLDL-loaded PMs from mice fed a ND. However, our results

did not show an effect of *M-Lipa^{KI}* on oxLDL uptake or free cholesterol accumulation.

OxLDL loading induced a modest increase in total cholesterol, in contrast to the

remarkable total cholesterol and cholesteryl ester accumulation in PMs isolated from

WD-fed mice. These results highlight that *in vitro* cultured macrophages loaded with

chemically modified oxLDL do not fully resemble the foamy macrophages formed in an

in vivo environment with WD-induced hyperlipidemia. Previous work suggests that

lysosomal CE hydrolysis leads to free cholesterol accumulation in the lysosome when

macrophages are loaded with oxLDL *in vitro*, disrupting lysosomal homeostasis,

including the maintenance of an acidic pH (56). It has also been observed that

macrophages form extracellular lysosomal synapses where the LIPA enzyme remains

active and mediates catabolism of aggregated-LDL, generating free cholesterol for

internalization (57). These *in vitro* studies align with our observation that *Lipa^{KI}*

increases free cholesterol accumulation. It is also worth noting that plaque

macrophages demonstrate a transcriptomic signature of upregulated integrin and cell-

matrix adhesion pathways, affirming increased monocyte infiltration and macrophage

content in the atherosclerotic plaques of *M-Lipa^{KI}* mice, while PMs of these mice

showed enhanced interferon signaling. The differences in the effects of *Lipa*

overexpression on the transcriptomic signature of PMs and plaque macrophages

highlight the importance of local microenvironment in shaping cellular phenotypes.

While our work establishes *LIPA*'s genetic contribution to CAD, its physiological role as

the only known acidic lipase in the lysosome, in intracellular lipid metabolism is equally

critical. Loss-of-function of *LIPA* is the cause of Mendelian disorders characterized by

lysosomal neutral lipid accumulation in multiple tissues (17). Enzyme replacement

therapy with recombinant LIPA protein has demonstrated efficacy and safety in patients

with *LIPA* deficiency (58). These findings have demonstrated the essential role of *LIPA*

in lipid homeostasis and the benefits of supplementing LIPA when there is a severe LIPA

deficiency. However, the effects of further increasing LIPA above the normal

physiological level and for a prolonged period remain elusive. Loss of function of *LIPA* in

humans led to severe dyslipidemia and hepatomegaly, prompting a hypothesis that

overexpressing *LIPA* in the liver would prevent hepatic lipid accumulation. Yet, a recent

study demonstrated that hepatic *LIPA* overexpression by AAV8 did not attenuate

steatosis but unexpectedly exacerbated inflammation and immune cell infiltration in the

liver of mice fed a WD (59). Our work using myeloid-specific *Lipa* overexpression mice

established that increased *Lipa* in monocytes/macrophages increases atherosclerosis.

Our study on the role of myeloid-specific *Lipa* overexpression in CAD prompts the

needs to assess the benefits and risks in potentially therapeutically targeting LIPA in

CAD. For example, the assumption that further enhancing LIPA will have a favorable

role in accelerating lysosomal lipid catabolism should be carefully evaluated in a cell

type and disease-specific manner.

In addition to the intracellular and cell-intrinsic effects observed with *Lipa*

overexpression, another unexplored aspect is the potential transferability of the LIPA

enzyme. This raises the question of how increased *Lipa* expression in myeloid cells

might impact other cell types within the plaque, as well as in organs with resident or

infiltrated macrophage populations. For instance, we observed altered SMC

heterogeneity and transcriptomic profile, though it remains unclear whether this could

be partly regulated by enzyme transfer that increases SMC LIPA levels. Additionally, we

noted a slight elevation in triglyceride levels among female *M-Lipa^{KI}* mice, while male

mice did not show this change. Exploring the potential impact of myeloid-*Lipa*

overexpression on liver and adipose biology stands as a future area of research.

Moreover, our mouse model for conditional overexpression of *Lipa* presents an

opportunity to explore the functions of *Lipa* and lysosomal lipid metabolism across

various tissue and cell types, both under normal homeostasis and disease contexts.

In conclusion, our results established that *LIPA* risk alleles drive increased myeloid LIPA

expression, which in turn aggravates atherosclerosis. These findings are consistent with

and support human functional genomic discoveries, highlighting the critical role of LIPA

in the progression of CAD. The workflow and functional genomic resources applied in

this study provide valuable tools that will accelerate the systematic interrogation of

genetic contributions to CAD risk, particularly through the roles of monocytes and

macrophages.

Methods

Detailed description of methods and materials are available in the online-only

Supplemental Material.

<u>Sex as a biological variable</u>

Our study examined male and female animals, and similar findings are reported for both

sexes.

<u>Statistics</u>

All statistical analyses were performed using GraphPad Prism 9 as described below,

unless otherwise specified in the figure legends. When sample size (n) ≥ 6, data were

tested for normality using Shapiro-Wilk test (when n < 8) or D'Agostino-Pearson test

(when n ≥ 8). F-test of equality of variances was performed to compare the two sample

variances. Data that passed normality tests are presented as mean $\pm$ standard error

of mean (SEM) and analyzed using two-tailed Student's t-test for comparison of two

groups and equal variances (or with Welch's correction if F-test was not satisfied), or

one-way ANOVA with Tukey's post-hoc tests for one independent variable with more

than two groups. Data that did not follow a normal distribution were analyzed using

nonparametric tests, *i.e.,* Mann-Whitney U test or Kruskal-Wallis test, and are presented

as median ±□95% confidence interval (CI). Nonparametric tests were also used when n

< 6. P < 0.05 was considered statistically significant. The number of independent

experiments and biological replicates is specified in the figure legends.

Study approval

All human study protocols were approved by the Human Subjects Research Institutional

Review Board at Columbia University (AAAQ6510). All animal experiments were

approved by the Institutional Animal Care and Use Committee at Columbia University

(AABN5560). Mice were cared for according to the NIH guidelines.

Data availability

Integrative genomic analyses have incorporated two data sources: GWAS summary

statistics from the UK BioBank (UKBB) (60) or Coronary Artery Disease Genome Wide

Replication and Meta-analysis plus The Coronary Artery Disease Genetics

(CARDIoGRAMplusC4D) (7) and tissue/cell-type specific eQTLs from the Genotype-

Tissue Expression project (GTEx) v8,(61) STARNET (15), or the BLUEPRINT project

(25). GTEx is a large multi-tissue eQTL dataset containing 48 tissue types from ~900

healthy subjects (61). STARNET is an eQTL dataset in nine disease-relevant metabolic

tissues/cells collected from ~600 CAD patients (15).

The bulk RNA-seq and single-cell RNA-seq datasets generated during the current study

are deposited in the Gene Expression Omnibus (GEO) with accession number

GSE243139 (secure token for the reviewers: kfwtsccihbufpgd). All code for data

analysis associated with the current submission will be available via the lab GitHub repository at https://github.com/hanruizhang/ upon acceptance of the manuscript.

Author contributions

FL and HZ conceived and designed the research. FL designed and conducted majority of the experiments. FL analyzed RNA-seq and scRNA-seq data. EF performed functional genomic analysis using GTEx and ENCODE data. HC performed functional genomic analysis using STARNET data. CX conducted imputation analysis and provided critical feedback on scRNA-seq data analysis. YZ performed and analyzed Tri-HiC data. PH, MNZ, JS, XW, ZW, YM, JC, AR, JC, AHO performed experiments. AJ, BR, MW, RCB, and YS provided input on experimental design. KH and TL were instrumental in the interpretation of functional genomic data. FL and HZ analyzed and interpreted results and wrote the paper. HZ directed and supervised the project and funding.

Acknowledgements

We would like to acknowledge the NIH funding sources to the Columbia Center for Translational Immunology (CCTi) Flow Cytometry Core by grant number S10OD020056 and S10RR027050 and P30DK063608; the NIH-supported microscopy resources in the Center for Biologic Imaging, specifically the confocal microscope supported by grant number S10OD019973-01; the NIH/NCI Cancer Center Support Grant P30CA013696

for the use of resources at the Columbia Genome Center; and the Columbia Stem Cell

Initiative (CSCI) Flow Cytometry Core under the leadership of Michael Kissner. We

thank Dr. Chyuan-Sheng (Victor) Lin, Manager of the Genetically Modified Mouse

Models shared resource at Columbia University Herbert Irving Comprehensive Cancer

Center (HICCC), for their expertise in generating the mouse model for *Lipa*

overexpression used in this study. The authors thank Drs. Towfique Raj and Jack

Humphrey (Icahn School of Medicine at Mount Sinai), and Drs. Johan Björkegren

(Karolinska Institutet) and Sean A. Bankier (University of Bergen) for access to pQTL

studies performed in STARNET plasma samples. The authors would like to

acknowledge Dr. Anna Ataran (Washington University in St. Louis) for support in

sample collection. Schematic figures were created with BioRender.com.

743 **Sources of funding**

The authors' research work has received funding from the National Institutes of Health

(NIH) (R00HL130574, R01HL151611, R01HL168174, and P01HL172741) and the

National Center for Advancing Translational Sciences (NCATS) (Irving Scholar Award

through UL1TR001873), the Marjorie and Lewis Katz Scholar Award, and the M. Iréne

Ferrer Scholar Award (to H.Z.), the American Heart Association Postdoctoral Fellowship

20POST35130003 and Career Development Award 23CDA1052177 (to F.L.), the

Russell Berrie Foundation Diabetes Scholar Program (to X.W.), and the American Heart

Association Postdoctoral Fellowship 21POST829654 (to X.W.), VIDI (917.15.350) and

Aspasia grants from the Netherlands Organization of Scientific Research (NWO) (to

M.W.), and a Rosalind Franklin Fellowship from the University of Groningen with EU Co-

Fund attached (to M.W.), the NIH R01MH106842 (to T.L.) and F31HG010580 (to E.F.).

The GTEx Project was supported by the Common Fund of the Office of the Director of

the NIH, and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS.

References

1. McPherson R, and Tybjaerg-Hansen A. Genetics of Coronary Artery Disease. *Circulation Research.* 2016;118(4):564-78.
2. Consortium IKC. Large-scale gene-centric analysis identifies novel variants for coronary artery disease. *PLoS Genet.* 2011;7(9):e1002260.
3. Wild PS, Zeller T, Schillert A, Szymczak S, Sinning CR, Deiseroth A, et al. A genome-wide association study identifies LIPA as a susceptibility gene for coronary artery disease. *Circ Cardiovasc Genet.* 2011;4(4):403-12.
4. Coronary Artery Disease Genetics C. A genome-wide association study in Europeans and South Asians identifies five new loci for coronary artery disease. *Nat Genet.* 2011;43(4):339-44.
5. Lee JY, Lee BS, Shin DJ, Woo Park K, Shin YA, Joong Kim K, et al. A genome-wide association study of a coronary artery disease risk variant. *J Hum Genet.* 2013;58(3):120-6.
6. Consortium CAD, Deloukas P, Kanoni S, Willenborg C, Farrall M, Assimes TL, et al. Large-scale association analysis identifies new risk loci for coronary artery disease. *Nat Genet.* 2013;45(1):25-33.
7. Nikpay M, Goel A, Won HH, Hall LM, Willenborg C, Kanoni S, et al. A comprehensive 1,000 Genomes-based genome-wide association meta-analysis of coronary artery disease. *Nat Genet.* 2015;47(10):1121-30.
8. van der Harst P, and Verweij N. Identification of 64 Novel Genetic Loci Provides an Expanded View on the Genetic Architecture of Coronary Artery Disease. *Circ Res.* 2018;122(3):433-43.
9. Matsunaga H, Ito K, Akiyama M, Takahashi A, Koyama S, Nomura S, et al. Transethnic Meta-Analysis of Genome-Wide Association Studies Identifies Three New Loci and Characterizes Population-Specific Differences for Coronary Artery Disease. *Circ Genom Precis Med.* 2020;13(3):e002670.
10. Koyama S, Ito K, Terao C, Akiyama M, Horikoshi M, Momozawa Y, et al. Population-specific and trans-ancestry genome-wide analyses identify distinct and shared genetic risk loci for coronary artery disease. *Nat Genet.* 2020;52(11):1169-77.
11. Tcheandjieu C, Zhu X, Hilliard AT, Clarke SL, Napolioni V, Ma S, et al. Large-scale genome-wide association study of coronary artery disease in genetically diverse populations. *Nat Med.* 2022;28(8):1679-92.
12. Vargas-Alarcon G, Posadas-Romero C, Villarreal-Molina T, Alvarez-Leon E, Angeles J, Vallejo M, et al. Single nucleotide polymorphisms within LIPA (Lysosomal Acid Lipase A) gene are associated with susceptibility to premature coronary artery disease. a replication in the genetic of atherosclerotic disease (GEA) Mexican study. *PLoS One.* 2013;8(9):e74703.
13. Aragam KG, Jiang T, Goel A, Kanoni S, Wolford BN, Atri DS, et al. Discovery and systematic characterization of risk variants and genes for coronary artery disease in over a million participants. *Nat Genet.* 2022;54(12):1803-15.
14. Willer CJ, Schmidt EM, Sengupta S, Peloso GM, Gustafsson S, Kanoni S, et al. Discovery and refinement of loci associated with lipid levels. *Nat Genet.* 2013;45(11):1274-83.
15. Franzen O, Ermel R, Cohain A, Akers NK, Di Narzo A, Talukdar HA, et al. Cardiometabolic risk loci share downstream cis- and trans-gene regulation across tissues and diseases. *Science.* 2016;353(6301):827-30.
16. Li F, and Zhang H. Lysosomal Acid Lipase in Lipid Metabolism and Beyond. *Arterioscler Thromb Vasc Biol.* 2019;39(5):850-6.

806   17.   Porto AF. Lysosomal acid lipase deficiency: diagnosis and treatment of Wolman and
807   Cholesteryl Ester Storage Diseases. *Pediatr Endocrinol Rev.* 2014;12 Suppl 1:125-32.
808   18.   Korbelius M, Kuentzel KB, Bradic I, Vujic N, and Kratky D. Recent insights into
809   lysosomal acid lipase deficiency. *Trends Mol Med.* 2023;29(6):425-38.
810   19.   Evans TD, Zhang X, Clark RE, Alisio A, Song E, Zhang H, et al. Functional
811   Characterization of LIPA (Lysosomal Acid Lipase) Variants Associated With Coronary
812   Artery Disease. *Arterioscler Thromb Vasc Biol.* 2019;39(12):2480-91.
813   20.   Williams JW, Zaitsev K, Kim KW, Ivanov S, Saunders BT, Schrank PR, et al. Limited
814   proliferation capacity of aortic intima resident macrophages requires monocyte
815   recruitment for atherosclerotic plaque progression. *Nat Immunol.* 2020.
816   21.   Hao K, Ermel R, Sukhavasi K, Cheng H, Ma L, Li L, et al. Integrative Prioritization of
817   Causal Genes for Coronary Artery Disease. *Circ Genom Precis Med.*
818   2022;15(1):e003365.
819   22.   Barbeira AN, Dickinson SP, Bonazzola R, Zheng J, Wheeler HE, Torres JM, et al.
820   Exploring the phenotypic consequences of tissue specific gene expression variation
821   inferred from GWAS summary statistics. *Nat Commun.* 2018;9(1):1825.
822   23.   Wen X, Pique-Regi R, and Luca F. Integrating molecular QTL data into genome-wide
823   genetic association analysis: Probabilistic assessment of enrichment and colocalization.
824   *PLoS Genet.* 2017;13(3):e1006646.
825   24.   Zhu Z, Zhang F, Hu H, Bakshi A, Robinson MR, Powell JE, et al. Integration of summary
826   data from GWAS and eQTL studies predicts complex trait gene targets. *Nat Genet.*
827   2016;48(5):481-7.
828   25.   Fernandez JM, de la Torre V, Richardson D, Royo R, Puiggros M, Moncunill V, et al. The
829   BLUEPRINT Data Analysis Portal. *Cell Syst.* 2016;3(5):491-5 e5.
830   26.   van de Geijn B, McVicker G, Gilad Y, and Pritchard JK. WASP: allele-specific software
831   for robust molecular quantitative trait locus discovery. *Nat Methods.* 2015;12(11):1061-3.
832   27.   Turner AW, Hu SS, Mosquera JV, Ma WF, Hodonsky CJ, Wong D, et al. Single-nucleus
833   chromatin accessibility profiling highlights regulatory mechanisms of coronary artery
834   disease risk. *Nat Genet.* 2022;54(6):804-16.
835   28.   Soehnlein O, and Libby P. Targeting inflammation in atherosclerosis - from experimental
836   insights to the clinic. *Nat Rev Drug Discov.* 2021;20(8):589-610.
837   29.   Zhang H, Shi J, Hachet MA, Xue C, Bauer RC, Jiang H, et al. CRISPR/Cas9-Mediated
838   Gene Editing in Human iPSC-Derived Macrophage Reveals Lysosomal Acid Lipase
839   Function in Human Macrophages-Brief Report. *Arterioscler Thromb Vasc Biol.*
840   2017;37(11):2156-60.
841   30.   Nurnberg ST, Zhang H, Hand NJ, Bauer RC, Saleheen D, Reilly MP, et al. From Loci to
842   Biology: Functional Genomics of Genome-Wide Association for Coronary Disease. *Circ
843   Res.* 2016;118(4):586-606.
844   31.   Pham TH, Benner C, Lichtinger M, Schwarzfischer L, Hu Y, Andreesen R, et al. Dynamic
845   epigenetic enhancer signatures reveal key transcription factors associated with
846   monocytic differentiation states. *Blood.* 2012;119(24):e161-71.
847   32.   Zhu Y, Rosenfeld MG, and Suh Y. Ultrafine mapping of chromosome conformation at
848   hundred basepair resolution reveals regulatory genome architecture. *Proc Natl Acad Sci
849   U S A.* 2023;120(45):e2313285120.
850   33.   Flynn ED, Tsu AL, Kasela S, Kim-Hellmuth S, Aguet F, Ardlie KG, et al. Transcription
851   factor regulation of eQTL activity across individuals and tissues. *PLoS Genet.*
852   2022;18(1):e1009719.
853   34.   Nerlov C, and Graf T. PU.1 induces myeloid lineage commitment in multipotent
854   hematopoietic progenitors. *Genes Dev.* 1998;12(15):2403-12.
855   35.   Kheradpour P, and Kellis M. Systematic discovery and characterization of regulatory
856   motifs in ENCODE TF binding experiments. *Nucleic Acids Res.* 2014;42(5):2976-87.

857   36.   Abram CL, Roberge GL, Hu Y, and Lowell CA. Comparative analysis of the efficiency
858   and specificity of myeloid-Cre deleting strains using ROSA-EYFP reporter mice. *J*
859   *Immunol Methods.* 2014;408:89-100.
860   37.   Wang X, Magkos F, and Mittendorfer B. Sex differences in lipid and lipoprotein
861   metabolism: it's not just about sex hormones. *J Clin Endocrinol Metab.* 2011;96(4):885-
862   93.
863   38.   Cai B, Thorp EB, Doran AC, Sansbury BE, Daemen MJ, Dorweiler B, et al. MerTK
864   receptor cleavage promotes plaque necrosis and defective resolution in atherosclerosis.
865   *J Clin Invest.* 2017;127(2):564-8.
866   39.   Potteaux S, Gautier EL, Hutchison SB, van Rooijen N, Rader DJ, Thomas MJ, et al.
867   Suppressed monocyte recruitment drives macrophage removal from atherosclerotic
868   plaques of Apoe-/- mice during disease regression. *J Clin Invest.* 2011;121(5):2025-36.
869   40.   Georgakis MK, van der Laan SW, Asare Y, Mekke JM, Haitjema S, Schoneveld AH, et al.
870   Monocyte-Chemoattractant Protein-1 Levels in Human Atherosclerotic Lesions Associate
871   With Plaque Vulnerability. *Arterioscler Thromb Vasc Biol.* 2021;41(6):2038-48.
872   41.   Boring L, Gosling J, Cleary M, and Charo IF. Decreased lesion formation in CCR2-/-
873   mice reveals a role for chemokines in the initiation of atherosclerosis. *Nature.*
874   1998;394(6696):894-7.
875   42.   Xue S, Tang H, Zhao G, Fang C, Shen Y, Yan D, et al. C-C motif ligand 8 promotes
876   atherosclerosis via NADPH oxidase 2/reactive oxygen species-induced endothelial
877   permeability increase. *Free Radic Biol Med.* 2021;167:181-92.
878   43.   Tsou CL, Peters W, Si Y, Slaymaker S, Aslanian AM, Weisberg SP, et al. Critical roles for
879   CCR2 and MCP-3 in monocyte mobilization from bone marrow and recruitment to
880   inflammatory sites. *J Clin Invest.* 2007;117(4):902-9.
881   44.   Kim K, Shim D, Lee JS, Zaitsev K, Williams JW, Kim KW, et al. Transcriptome Analysis
882   Reveals Nonfoamy Rather Than Foamy Plaque Macrophages Are Proinflammatory in
883   Atherosclerotic Murine Models. *Circ Res.* 2018;123(10):1127-42.
884   45.   Wang X, Collins HL, Ranalletta M, Fuki IV, Billheimer JT, Rothblat GH, et al. Macrophage
885   ABCA1 and ABCG1, but not SR-BI, promote macrophage reverse cholesterol transport
886   in vivo. *J Clin Invest.* 2007;117(8):2216-24.
887   46.   Willemsen L, Chen HJ, van Roomen C, Griffith GR, Siebeler R, Neele AE, et al.
888   Monocyte and Macrophage Lipid Accumulation Results in Down-Regulated Type-I
889   Interferon Responses. *Front Cardiovasc Med.* 2022;9:829877.
890   47.   Duewell P, Kono H, Rayner KJ, Sirois CM, Vladimer G, Bauernfeind FG, et al. NLRP3
891   inflammasomes are required for atherogenesis and activated by cholesterol crystals.
892   *Nature.* 2010;464(7293):1357-61.
893   48.   de la Roche M, Hamilton C, Mortensen R, Jeyaprakash AA, Ghosh S, and Anand PK.
894   Trafficking of cholesterol to the ER is required for NLRP3 inflammasome activation. *J*
895   *Cell Biol.* 2018;217(10):3560-76.
896   49.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.
897   Comprehensive Integration of Single-Cell Data. *Cell.* 2019;177(7):1888-902 e21.
898   50.   Sharma D, Worssam MD, Pedroza AJ, Dalal AR, Alemany H, Kim HJ, et al.
899   Comprehensive Integration of Multiple Single-Cell Transcriptomic Data Sets Defines
900   Distinct Cell Populations and Their Phenotypic Changes in Murine Atherosclerosis.
901   *Arterioscler Thromb Vasc Biol.* 2024;44(2):391-408.
902   51.   Lin PK, and Davis GE. Extracellular Matrix Remodeling in Vascular Disease: Defining Its
903   Regulators and Pathological Influence. *Arterioscler Thromb Vasc Biol.* 2023;43(9):1599-
904   616.
905   52.   Cybulsky MI, Iiyama K, Li H, Zhu S, Chen M, Iiyama M, et al. A major role for VCAM-1,
906   but not ICAM-1, in early atherosclerosis. *J Clin Invest.* 2001;107(10):1255-62.

53. Cochain C, Vafadarnejad E, Arampatzi P, Pelisek J, Winkels H, Ley K, et al. Single-Cell
RNA-Seq Reveals the Transcriptional Landscape and Heterogeneity of Aortic
Macrophages in Murine Atherosclerosis. *Circ Res.* 2018;122(12):1661-74.
54. Endo-Umeda K, Kim E, Thomas DG, Liu W, Dou H, Yalcinkaya M, et al. Myeloid LXR
(Liver X Receptor) Deficiency Induces Inflammatory Gene Expression in Foamy
Macrophages and Accelerates Atherosclerosis. *Arterioscler Thromb Vasc Biol.*
2022;42(6):719-31.
55. Morris GE, Braund PS, Moore JS, Samani NJ, Codd V, and Webb TR. Coronary Artery
Disease-Associated LIPA Coding Variant rs1051338 Reduces Lysosomal Acid Lipase
Levels and Activity in Lysosomes. *Arterioscler Thromb Vasc Biol.* 2017;37(6):1050-7.
56. Jerome WG, Cox BE, Griffin EE, and Ullery JC. Lysosomal cholesterol accumulation
inhibits subsequent hydrolysis of lipoprotein cholesteryl ester. *Microsc Microanal.*
2008;14(2):138-49.
57. Singh RK, Haka AS, Bhardwaj P, Zha X, and Maxfield FR. Dynamic Actin Reorganization
and Vav/Cdc42-Dependent Actin Polymerization Promote Macrophage Aggregated LDL
(Low-Density Lipoprotein) Uptake and Catabolism. *Arterioscler Thromb Vasc Biol.*
2019;39(2):137-49.
58. Balwani M, Breen C, Enns GM, Deegan PB, Honzik T, Jones S, et al. Clinical effect and
safety profile of recombinant human lysosomal acid lipase in patients with cholesteryl
ester storage disease. *Hepatology.* 2013;58(3):950-7.
59. Lopresti MW, Cui W, Abernathy BE, Fredrickson G, Barrow F, Desai AS, et al. Hepatic
lysosomal acid lipase overexpression worsens hepatic inflammation in mice fed a
Western diet. *J Lipid Res.* 2021;62:100133.
60. Nelson CP, Goel A, Butterworth AS, Kanoni S, Webb TR, Marouli E, et al. Association
analyses based on false discovery rate implicate new loci for coronary artery disease.
Nat Genet. 2017;49(9):1385-91.
61. Consortium GT. Human genomics. The Genotype-Tissue Expression (GTEx) pilot
analysis: multitissue gene regulation in humans. *Science.* 2015;348(6235):648-60.

Figures and figure legends

Figure 1. Genetic variants at the *LIPA* locus associated with higher risks of coronary artery disease (CAD) are also linked to higher *LIPA* mRNA, enzyme activity, and protein in human peripheral blood monocytes-derived macrophages (HMDM). A, LocusZoom plot to visualize the CARDIoGRAMplusC4D CAD GWAS signals at the *LIPA* locus. The GWAS lead SNP rs1412444 and the SNPs in linkage disequilibrium (LD) are color-coded based on pairwise r^2 with rs1412444. **B,** Colocalization analysis by integrating CARDIoGRAMplusC4D GWAS dataset with the

Genotype-Tissue Expression project (GTEx) eQTL in the blood shows that *LIPA* is the

only nearby gene at the locus (± 1 Mb flanking the *LIPA* gene) showing strong regional

colocalization probability (rcp = 0.657). **C,** eQTL analysis of *LIPA* in the BLUEPRINT

dataset shows strong eQTL signals in monocytes (blue), but not T-cells (green) or

neutrophils (red). Three eQTL methods, including WASP-corrected combined haplotype

test (WASP-CHT, triangle), WASP-corrected allele-specific expression (WASP-ASE,

square), and BLUEPRINT Work Package 10 (WP10, circle), were employed in the

analysis. **D-G,** LocusZoom plots to visualize *LIPA* eQTLs in atherosclerotic aortic wall

(Aorta, **D**), blood (**E**), HMDM (**F**), and foamy HMDM (HMDM loaded with acetylated low-

density lipoprotein, **G**) in the Stockholm-Tartu Atherosclerosis Reverse Network

Engineering Task (STARNET) dataset, which comprises nine cardiometabolic

tissues/cell types from subjects with CAD. The most significant SNP is marked by purple

diamond. Other SNPs in the region are color-coded by their LD with the lead SNP

according to pairwise r^2. **H-J**, The risk allele (T) of the CAD lead SNP rs1412444 is

associated with increased *LIPA* mRNA (**H**), enzyme activity (**I**), and protein (**J**) in HMDM.

n = the number of subjects. Data are presented as mean ± SEM in (**H**) and median ± 95%

CI in (**I**) and (**J**).

Figure 2. The non-coding genomic region containing rs1412444 is a myeloid-specific enhancer, interacting with the *LIPA* promoter and regulating *LIPA* expression. A, Genome Browser view of annotation tracks for the *LIPA* locus showing significant CAD GWAS SNPs, including the lead SNP (rs1412444) and SNPs in linkage disequilibrium (LD) as listed in **Table S2**, *LIPA* transcripts, and the regulatory landscape in human monocyte and HMDM. The potential enhancer regions showing H3K4me1 and H3K27ac modifications and hosting CAD GWAS SNPs, including the rs1412444 region and the rs2246883 region, are highlighted and prioritized for subsequent functional validation. **B,** The rs1412444 region, but not the rs2246833 region, shows

enhancer activity by luciferase assay in THP-1 monocytes (**left**). Neither region shows

enhancer activities in HEK 293 cells (**right**). (n = 3 independent experiments with 3

technical replicates, data are presented as median ± 95% CI.) **C,** High-resolution Tri-

HiC depicts chromatin interaction between the rs1412444 region and the *LIPA* promoter

in human CD14$^+$ monocytes, but not CD3$^+$ T-cells. **D,** CRISPRi targeting the rs1412444

enhancer region reduces *LIPA* expression. sgRNAs targeting the rs1412444 region

were designed and transduced to dCas9-KRAB expressing THP-1 cells for CRISPRi-

mediated silencing and mRNA expression of *LIPA* was determined by quantitative RT-

PCR. The sgRNA1 is 68 bp upstream of rs1412445 and is the highest scored sgRNA

within the region by CRISPick. The sgRNA2 span rs1320496. The sgRNA3 span

rs1412445. The rs1412444 SNP region lacks a PAM sequence for sgRNA design. The

sgRNA targeting the transcription start site of *LIPA* (*LIPA*-TSS) serves as the positive

control. (n = 6 replicates, data are presented as mean ± SEM.)

Figure 3. rs1320496 and rs1412444 represent the functional SNPs at the *LIPA*

locus, with risk alleles enhancing PU.1 binding and regulating *LIPA* expression. A,

Site-directed mutagenesis and luciferase assay confirmed that the risk haplotypes TTT

(rs1412445-T, rs1320496-T, and rs1412444-T) and CTC (rs1412445-C, rs1320496-T,

and rs1412444-C) lead to increased enhancer activities compared to the non-risk

haplotype CCC (rs1412445-C, rs1320496-C, and rs1412444-C). (n = 5 independent

experiments with 3 technical replicates, data are presented as median ± 95% CI.) **B,**

Conditional analysis was performed to assess the independent contribution of SNPs

(rs1412445, rs1320496, and rs141244) within the rs1412444-containing enhancer to

eQTL and GWAS signals. The perfect LD between rs1412444 and rs1412445 (r^2 = 0.99)

allows for the utilization of rs1412445 as a representative to capture the effects of both

variants. The high LD (r^2 = 0.45) between rs1320496 and the other two variants

prohibited full disentanglement. Both rs1412445 (blue) and rs1320496 (red) show

independent association with *LIPA* expression (left, in blood, P < 10^{-59}) and CAD risk

(right, P < 10^{-11}). After conditioning on rs1412445, rs1320496 remains showing

nominally significant effects on both *LIPA* expression (left, in blood, P = 3 × 10^{-10}) and

CAD risk (right, P = 0.031) (green, rs1320496 conditional), suggesting that both SNPs

independently contributing to the traits. In the context of eQTL analysis, the effect size is

quantified as the $\log_2$ allelic fold change in *LIPA* gene expression, while in GWAS, it

represents the log odds change in CAD risks. **C,** Allele-specific binding analysis of PU.1

ChIP-seq data in HMDM (GSM785501, n = 1 experiment) and LCL cell line (EBV-

transformed lymphoblastoid B-cell lines) (ENCODE dataset, n = 2 independent

experiments). Results suggest that PU.1 binds more favorably to the risk alleles (T). **D,**

Motif analysis shows that the risk allele (T, red) of rs1320496 creates a PU.1 binding

site. **E**, Electrophoretic Mobility Gel Shift Assay (EMSA) was conducted to determine the

effects of different alleles on PU.1 binding for sequences containing SNP rs1320496

(left) and rs1412444 (right), using recombinant human PU.1 protein and anti-PU.1

antibody. Data shown are representative images from n = 2 independent experiments. **F**,

siRNA-mediated knockdown of *SPI1* (encoding PU.1) in THP-1 cells reduces mRNA

expression of *LIPA*. (n = 3 independent experiments with 2 technical replicates, data are

presented as median ± 95% CI.) **G**, Knockdown of *SPI1* partly abolished the effects of

risk alleles on increasing the enhancer activity of the rs1412444 region. Data were

analyzed by two-way ANOVA followed by Tukey's multiple comparisons test. (n = 3

independent experiments with 3 technical replicates, data are presented as median ± 95%

CI.) **H**, Schematic figure summarizing results of functional genomic studies that the risk

alleles of the functional variants at *LIPA* risk locus increases the expression of *LIPA* by

enhancing PU.1 binding in monocytes/macrophages and via enhancer-promoter

interaction.

1026

1027

Figure 4. Myeloid overexpression of *Lipa* exacerbates atherosclerosis in *Ldlr*^{-/-} mice. A, The plasmid construction and breeding strategy of the mouse model. Mice for conditional overexpression of *Lipa* (NM_001111100) were generated by *Rosa26* knock-in (KI) of CAG-loxP-STOP-loxP-Lipa-IRES-eGFP. To achieve myeloid-specific overexpression, *Lipa*^{KI/KI} mice were bred with *LysMCre*^{+/-} mice (heterozygous for the Cre allele) to delete the *loxP-STOP-loxP* cassette, thereby enabling overexpression of *Lipa* in myeloid cells in *LysMCre*^{+/-}, *Lipa*^{KI/WT} mice. Littermates with the genotype *LysMCre*^{-/-}, *Lipa*^{KI/WT} without overexpression serve as controls. To assess the impact of myeloid-specific overexpression of *Lipa* on atherosclerosis, these mice were bred onto an *Ldlr*^{-/-} background. To induce atherosclerosis, *Ctrl* (*LysMCre*^{-/-}, *Lipa*^{KI/WT}, *Ldlr*^{-/-}) and *M-Lipa*^{KI} (*LysMCre*^{+/-}, *Lipa*^{KI/WT}, *Ldlr*^{-/-}) mice were fed a Western diet (WD) for 16 weeks. The atherosclerotic lesion size and features of plaque stability in the aortic sinus were quantified. **B,** Myeloid overexpression of *Lipa* modestly but significantly increased atherosclerotic lesion size in mice of both sexes (left: combined, n = 23 mice), as well as

in male mice (middle: n = 11) and female mice (right: n= 12) when analyzed separately.

Data are presented as median ± 95% CI. Scale bar = 200 μm. **C**, Atherosclerotic lesion

size is positively correlated with LIPA enzyme activity in peritoneal macrophages (PMs)

(two-tailed Pearson's correlation analysis, P = 0.014, r = 0.4499). **D**, No differences in

the percentage of necrotic core area between the two genotypes (n = 11 male mice and

12 female mice). **E**, Sirius red staining shows comparable fibrous cap thickness

between the two genotypes (n = 8 male mice). Data are presented as mean ± SEM.

Scale bar = 200 μm for (**D**) and (**E**).

1050

Figure 5. Myeloid overexpression of *Lipa* increases macrophage content in the atherosclerotic plaque by promoting monocyte recruitment. A, Immunofluorescence (IF) staining data suggest that myeloid overexpression of *Lipa* significantly increased the CD68+ [Magenta] macrophage area in the lesion in mice of both sexes (left: combined, n = 17 mice), as well as in male mice (middle: n = 12) and female mice (right: n = 5). Data are presented as median ± 95% CI. The white dashed contour indicates lesion area. Scale bar = 200 µm. **B-C**, IF staining and quantification for the proliferative macrophages (**B**, Ki67 [Red], CD68+ [Cyan], and DAPI [Blue]) and apoptotic macrophages (**C**, TUNEL [Red], CD68+ [Cyan], and DAPI [Blue]). (**B**, n = 13

male mice and 7 female mice; **C**, n = 13 male mice and 8 female mice). Data are presented as mean ± SEM. Scale bars = 50 µm. **D**, Monocyte infiltration was assessed by beads assay. Briefly, Ly6C^{hi} monocytes in the blood were pulse-labeled with fluorescent beads by transiently depleting circulating monocytes with clodronate liposome injection. Beads labeling efficiency was assessed by flow cytometry as the % of bead^{+}Ly6C^{hi} monocytes in Ly6C^{hi} monocytes 24 h after beads injection. The number of beads in the CD68^{+} area of atherosclerotic lesion was quantified in mice three days post-injection. Results are presented as the number of beads in the CD68^{+} area per section (left), the number of beads in the CD68^{+} area per section normalized by the percentage of bead^{+} monocytes in Ly6C^{hi} monocytes (middle), and representative IF staining showing beads^{+} [Green] CD68^{+} macrophages [Red] in the lesion (right). For *Ctrl*, n = 3 male mice and 8 female mice; for *M-Lipa^{KI}*, n = 3 male mice and 8 female mice; the average of two sections per mouse were reported. Data are presented as mean ± SEM. Scale bars = 200 µm.

1076

Figure 6. Myeloid overexpression of *Lipa* leads to reduced neutral lipid but increased free cholesterol accumulation in macrophages. A, Schematics of a flow cytometry-based method to characterize aortic macrophages (CD45⁺CD11b⁺CD64⁺) that are neutral lipid-enriched and foamy (SSCʰⁱLipidTOXʰⁱ) in aortas dissected from *Ctrl* and *M-Lipaᴷᴵ* mice fed a WD for 16 weeks. **B,** Percentage of foamy aortic macrophages in total aortic macrophages in *Ctrl* and *M-Lipaᴷᴵ* mice. **C**, Mean fluorescent intensity (MFI) of LipidTOX in foamy aortic macrophages in *Ctrl* and *M-Lipaᴷᴵ* mice. **D,** MFI of LipidTOX in GFP⁻ and GFP⁺ foamy aortic macrophages in *M-Lipaᴷᴵ* mice. **B-D**, n = 6 male mice for *Ctrl*; n = 7 male mice for *M-Lipaᴷᴵ*. Data are presented as mean ± SEM. **E,** Schematics of study design to characterize foamy (SSCʰⁱLipidTOXʰⁱ) PMs (CD45⁺F4/80⁺) in *Ctrl* and

1087 *M-Lipa^{KI}* mice fed a WD for 16 weeks. **F,** Percentage of foamy PMs in PMs of *Ctrl* and

1088 *M-Lipa^{KI}* mice. **G,** MFI of LipidTOX in foamy PMs in *Ctrl* and *M-Lipa^{KI}* mice. **F-G**, n = 6

1089 male mice. Data are presented as mean ± SEM. **H,** F4/80$^+$ PMs from *Ctrl* mice and

1090 F4/80$^+$GFP$^+$ PMs from *M-Lipa^{KI}* mice that fed a WD for 16 weeks were sorted for

1091 quantification of cellular cholesterol levels in (**I**). **I,** Quantification of total cholesterol (left),

1092 free cholesterol (second left), cholesteryl ester (second right), and ratio of free

1093 cholesterol/cholesteryl ester (right). (n = 5 female mice, with analysis based on the

1094 averages of two technical replicates. Data are presented as median ± 95% CI.)

1095

1096

Figure 7. Bulk RNA-seq analyses reveal distinct effects of myeloid overexpression of *Lipa* on aortic macrophages and PMs. A, Viable macrophages (CD45⁺CD11b⁺CD64⁺) were sorted from aortas of *Ctrl* and *M-Lipa^KI* mice fed a WD for 16 weeks and subjected to low-input RNA-seq. n = 4 biological replicates per genotype, with each replicate comprising pooled samples from 2-3 male mice, resulting a total of 10 male mice per genotype. **B**, Volcano plot to visualize the top DE genes (aortic macrophages isolated from *M-Lipa^KI* mice *versus* those from *Ctrl* mice, all fed a WD for 16 weeks). **C**, The top enriched pathways in the upregulated genes in aortic

macrophages of *M-Lipa^{KI}* mice as determined by Gene Set Enrichment Analysis (GSEA).

D, Heatmap visualization of top DE genes and the leading-edge subsets of the

associated integrin signaling and cell-matrix mediated adhesion pathways. **E**, F4/80$^+$

PMs from *Ctrl* mice and F4/80$^+$GFP$^+$ PMs from *M-Lipa^{KI}* mice fed a WD for 16 weeks

were sorted for bulk RNA-seq. n = 3 female mice. **F,** Volcano plot to visualize the top DE

genes (F4/80$^+$GFP$^+$ PMs from *M-Lipa^{KI}* mice *vs.* F4/80$^+$ PMs from *Ctrl* mice, all fed a

WD for 16 weeks). **G,** The top enriched pathways in the upregulated genes in *Lipa-*

overexpressing PMs as determined by GSEA. **H-I,** Heatmap visualization of top DE

genes and the leading-edge subsets of the associated interferon response pathways (**H**)

and antigen processing and presentation pathways (**I**). **F-I**, n = 3 female mice.

1115

1116

Figure 8. Single-cell RNA-seq (scRNA-seq) analyses suggest that myeloid

overexpression of *Lip*a alters both the proportions and transcriptomic profiles of

different aortic cell types in atherosclerosis, including smooth muscle cells. A,

Viable cells isolated from aortas of *Ctrl* and *M-Lipa^{KI}* mice fed a WD for 16 weeks were

subjected to scRNA-seq. Cells were obtained from a pooled sample of n = 5 male mice.

B, Uniform Manifold Approximation and Projection (UMAP) visualization of 6 cell types

identified from the analyses. **C**, Dot plot visualization of top marker genes of each cell

type. **D**, Stacked bar plot shows the proportion of each cell type in *Ctrl* (left) and *M-*

Lipa^{KI} (right) aortic cells. *M-Lipa^{KI}* mice show increased macrophages and fibroblasts,

while decreased smooth muscle cells (SMCs). Data were analyzed by Chi-square test

with Bonferroni correction. **E**, Sub-clustering analysis of SMCs and UMAP visualization

of 6 SMC subclusters. **F**, Stacked bar plot shows the proportion of each SMC subcluster

in *Ctrl* (left) and *M-Lipa^{KI}* (right) SMCs. Data were analyzed by Chi-square test with

Bonferroni correction. **G**, Top activated canonical pathways by Ingenuity Pathway

Analysis (IPA) in SMCs. **H**, Top activated canonical pathways by IPA in SMC1. **I**, Top

inhibited canonical pathways by IPA in Cxcl12 SMC.