Title:

Exploring efficiency landscapes in the acoustic-morphospace of crickets reveals two alternative calling strategies

Authors: Erin E. Brandt[1], Sarah Duke[1,2], Honglin Wang[1,3], and Natasha Mhatre[1]

Affiliations:

1. Department of Biology, Western University, London, Ontario. Canada
2. Present address: Michener Institute of Education, University Health Network, Toronto, Ontario. Canada
3. Present address: Schulich School of Medicine and Dentistry, Western University, London, Ontario. Canada

Abstract

Male crickets attract females by rubbing forewings together to produce calls. Louder calls are more effective, as they travel further, and attract more mates. However, crickets are small and produce calls inefficiently. Tree crickets make baffles, which reduce acoustic short-circuiting and increase efficiency. Previous work showed that tree cricket baffles function optimally (Mhatre et al., 2017). Here, we ask why baffle use is uncommon in crickets despite its advantages. We hypothesize that baffling may be rare because it is advantageous only for certain species. To test this, we modelled the calling efficiencies of cricket wings within their acoustic-morphospace (ie: the complete space of natural wing sizes and call frequency range). Using finite and boundary element analysis we calculated cricket acoustic efficiency in multiple of acoustic environments, incorporating both reflective and scattering surfaces like the ground and vegetation into our analyses. Within the efficiency landscapes generated from these data, we plotted the positions of 111 species across 7 phylogenetic clades. Using landscape and clade level analyses, we found that calling from the ground and using a baffle represent effective alternate strategies that both maximize calling efficiency.

Keywords: insect sound production | crickets | tool use | finite element modeling | boundary element modeling | sound propagation | acoustic efficiency

Introduction

Male crickets make loud advertisement calls to attract females who use these calls to locate mates (Regen, 1913). Louder calls travel further, cover more area, and attract more females (Deb et al., 2020; Farris et al., 1997; Römer et al., 1993). When faced with a choice, females prefer louder calls (Deb et al., 2020; Forrest, 1991). Being louder therefore has implications for mating success and evolutionary fitness in these singing insects. However, despite the apparent loudness of a nighttime chorus, cricket calls are acoustically constrained by a phenomenon known as 'acoustic short-circuiting' (Forrest, 1982; Mhatre et al., 2017). The smaller the wings of a cricket with respect to the wavelength of the sound it makes, the

35 higher the short-circuiting and associated loss of efficiency. Indeed, the few crickets that have been
36 studied are small and experience significant short-circuiting (Bennet-Clark, 1998).

Baffle use is an effective – but rare – means to overcome short-circuiting
To reduce the efficiency lost to acoustic short-circuiting, a few tree cricket species build and use baffles
(Forrest, 1991, 1982; Mhatre et al., 2017; Prozesky-Schulze et al., 1975). A baffle consists of a leaf with a
hole chewed by the cricket near the middle of the leaf. When the size of the leaf and hole are optimal,
such structures reduce acoustic short-circuiting and increase efficiency by as much as 10 dB compared to
unbaffled calling, reflecting a tripling of sound pressure levels (Mhatre et al., 2017). However, despite
their obvious benefits, only a handful of species among thousands make baffles, all within the sub-family
Oecanthinae (Collins, 2012; Forrest, 1991; Mhatre et al., 2017; Prozesky-Schulze et al., 1975).

Baffle use in crickets may be rare due to a "lack of utility"
Given the obvious benefits, why is acoustic baffle use rare in crickets? Some researchers believe that tree
cricket baffles fit the criteria for manufactured tools (Mhatre, 2018; Pierce, 1986) and, two hypotheses
from the tool use literature, the "cognitive capacity" and the "lack of utility" hypotheses offer two
different reasons. The "cognitive capacity" hypothesis suggests that complex tool use behaviors are less
likely to evolve in animals with smaller brains and lower cognitive capacity. This is an unlikely
explanation since many animals with relatively low cognitive capacities do use, and even make, tools
which themselves are not necessarily complex objects. A competing hypothesis is the "lack of utility"
hypothesis. This posits that tool and building behavior generally can evolve regardless of cognitive
capacity, but that its evolution requires an ecological context in which it confers sufficient selective
advantage (Hansell and Ruxton, 2008). Stated another way, only species that can achieve higher gains
from tool use than from other strategies (e.g., morphological features, site selection) are likely to evolve
tool using behavior. Indeed, few species use tools, whether crickets, other invertebrates or even
vertebrates (Hunt et al., 2013), but invertebrate tool use seems especially rare. For example, 56
independent occurrences of subsistence-related tool use were found in mammals, whereas only 13
occurrences were found in the significantly more speciose insects.

However, to truly test the lack of utility hypothesis, we must quantify tool utility and use of the tool must
have implications for evolutionary fitness. It is often difficult to meet these two conditions. Work in
chimpanzees has directly quantified tool utility by evaluating how much caloric value can be gained by
using a tool to exploit an otherwise unexploitable resource (Boesch and Boesch, 1983). Other studies
have made more indirect arguments about the utility of tools. Work in sea otters has shown that tools are
employed more frequently in populations in which tough prey require tools to access them (Fujii et al.,
2015). In capuchin monkeys, larger individuals who can more effectively use tools to crack nuts are more
likely to use tools (Spagnoletti et al., 2011). However, few studies directly assess the lack of utility
hypothesis, particularly outside the context of food.

Baffle use in crickets is an ideal system in which to test the lack of utility hypothesis. First, baffle use is
rare and second, we can directly measure its acoustic utility in terms of increase in calling efficiency (ref
Mhatre et al 2018). Finally, baffle use has been shown to have real fitness implications, by increasing the
number of mates attracted to a given male, and also by increase mating duration, both processes likely to
increase reproductive success (Deb et al., 2020).

Therefore, in this study, we tested the lack of utility hypothesis across this large group of singing insects,
the true crickets or Grylloidea. We used mathematical models to quantify baffle utility in two ways. First,
we ascertained the sizes of call radiator (cricket wings) and frequency ranges of the songs used by a large
number of crickets, spread over the cricket phylogeny. Then we quantified sound radiation efficiency
(SRE), with and without "perfect" baffles, for this complete parameter space which we call the acoustic-
morphospace. SRE captured the efficiency of the singer, by calculating local losses due to acoustic short
circuiting, similar to the previous study on tree cricket baffle efficiency (Mhatre et al., 2017). By plotting
SRE over the acoustic-morphospace, we were able to generate efficiency landscapes to fully investigate
the calling efficiency of crickets with and without baffles.

Finite element methods are known to be accurate at predicting sound levels and fields in acoustics
modeling, and have been used previously to model sound fields in crickets (Garud et al., 2017; Godthi et
al., 2022; Mhatre et al., 2017; Seybert et al., 1994). However, so far models have only considered very
idealized physical conditions. However, crickets live in real environments populated by objects such as
the ground and vegetation, which interact with sound and its radiator across spatial scales, which may
form the basis of an alternative strategy (Erregger and Schmidt, 2018; Muñoz and Halfwerk, 2022).
Therefore, we generated a second more ecological metric of utility, which we have called sound
propagation efficiency (SPE). By using the boundary element method, we quantified the effect of such
interacting acoustic surfaces and their effect on acoustic efficiency as sound propagates away from the
singer, under a range of environmental conditions. Since we considered calling from the ground, this
raised the issue of directionality. The spatial gradient of the sound field is also crucial to its utility, and
may be degraded preventing mate finding when animals call from near the ground (Embleton, 1996;
Kostarakos and Römer, 2010). Therefore, using the same models, we also quantified directionality to test
how efficiency might trade off with this biologically crucial feature.

Using these data, we asked whether the rarity of baffle use in crickets is explained by the lack of utility
hypothesis. We examined the differences between baffled and unbaffled calling in different realistic
scenarios. If the lack of utility hypothesis is supported, we predict that there would be at least two
alternative strategies to maximize calling efficiency. Known baffle-users will be animals who benefit
more from baffle use than non baffle-users who might instead use a morphological or environmental
means to maximize efficiency.

Results and Discussion

All crickets would benefit from baffle use in idealized conditions
To capture the natural range of wing sizes and calling frequencies among true crickets, we collected wing
surface area and call frequency data for 111 cricket species from a large range of sources (Figure 1,
Tables 1, 2, Supplement to Figure 1). Species were distributed across 7 clades as described by the most
recent phylogeny of the Grylloidae or "true cricket" super family (Chintauan-Marquier et al., 2016)
(Figure 1).

We then constructed finite element models which predicted the sound fields produced by wings of
different sizes at different call frequencies for 1086 different combinations which encompassed all the
observed frequencies and wing sizes , i.e. the full acoustic-morphospace (Figure 2). In all conditions,
wings were modelled as suspended in free space, vibrating with a uniform velocity perpendicular to the

wing plane (Figure Supplement 2-1, 2-3). The model then predicts the resulting sound field (see methods for details). We then calculate sound radiation efficiency (SRE (Pa/m/s)), by taking a volumetric average of the sound pressure level generated (Pa), over a sphere of radius 20 cm around the wing, divided by the time-space average of the wing vibration velocity (m/s). Since we do not have data on wing vibration velocities for most cricket species, this normalized measure of efficiency enabled comparison between species. This method of applying a velocity across the entire wing and the resulting normalized metric has also been used previously for tree crickets (Mhatre et al., 2017), and does not generate substantially different prediction from only vibrating part of the wing (Supplement to figure 2-4).

Next we plotted an SRE landscape (Figure 2) for the full acoustic-morphospace. On this landscape, we plotted the locations of the 111 species of crickets with known wing size and frequency allowing us to infer their SRE (Figure 2). These data indicate how efficient each wing should be at radiating sound, reflecting the match between radiator size and the wavelength of the radiated sound frequency. These data therefore demonstrate precisely how suboptimal crickets are in terms of their efficiency (Figure 2D), and how much they could gain through use of an optimal baffle (Figure 2E).

Previous work examining four species of crickets and a small number of other insects determined that they each perform less efficiently than the theoretical optimal level for dipole sound sources (Bennet-Clark, 1998). However, our finite element methods model more realistic geometries than these previous methods, which were restricted to analytically tractable shapes like circular pistons. In our dataset, there is a clear ridge of high efficiency running through the SRE landscape (red line in figure 2A, hereafter referred to as "optimal efficiency ridge"), which shows the optimal radiator size for every frequency. Despite sampling a wide range of wing areas ($0.4 - 258$ mm^2) and call frequencies ($1.6 - 27.9$ kHz), all species in our dataset occupy a region of the SRE landscape below this optimal efficiency ridge. The efficiency distribution among crickets is somewhat bimodal. The nine species with an SRE above 1 Pa/m/s all had calling frequencies above 14 kHz and belonged to the paraphyletic subfamily Eneopterinae, in clades F or G. Given the difference in their song radiation mechanics (Robillard et al., 2007) we excluded these individuals from subsequent analyses (see methods). After excluding these high frequency callers, we found that SRE ranges from 0.02 to 0.67 Pa/m/s, mean: 0.18 ± 0.01 SE, n = 101.

With baseline efficiencies measured above, we next calculated how much each species could gain by using a baffle, while singing at the same frequency. To do this, we used the optimal efficiency ridge, since optimal radiator efficiency is the same as optimal baffled efficiency (Hambric and Fahnline, 2007). If animals were to continue using the same call frequency, but used an ideal baffle, each species stood to gain between $1.7 - 35$ times ($5 - 30$ dB) above their baseline SRE (mean: 7.6 ± 0.41 times, 16 ± 0.37 dB, n = 99). Among those who stood to gain the most included animals in clade B, specifically in the subfamily Mogoplistinae (scaly crickets). These animals have very small wings (mostly under 5 mm^2), suggesting a poor match between radiator and wing size. On the other hand, the animal closest to the optimal efficiency ridge, was *Madasumma affinis*, belonging to the subfamily Podoscirtinae in clade F. This animal has the largest wing at 258 mm^2, however, even this animal stood to gain 0.5 Pa/m/s (5 dB or 1.6x increase) with the use of an ideal baffle. Taken together, these data suggest that all crickets could increase SRE, and therefore, stand to benefit from use of a baffle.

Grounded calling emerges as an alternative strategy to baffle use in complex environments

While analysis of SRE suggests that all crickets should use baffles, this prediction is based on sound fields travelling in free space and over short distances. It is possible that efficiency advantages from baffle use become negligible as sounds interact with objects in the cricket's local environment (such as the ground), or as the call propagates through space. Either of these scenarios would lend support to the lack of utility hypothesis.

To address whether and how the efficiency landscape is changed by realistic calling conditions, we used boundary element modeling. Specifically, we use this model to add a "ground" component to our existing models, where the ground could have different characteristics including vegetation cover. In these models, sound can be reflected and dissipated by the ground. We model the effect of the vegetation by an excess attenuation term (see methods). We used empirical measurements of different types of ground so although our modeled ground is flat and smooth, these measures take realistic ground variability into account. We also varied the height of the caller above the ground (grounded calling: 0 m, elevated calling: 1 m). We measured efficiency again by normalizing sound levels against radiator vibration levels (see methods). Sound levels were measured at two distances from the caller: near (averaged from 0.05 – 0.2 m away), and far (averaged from 0.8 – 0.9 m away). To simplify analysis, we always measured efficiency at the same height as the caller. To differentiate this measure of efficiency from SRE, we called it sound propagation efficiency or SPE (Pa/m/s).

The most striking result was that calling from the ground (Figure 3C, D) yielded much higher SPEs than calling from above it (Figure 3A, B). This is reflected in the efficiency landscapes by a general increase in efficiency values near the ground across the ranges of frequency and wing area that we measured (see scale bars). This is likely due to the ground effect, in which much of the sound energy that would propagate below the radiators is instead reflected upward from the ground, and significant constructive interference is responsible for increased SPE (Rossing, 2014). Indeed, the highest SPE observed with a grounded caller was 4 Pa/m/s (Figure 3C), two orders of magnitude higher than peak SPE with an elevated caller (0.06 Pa/m/s, Figure 3A). On the other hand, calling from far above the ground yields SPE values that are similar in level to SRE values calculated in the ideal free-field scenario modeled previously. At farther distances, the values decrease, as would be predicted by spreading in open air.

The SRE landscape for grounded calling was not smooth and became increasingly rougher as the distance from the caller increased. Whereas elevated calling yields landscapes with broad, smooth peaks and valleys, the landscapes of callers near the ground show high levels of variation in efficiency across small changes in frequency and radiator size. In fact, the irregularity of the landscapes for grounded calling did not allow us to calculate clean optimal efficiency ridges. However, given that grounded calling is generally much higher efficiency than elevated calling, grounded callers may not need to use a baffle in the first place. Taken together, our models incorporating the ground posit that grounded calling and elevated baffled calling are two potential alternative strategies to maximize efficiency.

Alternative strategies persist when ground properties and vegetation are varied

All grounds are not equivalent. For instance, soft grounds or those covered with vegetation would be much more dissipative and may eliminate the advantage accrued from ground calling. To test this possibility, we investigated whether this alternative strategy framework holds up when these properties of the environment are varied. We found few differences in SPE with different types of grounds.

(Supplement to figure 3-2). SPE tends to be slightly higher with a "soft" ground, which is better at
dissipating sound, similar to a freshly tilled agricultural field (see methods) and this effect is magnified
further away from the caller. With a harder, more reflective ground, similar to a tightly packed forest
floor, SPE is slightly lower. However, the differences between these two ground types occur at wing sizes
well outside the natural range for crickets. At close distances, and particularly above the ground, the
differences between ground types are very small. Therefore, all future analyses assume a "hard" ground.

Finally, we tested whether vegetation would reduce the predicted SPE landscapes for grounded calling.
Vegetation does slightly decrease the magnitude of SPE overall. However, we found that excess
attenuation due to vegetation does not significantly change the overall patterns of efficiency, by and large
shifting the efficiency landscape to a lower point at most points within the parameter space (Supplement
to figure 3-3 C, D) (Bashir et al., 2015). Therefore, it turns out baffled calling would be less efficient with
vegetation than without. This shift in efficiency is also not perfectly equal at all frequencies, and SPE is
lowered slightly more at high frequencies (Supplement to Figure 3-3). This suggests that high frequency
callers may be at an increased disadvantage when calling in vegetation as suggested before (Romer and
Lewald, 1992), and will also see diminishing returns when using a baffle. This effect is relatively small,
however (< 6 dB). Additionally, the effects of vegetation on SPE are undoubtedly more complicated than
an excess attenuation factor, so modeling plants explicitly, at a variety of shapes and sizes, would be a
useful extension to this study. However, overall, we conclude that calling from the ground remains an
effective alternative strategy, even if the ground is soft, or covered with some vegetation.

Alternative strategies are still viable when considering call directionality
So far, our analysis has used the loudness of calls to define efficiency. However, a call must be both loud
and directional to be effective. That is, the call must present a spatial gradient that a potential mate can
follow to the source. Previous data has suggested that such gradients are severely degraded in ground
calling crickets (Kostarakos and Römer, 2010; Mhatre and Balakrishnan, 2006; Römer, 2015; Romer and
Lewald, 1992), but not in elevated calling (Deb and Balakrishnan, 2014). This suggests that SPE gains
from grounded calling may trade off against call directionality. Since our models generate spatially
explicit predictions of sound fields, we can address this possibility. We analyzed call directionality by
measuring how difficult it would be for a female cricket to follow an acoustic gradient back to the call's
source. A value of one indicates that the gradient along a chosen transect perpendicular to the wing planes
is always in the "correct" direction, that is, sound pressure level increases as the female moves toward the
call in steps of ~ 2 body lengths (2 cm). A lower value means that over some stretches over this transect,
SPL increases and at other steps it decreases. A value of 0.5, for instance, means that the SPL decreases
over 50% of the steps as the female moves closer (see methods).

Directionality varies with respect to frequency, radiator size, and height from ground (Supplement to
Figure 3-4). Although grounded calling does experience a loss of directionality compared to elevated
calling, these losses are mostly minimal. Near a grounded caller, there is a strong cutoff at about 5 kHz,
below which all are strongly directional (> 0.9) except for very small wings. This cutoff decreases to
about 3.5 kHz far from the caller. Therefore, high frequency callers would be most susceptible to the
gradient effects. However, even below these cutoffs, directionality rarely drops below 0.5 in any
condition, and grounded calling remains a viable alternative. It should be noted that other studies have
found more substantial degradations in call directionality in sounds traveling along the ground, but over
greater distances than our current models (Kostarakos and Römer, 2010). However, data for both field

crickets and tree crickets suggest that the SPL of typical cricket calls drop below threshold at about 1 m

from the caller (Deb and Balakrishnan, 2014; Mhatre and Balakrishnan, 2006), and therefore we consider

this a biologically relevant distance over which to measure directionality.

Alternative calling strategies are likely in use by some cricket species

Based on the overall propagation efficiency landscape, grounded calling and baffled calling are potential

alternative strategies to maximize efficiency. However, these landscapes cover the full acoustic-

morphospace, i.e. all possible combinations of radiator (wing) size and call frequency, but most of these

combinations are not used by real crickets.

To shift our focus from the acoustics, and to make it more biologically relevant, we performed an analysis

to determine whether alternate calling strategies are in use by actual crickets. We divided all sampled

animals into their respective clades and calculated clade level SPE for each of three alternative strategies

as measured far from the caller: calling from the open hard ground (grounded calling), from within

vegetation 1 m off the ground with no baffle (unaided calling), and 1 m off the ground with a baffle

(baffled calling). We compared both baffled calling and grounded calling to unaided calling as a baseline.

It would then be ideal to determine whether the animals in fact use the strategy that we predict should

maximize efficiency based on known calling preferences. Unfortunately, we do not have data on calling

preferences of many sampled animals, however, some clade-level similarities have been observed in

calling behaviors, and different clades show some clustering in the wing size-frequency space (figures 2

and 3). We give three examples below of clades with some known information about calling behavior.

We turn first to clade F, consisting primarily of the Oecanthines, or tree crickets. Members of this group

stand to gain on average about 12 dB from grounded calling compared to unaided calling according to our

data (figure 4). However, they could gain ~19 dB it they baffled. Indeed, Oecanthine natural history bears

out our predictions; tree crickets are known to mostly call from vegetation, including vegetation that is

suitable for baffle building and use (Deb and Balakrishnan, 2014; Forrest, 1982). In fact, all known

cricket baffle users are in this clade, as predicted based on the lack of utility hypothesis. For clade G, on

the other hand, consisting mostly of the Gryllinae, or field crickets, we predict the opposite. On average,

grounded calling gives an advantage of 7x or 17 dB above unaided for species in this group, whereas

baffling only gives an advantage of 10x or 20 dB above unaided. Again, behavioral data suggests that

many field crickets indeed prefer to call from the open ground habitats that we predict would maximize

their efficiency (Weissman and Gray, 2019).

Clade A, the Gryllotalpidae or the mole crickets, represents an interesting exception to this alterative

strategy framework. This group stands to gain the most from baffled calling of all clades (13x, 22 dB)

compared to grounded calling (2x, 6 dB). Yet, species in this group are all known to exclusively call from

the ground and do not use baffles. However, they do use an acoustic aide. Mole crickets build and call

from burrows which function as resonators and convert them into monopole sound sources, eliminating

acoustic short-circuiting through a different mechanism than baffling (Bennet-Clark, 1987). Indeed, it is

possible that other acoustic means of maximizing call efficiency exist and could in the future add further

complexity to our hypotheses.

Finally, if an animal baffled, but its call propagated though vegetation compared with no vegetation, the

gains would be relatively small in most cases (< 6 dB) (Supplement to figure 4-2), We also performed a

similar analysis for call directionality (Figure 5). However, since directionality was quite high for all

calling conditions, we therefore suggest that directionality does not preclude one alternative strategy over
another.

Ideas and Speculation – why would baffle use evolve in the first place?

From our data, exploiting the ground effect with grounded calling emerges as a viable alternative to baffle
use. Grounded calling even exceeds the efficiency gains of baffled calling in some scenarios. Given that
baffle creation and use requires a combination of site selection preferences, a specialized behavioral
repertoire, and precise execution of these behaviors (Mhatre, 2018; Mhatre et al., 2017), the real question
becomes why a species would ever use this strategy if a fairly simple site selection preference for the
ground could give similar increases in efficiency.

Crickets have been calling for a long time. There is evidence that crickets with stridulatory mechanisms
(and therefore the ability to produce sound) existed as early as the Cretaceous period (Senter, 2008).
These early calling crickets were likely ground dwellers, with some species subsequently moving up into
vegetation as the group diversified (Song et al., 2015). We therefore suggest that baffle-using crickets
may have originally moved up into vegetation for non-acoustic reasons, whether it was to exploit
additional food resources or avoid predators. Baffle use would have then evolved secondarily to recover
some of the efficiency lost when abandoning grounded calling. Our biophysical modeling methods open
the door to testing such a hypothesis about baffle use.

Additionally, there are almost certainly undescribed instances of baffle use in crickets, but identifying
baffled calling involves careful observation of a species' calling behavior, followed by subsequent
measurements to confirm that a structure indeed has acoustic properties consistent with a baffle. Using
our biophysical models, we can help narrow the search for baffle-using crickets by identifying the clades
which would stand to benefit the most in terms of calling efficiency given their call frequency and wing
size. Boundary element models are particularly flexible in this regard, as their computational efficiency
allows the construction of larger, more complex habitats that can test a wide range of variables within
habitats. It is known, for example, that crickets use other acoustic aides to increase calling efficiency.
Some crickets call from burrows or cracks in the ground, from tree trunks or the walls of caves or even
use structures that have some baffling capability but do not function as fully optimal baffles (Weissman
and Gray, 2019). We show here that biophysical models can both test and generate hypotheses about such
rare behaviors in animals.

Another advantage of our modeling approach is that it does not require live, or even extant, animals. In
principle, we could model the wings of extinct crickets, and estimate their calling frequency based on the
stridulatory apparatus on the wing (Gu et al., 2012; Woodrow et al., 2022). By bringing extinct crickets
"back to life" in this way we could ask questions about the evolution of acoustic aides more broadly. We
suggest that biophysical modeling, grounded with data from real animals, can be a valuable tool for any
biologist wishing to better characterize and understand the diversity of animal communication.

Materials and Methods

Specimen Data

We collected data on wing surface area and call frequency for each of 111 cricket species distributed
across the 7 clades described by Chintauan-Marquier et al. (Chintauan-Marquier et al., 2016) (Figure 1).

We restrict our analysis to this group, since these species are known to raise their wings when singing
(Desutter-Grandcolas, 2003). This behavior means that they are dipole sources of sound, and acutely
affected by acoustic short-circuiting (Bennet-Clark, 1998; Forrest, 1991). Each species was assigned to a
clade based on the following two criteria: (1) the species itself was included in the Chintauan-Marquier et
al (2016) dataset or (2) the subfamily of the species was included in the dataset and it was found to be
monophyletic within one of the seven Chintauan-Marquier clades. Data were obtained from a variety of
databases including Orthoptera Species File (Cigliano, M. M. et al.), Crickets North of Mexico, and
numerous publications (all references are available in tables 1-2). For a few species of Oecanthines, wings
were provided by Nancy Collins and photographed in the lab under a dissecting microscope. All
specimens measured were adult males, identified by wing morphology and lack of ovipositor. We
measured surface area of the entire left forewing including the lateral field. Fitting an ellipse to the wing,
we calculated aspect ratio (length of ellipse/width of ellipse). All image data were gathered using ImageJ
version 1.53 (Schindelin et al., 2015). We next calculated the fundamental frequency of cricket
advertisement calls using Raven Lite version 2.0 (Cornell Lab of Ornithology, 2020). When multiple
specimens of a single species were analyzed, averages were used for wing size and call variables. To
better represent the full range of wing size and call frequency in our dataset, we included some specimens
in the histograms showing wing size and frequency (Figure 2) for which we had only one type of data.
Twelve animals had only wing size, but not call data, and 57 animals had call, but not wing size data (see
tables 1-2 for details).

Finite Element Models for Sound Radiation Efficiency
We first estimated the sound radiation efficiency of crickets calling in open air using finite element (FE)
analysis (Supplement to Figure 2-2). We built this model and all subsequent models in COMSOL
Multiphysics version 5.5. All models used the pressure acoustics module and were solved in the
frequency domain assuming a steady state. The Helmholtz equation was the governing equation.

Model geometry, boundary conditions, symmetry, and vibration
Animals were represented by two ellipses which modelled the forewings sitting next to each other along
the long axis, in the same plane (Supplement to Figure 2-2). These ellipses should accurately approximate
total radiated sound power and the true shape of the sound field, and it is the size and shape of the radiator
and frequency of the call that determines radiation efficiency, and not the radiator's material properties
(Hambric and Fahnline, 2007). Surrounding the wings was a spherical acoustic domain consisting of air
with a 20 cm radius. The acoustic domain was suspended inside a second spherical domain of 40 cm
radius (Supplement to figure 2-2. A perfectly-matched layer (PML) was applied to the area between the
acoustic domain and larger outer sphere. PMLs in finite element modelling are used to mimic an open and
non-reflecting infinite acoustic domain, which absorbs all sound energy from the acoustic domain
(Berenger, 1994). Thus PMLs mitigate modelling artefacts such as the effects of sound reflecting from the
edges of the acoustic domain (Mhatre et al., 2017). To reduce computational time, our model was
constructed using ¼ symmetry. That is, we modeled ½ of one wing and ¼ of the two spherical domains.
We then mirrored this model twice, once about the short axis of the wing to create a whole wing and ½ of
the spherical domains, and a second time about a line between and parallel to the long axis of the wings to
create a second wing and the whole spherical domains. We applied a time- and space-averaged velocity
normal to the entire surface of the wings at 0.13m/s. This was the value measured from the wings of
singing *Oecanthus henryi* and is the only known estimate for crickets (Mhatre et al., 2017). However,
given that we are calculating efficiency rather than reporting actual sound pressure levels, normalizing

using this known value allows comparison between species. We vibrated the wings at a frequency range
of 0.5 - 32 kHz, in increments of 0.25 kHz. SRE was subsequently calculated from each of these model
outputs as a volumetric average of the absolute pressure in the acoustic domain, divided by the time- and
space-averaged velocity of 0.13 m/s, resulting in units of Pa/m/s.

Finite Elements
3D tetrahedral elements were used in both the acoustic domain and PML. After undertaking a mesh size
sensitivity study (Supplement to figure 2-3), we chose the "extra fine" mesh setting in COMSOL, with
about 60000 elements in the acoustic domain. This number did vary somewhat with wing size, as fewer
elements are used with very small wings.

Model Parameters
We ran the finite element model at a range of wing surface areas from 0.4 – 4000 mm^2, scaled
logarithmically by the equation 4×10^x, where x ranges from -1 to 3). We used an aspect ratio of 2
(wings are twice as long as they were wide). Our chosen aspect ratio of 2 was well within the range of
most cricket species (median: 1.6, range: 0.7 - 3.7). Aspect ratio did not play a large role in sound
production, except at aspect ratios > 5 (length of wing 5x the width), which were not observed in real
wings (Supplement to Figure 2-4). For aspect ratios within the range of crickets, differences in SRE
between aspect ratios at a given wing area and frequency never exceeded 3 dB.

Other Modeling Considerations
The cricket body was not included in our models as it was found to have negligible effects on SRE at all
wing sizes and frequencies (mean difference: 0.05 ± 0.01 dB). We also evaluated whether applying
vibration to only a part of the wing (a "harp") influenced sound production. Some cricket species (though
not all) use this sound production method (Godthi et al., 2022). We found only minor increases in SRE
between vibrating only a harp or vibrating the entire wing (mean: 4 ± 0.08 dB), except at wing sizes well
outside the range of the real wings that we measured (Supplement to figure 2-4).

Boundary Element Models for sound propagation efficiency
To test hypotheses about how cricket calls interact with objects in the environment, we needed to include
an additional domain in the model: a "ground" with realistic parameterized acoustic impedance. To make
this model as realistic as possible and to minimize boundary effects, we needed to make the ground
element as large as possible relative to the size of the wings. The combination of the large size of ground
and the high sound frequencies of interest resulted in finite element models that were too computationally
intensive to run. We therefore turned to boundary element (BE) modeling as an alternative means of
assessing sound propagation efficiency.

Both acoustic boundary element models and finite element models are numerical methods for solving the
Helmholtz equation to capture a developing sound field within a medium. However, they differ in how
they discretize space within the model. Finite element models discretize volumes by partitioning into a 3-
dimensional mesh of finite elements. This allows detailed descriptions of the medium in which the field
develops. Boundary element models on the other hand reduce computational cost by discretizing only the
boundaries of the acoustic domain and assume a linear homogenous medium in all other spaces. The
boundary element formulation therefore trades off some specificity in exchange for computational
efficiency, allowing us to make relatively large, more biologically relevant models to assess sound

propagation in a spatially explicit manner. We ran our boundary element models using the pressure
acoustics, boundary elements module in COMSOL. All models were run in the frequency domain and
assumed steady-state behavior. The Helmholtz equation does not take attenuation due to damping into
account, which can become an issue at distances far from the source. However, at frequencies >500 Hz,
attenuation due to damping is only about 2 dB per kilometer (Embleton, 1996), so we considered this
effect to be negligible over the distances of interest for this study (0.2 – 1m).

Model geometry, forcing, and boundary conditions
Wings in the boundary element model were modeled the same way as in the finite element model
(Supplement to Figure 3-1), with no material properties and one-way coupling between the wings and
sound fields. Wings were positioned perpendicular to the top surface of the ground, with the flat surfaces
of the wings aligned with the short axis of the ground. The wings were centered with respect to ground.
The wings were placed above the ground at either 0 m, or 1 m. The same time- and space-averaged
velocity was applied as above, and the same set of wing surface areas were used. The ground was
modeled as a rectangular slab, 0.5 m wide, 2 m long, and 0.10 m thick. A sound-hard boundary was
applied to the bottom surface of the ground slab. Because we were interested in spatially-explicit
measures of efficiency as sound propagates across ground, we did not use symmetry conditions to create
this model. However, because the sound fields should be symmetric on either side of the wings, we only
measured the sound field on one side.

Model Parameters
We used a restricted frequency range for the boundary element models, ranging from of 0.5 – 10 kHz, in
increments of 0.25 kHz. We chose 10 kHz as the cutoff because very few of our measured animals call
above this frequency, and those that do were Eneopterine species who were likely using a vibrational
mode inconsistent with the piston mode that we have implemented (Robillard et al., 2007). High
frequency (> 10 kHz) callers were included in the finite element models to give a general sense of where
they might fit in with the other animals, but in reality no animals occupy this space in the landscape and
all analyses explicitly comparing species exclude them.

To model how a real ground interacts with sound, we applied an acoustic impedance to our modeled
ground. Acoustic impedance quantitatively describes how much sound energy is dissipated by the ground,
compared to the energy reflected. We used the Attenborough slit-pore model to implement ground
impedance. This model uses three parameters to capture both dissipative and reflective properties: flow
resistivity, pore density, and porous layer depth. We modeled two different types of ground, a "soft"
ground (flow resistivity: 2000 kPa×s/m^2, porosity: 0.6) which is less reflective and a "hard" ground (flow
resistivity: 9 kPa×s/m^2, porosity: 0.4), which is more dissipative. Porous layer depth was held constant for
both treatments, at 0.04 m. Both ground parameters were taken from empirical measurements of a "soft"
freshly-tilled field and a "hard" forest floor (Attenborough et al., 2011).

Sound propagation efficiency definition
In the finite element models, we calculated a volumetric average of absolute pressure within the acoustic
domain. However, such a measure would not be appropriate to assess propagation efficiency, as the sound
waves' interactions with the ground would accumulate as distance from the source increases. Therefore,
we calculated SPE in a spatially-explicit manner. We measured absolute pressure at 50 points along a 1m
long line parallel to the long axis of the ground, at the same height as the wings. The line originated at the

centre between the two elliptical 'wings'. We divided this line into "near" and "far" from the caller: near
= 0.05 – 0.2 m from wings, far = 0.8 – 0.9 m from wings. Efficiency was calculated as before, by dividing
sound pressure level (Pa) by 0.13 m/s, the space and time averaged velocity applied to the wings. We also
created an additional boundary element model with no ground, to allow for direct comparisons between
ground and no ground and to sanity check the boundary element method compared with the previous
finite element models.

Finite Elements
Tetrahedral elements were used on the surface of the ground and 2D triangular elements on the wings.
After performing a similar sensitivity study as with the finite element models, we decided on a maximum
element size for the wing surfaces of 0.5 cm and 1 cm for the surface of the ground. Since the sound wave
is not explicitly modelled, this element size is not related to sound frequency, and instead captures
boundary conditions and hence can be larger than in the finite element models.

Excess attenuation due to vegetation
To calculate the effect of vegetation, we used existing models to calculate how standing vegetation is
expected to impact call efficiency. We then subtracted this excess attenuation from the COMSOL result.
We calculated excess attenuation using the following empirically derived equation from Bashir et al 2015:

$$\frac{EA(Pa)}{FL} = 3[1 - \exp(0.3 - 0.5(ka))], ka > 0.6$$

Where EA(Pa) represents excess attenuation due to foliage, F = foliage area per unit volume, L = path
length, k = wavenumber, and a = average leaf size. We used values empirically derived for dense foliage
with relatively large leaves (Bashir et al., 2015), such as those used to construct baffles in known baffle-
using species. To approximate the foliage area and leaf size that a typical baffle user would prefer, we
used F = 6.3m^{-1} and a = 0.0784m in our measures of excess attenuation.

Efficiency of individual species and how much they gain with baffle
To test the lack of utility hypothesis, we used the output of our models to estimate the gains in efficiency
that each species could attain if it used a baffle. We did this for both the idealized measure of efficiency
(SRE) and the more realistic scenario involving a ground and vegetation (SPE). For each modeling
scenario, we first estimated the efficiency of each cricket species in our dataset, given their wing area and
call frequency. Next, we calculated the efficiency that each species would have if it used an ideal baffle.
To do this, we first determined the size at which the wings and baffle working together as a single radiator
would reach maximal efficiency. The optimal size is a function of the wavelength of that sound frequency
in that medium. We calculated the quantity ka for each surface area-frequency combination in the model,
where k is the wavenumber and a is the effective radius of the sound radiating plates (Hambric and
Fahnline, 2007). ka is a dimensionless quantity often used in acoustics, as it helps define when a radiator
of a particular size transitions from being inefficient sound radiator at low frequencies to an efficient high
frequency radiator. For instance, an optimally sized circular piston has ka = 1 (Hambric and Fahnline,
2007).

However, the radiators being considered here are two aligned ellipses which not perfectly circular in
shape. Hence the particular value of maximal efficiency ka will be different in this configuration. So, to
estimate optimal ka for cricket wings, we plotted ka versus efficiency as measured from our models, with

a separate trace for each frequency (Supplement to Figure 4-1). We then identified the ka at which
maximal efficiency was reached for all frequencies. This value represents the size at which highest
attainable efficiency is reached, consistent with an optimally baffled condition (Hambric and Fahnline,
2007) (Supplement to Figure 4-1). In our finite element models, we found optimal ka to be about 1.3. For
the boundary element models at the far distance, optimal ka was approximately 1.55 (Figure 4-1).

Next, we performed a simple linear regression between frequency and maximal efficiency at optimal ka,
then calculated the slope and y-intercept of this regression (Supplement to Figure 4-1). We used this
equation to calculate optimal baffled efficiency for each species. The relationship between frequency and
efficiency differed depending on condition (open ground vs ground + vegetation) (Supplement to figure
4-1), so this regression was performed separately for each environmental condition when calculating
optimal baffled efficiency for a given condition.

Directionality index

To address how difficult it would be for a female to localize a male call, we assessed the directionality of
the call in each modeling scenario. In an open field with no ground, the sound level is expected to
decrease smoothly following the inverse square law (Bradbury and Vehrencamp, 1998; Rossing, 2014). A
cricket moving toward the source of a call should therefore always experience either an increase in
loudness, or, if the increase is below the animal's difference threshold, no change in loudness. A cricket
should always move in the direction of increasing SPL to locate the singing male and therefore a mating
opportunity. However, in reality, sound fields become more complicated when they interact with the
ground, resulting in a noisy relationship between SPL and distance (Römer, 2021). In such sound fields,
female phonotaxis may fail as there is no clear acoustic gradient to follow to the source. To quantify this
degree of potential "confusion", we calculated a directionality index for each modeling scenario. First, we
calculated Δ SPL between each two adjacent points 2 cm apart (~ 2 body lengths for most animals in this
analysis). Δ SPL was calculated starting at 1 m away and moving toward the source. Next, we classified
each of these values as either consistent with expected change in SPL or inconsistent. Consistent values
represented either an increase, no change, or a decrease smaller than Δ 3dB (a factor of about 1.4), which
is thought to be close to the detectable threshold for crickets (Mhatre and Balakrishnan, 2007). See
Römer, 2021 for a more complex treatment of such thresholds. For our purposes, inconsistent values
represented a decrease in SPL greater than 3 dB. For each modeling scenario, we calculated the
proportion of Δ SPLs classified as consistent. This resulting value we call "Directionality" ranging from 0
to 1 (Supplement to figure 3-4). We calculated directionality for two different distance treatments, "near"
was calculated from 0.05 – 0.2 m from the wings, and "far" was calculated from 0.5 – 1m from the wings.

Acknowledgements

We wish to acknowledge a number of undergraduate students who assisted with the collection and
databasing of cricket acoustic and morphometric data: Morteza Al Rabya, Nancy Kim, Shanker
(Matthew) Nadarajah, and Daniel Xie. Nancy Collins provided specimens for several of the wing
measurements. Graduate students Hossein Asgari and Reese Gartly assisted with multiplexing dozens of
model runs on multiple machines. Emine Celiker and Damian Elias provided feedback on an earlier
version of this manuscript. This work would have not been possible without an extensive worldwide
network of specimen databases. We especially wish to thank those who contribute to, curate, or maintain

Orthoptera Species File, Singing Insects of North America, Museum D'Historie Naturelle, and numerous
university insect collections. We also wish to thank the photographers who graciously provided their
images free of charge for inclusion in this manuscript.

We would also like to thank the following funding sources for their support of this research: NSERC
Discovery (Grant no. 687216), and early career supplement (675248), and an NSERC Canada research
chair (Grant no. 693206) to NM; the Western University Postdoctoral Fellowship to EEB; Western
Undergraduate Work Study to SD and HW; Western Undergrad Summer Research Internship to HW.

References

Attenborough K, Bashir I, Taherzadeh S. 2011. Outdoor ground impedance models. *The Journal of the Acoustical Society of America* **129**:2806–2819. doi:10.1121/1.3569740

Bashir I, Taherzadeh S, Shin H-C, Attenborough K. 2015. Sound propagation over soft ground without and with crops and potential for surface transport noise attenuation. *J Acoust Soc Am* **137**:12.

Bennet-Clark HC. 1998. Size and scale effects as constraints in insect sound communication. *Phil Trans R Soc Lond B* **353**:407–419. doi:10.1098/rstb.1998.0219

Bennet-Clark HC. 1987. The Tuned Singing Burrow of Mole Crickets. *Journal of Experimental Biology* **128**:383–409.

Berenger J-P. 1994. A perfectly matched layer for the absorption of electromagnetic waves. *Journal of Computational Physics* **114**:185–200. doi:10.1006/jcph.1994.1159

Boesch C, Boesch H. 1983. Optimisation of Nut-Cracking With Natural Hammers By Wild Chimpanzees. *Behaviour* **83**:265–286. doi:10.1163/156853983X00192

Bradbury JW, Vehrencamp SL. 1998. Principles of Animal Communication, First edition. ed. Sunderland: Sinauer Associates, Inc.

Chintauan-Marquier IC, Legendre F, Hugel S, Robillard T, Grandcolas P, Nel A, Zuccon D, Desutter-Grandcolas L. 2016. Laying the foundations of evolutionary and systematic studies in crickets (Insecta, Orthoptera): a multilocus phylogenetic analysis. *Cladistics* **32**:54–81. doi:10.1111/cla.12114

Cigliano, M. M., Braun, D. C., Eades, D. C., Otte, D. n.d. Orthoptera Species File. *Orthoptera Species File Version 50/50*. http://orthoptera.speciesfile.org/HomePage/Orthoptera/HomePage.aspx

Collins N. 2012. A Citizen Scientist's Contributions to Tree Cricket Taxonomy. *American Entomologist* **58**:9–10. doi:10.1093/ae/58.1.0009

Deb R, Balakrishnan R. 2014. The opportunity for sampling: the ecological context of female mate choice. *Behavioral Ecology* **25**:967–974. doi:10.1093/beheco/aru072

Deb R, Modak S, Balakrishnan R. 2020. Baffling: a condition-dependent alternative mate attraction strategy using self-made tools in tree crickets. *Proceedings of the Royal Society B: Biological Sciences* **287**:20202229. doi:10.1098/rspb.2020.2229

Desutter-Grandcolas L. 2003. Phylogeny and the evolution of acoustic communication in extant Ensifera (Insecta, Orthoptera). *Zoologica Scripta* **32**:525–561. doi:10.1046/j.1463-6409.2003.00142.x

Embleton TFW. 1996. Tutorial on sound propagation outdoors. *The Journal of the Acoustical Society of America* **100**:31–48. doi:10.1121/1.415879

Erregger B, Schmidt AKD. 2018. Anthropogenic calling sites boost the sound amplitude of advertisement calls produced by a tropical cricket. *Animal Behaviour* **142**:31–38. doi:10.1016/j.anbehav.2018.05.021

Farris HE, Forrest TG, Hoy RR. 1997. The effects of calling song spacing and intensity on the attraction of flying crickets (orthoptera: Goryllidae: Nemobiinae). *J Insect Behav* **10**:639–653. doi:10.1007/BF02765384

Forrest TG. 1991. Power output and efficiency of sound production by crickets. *Behav Ecol* **2**:327–338. doi:10.1093/beheco/2.4.327

Forrest TG. 1982. Acoustic Communication and Baffling Behaviors of Crickets. *The Florida*
Entomologist **65**:33–44. doi:10.2307/3494144

Fujii JA, Ralls K, Tinker MT. 2015. Ecological drivers of variation in tool-use frequency across sea otter
populations. *Behavioral Ecology* **26**:519–526. doi:10.1093/beheco/aru220

Garud M, Godthi V, Reddy J, Dangi A, Pratap R. 2017. Cricket Inspired High Efficiency MEMS
Speakers. *Proceedings* **1**:345. doi:10.3390/proceedings1040345

Godthi V, Balakrishnan R, Pratap R. 2022. The mechanics of acoustic signal evolution in field crickets.
Journal of Experimental Biology **225**:jeb243374. doi:10.1242/jeb.243374

Gu J-J, Montealegre-Z F, Robert D, Engel MS, Qiao G-X, Ren D. 2012. Wing stridulation in a Jurassic
katydid (Insecta, Orthoptera) produced low-pitched musical calls to attract females. *Proceedings*
of the National Academy of Sciences **109**:3868–3873. doi:10.1073/pnas.1118372109

Hambric SA, Fahnline JB. 2007. Structural Acoustics Tutorial—Part 2: Sound—Structure Interaction.
Acou Today **3**:9. doi:10.1121/1.2961152

Hansell M, Ruxton G. 2008. Setting tool use within the context of animal construction behaviour. *Trends*
in Ecology & Evolution **23**:73–78. doi:10.1016/j.tree.2007.10.006

Hunt GR, Gray RD, Taylor AH. 2013. Why is tool use rare in animals? In: Boesch C, Sanz CM, Call J,
editors. Tool Use in Animals: Cognition and Ecology. Cambridge: Cambridge University Press.
pp. 89–118. doi:10.1017/CBO9780511894800.007

K. Lisa Yang Center for Conservation Bioacoustics at the Cornell Lab of Ornithology. 2020. Raven Lite:
Interactive Sound Analysis Software (Version 2.0).

Kostarakos K, Römer H. 2010. Sound transmission and directional hearing in field crickets:
neurophysiological studies outdoors. *J Comp Physiol A* **196**:669–681. doi:10.1007/s00359-010-
0557-x

Mhatre N. 2018. Tree cricket baffles are manufactured tools. *Ethology* **124**:691–693.
doi:10.1111/eth.12773

Mhatre N, Balakrishnan R. 2007. Phonotactic walking paths of field crickets in closed-loop conditions
and their simulation using a stochastic model. *Journal of Experimental Biology* **210**:3661–3676.
doi:10.1242/jeb.003764

Mhatre N, Balakrishnan R. 2006. Male spacing behaviour and acoustic interactions in a field cricket:
implications for female mate choice. *Animal Behaviour* **72**:1045–1058.
doi:10.1016/j.anbehav.2006.02.022

Mhatre N, Malkin R, Deb R, Balakrishnan R, Robert D. 2017. Tree crickets optimize the acoustics of
baffles to exaggerate their mate-attraction signal. *eLife* **6**:e32763. doi:10.7554/eLife.32763

Muñoz MI, Halfwerk W. 2022. Amplification of frog calls by reflective leaf substrates: implications for
terrestrial and arboreal species. *Bioacoustics* **31**:490–503. doi:10.1080/09524622.2021.1978319

Pierce JD. 1986. A Review of Tool Use in Insects. *The Florida Entomologist* **69**:95–104.
doi:10.2307/3494748

Prozesky-Schulze L, Prozesky OPM, Anderson F, Van Der Merwe GJJ. 1975. Use of a self-made sound
baffle by a tree cricket. *Nature* **255**:142–143. doi:10.1038/255142a0

Regen J. 1913. Über die Anlockung des Weibchens von Gryllus campestris L. durch telephonisch
übertragene Stridulationslaute des Männchens. *Pflügers Archiv European Journal of Physiology*
155:193–200.

Robillard T, Grandcolas P, Desutter-Grandcolas L. 2007. A shift toward harmonics for high-frequency
calling shown with phylogenetic study of frequency spectra in Eneopterinae crickets (Orthoptera,
Grylloidea, Eneopteridae). *Can J Zool* **85**:1264–1275. doi:10.1139/Z07-106

Römer H. 2021. Neurophysiology goes wild: from exploring sensory coding in sound proof rooms to
natural environments. *J Comp Physiol A* **207**:303–319. doi:10.1007/s00359-021-01482-6

Römer H. 2015. Directional hearing: from biophysical binaural cues to directional hearing outdoors. *J*
Comp Physiol A **201**:87–97. doi:10.1007/s00359-014-0939-6

Römer H, Butlin RK, Guilford T, Krebs JR. 1993. Environmental and biological constraints for the
evolution of long-range signalling and hearing in acoustic insects. *Philosophical Transactions of*

618 *the Royal Society of London Series B: Biological Sciences* **340**:179–185.
619  doi:10.1098/rstb.1993.0056
620 Romer H, Lewald J. 1992. High-frequency sound transmission in natural habitats: implications for the
621  evolution of insect acoustic communication. *Behav Ecol Sociobiol* **29**:437–444.
622  doi:10.1007/BF00170174
623 Rossing TD, editor. 2014. Springer Handbook of Acoustics. New York, NY: Springer New York.
624  doi:10.1007/978-1-4939-0755-7
625 Schindelin J, Rueden CT, Hiner MC, Eliceiri KW. 2015. The ImageJ ecosystem: An open platform for
626  biomedical image analysis. *Molecular Reproduction and Development* **82**:518–529.
627  doi:10.1002/mrd.22489
628 Senter P. 2008. Voices of the past: a review of Paleozoic and Mesozoic animal sounds: review. *Historical
629  Biology* **20**:255–287. doi:10.1080/08912960903033327
630 Seybert AF, Wu TW, Wu XF. 1994. Experimental validation of finite element and boundary element
631  methods for predicting structural vibration and radiated noise (No. NASA-CR-4561).
632 Song H, Amédégnato C, Cigliano MM, Desutter-Grandcolas L, Heads SW, Huang Y, Otte D, Whiting
633  MF. 2015. 300 million years of diversification: elucidating the patterns of orthopteran evolution
634  based on comprehensive taxon and gene sampling. *Cladistics* **31**:621–651. doi:10.1111/cla.12116
635 Spagnoletti N, Visalberghi E, Ottoni E, Izar P, Fragaszy D. 2011. Stone tool use by adult wild bearded
636  capuchin monkeys (Cebus libidinosus). Frequency, efficiency and tool selectivity. *Journal of
637  Human Evolution* **61**:97–107. doi:10.1016/j.jhevol.2011.02.010
638 Weissman DB, Gray DA. 2019. Crickets of the genus Gryllus in the United States (Orthoptera: Gryllidae:
639  Gryllinae). *Zootaxa* **4705**:1–277. doi:10.11646/zootaxa.4705.1.1
640 Woodrow C, Baker E, Jonsson T, Montealegre-Z F. 2022. Reviving the sound of a 150-year-old insect:
641  The bioacoustics of Prophalangopsis obscura (Ensifera: Hagloidea). *PLOS ONE* **17**:e0270498.
642  doi:10.1371/journal.pone.0270498
643

Figures

Figure 1. Crickets were sampled evenly across the cricket phylogeny covering most taxonomic groups known to produce calls. **A.** Representative images of cricket males with wings raised in calling posture. From upper left to lower right, species pictured are: *Hoplosphyrum boreale* (photo: James P. Bailey), *Phyllopalpus pulchellus* (photo: Wilbur Hershberger)*, Lerneca inalata* (photo: Richard C. Hoyer)*, Meloimorpha japonica* (photo: Ryosuke Kuwahara)*, Oecanthus quadripunctatus* (photo: James P. Bailey), and *Turanogryllus eous* (photo: Taewoo Kim). Boxes with letter indicate the clade to which each species belongs. **B.** Phylogeny illustrating the seven clades defined by Chintauan-Marquier et al (2016) along with subfamilies within each clade (branch lengths not to scale). Only subfamilies from which we sampled for this study are included. Subfamilies with asterisks are polyphyletic between multiple clades. **C.** Specimen sampling scheme. Top panel shows the distribution of all species described across each of the seven clades in Grylloidea (according to Orthoptera Species File, as of February 16, 2022). Bottom panel shows distribution of species that were sampled for this study.

Figure Supplement 1-1. Measuring acoustic-morphospace
of crickets. Wing area and call frequency were quantified.
A. Wing surface area was calculated as the area of the entire
forewing. **B.** The carrier frequency (sometimes called
fundamental frequency) of the call was identified from
spectrograms and recorded. If the fundamental frequency
occupied a sweep over a range of frequencies (as shown), an
average was taken.

Figure 2. All crickets could increase efficiency by baffled calling. Sound radiation efficiency (SRE) landscape across the acoustic-morphospace of crickets. **A.** SRE achieved by different combinations of wings sizes vibrating at different frequencies. Within this space of possibilities the wing areas and call frequencies of all sampled animals are shown overlaid as points on the SRE landscape. This SRE is calculated from finite element models. Red line indicates optimal efficiency ridge, or the size at each frequency that would produce an ideally baffled calling scenario. **B, C.** Distributions of the calling song frequency and wing size of different animals, respectively. Histograms include additional species for which only wing or call measurements were available. **D.** SRE of each species without use of a baffle. **E.** SRE of each species with use of an ideal baffle.

Figure Supplement 2-1. Sound fields produced by modeled wings vary with respect to call frequency and wing size. Fields are oriented such that wings are perpendicular to page and vibrate left to right, as indicated by the silhouette cricket. Sound fields are given for the following combinations of wing size and frequency: **A.** wing size = 0.04 mm^2, frequency = 10 kHz; **B.** wing size = 225 mm^2, frequency = 10 kHz; **C.** wing size = 0.4 mm^2, frequency = 1.5 kHz; **D.** wing size = 225 mm^2, frequency = 1.5 kHz. Spatial scale given in C applies to all sound fields. Efficiency (here, size and color of sound field) increases with improved match between wavelength of sound and size of radiator (wing). Cricket wings in general are small so this match is poor except at the extreme high end of radiator size and call frequency (B).

Figure Supplement 2-2. Geometry of finite element model. Inner blue sphere is the acoustic domain. The wings are represented by purple ellipses in the center. Outer sphere is a perfectly-matched layer, to mitigate boundary effects caused by the finite acoustic domain. SPL is averaged over the inner sphere for the calculation of SRE. Wings vibrate along the y axis, as indicated by the cricket silhouette.

Figure Supplement 2-3. Mesh sensitivity analysis for models based on the finite element method. Each line represents the SPL of a single wing size with a different number of mesh elements. Each panel shows this analysis at a different frequency: A. 30 kHz, B. 20 kHz, C. 10 kHz, D. 5 kHz. As the difference between the second-largest and largest number of elements was small, we proceeded with the largest number of elements shown here for the analysis.

Supplement to Figure 2-4. Wing aspect ratio and use of a "harp" resonator do not significantly impact SRE within biologically-relevant ranges of wing size and call frequency. **A.** The effect of wing aspect ratio on SRE at six different call frequencies. The aspect ratio that was used on all models in this study (2) is shown in red dotted line. **B.** Actual distribution of aspect ratios among species. Red line indicates aspect ratio that was used in our models (2). We see that while aspect ratio influences SRE, this effect is minor within the realistic range of aspect ratios (typically <3 dB for ARs from 1 to 3.3). **B.** The effect on SRE of vibration spread over a small area (harp) compared with the whole wing. Some species of crickets restrict the vibrating portion of the wing to a "harp" region. However, we find that this does not strongly affect SRE at any frequency within our range of interest.

Figure 3. Sound propagation efficiency when the cricket sings near the bare hard ground is orders of magnitude higher than in free space. Each panel represents a combination of caller height above ground (0 m or 1 m) and receiver distance from caller (0.05 – 02 m "near" and 0.8 – 0.9 m "far"). **A.** Distance = near, height = 1m, **B.** Distance = far, height = 1 m; **C.** Distance = near, height = 1 m, **D.** Distance = far, height = 0 m. Red lines indicate optimal efficiency ridge, or the size at each frequency that would produce an ideally baffled calling scenario. Note that the frequency range is reduced compared to figure 2, in order to exclude high frequency callers which likely use alternative vibrational modes. Each clade of animals is represented by a colored ellipse. **E.** Phylogeny representing each clade **F.** Key to clade represented by each ellipse.

Supplement to figure 3-1. Geometry of boundary element models. Grey shape represents a ground of defined acoustic impedance. Purple ellipses represent wings. Dotted line illustrates the line along which measurements were taken to assess SPE.

Supplement to figure 3-2. Acoustic hardness of ground does not significantly influence SPE at biologically relevant ranges of wing size and call frequency. Comparison of SPE over hard vs soft ground. Each panel represents a combination of caller height above ground (0 m or 1 m) and receiver distance from caller (0.05 – 02 m "near" and 0.8 – 0.9 m "far"). **A.** Distance = near, height = 1m, **B.** Distance = far, height = 1 m; **C.** Distance = near, height = 1 m, **D.** Distance = far, height = 0 m. Color indicates whether higher SRE is found with hard ground (red shades), soft ground (blue shades) or no difference (white). Data are presented as a $\log_2$ ratio instead of a straight proportion. $\log_2$ ratios are scaled such that the ranges above and below 1 are proportional, rather than values below 1 being compressed between 0 and 1. Each clade of animals is represented by a colored ellipse. **E.** Phylogeny

Supplement to figure 3-3. Vegetation decreases efficiency overall but does not substantially change the landscape pattern of efficiency. Each panel represents a combination of caller height above ground (0 m or 1 m) and receiver distance from caller (0.05 – 02 m "near" and 0.8 – 0.9 m "far"). In each height and distance scenario, an excess attenuation factor due to vegetation was also applied. **A.** Distance = near, height = 1m, **B.** Distance = far, height = 1 m; **C.** Distance = near, height = 1 m, **D.** Distance = far, height = 0 m. Red lines indicate optimal efficiency ridge, or the size at each frequency that would produce an ideally baffled calling scenario. Each clade of animals is represented by a colored ellipse. **E.** Phylogeny representing each clade **F**. Key to clade represented by each ellipse.

Supplement to figure 3-4. Call directionality decreases at higher frequencies, particularly with grounded calling. Each panel represents a combination of caller height above ground (0 m or 1 m) and receiver distance from caller (0.05 – 02 m "near" and 0.8 – 0.9 m "far"). **A.** Distance = near, height = 1m, **B.** Distance = far, height = 1 m; **C.** Distance = near, height = 1 m, **D.** Distance = far, height = 0 m. Each clade of animals is represented by a colored ellipse. **E.** Phylogeny representing each clade **F.** Key to clade represented by each ellipse.

Figure 4. The most effective calling strategy (grounded vs baffled) varies depending on clade. Shown is a comparison of SPE on bare ground with no baffle, and 1 m in vegetation with and without an ideal baffle. These measurements were taken far from the caller, i.e. an average of the SPL at a distance of 0.8-0.9 m from wings, directly in front of the wings was used. Upper panel indicates extreme outliers, which only occur when we consider animals calling from the bare ground.

Supplement to Figure 4-1. Calculating efficiency with optimal baffle. **A.** ka at which optimal efficiency occurs for each frequency. Each frequency is represented by a different line. Line at which efficiency is maximized for each frequency (maximum efficiency ridge) is shown by red dashed line. **B.** efficiency at this optimal ka (1.55) in open and vegetation conditions.

Supplement to figure 4-2. Vegetation somewhat decreases efficiency in baffled and grounded calling conditions. Differences in SPE by clade depending on vegetation and baffle use. For each clade, bars with black outline represent SPE without baffle and bars with red outline represent SPE with ideal baffle. Background color of the bars indicate vegetation or no vegetation.

Figure 5. Baffled and grounded calling do not substantially differ with respect to directionality, although it does vary somewhat by clade. **A.** Directionality in near condition, **B.** Directionality in far condition.

Table 1. Sources for morphological data. Asterisk after species name indicates species is represented in both frequency and wing size datasets

Clade	Subfamily	Genus	Species	Specimen	Relevant reference and/or collection specimen number
A	Gryllotalpinae	*Gryllotalpa*	*australis*[*]	1	Orthoptera Species File Specimen ID: 45466
			gryllotalpa[*]	1	Linnean Collection Specimen ID: LINN 8925
			orientalis[*]	1	Orthoptera Species File Taxon ID: 1128860
			permai[*]	1	(Tan and Kamaruddin, 2016)
			vineae[*]	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF4425
				2	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF4425
				3	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF4425
B	Mogoplistinae	*Cycloptiloides*	*canariensis*[*]	1	(Dambach and Gras, 1995)
		Cycloptilum	*irregularis*[*]	1	(Love and Walker, 1979)
			slossoni[*]	1	(Love and Walker, 1979)
			tardum[*]	1	(Love and Walker, 1979)
		Hoplosphyrum	*boreale*[*]	1	(Love and Walker, 1979)
		Ornebius	*bimaculatus*[*]	1	(Kim, 2011)
			kanetataki[*]	1	(Kim, 2011)
C	Nemobiinae	*Allonemobius*	*allardi*[*]	1	Orthoptera Species File Specimen ID: 40715
		Bobilla	*gullanae*[*]	1	(Su and Rentz, 2000)
			neobivittata[*]	1	(Su and Rentz, 2000)
		Hygronemobius	*guriri*	1	(Pereira et al., 2013)
			indaia[*]	1	(Pereira et al., 2013)
			iperoigae[*]	1	(Pereira et al., 2013)
		Nemobius	*sylvestris*[*]	1	(Barranco et al., 2013)
		Pteronemobius	*nigrovus*[*]	1	(McIntyre, 1977)
	Trigonidiinae	*Anaxipha*	*bradephona**	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF6482

Clade	Subfamily	Genus	Species	Specimen	Relevant reference and/or collection specimen number
			hyalicetra*	1	(Cole and Funk, 2019)
			tachephona*	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF6486
		Cranistus	colliurides*	1	(Martins et al., 2012)
		Phylloscirtus	amoenus*	1	(Martins et al., 2012)
D	Pteroplistinae	Singapuriola	separata*	1	(Gorochov and Tan, 2012)
E	Luzarinae	Lerneca	inalata*	1	(Lima et al., 2018)
		Luzaridella	susurra*	1	(Martins et al., 2013)
		Vanzoliniella	sambophila*	1	(Mello and Reis, 1994)
	Paragryllinae	Alcodes	chamocoru	1	Orthoptera Species File Specimen ID: 65179
			mococharu	1	Orthoptera Species File Specimen ID: 65181
		Aclogryllus	crybelos*	1	(Nischk and Otte, 2000)
		Escondacla	thymodes*	1	Orthoptera Species File Specimen ID: 65198
		Neoacla	clandestina*	1	Orthoptera Species File Specimen ID: 65199
		Silvastella	epiplatys*	1	Orthoptera Species File Specimen ID: 65196
	Phalangopsinae	Ceyloria	latissima	1	Orthoptera Species File Specimen ID: 2983
		Endecous	betariensis*	1	(Mello and Pellegatti-Franco, 1998)
			chape*	1	(Souza-Dias et al., 2017)
			didymus*	1	(Desutter-Grandcolas, 2009)
			itatibensis*	1	(Mello and Pellegatti-Franco, 1998)
			naipi	1	(Souza-Dias et al., 2017)
			troglobius*	1	(Castro-Souza et al., 2020)
		Lernecella	trinitatis	1	Orthoptera Species File Taxon ID: 1125930
		Pseudotrigonidium	personatum	1	(Desutter-Grandcolas, 2009)

Clade	Subfamily	Genus	Species	Specimen	Relevant reference and/or collection specimen number
E	Phalangopsinae	*Tremellia*	*timah**	1	(Gorochov and Tan, 2012)
	Phaloriinae	*Phaloria*	*anapina**	1	(Otte, 2007)
			*chopardi**	1	(Desutter-Grandcolas, 2009)
			*jerelynae**	1	(Gorochov and Tan, 2012)
		Trellius	*neesoon*	1	(Gorochov and Tan, 2012)
F	Eneopterinae	*Gnominthus*	*baitabagus**	1	(Robillard and Su, 2018)
		Pixibinthus	*sonicus**	1	(Anso et al., 2016)
	Hapithinae	*Hapithus*	*agitator**	1	Orthoptera Species File Specimen ID: 138599
			*vagus**	1	Orthoptera Species File Specimen ID: 65035
	Oecanthinae	*Neoxabea*	*bipunctata**	1	Image captured for present study in lab
			*brevipes**	1	(Zefa et al., 2018)
			*cerrojesusensis**	1	Image captured for present study in lab
			*oltei**	1	Image captured for present study in lab
		Oecanthus	*alexanderi**	1	Image captured for present study in lab
			*angustus**	1	PaDILspecies ID: *Oecanthus angustus*
			*argentinus**	1	University of British Columbia Insect Collection, Specimen: SEM-UBC GRY-0951
			*forbsei**	1	Image captured for present study in lab
			*fultoni**	1	Orthoptera Species File Specimen ID: 40710
			*latipennis**	1	University of Guelph Insect Collection: Specimen BIOUG44550-E07
				2	University of Guelph Insect Collection: Specimen BIOUG44550-E08
			*lineolatus**	1	(Zefa et al., 2012)
			major	1	Orthoptera Species File Specimen ID: 40712
		Oecanthus	*nigricornis**	1	Orthoptera Species File Taxon ID: 345166

Clade	Subfamily	Genus	Species	Specimen	Relevant reference and/or collection specimen number
F	Oecanthinae	*Oecanthus*	*niveus**	1	Orthoptera Species File Taxon ID: 345151
			*pallidus**	1	(Zefa et al., 2012)
			*pictus**	1	(Milach et al., 2015)
			*pini**	1	Image captured for present study in lab
			*quadripunctatus**	1	https://www.insectimages.org/browse/subthumb.cfm?sub=9113
				2	UBC Database ID: SEM-UBC GRY-0918
			*rileyi**	1	Orthoptera Species File Taxon ID: 1128127
			*rufescens**	1	NHM Specimen ID: 012497644
				2	NHM Specimen ID: 012497645
				3	NHM Specimen ID: 012497646
				4	PaDIL species ID: *Oecanthus rufescens*
			*texensis**	1	Image captured for present study in lab
			valensis	1	(Milach et al., 2016)
			*varicornis**	1	Image captured for present study in lab
	Podoscirtinae	*Madasumma*	*affinis**	1	(Otte, 2007)
		Truljalia	*formosa**	1	(He, 2012)
	Podoscirtinae	*Varitrella*	*suikei**	1	(Tan et al., 2020)
G	Eneopterinae	*Agnotecous*	*azurensis**	1	(Desutter-Grandcolas and Robillard, 2006)
			*brachypterus**	1	(Robillard et al., 2010)
			*meridionalis**	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF1775
			*pinsula**	1	(Robillard et al., 2010)
			*sarramea**	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF988
			*yahoue**	1	(Desutter-Grandcolas and Robillard, 2006)

Clade	Subfamily	Genus	Species	Specimen	Relevant reference and/or collection specimen number
G	Eneopterinae	*Arilpa*	*binderia**	1	(Otte, 2007)
			*gidya**	1	(Otte, 2007)
		Cardiodactylus	*guttulus**	1	(Robillard and Ichikawa, 2009)
			*novaeguinea**	1	(Robillard and Ichikawa, 2009)
		Eurepa	*bifasciata**	1	(Robillard and Su, 2018)
		Lebinthus	*bitaeniatus**	1	(Robillard et al., 2013)
			*luae**	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF3208
		Myara	*pakaria**	1	(Otte, 2007)
			*wintrena**	1	(Otte, 2007)
		Salmanites	*peekara**	1	(Otte, 2007)
		Xenogryllus	*eneopteroides**	1	(Jaiswara et al., 2019)
			*transversus**	1	(Jaiswara et al., 2019)
	Gryllinae	Eurepella	*mjobergi**	1	PaDIL species ID: *Eurepella mjobergi*
		Gryllus	*amarensis*	1	Museum D'Historie Naturelle ID: 7031
			*assimilis**	1	SINA species ID: *Gryllus assimilis*
			*bimaculatus**	1	Orthoptera Species File Taxon ID: 1122377
			*brevicaudus**	1	SINA species ID: *Gryllus brevicaudus*
			*campestris**	1	Need to figure out specific specimen
				2	Need to figure out specific specimen
			carvalhoi	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF7242
			chaldeus	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF7192
			*chappuisi**	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF7046
			*cohni**	1	(Weissman and Gray, 2019)

Clade	Subfamily	Genus	Species	Specimen	Relevant reference and/or collection specimen number
G	Gryllinae	*Gryllus*	*firmus**	1	(Weissman and Gray, 2019)
			*fultoni**	1	Orthoptera Species File Specimen ID: 40672
			*lineaticeps**	1	(Weissman and Gray, 2019)
			*multipulsator**	1	(Weissman and Gray, 2019)
			*pennsylvanicus**	2	Orthoptera Species File Specimen ID: 43773
				3	UBC Database ID: SEM-UBC GRY-0542
			*veletis**	1	Orthoptera Species File Specimen ID: 40674
				2	UBC Database ID: SEM-UBC GRY-0643
			*vocalis**	1	Orthoptera Species File Specimen ID: 64224
		Miogryllus	*itaquiensis**	1	(Orsini et al., 2017)
			*piracicabensis**	1	(Orsini et al., 2017)
		Teleogryllus	*commodus**	1	(Otte, 2007)
			*marini**	1	(Otte, 2007)
			*oceanicus**	1	(Otte, 2007)
	Itarinae	*Itara*	*kirejtshuki**	1	NMHUK 012497661
			*minor**	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF8162
	Landrevinae	*Striduleva*	*crepitans**	1	Museum D'Historie Naturelle ID: MNHN-EO-ENSIF2059

Clade	Subfamily	Genus	Species	Recording	Relevant reference and/or collection specimen number
A	Gryllotalpa	*Gryllotalpa*	*australis**	1-47	(Kavanagh and Young, 1989) (range of values given in publication)
			*fulvipes**	1	(Tan and Kamaruddin, 2016)
			*gryllotalpa**	1	Orthoptera Species File Sound ID: 1176
			*permai**	1	(Tan and Kamaruddin, 2016)
			*vineae**	1	Orthoptera Species File Sound ID: 1198
			*canariensis**	1	(Dambach and Gras, 1995)
B	Mogoplistinae	*Cycloptiloides*	*irregularis**	1	Crickets north of Mexico species Id: Key's scaly cricket
		Cycloptilum	*slossoni**	1	Crickets north of Mexico species Id: Slosson's scaly cricket
			*tardum**	1	(Otte, 2007)
			*boreale**	1	Crickets north of Mexico species Id: long-winged scaly cricket
		Hoplosphyrum	*bimaculatus**	1	(He et al., 2017)
		Ornebius	*kanetataki**	1	(He et al., 2017)
			*allardi**	1	Crickets north of Mexico species Id: Allard's ground cricket
C	Nemobiinae	*Allonemobius*	*gullanae**	1	(Su and Rentz, 2000)
		Bobilla	*neobivittata**	1	(Su and Rentz, 2000)
			*indaia**	1	(Pereira et al., 2013)
		Hygronemobius	*iperoigae**	1	(Pereira et al., 2013)
			*sylvestris**	1	Orthoptera Species File Sound ID: 1045

Clade	Subfamily	Genus	Species	Recording	Relevant reference and/or collection specimen number
C	Nemobiinae	*Nemobius*	*nigrovus**	1	(McIntyre, 1977)
		Pteronemobius	*bradephona**	1	Orthoptera Species File Sound ID: 1832
	Trigonidiinae	*Anaxipha*	*hyalicetra**	1	(Cole and Funk, 2019)
			tachephona*	1	Orthoptera Species File Sound ID: 1833
			colliurides*	1	(Martins et al., 2012)
		Cranistus	*amoenus**	1	(Martins et al., 2012)
		Phylloscirtus	*separata**	1	(Gorochov and Tan, 2012)
D	Pteroplistinae	*Singapuriola*	*inalata**	1	(Lima et al., 2018)
E	Luzarinae	*Lerneca*	*susurra**	1	(Martins et al., 2013)
		Luzaridella	*sambophila**	1	(Mello and Reis, 1994)
		Vanzoliniella	*chamocoru**	1	(Nischk and Otte, 2000)
	Paragryllinae	Aclodes	*mococharu**	1	(Nischk and Otte, 2000)
			crybelos*	1	(Nischk and Otte, 2000)
		Aclogryllus	*thymodes**	1	(Nischk and Otte, 2000)
		Escondacla	*clandestine**	1	(Nischk and Otte, 2000)
		Neoacla	*epiplatys**	1	(Nischk and Otte, 2000)
		Silvastella	*betariensis**	1	(He, 2012)
	Phalangopsinae	Endecous	*chape**	1	(Souza-Dias et al., 2017)

Clade	Subfamily	Genus	Species	Recording	Relevant reference and/or collection specimen number
			*didymus**	1	(Castro-Souza et al., 2020)
E	Phalangopsinae	*Endecous*	*itatibensis**	1	(Mello and Pellegatti-Franco, 1998)
			*troglobius**	1	(Castro-Souza et al., 2020)
			*timah**	1	(Gorochov and Tan, 2012)
		Tremellia	*anapina**	1	(Su and Rentz, 2000)
	Phaloriinae	*Phaloria*	*chopardi**	1	(Desutter-Grandcolas, 2009)
			*jerelynae**	1	(Gorochov and Tan, 2012)
			*baitabagus**	1	(Vicente et al., 2015)
F	Eneopterinae	*Gnominthus*	*sonicus**	1	(Anso et al., 2016)
		Pixibinthus	*agitator**	1	Crickets north of Mexico species ID: *Hapithus agitator*
	Hapithinae	*Hapithus*	*melodius*	1	Handbook of crickets and katydids
			*vagus**	1	Macaulay Library asset: 114470
			diplastes	1	Handbook of crickets and katydids
		Orocharis	*gryllodes*	1	Handbook of crickets and katydids
			luteolira	1	Handbook of crickets and katydids
			nigrifrons	1	Handbook of crickets and katydids
			saltator	1	Handbook of crickets and katydids
			tricornis	1	Handbook of crickets and katydids

Clade	Subfamily	Genus	Species	Recording	Relevant reference and/or collection specimen number
F	Hapithinae	*Orocharis*	*bipunctata**	1	Crickets north of Mexico species ID: *Neoxabea bipunctata*
	Oecanthinae	*Neoxabea*	*brevipes**	1	(Zefa et al., 2018)
			*cerrojesusensis**	1	Orthoptera Species File Sound ID: 2345
			*ottei**	1	Orthoptera Species File Sound ID: 2346
			*alexanderi**	1	Crickets North of Mexico Species ID: *Oecanthus alexanderi*
		Oecanthus	*angustus**	1	(Otte, 2007)
			*argentinus**	1	Crickets North of Mexico Species ID: *Oecanthus argentinus*
			*argentinus** *californicus*	2	Orthoptera Species File Sound ID: 1535
				1	Crickets North of Mexico Species ID: *Oecanthus californicus*
			californicus *forbsei**	2	Orthoptera Species File Sound ID: 1536
				1	Crickets North of Mexico Species ID: *Oecanthus forbsei*
			*fultoni**	1	Crickets North of Mexico Species ID: *Oecanthus fultoni*
			*latipennis**	1	Crickets North of Mexico Species ID: *Oecanthus latipennis*
			*latipennis**	2	Orthoptera Species File Sound ID: 1002
			*lineolatus**	1	(Zefa et al., 2012)
			*nigricornis**	1	Crickets North of Mexico Species ID: *Oecanthus nigricornis*
			*niveus**	1	Crickets North of Mexico Species ID: *Oecanthus niveus*
			*pallidus**	1	(Zefa et al., 2012)

Clade	Subfamily	Genus	Species	Recording	Relevant reference and/or collection specimen number
F	Oecanthinae	*Oecanthus*	*pictus**	1-9	Orthoptera Species File Taxon ID: 1223417 (9 songs from different temperatures)
			*pini**	1	Crickets North of Mexico Sound File: 587sl
			*quadripunctatus**	1	Orthoptera Species File Sound File 1531
			*rileyi**	1	Orthoptera Species File Sound File: 1540
			*rufescens**	1	(Otte, 2007)
			*texensis**	1	(Symes and Collins, 2013)
			*varicornis**	1	Crickets North of Mexico Sound File: 593sl
			walker	1	Crickets North of Mexico Species ID: *Oecanthus walkeri*
			*affinis**	1	(Otte, 2007)
	Podoscirtinae	*Madasumma*	*jirranda*	1	(Otte, 2007)
			kanina	1	(Otte, 2007)
			loorea	1	(Otte, 2007)
			*formosa**	1	(He, 2012)
		Truljalia	*suikei**	1	(Tan et al., 2020)
		Varitrella	*azurensis**	1	Museum D'Historie Naturelle ID: MNHN-SO-2018-100
G	Eneopterinae	*Agnotecous*	*brachypterus**	1	(Robillard et al., 2010)
			clarus	1	Museum D'Historie Naturelle ID: MNHN-SO-2018-102
			*meridionalis**	1	Museum D'Historie Naturelle ID: MNHN-SO-2018-99

Clade	Subfamily	Genus	Species	Recording	Relevant reference and/or collection specimen number
G	Eneopterinae	*Agnotecous*	*pinsula**	1	(Robillard et al., 2010)
			*sarramea**	1	(Robillard and Desutter-Grandcolas, 2004)
			*yahoue**	1	(Robillard and Desutter-Grandcolas, 2004)
			*binderia**	1	(Otte, 2007)
		Arilpa	*gidya**	1	(Otte, 2007)
			wirrilla	1	(Otte, 2007)
			*guttulus**	1	(Robillard and Ichikawa, 2009)
		Cardiodactylus	*novaeguinea**	1	(Otte, 2007)
			*bifasciata**	1	(Otte, 2007)
		Eurepa	*eeboolaga*	1	(Otte, 2007)
			marginipennis	1	(Otte, 2007)
			noarana	1	(Otte, 2007)
			nurndina	1	(Otte, 2007)
			wirkutta	1-2	(Otte, 2007) (range of values given in publication)
			woortooa	1	(Otte, 2007)
			yumbena	1	(Otte, 2007)
			*bitaeniatus**	1	(Robillard and Tan, 2013)
		Lebinthus	*luae**	1	(Robillard and Tan, 2013)

Clade	Subfamily	Genus	Species	Recording	Relevant reference and/or collection specimen number
G	Eneopterinae	*Lebinthus*	*aperta*	1	(Otte, 2007)
		Myara	*marimbula*	1	(Otte, 2007)
			muttaburra	1	(Otte, 2007)
			pakaria*	1	(Otte, 2007)
			sordida	1	(Otte, 2007)
			unicolor	1-2	(Otte, 2007) (range of values given in publication)
			wintrena*	1	(Robillard and Desutter-Grandcolas, 2004)
			yurgama	1	(Otte, 2007)
			vittatus	1	(Robillard and Desutter-Grandcolas, 2004)
		Nisitrus	*allaris*	1	(Otte, 2007)
		Salmanites	*ninbella*	1	(Otte, 2007)
			noccundris	1	(Otte, 2007)
			noonamina	1	(Otte, 2007)
			peekara*	1	(Otte, 2007)
			poene	1	(Otte, 2007)
			taltantris	1	(Otte, 2007)
			terba	1-2	(Otte, 2007) (range of values given in publication)
			wittilliko	1	(Otte, 2007)

Clade	Subfamily	Genus	Species	Recording	Relevant reference and/or collection specimen number
G	Eneopterinae	*Salmanites*	*eneopteroides**	1	(Jaiswara et al., 2019)
		Xenogryllus	*maichauensis*	1	(Jaiswara et al., 2019)
			marmoratus	1	(Jaiswara et al., 2019)
			mozambicus	1	(Jaiswara et al., 2019)
			*transversus**	1	Database found within http://www.biologie.uni-ulm.de, no longer exists
				2	Database found within http://www.biologie.uni-ulm.de, no longer exists
			ululiu	1	Database found within http://www.biologie.uni-ulm.de, no longer exists
			ballina	1	(Otte, 2007)
	Gryllinae	*Eurepella*	*iando*	1	(Otte, 2007)
			jillangolo	1	(Otte, 2007)
			kulkawirra	1	(Otte, 2007)
			lewara	1	(Otte, 2007)
			mataranka	1	(Otte, 2007)
			meda	1	(Otte, 2007)
			*mjobergi**	1-2	(Otte, 2007) (range of values given in publication)
			moojerra	1	(Otte, 2007)
			oana	1	(Otte, 2007)
			quarriana	1	(Otte, 2007)

Clade	Subfamily	Genus	Species	Recording	Relevant reference and/or collection specimen number
G	Gryllinae	*Eurepella*	*tinga*	1	(Otte, 2007)
			tjairaia	1	(Otte, 2007)
			torowatta	1	(Otte, 2007)
			wanga	1	(Otte, 2007)
			waninga	1	(Otte, 2007)
		Gryllus	*assimilis**	1	Crickets North of Mexico sound file: 483sl
				2	Crickets North of Mexico sound file: 483ss2
			*bimaculatus**	1	Orthoptera Species File Sound ID: 1295
			*brevicaudus**	1	Crickets North of Mexico sound file: 465sldw
				2	Crickets North of Mexico sound file: 465ss2wg
			*campestris**	1	Orthoptera Species File sound ID: 1741
			*chappuisi**	1	Orthoptera Species File sound ID: 1739
			*cohni**	1	Crickets North of Mexico sound file: 722sl
			*firmus**	1	Crickets North of Mexico sound file: 481sl
			*fultoni**	1	Crickets North of Mexico sound file: 484sl
				2	Crickets North of Mexico sound file: 484slc
			*lineaticeps**	1	Crickets North of Mexico sound file: 467sldw
			*multipulsator**	1	Crickets North of Mexico sound file: 499sl

Clade	Subfamily	Genus	Species	Recording	Relevant reference and/or collection specimen number
G	Gryllinae	*Gryllus*	*multipulsator**	2	Crickets North of Mexico sound file: 499slwg
			*pennsylvanicus**	1	Orthoptera Species File sound ID: 1258
			texensis	1	Crickets North of Mexico sound file: 479sl
			*veletis**	1	Crickets North of Mexico sound file: 488sl
			*vocalis**	1	Crickets North of Mexico sound file: 466sldw
			*itaquiensis**	1	(Otte, 2007)
		Miogryllus	*piracicabensis**	1-30	(Orsini et al., 2017) (range of values given in publication)
			*commodus**	1	(Otte, 2007)
		Teleogryllus	*marini**	1-2	(Otte, 2007) (range of values given in publication)
			*oceanicus**	1	(Otte, 2007)
		Itara	*kirejtshuki**	1	Orthoptera Species File sound ID: 1796
			*minor**	1	Database found within http://www.biologie.uni-ulm.de, no longer exists
		Striduleva	*crepitans**	1-2	(Hugel, 2009) (range of values given in publication)

Full citations for Specimen data:

Anso J, Barrabé L, Desutter-Grandcolas L, Jourdan H, Grandcolas P, Dong J, Robillard T. 2016. Old Lineage on an Old Island: Pixibinthus, a New Cricket Genus Endemic to New Caledonia Shed Light on Gryllid Diversification in a Hotspot of Biodiversity. *PLOS ONE* **11**:e0150920. doi:10.1371/journal.pone.0150920

Barranco P, Gilgado JD, Ortuño VM. 2013. A new mute species of the genus Nemobius Serville (Orthoptera, Gryllidae, Nemobiinae) discovered in colluvial, stony debris in the Iberian Peninsula: A biological, phenological and biometric study. *Zootaxa* **3691**:201–219. doi:10.11646/zootaxa.3691.2.1

Castro-Souza RA, Zefa E, Ferreira RL. 2020. New troglobitic and troglophilic syntopic species of Endecous (Orthoptera, Grylloidea, Phalangopsidae) from a Brazilian cave: a case of sympatric speciation? *Zootaxa* **4810**:271–304. doi:10.11646/zootaxa.4810.2.3

Cole JA, Funk DH. 2019. Anaxipha hyalicetra sp. n. (Gryllidae: Trigonidiinae), a new sword-tailed cricket species from Arizona. *Journal of Orthoptera Research* **28**:3–9.

Dambach M, Gras A. 1995. Bioacoustics of a miniature cricket, Cycloptiloides canariensis (Orthoptera: Gryllidae: Mogoplistinae). *Journal of Experimental Biology* **198**:721–728. doi:10.1242/jeb.198.3.721

Desutter-Grandcolas L. 2009. New and little known crickets from Espiritu Santo Island, Vanuatu (Insecta, Orthoptera, Grylloidea, Pseudotrigonidium Chopard, 1915, Phaloriinae and Nemobiinae p.p.). *zoos* **31**:619–659. doi:10.5252/z2009n3a12

Desutter-Grandcolas L, Robillard T. 2006. Phylogenetic systematics and evolution of Agnotecous in New Caledonia (Orthoptera: Grylloidea, Eneopteridae). *Systematic Entomology* **31**:65–92. doi:10.1111/j.1365-3113.2005.00299.x

Gorochov AV, Tan MK. 2012. New crickets of the subfamilies Phaloriinae and Pteroplistinae (Orthoptera: Gryllidae) from Singapore. *Zootaxa* **3525**:18–34. doi:10.11646/zootaxa.3525.1.2

He Z. 2012. A new species of Truljalia Gorochov 1985 from Taiwan (Orthoptera: Gryllidae; Podoscirtinae; Podoscirtini). *Zootaxa* **3591**:79–83. doi:10.11646/zootaxa.3591.1.5

He Z, Lu H, Liu Y, Li K. 2017. A new species of Ornebius Guérin-Méneville, 1844 from East China (Orthoptera: Mogoplistidae: Mogoplistinae). *Zootaxa* **4303**:445–450. doi:10.11646/zootaxa.4303.3.10

Hugel S. 2009. New Landrevinae from Mascarene islands and little known Landrevinae from Africa and Comoros (Grylloidea: Landrevinae). *Annales de la Société entomologique de France (NS)* **45**:193–215. doi:10.1080/00379271.2009.10697602

Jaiswara R, Dong J, Ma L, Yin H, Robillard T. 2019. Taxonomic revision of the genus Xenogryllus Bolívar, 1890 (Orthoptera, Gryllidae, Eneopterinae, Xenogryllini). *Zootaxa* **4545**:301–338. doi:10.11646/zootaxa.4545.3.1

Kavanagh MW, Young D. 1989. Bilateral symmetry of sound production in the mole cricket, Gryllotalpa australis. *J Comp Physiol A* **166**:43–49. doi:10.1007/BF00190208

Kim T. 2011. A Taxonomic Study of the Scaly Cricket Family Mogoplistidae (Orthoptera: Ensifera: Grylloidea) in Korea. *Zootaxa* **2928**:41–48. doi:10.11646/zootaxa.2928.1.4

Lima RM, Schuchmann K-L, Tissiani AS, Nunes LA, Jahn O, Ganchev TD, Lhano MG, Marques MI. 2018. Tegmina-size variation in a Neotropical cricket with implications on spectral song properties. *Journal of Natural History* **52**:1225–1241. doi:10.1080/00222933.2018.1457728

Love RE, Walker TJ. 1979. Systematics and Acoustic Behavior of Scaly Crickets (Orthoptera: Gryllidae: Mogoplistinae) of Eastern United States. *Transactions of the American Entomological Society (1890-)* **105**:1–66.

Martins L de P, Redü DR, Oliveira GL de, Zefa E. 2012. Recognition characters and new records of two species of Phylloscyrtini (Orthoptera, Gryllidae, Trigonidiinae) from southern Brazil. *Iheringia, Sér Zool* **102**:95–98. doi:10.1590/S0073-47212012000100013

Martins LDP, Silva LGD, Henriques AL, Zefa E. 2013. First record of the genera Luzarida Hebard, 1928 and Luzaridella Desutter-Grandcolas, 1992 (Orthoptera, Gryllidae, Phalangopsinae) from Brazil, including a new species and description of the female of Luzarida lata Gorochov, 2011. *Zootaxa* **3609**:421–430. doi:10.11646/zootaxa.3609.4.4

McIntyre ME. 1977. Acoustical communication in the field crickets Pteronemobius nigrovus and P. bigelowi (Orthoptera: Gryllidae). *New Zealand Journal of Zoology* **4**:63–72. doi:10.1080/03014223.1977.9517937

Mello F de AG de, Pellegatti-Franco F. 1998. A New Cave Cricket of the Genus Endecous from Southeastern Brazil and Characterization of Male and Female Genitalia of E. itatibensis Rehn, 1918 (Orthoptera: Grylloidea: Phalangopsidae: Luzarinae). *Journal of Orthoptera Research* 185–188. doi:10.2307/3503517

Mello F de AG de, Reis JC dos. 1994. Substrate Drumming and Wing Stridulation Performed during Courtship by a New Brazilian Cricket (Orthoptera: Grylloidea: Phalangopsidae). *Journal of Orthoptera Research* 21–24. doi:10.2307/3503603

Milach EM, Costa MKMD, Martins LDP, Nunes LA, Silva DSM, Garcia FRM, Oliveira ECD, Zefa E. 2016. New species of tree cricket Oecanthus Serville, 1831 (Orthoptera: Gryllidae: Oecanthinae) from Reserva Natural Vale, Espírito Santo, Brazil, with chromosome complement. *Zootaxa* **4173**:137–146. doi:10.11646/zootaxa.4173.2.4

Milach EM, Martins LDP, Costa MKMD, Gottschalk MS, Oliveira GLD, Redü DR, Neutzling AS, Dornelles JEF, Vasconcellos LA, Zefa E. 2015. A new species of tree crickets Oecanthus (Orthoptera, Gryllidae, Oecanthinae) in tobacco plantation from Southern Brazil, with body color variation. *Zootaxa* **4018**:266–278. doi:10.11646/zootaxa.4018.2.6

Nischk F, Otte D. 2000. Bioacoustics, Ecology and Systematics of Ecuadorian Rainforest Crickets (Orthoptera: Gryllidae: Phalangopsinae), with a Description of Four New Genera and Ten New Species. *Journal of Orthoptera Research* 229–254. doi:10.2307/3503651

Orsini MP, Costa MKMD, Szinwelski N, Martins LDP, Corrêa RC, Timm VF, Zefa E. 2017. A new species of Miogryllus Saussure, 1877 and new record of Miogryllus piracicabensis Piza, 1960 (Orthoptera: Gryllidae) from State of Rio Grande do Sul, Brazil, with calling song and chromosome complement. *Zootaxa* **4291**:361–372. doi:10.11646/zootaxa.4291.2.8

Otte D. 2007. Australian Crickets (Orthoptera: Gryllidae). Academy of Natural Sciences.

Pereira MR, Miyoshi AR, Martins LDP, Fernandes ML, Sperber CF, Mesa A. 2013. New Neotropical species of Hygronemobius Hebard, 1913 (Orthoptera: Grylloidea: Nemobiinae), including a brief discussion of male genitalia morphology and preliminary biogeographic considerations of the genus. *Zootaxa* **3641**:1–20. doi:10.11646/zootaxa.3641.1.1

Robillard T, Desutter-Grandcolas L. 2004. High-frequency calling in Eneopterinae crickets (Orthoptera, Grylloidea, Eneopteridae): adaptive radiation revealed by phylogenetic analysis. *Biological Journal of the Linnean Society* **83**:577–584. doi:10.1111/j.1095-8312.2004.00417.x

Robillard T, Ichikawa A. 2009. Redescription of Two Cardiodactylus Species (Orthoptera, Grylloidea, Eneopterinae): The Supposedly Well-Known C. novaeguineae (Haan, 1842), and the Semi-Forgotten C. guttulus (Matsumura, 1913) from Japan. *jzoo* **26**:878–891. doi:10.2108/zsj.26.878

Robillard T, Nattier R, Desutter-Grandcolas L. 2010. New species of the New Caledonian endemic genus Agnotecous (Orthoptera, Grylloidea, Eneopterinae, Lebinthini). *Zootaxa* **2559**:17–35. doi:10.11646/zootaxa.2559.1.2

Robillard T, Su YN. 2018. New lineages of Lebinthini from Australia (Orthoptera: Gryllidae: Eneopterinae). *Zootaxa* **4392**:241–266. doi:10.11646/zootaxa.4392.2.2

Robillard T, Tan MK. 2013. A taxonomic review of common but little knwon crickets from Singapore and the Philippines (Insecta: Orthoptera: Eneoptera: Eneopterinae) 21.

Robillard T, Yap S, Yngente MV. 2013. Systematics of cryptic species of Lebinthus crickets in Mount Makiling (Grylloidea, Eneopterinae). *Zootaxa* **3693**:49–63. doi:10.11646/zootaxa.3693.1.3

Souza-Dias PGB, Szinwelski N, Fianco M, Oliveira ECD, Mello FD a. GD, Zefa E. 2017. New species of Endecous (Grylloidea, Phalangopsidae, Luzarinae) from the Iguaçu National Park (Brazil), including bioacoustics, cytogenetic and distribution data. *Zootaxa* **4237**:454–470. doi:10.11646/zootaxa.4237.3.2

Su YN, Rentz DCF. 2000. Australian Nemobiine Crickets: Behavioral Observations and New Species of Bobilla Otte & Alexander (Orthoptera: Gryllidae: Nemobiinae). *Journal of Orthoptera Research* 5–20. doi:10.2307/3503626

Symes LB, Collins N. 2013. Oecanthus Texensis: A New Species of Tree Cricket from the Western United States. *orth* **22**:87–91. doi:10.1665/034.022.0203

Tan MK, Japir R, Chung AYC, Wahab RBHA. 2020. New taxa of crickets (Orthoptera: Grylloidea: Phaloriinae, Phalangopsinae, Itarinae and Podoscirtinae) from Borneo (Brunei Darussalam and Sandakan). *Zootaxa* **4810**:244–270. doi:10.11646/zootaxa.4810.2.2

Tan MK, Kamaruddin KN. 2016. A new species of Gryllotalpa mole cricket (Orthoptera: Gryllotalpidae: (Gryllotalpinae) from Peninsular Malaysia. *Zootaxa* **4066**:552–560. doi:10.11646/zootaxa.4066.5.3

Vicente NM, Olivero P, Lafond A, Dong J, Robillard T. 2015. Gnominthus gen. nov., a new genus of crickets endemic to Papua New Guinea with novel acoustic and behavioral diversity (Insecta, Orthoptera, Gryllidae, Eneopterinae). *Zoologischer Anzeiger - A Journal of Comparative Zoology* **258**:82–91. doi:10.1016/j.jcz.2015.06.005

Weissman DB, Gray DA. 2019. Crickets of the genus Gryllus in the United States (Orthoptera: Gryllidae: Gryllinae). *Zootaxa* **4705**:1–277. doi:10.11646/zootaxa.4705.1.1

Zefa E, Acosta RC, Timm VF, Szinwelski N, Marinho MAT, Costa MKMD. 2018. The Tree Cricket Neoxabea brevipes Rehn, 1913 (Orthoptera: Gryllidae: Oecanthinae) from the Brazilian southern Atlantic Forest: morphology, bioacoustics and cytogenetics. *Zootaxa* **4531**:554–566. doi:10.11646/zootaxa.4531.4.6

Zefa E, Neutzling AS, Redü DR, Oliveira GLD, Martins LDP. 2012. A new species of Oecanthus and Oecanthus lineolatus Saussure, 1897 from Southern Brazil: species description, including phallic sclerites, metanotal glands and calling song (Orthoptera: Gryllidae: Oecanthinae). *Zootaxa* **3360**:53–67. doi:10.11646/zootaxa.3360.1.2