1. **Manuscript Title:** Chronic Alcohol Drinking Drives Sex-Specific Differences in Affective Behavior and Medial Prefrontal Cortex Activity in CRF1:Cre:Tdtomato Transgenic Rats

2. **Abbreviated Title:** EtOH Alters Affect & PFC Activity in CRF1-cre rats

3. **Author names & Affiliations:** Quadir SG[1], Arleth GM[1], Cone MG[1], High MW[1], Ramage MC[1], Effinger DP[2], Echeveste-Sanchez M[1], Herman MA[1,2]
Affiliations:
[1] Bowles Center for Alcohol Studies, University of North Carolina Chapel Hill
[2] Department of Pharmacology, University of North Carolina Chapel Hill

4. Author Contributions:
SGQ: designed research, performed research, analyzed data, wrote the paper; **GMA**: designed research, performed research; **MGC** performed research; **MWH:** performed research; **DPE:** performed research; **MCR**: performed research **ME:** performed research **MAH**: designed research, wrote the paper, funding acquisition

5. Correspondence should be addressed to
Melissa A Herman
1027 Thurston-Bowles Building CB 7178
Chapel Hill, NC 27599-7178
Email: melissa_herman@unc.edu

6. Number of Figures: 6

7. Number of Tables: 1

8. Number of Multimedia: 0

9. Number of Words for Abstract: 240

10. Number of Words for Significance: 97

11. Number of Words for Introduction: 508

12. Number of Words for Discussion: 1842

13. Acknowledgements: We thank Jasmine J Jahad for their help in early intermittent access drinking experiments.

14. Conflicts of Interest: Authors report no conflict of Interest.

15. Funding Sources: This work was supported by National Institute of Health grants AA011605 (MAH), AA026858 (MAH), AA007573 (SGQ), AA030493 (SGQ) and GM135095 (DPE).

ABSTRACT

Alcohol use disorders (AUDs) are characterized by compulsive alcohol use, loss of control over intake, and a negative emotional state during abstinence. While AUDs are associated with both mood and chronic pain disorders, the relationship between these associations remains unclear. Corticotropin releasing factor-1 receptor (CRF1) has been implicated in alcohol (EtOH) use, affective states, and pain sensitivity; often in a sex-dependent manner. Using CRF1-cre transgenic rats, we found no sex differences in basal affective behavior with the exception of mechanical sensitivity, where females were more sensitive to mechanical stimuli. Following baseline testing, rats began EtOH (or water) drinking under intermittent access conditions. Females consumed more alcohol in the first week, but overall EtOH intake was not significantly different between males and females. Following 3-4 weeks of drinking, rats were tested again for negative affect. EtOH drinking decreased mechanical sensitivity, but no other group effects were observed. However, individual EtOH intake was directly correlated with anxiety- and depressive-like behavior in both sexes. Interestingly, EtOH intake inversely correlated with thermal sensitivity in males only. There were no group differences in CRF1+ neuronal activity in either prelimbic or infralimbic cortices, but final session EtOH intake was significantly correlated with activity in CRF1+ neurons in the infralimbic cortex. Together, our results suggest complex interplay between affective state, EtOH drinking, and the role of prefrontal cortex CRF1-containing neurons in mediating these behaviors. Additionally, these results highlight the importance of examining individual differences in AUD-related behaviors.

Keywords: alcohol, anxiety, pain, corticotropin releasing factor receptor 1, prefrontal cortex, prelimbic, infralimbic

SIGNIFICANCE STATEMENT

Despite alcohol use disorders being extremely comorbid with mood and pain disorders, there is still a limited understanding of the interaction and directionality between the them. To investigate this problem, rats were tested for affective behavior before and after being allowed to drink alcohol for 6 weeks. While baseline behavior did not predict subsequent intake, alcohol intake predicted both anxiety- and depressive-like behavior. These findings were accompanied by increased activity of the corticotropin releasing factor 1 containing neurons in the infralimbic region of the prefrontal cortex. Together, these findings reveal a new mechanism for understanding alcohol use.

1. INTRODUCTION

In 2021, 28.6 million adult Americans met the criteria for having an alcohol use disorder (AUD)[1]. AUDs are generally characterized by excessive drinking and negative affective states upon drinking cessation[2]. These states can promote dependence as individuals drink to alleviate withdrawal symptoms such as increased anxiety and anhedonia[3, 4]. Conversely, anxiety also often precedes alcohol use and is linked to a more rapid progression of AUD in both humans[5-8] and rodents[9-11]. AUDs and mood disorders are extremely co-morbid [1, 12-14]; however, our understanding of this relationship remains incomplete as both are influenced by a variety of other factors including sex. For example, genetic models demonstrated that rats bred for high anxiety-phenotypes or congenital learned helplessness drink more alcohol (EtOH) than rats bred for low-anxiety or non-congenital learned helplessness phenotypes; but both differences are only present in females[15, 16]. Other studies found that anxiety-like behavior in the elevated plus maze predicts EtOH intake in male, but not female, rats[17]. In humans, AUD is more prevalent among men[18]; however, recent trends on EtOH consumption indicate that the prevalence of AUD among women is increasing at a faster rate than in men[19] and the sex-related differences are narrowing[20]. Additionally, mood disorders are more common in women[21, 22] and women are more likely to develop comorbid anxiety and AUD[23]. Together, these studies highlight a major gap in our understanding of AUDs: how sex influences affective behaviors that both precede and follow AUD development.

Affective behavior and AUDs are not only co-morbid in incidence, but also share similar neurobiological underpinnings. More specifically, both are mediated by the stress-related corticotropin releasing factor (CRF) system. While hypothalamic release of CRF results in activation of the hypothalamic-pituitary-adrenal axis, extrahypothalamic CRF acts as an important neuromodulator in the brain through activity at two G-protein coupled receptors, CRF1 and CRF2. In humans, polymorphisms in the gene encoding CRF1 are associated with AUD and AUD-related behaviors such as binge drinking, EtOH intake, and intoxication[24, 25]. Polymorphisms in CRF1 are also associated with affective behaviors such as post-traumatic stress disorder, panic disorder, depression, suicidality, and loneliness[26-32]. In rodents, systemically-administered CRF1 antagonists decrease anxiety-like behavior, EtOH intake, EtOH-induced negative affect and EtOH-induced hyperalgesia[33-39]. CRF1 is found extensively throughout the brain and has been studied in the context of both AUD and affective behaviors. However, most studies focus only on the classical stress regions, such the bed nucleus of the stria terminalis[40-42] and the amygdala[41, 43-45] despite CRF1 being abundantly expressed in cortical regions[46-48]. Indeed, alcohol preferring rats have increased CRF1 levels in the cingulate cortex, motor cortex, and somatosensory cortex[49]. CRF1 is also abundantly expressed in the medial prefrontal cortex,[50, 51] but there are limited studies examining the effects of chronic EtOH drinking on prefrontal CRF1-containing neurons.

The present study uses behavioral testing and voluntary EtOH drinking in a CRF1 transgenic rat line to investigate the interaction between EtOH drinking and affective behavior in male and female rats. This study also examines the impact of chronic EtOH drinking on activity in the prefrontal cortex overall and specific to the CRF1 population.

EtOH Alters Affect & PFC Activity in CRF1-cre rats

2. METHODS

2.1 Subjects

32 adult male and female (n=16/sex) CRF1:cre: [td]Tomato rats[52] (8-11 weeks old, bred in-house) were housed in temperature and humidity-controlled rooms with 12h light/dark cycle (7:00 am lights ON). Rats had *ad libitum* access to food and water unless otherwise stated. Rats were handled before behavioral testing intake began. All animal procedures were performed in accordance with the [Author University] animal care committee's regulations.

2.2 Splash Test:

Rats were allowed to habituate in the behavioral testing room for 1h prior to testing. Rats were sprayed with 10% w/v sucrose solution, placed into a plexiglass/textured flooring behavioral testing chamber [50cm(*l*)×50cm(*w*)×38cm(*h*)] and recorded for 10min in red light conditions. Latency to groom and total time spent grooming were scored by an experimenter blind to experimental group using the Behavioral Observation Research Interactive Software (BORIS) program[53]. Latency to groom was used as a measure of anxiety-like behavior, whereas time spent grooming was used as a measure of depressive-like behavior. One subject was excluded from analysis due to equipment malfunction during testing.

2.3 Mechanical Sensitivity Testing (Von Frey):

Rats were brought into behavioral testing room 1h prior to testing and were habituated for 15min to a stainless-steel table [(78cm(*l*)×32cm(*w*)×19.5cm(*h*)] with a perforated sheet containing staggered holes (0.28cm diameter). Plastic filaments of increasing force (Bioseb EB2-VFF) were applied perpendicularly to hind paws until bending, and nocifensive responses defined as shaking, licking, or withdrawing the paw were recorded. Each paw was prodded 3 times/filament, beginning with the 2g filament and increasing until the rat exhibited a nocifensive response during at least 2 trials. Withdrawal thresholds for each hind paw were then averaged for each animal to determine the paw withdrawal threshold.

2.4 Novelty Suppressed Feeding (NSF):

Rats were given 1 Froot Loop in the homecage 48h prior to testing to prevent neophobia. 24h prior to testing food was removed from the cage. On test day, rats were habituated to behavioral testing room for 1h prior to testing. During testing, rats were individually placed into a brightly-lit (150 lux) behavioral testing chamber [50cm(*l*)×76cm(*w*)×40cm(*h*)] containing one Froot Loop on filter paper in the center. Latency to eat was recorded and used as a measure of anxiety-like behavior. After feeding initiation (or after 10min), rats were removed from the chamber and returned to the homecage where post-test consumption of pre-weighed Froot loops was recorded for 10min. Post-test consumption was used as a measure of motivated behavior.

2.5 Thermal Sensitivity Testing (Hargreaves Test)

Rats were assessed for thermal sensitivity using the Plantar Analgesia Meter (IITC). Rats were brought into the testing room 1h prior to testing and habituated to the heated glass (temperature: 32°C) apparatus for 20min. Infrared light (artificial intensity: 40) was

focused onto each hindpaw and withdrawal latency was recorded. A maximum cutoff of 20s was used to prevent tissue damage. Each hindpaw was tested twice. If the latencies differed by more than 1s, then the paw was tested one additional time. Withdrawal latencies were averaged per paw then per animal.

2.6 Intermittent Access to Two Bottle Choice (IA2BC)
After 4d acclimatization to drinking out of two water bottles, one bottle containing 20% EtOH and one bottle of water were placed on cages for 24 hr on alternating days (MWF). Water drinking rats had their water bottles changed every 24h to ensure comparable treatment. Bottle placement was alternated to eliminate side preference. 20% v/v EtOH was prepared by diluting 95% EtOH (Pharmco Products Inc) in tap water.

2.7 Immunohistochemistry
24h after the last alcohol presentation, rats were anesthetized with isoflurane and transcardially perfused using phosphate-buffered saline (PBS) followed by 4% paraformaldehyde. Brains were post-fixed in 4% PFA at 4°C overnight and then transferred to 30% sucrose in PBS at 4°C until brains sank. Brains were serially sectioned at 40 μm and stored in cryoprotectant (30% sucrose, 30% ethylene glycol, 0.01% sodium azide in PBS) at 4°C.

For each animal, 4-6 sections containing the medial prefrontal cortex (ranging from 3.72 to 2.52mm anterior of bregma based on Paxinos & Watson[54]) were used. These 40 μm sections underwent PBS washes and were incubated in 50% methanol for 30 minutes. Slices were blocked in 3% hydrogen peroxide for 10min then 3% normal goat serum, 1% bovine serum albumin, 0.3% Triton X in PBS for 1h (with PBS washes in-between). Slices were incubated at 4°C for 48h with rabbit anti-cFos primary antibody (1:3000, Millipore Sigma ABE457) and mouse anti-RFP primary antibody (1:500, Invitrogen MA5-15257) in blocking solution. Slices were then washed with 0.1% Tween-20 in Tris-buffered saline (TNT) before 30min incubation in TNB blocking buffer (Perkin-Elmer FP1012). Next, slices were incubated with goat anti-rabbit horseradish peroxidase (1:400, Abcam ab6721) and goat anti-mouse 555 (1:800, Invitrogen A2123) in TNB buffer for 2h before TNT washes. Lastly, slices were incubated in tyramide conjugated fluorescein (Akoya Biosciences, NEL741001KT, 1:50) for 10 min. Following TNT washes, slices were mounted and sealed with mounting medium (Vector Laboratories H1500, Burlingame, CA). Slides were stored at 4°C until imaging with the Keyence BZ-X800 fluorescence microscope.

Imaging and quantification were performed by an experimenter blind to the condition and sex of each animal. Images were taken at 20X and stitched together using the BZ-X800 Analyzer program. One hemisphere was selected at random for quantification, where the area of interest was outlined and counted manually using the FIJI multi-point counter tool[55]. Cell counts were averaged per animal.

2.8 Statistical Analysis
All statistical analysis was performed using Graphpad Prism 8.0 (San Diego, CA). In all studies, the threshold for significance was set to $p < 0.05$. Data was assessed for

EtOH Alters Affect & PFC Activity in CRF1-cre rats

normality using the Shapiro–Wilk test or D'Agostino & Pearson. For data that did not follow a normal distribution, nonparametric tests were used as described below. Detailed statistics for each experiment can be found in **Table 1**, and are referenced by superscripts in the results section.

Baseline behavioral tests were analyzed using an unpaired Student's t test (parametric data with equal sample variances), Welch's t test (parametric data with unequal sample variances), or a Mann-Whitney U test (nonparametric data). For cumulative incidence graphs, Kaplan-Meier survival curves were generated and were followed by Mantel-Cox log-rank test (followed by Bonferonni when applicable). Fluid intake and preference were analyzed by a mixed model 2-way ANOVA, using session as a within-subjects factor and sex as a between-subjects factor. Cumulative intake data were analyzed using an unpaired Student's t test (equal sample variances) or an unpaired Welch's t test (unequal sample variances). When behavioral tests were repeated (following drinking), data was analyzed using a 3-way mixed ANOVA (using time as a within-subjects factor while EtOH and sex were analyzed as between-subjects factors). When applicable, Bonferonni's correction was used for multiple comparisons. Correlational analyses were performed using Pearson's correlation (parametric data) or Spearman's correlation (nonparametric data). To compare correlations between males and females, Fisher's r to z transformation was used:

$$\text{Fisher's Z: } z_r = 0.5\ln((1+r)/(1-r)) \text{ where r = Pearson's r (or Spearman's } r_s)$$

Z_r from males and females (denoted z_m and z_f, respectively) was used to compute an observed z value, z_{obs} with the following calculation:

$$z_{obs} = \frac{z_m - z_f}{\sqrt{1/(N_m-3)+1/(N_f-3)}} \quad \text{where N = number of males (} N_m \text{) or females (} N_f \text{).}$$

In each set of correlations, the same analysis (Pearson's or Spearman's) was used for both sexes to calculate a z_{obs}.

Immunofluorescence data was analyzed by a 2-way ANOVA using sex and EtOH as between-subjects factors.

EtOH Alters Affect & PFC Activity in CRF1-cre rats

3. RESULTS

3.1 Experimental Timeline

As shown in **Fig. 1A**, rats were single-housed for 1 week prior to the onset of behavioral tests, which were conducted on separate but consecutive days. After baseline testing was complete, rats were left undisturbed for 3 days before being split into 2 groups, one allowed access to a bottle containing water and a bottle containing 20% EtOH under an intermittent access paradigm, and the other allowed access to only water bottles. After 3 weeks of drinking, post-drinking behavioral testing begun. These tests were separated by at least two drinking sessions in order to ensure testing did not disrupt EtOH consumption. During each test, rats were run in a pseudo-randomized subject order to minimize effects of circadian cycle and withdrawal time. After the last behavioral test, rats underwent a final EtOH drinking session. 24h after the final drinking session, rats were perfused and tissue was collected as described above.

3.2 Baseline Affective Behavior

Rats underwent a series of baseline behavioral tests to probe potential sex differences in behavior prior to EtOH drinking. In particular, we found sex differences in the Von Frey test of mechanosensitivity where female rats exhibited decreased withdrawal thresholds[a], suggesting increased mechanical sensitivity (**Fig. 1B**). However, these differences were unique to mechanosensation, as there were no sex differences in the Hargreaves test of thermosensitivity[b] (**Fig. 1C). ** Additionally, there were no sex differences in the Splash test parameters including latency to groom[c] **(Fig. 1D)** or time spent grooming **(Fig. 1E)**[d]. Some animals never initiated grooming and were thus assigned artificial latency values of 600s; however, cumulative incidence of grooming initiation curves **(Extended Data 1-1)** revealed no differences between sexes. There were also no sex differences in latency to eat[e] (**Fig. 1F**) or post-test home-cage chow consumption[f] **(Fig. 1G)** in the NSF test. Similarly, there were no differences in cumulative initiation of feeding **(Extended Data 1-2)**. Together, these results indicate that while female CRF1-cre rats display increased basal mechanosensitvity, there are no sex differences in basal anxiety-like or depressive-like behavior.

EtOH Alters Affect & PFC Activity in CRF1-cre rats

Figure 1.
Experimental timeline and baseline behavior. A) Timeline of entire experiment. **B**) Mechanical sensitivity in Von Frey test, as measured by paw withdrawal threshold. **C**) Thermal sensitivity in the Hargreaves test, as measured by latency to withdraw. **D)** Anxiety-like behavior in the splash test, as measured by latency to groom. **E)** Depressive-like behavior in the splash test, as measured by time spent grooming. **F)** Anxiety-like behavior in the Novelty Suppressed Feeding test, as measured by latency to eat. G) Motivation to eat in the Novelty Suppressed Feeding test, as measured by post-test consumption

Extended Figure 1-1
Cumulative Incidence of Grooming Initiation in the Splash Test. Even when accounting for the rats that never groomed (rather than artificially assigning them a maximum value or excluding them from studies), there was no sex difference in grooming initiation (Mantel-Cox log-rank test: $\chi^2=1.430$, df=1, $p=0.2318$). There was also no sex difference in total time spent grooming (**C;** Mann-Whitney test, $p=0.8901$).

Extended Figure 1-2
Cumulative Incidence of Feeding Initiation in the Novelty Suppressed Feeding Test. There was no effect of sex on cumulative occurrence of feeding initiation (**B**; Mantel-Cox log-rank test, $\chi^2=0.3366$, df=1, $p=0.5618$).

3.3 Voluntary EtOH Drinking

After assessing baseline affective behavior, rats were divided into two groups. The rats in the EtOH group were presented with 20% EtOH and water under an intermittent access two-bottle-choice procedure, whereas the rats in the water group received two water bottles. While there were no significant sex differences in overall EtOH intake[g] (**Fig. 2A**); females consumed significantly more EtOH in the first week of drinking[h] (**Fig. 2B**). Interestingly, there were no sex differences in water intake overall[i] (**Fig. 2C**) or during the first week[j] (**Fig 2D**), but water intake decreased over time in both sexes. We also looked at EtOH preference, calculated as volume of 20% EtOH consumed divided by volume of total fluid consumed during the session x 100%. While EtOH preference increased over time in a non-sex-dependent manner[k] (**Fig. 2E**), females exhibited a stronger EtOH preference during the first week[l] (**Fig. 2F**). Additionally, females consumed significantly more total fluid compared to males overall[m] (**Fig 2G**) and during the first week[n] (**Fig. 2H**). There were no significant correlations between any of the affective behaviors tested and initial EtOH intake (**Extended Figure 2-1**).

Figure 2.
Voluntary EtOH drinking under an intermittent access paradigm. A) EtOH intake over time. **B**) Cumulative EtOH intake during the first week of EtOH access. **C**) Water intake over time. **D**) Cumulative water intake during the first week of EtOH access. **E**) EtOH preference over time. **F**) Cumulative EtOH preference during the first week of EtOH access. **G**) Total fluid intake over time. **H**) Cumulative total fluid intake during the first week of EtOH access.

** $p<0.01$, * $p<0.01$ (unpaired Student's t test); & $p<0.05$ (2-way ANOVA, Main Effect of Sex); @ $p<0.05$ (unpaired Welch's t test)

EtOH Alters Affect & PFC Activity in CRF1-cre rats

Extended Figure 2-1
Behavioral measures did not correlate with first day EtOH intake in both males (M) and females (F). There was a trend where the rats with lower withdrawal thresholds drank more EtOH (**A**). Interestingly, there was no association between initial EtOH intake and basal thermal withdrawal (**B**), or basal latency to groom (**C**) and time spent grooming (**D**) in the splash test. There was also no significant relationship between initial EtOH intake and latency to eat (**E**) or post-test consumption (**F**) in the NSF test.

Pearson's Correlations:
A) M&F: r(14)= -0.4742, p=0.0635; Males: r(6)=0.5673, p=0.1425; Females: r(6)=-0.2699, p=0.5179
B) M&F r(14)= 0.1836, p=0.4962; Males: r(6)=0.3344, p=0.4162; Females: r(6)=0.1388 p=0.7431
F) M&F: r(14)=0.1751, p=0.5167; Males: r(6)=0.4654, p=0.2451; Females: r(6)=-0.01082, p=0.9797

Spearman's Correlations:
C) M&F: r_s(14)=0.3991, p=0.1266; Males: r_s(6)=0.03593, p=0.9434; Females: r_s(6)=0.4637, p=0.2619
D) M&F together: r_s(14)=-0.2088, p=0.4355, Males: r_s(6)=0.3234, p=0.4317; Females: r_s(6)=-0.4364, p=0.3036
E) M&F: r_s(14)=0.1177, p=0.6631; Males: r_s(6)=0.3571; p=0.3894; Females: r_s(6)=-0.2857, p=0.5008

3.4 EtOH Drinking & Associated Pain Sensitivity

After 3 weeks of EtOH access, the rats began post-drinking behavioral testing beginning with the Von Frey test of mechanosensitivity. There was a main effect of time, where paw withdrawal threshold decreased in all 4 groups[o] (**Fig. 3A**), suggesting that all rats became more sensitive to mechanical stimuli over time. A main effect of EtOH[o] also emerged, where EtOH-drinking rats had increased paw withdrawal thresholds, indicative of decreased mechanical sensitivity. Furthermore, there was an EtOH by time interaction, where Bonferonni posthoc analysis revealed that the decrease in paw withdrawal threshold over time was significantly stronger in water drinking rats compared to EtOH-drinking rats. Bonferroni posthoc analysis also indicated that the main effect of EtOH was driven predominantly by the post-test rather than pre-test, as there was no significant difference between water and EtOH-drinking rats at baseline. Finally, there was a main effect of sex[o], where females exhibited decreased paw withdrawal thresholds suggesting increased mechanical sensitivity (consistent with **Fig. 1B**). Interestingly, paw withdrawal threshold did not correlate with previous day[p] (**Fig. 3B**) or subsequent day EtOH intake[q] (**Fig. 3C**).

Figure 3.
Mechanical and thermal sensitivity following chronic EtOH drinking in males and females. The Von Frey test was repeated and mechanical sensitivity (**A**) was correlated with previous day EtOH intake(**B**) and subsequent EtOH intake (**C**). The Hargreaves test was also repeated and thermal sensitivity (**D**) was correlated with previous day EtOH intake (**E**) and subsequent EtOH intake (**F**).

$p<0.0001$ (3-way ANOVA, Main Effect of Time); @ $p<0.05$ (3-way ANOVA, Main effect of EtOH); $$ $p<0.01$ (Pearson's correlation); && $p<0.01$ (Fisher's r to z transformation).

Pearson's Correlations:
B) M&F: r(14)=0.3155, p=0.2340; Males: r(6)=0.04155, p=0.9222; Females: r(6)=0.6262, p=0.0967
C) M&F: r(14)=0.2660, p=0.3193; Males: r(6)=0.03441, p=0.9355; Females: r(6)=0.5708, p=0.1395
E) M&F: r(14)=0.2846, p=0.2853; Males: r(6)=0.9043, p=0.0020; Females: r(6)=-0.1378, p=0.7448
F) M&F: r(14)=-0.03709, p=0.8915; Males: r(6)=0.8494, p=0.0075; Females: r(6)=-0.6227, p=0.0991

EtOH Alters Affect & PFC Activity in CRF1-cre rats

Next, the Hargreaves test was repeated. In this test, latency to withdraw the paw is used as an index of thermal sensitivity, where increased withdrawal latency is indicative of decreased thermal sensitivity. There were no significant effects of time, EtOH, or sex on withdrawal latency[r] (**Fig. 3D**). Next, correlations examining the relationship between EtOH intake and withdrawal latency were performed. Previous day EtOH intake was directly correlated with withdrawal latency in males but not females[s] (**Fig. 3E**). This association was also seen when examining next day EtOH intake, where withdrawal latency predicted subsequent EtOH intake in males but not females[t] (**Fig. 3F**). The correlation of paw withdrawal threshold and EtOH intake in males as compared to females was statistically significant[s,t] (**Fig. 3E-F**). Together, these findings reveal that increased EtOH intake is bidirectionally associated with decreased thermal sensitivity in a sex-specific manner. Finally, there were no changes in 24h EtOH intake produced by behavioral testing (**Extended Figure 3-1**).

Extended Figure 3-1
Pain testing does not affect EtOH Intake. There was no effect of Von Frey testing (**A**) or Hargreaves testing (**B**) on EtOH consumption.

2-way ANOVAs:
A) Sex × Day: $F_{(1,14)}=0.1799$, $p=0.6779$; Sex: $F_{(1,14)}=0.01875$, $p=0.8930$; Day: $F_{(1,14)}=1.244$, $p=0.2835$
B) Sex × Day: $F_{(1,14)}=0.01586$, $p=0.9016$; Sex: $F_{(1,14)}=5.719e005$, $p=0.9941$; Day: $F_{(1,14)}=0.2869$, $p=0.6006$

There was no effect of either EtOH drinking or sex on latency to groom[u] (**Fig. 4A**) or total time spent grooming[v] (**Fig. 4B**). However, a main effect of time emerged on total time spent grooming, where all rats spent significantly less time grooming on the post-test. This suggests that regardless of sex or history of EtOH drinking, rats exhibited increased depressive-like behavior over time.

To account for animals that failed to groom during the 10 min test period, the cumulative incidence of grooming was plotted using the Kaplan-Meier survival curve followed by the Mantel-Cox log-rank test. There were no differences in cumulative incidence of grooming during the pre-test[w] (**Fig. 4C**). However, there was a significant effect during the post-test[x] (**Fig. 4D**), specifically where there was an increased cumulative incidence of grooming in EtOH-drinking males compared to EtOH-drinking females[x]. This suggests sex differences in anxiety-like behavior occur only in rats with a history of EtOH drinking (and not in EtOH-naïve rats).

To further investigate the association between EtOH intake and affective behavior, individual EtOH intake during the drinking session immediately before and after the Splash test were compared to latency to groom and total time spent grooming. Rats that consumed more EtOH prior to testing took significantly longer to begin grooming[y] (**Fig. 4E)**, an effect that was only seen when sexes were combined. This relationship was

EtOH Alters Affect & PFC Activity in CRF1-cre rats

unidirectional, as latency to groom did not predict subsequent drinking[z] (**Fig. 4F**). Total time spent grooming was also associated with EtOH intake prior to testing such that rats that consumed more EtOH groomed less[aa] (**Fig. 4G**); however, this effect was primarily driven by females[w]. Additionally, there was no association between total time spent grooming and subsequent EtOH intake[ab] (**Fig. 4H**) and the Splash test did not affect EtOH intake (**Extended Figure 4-1**). Together, these data suggest that previous day EtOH intake can predict both anxiety-like and depressive-like behavior in the Splash test.

Figure 4.
Anxiety- and depressive-like behavior in the splash test following chronic EtOH drinking in males and females. Latency to groom **(A)** and time spent grooming **(B)** were used as measured of anxiety- and depressive-like behavior. To account for animals that failed to groom within the 10min testing period, cumulative incidence curves for the pre-test and post-test are shown in **(C)** and **(D)**, respectively. The splash test was repeated and anxiety-like behavior was correlated with previous day EtOH intake (**E**) or subsequent EtOH intake (**F**). Depressive-like behavior in the splash test was correlated with previous day EtOH intake (**G**) or subsequent EtOH intake (**H**).

# $p<0.05$ (3-way ANOVA, Main Effect of Time); ^$p<0.05$, Mantel-Cox log-rank test followed by Bonferroni post-hoc, $ $p<0.05$ (Pearson's correlation); * $p<0.05$ (Spearman's correlation)

Pearson's Correlations:
E) M&F: r(14)=0.6108, p=0.0120; Males: r(6)=0.4120, p=0.3105; Females: r(6)=0.6172, p=0.1030
F) M&F: r(13)=0.2592, p=0.3508; Males: r(6)=0.4774, p=0.2269; Females: r(5)=0.1107, p=0.3132

Spearman's Correlations:
G) M&F: r_s(14)=-0.5859, p=0.0193; Males: r_s(6)=-0.8456, p=0.0179; Females: r_s(6)=0.07143, p=0.0179
H) M&F: r_s(13)=-0.02951, p=0.9180; Males: r_s(6)=0.09524, p=0.8401; Females: r_s(5)=-0.1782, p=0.7182

Extended Figure 4-1
Splash test testing does not affect EtOH Intake.
2-way ANOVA: Sex × Day: F(1,13)=1.958, p=0.1852; Sex: F(1,13)=1.883, p=0.1932; Day: F(1,13)=1.989, p=0.1819

EtOH Alters Affect & PFC Activity in CRF1-cre rats

The NSF test was also repeated to investigate changes in anxiety-like and motivated behavior following EtOH drinking. While there was no effect of sex or EtOH drinking on latency to eat[ac], there was an effect of time[ac] where all rats decreased their latency to eat on the second test (**Fig. 5A**). Post-test home-cage consumption was monitored for 10 minutes following the test and was used as a measure of motivation to eat. Compared to baseline, all the rats ate significantly less in their home-cage, suggesting a reduced motivation[ad] (**Fig. 5B**). There were no differences in cumulative incidence of feeding initiation during baseline or after chronic drinking (**Extended Figure 5-1)** suggesting that while rats decreased their anxiety-like behavior in the NSF, there were no group differences.

Figure 5.
Anxiety-like behavior in the Novelty Suppressed Feeding test following chronic EtOH drinking in males and females. Latency to eat **(A)** and post-test consumption **(B)** were used as measured of anxiety-like and motivated behavior. Anxiety-like behavior was correlated with previous day EtOH intake **(C)** or subsequent EtOH intake **(D)**. Motivated behavior was correlated with previous day EtOH intake **(F)** or subsequent EtOH intake **(G)**.

# $p<0.05$, ## $p<0.01$ (3-way ANOVA, Main Effect of Time); $ $p<0.05$ (Pearson's correlation); * $p<0.05$ (Spearman's correlation)

Spearman's Correlations:
C) M&F: $r_s(14)=0.5647$, $p=0.0248$; Males: $r_s(6)=0.7381$, $p=0.0458$; Females: $r_s(6)=0.4286$, $p=0.2992$
D) M&F: $r_s(14)=0.1912$, $p=0.4769$; Males: $r_s(6)=0.6667$, $p=0.0831$; Females: $r_s(6)=-0.1905$, $p=0.6646$

Pearson's Correlations:
E) M&F: $r(14)=0.4202$, $p=0.1051$; Males: $r(6)=0.1630$, $p=0.6997$; Females: $r(6)=-0.6833$ $p=0.0617$
F) M&F: $r(14)=0.1875$, $p=0.4867$; Males: $r(6)=-0.1263$, $p=0.7657$; Females: $r(6)=0.4296$ $p=0.2881$

Extended Figure 5-1
Cumulative occurrence of behavioral initiation in the NSF is not affected by EtOH drinking, nor is EtOH intake affected by NSF testing. There was no effect of group on cumulative occurrence of feeding initiation in the pre-test (**A**; Mantel-Cox log-rank test; $\chi^2=0.5295$, df=3, $p=0.9124$) or post-test (**B**; Mantel-Cox log-rank test; $\chi^2=3.717$, df=3, $p=0.2937$). Lastly, there was no effect of testing on EtOH intake (**C**; 2-way ANOVA; Sex × Day: $F(1,13)=0.5462$, $p=0.4721$; Sex: F(1,14)=1.901, $p=0.3442$; Day: F(1,14)=1.901, $p=0.1896$).

EtOH Alters Affect & PFC Activity in CRF1-cre rats

Associations between NSF behavioral measures and EtOH drinking were performed to examine potential individual effects of EtOH drinking on behavior. Rats that consumed more EtOH prior to the NSF test had increased latencies to eat,[ae] and this effect was driven primarily by males (**Fig. 5C**). There were no significant correlations between latency to eat and subsequent EtOH intake[af] (**Fig. 5D**). Post-test home-cage consumption did not correlate with previous day[ag] (**Fig. 5E**) or subsequent EtOH intake[ah] (**Fig. 5F**). Similar to the Splash test, there was no effect of testing on EtOH intake in either sex (**Extended Figure 5-1**). Together, these correlations suggest that EtOH intake is correlated with subsequent anxiety-like behavior on the NSF test.

3.7 EtOH drinking and Medial Prefrontal Cortex Activity
Following the NSF test, rats were allowed one final drinking session before brains were collected and the medial prefrontal cortex was stained for RFP (to identify CRF1+ cells) and cFos (a marker of neuronal activity). The medial prefrontal cortex was divided into prelimbic (PrL) and infralimbic (IL) cortices, based on the Paxinos and Watson atlas[54].

Prefrontal correlates of EtOH Drinking in males and females . Slices containing both the prelimbic cortex (PrL) and infralimbic cortex (IL) were stained for RFP (as a measure of CRF1+ neurons, shown in red) and cFos (as a measure of cellular activity, shown in green). PrL representative images **(A)** showing CRF1+ **(B)**, cFos+ **(C)** and total double labeled cells **(D)**. Double labeled cells were also expressed as % total CRF1+ cells **(E)**. IL representative images **(F)** showing CRF1+ **(G)**, cFos+ **(H)** and total double labeled cells **(I)**. Double labeled cells were also expressed as % total CRF1+ cells **(J)**. We also correlated EtOH intake on the last day with IL labeling of CRF1+ cells **(K)**, cFos+ cells **(L)**, total double labeled cells **(M)** and double labeled cells as %CRF1+ **(N)**.

Scale bars= 100μm; %% *p*<0.01 (2-way ANOVA; Main effect of Sex); $ *p*<0.05 (Pearson's correlation); && *p*<0.01 (Fisher's r to z transformation)

Pearson's Correlations:
K) M&F: r(14)=-0.07998, *p*=0.7684; Males: r(6)=-0.7277, *p*=0.0407; Females: r(6)=0.8307, *p*=0.0106
L) M&F: r(14)=-0.1695, *p*=0.5403; Males: r(6)=-0.7775, *p*=0.0231; Females: r(6)=0.2192, *p*=0.6022
M) M&F: r(14)=-0.1909, *p*=0.4787; Males: r(6)=-0.7752, *p*=0.0238; Females: r(6)=0.5508, *p*=0.1571
N) M&F: r(14)=-0.3485, *p*=0.1858; Males: r(6)=-0.5444, *p*=0.1630; Females: r(6)=-0.4090, *p*=0.3144

EtOH Alters Affect & PFC Activity in CRF1-cre rats

When investigating PrL (**Fig. 6A**), there were no group differences in CRF1+ cells[ai] (**Fig. 6B**), cFos+ cells[aj] (**Fig. 6C**), total double-labeled cells[ak] (**Fig. 6D**) or double-labeled cells as a percentage of total CRF1+ cells[al] (**Fig 6E**). Additionally, final session EtOH intake was not associated with any of these cell counts (**Extended Figure 6-1**). In the IL (**Fig. 6F**), there was no effect of sex or EtOH on CRF1+ cells[am] (**Fig. 6G**). However, females had increased cFos+ cells[an] (**Fig. 6H**). There were no significant differences in total double-labeled cells[ao] (**Fig. 6I**), but the sex difference seen in **Fig. 6H** persisted when double-labeled cells were expressed as %CRF1+ cells[ap] (**Fig. 6J**). The number of CRF1+ cells was directly correlated with EtOH intake in females[aq], but was inversely correlated with EtOH intake in males[aq] (**Fig. 6K**). Additionally, the number of cFos+ cells was inversely correlated with EtOH intake in males only[ar] (**Fig. 6L**). Consistent with the correlations seen thus far, total double-labeled cells was inversely correlated with EtOH intake in males only[as] (**Fig. 6M**), however this correlation disappeared when double labeled cells were normalized to %CRF1+ cells[at] (**Fig. 6N**). Taken together, these findings highlight differential effects of EtOH drinking on medial prefrontal cortex subdivisions, and suggest sex- and dose-dependent effects of EtOH on IL CRF1+ and cFos+ cells.

Extended Figure 6-1
Correlations between EtOH intake and activity in the prelimbic cortex. EtOH intake did not correlate with CRF1+ expression (**A**), cFos+ cells (**B**), total double-labeled cells (**C**) or double-labeled cells when expressed as %CRF1+ cells (**D**).

Spearman's Correlations
A) M&F: $r_s(14)=-0.1735$, $p=0.5193$; Males: $r_s(6)=-0.5476$, $p=0.1710$; Females: $r_s(6)=0.09524$, $p=0.8401$

Pearson's Correlations
B) M&F: $r(14)=-0.1997$, $p=0.4582$; Males: $r(6)=-0.25643$, $p=0.5271$; Females: $r(6)=-0.3325$, $p=0.4210$),
C) M&F: $r(14)=-0.07430$, $p=0.7842$; Males: $r(6)=-0.2966$, $p=0.4757$; Females: $r(6)=-0.02222$, $p=0.9583$
D) M&F: $r(14)=-0.01427$, $p=0.9582$; Males: $r(6)=-0.1604$, $p=0.7043$; Females: $r(6)=-0.09422$, $p=0.82$

EtOH Alters Affect & PFC Activity in CRF1-cre rats

4. DISCUSSION

The current study examined the dynamic relationship between affective states, pain sensitivity, and alcohol intake in male and female CRF1:cre: [td]Tomato rats. While female rats displayed increased basal sensitivity to mechanical stimuli, there were no other sex differences in baseline behavior. Females consumed significantly more EtOH during the first week of intermittent access, but overall EtOH intake was not significantly different between males and females. EtOH drinking decreased mechanical sensitivity in both male and female rats, but mechanical sensitivity did not correlate with previous or subsequent EtOH intake. Conversely, there were no group effects of EtOH on thermal sensitivity; but in males, decreased thermal sensitivity was associated with increased EtOH intake before and after the Hargreaves test. There were no group effects of EtOH on anxiety- or depressive-like behavior; however, both anxiety- and depressive-like behavior were directly correlated with EtOH intake prior to testing, but were not associated with subsequent EtOH intake. Additionally, all groups displayed increased mechanical sensitivity and depressive-like behavior as well as decreased motivated behavior compared to baseline levels. Finally, we investigated potential effects of EtOH intake on the activity of CRF1-containing neurons in the prelimbic and infralimbic prefrontal cortex. While there were no significant group effects of alcohol on overall or CRF1-specific activity, there were significant sex-specific associations between EtOH intake and activity in the infralimbic region. These findings provide new insight into individual differences in EtOH intake and its association with affective state, pain sensitivity, and neuronal activation of CRF1-containing neurons in key regions of the prefrontal cortex.

One important goal of this study was to examine affective behavior and EtOH drinking in male and female CRF1:cre: [td]Tomato rats. There was no effect of sex on basal affective behavior with the exception of mechanical sensitivity, where female rats displayed increase mechanical sensitivity. One previous study examined pain sensitivity in male and female CRF1:cre: [td]Tomato rats; however, data were not reported by sex[52]. Studies involving Wistar rats (the background strain for the CRF1:cre:[td]Tomato rats) also found no sex differences in baseline anxiety- and depressive-like behavior in the Splash Test[56, 57]. Sex differences in the NSF test in Wistar rats are less clear: one study found females exhibited increased anxiety-like behavior[58]; however, other studies report no sex differences[59-61]. Following baseline behavioral testing, rats were allowed to voluntarily drink EtOH under an intermittent access paradigm. In this study, male and female CRF1:cre: [td]Tomato rats drank 2-4g/kg EtOH with an approximate 15-30% EtOH preference, consistent with studies in Wistar rats[62-68]. While there were no sex differences in EtOH intake throughout the experiment, females drank significantly more EtOH during the first week. Studies comparing male and female EtOH intake under intermittent access conditions in Wistar rats are limited and inconclusive: some researchers found females drank more EtOH whereas others found no sex differences[67, 69]. Future studies will examine EtOH drinking in other contexts, such as binge-drinking or drinking following abstinence from EtOH. The studies presented here expand our knowledge of baseline behavior and EtOH intake in male and female CRF1:cre: [td]Tomato rats.

EtOH Alters Affect & PFC Activity in CRF1-cre rats

Although the behavioral tests used in these studies examine different aspects of affective behavior, one common theme emerged in the post-drinking tests: negative affect increased in both EtOH drinking animals as well as EtOH-naïve animals. Specifically, we found increased negative affect in both sexes in the Von Frey test (decreased withdrawal thresholds, suggesting increased mechanical sensitivity), Splash test (decreased time spent grooming, suggestive of decreased motivation), and NSF test (decreased post-test consumption, suggestive of decreased motivated behavior). Our study aimed to investigate the directionality of the relationship between negative affect and EtOH drinking, and therefore having baseline measures were crucial. However, affective tests are rarely repeated in the same subjects; thus, there are limited findings regarding the effects of time and re-testing on behavior. In one study, researchers found that male rats, but not female rats, spent significantly less time grooming on the second splash test[70]. While the current findings indicate decreased motivation regardless of sex, strain differences may account for this discrepancy. One potential explanation for the increased negative affect is increased stress due to prolonged social isolation. Indeed, prolonged social isolation during adulthood increases pain and negative affect in both humans and animals[71-78]. Together, these studies add to the literature by suggesting prolonged social isolation increases mechanical sensitivity and depressive-like behavior in both males and females regardless of EtOH exposure.

Many studies have reported increased negative affect following chronic EtOH drinkingt[36, 79-82]. However, there were no group effects of EtOH on thermal sensitivity or anxiety- and depressive-like behavior in the studies presented here. One potential explanation is that rats in the present study only consumed EtOH for 3-4 weeks before behavioral testing and thus did not consume enough EtOH to produce behavioral deficits. Studies in Wistar rats report escalated intake after ~20 EtOH drinking sessions[64, 83, 84]; rats in this study only drank for 16 sessions. Others have found that various aspects of EtOH withdrawal (including increased anxiety-like behavior, seizures, and EtOH intake during withdrawal) increase with repeated EtOH withdrawal cycles[85-90]. Another possibility is that the environment was too anxiogenic, creating a ceiling effect on anxiety-like behavior. Recent studies have developed methods to test anxiety-like behavior in the home-cage[91] to remove this potential confound and will be considered in the future.

Despite the lack of group effects, there were significant individual correlations between affective behavior and EtOH intake, thus highlighting the importance of examining individual differences in the context of EtOH-related behaviors. More specifically, this study found previous EtOH correlated with thermal sensitivity as well as anxiety- and depressive-like behavior. There was an inverse relationship between EtOH intake and thermal sensitivity, where males that consumed more EtOH were less sensitive to thermal stimuli and males with increased thermal sensitivity consumed less EtOH on the next day. These findings are in accordance with a previous study reporting that EtOH withdrawal increases nociceptive threshold[92]. However, it is important to note that other studies have demonstrated EtOH-induced decreases in pain threshold [93-97]. One potential explanation is that EtOH-induced peripheral neuropathy blunted the ability of

EtOH Alters Affect & PFC Activity in CRF1-cre rats

the rats to feel the thermal stimulus. This would be consistent with the group effect of EtOH on mechanical sensitivity, where the EtOH group had increased paw withdrawal thresholds, corresponding to decreased mechanical sensitivity. An alternative possibility is that CRF1:cre: [td]Tomato rats may be resistant to the analgesic properties of EtOH; however, additional studies are required to examine this in more detail. In addition to associations with thermal sensitivity, previous day EtOH intake was also associated with anxiety- and depressive-like behavior in the Splash test and NSF test. These findings are consistent with previous studies that demonstrate increased fear-potentiated startle in various genetic lines of alcohol preferring rats[98] and mice[99]. While all of these correlations held true when both sexes were analyzed together, analyzing the sexes separately revealed that the EtOH intake associations with anxiety-like behavior in the NSF were primarily driven by males whereas EtOH intake associations with depressive-like in the Splash test were primarily driven by females. These findings suggest test-specific contributions of sex in EtOH-induced affective behavior. Interestingly, anxiety-like and depressive-like behavior were unable to predict subsequent EtOH intake. Other studies have mixed findings regarding associations of anxiety-like behavior and subsequent drinking with significant correlations found in some studies[9-11] but not others[17, 100-102]. This work expands on these findings by demonstrating that neither baseline nor post-drinking anxiety- or depressive-like behavior predicts EtOH intake in male or female rats.

The effects of alcohol on affective behaviors are known to be mediated by key brain regions including the medial prefrontal cortex. Many studies fail to differentiate the prelimbic (PL) and infralimbic (IL) subdivisions of this region, but it is clear that these regions often play opposing roles in mediating behavior. For example, dopamine blockade in the PL reduces place conditioning without affecting cue conditioning but dopamine blockade in the IL reduces cue conditioning without affecting place conditioning[103]. While both PL and IL are active during cue-induced reinstatement of EtOH seeking, only ablation of activity-dependent IL neurons inhibited EtOH seeking[104]. Another study found EtOH drinking during adolescence increased Arc immunoreactivity in PL (but not in IL) whereas EtOH drinking during adulthood increased Arc immunoreactivity in IL (but not PL)[105]. While the lack of effect in PrL is consistent with the present findings, the effects in IL were divergent in that our study found decreased activity in the IL whereas their study found increased IL activity. In addition to species differences (mice vs rats), the timing of tissue collection may explain the differences seen. In the referenced study, brains were collected 25d after the last drinking session, but in the studies presented here, brains were collected 24h after the last drinking session. It is possible that increased length of forced abstinence results in compensatory upregulation of neuronal activity. Indeed, many studies have demonstrated opposing effects during short-term abstinence (~24h) and long-term abstinence (>7d)[106, 107].PL and IL also exert opposing effects on stress reactivity. For example, PL lesions result in augmented stress responses whereas IL lesions result in attenuated stress responses[108]. These are just a few studies that highlight the necessity to investigate PL and IL as separate regions, rather than combining them into one. In the studies presented here, there were no group differences or individual differences in the effects of chronic EtOH drinking on neuronal activation (cFos+ cells) or activation of

EtOH Alters Affect & PFC Activity in CRF1-cre rats

CRF1+ cells (double labeled cells) in the PL. However, individual differences emerged when examining the association between EtOH intake and cFos+ cells as well as EtOH intake and activated CRF1+ cells in the IL. Not only were these associations region-specific, but also sex-dependent as the correlations only occurred in male rats. There are limited studies investigating the effects of EtOH drinking in CRF1+ cells in the PL and IL subdivisions of the PFC. An important consideration is that most prior studies examined changes in CRF1 transcript levels; however, our studies examined the effects of EtOH on cells that contain CRF1 and can thus not be compared directly. One important study found that EtOH vapor withdrawal decreases excitability and excitatory neurotransmission in CRF1+ neurons in the prefrontal cortex[109]; however, this study was only performed in male mice and did not separate PL from IL. Collectively, these studies highlight the relevance of including both sexes and sub-regions while examining the effects of chronic EtOH drinking.

Together, our results illustrate the complex interplay between affective state, EtOH drinking, and the role of prefrontal cortex CRF1+ neurons in mediating these behaviors. Moreover, these data highlight the importance of examining individual differences in addition to group averages when investigating AUD and affective states. There are currently only 4 FDA-approved medications to treat AUD, and persistent issues with efficacy and compliance complicate effective treatment. Clinical trials have demonstrated differential efficacies of medications when stratified by factors such as sex, age of AUD onset, severity of AUD, and impulsivity levels[110, 111]. However, co-morbidity of AUD and affective disorders remains an underexamined issue. Thus, additional studies that consider individual differences in affective behavior and population-specific neurobiological changes in the context of AUD are warranted.

EtOH Alters Affect & PFC Activity in CRF1-cre rats

Letter	Figure	Distribution (Shapiro-Wilk test or D'Agostino & Pearson test (for repeated values))	Statistical Test	95% CI	Statistical Statement
a	1B; Baseline VF	Males: W=0.8693, *p*=0.025; Females: W=0.7876, *p*=0.0019)	Mann Whitney Test	[-3.0, -1.0]	U=15, *p*<0.0001
b	1C; Baseline Hargreaves	Males: W=0.9866, *p*=0.9952; Females: W=0.9324, *p*=0.2660	Unpaired Welch's t test	[-1.754, 3.054]	t(23.55)=0.5586, *p*=0.5817
c	1D; Baseline Splash Latency to Groom	Males: W=0.9147, *p*=0.16; Females: W=0.7800, *p*=0.0003	Mann Whitney Test	[-104.6, 204.9]	U=103, *p*=0.5140
d	1E; Baseline Splash Time Spent Grooming	Males: W=0.6525, *p*<0.0001; Females: W=0.8400, *p*=0.0097	Mann Whitney Test	[-13.39, 21.96]	U=116, *p*=0.8901
e	1F; Baseline NSF Latency to Eat	Males: W=0.8124, *p*=0.0040; Females: W=0.8299, *p*=0.0070	Mann Whitney Test	[-90.00, 107.0]	U=119, *p*=0.7520
f	1G; Baseline NSF Post-test	Males: W=0.9618, *p*=0.6937; Females: W=0.9651, *p*=0.7552	Student's t test	[-0.7823, 0.8027]	t(30)=0.02634
g	2A; EtOH Intake	Males: W=0.9716, *p*=0.8644; Females: W=0.9364, *p*=0.3074	2-way ANOVA	[-2.447, 0.7983]	Day ✕ Time: F(15,207)=1.352, *p*=0.1740; Sex: F(1,14)=1.187, *p*=0.2943; Time: F(15,207)=1.612, *p*=0.0726
h	2B; EtOH Intake 1 week	Males: W=0.9635, *p*=0.8424; `Females: W=0.9210, *p*=0.4830	Student's t test	[2.653, 11.42]	t(14)=3.442, *p*=0.0040

i	2C; H_2O Intake	Males: W=0.8956, p=0.0684; Females: W=0.9630, p=0.7156	2-way ANOVA	[-37.59, 8.638]	Sex × Time: F(15,205)=1.026; p=0.4302; Sex: F(1,14)=1.804, p=0.2006; Time: F(15,205)=2.882, p=0.0004
j	2D; H_2O Intake 1 week	Males: W=0.9289, p=0.5085; Females: W=0.8810, p=0.1924	Student's t test	[-22.85, 218.9]	t(14)=1.739, p=0.1039
k	2E; EtOH Preference	Males: W=0.9617, p=0.06923; Females: W=0.9002, p=0.0811	2-way ANOVA	[-11.23, 10.85]	Sex × Time: F(15,204)=1.344, p=0.1784; Sex: F(1,14)=0.001386, p=0.9708; Time: F(15,204)=3.382, p<0.0001
l	2F; EtOH Preference 1 week	Males: W=0.9040, p=0.3138; Females: W=0.8657; p=0.1368	Student's t test	[0.1420, 12.98]	t(14)=2.192, p=0.0458
m	2G; Total Fluid Intake	Males: W=0.9242 p=0.1970; Females: W=0.9192 p=0.1638	2 Way ANOVA	[-42.90, -1.558]	Sex × Time: F(15,206)=0.7308, p=0.7516; Sex: F(1,14)=5.320, p=0.0369; Time: F(15,206)=1.269, p=0.2240
n	2H; Total Fluid Intake 1 week	Males: W=0.9372, p=0.5836; Females: W=0.9236, p=0.4599	Welch's t test	[3.221, 282.2]	t(9.424)=2.299, p=0.0459
o	3A; Von Frey	H_2O Males pre-test: K^2=0.6402, p=0.7261; H_2O Females pre-test: K^2=0.6402, p=0.7261; EtOH Males pre-test: K^2=0.2973, p=0.8619; EtOH Females pre-test: K^2=1.227, p=0.5414; H_2O Males post-test: K2=9967, p=0.6075; H_2O Females post-test: K^2=2.182, p=0.3358; EtOH Males post-test: K^2=0.6506, p=0.7223; EtOH Females post-test: K^2=2.497, p=0.2870	3-way ANOVA	NA	Sex × Time × EtOH: F(1,28)=0.000, p>0.999; Time × EtOH: F(1,28)=5.628, p=0.0248; Sex × EtOH: F(1,28)=0.04308, p=0.8371; Sex × Time: F(1,28)=1.674, p=0.2062; Time: F(1,28)=39.12, p<0.0001, Sex: F(1,28)=4.308, p=0.0472; EtOH: F(1,28)=26.92; p<0.0001
p	3B; drinking before and Von Frey	EtOH intake (Males & Females): K^2=2.8323, p=0.2438; EtOH intake (Males): K^2=0.3602, p=0.8352; EtOH intake (Females): K^2=3.753, p=0.1531; Von Frey (Males & Females): K^2=1.085, p=0.5814; Von Frey (Males):	Pearson's Correlation	Males & Females: [-0.2136, 0.701] Males: [-0.6831, 0.7250]; Females: [-	Males & Females: r(14)=0.3155, p=0.2340; Males: r(6)=0.04155, p=0.9222; Females: r(6)=0.6262, p=0.0967

EtOH Alters Affect & PFC Activity in CRF1-cre rats

		K^2=2.182, p=0.3358; Von Frey (Females): K^2=2.497, p=0.2870		0.1405, 0.9234]	
q	3C; Von Frey and drinking after	Von Frey (Males & Females): K^2=1.085, p=0.5814; Von Frey (Males): K^2=2.182, p=0.3358; Von Frey (Females): K^2=2.497, p=0.2870; EtOH intake (Males & Females): K^2=1.487, p=0.4756; EtOH intake (Males): K^2=0.5575, p=0.7567; EtOH intake (Females): K^2=1.038, p=0.5951	Pearson's Correlation	Males & Females: [-0.2646, 0.6730]; Males:[-0.6869, 0.7216]; Females:-[0.2240, 0.9096]	Males & Females: r(14)=0.2660, p=0.3193, Males: r(6)=0.03441, p=0.9355; Females: r(6)=0.5708, p=0.1395
r	3D; Hargreaves	H_2O Males pre-test: K^2=0.5751, p=0.7501; H_2O Females pre-test: K^2=0.7428, p=0.6898; EtOH Males pre-test: K^2=0.07525, p=0.9631; EtOH Females pre-test: K^2=1.759, p=0.4149; H_2O Males post-test: K^2=1.255, p=0.0.5339; H_2O Females post-test: K^2=0.05513, p=0.09728; EtOH Males post-test: K^2=5.464, p=0.0651; EtOH Females post-test: K^2=5.464, p=0.0651	3-way ANOVA	NA	Sex × Time × EtOH: $F_{(1,28)}$=0.5697, p=0.4567; Time × EtOH: $F_{(1,28)}$=3.394, p=0.0760; Sex × EtOH: $F_{(1,28)}$=0.2055, p=0.6538; Sex × Time: $F_{(1,28)}$=0.2997, p=0.5884; EtOH: $F_{(1,28)}$=1.367; p=0.2521; Sex: $F_{(1,28)}$=3.529 p=0.0707; Time: $F_{(1,28)}$=1.224, p=0.2780
s	3E; drinking before and Hargreaves	EtOH Intake (Males & Females): K^2=2.074, p=0.3456; EtOH intake (Males): K^2=.288, p=0.5251; EtOH intake (Females): K^2=3.507, p=0.1732; Hargreaves (Males & Females): K^2=4.744, p=0.0933; Hargreaves (Males): K^2=5.464, p=0.0651; Hargreaves (Females): K^2=1.582, p=0.4534	Pearson's Correlation followed by Fisher's r to z transformation	Males & Females:[-0.2458, 0.6838]; Males: [0.5502, 0.9827]; Females: -[0.7679, 0.6278]	Pearson's: Males & Females: r(14)=0.2846, p=0.2853; Males: r(6)=0.9043, p=0.0020; Females: r(6)=-0.1378, p=0.7488 Fishers r to z: z_{obs}=2.5836, p=0.0098
t	3F; Hargreaves and drinking after	Hargreaves (Males & Females): K^2=4.744, p=0.0933; Hargreaves (Males): K^2=5.464, p=0.0651; Hargreaves (Females): K^2=1.582, p=0.4534; EtOH Intake (Males & Females): K^2=1.758, p=0.4152; EtOH Intake (Males): K^2=0.5217, p=0.7704, EtOH intake (Females): K^2=2.740, p=0.2541	Pearson's Correlation followed by Fisher's r to z transformation	Males & Females: [-0.5232, 0.4672]; Males: [0.3605, 0.9722]; Females: -[0.9226, 0.1460]	Pearson's: Males & Females together: r(14)=-0.03709, p=0.8915; Males: r(6)=0.8494, p=0.0075; Females: r(6)=-0.6227, p=0.0991 Fisher's r to z; z_{obs}=3.1360, p=0.0017

EtOH Alters Affect & PFC Activity in CRF1-cre rats

u	4A; Splash Test: Latency to Groom	H_2O Males pre-test: W=0.8928, p=0.3121; H_2O Females pre-test: K^2=2.030, p=0.3624; EtOH Males pre-test: K^2=1.950, p=0.3773; EtOH Females pre-test: K2=3.028, p=0.2200; H_2O Males post-test: W=0.9594, p=0.9138; H_2O Females post-test: K^2=4.792, p=0.0911; EtOH Males post-test: K^2=2.392, p=0.3025; EtOH Female post-test: K^2=3.433, p=01797	3-way ANOVA	NA	Sex × Time × EtOH: F(1,27)=0.03307, p=0.8570; Time × EtOH: F(1,27)=0.3165, p=0.5784; Sex × EtOH: F(1,27)=4.164, p=0.0512; Sex × Time: F(1,27)=2.776, p=0.1073; EtOH: F(1,27)=0.6438; p=0.5923; Sex: F(1,27)=3.782, p=0.0623; Time: F(1,27)=0.9851, p=0.3298
v	4B: Splash Test: Time Spent Grooming	H_2O Males pre-test: W=0.9455, p=0.5887; H_2O Females pre-test: K^2=1.531, p=0.4652; EtOH Males pre-test: K^2=13.46, p=0.0012; EtOH Females pre-test: K^2=4.000, p=0.153; H_2O Males post-test: W=0.8747, p=0.2038; H_2O females post-test: K^2=3.047, p=0.2179; EtOH-Males post-test: K^2=4.314, p=0.1156; EtOH Females post-test: K^2=2.577, p=0.2756	3-way ANOVA	NA	Sex × Time × EtOH: F(1,27)=2.150, p=0.1551; Time × EtOH: F(1,27)=0.1993, p=0.6589; Sex × EtOH: F(1,27)=1.272, p=0.2694; Sex × Time: F(1,27)=0.07479, p=0.7866; EtOH: F(1,27)=0.1483e-007; p=0.9997; Sex: F(1,27)=0.03729, p=0.8483; Time: F(1,27)=4.743, p=0.0383
w	4C: Cumulative Incidence (pre-test)	NA	Mantel-Cox log-rank	NA	χ^2=4.469, df=3, p=0.2151
x	4D: Cumulative Incidence (post-test)	NA	Mantel-Cox log-rank test followed by Bonferroni Correction	NA	χ^2=6.968, df=3, p=0.0467
y	4E: Drinking before & latency to groom	EtOH Intake (Males & Females): K^2=1.986, p=0.3704; EtOH intake (Males): K^2=0.7800, p=0.6671; EtOH intake (Females): K^2=0.7666, p=0.6816; Splash Latency (Males & Females): K^2=2.543, p=0.2804; Splash Latency (Males): K^2=2.392, p=0.3025; Splash Latency (Females): K^2=3.433, p=0.1797	Pearson's correlation	Males & Females: [0.1649, 0.8493]; Males: [-0.4124, 0.8654]; Females: [-0.1547, 0.9212]	Males & Females together: r(14)=0.6108, p=0.0120; Males: r(6)=0.4120, p=0.3105; Females: r(6)=0.6172, p=0.1030

EtOH Alters Affect & PFC Activity in CRF1-cre rats

z	4F: latency to groom & drinking after	Splash Latency (Males & Females): K^2=3.211, p=0.2008; Splash Latency (Males): K^2=2.392, p=0.3205; Splash Latency (Females): n too small; EtOH intake (Males & Females): K^2=2.408, p=0.3000; EtOH intake (Males): K^2=0.5217, p=0.7704; EtOH intake (Females): n too small	Pearson's correlation	Males & Females: [-0.2918, 0.6811]; Males: [-0.3425, 0.8845]; Females: [-0.7008, 0.7973]	Males & Females together: r(13)=0.2592, p=0.3508; Males: r(6)=0.4774, p=0.2269; Females: r(5)=0.1107, p=0.3132
aa	4G: drinking before & time spent grooming	EtOH Intake (Males & Females): K2=1.986, p=0.3704; EtOH intake (Males): K^2=0.7800, p=0.6671; EtOH intake (Females): K^2=0.7666, p=0.6816; Splash Grooming (Males & Females): K^2=8.434, p=0.0147; Splash Grooming (Males): K^2=4.314, p=0.1156; Splash Grooming (Females): K^2=2.577, p=0.2756	Spearman's Correlation	Males & Females: [-0.8429, -0.1112]; Males: n too small; Females: n too small	Males & Females: r_s(14)=-0.5859, p=0.0193; Males: r_s(6)=-0.8456, p=0.0179; Females: r_s(6)=0.07143, p=0.0179
ab	4H: time spent grooming & drinking after	Splash Grooming (Males & Females): K^2=15.70, p=0.0004, Splash Grooming (Males): K^2=4.314, p=0.1156; Splash Grooming (Females): n too small; EtOH intake (Males & Females): K^2=2.408, p=0.3000; EtOH intake (Males): K^2=0.5217, p=0.7704; EtOH intake (Females): n too small	Spearman's Correlation	Males & Females: [-0.5456, 0.5028]; Males: n too small; Females: n too small	Males & Females: r_s(13)=-0.02951, p=0.9180; Males: r_s(6)=0.09524, p=0.8401; Females: r_s(5)=-0.1782, p=0.7182
ac	5A: NSF Latency	**On log-transformed values:** H_2O Males pre-test: K^2=1.272, p=0.5293; H_2O Females pre-test: K^2=0.8888, p=0.6412; EtOH Males pre-test: K^2=1.481, p=0.4769; EtOH Females pre-test: K^2=4.336, p=0.1144; H_2O Males post-test: K^2=1.460, p=0.4819; H_2O Females post-test: K^2=1.083, p=0.5819; EtOH Males post-test: K^2=1.342, p=0.5112; EtOH Females post-test: K^2=2.836, p=0.2421	3-way ANOVA	NA	Sex × Time x EtOH: F(1,28)=0.09325, p=0.7623; Time × EtOH: F(1,28)=0.01534, p=0.9023; Sex × EtOH: F(1,28)=0.9187, p=0.3460; Sex × Time: F(1,28)=1.512, p=0.2291; EtOH: F(1,28)=0.3362; p=0.5667; Sex: F(1,28)=0.4367, p=0.5141; Time: F(1,28)=7.069, p=0.0128
ad	5B: NSF Post-test	H_2O Males pre-test: K^2=0.9742, p=0.6144; H_2O Females pre-test: K^2=0.7841, p=0.6757; EtOH Males	3-way ANOVA	NA	Sex × Time x EtOH: F(1,28)=0.2294, p=0.6357; Time × EtOH: F(1,28)=0.1.494, p=0.2317; Sex ×

EtOH Alters Affect & PFC Activity in CRF1-cre rats

		pre-test: K^2=0.2186, p=0.989; EtOH Females pre-test: K^2=0.09536, p=0.9534; H_2O Males post-test: K^2=2.543, p=0.2805; H_2O Females post-test: K^2=1.92, p=0.3786; EtOH Males post-test: K^2=0.175, p=0.9289; EtOH Females post-test: K^2=0.1529, p=0.9264			EtOH: F(1,28)=0.8227, p=0.3727; Sex × Time: F(1,28)=0.001506, p=0.9693; EtOH: F(1,28)=1.248; p=0.2734; Sex: F(1,28)=0.006310, p=0.9372; Time: F(1,28)=29.96 p<0.0001
ae	5C: drinking before & NSF Latency	EtOH intake (Males & Females): K^2=2.823, p=0.2438; EtOH intake (Males)=0.3602, p=0.8352; EtOH intake (Females): K^2=3.751, p=0.1531; NSF Latency (Males & Females): K^2=18.87, p<0.0001; NSF latency (Males): K^2=5.519, p=0.0633; NSF Latency (Females): K^2=21.53; p<0.0001	Spearman's Correlation	Males & Females: [0.07988, 0.8335]; Males: n too small; Females: n too small	Males & Females: r_s(14)=0.5647, p=0.0248; Males: r_s(6)=0.7381, p=0.0458; Females: r_s(6)=0.4286, p=0.2992
af	5D: NSF latency & drinking after	NSF Latency (Males & Females): K^2=18.87, p<0.0001; NSF latency (Males): K^2=5.519, p=0.0633; NSF Latency (Females): K^2=21.53; p<0.0001; EtOH intake (Males & Females): K^2=1.068, p=0.5862; EtOH intake (Males): K^2=1.074, p=0.5844; EtOH intake (Females): K^2=0.3717, p=0.8304	Spearman's Correlation	Males & Females: [-0.3506, 0.6371]; Males: n too small; Females: n too small	Males & Females: r_s(14)=0.1912, p=0.4769; Males: r_s(6)=0.6667, p=0.0831; Females: r_s(6)=-0.1905 p=0.6646
ag	5E: drinking before & NSF post-test	EtOH intake (Males & Females): K^2=2.823, p=0.2438; EtOH intake (Males)=0.3602, p=0.8352; EtOH intake (Females): K^2=3.751, p=0.1531; NSF post-test (Males & Females): K^2=0.4805, p=0.7864; NSF post-test (Males): K^2=0.1475, p=0.9289; NSF post-test (Females): K^2=14.75, p=0.2438	Pearson's Correlation	Males & Females: [-0.09538, 0.7580]; Males: [-0.6119, 0.7783]; Females: [-0.04114, 0.9369]	Males & Females: r(14)=0.4202, p=0.1051; Males: r(6)=0.1630, p=0.6997; Females: r(6)=-0.6833 p=0.0617
ah	5F: NSF post-test & drinking after	NSF post-test (Males & Females: K^2=0.4805, p=0.7864; NSF post-test (Males): K^2=0.1475, p=0.9289; NSF post-test (Females): K^2=14.75, p=0.2438; EtOH intake (Males &	Pearson's Correlation	Males & Females: [-0.3398, 0.6251]; Males: [-0.7631,	Males & Females: r(14)=0.1875, p=0.4867; Males: r(6)=-0.1263, p=0.7657; Females: r(6)=0.4296 p=0.2881

EtOH Alters Affect & PFC Activity in CRF1-cre rats

		Females): K^2=1.068, p=0.5862; EtOH intake (Males): K^2=1.074, p=0.5844; EtOH intake (Females): K^2=0.3717, p=0.8304		0.6349]; Females: [-0.3945, 0.8707]	
ai	6B: Prelimbic CRF1+ cells	H$_2$O Males: W=0.9387, p=0.5982; EtOH Males: W=0.8026, p=0.0305; H$_2$O Females: W=0.9307, p=0.5227; EtOH Females: W=0.9863; p=0.9871	2-way ANOVA	[-98.40, 66.86]	Sex × EtOH: F(1,28)=0.0005307, p=0.9818; EtOH: F(1,28)=0.002376, p=0.9615; Sex: F(1,28)=0.1528, p=0.6988
aj	6C: Prelimbic cFos+ cells	H$_2$O Males: W=0.9863, p=0.9871; EtOH Males: W=0.9250, p=0.4715; H$_2$O Females: W=0.8285, p=0.0573; EtOH Females: W=0.2578; p=0.1403	2-way ANOVA	[-413.1, 70.23]	Sex × EtOH: F(1,28)=0.0003888, p=0.9844; EtOH: F(1,28)=3.972, p=0.0561; Sex: F(1,28)=2.112 p=0.1573
ak	6D: Prelimbic total double labeled cells	H$_2$O Males: W=0.8668, p=0.1403; EtOH Males: W=0.9094, p=0.3501; H$_2$O Females: W=0.8506, p=0.0966; EtOH Females: W=0.9591; p=0.8018	2-way ANOVA	[-46.90, 5.043]	Sex × EtOH: F(1,28)=0.1194, p=0.7323; EtOH: F(1,28)=1.108, p=0.3015; Sex: F(1,28)=2.725 p=0.1100
al	6E: Prelimbic double labeled (%CRF1 cells)	H$_2$O Males: W=0.9421, p=0.6316; EtOH Males: W=0.9044, p=0.3165; H$_2$O Females: W=0.9501, p=0.7127; EtOH Females: W=0.9399; p=0.61	2-way ANOVA	[-11.99, 0.1321]	Sex × EtOH: F(1,28)=0.1477, p=0.7037; EtOH: F(1,28)=2.757, p=0.1080; Sex: F(1,28)=4.015 p=0.0549,
am	6G: Infralimbic CRF1+ cells	H$_2$O Males: W=0.9538, p=0.7494; EtOH Males: W=0.9358, p=0.57; H$_2$O Females: W=0.9501, p=0.7125; EtOH Females: W=0.9718, p=0.9121	2-way ANOVA	[-7.164, 19.40]	Sex × EtOH: F(1,28)=0.7886, p=0.3821; EtOH: F(1,28)=0.01444, p=0.9052; Sex: F(1,28)=0.8904, p=0.3534
an	6H: Infralimbic cFos+ cells	H$_2$O Males: W=0.9011, p=0.2954; EtOH Males: W=0.9654, p=0.8593; H$_2$O Females: W=0.9354, p=0.5661; EtOH Females: W=0.8903, p=0.2355	2-way ANOVA	[-104.3, -14.12]	Sex × EtOH: F(1,28)=0.000825, p=0.9773; EtOH: F(1,28)=1.639, p=0.2110; Sex: F(1,28)=7.233, p=0.0119
ao	6I: Infralimbic double labeled cells (total)	H$_2$O Males: W=0.9146, p=0.3874; EtOH Males: W=0.9709, p=0.9052; H$_2$O Females: W=0.9709, p=0.9052; EtOH Females: W=0.8742, p=0.1657	2-way ANOVA	[-3.729, 1.057]	Sex × EtOH: F(1,28)=2.961, p=0.0963; EtOH: F(1,28)=2.026, p=0.1657; Sex: F(1,28)=1.307, p=0.2626
ap	6J: Infralimbic double labeled (% CRF1+)	H$_2$O Males: W=0.9810, p=0.9679; EtOH Males: W=0.9193, p=0.4240; H$_2$O Females: W=0.9799, p=0.9625; EtOH Females: W=0.9043, p=0.3159	2-way ANOVA	[-2.276, 7.709]	Sex × EtOH: F(1,28)=0.7972, p=0.3795; EtOH: F(1,28)=1.242, p=0.2746; Sex: F(1,28)=7.356, p=0.0113
aq	6K: Infralimbic CRF1+ correlations	EtOH intake (Males & Females): W=0.9551, p=0.5745; EtOH Intake (Males): W=0.9177, p=0.4116; EtOH Intake (Females): W=0.9610, p=8197;	Pearson's correlation	Males & Females: [-0.5537, 0.4329]; Males:	Pearson's: Males & Females: r(14)=-0.07998, p=0.7684; Males: r(6)=-0.7277, p=0.0407; Females: r(6)=0.8307, p=0.0106;

EtOH Alters Affect & PFC Activity in CRF1-cre rats

	with EtOH intake	IHC (Males & Females): W=0.9490, p=0.4744; IHC (Males): W=0.9358, p=0.57; IHC (Females): W=0.9718, p=0.9121	followed by Fisher's r to z transformation	[-0.9468, -0.04716]; Females: [0.3041, 0.9685]	Fisher's r to z: z_{obs}=-3.3429, p=0.0008
ar	6L: infralimbic cFos+ correlations with EtOH intake	EtOH intake (Males & Females): W=0.9551, p=0.5745; EtOH Intake (Males): W=0.9177, p=0.4116; EtOH Intake (Females): W=0.9610, p=8197; IHC (Males & Females): W=0.9130, p=0.1310, IHC (Males): W=0.9654, p=0.8593; IHC (Females): W=0.9654, p=0.8593	Pearson's correlation followed by Fisher's r to z transformation	Males & Females: [-0.6137, 0.3561]; Males: [-0.9576, -0.1612]; Females: [-0.5743, 0.8002]	Pearson's: Males & Females: r(14)=-0.1695, p=0.5403; Males: r(6)=-0.7775, p=0.0231; Females: r(6)=0.2192, p=0.6022 Fisher's r to z: z_{obs}=-1.19950, p=0.0460
as	6M: infralimbic total double labeled correlations with EtOH intake	EtOH intake (Males & Females): W=0.9551, p=0.5745; EtOH Intake (Males): W=0.9177, p=0.4116; EtOH Intake (Females): W=0.9610, p=8197; IHC (Males & Females): W=0.9678, p=0.8015; IHC (Males): W=0.9311, p=0.5265; IHC (Females): W=0.8742, p=0.1657	Pearson's correlation followed by Fisher's r to z transformation	Males & Females: [-0.6273, 0.3366]; Males: [-0.9571, -0.1554]; Females: [-0.2515, 0.9044]	Pearson's: Males & Females: r(14)=-0.3485, p=0.1858; Males: r(6)=-0.5444, p=0.1630; Females: r(6)=-0.4090, p=0.3144 Fisher's r to z: z_{obs}=-2.6132, p=0.0090
at	6N: infralimbic double labeled (%CRF1+) correlations with EtOH intake	EtOH intake (Males & Females): W=0.9551, p=0.5745; EtOH Intake (Males): W=0.9177, p=0.4116; EtOH Intake (Females): W=0.9610, p=8197; IHC (Males & Females): W=0.8705, p=0.0277; IHC (Males): W=0.9193, p=0.4240; IHC (Females): W=0.9043, p=0.3159	Pearson's correlation	Males & Females: [-0.7199, 0.1779]; Males: [-0.9027, 0.2601]; Females: [-0.8645, 0.4154]	Males & Females: r(14)=-0.3485, p=0.1858; Males: r(6)=-0.5444, p=0.1630; Females: r(6)=-0.4090, p=0.3144

Table 1: Statistics Table

EtOH Alters Affect & PFC Activity in CRF1-cre rats

References

1. SAMHSA. *2021 National Survey on Drug Use and Health (NSDUH)*. 2022; Available from: https://www.samhsa.gov/data/report/2021-nsduh-annual-national-report.
2. Koob, G.F. and M. Le Moal, *Neurobiology of Addiction*. 2005, Academic Press.
3. Howe, L.K., et al., *Symptoms of anxiety, depression, and borderline personality in alcohol use disorder with and without comorbid substance use disorder.* Alcohol, 2021. **90**: p. 19-25.
4. Martinotti, G., et al., *Alcohol protracted withdrawal syndrome: the role of anhedonia.* Subst Use Misuse, 2008. **43**(3-4): p. 271-84.
5. Boschloo, L., et al., *Depressive and anxiety disorders predicting first incidence of alcohol use disorders: results of the Netherlands Study of Depression and Anxiety (NESDA).* J Clin Psychiatry, 2013. **74**(12): p. 1233-40.
6. Sloan, T.B., J.D. Roache, and B.A. Johnson, *The role of anxiety in predicting drinking behaviour.* Alcohol Alcohol, 2003. **38**(4): p. 360-3.
7. Kushner, M.G., et al., *Vulnerability to the rapid ("telescoped") development of alcohol dependence in individuals with anxiety disorder.* J Stud Alcohol Drugs, 2011. **72**(6): p. 1019-27.
8. Sartor, C.E., et al., *The role of childhood risk factors in initiation of alcohol use and progression to alcohol dependence.* Addiction, 2007. **102**(2): p. 216-25.
9. Pelloux, Y., J. Costentin, and D. Duterte-Boucher, *Differential involvement of anxiety and novelty preference levels on oral ethanol consumption in rats.* Psychopharmacology (Berl), 2015. **232**(15): p. 2711-21.
10. Spanagel, R., et al., *Anxiety: a potential predictor of vulnerability to the initiation of ethanol self-administration in rats.* Psychopharmacology (Berl), 1995. **122**(4): p. 369-73.
11. Hayton, S.J., M.K. Mahoney, and M.C. Olmstead, *Behavioral traits predicting alcohol drinking in outbred rats: an investigation of anxiety, novelty seeking, and cognitive flexibility.* Alcohol Clin Exp Res, 2012. **36**(4): p. 594-603.
12. Anker, J.J. and M.G. Kushner, *Co-Occurring Alcohol Use Disorder and Anxiety: Bridging Psychiatric, Psychological, and Neurobiological Perspectives.* Alcohol Res, 2019. **40**(1).
13. Schellekens, A.F., et al., *Co-morbid anxiety disorders predict early relapse after inpatient alcohol treatment.* Eur Psychiatry, 2015. **30**(1): p. 128-36.
14. McHugh, R.K. and R.D. Weiss, *Alcohol Use Disorder and Depressive Disorders.* Alcohol Res, 2019. **40**(1).
15. Izidio, G.S. and A. Ramos, *Positive association between ethanol consumption and anxiety-related behaviors in two selected rat lines.* Alcohol, 2007. **41**(7): p. 517-24.
16. Vengeliene, V., et al., *Voluntary alcohol intake in two rat lines selectively bred for learned helpless and non-helpless behavior.* Psychopharmacology (Berl), 2005. **178**(2-3): p. 125-32.
17. McNamara, T.A. and R. Ito, *Relationship between voluntary ethanol drinking and approach-avoidance biases in the face of motivational conflict: novel sex-dependent associations in rats.* Psychopharmacology (Berl), 2021. **238**(7): p. 1817-1832.
18. Grant, B.F., et al., *Epidemiology of DSM-5 Alcohol Use Disorder: Results From the National Epidemiologic Survey on Alcohol and Related Conditions III.* JAMA Psychiatry, 2015. **72**(8): p. 757-66.
19. Grant, B.F., et al., *Prevalence of 12-Month Alcohol Use, High-Risk Drinking, and DSM-IV Alcohol Use Disorder in the United States, 2001-2002 to 2012-2013: Results From the*

National Epidemiologic Survey on Alcohol and Related Conditions. JAMA Psychiatry, 2017. **74**(9): p. 911-923.

20. White, A., et al., *Converging Patterns of Alcohol Use and Related Outcomes Among Females and Males in the United States, 2002 to 2012.* Alcohol Clin Exp Res, 2015. **39**(9): p. 1712-26.

21. Salk, R.H., J.S. Hyde, and L.Y. Abramson, *Gender differences in depression in representative national samples: Meta-analyses of diagnoses and symptoms.* Psychol Bull, 2017. **143**(8): p. 783-822.

22. Franceschini, A. and L. Fattore, *Gender-specific approach in psychiatric diseases: Because sex matters.* Eur J Pharmacol, 2021. **896**: p. 173895.

23. Kessler, R.C., et al., *Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication.* Arch Gen Psychiatry, 2005. **62**(6): p. 593-602.

24. Chen, A.C., et al., *Single-nucleotide polymorphisms in corticotropin releasing hormone receptor 1 gene (CRHR1) are associated with quantitative trait of event-related potential and alcohol dependence.* Alcohol Clin Exp Res, 2010. **34**(6): p. 988-96.

25. Treutlein, J., et al., *Genetic association of the human corticotropin releasing hormone receptor 1 (CRHR1) with binge drinking and alcohol intake patterns in two independent samples.* Mol Psychiatry, 2006. **11**(6): p. 594-602.

26. Keck, M.E., et al., *Combined effects of exonic polymorphisms in CRHR1 and AVPR1B genes in a case/control study for panic disorder.* Am J Med Genet B Neuropsychiatr Genet, 2008. **147B**(7): p. 1196-204.

27. Weber, H., et al., *Allelic variation in CRHR1 predisposes to panic disorder: evidence for biased fear processing.* Mol Psychiatry, 2016. **21**(6): p. 813-22.

28. Kranzler, H.R., et al., *A CRHR1 haplotype moderates the effect of adverse childhood experiences on lifetime risk of major depressive episode in African-American women.* Am J Med Genet B Neuropsychiatr Genet, 2011. **156B**(8): p. 960-8.

29. Bradley, R.G., et al., *Influence of child abuse on adult depression: moderation by the corticotropin-releasing hormone receptor gene.* Arch Gen Psychiatry, 2008. **65**(2): p. 190-200.

30. Grabe, H.J., et al., *Childhood maltreatment, the corticotropin-releasing hormone receptor gene and adult depression in the general population.* Am J Med Genet B Neuropsychiatr Genet, 2010. **153B**(8): p. 1483-93.

31. Wasserman, D., et al., *Depression in suicidal males: genetic risk variants in the CRHR1 gene.* Genes Brain Behav, 2009. **8**(1): p. 72-9.

32. Chou, K.L., et al., *Influence of social environment on loneliness in older adults: Moderation by polymorphism in the CRHR1.* Am J Geriatr Psychiatry, 2014. **22**(5): p. 510-8.

33. Wang, H., et al., *Blockade of CRF1 and CCK2 receptors attenuated the elevated anxiety-like behavior induced by immobilization stress.* Pharmacol Biochem Behav, 2011. **98**(3): p. 362-8.

34. Sparta, D.R., et al., *Blockade of the corticotropin releasing factor type 1 receptor attenuates elevated ethanol drinking associated with drinking in the dark procedures.* Alcohol Clin Exp Res, 2008. **32**(2): p. 259-65.

35. Lowery, E.G., et al., *CRF-1 antagonist and CRF-2 agonist decrease binge-like ethanol drinking in C57BL/6J mice independent of the HPA axis.* Neuropsychopharmacology, 2010. **35**(6): p. 1241-52.

36. Overstreet, D.H., D.J. Knapp, and G.R. Breese, *Pharmacological modulation of repeated ethanol withdrawal-induced anxiety-like behavior differs in alcohol-preferring P and Sprague-Dawley rats.* Pharmacol Biochem Behav, 2005. **81**(1): p. 122-30.

EtOH Alters Affect & PFC Activity in CRF1-cre rats

37. Valdez, G.R., et al., *Increased ethanol self-administration and anxiety-like behavior during acute ethanol withdrawal and protracted abstinence: regulation by corticotropin-releasing factor.* Alcohol Clin Exp Res, 2002. **26**(10): p. 1494-501.

38. Breese, G.R., et al., *Prior multiple ethanol withdrawals enhance stress-induced anxiety-like behavior: inhibition by CRF1- and benzodiazepine-receptor antagonists and a 5-HT1a-receptor agonist.* Neuropsychopharmacology, 2005. **30**(9): p. 1662-9.

39. Edwards, S., et al., *Development of mechanical hypersensitivity in rats during heroin and ethanol dependence: alleviation by CRF(1) receptor antagonism.* Neuropharmacology, 2012. **62**(2): p. 1142-51.

40. Rinker, J.A., et al., *Extended Amygdala to Ventral Tegmental Area Corticotropin-Releasing Factor Circuit Controls Binge Ethanol Intake.* Biol Psychiatry, 2017. **81**(11): p. 930-940.

41. Huang, M.M., et al., *Corticotropin-releasing factor (CRF) sensitization of ethanol withdrawal-induced anxiety-like behavior is brain site specific and mediated by CRF-1 receptors: relation to stress-induced sensitization.* J Pharmacol Exp Ther, 2010. **332**(1): p. 298-307.

42. Faria, M.P., et al., *Anxiety-like responses induced by nitric oxide within the BNST in mice: Role of CRF1 and NMDA receptors.* Horm Behav, 2016. **79**: p. 74-83.

43. Kreifeldt, M., et al., *Central amygdala corticotropin-releasing factor neurons promote hyponeophagia but do not control alcohol drinking in mice.* Mol Psychiatry, 2022. **27**(5): p. 2502-2513.

44. Rodriguez, L., et al., *Alcohol Dependence Induces CRF Sensitivity in Female Central Amygdala GABA Synapses.* Int J Mol Sci, 2022. **23**(14).

45. Koob, G.F., *The role of CRF and CRF-related peptides in the dark side of addiction.* Brain Res, 2010. **1314**: p. 3-14.

46. Rieger, N.S., et al., *Insular cortex corticotropin-releasing factor integrates stress signaling with social affective behavior.* Neuropsychopharmacology, 2022. **47**(6): p. 1156-1168.

47. Gozzi, A., et al., *Differential effect of orexin-1 and CRF-1 antagonism on stress circuits: a fMRI study in the rat with the pharmacological stressor Yohimbine.* Neuropsychopharmacology, 2013. **38**(11): p. 2120-30.

48. Hubbard, C.S., et al., *Corticotropin-releasing factor receptor 1 antagonist alters regional activation and effective connectivity in an emotional-arousal circuit during expectation of abdominal pain.* J Neurosci, 2011. **31**(35): p. 12491-500.

49. Hansson, A.C., et al., *Variation at the rat Crhr1 locus and sensitivity to relapse into alcohol seeking induced by environmental stress.* Proc Natl Acad Sci U S A, 2006. **103**(41): p. 15236-41.

50. Weera, M.M., et al., *Generation and validation of a CRF1:Cre transgenic rat: The role of central amygdala CRF1 in nociception and anxiety-like behavior.* bioRxiv, 2021: p. 2021.02.23.432551.

51. Brufatto, J.P.T., et al., *Ventromedial prefrontal cortex CRF1 receptors modulate the tachycardic activity of baroreflex.* Pflugers Arch, 2021. **473**(4): p. 697-709.

52. Weera, M.M., et al., *Generation of a CRF1-Cre transgenic rat and the role of central amygdala CRF1 cells in nociception and anxiety-like behavior.* Elife, 2022. **11**.

53. Friard, O. and M. Gamba, *BORIS: a free, versatile open-source event-logging software for video/audio coding and live observations.* Methods in Ecology and Evolution, 2016. **7**(11): p. 1325-1330.

54. Paxinos, G. and C. Watson, *The Rat Brain in Stereotaxic Coordinates: Hard Cover Edition.* 2007: Elsevier Science.

55. Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nat Methods, 2012. **9**(7): p. 676-82.

EtOH Alters Affect & PFC Activity in CRF1-cre rats

56.   Abelaira, H.M., et al., *Combination of electroconvulsive stimulation with ketamine or escitalopram protects the brain against inflammation and oxidative stress induced by maternal deprivation and is critical for associated behaviors in male and female rats.* Mol Neurobiol, 2022. **59**(3): p. 1452-1475.

57.   Keledjian, K., et al., *The peroxisome proliferator-activated receptor gamma (PPARgamma) agonist, rosiglitazone, ameliorates neurofunctional and neuroinflammatory abnormalities in a rat model of Gulf War Illness.* PLoS One, 2020. **15**(11): p. e0242427.

58.   De Oliveira Sergio, T., et al., *Sex differences in specific aspects of two animal tests of anxiety-like behavior.* Psychopharmacology (Berl), 2021.

59.   Enayati, M., et al., *Prenatal maternal stress alters depression-related symptoms in a strain - and sex-dependent manner in rodent offspring.* Life Sci, 2020. **251**: p. 117597.

60.   Francis-Oliveira, J., et al., *Fluoxetine exposure during pregnancy and lactation: Effects on acute stress response and behavior in the novelty-suppressed feeding are age and gender-dependent in rats.* Behav Brain Res, 2013. **252**: p. 195-203.

61.   Cabbia, R., A. Consoli, and D. Suchecki, *Association of 24 h maternal deprivation with a saline injection in the neonatal period alters adult stress response and brain monoamines in a sex-dependent fashion.* Stress, 2018. **21**(4): p. 333-346.

62.   Scott, H., N. Tjernstrom, and E. Roman, *Effects of pair housing on voluntary alcohol intake in male and female Wistar rats.* Alcohol, 2020. **86**: p. 121-128.

63.   Momeni, S. and E. Roman, *Subgroup-dependent effects of voluntary alcohol intake on behavioral profiles in outbred Wistar rats.* Behav Brain Res, 2014. **275**: p. 288-96.

64.   Kimbrough, A., et al., *Intermittent Access to Ethanol Drinking Facilitates the Transition to Excessive Drinking After Chronic Intermittent Ethanol Vapor Exposure.* Alcohol Clin Exp Res, 2017. **41**(8): p. 1502-1509.

65.   Adermark, L., et al., *Intermittent ethanol consumption depresses endocannabinoid-signaling in the dorsolateral striatum of rat.* Neuropharmacology, 2011. **61**(7): p. 1160-5.

66.   Cippitelli, A., et al., *Pharmacological blockade of corticotropin-releasing hormone receptor 1 (CRH1R) reduces voluntary consumption of high alcohol concentrations in non-dependent Wistar rats.* Pharmacol Biochem Behav, 2012. **100**(3): p. 522-9.

67.   Albrechet-Souza, L., C.L. Schratz, and N.W. Gilpin, *Sex differences in traumatic stress reactivity in rats with and without a history of alcohol drinking.* Biol Sex Differ, 2020. **11**(1): p. 27.

68.   Moench, K.M. and M.L. Logrip, *Housing Condition Differentially Impacts Escalation of Alcohol Intake, Relapse-Like Drinking, Anxiety-Like Behavior, and Stress History Effects by Sex.* Alcohol Clin Exp Res, 2021. **45**(2): p. 480-489.

69.   Priddy, B.M., et al., *Sex, strain, and estrous cycle influences on alcohol drinking in rats.* Pharmacol Biochem Behav, 2017. **152**: p. 61-67.

70.   Quadir, S.G., et al., *Sex differences in affective states and association with voluntary ethanol intake in Sprague-Dawley rats.* Psychopharmacology (Berl), 2022. **239**(2): p. 589-604.

71.   Yamada, K., et al., *Loneliness, social isolation, and pain following the COVID-19 outbreak: data from a nationwide internet survey in Japan.* Sci Rep, 2021. **11**(1): p. 18643.

72.   Bannon, S., et al., *The role of social isolation in physical and emotional outcomes among patients with chronic pain.* Gen Hosp Psychiatry, 2021. **69**: p. 50-54.

73.   Baker, T., et al., *The Intersection of Pain Outcomes and Social Isolation Among African Americans.* J Racial Ethn Health Disparities, 2022. **9**(5): p. 1905-1911.

74.   Santini, Z.I., et al., *Social disconnectedness, perceived isolation, and symptoms of depression and anxiety among older Americans (NSHAP): a longitudinal mediation analysis.* Lancet Public Health, 2020. **5**(1): p. e62-e70.

EtOH Alters Affect & PFC Activity in CRF1-cre rats

75. Taheri Zadeh, Z., et al., *Depresssion, anxiety and other cognitive consequences of social isolation: Drug and non-drug treatments.* Int J Clin Pract, 2021. **75**(12): p. e14949.
76. Mazur, F.G., et al., *Effects of physical exercise and social isolation on anxiety-related behaviors in two inbred rat strains.* Behav Processes, 2017. **142**: p. 70-78.
77. Blass, E., E. Fitzgerald, and P. Kehoe, *Interactions between sucrose, pain and isolation distress.* Pharmacol Biochem Behav, 1987. **26**(3): p. 483-9.
78. Tuboly, G., G. Benedek, and G. Horvath, *Selective disturbance of pain sensitivity after social isolation.* Physiol Behav, 2009. **96**(1): p. 18-22.
79. Quadir, S.G., et al., *Antagonism of Sigma-1 receptor blocks heavy alcohol drinking and associated hyperalgesia in male mice.* Alcohol Clin Exp Res, 2021. **45**(7): p. 1398-1407.
80. Xu, H., et al., *Chronic Voluntary Alcohol Drinking Causes Anxiety-like Behavior, Thiamine Deficiency, and Brain Damage of Female Crossed High Alcohol Preferring Mice.* Front Pharmacol, 2021. **12**: p. 614396.
81. Gong, M.F., et al., *Attenuation of ethanol abstinence-induced anxiety- and depressive-like behavior by the phosphodiesterase-4 inhibitor rolipram in rodents.* Psychopharmacology (Berl), 2017. **234**(20): p. 3143-3151.
82. Van Skike, C.E., J.L. Diaz-Granados, and D.B. Matthews, *Chronic intermittent ethanol exposure produces persistent anxiety in adolescent and adult rats.* Alcohol Clin Exp Res, 2015. **39**(2): p. 262-71.
83. George, O., et al., *Recruitment of medial prefrontal cortex neurons during alcohol withdrawal predicts cognitive impairment and excessive alcohol drinking.* Proc Natl Acad Sci U S A, 2012. **109**(44): p. 18156-61.
84. Carnicella, S., R. Amamoto, and D. Ron, *Excessive alcohol consumption is blocked by glial cell line-derived neurotrophic factor.* Alcohol, 2009. **43**(1): p. 35-43.
85. Holter, S.M., et al., *Long-term ethanol self-administration with repeated ethanol deprivation episodes changes ethanol drinking pattern and increases anxiety-related behaviour during ethanol deprivation in rats.* Behav Pharmacol, 1998. **9**(1): p. 41-8.
86. Becker, H.C., *The alcohol withdrawal "kindling" phenomenon: clinical and experimental findings.* Alcohol Clin Exp Res, 1996. **20**(8 Suppl): p. 121A-124A.
87. Rimondini, R., W. Sommer, and M. Heilig, *A temporal threshold for induction of persistent alcohol preference: behavioral evidence in a rat model of intermittent intoxication.* J Stud Alcohol, 2003. **64**(4): p. 445-9.
88. Lopez, M.F. and H.C. Becker, *Effect of pattern and number of chronic ethanol exposures on subsequent voluntary ethanol intake in C57BL/6J mice.* Psychopharmacology (Berl), 2005. **181**(4): p. 688-96.
89. Overstreet, D.H., D.J. Knapp, and G.R. Breese, *Modulation of multiple ethanol withdrawal-induced anxiety-like behavior by CRF and CRF1 receptors.* Pharmacol Biochem Behav, 2004. **77**(2): p. 405-13.
90. Zhang, Z., et al., *Dose- and time-dependent expression of anxiety-like behavior in the elevated plus-maze during withdrawal from acute and repeated intermittent ethanol intoxication in rats.* Alcohol Clin Exp Res, 2007. **31**(11): p. 1811-9.
91. Neira, S., et al., *Chronic alcohol consumption alters home-cage behaviors and responses to ethologically relevant predator tasks in mice.* Alcohol Clin Exp Res, 2022. **46**(8): p. 1616-1629.
92. Schunck, R.V., et al., *Protracted alcohol abstinence induces analgesia in rats: Possible relationships with BDNF and interleukin-10.* Pharmacol Biochem Behav, 2015. **135**: p. 64-9.
93. Avegno, E.M., et al., *Central Amygdala Circuits Mediate Hyperalgesia in Alcohol-Dependent Rats.* J Neurosci, 2018. **38**(36): p. 7761-7773.
94. Walcott, A.T., et al., *Social transfer of alcohol withdrawal-induced hyperalgesia in female prairie voles.* Soc Neurosci, 2018. **13**(6): p. 710-717.

EtOH Alters Affect & PFC Activity in CRF1-cre rats

95.   Kang, S., et al., *Downregulation of M-channels in lateral habenula mediates hyperalgesia during alcohol withdrawal in rats.* Sci Rep, 2019. **9**(1): p. 2714.

96.   Kononoff, J., et al., *Systemic and Intra-Habenular Activation of the Orphan G Protein-Coupled Receptor GPR139 Decreases Compulsive-Like Alcohol Drinking and Hyperalgesia in Alcohol-Dependent Rats.* eNeuro, 2018. **5**(3).

97.   You, D.S., et al., *Hyperalgesia after a Drinking Episode in Young Adult Binge Drinkers: A Cross-Sectional Study.* Alcohol Alcohol, 2020. **55**(6): p. 608-615.

98.   McKinzie, D.L., et al., *Acoustic startle and fear-potentiated startle in alcohol-preferring (P) and -nonpreferring (NP) lines of rats.* Pharmacol Biochem Behav, 2000. **65**(4): p. 691-6.

99.   Barrenha, G.D. and J.A. Chester, *Genetic correlation between innate alcohol preference and fear-potentiated startle in selected mouse lines.* Alcohol Clin Exp Res, 2007. **31**(7): p. 1081-8.

100.   Correia, D., et al., *Trait anxiety and ethanol: anxiolysis in high-anxiety mice and no relation to intake behavior in an addiction model.* Prog Neuropsychopharmacol Biol Psychiatry, 2009. **33**(5): p. 880-8.

101.   Da Silva, G.E., A. Ramos, and R.N. Takahashi, *Comparison of voluntary ethanol intake by two pairs of rat lines used as genetic models of anxiety.* Braz J Med Biol Res, 2004. **37**(10): p. 1511-7.

102.   Henniger, M.S., et al., *Alcohol self-administration in two rat lines selectively bred for extremes in anxiety-related behavior.* Neuropsychopharmacology, 2002. **26**(6): p. 729-36.

103.   Hayen, A., et al., *Opposing roles of prelimbic and infralimbic dopamine in conditioned cue and place preference.* Psychopharmacology (Berl), 2014. **231**(12): p. 2483-92.

104.   Pfarr, S., et al., *Losing Control: Excessive Alcohol Seeking after Selective Inactivation of Cue-Responsive Neurons in the Infralimbic Cortex.* J Neurosci, 2015. **35**(30): p. 10750-61.

105.   Lawson, K., et al., *Adolescence alcohol exposure impairs fear extinction and alters medial prefrontal cortex plasticity.* Neuropharmacology, 2022. **211**: p. 109048.

106.   Smith, R.J., et al., *Dynamic c-Fos changes in mouse brain during acute and protracted withdrawal from chronic intermittent ethanol exposure and relapse drinking.* Addict Biol, 2020. **25**(6): p. e12804.

107.   Cannady, R., et al., *Distinct Region- and Time-Dependent Functional Cortical Adaptations in C57BL/6J Mice after Short and Prolonged Alcohol Drinking.* eNeuro, 2020. **7**(3).

108.   Radley, J.J., C.M. Arias, and P.E. Sawchenko, *Regional differentiation of the medial prefrontal cortex in regulating adaptive responses to acute emotional stress.* J Neurosci, 2006. **26**(50): p. 12967-76.

109.   Patel, R.R., et al., *Ethanol withdrawal-induced adaptations in prefrontal corticotropin releasing factor receptor 1-expressing neurons regulate anxiety and conditioned rewarding effects of ethanol.* Mol Psychiatry, 2022.

110.   Leggio, L., et al., *Medication Development for Alcohol Use Disorder: A Focus on Clinical Studies.* Handb Exp Pharmacol, 2020. **258**: p. 443-462.

111.   Zorrilla, E.P., et al., *Behavioral, biological, and chemical perspectives on targeting CRF(1) receptor antagonists to treat alcoholism.* Drug Alcohol Depend, 2013. **128**(3): p. 175-86.

Experimental Timeline

A

Splash Test | Von Frey | NSF | Hargreaves

single housed — baseline tests — 3d — 3 weeks IA — post-drinking tests — IA — 1d — brains collected

Von Frey | IA | IA | Hargreaves | IA | IA | Splash Test | IA | IA | NSF

Von Frey

B ****

Paw Withdrawal Threshold (g)

Males   Females

Hargreaves

C

Paw Withdrawal Latency (s)

Males   Females

Splash Test

D

Latency to Groom (s)

Males   Females

E

Time Spent Grooming (s)

Males   Females

Novelty Suppressed Feeding

F

Latency to Eat (s)

Males   Females

G

Post-test Consumption (g/kg)

Males   Females

Behavioral Responses

Correlations with EtOH Intake

Von Frey

A pre-test / post-test **###**

B M & F, Males, Females

C M & F, Males, Females

Hargreaves

D pre-test / post-test

E M & F, Males **$$**, Females **&&**

F M & F, Males **$$**, Females **&&**

Behavioral Responses

A

Latency to Groom (s)

☐ pre-test   ▨ post-test

| Male | Female | Male | Female |

H₂O Drinking   EtOH Drinking

B

Time Spent Grooming (s)

☐ pre-test   ▨ post-test #

| Male | Female | Male | Female |

H₂O Drinking   EtOH Drinking

Cumulative Incidence Curves

C

% of Animals Who Have Initiated Grooming

pre-test

H₂O-males
H₂O-females
EtOH-males
EtOH-females

0   200   400   600

Time (s)

D

% of Animals Who Have Initiated Grooming

post-test

H₂O-males
H₂O-females
EtOH-males
EtOH-females ^

0   200   400   600

Time (s)

Correlations with EtOH Intake

E

Latency to Groom (s)

0   5   10   15

Previous Day EtOH Intake (g/kg)

M & F $
Males
Females

F

Next Day EtOH Intake (g/kg)

0   200   400   600

Latency to Groom (s)

M & F
Males
Females

G

Time Spent Grooming (s)

0   5   10   15

Previous Day EtOH Intake (g/kg)

M & F ★
Males
Females ★

H

Next Day EtOH Intake (g/kg)

0   20   40   60

Time Spent Grooming (s)

M & F
Males
Females

Behavioral Responses

A

Latency to Eat (s)

pre-test / post-test #

H₂O Drinking | EtOH Drinking
Male | Female | Male | Female

B

Post-test Consumption (g/kg)

pre-test / post-test ##

H₂O Drinking | EtOH Drinking
Male | Female | Male | Female

Correlations with EtOH Intake

C

Latency to Eat (s)

··· M & F ★
▫ Males ★
▲ Females

Previous Day EtOH Intake (g/kg)

D

Next Day EtOH Intake (g/kg)

··· M & F
▫ Males
▲ Females

Latency to Eat (s)

E

Post-test Consumption (g/kg)

··· M & F
▫ Males
▲ Females

Previous Day EtOH Intake (g/kg)

F

Next Day EtOH Intake (g/kg)

··· M & F
▫ Males
▲ Females

Post-test consumption (g/kg)