Deciphering divergent trypanosomatid nuclear complexes by analyzing interactomic datasets with AlphaFold2 and genetic approaches

Rodriguez Araya, Elvio[1,2]; Cribb, Pamela[1,2]; Merli, Marcelo L.[1,2]; Serra, Esteban[1,2]

[1]Instituto de Biología Molecular y Celular de Rosario, CONICET, Suipacha 590, CP200 Rosario, Argentina.
[2]Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Suipacha 531, CP2000 Rosario, Argentina.

*Email: serra@ibr-conicet.gov.ar
*Email: rodriguezaraya@ibr-conicet.gov.ar

Abstract

Little is known about acetylation signaling pathways in early branching organisms such as trypanosomatids in comparison to other unicellular eucaryotic species like yeast. In addition to important biological differences described in this deep branched organisms, serious bioinformatic limitations arise when facing the highly divergent sequences of the proteins involved, making it difficult to perform homology-based prediction of protein functions. The availability of several interactomic datasets, like those recently generated by Staneva *et al* for *Trypanosoma brucei* acetyl-lysine readers, writers and erasers and the public release of AlphaFold2 (AF2) and AF2-multimer could be envisaged as a shortcut to address this problem. These tertiary and quaternary structure predictions could be used as a tool with high predictive value when inferring protein function. In this work, we made use of public interactomic datasets to predict direct interactions using AF2-multimer and validated them by yeast two-hybrid (Y2H) assays. We focused on MRG domain-containing proteins of *Trypanosoma cruzi* and their interactions, typically found as structural part of histone acetyl transferase (HAT) and histone deacetylase (HDAC) complexes. Our results led us to identify *Tc*TINTIN, a structurally conserved complex that is orthologous to the human and yeast TINTIN complexes, that cannot be identified by homology searches based only in the primary sequences. Our *in silico* and wet lab approach led us to determine a specific sequential assembly of this complex, involving the proteins *Tc*MRGx (MORF4 Related Gene X), *Tc*MRGBP (MRG Binding Protein) and *Tc*BDF6 (Bromodomain Factor 6). We also found another MRG domain assembled in a trimeric complex formed between *Tc*BDF5, *Tc*BDF5BP (BDF5 Binding Protein) and *Tc*BDF8. In this complex we describe a novel way in which an MRG domain participates in the interaction with two proteins binding two different surfaces, instead of just one as previously reported. Together, these results show that AF2-processed interactomic datasets can be used to identify chromatin-remodeling protein complexes previously unknown in deeply branched eukaryotes.

Table of Contents Graphic

Introduction

Trypanosomatids are protozoans that diverged early in the eukaryotic tree of life, reflected in their highly divergent coding sequences (CDS), sometimes making the search for homology by sequence similarity extremely difficult[1]. Many of them cause human diseases such as sleeping sickness (*T. brucei*)[2], Chagas disease (*T. cruzi*)[3], and leishmaniasis (*Leishmania* spp)[4]. They have complex life cycles with different evolutionary stages that range from mammalian hosts to insect vectors. This formidable biological versatility is achieved despite the fact that these organisms are devoid of clear gene transcription regulation mechanisms, since they lack promoter sequences and associated factors typically found in other eukaryotes[5]. Based on these observations and other evidence, has led to hypothesize that transcription regulation in trypanosomatids is mainly epigenetic and gene expression would depend on the degree of chromatin accessibility[6]. Chromatin structure can be altered either by post-translational modifications of the N-terminal tails of histones, by the substitution of one or more histones by their variants, or by the combination of these events. These changes not only regulate the ability of chromatin to be transcribed but also other DNA-related processes such as recombination, replication, repair and chromosomal segregation during metaphase[7]. In recent years, a wide range of histone modifications has been described, as well as a limited repertoire of enzymes and proteins that serve to direct the placement (acetylases, methylases, kinases), the removal (deacetylases, demethylases, phosphatases) or the recognition (bromodomains, chromodomains, SANT, PHD fingers, WD40 repeats) of these modifications or "marks", in what has been called the "histone code"[8].

One of the most important post transcriptional modifications (PTM) that histone tails undergo is lysine acetylation[9], a chemical event that is mediated by multiprotein complexes with histone acetyl transferase (HAT) and/or histone deacetylase (HDAC) activities. Many of these complexes also contain bromodomain factors (BDF)[10,11], proteins with a small domain, the bromodomain (BD), formed by 4 α-helices connected by loops that forms a hydrophobic cavity that recognize acetyl-lysines. Despite components of these signaling pathways have been validated as therapeutic targets for many pathologies, including parasitic diseases[12,13], the identity of these complexes is just beginning to be revealed and there is little knowledge of their structures.

Recently, Staneva *et al* systematized the analysis of the proteins that compose many putative chromatin regulators, using co-immunoprecipitation (Co-IP) followed by mass spectrometry assays in the process to identify protein networks[14]. They performed Co-IP assays of the 7 BDFs recognizable in the *T. brucei* genome, 9 proteins with acetyl transferase domains and 7 proteins identified as potential HDACs, among others. Another approach to this problem was that of Jones *et al*, who used a proximity labeling methodology to generate an interactomic dataset of BDF5 from *L. mexicana* (*Lm*BDF5)[15]. Also, Vellmer *et al*[16], while trying to unveil the identity and composition of a novel SNF2 complex involved in H2A.Z deposition, derived in the conclusion that some of the proteins they identified were forming two different HAT complexes, involving *Tb*HAT1 and *Tb*HAT2. In Staneva´s and Jones´ interactomic datasets, also two distinct MRG (MORF4 related gene) domain-containing proteins were identified. These domains form a structural part of several complexes with HAT and HDAC activity, as described in yeasts and humans[17]. MRG domains seem to act as a "core" or "anchor" to nucleate the protein complex, where multiple proteins compete for a single binding site on the surface of MRG domains. MRGs interact via intrinsically disordered regions (IDR) that contain an FxLP

motif, vital for such interactions[18]. Upon binding, these IDRs acquire an extended conformation, fitting the phenylalanine of the FxLP motif within a small cavity of its partner MRG domain, resting on a conserved arginine[19]. A second segment, more variable, contributes to form a bipartite interaction. While the above cited works detected MRG domain containing proteins in *T. brucei* and *L. mexicana*, no MRG binding protein (MRGBP) was reported there.

On the other hand, recent years have been characterized by an explosion in the development of artificial intelligence algorithms that are contributing to solve previously unsolvable biological problems[20]. One outstanding milestone was the release of the deep learning model AlphaFold2 (AF2), trained to solve the three-dimensional structure of proteins with an unprecedented degree of detail[21]. More recently, after noticing the emerging property of AF2 in predicting protein complexes, AF2-multimer was built and trained using protein complexes as training database, resulting in a higher degree of accuracy [22].

In this work, we identified the *T. cruzi* homologs of the proteins present in public interactomic datasets, analyzed their sequences and focused mainly on the search for proteins that interact trough MRG domains. We then generated the structural models of their complexes with AF2-multimer, identifying the presence of two distinct MRG domain-driven assembled complexes. One is the ortholog of the yeast and human TINTIN complexes[23,24] (that we named *Tc*TINTIN), formed by the protein *Tc*BDF6 and two conserved hypothetical proteins, that we named *Tc*MRGx and *Tc*MRGBP. The second trimeric complex is formed between the MRG domain of *Tc*BDF5 (*Tc*BDF5-MRG), the protein that we named *Tc*BDF5BP (*Tc*BDF5 Binding Protein), and the bromodomain factor *Tc*BDF8, the ortholog of *Lm*BDF8 described by Jones *et al* in *L. mexicana*[25]. The protein-protein interactions predicted by the quaternary structure models of these two complexes, as well some details about their interactions, were validated using yeast hybrid assays and mutagenesis. Altogether, our results demonstrate that the models produced by AF2-multimer can be used to predict protein-protein interactions in trypanosomatids and are a tool with high predictive value when it comes to inferring protein function.

Results

Identification of putative trypanosomatid MRG binding proteins (MRGBPs)

In order to identify the MRG binding proteins, we took as reference the IDs of the interactomic datasets generated by CoIP-MS of *Tb*HAT1 and *Tb*BDF5 produced by Staneva *et al*[26]. *Tb*HAT1 interactome contains the protein Tb927.1.650, which has an HHblits-identifiable MRG domain; along with other 11 significantly enriched proteins, 4 of them annotated as "hypothetical proteins". By using HHblits[27], some of them could be identified as constituents of a complex orthologous to the yeast NuA4 (<u>Nu</u>cleosome <u>A</u>cetyltransferase of H<u>4</u>) complex. This is a 13-subunit complex that acts as a chromatin remodeler via its HAT activity and is involved in many processes like transcription and double-strand DNA break repair, among others[28]. However, several of the yeast NuA4 components could not be identified, even among the proteins enriched during reciprocal CoIPs of other components (*Tb*BDF6 and *Tb*EAF6). Given the high evolutionary distance between trypanosomatids and the model organisms where many chromatin remodeling complexes were described, we considered that some proteins apparently absent from the NuA4 complex could be hidden among the proteins identified as hypothetical in the CoIPs, being impossible to associate them as orthologs using algorithms such as HHblits due to their high divergence. MRG-binding proteins (MRGBPs) are part of the apparent missing proteins. These proteins are characterized by possessing an intrinsically

disordered region (IDR) involved in protein-protein interaction, with a linear motif that possesses an aromatic residue in the first position and a proline in the fourth, which in yeast and humans is an FxLP motif[18].

On the other hand, *Tb*BDF5 is a protein that has a C-terminal MRG domain. Among the proteins enriched in its interactome, there are two proteins associated with chromatin remodeling: *Tb*HAT2, known to acetylate histone H4 in K2, K5 and K10[29]; and the bromodomain factor *Tb*BDF8. *Tb*BDF5 interactome dataset also contains several hypothetical proteins. One of them is Tb927.9.13320, which possesses an FHA (forkhead associated) domain, a module involved in phospho-threonine recognition[30]. This protein network was proposed to be forming a complex only found in trypanosomatids that would act as a central regulator of transcription and RNA processing[15]. No MRGBPs were identified in this interaction network either.

We set out to find the unidentified MRGBPs among the hypothetical proteins by analyzing their sequence, *i.e.*, by searching for proteins with a predicted IDR that also contain something similar to a FxLP motif. Once identified, we thought that it would be possible to use the state-of-the-art protein-protein interaction structure prediction algorithm, AF2-multimer[22], to generate a quaternary structure model of these potential MRGBPs complexed with the identified MRG of the dataset. By looking at the models and their confidence statistics (pLDDT and PAE), we can filter those with bad statistics or that do not match the structural descriptions of MRGBPs. Finally, to validate these interactions we used a yeast two-hybrid (Y2H) approach using *T. cruzi* protein sequences[31]. In order to clone the genes and perform experiments in *T. cruzi*, we used OrthoMCL[32] to convert by orthology *T. brucei* IDs from the interactomic dataset of *Tb*HAT1 and *Tb*BDF5 to the corresponding ones from *T. cruzi*. **Table 1** and **Table 2** contain the IDs of the orthologs from *T. brucei*, *T. cruzi* and *L. infantum* reference strains.

To identify conserved sequences, we retrieved all trypanosomatid orthologs of each protein using TriTrypDB and aligned each orthology group separately using ClustalO. Because we were interested in IDRs, probability of disorder was predicted with the disorder predictor PrDOS. We also used the pLDDT per residue extracted from the precalculated protein structures of *T. cruzi* in AF2 database, since there is a direct correlation between a low pLDDT value and the presence of IDRs[33]. We mapped these annotations on AF2 predicted monomers and inspected the proteins´ structures and alignments one by one. In the *Tb*BDF5 CoIP dataset, we identified a conserved Fx[LIV]P motif over an IDR at the N-terminus of BDF8 (TcCLB.506559.310) and a conserved [FY]EPP motif at an N-terminal IDR of the FHA domain containing protein TcCLB.510747.120 (**Fig. 1a** and **b**, respectively), as the potential interactors of *Tc*BDF5-MRG (TcCLB.506755.120). BDF8 displays a domain architecture consisting of an N-terminal IDR (residues 1-60) containing the sequence FAIP, a central globular domain of unknown function (DUF) and a C-terminal BD (BD8) (**Fig. 1a**). TcCLB.510747.120 has three identifiable regions: an IDR N-terminal containing the sequence YEPP, a potential globular DUF with unclear boundaries and an FHA C-terminal (**Fig. 1b**). However, we did not find any protein with a conserved motif containing an aromatic aminoacid in the first position and a proline in the fourth among the IDR regions predicted for the proteins on HAT1 interactome. Therefore, we decided to compare their structures with AF2-predicted structures of known MRGBPs to search for structural similarities. We found that the hypothetical protein TcCLB.511217.190 has a globular domain followed by an IDR region similar to that of the human MRGBP protein

(*Hs*MRGBP). In addition, the region of similarity corresponds to residues involved in the interaction with the human MRG-containing proteins *Hs*MRG15 and *Hs*MRGx (**Fig. 1c**), indicating that TcCLB.511217.190 could be a good candidate for binding the protein with MRG domain TcCLB.505999.50. With this reduced set of candidates, we used AF2-multimer to predict the structures of the pairs TcCLB.505999.50/TcCLB.511217.190, TcCLB.506755.120/TcCLB.510747.120 and TcCLB.506755.120/TcCLB.506559.310. The structures obtained, the derived structure predictions and the experimental evidence that validates them are described in the following sections.

*Tc*TINTIN is a partially conserved eukaryotic complex formed by the sequential assembly of *Tc*MRGx, *Tc*MRGBP and *Tc*BDF6

The structural similarity of TcCLB.511217.190 with *Hs*MRGBP led us to think that it could be its ortholog. In humans, *Hs*MRGBP is involved in direct binding to the C-terminal MRG domain of *Hs*MRG15 or to the MRG domain of its ortholog *Hs*MRGx, which is smaller, as it lacks the N-terminal chromodomain present in *Hs*MRG15. Furthermore, *Hs*MRGBP also associates with the bromodomain protein *Hs*BRD8. Together with *Hs*MRG15 (or *Hs*MRGx), they form a complex known as Trimer Independent of NuA4 for Transcription Interactions (TINTIN)[24], which is also present in *Saccharomyces cerevisiae* formed by *Sc*Eaf3, *Sc*Eaf5 and *Sc*Eaf7[34]. As its name suggests, this trimer is part of the NuA4 complex, but it has transcriptional regulation functions that are independent of NuA4, since it is capable of being assembled independently. Given that BDF6 was proposed as an ortholog of *Hs*BRD8[14], we thought that the set TcCLB.505999.50 (*Tc*MRGx from here, given it domain architecture similarity to *Hs*MRGx), TcCLB.511217.190 (*Tc*MRGBP from here) and TcCLB.506529.660 (*Tc*BDF6) (**Fig. 2a**) could be forming a TINTIN-like complex in trypanosomatids.

Fig. 2b shows the statistics for the AF2-multimer prediction of the *Tc*MRGx/*Tc*MRGBP pair. The Predicted aligned error (PAE) plot indicates a low predicted error between the relative positions of the amino acids of both proteins (upper right and lower left quadrants), so it is possible to consider the generated model (**Fig. 2c**) as the structural prediction of this dimer. The pLDDT plot, which represents the confidence per-residue, indicates good confidence in the individual domains generated, with values below 60 only for some loops connecting α-helices. Values greater than 80 can even be observed at IDR zone that contacts *Tc*MRGx, indicating that AF2 is confident in its local structure. The mode of interaction is similar to the bipartite interaction observed for the NMR structure of the *Hs*MRG15/*Hs*MRGBP pair (PDB: 2N1D)[18], involving multiple hydrophobic amino acids, both in the globular region, and in the IDR (Supplementary Fig. 1). Regarding the residues where the FxLP motif should be, the I112, conserved among trypanosomatid orthologs, is found in *Tc*MRGBP, while *Tc*MRGx does not display any clear hydrophobic pocket over its conserved arginine R138 (on which I112 is resting). This suggests that this interaction may not be as important as in other MRGBPs (see below).

The statistics and the model for the prediction of the *Tc*MRGBP/*Tc*BDF6 pair are shown in **Fig. 2d** and **Fig. 2e**, respectively. The PAE plot shows values with low relative error only between the residue pairs involving the globular domain of *Tc*MRGBP and the BD of *Tc*BDF6, suggesting that the IDR of *Tc*MRGBP and the N-terminal domain of *Tc*BDF6 are not involved in the interaction. Consistently, a decrease in pLDDT values over the IDR of *Tc*MRGBP can also be seen on this model when compared to the model of *Tc*MRGx/*Tc*MRGBP dimer from **Fig. 2c**, suggesting that the IDR is only involved in the interaction with *Tc*MRGx. A low pLDDT value was

also observed over the IDR residues of *Tc*MRGBP monomer, indicating that *Tc*MRGBPmay be acquiring conformational stability only after binding to *Tc*MRGx. There is also a slight decrease in pLDDT values of other *Tc*MRGBP residues that contact *Tc*MRGx but not *Tc*BDF6.

Finally, we predicted the structure of the three-component complex (*Tc*MRGx/*Tc*MRGBP/*Tc*BDF6), which we will refer to as *Tc*TINTIN from now on. We did the same for the proteins *Sc*EAF3, *Sc*EAF7 and *Sc*EAF5, which together form the yeast TINTIN complex (*Sc*TINTIN); and with the proteins *Hs*MRG15, *Hs*MRGBP and *Hs*BRD8, which assemble *Hs*TINTIN. A comparison between these predictions can be seen in **Fig. 2f**. The PAE of the three complexes show good confidence between the relative positions of the atoms of all the domains of the three proteins, except for the N-terminal domains of *Sc*EAF5 in *Sc*TINTIN and *Tc*BDF6 in *Tc*TINTIN. This suggests that the N-terminal domains are not involved in the interaction. Due to video memory size limitations, we restricted the length of *Hs*BRD8 only to its BD residues, since it has a long insertion between the BD and its N-terminal domain, the latter being homologous to the N-terminal domains of *Tc*BDF6 and *Sc*EAF5 (Supplementary Fig. 2). So, it was not included in the prediction of *Hs*TINTIN, but we would expect to see a similar chart with low PAE values relative to the rest of the domains if it could be included. A good general correlation between high values of pLDDT and low values of intraprotein PAE is also observed, indicating good packing of the individual domains. From the structural point of view, the models present a similar spatial arrangement of the proteins that compose them. ScEAF7, TcMRGBP and HsMRGBP show the classic bipartite interaction described for MRGBPs, extending their IDR over the surface of their partner MRG domain, also binding via their globular domains. Meanwhile, *Sc*EAF5, *Tc*BDF6 and *Hs*BRD8 mediate their interaction with *Sc*EAF7, *Tc*MRGBP and *Hs*MRGBP, respectively, via two of their α-helices, exposing a hydrophobic pocket to the solvent that can potentially recognize acetylated lysines.

Although these models are consistent and clearly describe homologous structures, AF2-multimer has not yet been widely validated, nor is there experimental evidence indicating that its predicted quaternary structures can be assumed legitimate in trypanosomatids. Therefore, we set out to validate the formation of these complexes *in vivo* using the Y2H methodology previously used in our lab to test protein-protein interactions in multicomponent complexes[31]. In this methodology, protein-protein interaction is measured in yeasts by the expression level of reporter genes which confer the ability to grow in absence of essential molecules, together with the expression of β-*gal* whose expression can be easily seen through β-galactosidase-mediated colorimetric reactions[35].

Initially, we tested *Tc*MRGx/*Tc*MRGBP and *Tc*MRGBP/*Tc*BDF6 interactions. As expected, yeast clones expressing the *Tc*MRGx/*Tc*MRGBP interaction pair showed β-galactosidase activity (**Fig. 3a**). The blue color produced after X-Gal hydrolysis could be observed even one-hour post-incubation, suggesting that *Tc*MRGx/*Tc*MRGBP are strong interactors. No *Tc*MRGx, neither *Tc*MRGBP alone were able to induce reporter genes' expression (autoactivating controls). However, no β-galactosidase activity was seen for *Tc*MRGBP/*Tc*BDF6 pair, an unexpected result given the high confidence observed in the statistics of the models produced by AF2-multimer for this pair of proteins (**Fig. 2**). We thought that it was because *Tc*MRGBP/*Tc*BDF6 interaction was too weak to see with β-gal as reporter. Therefore, we tested *HIS3* reporter, which has a stronger promoter, but the spot growth assays performed did not show interaction phenotype either (Supplementary Fig. 3). We also checked the presence of the proteins by western blot to discard any defect in protein expression in yeasts (Supplementary

Fig. 4). Finally, we reanalyzed the interaction models, and realized that the motility of the free IDR of *Tc*MRGBP could only be retained after binding to *Tc*MRGx. Due to this, in a scenario where *Tc*MRGx is absent, the IDR chaotic movement could prevent the formation of a stable complex with *Tc*BDF6. But if *Tc*MRGx is present in the system, the IDR would stabilize (maybe inducing a conformational change on the globular domain of *Tc*MRGBP), allowing the assembly of the complete complex. If this is true, it would mean that *Tc*TINTIN should assemble in a sequential fashion. To test this hypothesis (**Fig. 3b**), we generated a third plasmid coding for *Tc*MRGx fused to GFP and introduced it into the strain expressing *Tc*MRGBP/*Tc*BDF6 interaction pair to perform a yeast-three hybrid (Y3H)-like experiment. Correct expression of GFP-*Tc*MRGx in the nucleus was verified by fluorescence microscopy (Supplementary Fig. 5). Fluorescent colonies were streaked into a master plate to evaluate their β-galactosidase activity, together with colonies from the strain expressing the pair *Tc*MRGx/*Tc*MRGBP to compare (**Fig. 3c**). After 24 hours, clones expressing full *Tc*TINTIN (*Tc*MRGx/*Tc*MRGBP/*Tc*BDF6) showed β-galactosidase activity, while the autoactivation controls did not show it at any time. The same result was observed using *HIS3* as reporter gene by spot growth assay (Supplementary Fig. 6).

These results do not only validate the existence of *Tc*TINTIN complex but also gives information about the dynamic of its assembly. This sequential assembly can potentially have regulatory effects on the function of the complex, perhaps modifying its affinity for NuA4 or even for the acetylated lysine(s) recognized by the bromodomain of *Tc*BDF6, since the loop between its B and C helices (BC loop), involved in acetyl lysine recognition in other bromodomains[36], is also involved in the interaction with *Tc*MRGBP. In this sense, the aminoacids on the hydrophobic pocket can potentially undergo conformational changes during binding or releasing from *Tc*TINTIN.

The MRG domain of *Tc*BDF5 forms a complex only observed in trypanosomatids and displays novel MRGBPs interaction modes

We also used the process described above for the predicted *Tc*BDF5 interactors. A representation of the domain architecture of *Tc*BDF5 and the proteins predicted as potential interactors of its C-terminal MRG domain is shown in **Fig. 4a**. We made several structural predictions for the pair constituted by *Tc*BDF5 and the FHA domain-containing protein TcCLB.510747.120, identifying low relative PAE values only between the amino acids of the MRG domain of *Tc*BDF5 and the IDR region of TcCLB.510747.120 (**Fig. 4b**). Comparing the pLDDT values for the IDR region observed in the TcCLB.510747.120 monomer of **Fig. 1b** and those of the dimer, we saw an increase in the maximum values from ~60 in the former to ~95 in the latter, pointing to a greater confidence in the prediction of its local structure in the dimer. The binding of this IDR shows the typical extended conformation, positioning the aromatic residue of the [FY]EPP motif inside a cavity formed on the surface of its partner MRG domain *Tc*BDF5-MRG (**Fig. 4c**), so we decided to name this protein as *Tc*BDF5BP (*Tc*BDF5 Binding Protein). Nonetheless, it also displays a novel way of interaction for MRGBPs, with the IDR in an extended-turn-extended conformation. The models indicate that the DUF and the FHA domain were not involved in the formation of the dimer (Supplementary Fig. 7). In an attempt to find potential homologs of *Tc*BDF5BP, we constructed a phylogenetic tree using the seed sequences of Pfam[37] FHA protein family (PF00498) and the FHA domain sequences of BDF5BP from *T. cruzi*, *T. brucei* and *L. infantum* (Supplementary Fig. 8). In this tree, the FHA clusters in a clade of proteins with similar size and domain architecture to that of *Tc*BDF5BP,

i.e., around 500 aminoacids, an N-terminal IDR, a central globular domain and a C-terminal FHA domain. They even had a coiled-coil between their central globular domain and the FHA, also observed between the DUF and FHA domains of *Tc*BDF5BP. Among them are *Hs*MCRS1 and *Dm*Rcd5 (*Drosophila melanogaster* Rcd5), which are part of various chromatin remodeling complexes. In particular, they are part of the NSL (Non-Specific Lethal) complex, with a fundamental role in dosage compensation of X chromosome through the acetylation of histone H4 in lysines 5, 8 and 16[38]. This complex shares its HAT enzyme (MOF) with another complex named MSL (Male-Specific Lethal). MOF modifies its substrate specificity when it is in MSL, losing the ability to acetylate lysines 5 and 8. Contrary to NSL, MSL contains an MRG protein, paralogous to MRG15, known as MSL-3[39]. As far as we know, there is no report about *Hs*MCRS1 or *Dm*Rcd5 binding any MRG domain.

For the predictions of the *Tc*BDF5/*Tc*BDF8 pair, we find models describing interaction not only through the predicted IDR of *Tc*BDF8, but also involving part of the DUF (**Fig. 4d** y **e**). As occurred with *Tc*BDF5BP, the pLDDT values for the IDR in *Tc*BDF8 monomer (**Fig. 1a**) are lower (~40) than those reached in the dimer (~85), indicating an improvement in the confidence of the packaging of these residues when the MRG of BDF5 is present in the prediction. In particular, the IDR peaks in pLDDT over the region containing the Fx[LIV]P motif. Structurally, the phenylalanine deepens into a hydrophobic cavity formed between two α-helices of *Tc*BDF5-MRG. This hydrophobic pocket has never been reported as a binding site for the conserved motif of any MRGBP. Another region of interaction occurs on the opposite side of the MRG domain, where several DUF α-helices wrap around one *Tc*BDF5-MRG α-helix. Taken together, this bipartite interaction of *Tc*BDF8 does not occupy the same cavities involved in *Tc*BDF5BP binding. Typically, several MRGBPs compete for the same binding site on the MRG[40], assembling different complexes depending on the context. In this case, AF2-multimer prediction suggest that *Tc*BDF5 is forming a trimer through its MRG domain with *Tc*BDF5BP and *Tc*BDF8 (**Fig. 4f**).

While reviewing the structures, we noticed a structural similarity between the folds of the α-helices present in *Tc*BDF8-DUF domain with the bromodomain fold (**Fig. 4g**). We made several structural alignments with different BD crystal structures and concluded that DUF is composed of two tandem bromodomain-like structures packaged into one globular domain. The only residues clearly conserved in *Tc*BDF8-DUF are mainly hydrophobic and located in opposed pairs at the core of the bromodomain-like structures, as if they had only a structural function in keeping the domain folded (Supplementary Fig. 9a). We saw no signs of conserved residues or hydrophobic pocket potentially involved acetyl-lysine recognition. On the contrary, this do not happen with the C-terminal bromodomain (BD8), which, although it does not have an essential asparagine involved in acetyl-lysine recognition, it has multiple clearly conserved hydrophobic residues on a small pocket (Supplementary Fig. 9b). Therefore, even though the bromodomain-like structures of DUF would have mainly a structural function, we cannot rule out the possibility that BD8 recognizes acetylated lysines.

To perform the Y2H assays, we fragmented *Tc*BDF5BP and *Tc*BDF8 to compare eventual positive interactions with AF2-multimer predictions. We generated truncated versions of *Tc*BDF5BP: one that only contains the IDR (IDR), another that contains the rest but not the IDR (NoIDR) and a third with the complete CDS (Full). To avoid the potential autoinduction caused by the BDs of *Tc*BDF5 binding to chromatin, we performed the tests only using its MRG domain. The results for the pair *Tc*BDF5-MRG/*Tc*BDF5BP are shown in **Fig. 5a**. As predicted by

the models, we only found β-galactosidase activity in yeasts that contain the MRG domain of *Tc*BDF5 together with the *Tc*BDF5BP versions that contains the IDR (Full and IDR on the figure). The blue color could be detected after one hour of incubation, suggesting a strong interaction. For *Tc*BDF8 we generated truncated versions: only its IDR (IDR), only DUF (DUF), only BD8 (BD8) and the complete CDS (Full). Again, we tested them against the MRG domain of *Tc*BDF5 (**Fig. 5b**). We were expecting to see a week interaction for *Tc*BDF5-MRG with both IDR and DUF, along no interaction with BD8 and a strong one with the full protein. Instead, we saw β-galactosidase activity only for the full protein only after 24 hours of incubation. This means that the dimer can only be formed if both IDR and DUF are present in the CDS of *Tc*BDF8 and it is not a strong interaction.

The number of predicted contacts with the MRG conserved arginine correlates with its importance in maintaining MRGBPs interaction

As discussed above, *Tc*MRGBP do not have any sequence in its IDR region that fit the FxLP motif described in other organisms (**Fig. 2c**). Instead, it has a conserved isoleucine that contacts loosely with the conserved arginine R138 of *Tc*MRGx (**Fig. 6a**). Furthermore, contrary to what is described for other MRGs[41], the surface exposed to the solvent around R138 has no clear hydrophobic pocket. Meanwhile, *Tc*BDF5BP has a conserved tyrosine (Y28) in its [FY]EPP motif that closely contacts the aliphatic portion of the conserved Arg R623 of *Tc*BDF5. Also, a clear cavity is present in this region, in which Y28 deepens. By comparison, *Tc*MRGx-R138 has 1 predicted atom to atom contact with *Tc*MRGBP-I112, while *Tc*BDF5-R623 shares 9 with *Tc*BDF5BP-Y28 (**Fig. 6a**).

In order to see if these structural observations were correlated to the strength of the interaction established between the MRGs and their corresponding MRGBPs, we mutated the conserved Arg in both MRG domains to Ala and tested the interactions again. Our hypothesis was that *Tc*MRGx/*Tc*MRGBP interaction would be less affected by changing R138 than *Tc*BDF5-MRG/*Tc*BDF5BP by changing R623. We explored this intervention by Y2H and the results are shown in the **Fig. 6b**. As predicted, the interaction was completely abolished for *Tc*BDF5-MRG-R623A/*Tc*BDF5BP, while *Tc*MRG-R138A/*Tc*MRGBP interaction was almost unchanged. This suggests that the [FY]EPP motif is essential to maintain interaction by contacting the R623 of *Tc*BDF5.

In general, this again shows the level of detail reached by the models produced by AF2-multimer in trypanosomatids, since we can perform predictions with them, allowing us to get information about the importance of conserved sequences in interactions.

Discussion

In this work, we addressed the problem of structurally define the chromatin remodeling complexes from trypanosomatids. We identify two heterotrimers that assemble through interactions with MRG domains: One partially conserved, TcTINTIN; and another completely novel, the one formed by *Tc*BDF5. We achieve this using several bioinformatics methods and subsequent validation by yeast hybrid approaches. We took particular advantage from having AF2 monomer database available as well as AF2-multimer available with ColabFold. This allows to overcome limitations due to very low similarity (less than 30%) and original domain architecture, as found in proteins from deep branched protists, like trypanosomatids. For example, *Tc*MRGBP has a 17% sequence identity with its ortholog *Hs*MRGBP, making its identification impossible using highly performant algorithms such as HHblits[27]. However, AF2

was able to model the conserved fold of its globular domain, helping us in the identification. In this case, we approached the problem using the expert's eye, since we extensively reviewed the literature describing the structures of complexes formed through MRG domains. Therefore, after reducing the number of potential MRGBPs by sequence analysis, we identified immediately that we were in presence of an MRGBP while reviewing the structures predicted with AF2-multimer. In this sense, we propose to use the predicted structures of AF2 to annotate sequences by structural similarity, automating the process of homology detection. For example, Foldseek[42] enables fast and sensitive comparisons of large structure sets and has a public server that finds structural similarities between any input PDB and the AF2 database.

In addition to the identification of the complexes, we were able to infer new biological features for them. In *Tc*TINTIN, given that the BC loop of BD6 is involved in the interaction with *Tc*MRGBP, it is possible that the activation of this interaction by *Tc*MRGx regulates the acetyl-lysine(s) recognition capability of the hydrophobic pocket of BD6, either by enhancing its release from acetylated nucleosomes or by promoting its binding. For example, in humans, MRG15 mediates the activation of ASH1L histone methyltransferase by releasing an autoinhibitory loop upon interaction[19]. Thus, it is remarkable that AF2 predicts interactions that can only happen under certain circumstances. Complementing this information with molecular dynamics calculations would add a key aspect in describing molecular interactions that is not considered by AF2.

On the other hand, the interaction modes of the structures predicted for the heterotrimer assembled through *Tc*BDF5-MRG are totally novel. The way in which the IDR of *Tc*BDF5BP folds into an extended-turn-extended conformation has not been described before. Even the MRG is quite divergent, possessing several large-disordered loops and being bigger than usual. Also, the cavity where the Fx[LIV]P motif of *Tc*BDF8 binds had never been described for an MRGBP. The latter may be a unique trypanosomatid feature, due to the divergence of this complex, or a yet undescribed interaction way present in eukaryotes. Given these observations, it is possible that a duplication of an MRG domain containing gene occurred at some point in evolution, opening the way to new of epigenetic regulation mechanisms through genetic drift. Then, the MRG domain of BDF5 would be the consequence of said duplication and traces of this event could be found in other organisms. Supporting this, we identified an MRG-like fold in KAT2A residues 227-370 (Supplementary Fig. 10), one of the best-studied proteins with HAT activity in the human genome[43], never reported to have an MRG-like structure. Far from stablishing that this domain is the ortholog of the MRG domain of BDF5, observations like this points to a picture more complicated than what it seems, and AF2 proves to be a great tool to clear it up.

We also explored the implication of the conserved arginine of *Tc*BDF5 and *Tc*MRG in maintaining the interaction with the IDRs of *Tc*BDF5BP and *Tc*MRGBP, respectively. We verified through mutagenesis assays that the union between the aromatic ring of *Tc*BDF5BP-Y28 and the aliphatic chain of *Tc*BDF5-R623 is essential to sustain the union. Given the essentiality of BDF5[15], an inhibitor could be developed that binds to the cavity formed on *Tc*BDF5-R623, where *Tc*BDF5BP-Y28 deepens, which is relatively different from that of *Hs*MRG15 and *Hs*MRGx. Alternatively, a PROTAC structure can be synthetized by linking the YEPP peptide to known E3 ligase ligands and assess whether this molecule manages to direct *Tc*BDF5 to its degradation[44]. Furthermore, with R623A mutant, we now have a model to study the biological implications of the TcBDF5-*Tc*BDF5BP interaction and investigate its essentiality.

Finally, during the search, the structures of many other protein monomers and complexes present in the datasets were also analyzed. Some of them could be in charge of assembling the great complex of which TINTIN or BDF5 are a part, but exploring this idea was beyond the objective of this work. However, we realize that with enough computing power, it is possible to compute the structure of the hypothetical megadalton complex of the BDF5-HAT2 network, as well as the structure of the entire HAT1-containing NuA4 complex. For that, AF2Complex could be used, designed to predict the structure of large multimeric complexes using high-performance computing (HPC)[45]. In spite of the implementation to be used, we proved that AF2 algorithm can be used in complex systems to decipher the structure of complexes form by very low, or even undetectable, sequence identity protein-components.

Materials and methods

Ortholog identification, alignments and disorder prediction

The identification of orthologous proteins was performed using the OrthoMCL[46] and TriTrypDB[32] databases, with the IDs of the sequences present in the *Tb*HAT1 and *Tb*BDF5 interactomes as search input (Table 1 and Table 2, respectively), filtering by organism in the cases in which it was required. Multiple sequence alignment was carried out using ClustalO with default parameters. Disorder prediction was done with the software PrDOS[47], using as query the sequences identified as orthologs from *T. cruzi* CL Brener strain.

Multiple sequence alignment, structure prediction, visualization and comparison

Multiple sequence alignments visualization was carried out using Jalview[48]. The monomer structures for each *T. cruzi* CL Brener ortholog were downloaded from AF2 protein structure database[49]. To map the predicted disorder and the degree of conservation into the structures, we setup Jalview to display them using PyMOL[50], representing these annotations as color codes. The conservation annotation was performed by setting the threshold for color display using a sequence conservation degree of at least 30%.

Structure predictions of the complexes were preformed using ColabFold-MMseqs2 notebook[51]. `msa_mode` was set to `MMseqs2(UniRef+Environmental)`, `pair_mode` to `unpaired+paired`, `model_type` to `AlphaFold2-multimer-v2` and `num_recycles` to 3.

Structure alignment and comparison was performed in ChimeraX using matchmaker algorithm, guided by Smith and Waterman algorithm, which also provides the final RMSD values.

CDS, cloning and expression vectors

All the cloning experiments were made using the Gateway cloning system[52]. Although the structures were predicted using the reference strain, the CDSs were cloned from the genomic DNA of Dm28c strain. Sequences ID and primers are shown in Supplementary Table 1. The sequences were amplified by PCR, purified and digested with the corresponding restriction enzymes, ligated into pENTR3C, transformed into electrocompetent DH5α cells and sequenced. The CDSs of *Tc*BDF5-MRG and *Tc*MRGBP were transferred to the Y2H vector pGBKT7-GW, leaving them in frame with the BD domain of GAL4 and a Myc-tag. The rest of the CDSs were transferred to the Y2H vector pGADT7-GW, leaving them in frame with the AD domain of GAL4 and an HA-tag. In order to perform the Y3H, we converted the yeast expression plasmid p426GPD to the gateway system, inserting the   Gateway   Cassette

frame B (Invitrogen) into its SmaI site. In order to track whether *Tc*MRGx was sent to the nucleus, we N-terminal inserted the CDS of GFP in frame to *Tc*MRGx into the vector pENTR3C_*Tc*MRGx. Finally, we transferred the GFP-*Tc*MRGx fusion to the pGPD426-GW vector. The mutations of the MRG domains were made by site directed mutagenesis of whole plasmids[53]. The expected sequences were confirmed in the resulting plasmids by sequencing.

Yeast strains

For the Y2H and Y3H experiments, *S. cerevisiae* MaV203 strain was used (*MAT*α, *leu*2-3,112, *trp*1-901, *his*3Δ200, *ade*2-101, *gal*4Δ, *gal*80Δ, *SPAL*10::URA3, *GAL1*::*lacZ*, $HIS3_{UAS}$ $_{GAL1}$::*HIS3@LYS2*, *can*1^R, *cyh*2^R). For transformations, electrocompetent cells were prepared from a 1/10,000 dilution of a saturated culture that was inoculated into 500 ml liquid YPAD medium at 30°C and grown ON to OD_{600}=1,0. Cells were cooled and centrifuged at 4000 g for 5 min. The supernatant was discarded, and the cells were washed twice with sterile distilled water and twice with 10% sorbitol. Cells were resuspended in 2 ml of 10% sorbitol and 40 µl per transformation mixed with up to 5 µl of plasmids were electroporated in 2 mm cuvettes (Gene Pulser Xcell BIO-RAD). For autoactivation controls, only one of the plasmids containing protein fusions was mixed with either empty pGBKT7 or empty pGADT7 (*e.g.*, pGBKT7/pGADT7-GW_*Tc*MRGx). For the interaction tests cells, a combination of two plasmids containing two fusion proteins was used (*e.g.*, pGBKT7-GW_*Tc*BDF6/pGADT7-GW_*Tc*MRGx). Electroporated cells were recovered in 1 ml of liquid YPAD with 10% sorbitol for 1 h at 30°C - 200 rpm and plated in Synthetic Complete (SC) medium lacking Trp to select pGBKT7 and lacking Leu for pGADT7 (SC-Leu-Trp).

The strains used in the Y3H experiments were derived from the parental strains MaV203 pGADT7-GW_*Tc*MRGBP pGBKT7-GW_*Tc*BDF6 (interaction test) and MaV203 pGADT7-GW_*Tc*MRGBP pGBKT7 (autoactivation control). They were generated using protocols mentioned above. Electroporated cells were selected using SC-Leu-Trp-Ura, to select for p426GPD-GW_GFP-*Tc*MRGx by also removing Ura. Only fluorescent colonies were used in the interaction assays.

Y2H and Y3H assays

Master plating: For each experiment, desired colonies were first streaked onto master plates with SC-Leu-Trp for Y2H experiments or SC-Leu-Trp-Ura for Y3H and grown for 24 h at 30°C. Four different clones were streaked for the interaction tests and two for each of the controls. **Replica plating:** The master plates were replicated using autoclaved velvets, transferring the streaks onto a nitrocellulose membrane supported on a YPAD plate to evaluate the expression of β-galactosidase. **X-Gal assay:** After incubation, nitrocellulose membranes were removed from YPAD plates and immersed in liquid N_2 for 20-30 s. Membranes were then transferred on top of round Whatman filters soaked with X-gal buffer (10 mg X-Gal, 100 µl DMF, 10 ml Z-buffer) colonies side up. The appearance of blue color was monitored over a 24-h period at 37°C. **Spot growth assay**. Spot growth assays were conducted to use *HIS3* as reporter gene. MaV203 cells transformed with plasmids derived from pGBKT7-GW and pGADT7-GW were grown overnight (ON) in liquid SC-Leu-Trp medium at 30°C with vigorous shaking. Yeast cells also transformed with p426GPD-GW_GFP-*Tc*MRGx were grown ON in liquid SC-Leu-Trp-Ura medium. The cells were suspended at OD_{600}=1,000 and serially diluted by 1/10. 7 µl of each culture dilution was spotted on solid SC medium alone or solid SC-His supplemented with 50 mM of 3-amino-1,2,4-triazole (3-AT) to titrate basal *HIS3* expression. Plates were incubated at 30°C and followed for 4 to 12 days.

Western blot

Cells were inoculated and grown for 48 h at 200 rpm and 30°C in 5 ml of the adequate liquid SC medium required to keep selective pressure. Cells were harvested by centrifugation (8000 g, 5 min) and the pellet was resuspended in 200 µl of extraction buffer (0.1 M NaOH, 0.05 M EDTA, 2% SDS, 2% β-mercaptoethanol). After 10 min at 96°C, 5 µl of 4 M acetic acid was added, followed by 50 µl of loading buffer. Eight µL of each sample was loaded per lane onto a 15% SDS-PAGE gel. After electrophoresis, the proteins were transferred to a nitrocellulose membrane and stained with Ponceau Red stain to ensure even loading and proper transfer. Blocking was done ON with PBS - 0.1% Tween20 (PBST) - 5% non-fat milk at 4°C. 3.3 µg/ml of rabbit anti-Myc primary antibody (Calbiochem) was incubated for 1 h at room temperature (RT) in PBST 5% milk. Three 10 min washes were performed with 5 ml of PBST. Goat secondary antibody anti-rabbit (GE) conjugated to HRP was incubated for 1 hour at RT (1/20000 dilution) in PBST 5% milk. The membrane was then stripped and incubated with rat anti-HA primary antibody (Roche) at 1/2000 dilution in PBST 5% milk and washed 3 times. Goat secondary antibody anti-rat (Pierce) conjugated to HRP was incubated for 1 hs at RT (1/10000 dilution) in PBST 5% milk.

Phylogenetic analysis

The FHA protein sequences for constructing the phylogenetic tree were retrieved from Pfam[37] entry PF00498, restricting the download only to seed sequences and requesting them pre-aligned. Residues 310-472 from TcCLB.510747.120, 328-507 from Tb927.9.13320 and 342-555 from LINF_350030200, corresponding to the FHA domains, were added to the multiple sequence alignment using MAFFT[54] version 7 setting the option `--add` to maintain the original alignment. Positions with low phylogenetic information were filtered using ClipKIT[55] using the trimming mode `-m kpic-gappy`. The phylogenetic tree was inferred by maximum-likelihood using IQ-TREE[56], setting a bootstrap of 1000 and using ModelFinder[57] to find the best substitution model. Tree visualizations were made with Dendroscope[58].

Data availability

Complex structure predictions have been uploaded as comments into the corresponding gene entries from TriTrypDB *T. cruzi* CL Brener strain. *Tc*TINTIN complexes: TcCLB.505999.50 (*Tc*MRGx), TcCLB.511217.190 (*Tc*MRGBP) and TcCLB.506529.660 (*Tc*BDF6). *Tc*BDF5-MRG complexes: TcCLB.506755.120 (*Tc*BDF5), TcCLB.510747.120 (*Tc*BDF5BP) and TcCLB.506559.310 (*Tc*BDF8).

Acknowledgments

This work was funded by the National Research Council, CONICET (PIP 2021-0848), the National Agency for Science, ANPCyT (PICT 2017-1978, 2020-01704) and Universidad Nacional de Rosario (0001-00285485). We want to thank Dr. Antonio Uttaro, who kindly give us the p426GPD plasmid.

References

1.   Gazestani, V. H., Yip, C. W., Nikpour, N., Berghuis, N. & Salavati, R. TrypsNetDB: An integrated framework for the functional characterization of trypanosomatid proteins. *PLoS Negl Trop Dis* **11**, 1–11 (2017).

2.   Büscher, P., Cecchi, G., Jamonneau, V. & Priotto, G. Human African trypanosomiasis. *The Lancet* **390**, 2397–2409 (2017).

3.   Pérez-Molina, J. A. & Molina, I. Chagas disease. *The Lancet* **391**, 82–94 (2018).

4.   Burza, S., Croft, S. L. & Boelaert, M. Leishmaniasis. *The Lancet* **392**, 951–970 (2018).

5.   De Gaudenzi, J. G., Noé, G., Campo, V. A., Frasch, A. C. & Cassola, A. Gene expression regulation in trypanosomatids. *Essays Biochem* **51**, 31–46 (2011).

6.   Saha, S. Histone Modifications and Other Facets of Epigenetic Regulation in Trypanosomatids: Leaving Their Mark. *mBio* **11**, 1–19 (2020).

7.   Kouzarides, T. Chromatin Modifications and Their Function. *Cell* **128**, 693–705 (2007).

8.   Biswas, S. & Rao, C. M. Epigenetic tools (The Writers, The Readers and The Erasers) and their implications in cancer therapy. *Eur J Pharmacol* **837**, 8–24 (2018).

9.   Yang, X. J. & Seto, E. HATs and HDACs: From structure, function and regulation to novel strategies for therapy and prevention. *Oncogene* **26**, 5310–5318 (2007).

10. Doyon, Y. & Côté, J. The highly conserved and multifunctional NuA4 HAT complex. *Curr Opin Genet Dev* **14**, 147–154 (2004).

11. Joshi, P. *et al.* The functional interactome landscape of the human histone deacetylase family. *Mol Syst Biol* **9**, 1–21 (2013).

12. Ye, F., Huang, J., Wang, H., Luo, C. & Zhao, K. Targeting epigenetic machinery: Emerging novel allosteric inhibitors. *Pharmacol Ther* **204**, 107406 (2019).

13. Pezza, A. *et al.* Essential Bromodomain TcBDF2 as a Drug Target against Chagas Disease. *ACS Infect Dis* (2022) doi:10.1021/acsinfecdis.2c00057.

14. Staneva, D. P. *et al.* A systematic analysis of Trypanosoma brucei chromatin factors identifies novel protein interaction networks associated with sites of transcription initiation and termination . *Genome Res* **31**, 2138–2154 (2021).

15. Jones, N. G. *et al.* Bromodomain factor 5 is an essential regulator of transcription in Leishmania. *Nat Commun* **13**, 4071 (2022).

16. Vellmer, T. *et al.* A novel SNF2 ATPase complex in Trypanosoma brucei with a role in H2A.Z-mediated chromatin remodelling. *PLoS Pathog* **18**, 1–29 (2022).

17. Bowman, B. R. *et al.* Multipurpose MRG domain involved in cell senescence and proliferation exhibits structural homology to a DNA-interacting domain. *Structure* **14**, 151–158 (2006).

18. Xie, T., Zmyslowski, A. M., Zhang, Y. & Radhakrishnan, I. Structural Basis for Multi-specificity of MRG Domains. *Structure* **23**, 1049–1057 (2015).

19. Lee, Y. *et al.* Structural Basis of MRG15-Mediated Activation of the ASH1L Histone Methyltransferase by Releasing an Autoinhibitory Loop. *Structure* **27**, 846-852.e3 (2019).

20. Sapoval, N. *et al.* Current progress and open challenges for applying deep learning across the biosciences. *Nat Commun* **13**, 1–12 (2022).

21. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

22. Evans, R. *et al.* Protein complex prediction with AlphaFold2-Multimer. *Methods in Molecular Biology* **804**, 297–312 (2021).

23. Bhat, W., Ahmad, S. & Côté, J. TINTIN, at the interface of chromatin, transcription elongation, and mRNA processing. *RNA Biol* **12**, 486–489 (2015).

24. Devoucoux, M. *et al.* MRG proteins are shared by multiple protein complexes with distinct functions. *Molecular & Cellular Proteomics* **21**, 100253 (2022).

25. Jones, N. G. *et al.* Bromodomain factor 5 is an essential regulator of transcription in Leishmania. *Nat Commun* **13**, 4071 (2022).

26. Staneva, D. P. *et al.* A systematic analysis of Trypanosoma brucei chromatin factors identifies novel protein interaction networks associated with sites of transcription initiation and termination. *bioRxiv* 2021.02.09.430399 (2021).

27. Cheung, A. C. M. & Díaz-Santín, L. M. Share and share alike: the role of Tra1 from the SAGA and NuA4 coactivator complexes. *Transcription* **10**, 37–43 (2019).

28. Kraus, A. J. *et al.* Distinct roles for H4 and H2A.Z acetylation in RNA transcription in African trypanosomes. *Nat Commun* **11**, (2020).

29. Almawi, A. W., Matthews, L. A. & Guarné, A. FHA domains: Phosphopeptide binding and beyond. *Prog Biophys Mol Biol* **127**, 105–110 (2017).

30. Cribb, P. & Serra, E. One- and two-hybrid analysis of the interactions between components of the Trypanosoma cruzi spliced leader RNA gene promoter binding complex. *Int J Parasitol* **39**, 525–532 (2009).

31. Amos, B. *et al.* VEuPathDB⊡: the eukaryotic pathogen , vector and host bioinformatics resource center. 1–14 (2021).

32. Wilson, C. J., Choy, W. Y. & Karttunen, M. AlphaFold2: A Role for Disordered Protein/Region Prediction? *Int J Mol Sci* **23**, 1–14 (2022).

33. Wang, X., Ahmad, S., Zhang, Z., Côté, J. & Cai, G. Architecture of the Saccharomyces cerevisiae NuA4/TIP60 complex. *Nat Commun* **9**, 1–11 (2018).

34. Invitrogen. *ProQuest^TM Two-Hybrid System with Gateway ® Technology*. (2002).

35. Ferri, E., Petosa, C. & McKenna, C. E. Bromodomains: Structure, function and pharmacology of inhibition. *Biochem Pharmacol* **106**, 1–18 (2016).

36. Mistry, J. *et al.* Pfam: The protein families database in 2021. *Nucleic Acids Res* **49**, D412–D419 (2021).

37. Cai, Y. *et al.* Subunit composition and substrate specificity of a MOF-containing histone acetyltransferase distinct from the Male-specific Lethal (MSL) complex. *Journal of Biological Chemistry* **285**, 4268–4272 (2010).

38. Kadlec, J. *et al.* Structural basis for MOF and MSL3 recruitment into the dosage compensation complex by MSL1. *Nat Struct Mol Biol* **18**, 142–150 (2011).

39. Redington, J., Deveryshetty, J., Kanikkannan, L., Miller, I. & Korolev, S. Structural insight into the mechanism of palb2 interaction with mrg15. *Genes (Basel)* **12**, (2021).

40. Hou, P. *et al.* Structural Insights into Stimulation of Ash1L's H3K36 Methyltransferase Activity through Mrg15 Binding. *Structure* **27**, 837-845.e3 (2019).

41. Remmert, M., Biegert, A., Hauser, A. & Söding, J. HHblits: lightning-fast iterative protein sequence searching by HMM-HMM alignment. *Nat Methods* **9**, 173–175 (2012).

42. Kempen, M. van *et al.* Foldseek: fast and accurate protein structure search. *bioRxiv* **1**, 2022.02.07.479398 (2022).

43. Haque, Md. E. *et al.* The GCN5: its biological functions and therapeutic potentials. *Clin Sci* **135**, 231–257 (2021).

44. Békés, M., Langley, D. R. & Crews, C. M. PROTAC targeted protein degraders: the past is prologue. *Nat Rev Drug Discov* **21**, 181–200 (2022).

45. Gao, M., Nakajima An, D., Parks, J. M. & Skolnick, J. AF2Complex predicts direct physical interactions in multimeric proteins with deep learning. *Nat Commun* **13**, 1744 (2022).

46. Ishida, T. & Kinoshita, K. PrDOS: Prediction of disordered protein regions from amino acid sequence. *Nucleic Acids Res* **35**, 460–464 (2007).

47. Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M. & Barton, G. J. Jalview Version 2-A multiple sequence alignment editor and analysis workbench. *Bioinformatics* **25**, 1189–1191 (2009).

48. Varadi, M. *et al.* AlphaFold Protein Structure Database: Massively expanding the structural coverage of protein-sequence space with high-accuracy models. *Nucleic Acids Res* **50**, D439–D444 (2022).

49. Mirdita, M. *et al.* ColabFold: making protein folding accessible to all. *Nat Methods* **19**, 679–682 (2022).

50. Reece-Hoyes, J. S. & Walhout, A. J. M. Gateway recombinational cloning. *Cold Spring Harb Protoc* **2018**, 1–6 (2018).

51. Laible, M. & Boonrod, K. Homemade site directed mutagenesis of whole plasmids. *Journal of Visualized Experiments* 1–3 (2009) doi:10.3791/1135.

52. Katoh, K., Rozewicki, J. & Yamada, K. D. MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. *Brief Bioinform* **20**, 1160–1166 (2018).

53. Steenwyk, J. L., Buida, T. J., Li, Y., Shen, X. X. & Rokas, A. ClipKIT: A multiple sequence alignment trimming software for accurate phylogenomic inference. *PLoS Biol* **18**, 1–17 (2020).

54. Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530–1534 (2020).

55. Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**, 587–589 (2017).

56. Huson, D. H. & Scornavacca, C. Dendroscope 3: An interactive tool for rooted phylogenetic trees and networks. *Syst Biol* **61**, 1061–1067 (2012).

Figures

Fig. 1: **Proteins identified as candidates to be MRGBPs of _T. cruzi_.** **(a)** AF2 predicted structure of TcCLB.506559.310 (_Tc_BDF8), disorder probability per residue graph, pLDDT per residue graph and fragment of the multiple sequence alignment for the predicted IDR. Different colors represent the identified domains in both structure and graphs. The dashed orange rectangle points to a zone with low pLDDT and high disorder probability which contains an Fx[LIV]P motif, shown in a green square in the multiple sequence alignment. **(b)** AF2 predicted structure of TcCLB.510746.120 (FHA domain containing protein), disorder probability per residue graph, pLDDT per residue graph and fragment of the multiple sequence alignment for its predicted IDR. As before, each identified domain was colored differently and the dashed orange rectangle indicates the zone in which a conserved [FY]EPP motif was identified. We included the ortholog of _B. saltans_, who contains a phenylalanine in the first position. A disordered loop was removed between the DUF and the FHA domain to enhance visualization. **(c)** Structure comparison between TcCLB.511217.190 and the region of _Hs_MRGBP known to interact with _Hs_MRG15. Both share a globular domain with the same fold, followed by an IDR with a α-helix. The residues corresponding to the FxLP motif of _Hs_MRGBP are shown, along with the sequence identity computed by Needleman-Wunsch algorithm and the RMSD calculation for the globular domain.

Fig. 2: Complex structure prediction of *Tc*TINTIN and its subunits compared to its homologs. **(a)** Domain representations, IDs and length of the proteins involved in the complex. **(b)** Statistics for the AF2-multimer structure prediction of the *Tc*MRGx/*Tc*MRGBP pair. On the left is the pLDDT per residue with the corresponding domains represented over the graph. On the right is the PAE plot with the domain correspondence remarqued in dashed squares. **Glob:** Globular domain **(c)** Complex structure prediction of the heterodimer *Tc*MRGx/*Tc*MRGBP. *Tc*MRGx is in green and *Tc*MRGBP is colored by pLDDT (pLDDT color labels are in panel **e**). An arrow points to the predicted IDR in *Tc*MRGBP. The residues of the IDR structure where the FxLP motif is expected to be is enlarged in a rectangle showing the sidechains E111, I112 and R113 of *Tc*MRGBP, with the surface exposed to the solvent of *Tc*MRGx colored by hydrophobicity (yellow: hydrophobic, blue: hydrophilic). **(d)** Statistics for the prediction of the *Tc*MRGBP/*Tc*BDF6 pair. **(e)** Predicted structure for the heterodimer *Tc*MRGBP/*Tc*BDF6. An arrow points to the predicted IDR of *Tc*MRGBP. The N-terminal domain of *Tc*BDF6 was not included in the visualization because it was not involved in the interaction, but was included in the prediction. Also, a disordered loop on BD6 was removed to enhance visualization. **(f)** Predicted structures of the heterotrimers of *S. cerevisiae* TINTIN (*Sc*TINTIN), *T. cruzi* TINTIN (*Tc*TINTIN) and *H. sapiens* TINTIN (*Hs*TINTIN) and its statistics (pLDDT and PAE). All five models of each prediction were aligned to the rank_1 model of *Tc*TINTIN in ChimeraX. Disordered loops and the non-interacting N-terminal domains from *Sc*EAF5 and *Tc*BDF6 were removed to facilitate visualization. Notice the lack of low interprotein PAE values in the PAE plot of *Hs*TINTIN. This is because we were computationally unable to include the N-terminal domain of *Hs*BRD8 in the prediction due to a big insertion between it and the bromodomain. The curved black arrows between panels **c** and **e** indicates that the combination *Tc*MRG/*Tc*MRGBP/*Tc*BDF6 forms *Tc*TINTIN. The hydrophobic pockets of *Sc*EAF5, *Tc*BDF6 and *Hs*BRD8 are shown in an inset for each trimer.

Fig. 3: Y2H and Y3H-like assays validate the formation of *Tc*TINTIN *in vivo* and suggests that is assembled sequentially. **(a)** Y2H assay to test the interaction pairs *Tc*MRG/*Tc*MRGBP and *Tc*MRGBP/*Tc*BDF6 using β-galactosidase as reporter gene. The blue circle has a reference to the yeast strains streaked on the master plate (SC-Leu-Trp). Control strains were streaked below the horizontal line, autoactivation controls were streaked at the right of the vertical line and the actual interaction tests to the left. **(b)** Cartoon of *Tc*TINTIN sequential assembly hypothesis. Left: *Tc*BDF6 cannot interact with *Tc*MRGBP in absence of *Tc*MRGx. Right: *Tc*MRGBP binds to *Tc*MRGx and changes conformation, activating the interaction with *Tc*BDF6 and transcribing the reporter gene. **(c)** Results from the Y3H-like experiment. The reference indicates the yeast strains that were streaked on the master plate (SC-Leu-Trp-Ura). The negative interaction control strain (-) was not included because it does not growth in absence of Ura, used to select GFP-*Tc*MRGx expression plasmid. A red rectangle indicates the yeast clones that expresses the three proteins from panel b-right, showing β-galactosidase signal at 24 hours.

Fig. 4: Predicted complex formed via the MRG domain of *Tc*BDF5. (a) Domain representation, IDs and length of the proteins involved in the complex. Positions of the YEPP sequence corresponding to the [FY]EPP motif and FAIP sequence corresponding to the Fx[LIV]P motif are indicated with a triangle over the IDRs of *Tc*BDF5BP and *Tc*BDF8, respectively. Solid black lines indicate the interaction regions with *Tc*BDF5-MRG predicted by AF2-multimer. FHA: Forkhead associated domain. DUF: Domain of unknown function. BD8: Bromodomain of *Tc*BDF8. **(b)** Model confidence statistics for the *Tc*BDF5-MRG/*Tc*BDF5BP-IDR heterodimer. **(c)** Structure of the complex formed between the IDR of *Tc*BDF5BP and the MRG of *Tc*BDF5. *Tc*BDF5 is colored in green and *Tc*BDF5BP is colored by pLDDT. The mode of interaction of the YEPP sequence is seen in the box, which is analogous to the FxLP motif observed in other MRG-domain partners. **(d)** Model confidence statistics for the *Tc*BDF5-MRG/*Tc*BDF8 heterodimer. **(e)** Structure prediction of the *Tc*BDF5-MRG/*Tc*BDF8 heterodimer. *Tc*BDF5 is colored in green and *Tc*BDF8 is colored by pLDDT (pLDDT color labels are in panel **c**). An arrow points to the IDR of *Tc*BDF8. The mode of interaction of the FAIP sequence is seen in the box with a 180˚ rotated point of view, with the surface exposed to the solvent of *Tc*BDF5-MRG colored by hydrophobicity. A black arrow points to the IDR domain of *Tc*BDF8. **(f)** Structure of the trimer formed by *Tc*BDF8/*Tc*BDF5/*Tc*BDF5BP, predicted with AF2-multimer. **(g)** Structure alignment between the two bromodomain-like structures of *Tc*BDF8-DUF in purple and the crystal structure of the BD of *Tc*BDF2 in light brown (PDB: 6NP7). RMSD calculations are shown below.

Fig. 5: Y2H assays validate the complexes predicted to be formed by the MRG domain of *Tc*BDF5. Y2H assay to test the interactions of both full length and fragments of *Tc*BDF5BP and *Tc*BDF8 with *Tc*BDF5-MRG, using β-galactosidase as reporter gene. A reference to the yest strains streaked in each master plate can be seen in the blue circle. An interpretation of these interactions is shown on the right. **(a)** Y2H between fragments of *Tc*BDF5BP and the MRG domain of *Tc*BDF5. Control strains are shown below the horizontal line, autoactivation strains where streaked to the right of the vertical line and interaction tests strains expressing TcBDF5-MRG and the fragments of TcBDF5BP are to the left of the vertical line. IDR: CDS containing just the IDR of *Tc*BDF5BP. NoIDR: CDS containing *Tc*BDF5BP without its IDR. Full: Full length of *Tc*BDF5BP. **(b)** Y2H between fragments of *Tc*BDF8 and the MRG domain of *Tc*BDF5. IDR: CDS containing just the IDR of *Tc*BDF8. DUF: CDS containing just the DUF of *Tc*BDF8. BD8: CDS containing just the BD of *Tc*BDF8. Full: Full length of *Tc*BDF5BP.

Fig. 6: Number of MRGBPs atoms in contact with the conserved arginine of its partner MRG correlates with the loss of interaction when mutated by alanine. (a) The I112 of *Tc*MRGBP that is in the position where the aromatic residue of the conserved FxLP motif should be, shows only has a single contact with the conserved arginine R138 of *Tc*MRGx in the AF2-multimer prediction. In contrast, the aromatic residue Y28 of the [FY]EPP motif of *Tc*BDF5BP is in close contact (9 atom to atom contacts) with the conserved arginine R623 of *Tc*BDF5. Note: R623 is the conserved arginine in the position 623 of Dm28c *Tc*BDF5, that is R634 for CL Brener (see materials and methods). **(b)** Spot growth assay of MaV203 strains expressing a combination of DB, AD, DB-*Tc*MRGBP, DB-*Tc*BDF5-MRG, DB-*Tc*BDF5-MRG-R623A, AD-*Tc*MRGx, AD-*Tc*MRGx-R138A and AD-*Tc*BDF5BP. The table on the left indicates the combinations tested in each row. Positive and negative interaction controls were also included. Dilutions of OD_{600}=1.000, 0.100, 0.010 and 0.001 were spotted on solid SC medium or solid SC medium without histidine (-His) supplemented with 50mM 3-AT to titrate basal HIS3 expression. Cells were incubated at 30 °C for 8 days.

Tables

Table 1: Identified orthologs of the proteins significatively enriched in *Tb*HAT1 CoIP from Staneva *et al*. The identification column contains the protein names or domains identified by Staneva *et al*. **Tb*YEA2 and *Tb*EAF6 are fused forming a single protein in *L. infantum*.

T. brucei ID	Identification	*T. cruzi* ID	*L. infantum* ID
Tb927.7.5310	*Tb*YEA2	TcCLB.506825.80	*LINF_060012900
Tb927.10.14190	*Tb*EPL1	TcCLB.506525.30	LINF_320005600
Tb927.8.5320	hypothetical protein, conserved	TcCLB.511217.190	LINF_160016800
Tb927.1.650	MRG domain protein	TcCLB.505999.50	LINF_200005700
Tb927.1.3400	*Tb*BDF6	TcCLB.506529.660	LINF_120009500
Tb927.11.3430	hypothetical protein, conserved	TcCLB.507099.100	LINF_130019500
Tb927.7.4560	*Tb*HAT1	TcCLB.506605.160	LINF_140006300
Tb927.9.2910	*Tb*EAF6	TcCLB.510565.170	*LINF_060012900
Tb927.6.1240	hypothetical protein, conserved	TcCLB.507603.54	LINF_120005400
Tb927.11.9640	glycyl-tRNA synthetase, putative	TcCLB.504017.79	LINF_360047300
Tb927.2.4580	UNC119	TcCLB.506979.50	LINF_270027100
Tb927.10.12040	mitogen-activated protein kinase 11, putative	TcCLB.510123.20	LINF_330022100, LINF_190020900

Table 2: Identified orthologs of the proteins significatively enriched in *Tb*BDF5 CoIP from Staneva *et al*. The identification column contains the protein names or domains identified by Staneva *et al*. *Found by BLAST. **IDs of the syntenic orthologs.

T. brucei ID	Identification	*T. cruzi* ID	*L. infantum* ID
Tb927.11.10070	*Tb*BDF3	TcCLB.510719.70	LINF_360042200
Tb927.11.5230	EMSY ENT domain	TcCLB.503981.50	LINF_240010100
Tb927.9.13320	FHA domain	TcCLB.510747.120	LINF_350030200
Tb927.4.2340	*Tb*BDF8 (similar to TFIID TAF1)	TcCLB.506559.310	LINF_340028400
Tb927.11.13400	*Tb*BDF5	TcCLB.506755.120	LINF_090020000
Tb927.7.2770	hypothetical protein, conserved	TcCLB.510525.30	LINF_220014800
Tb927.6.1070	hypothetical protein, conserved	TcCLB.507603.240	*LINF_120007400
Tb927.3.4140	Poly ADP Ribose Polymerase (PARP)	TcCLB.506175.70	LINF_290022100
Tb927.11.11530	*Tb*HAT2	TcCLB.509203.60	LINF_280029500
Tb927.11.18680	dynein light chain LC8	TcCLB.506925.104	LINF_320007400
Tb927.2.4580	UNC119	TcCLB.506979.50	LINF_270027100
Tb927.10.3280	60S ribosomal proteins L38	TcCLB.503575.34	LINF_030007300, LINF_260027600
Tb927.5.3210	small ubiquitin-related modifier	TcCLB.507809.70	LINF_080009800
Tb927.1.2230	small myristoylated protein.1-1	TcCLB.509003.30	LINF_200018300
Tb927.11.11290	heat shock protein 70	**TcCLB.511211.220	**LINF_280036400

Supplementary information

*Tc*MRGx/*Tc*MRGBP
(AF2-multimer dimer prediction)

Globular domain

IDR

*Tc*MRGBP
*Tc*MRGx

*Hs*MRGx/*Hs*MRGBP
(NMR structure, PDB 2N1D)

Globular domain
(fragment)

IDR

FxLP

*Hs*MRGBP
*Hs*MRG15

Supplementary Fig. 1: Comparison between the predicted *Tc*MRGx/*Tc*MRGBP dimer and the solution structure 2N1D. The best ranked predicted structure of the *Tc*MRGx/*Tc*MRGBP pair was aligned in ChimeraX using matchmaker algorithm with one of the conformers of the 2N1D solution structure and then displayed in two different panels. Inset shows the structure of the FxLP motif positioned on a pocket formed on the solvent-exposed surface of *Hs*MRG15 colored according to its degree of hydrophobicity (yellow = hydrophobic, light blue = hydrophilic). The arrow points to its position on the IDR of *Hs*MRGBP. The HsMRGBP sequence chosen to resolve by NMR in 2N1D omitted a fragment of the globular domain.

TcBDF6
HsBRD8
ScEAF5

C

N

Supplementary Fig. 2: The structure alignment of the N-terminal domains of *Tc*BDF6, *Hs*BRD8 and *Sc*EAF5 indicates that they are homologous. On the left are the models of these domains colored my chain. On the right are the models colored in rainbow colors by position, from blue in the N-terminal residue to red in the C-terminal residue.

Supplementary Fig. 3: Interaction tests by spot growth assay between *Tc*MRGBP and *Tc*BDF6 or *Tc*MRGx using HIS3 as reporter gene. Spot growth assay of MaV203 strains expressing a combination of AD, DB, DB-*Tc*MRGBP, AD-*Tc*BDF6 and AD-*Tc*MRGx. The table on the left indicates the combinations tested in each row. Positive and negative interaction controls were also included. Dilutions of OD_{600}=1.000, 0.100, 0.010 and 0.001 were spotted on solid SC medium or solid SC medium without histidine (-His) supplemented with 50mM 3-AT to titrate basal HIS3 expression. Cells were incubated at 30 °C for 4 to 12 days.

Supplementary Fig. 4: Protein expression assessed by western blot. Western blots of total extracts of the yeast strains from the master plate on *Fig. 3*a. The a-HA blot shows the bands corresponding to constructs containing the AD domain and the a-Myc blot shows the bands corresponding to constructs containing the DB domain. Ponceau stain (Load) displays even loading of the samples. Red arrows points to the bands corresponding to the expected sizes.

Supplementary Fig. 5: Direct fluorescence of GFP-*Tc*MRGx expressing cells. Nuclear GFP signal, coming from GFP-*Tc*MRG expression on both autoactivation control strain (that lacks *Tc*MRGBP) and the strain containing the three fusions from ***Fig. 3*c** (Full *Tc*TINTIN).

Protein 1 (DB fusion)	Protein 2 (AD fusion)	Protein 3 (GFP fusion)	SC day 6	SC – His+3AT 50mM day 6
*Tc*MRGBP	*Tc*BDF6	*Tc*MRGx		
-	*Tc*MRGx	*Tc*MRGx		
*Tc*MRGBP	*Tc*BDF6	-		
*Tc*MRGBP	*Tc*MRGx	-		
Control (++)				
Control (+)				
Control (-)				

Supplementary Fig. 6: *Tc*MRGx activates the interaction between *Tc*MRGBP and *Tc*BDF6. Spot growth assay of MaV203 strains expressing a combination of DB, AD-*Tc*BDF6, DB-*Tc*MRGBP, and AD-*Tc*MRGx. GFP-*Tc*MRGx was also expressed in some of the cells to test its effect on the interaction between TcMRGBP and TcBDF6. The table on the left indicates the combinations tested in each lane. Positive and negative interaction controls were also included. Dilutions of OD_{600}=1.000, 0.100, 0.010 and 0.001 were spotted on solid SC medium or solid SC medium without histidine (-His) supplemented with 50mM 3-AT to titrate basal HIS3 expression. Plates were incubated at 30 °C for 6 days. In this case we were able to grow the parental strain because of the experimental design, allowing us to observe the relative increase in HIS3 expression between the full TcTINTIN expressing cells compared to the parental.

Supplementary Fig. 7: Statistics for the prediction of the dimer *Tc*BDF5-MRG/*Tc*BDF5BP using the complete sequence of *Tc*BDF5BP. The PAE graph shows that only the IDR of *Tc*BDF5BP is involved in the interaction.

Supplementary Fig. 8: Phylogenetic tree of FHA domains. (a) Complete phylogenetic tree of the FHA domains sequences from the Pfam entry PF00498, adding the sequences of FHA domains of BDF5BPs from *T. brucei*, *T. cruzi* and *L. infantum*. The orange rectangle encloses the subtree from figure b. **(b)** IDs and branches from trypanosomatids BDF5BPs are shown in red. Green branches correspond to FHA proteins that clusters with trypanosomatids BDF5BP. The bold green IDs are characterized proteins similar in size and architecture to trypanosomatids BDF5BP. The IDs are followed by an underscore and the limits used for each protein to generate the tree.

Supplementary Fig. 9: Conserved residues of TcBDF8 domains. BD and BD-like structures present in *Tc*BDF8. Conserved residues are colored by conservation and by physicochemical properties. Not conserved residues are in white. Brighter colors refer to a higher degree of conservation. (**a**) Bromodomain-like structures from the DUF domain. The pairs of blue lines coming from the blue dots shows the pairs of conserved hydrophobic residues that may be involved in keeping the fold of the domain. The yellow circles points to the region in which must be located the conserver residues involved in acetyl-lysine recognition. (**b**) C-terminal bromodomain (BD8). The region involved in acetyl-lysine recognition has many conserved residues that could be involved in this function. The position 489 is expected to have a conserved asparagine, but instead it has a conserved tyrosine.

Supplementary Fig. 10: Structural similarity between residues 227 to 370 of human KAT2A and the MRG domain of human MRG15. The residues 227-370 of KAT2A predicted structure (AF2) were structurally aligned to the crystal structure of human MRG15 (PDB: 2F5J). An RMSD of 1.386 Å was obtained between 26 pruned atom pairs and of 7.737 Å for the entire structures.

Supplementary Table 1: Primers used in the study.

Primer	Gene name	Gene ID	Sequence
TcMRGx_BamHI-Fw	*Tc*MRGx	C4B63_22g316	AACGGATCCATGGAGGGAGTTGACTGG
TcMRGx_XhoI-Rv	*Tc*MRGx	C4B63_22g316	CCACTCGAGTCATTTCAAAGGAATATACGC
TcMRGx-R138A-Fw	*Tc*MRGx	C4B63_22g316	GTGGAGTATCTTCTTGCCTTTTTTATGGCTTTGC
TcMRGx-R138A-Rv	*Tc*MRGx	C4B63_22g316	GCAAAGCCATAAAAAAGGCAAGAAGATACTCCAC
TcMRGBP_SalI-Fw	*Tc*MRGBP	C4B63_55g378c	AAAGTCGACAATGCGATTGGCTACAAGTGG
TcMRGBP_XbaI-Rv	*Tc*MRGBP	C4B63_55g378c	AAATCTAGATTATCACTTTTCCTCACCCAGAAGC
TcBDF6-HA_BamHI-Fw	*Tc*BDF6	BCY84_19389	AAAGGATCCATGTATCCGTATGATGTGCCGGATTATGCTCGGCGGGAAGATTTACTGC
TcBDF6_EcoRV-Rv	*Tc*BDF6	BCY84_19389	AAAGATATCTCATGCACCACGCAAATGC
TcBDF5-MRG_BamHI-Fw	*Tc*BDF5	C4B63_1g51	TTGGATCCCCTCCGCTACGGGCTCC
TcBDF5-MRG_XhoI-Rv	*Tc*BDF5	C4B63_1g51	AAACTCGAGCTACGATATTTCTTTTCCAATTTCCG
TcBDF5-R623A-Fw	*Tc*BDF5	C4B63_1g51	CTTGTGTTATCTTGTTGCTTTTTTGCAGCACCTTC
TcBDF5-R623A-Rv	*Tc*BDF5	C4B63_1g51	GAAGGTGCTGCAAAAAAGCAACAAGATAACACAAG
TcBDF5BP_SalI-Fw	*Tc*BDF5BP	C4B63_2g92	AAAGTCGACAATGGATGGGTCGCGGTTTGG
TcBDF5BP_XbaI-Rv	*Tc*BDF5BP	C4B63_2g92	AAATCTAGATTATTAATCCTCCATCAACACCTCGG
TcBDF5BP-IDR_XbaI-Rv	*Tc*BDF5BP	C4B63_2g92	AAATCTAGATTATGTCGTCATGGGTGCTGCAG
TcBDF5BP-NoIDR_XbaI-Fw	*Tc*BDF5BP	C4B63_2g92	AAAGTCGACAATGCTGGCCTCATTGTTGGCCG
TcBDF8_SalI-Fw	*Tc*BDF8	C4B63_7g80	AAAGTCGACAATGGATTTGATTTTTGCTGAGGGAG
TcBDF8_XbaI_Rv	*Tc*BDF8	C4B63_7g80	AAATCTAGATTATCACAACAGGCCCACCTCG
TcBDF8-Peptide_XbaI-Rv	*Tc*BDF8	C4B63_7g80	AAATCTAGATTATGGCAGGACTCGTGCCGC
TcBDF8-DUF_SalI_Fw	*Tc*BDF8	C4B63_7g80	AAAGTCGACAATGGCGGCACGAGTCCTGCCAG
TcBDF8-DUF_XbaI_Rv	*Tc*BDF8	C4B63_7g80	AAATCTAGATTAAGGAAGTGGAGTCGGCGTC
TcBDF8_BD8_SalI-Fw	*Tc*BDF8	C4B63_7g80	AAAGTCGACAATGTCCTGCACTGATACACAGC