Diversified repertoire of phage defenses in *Klebsiella pneumoniae*: Bi-

directional steering effects impacting antibiotic susceptibility

Sue C. Nang[1], Jing Lu[1], Heidi H. Yu[1], Hasini Wickremasinghe[1], Mohammad A. K. Azad[1], Meiling

Han[1], Jinxin Zhao[1], Gauri Rao[2], Phillip J. Bergen[1], Tony Velkov[1], Norelle Sherry[3], David T.

McCarthy[4], Saima Aslam[5], Robert T. Schooley[5], Benjamin P. Howden[3], Jeremy J. Barr[6], Yan

Zhu[1], Jian Li[1*]

[1]Biomedicine Discovery Institute, Infection Program, Department of Microbiology, Monash

University, Clayton, Victoria, Australia.

[2]Division of Pharmacotherapy and Experimental Therapeutics, Eshelman School of Pharmacy,

University of North Carolina, Chapel Hill, North Carolina, USA.

[3]Microbiological Diagnostic Unit Public Health Laboratory, Department of Microbiology and

Immunology, The University of Melbourne at The Peter Doherty Institute for Infection and

Immunity, Melbourne, Victoria, Australia.

[4]Department of Civil Engineering, Monash University, Clayton, Victoria, Australia.

[5]Division of Infectious Diseases and Global Public Health, Department of Medicine, University

of California San Diego, La Jolla, California, USA.

[6]School of Biological Sciences, Monash University, Clayton, Victoria, Australia.

*Corresponding author. Email: jian.li@monash.edu

Abstract

Phage therapy is a promising alternative treatment for multidrug-resistant bacterial infections. Unfortunately, phage resistance represents a major barrier hindering the clinical utility. Mechanistic insights into bacterial phage defense mechanisms are critical to optimize phages therapy. Here, we discovered a repertoire of phage resistance mechanisms in a model strain *Klebsiella pneumoniae* MKP103, including the disruption of phage binding site (*fhuA*::Tn and *tonB*::Tn), extension of phage latent period (*mnmE*::Tn and *rpoN*::Tn) and increased mutation frequency (*mutS*::Tn and *mutL*::Tn). Different from the prevailing view that phage resistance re-sensitizes antibiotic-resistant bacteria, we demonstrated a bi-directional steering effect on antibiotic susceptibility. Specifically, it was uncovered that, while *rpoN*::Tn became more susceptible to colistin, *mutS*::Tn and *mutL*::Tn caused increased resistance to rifampicin and the last-line colistin. Our findings highlight the diversified strategies utilized by *K. pneumoniae* to overcome phage infection. Overall, mechanism-guided phage steering represents a vital strategy to maximize the success of phage therapy.

Introduction

Bacteriophages (phages) are viruses that specifically target bacteria[1, 2]. The earliest recorded therapeutic practice of phages for the treatment of bacterial infections took place almost a century ago[3, 4, 5], prior to the discovery of penicillin[6]. Unfortunately, as the golden age of antibiotic was on the rise, phage therapy was abandoned mainly due to technical and logistical hurdles[3]. In recent times, the 'perfect storm' of widespread antimicrobial resistance (AMR) combined with the sparse antibiotic development pipeline has dramatically reduced the antibiotic armamentarium available to physicians for treating multidrug-resistant (MDR) infections[7, 8]. Of particular concern are the ESKAPE pathogens which 'escape' the antibacterial effects of many antibiotics through a wide range of resistance mechanisms[9, 10]. With the dawn of a post-antibiotic era rapidly emerging, the World Health Organization (WHO) and the Centers for Disease Control and Prevention (CDC) have declared AMR as a major global health threat[11, 12]. Clearly, alternative antimicrobial therapies are urgently required and therefore, significant interest has risen in the use of phages for MDR bacterial infections[3, 4, 13]. To date, the clinical utility of phages has been hampered by a poor knowledge base on their biology, resistance mechanisms and combination use with antibiotics[3]. A concept receiving significant favorable attention recently in the AMR field is 'phage steering' which involves the use of phages to kill susceptible bacterial cells while re-sensitizing the remaining phage-resistant bacterial population to antibiotics[14, 15]. The utility of phage steering lies in the prospect of eliminating the entire bacterial populations using phage-antibiotic combinations[14, 15]; however, the intricacies and limitations of phage steering have yet to be explored. Here, we discovered an unprecedented bi-directional phage steering effect with decreased and increased antibiotic susceptibility associated with the repertoire of phage resistance mechanisms.

Results

Phage resistome screening reveals involvement of diverse genes

Most phage resistance studies in the literature have relied on the use of a population-based whole-genome sequencing method, which often limits the detection of phage resistance mechanisms to those associated with phage binding sites such as the capsule[15, 16], lipopolysaccharide (LPS) and outer membrane protein components[16, 17, 18]. Utilizing a strictly lytic *Siphoviridae* phage pKMKP103_1 (**Fig. 1**), we conducted a genome-wide resistome study by investigating the susceptibility of individual transposon mutants of *K. pneumoniae* MKP103[19]. Screening of phage resistome identified six genes, *rpoN*, *mutS*, *mutL*, *mnmE*, *fhuA* and *tonB*, in *K. pneumoniae* MKP103 that are associated with effective pKMKP103_1 infection (**Fig. 2a and Supplementary Table 1**). Subsequently, we proceeded to confirm the phage susceptibility of these candidate transposon mutants by evaluating: (i) the ability of pKMKP103_1 to form clear plaques on soft agar containing the respective mutants (**Fig. 2b**), and (ii) the killing kinetic profiles of pKMKP103_1 against the respective mutants in broth (**Fig. 2c**). Against *fhuA*::Tn and *tonB*::Tn, pKMKP103_1 did not show any lysis zone on agar plates (**Fig. 2b**) nor killing in broth (**Fig. 2c**), potentially attributed by the inability of pKMKP103_1 to infect these mutants. The efficiency of plating (EOP) of pKMKP103_1 on *mnmE*::Tn was reduced (**Fig. 2b**), showing that the ability of pKMKP103_1 to form plaque on this mutant was restricted. A reduced bacterial killing was also illustrated with *mnmE*::Tn over the initial 4 h (**Fig. 2c**). Interestingly, although the EOP of pKMKP103_1 on *rpoN*::Tn was unaffected (**Fig. 2b**), no bacterial reduction was observed in the time-kill studies over 24 h (**Fig. 2c**). With *mutS*::Tn and *mutL*::Tn, the EOP of pKMKP103_1 was similar to the WT (**Fig. 2b**); however, a weaker initial killing activity was evident in broth (**Fig. 2c**).

Phage resistance is associated with multiple mechanisms

As different phage susceptibility profiles were demonstrated by the aforementioned mutants, we proceeded to investigate the mechanisms causing these differential degrees of resistance. The adsorption of pKMKP103_1 to the bacterial surface of *fhuA*::Tn and *tonB*::Tn was inhibited (**Fig. 3a–b**). FhuA gating loop has been previously reported to be the binding site for phages T1, T5 and Φ80 in *Escherichia coli*[21], irrespective of conformational changes induced by TonB[22, 23]. In contrary to these earlier studies, pKMKP103_1 in our study potentially have a higher specificity for its binding to *K. pneumoniae* MKP103 as the disruption of either *fhuA* or *tonB* alone was sufficient to prevent the binding of pKMKP103_1 to this strain.

As the adsorption of pKMKP103_1 onto the bacterial surface of *mnmE*::Tn, *rpoN*::Tn, *mutS*::Tn and *mutL*::Tn was unaffected (**Fig. 3b**), we subsequently examined the impact for the disruption of these genes on the lytic lifecycle of pKMKP103_1 by examining the phage one-step growth profiles with these respective mutants. With *mutS*::Tn and *mutL*::Tn as the host strains, similar one-step growth profiles of pKMKP103_1 to that observed with the WT were demonstrated (**Fig. 3c**). Interestingly, a longer latent period (40 min) compared to the WT (30 min) was displayed by pKMKP103_1 when *mnmE*::Tn and *rpoN*::Tn were employed as the host strains; however, there were no substantial differences in the burst size (**Fig. 3c**). To the best of our knowledge, this study is the first to identify the association of phage resistance with disrupted bacterial *mnmE*, in which this gene encodes a GTPase involved in the modification of tRNA wobble uridine[24]. Our results indicate that mutation of *mnmE* can substantially delay phage replication; however, the exact mechanism warrants further investigation. For RpoN, an RNA polymerase factor sigma-54 which regulates the transcription of genes across diverse cellular functions[25, 26, 27, 28], a previous study reported its association

with phage resistance in *Pseudomonas aeruginosa*[29]. With the diverse regulatory roles played

by RpoN, the specific mechanism for its association with phage infection is unknown. Phage

resistance due to *rpoN* mutation was postulated to be associated with changes in LPS

structure as RpoN has been well known to regulate LPS biosynthetic pathway[29, 30]. We

conducted lipid A profiling and revealed that, compared to the WT, there were decreased

abundance of L-Ara4N-modified lipid A and increased abundance of 2-hydroxymyristate lipid

A in *rpoN*::Tn (**Fig. 4a**), confirming the regulatory role of RpoN on the lipid A modification

pathways in *K. pneumoniae* MKP103. Further examinations on the transposon mutants of

arnT (required for L-Ara4N modification of lipid A) and *lpxO* (required for 2-hydroxylation of

the myristate group on lipid A) demonstrated that the infectivity of pKMKP103_1 on these

mutants was similar to that on the WT. Our result shows that these lipid A species are not

involved in pKMKP103_1 infection.

Notably, we demonstrate for the first time that the rapid emergence of phage

resistance can occur in *K. pneumoniae* with disrupted *mutS* and *mutL*. Bacterial isolates with

interrupted *mutS* and *mutL* are commonly known as hypermutators[31, 32, 33, 34] and are

frequently associated with antibiotic resistance[35]. Compared to the WT, a significant increase

in mutation frequency to pKMKP103_1 was discovered with *mutS*::Tn and *mutL*::Tn (**Fig. 3d**).

Furthermore, complementation of *mutL* fully restored the mutation frequency of *mutL*::Tn.

While for *mutS*::Tn, a reduced mutation frequency was observed following complementation;

however, it did not reach the WT level. Altogether, these findings highlight the important role

of the DNA mismatch repair system in preventing the rapid emergence of resistance during

phage therapy. Different from our result in *K. pneumoniae* where MutL played a preventive

role against phage resistance, MutL caused phage resistance in *P. aeruginosa* by promoting

the deletion of a large chromosomal fragment containing *galU*, leading to a lack of *O*-antigen

(the binding site for phage PaoP5)[36]. Taken together, this highlights the complexity of the

functionality of this DNA mismatch repair system in relation to phage resistance among

different bacterial species.

Bi-directional steering effects on polymyxin susceptibility in phage-resistant mutants

In the contemporary literature, 'phage steering' has been portraited as a promising

antibacterial strategy, having the potential to eliminate the entire bacterial population with

phage-antibiotic combination[14, 15, 37]. There have been reports on increased susceptibility to

beta-lactams of a phage-resistant *A. baumannii* due to capsule loss[15] and an improved

susceptibility of *P. aeruginosa* to erythromycin, tetracycline and ciprofloxacin, following the

exposure to phage OMKO1 (known to interact with efflux pumps)[14]. In the case of *K.*

pneumoniae, the loss of the multidrug-resistance phenotype was observed following the

emergence of phage resistance, caused by the loss of antimicrobial resistance genes [*bla*$_{CTX-M}$,

ant(3"), *sul2*, *folA*, *mph(E)/mph(G)*] on a plasmid[37]. Hereby, we examined whether the phage

resistome repertoire identified in this study would affect the antibiotic susceptibility of

K. pneumoniae MKP103 (**Fig. 4b and Supplementary Table 2**). For the first time, we revealed

a bi-directional antibiotic susceptibility effect caused by different bacterial mechanisms of

resistance to a single phage. In comparison to the WT (rifampicin minimum inhibitory

concentration [MIC]=64 mg/L; colistin MIC=32 mg/L), phage-resistant *K. pneumoniae* mutants

with the hypermutator phenotype (*mutS*::Tn and *mutL*::Tn) exhibited increased resistance to

rifampicin (MIC>128 mg/L) and the last-line antibiotic colistin (MIC=128 mg/L); whereas

rpoN::Tn showed decreased resistance to colistin (MIC=4 mg/L; **Fig. 4b**). The

complementation of mutants with the respective WT genes restored the antibiotic

susceptibilities to the WT levels, confirming that the disruption of these genes was indeed

directly associated with the change in bacterial susceptibility to rifampicin and/or colistin in

K. pneumoniae MKP103 (**Fig. 4b**). Moreover, we conducted population analysis profiles (PAPs)

of colistin with *mutS*::Tn, *mutL*::Tn and *rpoN*::Tn to further examine the susceptibility change.

PAPs revealed an ~100-fold increase in the subpopulation of *mutS*::Tn and *mutL*::Tn on

Mueller Hinton (MH) agar containing 128 mg/L colistin, when compared to the WT (**Fig. 4c**).

In the case of *rpoN*::Tn, compared to the WT, there was ~1,000-fold reduction in the

subpopulation growing on MH agar containing 16 and 32 mg/L colistin and ~100-fold

reduction in the subpopulation growing on MH agar containing 64 mg/L colistin (**Fig. 4c**). The

reduced abundance of L-Ara4N-modified lipid A in *rpoN*::Tn (**Fig. 4a**) explained the reduced

colistin resistance in this mutant.

In conclusion, the bi-directional steering effects of phage therapy we discovered here

(**Fig. 5**) highlight the need for extra caution with phage steering and phage-antibiotic

combination therapy. Although multiple recent studies have demonstrated the potential of

phage steering as a promising approach to restore antibiotic susceptibility in bacteria[14, 15, 37],

our findings stress that the importance of understanding the underlying mechanisms.

Systematic mechanistic investigations on phage resistance and phage steering are required

to facilitate efficacious phage therapy in patients and minimize the risk for the emergence of

antibiotic resistance.

Methods

Bacteria and phage

K. pneumoniae MKP103 and the transposon library mutants were used in this study[19].

Transposon mutants were selected from Luria Bertani (LB) agar containing 100 mg/L

chloramphenicol during initial isolation[19]. All experiments were conducted in nutrient broth

(Oxoid) unless stated otherwise. Phage pKMKP103_1 was isolated from sewage water

obtained from a water treatment plant in Melbourne (Victoria, Australia) with *K. pneumoniae*

MKP103 as the host strain. The isolation, purification and amplification of phages were

conducted according to the Phage on Tap protocol[38].

Phage DNA extraction and genome sequencing

Phage pKMKP103_1 was propagated to a titer of ~10^{10} pfu/mL using the plate lysate

method[38] and DNA was extracted using the phenol-chloroform method[39]. Briefly, the phage

was treated with DNase I (Qiagen) and RNase A (Roche) to remove exogenous bacterial DNA

and RNA prior to the addition of sodium dodecyl sulphate and proteinase K to release the

DNA from phage particles. Phage DNA was isolated using phenol-chloroform followed by

precipitation with isopropanol and 3.0 M sodium acetate (pH 5.2). After two washes with 70%

ethanol, DNA was resuspended in DNase-free water. The quality and quantity of DNA were

determined using Nanodrop. Sequencing was performed by Genewiz (Suzhou, China) on the

Illumina HiSeq. Raw reads (150 bp paired-end) were trimmed and used for *de novo* assembly

using SPAdes 3.15.3[40]. Annotation was performed using RAST[41] and PHASTER[42], and

hypothetical proteins were further annotated using Blastp (≥90% coverage and ≥80%

similarity). Blast Ring Image Generator (BRIG) was used to create a representative

visualization of the phage pKMKP103_1 genome[20].

Transmission electron microscopy

High purity pKMKP103_1 was prepared using the cesium chloride gradient method[43].

For TEM imaging, 5 µL of the phage solution was applied to a glow-discharged carbon-coated

copper grid and stained with 1% uranyl acetate. Imaging was conducted on a JEOL JEM-

1400Plus TEM (Ramaciotti Centre for Cryo-Electron Microscopy, Monash University). ImageJ

(https://imagej.nih.gov/ij/) was used for visualizing and measuring the dimensions of the

phage.

Genome-wide transposon screening of phage-resistant mutants

The transposon mutant library of *K. pneumoniae* MKP103[19] was utilized for the

screening of phage pKMKP103_1-resistant mutants. A 96-pin replicator was used to inoculate

the mutants into 96-well plates (each well containing 200 µL of nutrient broth) which were

then incubated at 37°C for 18 h to reach $\sim 10^9$ cfu/mL. A 10-fold dilution (20 µL in 180 µL) was

conducted twice using 96-well plates to achieve a starting bacterial inoculum of $\sim 10^7$ cfu/mL.

Phage pKMKP103_1 was added to the wells to achieve a multiplicity of infection (MOI) of 1

($\sim 10^7$ pfu/mL). *K. pneumoniae* MKP103 (WT) was included as the control in each round of the

screening experiments. OD_{600nm} was measured at 8 and 10 h with a SPECTROstar® Nano

Microplate Reader (BMG Labtech); these time-points were chosen based on preliminary

experiments conducted with the WT strain which demonstrated an increase in OD_{600nm} after

10 h of incubation. Consequently, mutants with greater OD_{600nm} in comparison to the WT at

8 and/or 10 h ($OD_{600nm} \geq 0.12$) were potential resistant mutants. This transposon mutant

library consists of 1 to 6 individual transposon-inserted mutant(s) of each non-essential

gene[19]. The screening process was narrowed to the genes of which ≥2 individual transposon

mutants were identified; additionally, sole transposon mutants were also included to prevent

missing out of any potential candidates. With these selected mutants, turbidimetric assays

were conducted in three biological replicates followed by time-kill analyses to confirm the

phage-resistant phenotype.

Complementation of mutants

Sanger sequencing was first conducted to confirm that the mutants contain the

transposon at the indicated site (**Supplementary Table 3**). To complement the gene into the

respective transposon mutant, the gene was amplified by PCR from the WT strain

(**Supplementary Table 3**). PCR products and pBBR1MCS-5[44] (carrying the gentamicin

resistance cassette) were digested over 4 h with the restriction enzymes targeting the

relevant restriction sites (**Supplementary Table 3**). Following clean-up of the digested

products, the PCR product and pBBR1MCS-5 were ligated overnight at 4°C. The ligation

mixture was then transformed into chemically competent *Escherichia coli* DH5α and plated

on LB agar containing 20 mg/L gentamicin. Colony PCR was conducted to identify the colonies

with successful ligation of the gene of interest to pBBR1MCS-5. The plasmid was extracted

from *E. coli* DH5α using a QIAprep® Miniprep Kit and Sanger sequencing was conducted to

confirm the correct ligation. The complemented plasmids were electroporated into the

electrocompetent transposon mutants of the respective gene. An empty pBBR1MCS-5 vector

(p$_{Vector}$) was also electroporated into WT and all mutants as control strains for the experiments

to exclude vector effects. Successful transformation was confirmed with the growth of

colonies on LB agar containing 20 mg/L gentamicin, followed by PCR.

Turbidimetric assays

K. pneumoniae strains were grown in nutrient broth at 37°C with shaking (200 rpm)

for 18 h. The overnight bacterial culture was inoculated in fresh nutrient broth (1:100 dilution)

and incubated until log-phase with an OD_{600nm} of 0.5 (equivalent to ~10^8 cfu/mL). The culture

was then diluted 1:10 with fresh nutrient broth to reach a final inoculum of ~10^7 cfu/mL.

Phage pKMKP103_1 was added at an MOI of 1 (final titer of ~10^7 pfu/mL) or 10 (final titer of

~10^8 pfu/mL). Continuous OD_{600nm} readings were taken hourly for 20 h with a SPECTROstar®

Nano Microplate Reader (BMG Labtech).

Time-kill kinetics

K. pneumoniae strains were grown in nutrient broth at 37°C with shaking (200 rpm)

for 18 h. The overnight bacterial culture was inoculated in fresh nutrient broth (1:100 dilution)

and incubated until log-phase with an OD_{600nm} of 0.5 (equivalent to ~10^8 cfu/mL). The culture

was then diluted 1:10 with fresh nutrient broth to reach a final inoculum of ~10^7 cfu/mL.

Phage pKMKP103_1 was added at an MOI of 10 (final titer of ~10^8 pfu/mL). For mutants

carrying the empty vector and complemented strains, 5 mg/L gentamicin was added as the

selection pressure to maintain the plasmid throughout the experiment. An aliquot of 200 μL

of culture was collected at 0, 1, 2, 4 and 24 h, centrifuged at 10,000 × *g*, and the bacterial

pellet was resuspended in an equal volume of 0.9% NaCl. Following 10-fold dilutions with 0.9%

NaCl, 10-μL spots were plated on nutrient agar for viable counting.

Efficiency of plating (EOP)

The EOP was conducted to measure the lysis ability of the phage[45]. An overnight

bacterial culture was prepared as described above, with 100 µL of the culture subsequently

mixed with 4 mL soft agar and poured onto a nutrient agar plate to form double-layered agar

containing bacteria. Phage pKMKP103_1 at 10^9 pfu/mL (determined with the WT) was

subjected to a series of 10-fold dilutions using SM buffer (50 mM Tris-HCl, 100 mM NaCl, 8

mM MgSO$_4$, pH7.4) and plated onto the double-layered agar as 10-µL spots. Plates were then

incubated overnight at 37°C before enumeration of visible plaque.

Phage adsorption assay

K. pneumoniae was grown to log-phase as described above and diluted to ~10^7 cfu/mL

(final volume of 1 mL). Phage pKMKP103_1 was then added to reach a final titer of 2.5×10^4

pfu/mL (MOI of 0.0025). At 2-, 4- and 10-min following co-incubation, 100-µL was collected

and centrifuged at 16,000 × g for 1 min. The supernatant containing the unabsorbed phage

was then spotted in three technical replicates on double-layered agar containing either *K.*

pneumoniae MKP103 or *K. pneumoniae* MKP103 (p$_{Vector}$); strains carrying the plasmid (empty

vector and complemented) were plated with *K. pneumoniae* MKP103 (p$_{Vector}$). Plaques were

enumerated following overnight incubation at 37°C. Three biological replicates were

conducted for each group.

Fluorescent and confocal microscopic assay

Phage pKMKP103_1 was incubated with 100× SYBR™ Gold Nucleic Acid stain

(Invitrogen™) at 4°C for 4 h. The stained pKMKP103_1 was then transferred to an Amicon®

Ultra-15 centrifugal filter tube (MWCO 100 kDa) and centrifuged at 3,220 × g and 4°C to

remove excess stain. Two further washes with SM buffer were conducted prior to

resuspension of the stained pKMKP103_1 in an equal volume of SM buffer. Enumeration of

the plaques was performed to determine the titer of stained pKMKP103_1 for the microscopic

assays. The log-phase bacterial culture was grown and diluted to 2×10^7 cfu/mL, and

incubated with FM™ 4-64FX membrane stain (Invitrogen™) at room temperature for 5 min.

The SYBR™ Gold stained pKMKP103_1 was added to the FM™ 4-64FX stained *K. pneumoniae*

at an MOI of 100 (fluorescent microscopy) and 1,000 (confocal microscopy). For fluorescent

microscopy, the mixture was applied directly onto the glass slide. For confocal microscopy,

the mixture was mixed with molten 2% agarose (55°C) to fix the sample for imaging.

One-step growth curve

Log-phase *K. pneumoniae* culture was diluted to ~10^7 cfu/mL and phage pKMKP103_1

added to achieve a final titer of 10^5 pfu/mL (MOI 0.01). Following 2-min incubation at 37°C

the culture was centrifuged at $16,000 \times g$ for 1 min, the supernatant containing the free phage

was removed and the bacterial pellet was resuspended in an equal volume of nutrient broth.

Aliquots (100 µL) were then removed at 10-min intervals and added to a glass tube containing

4 mL soft agar with 100 µL bacterial culture of *K. pneumoniae* MKP103 or *K. pneumoniae*

MKP103 (p$_{Vector}$). Following a rapid vortex, the entire content of the tube was poured onto a

nutrient agar plate to form a double-layered agar. Plaques were enumerated following

overnight incubation at 37°C.

Mutation frequency to pKMKP103_1

The mutation frequency of *K. pneumoniae* strains to pKMKP103_1 was examined

using the agar overlay method[46] with modifications. *K. pneumoniae* was grown to log-phase

as described above to an OD_{600nm} of 0.5 (equivalent to ~10^8 cfu/mL). For mutants carrying the

empty vector and complemented strains, 5 mg/L gentamicin was added as the selection

pressure to maintain the plasmid. Bacterial culture was centrifuged at 16,000 × g at 4°C and

the pellet was resuspended in an equal volume of 0.9% NaCl. The bacterial culture was serially

diluted and 100 μL was added to a glass tube containing 4 mL soft agar with pKMKP103_1

(10^9 pfu/mL). Control samples without phage were included to obtain the total number of

colonies. A double-layered agar was prepared as described above. Colonies were counted

following a 20-h incubation at 37°C. Mutation frequency was calculated by dividing the

number of colonies formed on the agar containing pKMKP103_1 over the total number of

colonies. Statistical analysis was conducted using the multiple comparisons with uncorrected

Dunn's test on GraphPad Prism 9.3.1.

Antibiotic susceptibility testing and population analysis profiles (PAPs)

329   Susceptibility of *K. pneumoniae* strains to a range of antibiotics was determined using

broth microdilution (Trek Sensititre®, ThermoFisher Scientific) as per the manufacturer's

instructions. The lowest antibiotic concentration that inhibited the visible growth of bacteria

was the MIC. For mutants that showed a 2-fold difference in MIC compared to the WT strain

and demonstrated restoration of the MIC following complementation (**Supplementary Table**

2), additional broth microdilution assays were conducted in two technical replicates and

confirmed on two separate occasions. All MIC assays were conducted in cation-adjusted (22.5

mg/L Ca^{2+}, 11.25 mg/L Mg^{2+}) Mueller Hinton broth (CAMHB).

337   PAPs were undertaken as described previously with minor modifications[47]. Briefly,

bacterial culture was grown in CAMHB until log-phase and following 10-fold serial dilutions,

cultures were plated onto MH agar containing colistin at 2, 4, 8, 16, 32, 64 or 128 mg/L. Viable

colonies were enumerated following overnight incubation at 37°C.

Lipid A profiling

A single bacterial colony was inoculated in CAMHB and incubated at 37°C with shaking

(200 rpm) for 18 h. The overnight culture was then diluted 1:100 with fresh CAMHB (final

volume, 200 mL) and grown to an OD_{600nm} of ~0.7 (8.35±0.13 log_{10} cfu/mL). Bacterial pellets

were collected via centrifugation at 9,000 × g for 10 min, washed twice with 0.9% NaCl, and

resuspended in 4 mL of 0.9% NaCl. Lipid A extraction was conducted with minor

modifications[48]. Methanol (10 mL) and chloroform (5 mL) were then added to prepare a

single-phase Bligh-Dyer mixture (chloroform:methanol:water, 1:2:0.8, v/v) for the extraction

of LPS. LPS pellet was obtained by centrifugation at 3,220 × g for 10 min and washed once

with single-phase Bligh-Dyer mixture. The washed pellet was resuspended in 5.4 mL of

hydrolysis buffer (50 mM sodium acetate, pH 4.5) and incubated in a boiling water bath for 1

h. Following hydrolysis, methanol (6 mL) and chloroform (6 mL) was added to make a two-

phase Bligh-Dyer mixture (chloroform:methanol:water, 1:1:0.9, v/v). Following centrifugation

at 3,220 × g for 10 min, the lower phase of the mixture containing the lipid A was collected,

dried and stored at -20°C. Prior to LC-MS analysis, the dried lipid A was resuspended in 200

µL of chloroform:methanol (1:1, v/v), centrifuged at 14,000 × g for 10 min, and 150 µL of

supernatant was collected. An equal volume of isopropanol:water (2:1, v/v) was then added

prior to centrifugation at 14,000 × g for 10 min, and 200 µL of supernatant was obtained for

LC-MS analysis on the Dionex U3000 high-performance liquid chromatography (HPLC) system

in tandem with a Q-Exactive Orbitrap high-resolution mass spectrometer (Thermo Fisher).

Acknowledgements

We would like to express our gratitude to the Ramaciotti Centre for Cryo-Electron Microscopy

(Monash University) and Micromon Genomics (Monash University) for their facilities to

conduct TEM imaging and Sanger sequencing. This study was supported by the National

Institute of Allergy and Infectious Diseases of the National Institutes of Health (R21 AI156766)

and a Monash-UCSD Seed Fund. J.L. is an NHMRC Principal Research Fellow (APP1157909).

The content is solely the responsibility of the authors and does not necessarily represent the

official views of the National Institute of Allergy and Infectious Diseases or the National

Institutes of Health.

Contributions

S.C.N. and J.L. conceived the study. D.T.M provided the sewage sample for phage isolation.

S.C.N, J.L., H.H.Y., H.W., M.A.K.A, M.H., J.Z., N.S., B.P.H and Y.Z. conducted the experiments.

S.C.N, P.J.B. T.V. and J.L. wrote and edited the manuscript. G.R., T.V., S.A., R.T.S., J.J.B and J.L.

supervised and funded the project. All authors reviewed and agreed to the final manuscript.

References

379   1.   Taylor MW. *Viruses and man: A history of interactions*. Springer, Cham (2014).
380

381   2.   Campbell A. The future of bacteriophage biology. *Nat Rev Genet* **4**, 471-477 (2003).
382

383   3.   Lin DM, Koskella B, Lin HC. Phage therapy: An alternative to antibiotics in the age of
384   multi-drug resistance. *World J Gastrointest Pharmacol Ther* **8**, 162-173 (2017).
385

386   4.   Brives C, Pourraz J. Phage therapy as a potential solution in the fight against AMR:
387   Obstacles and possible futures. *Palgrave Commun* **6**, 100 (2020).
388

389   5.   Chanishvili N. Phage therapy--history from Twort and d'Herelle through Soviet
390   experience to current approaches. *Adv Virus Res* **83**, 3-40 (2012).
391

392   6.   Fleming A. Classics in infectious diseases: On the antibacterial action of cultures of a
393   penicillium, with special reference to their use in the isolation of *B. influenzae* by
394   Alexander Fleming, Reprinted from the British Journal of Experimental Pathology
395   10:226-236, 1929. *Rev Infect Dis* **2**, 129-139 (1980).
396

397   7.   Butler MS, Blaskovich MA, Cooper MA. Antibiotics in the clinical pipeline at the end of
398   2015. *J Antibiot (Tokyo)* **70**, 3-24 (2017).
399

400   8.   Rossolini GM, Arena F, Pecile P, Pollini S. Update on the antibiotic resistance crisis.
401   *Curr Opin Pharmacol* **18**, 56-60 (2014).
402

403   9.   Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial
404   resistance in 2019: A systematic analysis. *Lancet* **399**, 629-655 (2022).
405

406   10.   Rice LB. Federal funding for the study of antimicrobial resistance in nosocomial
407   pathogens: no ESKAPE. *J Infect Dis* **197**, 1079-1081 (2008).
408

409   11.   World Health Organization. Antibiotic resistance. https://www.who.int/news-
410   room/fact-sheets/detail/antibiotic-resistance (2020).
411

412   12.   Centers for Disease Control and Prevention. Antibiotic/Antimicrobial resistance
413   (AR/AMR). https://www.cdc.gov/drugresistance/index.html (2021).
414

415   13.   Kortright KE, Chan BK, Koff JL, Turner PE. Phage therapy: A renewed approach to
416   combat antibiotic-resistant bacteria. *Cell Host Microbe* **25**, 219-232 (2019).
417

418   14.   Gurney J*, et al.* Phage steering of antibiotic-resistance evolution in the bacterial
419   pathogen, *Pseudomonas aeruginosa*. *Evol Med Public Health* **2020**, 148-157 (2020).
420

421   15.   Gordillo Altamirano F*, et al.* Bacteriophage-resistant *Acinetobacter baumannii* are
422   resensitized to antimicrobials. *Nat Microbiol* **6**, 157-161 (2021).
423

424   16.   Hesse S, *et al.* Phage resistance in multidrug-resistant *Klebsiella pneumoniae* ST258
425   evolves via diverse mutations that culminate in impaired adsorption. *mBio* **11**,
426   e02530-02519 (2020).

428   17.   Burmeister AR, *et al.* Pleiotropy complicates a trade-off between phage resistance and
429   antibiotic resistance. *Proc Natl Acad Sci U S A* **117**, 11207-11216 (2020).

431   18.   Li P, Lin H, Mi Z, Xing S, Tong Y, Wang J. Screening of polyvalent phage-resistant
432   *Escherichia coli* strains based on phage receptor analysis. *Front Microbiol* **10**, 850
433   (2019).

435   19.   Ramage B, *et al.* Comprehensive arrayed transposon mutant library of *Klebsiella*
436   *pneumoniae* outbreak strain KPNIH1. *J Bacteriol* **199**, e00352-00317 (2017).

438   20.   Alikhan NF, Petty NK, Ben Zakour NL, Beatson SA. BLAST Ring Image Genarator (BRIG):
439   Simple prokaryote genome comparisons. *BMC Genomics* **12**, 402 (2011).

441   21.   Killmann H, Videnov G, Jung G, Schwarz H, Braun V. Identification of receptor binding
442   sites by competitive peptide mapping: phages T1, T5, and phi 80 and colicin M bind to
443   the gating loop of FhuA. *J Bacteriol* **177**, 694-698 (1995).

445   22.   Killmann H, Braun V. Energy-dependent receptor activities of *Escherichia coli* K-12:
446   mutated TonB proteins alter FhuA receptor activities to phages T5, T1, $\oslash$80 and to
447   colicin M. *FEMS Microbiol Lett* **119**, 71-76 (1994).

449   23.   Langenscheid J, Killmann H, Braun V. A FhuA mutant of *Escherichia coli* is infected by
450   phage T1-independent of TonB. *FEMS Microbiol Lett* **234**, 133-137 (2004).

452   24.   Armengod ME, *et al.* Enzymology of tRNA modification in the bacterial MnmEG
453   pathway. *Biochimie* **94**, 1510-1520 (2012).

455   25.   Liu X, *et al.* Involvement of RpoN in regulating motility, biofilm, resistance, and
456   spoilage potential of *Pseudomonas fluorescens*. *Front Microbiol* **12**, 641844 (2021).

458   26.   Mahmud A, *et al.* Genome-scale mapping reveals complex regulatory activities of
459   RpoN in *Yersinia pseudotuberculosis*. *mSystems* **5**, e01006-01020 (2020).

461   27.   Leang C, *et al.* Genome-wide analysis of the RpoN regulon in *Geobacter sulfurreducens*.
462   *BMC Genomics* **10**, 331 (2009).

464   28.   Burgess RR. *Encyclopedia of genetics: Sigma factors*. Academic Press (2001).

466   29.   Wright RCT, Friman VP, Smith MCM, Brockhurst MA. Cross-resistance is modular in
467   bacteria-phage interactions. *PLoS Biol* **16**, e2006057 (2018).

469   30.   Lloyd MG, *et al.* Targeting the alternative sigma factor RpoN to combat virulence in
470   *Pseudomonas aeruginosa*. *Sci Rep* **7**, 12615 (2017).

31. Prunier AL, Leclercq R. Role of *mutS* and *mutL* genes in hypermutability and
recombination in *Staphylococcus aureus*. *J Bacteriol* **187**, 3455-3464 (2005).

32. Willems RJ, Top J, Smith DJ, Roper DI, North SE, Woodford N. Mutations in the DNA
mismatch repair proteins MutS and MutL of oxazolidinone-resistant or -susceptible
Enterococcus faecium. *Antimicrob Agents Chemother* **47**, 3061-3066 (2003).

33. Oliver A, Baquero F, Blázquez J. The mismatch repair system (*mutS*, *mutL* and *uvrD*
genes) in *Pseudomonas aeruginosa*: Molecular characterization of naturally occurring
mutants. *Mol Microbiol* **43**, 1641-1650 (2002).

34. Galán JC*, et al.* Mutation rate is reduced by increased dosage of *mutL* gene in
Escherichia coli K-12. *FEMS Microbiol Lett* **275**, 263-269 (2007).

35. Woodford N, Ellington MJ. The emergence of antibiotic resistance by mutation. *Clin
Infect Dis* **13**, 5-18 (2007).

36. Shen M*, et al. Pseudomonas aeruginosa* MutL promotes large chromosomal deletions
through non-homologous end joining to prevent bacteriophage predation. *Nucleic
Acids Res* **46**, 4505-4514 (2018).

37. Majkowska-Skrobek G, Markwitz P, Sosnowska E, Lood C, Lavigne R, Drulis-Kawa Z.
The evolutionary trade-offs in phage-resistant *Klebsiella pneumoniae* entail cross-
phage sensitization and loss of multidrug resistance. *Environ Microbiol* **23**, 7723-7740
(2021).

38. Bonilla N, Rojas MI, Netto Flores Cruz G, Hung SH, Rohwer F, Barr JJ. Phage on tap-a
quick and efficient protocol for the preparation of bacteriophage laboratory stocks.
PeerJ **4**, e2261 (2016).

39. Dunstan RA*, et al.* The flagellotropic bacteriophage YSD1 targets *Salmonella* Typhi with
a Chi-like protein tail fibre. *Mol Microbiol* **112**, 1831-1846 (2019).

40. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes de novo
assembler. *Curr Protoc Bioinformatics* **70**, e102 (2020).

41. McNair K, Aziz RK, Pusch GD, Overbeek R, Dutilh BE, Edwards R. Phage genome
annotation using the RAST pipeline. *Methods Mol Biol* **1681**, 231-238 (2018).

42. Arndt D*, et al.* PHASTER: A better, faster version of the PHAST phage search tool.
Nucleic Acids Res **44**, W16-W21 (2016).

43. Thung TY*, et al.* Component parts of bacteriophage virions accurately defined by a
machine-learning approach built on evolutionary features. *mSystems* **6**, e0024221
(2021).

44.   Kovach ME*, et al.* Four new derivatives of the broad-host-range cloning vector
519   pBBR1MCS, carrying different antibiotic-resistance cassettes. *Gene* **166**, 175-176
520   (1995).

522   45.   Kutter E. Phage host range and efficiency of plating. *Methods Mol Biol* **501**, 141-149
523   (2009).

525   46.   Blasco L*, et al.* Combined use of the Ab105-2φΔCI lytic mutant phage and different
526   antibiotics in clinical isolates of multi-resistant *Acinetobacter baumannii*.
527   *Microorganisms* **7**, 556 (2019).

529   47.   Li J*, et al.* Heteroresistance to colistin in multidrug-resistant *Acinetobacter baumannii*.
530   *Antimicrob Agents Chemother* **50**, 2946-2950 (2006).

532   48.   Henderson JC, O'Brien JP, Brodbelt JS, Trent MS. Isolation and chemical
533   characterization of lipid A from Gram-negative bacteria. *J Vis Exp*, e50623 (2013).

Figure 1. Basic characterization of phage pKMKP103_1. **(a)** Time-kill kinetics of phage pKMKP103_1 against *K. pneumoniae* MKP103 at an MOI of 10 (*n*=3). **(b)** One-step growth curve of pKMKP103_1 with *K. pneumoniae* MKP103 (*n*=3). **(c)** A representative TEM image of pKMKP103_1. The scale bar represents 100 nm. **(d)** The pKMKP103_1 genome visualized using BRIG[20]. The colors represent the functions of the genes: orange (DNA packaging, replication, and modification), lime (transcription), green (host lysis), blue (head structural components), red (tail structural components), gray (others/unknown functions), and black (hypothetical).

Figure 2. Identification of *K. pneumoniae* MKP103 genes that are correlated with phage

pKMKP103_1 resistance and the associated resistance phenotype. **(a)** Workflow for the

genome-wide transposon screening of pKMKP103_1 resistome using the transposon mutant

library. **(b)** The plaque forming unit (pfu) when pKMKP103_1 was plated on the respective *K.*

pneumoniae MKP103 strains. Individual plaque counts were not available for *mnmE*::Tn and

mnmE::Tn (p$_{vector}$); the count was calculated based on the highest dilution in which lysis zone

could be observed. Data are presented as mean ± s.d. (*n*=3). **(c)** Time-kill kinetics of

pKMKP103_1 against *K. pneumoniae* MKP103 and mutant strains at an MOI of 10 (*n*=3).

Figure 3. Phage pKMKP103_1 adsorption, one-step growth profiles and bacterial mutation

frequency towards pKMKP103_1. **(a)** Visualization of pKMKP103_1 adsorption determined

using fluorescent (top panels) and confocal (bottom panels) microscopy. The pKMKP103_1

was stained with SYBR^{TM} Gold strain (green) and *K. pneumoniae* MKP103 strains were stained

with FM^{TM} 4-64FX (red). **(b)** pKMKP103_1 adsorption profile with respective *K. pneumoniae*

strains (*n*=3). **(c)** One-step growth profile of pKMKP103_1 in the presence of different *K.*

pneumoniae strains. Data is presented as the mean ± s.d. (*n*=3). **(d)** The mutation frequency

of *K. pneumoniae* strains to phage pKMKP103_1 (*n*=3). Statistical significance is denoted with

*p<0.05.

Figure 4. Lipid A and antibiotic susceptibility profiles. **(a)** Analysis of the individual lipid A species abundance in relative to the total abundance (n=3). Representative mass spectrum profile and chemical structure of each lipid A species. **(b)** Minimum inhibitory concentrations (MICs) of rifampicin and colistin. Increased and decreased MICs compared to *K. pneumoniae* MKP103 (WT) were represented by red and blue font colors, respectively. **(c)** Population analysis profiles of *K. pneumoniae* strains on MH agar containing colistin at 2, 4, 8, 16, 32, 64 and 128 mg/L (n=3).

Figure 5. Schematic diagram representing the proposed mechanisms of phage resistance in *K. pneumoniae* MKP103, and the interrelationships with rifampicin and colistin resistance. **(a)** A TonB-dependent FhuA is required for the adsorption of phage. **(b)** MnmE potentially involves in the modification of tRNA, enhancing the protein synthesis for phage particles. **(c)** RpoN could affect the transcription of the phage genome. **(d)** RpoN regulates bacterial transcription for cationic lipid A modifications that confer colistin resistance. MutS and MutL mediate DNA mismatch repair prevent chromosomal mutations associated with resistance to **(e)** phage, **(f)** rifampicin and colistin.