A Global Survey of Host, Aquatic, and Soil Microbiomes Reveals Ecological Properties Shared between Bacterial and Fungal Generalists

Daniel Loos (1), Ailton Pereira da Costa Filho (2),

Amelia E. Barber (2), Gianni Panagiotou (1,3,4)

(1) Research Group Systems Biology and Bioinformatics, Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute. Jena, Germany.

(2) Junior Research Group Fungal Informatics, Institute of Microbiology, Friedrich Schiller University. Jena, Germany.

(3) Institute of Microbiology, Friedrich Schiller University. Jena, Germany.

The State Key Laboratory of Pharmaceutical Biotechnology, The University of Hong Kong. Hong Kong SAR, China.

(4) Department of Medicine, The University of Hong Kong. Hong Kong SAR, China.

correspondence: gianni.panagiotou@leibniz-hki.de or amelia.barber@uni-jena.de

Abstract

Microbiome engineering is a fast-evolving area with relevance for human health, agriculture, and climate management solutions. Despite significant efforts in engineering microbiomes to repair dysbiotic communities, new microbes often fail to establish and/or alter ecosystem function. To identify bacterial and fungal genera with the desired ability to stably incorporate in diverse microbial communities, we retrieved paired 16S and ITS rRNA amplicon sequence data from 1,580 host, soil, and aquatic samples and explored the ecological patterns of the 2,977 bacteria and 1,740 fungal genera detected across all samples. Through this large-scale analysis, we revealed that a small number of bacterial and fungal generalists with high

prevalence across all environments positively contribute to the taxonomic diversity of their respective kingdom and explain a large percentage of the variation in the cross-kingdom community structure. We also observed that bacterial and fungal generalists have a significantly higher abundance compared to specialists, or genera whose prevalence was strongly associated with a single habitat - possibly due to their ability to stimulate positive associations with other highly prevalent genera. These findings can streamline existing strategies to identify bacterial and fungal inoculants with a higher probability to establish in recipient ecosystems and confer noticeable changes in their structure and function.

Introduction

With only a handful of exceptions, environments, plants, and animals are colonized with communities of microbial organisms, termed the microbiome. These microbial consortia play critical roles in the function and health of their hosts and habitats. For example, the microbiota contributes remarkably to the earth's biomass and plays an important role in shaping soil and oceanic biogeochemical processes.[1–3] Microbial communities also play critical roles in host's health. In humans, the microbiome helps digest food, produces essential vitamins, educates, and regulates the immune system, and protects against more harmful microbes.[4–11] Similarly, in plants, mycorrhizal fungi are critical symbionts for growth[12] and mediate disease-resistance.[13] However, these microbial communities can be altered or disturbed, leading to functional changes and dysbiosis.

An exciting new frontier in microbiome research is the rational editing of disbalanced microbial communities. This has been demonstrated in humans through both fecal microbiota transplants that cause global changes in the microbiome structure[14] and more targeted approaches such as the administration of probiotics that shift the gut microbiome towards a beneficial state.[15] Similar strategies are also being explored in agriculture through biological control agents and the direct application of non-pathogenic microbes to fight plant disease caused by *Pythium ultimum*, *Gaeumannomyces graminis*, and *Fusarium oxysporum*,[16] as well as the application of bioactive signaling compounds that attract beneficial microbes such as

Bacillus and *Lysinibacillus*.[17] However, the selection of species to be introduced
directly or serve as chassis strains to modify the structure and function of a dysbiotic
microbiome is challenging and in many cases based largely on their ability to be
easily cultivated and known phenotypic properties. Moreover, their ecological
properties are rarely considered in bacteria and are completely ignored in fungi.

To this end, we performed a large-scale meta-analysis of community sequencing
data sets from host, soil, and aquatic environments with paired bacterial and fungal
characterization to shed light on the ecological properties of the genera present and
their potential use as global or habitat-specific modifiers of microbial communities.
We focused specifically on three main properties: (i) the capacity of bacterial and
fungal genera to adapt to diverse environments (generalists) or specific
environments (specialists), which may be important for long-term establishment
when introducing new species into an environment; (ii) the relative abundance of
bacterial and fungal generalists and specialists which reflects their competitive
colonization potential and is a critical factor for both native and chassis strains to
make a functional impact on the ecosystem; and (iii) whether their presence in a
habitat might trigger global changes in inter- and cross-kingdom population
structure. These global changes can have the benefit of completely restructuring a
disbalanced community when this is the desired outcome. However, they also pose
the risk of establishing a completely new microbiome state with unpredictable
consequences.

Results

Environmental specificity of bacterial and fungal communities

To perform a cross-habitat comparison of bacterial-fungal communities, we
analyzed paired 16S and ITS rRNA amplicon sequence data from 1,580 aquatic, host-
, and soil - associated samples originating from 27 habitats deposited in NCBI
databases.

Samples were collected from Europe, Asia, and the Americas between 2010 and
2018 (Figure 1A & B). Of the 1,580 samples, 871 originated from host- (including

both mammalian and non-mammalian hosts, such as insects), 494 from soil-, and

215 from aquatic-associated environments. The habitats that contributed largest

number of samples for each environment were gut (N=287) and skin (N=68) for the

host, temperate (N=498) and conifer forests (N=147) for the soil and large lakes

(N=87) followed by other freshwater (N=71) for the aquatic (see Supplementary

table 1 for details of all projects). Taxonomic annotation of bacterial and fungal

communities was performed using SILVA and UNITE databases, respectively.

Rarefaction curves of each habitat indicated that the majority of habitats adequately

capture the diversity of both the bacterial and fungal communities (Figure 1C). In

total 2,977 bacteria and 1,740 fungal genera were detected across all samples

(Figure 1D). We next examined the overlap of genera between environments, where

a genus was considered shared if it was detected in at least one habitat in each of the

three different environments (host, aquatic, and soil). Soil and aquatic environments

had the highest number of shared bacterial genera (N=1,662), followed by host-soil

(N=1,483) and host-aquatic (1,226). The pattern was different for shared fungal

genera, with host-soil sharing the most (N=884), followed by aquatic-soil (N=205)

and host-aquatic (N=189). This trend remained after controlling for the different

number of samples across the three environments in 842 and 998 out of 1000

random down sampled subsets for bacteria and fungi, respectively.

While 40% of the total bacterial genera were found in all three environments, the

percentage dropped to only 11% for fungal genera, indicating a higher degree of

environmental specificity (Figure 1E & F). The most prevalent phyla that were

detected in all three environments were *Proteobacteria* followed closely by

Firmicutes for bacteria, and *Ascomycota* for fungi. For both bacteria and fungi, soil

was the environment with the highest percentage of uniquely detected genera

(i.e. genera not detected in any sample from host or aquatic origin), with 23% and

38%, respectively for each two kingdom. While aquatic-specific bacteria accounted

for 7% of the total number of detected genera, the percentage of unique fungi in

aquatic samples was only 2% (Figure 1 E & F). The opposite trend was observed for

host-associated microbes, with 3% and 8% of unique bacteria and fungi, respectively

in this environment.

We subsequently compared the relative abundance of genera that were found in all

environments or were uniquely detected in soil-, host- or aquatic-associated

environments. Bacterial genera detected in all three environments were significantly

more abundant (Wilcoxon Rank Sum test, $p < 0.001$) than genera uniquely detected

in one of the environments (Figure 1G). A similar pattern was observed with fungi.

However, a notable exception was the relative abundance of fungi uniquely detected

in aquatic samples, which was the highest among both the common genera as well as

those uniquely detected in soil- or host-associated environments (Figure 1H). Taken

together, we find that soil bacteria and fungi show a higher degree of specificity and

genera commonly detected in all environments were also more abundant in these

microbial communities.

Bacterial and fungal generalists are more abundant than specialists

Although a classical concept in ecology, generalists and specialists have remained

challenging to objectively identify and quantify. To define multi-kingdom generalists

and specialists, we set the following criteria: generalists are genera found with high

prevalence (>40%) in at least one habitat from each of the three environments (host,

aquatic, soil). Conversely, specialists are genera with a high prevalence (>40%) in

one habitat and low prevalence (<5%) in every other. A relative abundance (RA) cut-

off of 0.01% was used for determining whether a genus was present in a sample.

Using this approach, we detected 48 bacterial generalists and 30 specialists (Figure

2A). To confirm our definition of generalists and specialists, we calculated Levins'

niche breadth indices. Generalists showed significantly higher Bn values compared

to specialists (Wilcoxon Rank Sum test, $p < 0.001$; Supplementary Figure 1). All

specialists and all generalists, with the exception of Christensenellaceae R7, were

above the detection limit and had a significant niche breadth signal after Benjamini-

Hochberg adjustment.[18]

We did not observe a clear phylogenetic origin for generalists or specialists (Chi

squared test, $p > 0.05$), indicating their roles as generalists and specialists evolved

independently. Each of the top five bacterial generalists were detected in more than

50% of the 1,580 samples. Among them, the most prevalent was *Pseudomonas* which

was detected in 52%, 70% and 89% of host, soil and aquatic samples respectively,

followed by *Bacillus* (33%, 71%, 35%) and *Bradyrhizobium* (17%, 73%, 35%)

(Supplementary table 2). Concerning the specialists, the most extreme bacterial

genera came from the soil. While *Gryllotalpicola* and *Anaerovibrio* were found in

>91% of biochar samples, the prevalence dropped to 0.1% on average for non-soil

environments (Supplementary table 2). Specialists were also found in host- and

aquatic-associated environments. For example, *Acetatifactor* was found in 80% of

samples from the murine gut, but had a prevalence less than 3% in all other habitats.

The genus *Leptospira* was found in 83% of samples of the Cuyahoga River, but had a

prevalence less than 2% in all other habitats. Interestingly, when comparing the

relative abundance of generalists and specialists, we observed that both bacterial

and fungal generalists had a significantly higher abundance (Figure 2B, Wilcoxon

Rank Sum test, $p < 0.001$). This pattern remained even when we used stricter and

looser thresholds to define generalists and specialists (Supplementary Figure 2).

This finding confirms the pattern observed above (Figure 1G & H), suggesting that

independently of how groups are defined, genera that can colonize diverse

environments are usually able to outcompete niche-specific genera.

When looking at the fungal kingdom, the number of generalists was much lower and

only *Aspergillus*, *Malassezia*, *Aureobasidium*, and *Cortinarius* satisfied the criteria of a

generalist (Figure 2A). Among these, *Aspergillus* had the highest overall prevalence

among all samples with 38%, 52% and 12% in the host, soil, and aquatic samples,

respectively. From the 19 fungal specialists, *Chrysanthotrichum* and

Mycocentrospora were the most niche-specific, with prevalences of 68% and 48% in

temperate and conifer forests, but a mean prevalence of ≤0.1% in all other habitats.

Only two of the 19 fungal specialists (11%) originated from outside soil

environments (*Vuilleminia* and *Seimatosporium*). As with bacterial genera, the

relative abundance of fungal generalists was significantly higher than that of fungal

specialists (Figure 2C, Wilcoxon Rank Sum test, $p < 0.001$).

To gain insight into the ecological lifestyle that allows generalists to dominate

microbial communities in abundance, even in diverse environments, we further

explored their intra and inter-kingdom interaction patterns. Using the 1,188 bacteria

and 184 fungal genera detected in all three environments, we constructed individual

co-abundance networks for soil, host, and aquatic environments (see Methods for

details). Despite only using genera commonly detected in all three environments, the

topological characteristics of the networks were highly distinct, as measured by

significant differences in betweenness and Kleinberg's hub node centrality scores

(Wilcoxon test, $p < 0.00$; Supplementary Figure 3). Despite the differences in

topology, we could still compile subnetworks of inter- and intra-kingdom

correlations found jointly among host-soil, host-aquatic and/or soil-aquatic

environments. Strikingly, 45 of the 48 bacterial generalists and all 4 fungal

generalists were part of those subnetworks, which are characterized by a higher

number of positive than negative edges (Figure 2D). The ratio of positive to negative

edges was higher in correlations involving a generalist (2.5) compared to all other

edges (2.2). When we looked for interactions between genera found in all three

environments, we identified 43 such edges that all represented positive interactions

between bacteria and included 21 generalists. These findings suggest that the

success of generalists in colonizing diverse environments and achieving high

abundances may be attributable to their ability to avoid competitive and elicit

positive interactions with other highly prevalent genera.

Bacterial generalists exert a strong influence on the intra- and inter-

kingdom community structure

We subsequently explored whether the presence of generalists and specialists had

an impact on the diversity of a community. Interestingly, alpha diversity, as

measured as Chao 1 and Shannon, was significantly lower in samples where no

generalist was detected compared to samples with generalists present for both

bacterial (Figure 3A) and fungal (Figure 3B) communities (Permutation test of

samples lacking any of the N generalists compared with samples lacking any N

random taxa, $10*10^3$ permutations, $p < 0.03$). Conversely, the impact of specialists

on alpha diversity in their specific habitat was much less profound and varied by

habitat without a clear trend (Supplementary Figure 4).

We subsequently shifted our focus to inter-kingdom interactions which are often

overlooked in microbial ecology studies and examined bacterial generalists for a role

in shaping the mycobiome community structure and *vice versa*. As expected, we

observed a significant separation between the soil, host, and aquatic micro- and

mycobiome beta diversity by Bray-Curtis dissimilarity (Figure 3C, PERMANOVA, p <

0.001 for both bacteria and fungi). Constrained ordination revealed a significant,

linear relationship between bacterial Bray-Curtis dissimilarity and fungal

community composition and *vice versa* (dbRDA, ANOVA p < 0.04 for all explanatory

genera of the other kingdom in a multivariate model). Bacteria genera could explain

an extensive part of the mycobiome variation observed in the three environments

with a partial R2 of 25% by dbRDA. Of the bacterial genera, *Conexibacter*, *Bacillus*,

and *Lysobacter* had the highest explanatory power on mycobiome variation (Fig 3D).

Interestingly, six out of the top ten explanatory bacteria genera in the dbRDA were

generalists. Similarly, fungal genera explained 26% of the microbiome variation

between host, soil, and aquatic samples, with *Mortierella*, *Trichocladium*, and

Candida having the highest explanatory power (Fig 3D). Among the top ten

explanatory genera was one of the four fungal generalists - *Malassezia*.

Altogether, our analysis indicates that bacterial and fungal generalists are attractive

candidates for re-structuring microbiome communities given that they can reach

high abundances, they contribute positively to the taxonomic (alpha) diversity of

their kingdom- an ecological characteristic often associated with healthy

environments, and they can also contribute to shaping cross-kingdom microbial

structures.

Discussion

Microbiome engineering is a fast-growing field that aims to improve ecosystem

function through the manipulation of its microbial composition. Microbiome

engineering has had several significant successes in human health, including in the

infant gut,[19,20] bioremediation,[21] wastewater,[22] and agriculture.[23] However, the

discipline still faces marked challenges - namely that inoculants often fail to

establish or modify ecosystem functioning over extended periods of time.[24] These

failures are likely due to microbial dispersal, as well as environmental and biotic

constraints to organism establishment.[25] A widely-accepted ecological framework

posits that inoculant success is determined by its dispersal potential, compatibility

with the new environment, and the range of interactions that occur between the

introduced and resident organisms. Given the broad range of potential barriers,

complementing existing microbiome engineering solutions with an ecology-

motivated decision tree that prioritizes inoculants is an urgent priority.[25]

With this in mind, we performed a global survey of host, aquatic, and soil

microbiomes to reveal ecological properties of bacterial and fungal genera that may

promote or deter their establishment in an environment and/or restore the richness

and diversity of microbial communities in diseased ecosystems. The metagenomic

analysis of >1,580 paired host, soil, and aquatic microbiomes and mycobiomes

identified approximately 3,000 bacterial and 1,700 fungal genera – highlighting the

daunting task of selecting organisms as potential engineering solutions. From this,

we used the concept of generalists and specialists to limit the selection space for

microbial community engineering.

While the concept of generalists and specialists in ecology is not new, it has mostly

been applied in specific habitats[26–30] and not on a global scale such as this. Moreover,

it has rarely included eukaryotic organisms such as fungi[27] and has not been used

towards a cross-kingdom understanding of the biotic interactions shaping microbial

communities. We demonstrate that many of the important barriers in microbiome

engineering are more likely to be overcome when selecting inoculants from a short

list of 48 bacterial and 4 fungal generalists. Besides their global presence, these

genera reach significantly higher abundances and contribute positively to the

richness and diversity of their respective kingdom. Moreover, six bacterial

generalists, including *Bacillus*, *Lysobacter*, *Escherichia* and *Gemmatimonas* and one

fungal generalist, *Malassezia*, harbor additional ecological properties and appear to

play a significant role in shaping cross-kingdom microbial composition (Figure 3D).

In summary, we put forward a strategy to improve microbiome engineering, with a

focus on microbial establishment and community diversity in both the bacterial and

fungal kingdoms. Our findings on the environment-agnostic ability of specific

bacterial and fungal genera to reach sufficient colonization levels to impact the

ecosystem function can be integrated with other properties (e.g. ease of cultivation,

genetic engineering) in a decision tree framework to accelerate microbiome

engineering solutions. The challenge ahead will be understanding the functional

characteristics that differentiate generalists from other microbes and dictates their

ability to flourish in highly diverse environmental niches. While for bacteria

generalists this can be likely explored by deep functional characterization at the

pathway and enzyme level using shotgun metagenomics datasets, the functional

characterization of fungal generalists will be a much greater challenge. Additionally,

care will need to be used when selecting microbes as the ability of generalists to

thrive in diverse environments is also a hallmark of many opportunistic

pathogens.[31,32] Nevertheless, we believe large-scale computational analyses

combined with multifactorial experiments in cross-disciplinary approaches will be

able to overcome these challenges and accelerate microbiome engineering solutions.

Materials and Methods

Sample Selection

Included studies were retrieved by querying NCBI BioProject with the terms

'bacteria' and 'fungi' in any field. Only Biosamples with both 16S rRNA and ITS

amplicon sequencing data were considered for the concurrent analysis of both

kingdoms. We used both the identifier and attributes of the biosample, such as

aliases and library names, to map fungal and bacterial read files to a sample using a

custom script. Samples were associated to an environment (aquatic, host, or soil)

using manual curation of associated publications and biosample attributes provided

by the depositor. The three environments were further subdivided into 17 habitat

groups based on the body part and/or the ecoregion of the sampling location for

host and other samples, respectively (Olson et al., 2001). Habitats with less than five

samples were pooled together.

Generation of Genus-level Abundance Profiles

Genus-level abundance profiles were calculated using a custom nextflow pipeline (Di

Tommaso et al., 2017). Briefly, reads were downloaded from NCBI SRA using

grabseqs, except for the American Gut Project, which was downloaded from Qiita

(Gonzalez et al., 2018; Taylor et al., 2020). Paired-end reads were merged using

NGmerge (Gaspar, 2018). Quality Control (QC) and adapter removal was performed

using trimmomatic with a minimum Phread quality of 20 and a minimal read length

of 100 (Bolger et al., 2014). Quality was assessed using FastQC and MultiQC (Ewels

et al., 2016). Subsequent steps were performed using QIIME2 (Bolyen et al., 2019).

Reads were dereplicated following closed-reference OTU picking for both kingdoms

separately using VSEARCH with a 97% identity threshold (Rognes et al., 2016). For

taxonomic annotation, SILVA 132 97% consensus and UNITE 8.2 dynamic databases

were used for bacteria and fungi, respectively (Nilsson et al., 2019; Quast et al.,

2013). Following quality control, a total of 1,580 samples were selected for

downstream analyses.

Workflow and Statistical Analysis

310 Analyses were performed using a custom drake pipeline (Landau, 2018) built using

311 the programming language R 4.0.2. Briefly, abundances obtained from OTU profiling

312 were total-sum-scaled (TSS) and pooled at genus rank. All tools were used with

313 default parameters if not explicitly specified.

Diversity

Alpha diversity was estimated using Shannon and Chao1 metrics with the phyloseq

and vegan packages (Dixon, 2003; McMurdie and Holmes, 2013). To quantify the

contributions of a bacterial community profile with the fungal one and *vice versa*, we

used linear and unsupervised Canonical Correlation Analysis, as implemented in the

function CCorA of the vegan R package (Dixon, 2003). P-values were obtained using

blocked permutations to control for the habitat and to reduce assumptions of the

test. Supervised constrained ordination was performed using stepwise Distance-

based Redundancy Analysis (dbRDA) adapted from (Vieira-Silva et al., 2020). This

analysis shows linear relationships between bacterial dissimilarities and

abundances of selected explanatory fungal genera (and *vice versa*). An optimal

subset of up to 50 explanatory genera of the other kingdom was computed using a

stepwise feed-forward approach, as implemented in the ordistep function of the

vegan R package (Dixon, 2003).

Co-abundance Networks

SparCC, as implemented in FastSpar, was used to assess correlation between taxa pairs for each environment separately.[33,34] Both kingdoms were pooled together, allowing for the identification of interkingdom correlations. Only genera found in all three environments were considered for pairwise correlation. Node topology metrics were calculated using the R package igraph.

Generalists and Specialists

Taxa were defined as generalists if they were found in at least 40% of samples in at least one habitat from each environment (host, soil, aquatic) with a relative abundance of at least 0.01%. Complementary, taxa were defined as specialists if they were found in at least 40% of samples in one habitat and less than 5% of samples in all other habitats using the same abundance threshold as for generalists. Levins' niche breadth index was calculated as implemented in the R package MicroNiche.

Code and Data Availability

Scripts created for data processing and statistical analysis are available at https://github.com/bioinformatics-leibniz-hki/its-16s. Raw sequence data can be downloaded from any International Nucleotide Sequence Database Collaboration (INSDC) server using accessions as provided in Supplemental Table 2 and the git repository.

References

1.   Bar-On, Y. M., Phillips, R. & Milo, R. The biomass distribution on Earth. *Proc Natl Acad Sci U S A* **115**, 6506–6511 (2018).

2.   Widder, S. *et al.* Challenges in microbial ecology: building predictive understanding of community function and dynamics. *ISME J* **10**, 2557–2568 (2016).

3.   Sunagawa, S. *et al.* Ocean plankton. Structure and function of the global ocean microbiome. *Science* **348**, 1261359 (2015).

4.   Oak, S. J. & Jha, R. The effects of probiotics in lactose intolerance: A systematic review. *Crit Rev Food Sci Nutr* **59**, 1675–1683 (2019).

5.   Mohajeri, M. H. *et al.* The role of the microbiome for human health: from basic

science to clinical applications. *Eur J Nutr* **57**, 1–14 (2018).

6.   Holscher, H. D. Dietary fiber and prebiotics and the gastrointestinal

microbiota. *Gut Microbes* **8**, 172–184 (2017).

7.   Fang, H., Kang, J. & Zhang, D. : a review and future perspectives. *Microb Cell*

Fact **16**, 15 (2017).

8.   Rossi, M., Amaretti, A. & Raimondi, S. Folate production by probiotic bacteria.

Nutrients **3**, 118–134 (2011).

9.   Conly, J. M. & Stein, K. The production of menaquinones (vitamin K2) by

intestinal bacteria and their role in maintaining coagulation homeostasis. *Prog Food*

Nutr Sci **16**, 307–343 (1992).

10.   Souak, D. *et al.* Challenging Cosmetic Innovation: The Skin Microbiota and

Probiotics Protect the Skin from UV-Induced Damage. *Microorganisms* **9**, (2021).

11.   Singh, R. *et al.* Donor feces infusion for eradication of Extended Spectrum

beta-Lactamase producing Escherichia coli in a patient with end stage renal disease.

Clin Microbiol Infect **20**, O977–978 (2014).

12.   Heijden, M. G. A. van der, Martin, F. M., Selosse, M. A. & Sanders, I. R.

Mycorrhizal ecology and evolution: the past, the present, and the future. *New Phytol*

205, 1406–1423 (2015).

13.   Vannier, N., Agler, M. & Hacquard, S. Microbiota-mediated disease resistance

in plants. *PLoS Pathog* **15**, e1007740 (2019).

14.   Khoruts, A. & Sadowsky, M. J. Understanding the mechanisms of faecal

microbiota transplantation. *Nat Rev Gastroenterol Hepatol* **13**, 508–516 (2016).

15.   Li, J. *et al.* Probiotics modulated gut microbiota suppresses hepatocellular

carcinoma growth in mice. *Proc Natl Acad Sci U S A* **113**, E1306–1315 (2016).

16.   Oku, S., Komatsu, A., Nakashimada, Y., Tajima, T. & Kato, J. Identification of

Pseudomonas fluorescens chemotaxis sensory proteins for malate, succinate, and

fumarate, and their involvement in root colonization. *Microbes Environ* **29**, 413–419

(2014).

17.   Carvalhais, L. C. *et al.* Activation of the jasmonic acid plant defence pathway

alters the composition of rhizosphere bacterial communities. *PLoS One* **8**, e56457

(2013).

18.   Feinsinger, P., Spears, E. E. & Poole, R. W. A simple measure of niche breadth.
Ecology **62**, 27–32 (1981).
19.   Frese, S. A. *et al.* EVC001 in Breastfed Infants. *mSphere* **2**, (2017).
20.   Panigrahi, P. *et al.* A randomized synbiotic trial to prevent sepsis among
infants in rural India. *Nature* **548**, 407–412 (2017).
21.   Azubuike, C. C., Chikere, C. B. & Okpokwasili, G. C. Bioremediation techniques-
classification based on site of application: principles, advantages, limitations and
prospects. *World J Microbiol Biotechnol* **32**, 180 (2016).
22.   Henze, M., Gujer, W., Mino, T. & Loosdrecht, M. C. van. Activated sludge
models ASM1, ASM2, ASM2d and ASM3. *IWA Scientific and Technical Report* **9**,
(2000).
23.   Orozco-Mosqueda, M. D. C., Rocha-Granados, M. D. C., Glick, B. R. & Santoyo, G.
Microbiome engineering to improve biocontrol and plant growth-promoting
mechanisms. *Microbiol Res* **208**, 25–31 (2018).
24.   Kaminsky, L. M., Trexler, R. V., Malik, R. J., Hockett, K. L. & Bell, T. H. The
Inherent Conflicts in Developing Soil Microbial Inoculants. *Trends Biotechnol* **37**,
140–151 (2019).
25.   Albright, M. B. N. *et al.* Solutions in microbiome engineering: prioritizing
barriers to organism establishment. *ISME J* **16**, 331–338 (2022).
26.   Liao, J. *et al.* The importance of neutral and niche processes for bacterial
community assembly differs between habitat generalists and specialists. *FEMS*
Microbiol Ecol **92**, (2016).
27.   Liu, L., Wang, S. & Chen, J. Transformations from specialists to generalists
cause bacterial communities are more stable than micro-eukaryotic communities
under anthropogenic activity disturbance. *Sci Total Environ* **790**, 148141 (2021).
28.   Garrison, C. E. & Field, E. K. Introducing a "core steel microbiome" and
community functional analysis associated with microbially influenced corrosion.
FEMS Microbiol Ecol **97**, (2020).
29.   Walter, J. & Ley, R. The human gut microbiome: ecology and recent
evolutionary changes. *Annu Rev Microbiol* **65**, 411–429 (2011).
30.   Chen, Y. J. *et al.* Metabolic flexibility allows bacterial habitat generalists to
become dominant in a frequently disturbed ecosystem. *ISME J* **15**, 2986–3004
(2021).

31.   Zajc, J., Gostinčar, C., Černoša, A. & Gunde-Cimerman, N. Stress-Tolerant

Yeasts: Opportunistic Pathogenicity Versus Biocontrol Potential. *Genes (Basel)* **10**,

(2019).

32.   Gostinčar, C. *et al.* Fungi between extremotolerance and opportunistic

pathogenicity on humans. *Fungal diversity* **93**, 195–213 (2018).

33.   Friedman, J. & Alm, E. J. Inferring correlation networks from genomic survey

data. *PLoS Comput Biol* **8**, e1002687 (2012).

34.   Watts, S. C., Ritchie, S. C., Inouye, M. & Holt, K. E. FastSpar: rapid and scalable

correlation estimation for compositional data. *Bioinformatics* **35**, 1064–1066 (2019).

 Figures

*Figure 1: A global analysis of microbial communities reveals differences in environmental specificities between bacteria and fungi. Distribution of samples used in this study (N=1,580) shown over time (**A**) and geographic location (**B**). **C:** Rarefaction curves of Shannon alpha diversity for each study demonstrate sufficient sampling depth. Curves are shown as LOESS regressions from 10 independent sampling trials at 10 given sampling subset sizes. Lines are colored by environment and are surrounded by ribbons indicating the 95% confidence interval across the trails. **D:** Intersection of bacterial and fungal genera found in at least one sample in each environment as Venn diagrams. **E, F**: Percentage of genera found in all three or only one environment. **G, H**: Abundance comparisons of common and unique genera. A genus was considered present in a sample using a threshold of abundance > .01%. Significance determined by Wilcoxon rank sum test; *** denotes p < 0.001.*

Figure 2: Abundance and co-abundance of generalists and specialists. A:

Abundance heatmap of bacteria and fungal genera. Samples are arranged by

environment (top bar) with only abundance values >0.01% displayed. Right bars

indicate phylum and prevalence among all samples, where only taxa with >5%

prevalence are displayed. Generalists (>40% prevalence in at least one habitat in all

three environments) and specialists (>40% prevalence in one habitat and <5%

prevalence in all others) are annotated in blue and yellow, respectively. The

dendrogram was calculated using average linkage of Manhattan distances between

taxa abundance profiles. **B, C**: Abundances of bacterial (**B**) and fungal (**C**) generalists

and specialists. Values were averaged by project to account for different cohort sizes.

Statistical significance calculated using Wilcox rank-sum test (*** denotes $p<0.001$).

D: Networks of genera found in all three environments and significantly co-abundant

in the majority of environments (SparCC FDR $p < 0.05$, $|r| > 0.2$)

Figure 3: Generalists significantly impact diversity and cross-kingdom variation.

A,B: Shannon and Chao1 alpha diversity were calculated for bacteria (A) and fungi (B).

Samples were grouped by whether they contained any generalist (genera with >40%

prevalence in at least one habitat from every environment; abundance > 0.01%), or

not. Significance bars indicate permutation test compared to samples without random

taxa instead of generalists (q<0.05, ** q<0.01, *** q<0.001). C-D: Bacterial and fungal*

Bray-Curtis dissimilarities constrained by explanatory genus abundances of the other

kingdom using distance-based Redundancy Analysis (dbRDA). C: Explaining genera

were selected using a feedforward approach. Effect size of most explanatory taxa is

shown in D by multivariate model (as displayed in C) or univariate model containing

only the taxon of interest. Generalists are indicated with blue text. Stars indicate

*significance by ANOVA (** p<0.01).*