Relationship between thyroid hormones and associated metabolites and gene expression bioindicators in the serum of *Rana [Lithobates] catesbeiana* tadpoles and frogs during metamorphosis

Rikke Poulsen [1,3,*], Shireen H. Jackman[2], Martin Hansen[1], Caren C. Helbing[2]

[1] Department of Environmental Science, Aarhus University, Frederiksborgvej 399, 4000 Roskilde,
Denmark
[2] Department of Biochemistry and Microbiology, University of Victoria,Victoria, BC V8P 5C2,
Canada
[3] Department of Biology, University of Southern Denmark, 5230 Odense M, Denmark

* Correspondence:
Rikke Poulsen
rikkepoulsen@envs.au.dk

Keywords: thyroid hormone, amphibian metamorphosis, *Rana [Lithobates] catesbeiana*,
bioindicators, untargeted metabolomics, gene expression, *Xenopus* comparison

Abstract

Anuran metamorphosis is characterized by profound morphological changes including remodeling of
tissues and organs. This transition is initiated by thyroid hormones (THs). However, the current
knowledge of baseline levels of THs during metamorphosis relies on pooled samples using methods
known for high variability with little reporting of measured variation. Moreover, establishing a clear
linkage between key gene expression bioindicators and TH levels throughout the metamorphic
process is needed. Using state-of-the-art ultra-high performance liquid chromatography isotope-
dilution tandem mass spectrometry, we target 12 THs -and metabolites in the serum of *Rana*
[Lithobates] catesbeiana (n=5-10) across seven distinct postembryonic stages beginning with
premetamorphic tadpoles (Gosner stage 31-33) and continuing through metamorphosis to a juvenile
frog (Gosner stage 46). TH levels are then related to TH-relevant gene transcripts (*thra*, *thrb*, and
thibz) in back skin of the same individual animals. Significant increases from basal levels were
observed for thyroxine (T4) and 3,3',5-triiodothyronine (T3) at Gosner stage 41, reaching maximal
levels at Gosner stage 44 (28 ± 10 and 2.3 ± 0.5 ng/mL, respectively), and decreasing to basal levels in
juvenile frogs. In contrast, 3,5-diiodothyronine (T2) increased significantly at Gosner stage 40 and
was maintained elevated until stage 44. While *thra* transcript levels remained constant and then
decreased at the end of metamorphic climax, *thrb* and *thibz* were induced to maximal levels at
Gosner stage 41, followed by a decrease to basal levels in the froglet. This exemplifies the exquisite
timing of events during metamorphosis as classic early response genes are transcribed in anticipation
of peak TH concentrations. Interestingly T2 showed a distinct concentration profile suggesting a
specific biological role of this biomolecule in anuran postembryonic development. This finding also
underlines the sensitivity of the developmental process, and it is, indeed, well-known that external
factors such as anthropogenic chemicals can disrupt anuran metamorphosis. Hence, as a second aim
of the study, we set out to find additional bioindicators of metamorphosis, which can aid future
investigations of developmental disruption. Using a sensitive nanoLC-Orbitrap system an untargeted
analysis workflow was applied. Among 6,062 endogenous metabolites, 421 showed metamorphosis-

dependent concentration dynamics. These potential bioindicators included several carnitines,
prostaglandins and some steroid hormones.

1   Introduction

Thyroid hormones (THs) are essential for the normal growth, development and health of all
vertebrates (1–3) Among many other biological processes, THs control metabolism, cell
differentiation and migration and also play a pivotal role for e.g., neurodevelopment (3,4). One of the
most remarkable examples of TH action is anuran metamorphosis where a tadpole transitions into a
frog with extensive reprogramming and reorganization of organs and tissues (5–8). This remodeling
has direct parallels to the developing human during the last trimester and the first weeks of life where
a surge of THs is essential for proper development (8–10). Hence, anuran metamorphosis is an
excellent model system to study developmental endocrinology and the dynamics of the TH system.
This has been acknowledged in several publications (5,7–9,11). The potential of amphibian
metamorphosis as a screening tool for chemical disruption of the hypothalamus-pituitary-thyroid axis
has furthermore been recognized in the OECD guideline 231 for the testing of chemicals i.e., the
Amphibian Metamorphosis Assay (12).

Despite this, the current knowledge of baseline levels of THs during metamorphosis largely relies on
pooled samples using methodologies known for high variability with sparse reporting of measured
variation (13–23)(Table 2). In fact, (radio)immunoassays are still one of the most commonly used
methods for quantification of THs despite known limitations, e.g., cross-reactivity and interference
from proteins as summarized by Spencer (24). The current study uses a highly robust, selective and
sensitive liquid chromatography tandem mass spectrometry (LC-MS/MS) method based on isotopic
dilution to quantify 12 THs and metabolites (16,25).

THs exert their chemical messenger effect mainly through nuclear TH receptors, although growing
evidence shows that non-genomic signaling also plays a role (8,26). In the genomic pathways,
nuclear TH receptors (THRs) bind directly to DNA or indirectly by interacting with chromatin
associated proteins to modulate expression levels of hormone-responsive genes (26). Two THR
genes, α and β, code for these receptors. Both have been found to show specific expression patterns
according to cell type and developmental stage (27). In addition to TH concentrations, the
transcription levels of THR genes therefore serve as robust indicators of TH signaling and the
endocrinological processes during metamorphosis. Earlier studies have shown that an abundance of
transcripts encoding THR α and β (*thra*, *thrb*) as well as TH-induced bZip protein (*thibz*, Fig. 1F)
show high correlation with developmental stage in anuran tadpoles (27–29) indicating their potential
to serve as bioindicators of metamorphosis in addition to THs. However, these data and their
relationships to changes in TH levels *in the same subjects* have not been established in amphibians.

American bullfrog (*Rana [Lithobates] catesbeiana*) is a great alternative model species as it is a
representative of the largest amphibian group, Ranids or "true frogs", and is found worldwide. Their

large-sized tadpoles enable the analysis of individual animals rather than pooled samples.
Furthermore, this amphibian compared to e.g., *Xenopus laevis* tadpoles, also has a greater similarity
to human life stages as they completely transition from aquatic to terrestrial living (*Xenopus* remain
fully aquatic as adults). By using wild caught animals instead of laboratory manipulated individuals,
we ensure that the natural variation of endocrinological state is represented. By using a high sample
replication we also capture the natural biological variation without losing statistical power.

Although anuran metamorphosis is initiated solely by THs, many other biochemical pathways
undergo extensive remodelling during this transition (30) and the regulation of metamorphosis is a
complex process of interacting mechanisms (11). The complexity, tight regulation, and
interconnection of biological networks and pathways can be conceptualized using untargeted
metabolomics. In contrast to traditional hypothesis-driven investigations, untargeted approaches such
as metabolomics interrogate the whole biological system and let the organism-response guide the
investigations (31–35). Such knowledge is very powerful when trying to make new discoveries about
a biological system and can aid the search for new robust biological markers. In the case of anuran
metamorphosis, this type of analysis provides a holistic view of the biological processes and allows
for a search of robust bioindicators of the molecular dynamics during the transitions. Such
bioindicators are useful for assessing the state of the endocrine system and its disruption by external
environmental factors (such as endocrine disrupting chemicals or climate change (i.e., temperature)).

The first aim of the present study is to map the dynamics of known bioindicators, ie. TH
concentration and transcription of TH sensitive genes in wild-caught *R. catesbeiana* tadpoles and
frogs during metamorphosis. This includes the quantification of measurable THs and metabolites in
serum as well as abundance of gene transcripts of *thra, thrb* and *thibz* in back skin of the same
animals. As a second aim, we use untargeted metabolomics to reveal new bioindicators of
metamorphosis.
105

106 2   Materials and methods

107 2.1   Sampling

Rana [Lithobates] catesbieana tadpoles of unknown sex were caught locally in Victoria, British
Columbia, Canada and housed in the University of Victoria Outdoor Aquatics Unit. Animals were
maintained with a 12 h light/12 h dark photoperiod in 340 L fiberglass tanks containing 15°C
recirculated and dechlorinated municipal water and fed daily with Spirulina flakes (Dynamic Aqua
Supply, Surrey, BC, Canada). Treatment of animals was in accordance with the guidelines of the
Canadian Council on Animal Care and experimental protocol #2015-028 was approved by the
University of Victoria Animal Care Committee.
115
Tadpoles and recently metamorphosed froglets were euthanized at distinct postembryonic
developmental stages (36). The mean ± standard error weights and number of animals are shown in
Table S1.(36). Animals were euthanized in 0.1% (w/v) tricaine methanesulfonate (Syndel
Laboratories Ltd., Vancouver, BC, Canada) buffered with 25 mM sodium bicarbonate in
dechlorinated municipal water. Blood was collected by tail muscle incision for tadpoles and severing

of the aorta for froglets. The collected blood was allowed to coagulate at room temperature for 10-15
min. The blood from each animal was kept separate and never pooled. The coagulated blood was
centrifuged at 10,000 xg for 10 minutes at 4°C, the serum was collected, immediately flash frozen in
liquid nitrogen and stored at -80°C until use. On average, 150 µL serum per individual animal was
collected using this procedure.
126
The back skin from the same animals was harvested from each individual tadpole or frog from
between the eyes to the tip of the rostrum to permit the most consistent comparison of skin tissue
across developmental stages. The skin was immediately immersed in RNAlater (Invitrogen, Thermo
Fisher Scientific) and stored at -20°C until RNA extractions were performed.

2.2   Gene expression bioindicators

RNA extraction, cDNA synthesis, and quantitative real time polymerase chain reaction (qPCR)
analyses were performed as previously described (37). Three normalizer gene transcripts encoding
ribosomal protein L8 (*rpl8*), ribosomal protein S10 (*rps10*), and elongation factor 1 alpha (*eef1a*)
were used, and three classical TH-responsive transcripts were analysed encoding thyroid hormone
receptor α (*thra*), thyroid hormone receptor β (*thrb*), and thyroid hormone induced basic region
leucine-zipper containing transcription factor (*thibz*). The exact run conditions and primer sequences
have been previously reported (29,37,38). RNA extraction, cDNA synthesis, and quantitative real
time polymerase chain reaction (qPCR) analyses were performed as previously described (37). Three
normalizer gene transcripts encoding ribosomal protein L8 (*rpl8*), ribosomal protein S10 (*rps10*) and
elongation factor 1 alpha (*eef1a*) were used, and three classical TH-responsive transcripts were
analysed encoding thyroid hormone receptor α (*thra*), thyroid hormone receptor β (*thrb*), and thyroid
hormone induced basic region leucine-zipper containing transcription factor (*thibz*). The exact run
conditions and primer sequences have been previously reported (29,37,38).
145

2.3   TH and metabolite extraction

Serum samples were extracted for targeted TH determination as previously described (39) and
untargeted metabolomics using the same extract. In brief, 50 µL serum was spiked with isotopic-
labelled (^{13}C)-thyroid hormone standards (internal standards (IS), cT2, cT3 and cT4) and
homogenized. After antioxidant treatment (100 µL, 25 mg/mL ascorbic acid, R,R-dithiothreitol and
citric acid solution) and protein denaturation by urea (8 M in 1% ammonium hydroxide), the samples
were enriched using solid-phase micro-extraction (SOLAµ HRP 10 mg/1 mL 96 well plate,
ThermoFisher Scientific, Germany) and reconstituted in 100 µL 5% methanol containing an
instrument control standard (ICS, crT3). Procedural blanks containing water instead of plasma were
included from the beginning. Detailed descriptions of chemicals and reagents are found in the
supplementary material (S1.1).

2.4   TH quantification

The targeted analysis of THs was performed on an Agilent 6495c triple-quadrupole system with a
hyphenated Agilent 1290 Infinity II ultra-high performance liquid chromatography (UHPLC) system
(binary pump, degasser, and autosampler; Agilent Technologies) as previously described (39).
Targeted analytes were thyroxine (T4), 3,3',5-triiodothyronine (T3), 3,3',5'-triiodothyronine (rT3),
3,5-diiodothyronine (T2), 3,3'-diiodothyronine (3,3'-T2), 3-iodothyronine (T1), thyronine (T0), 3-
iodothyronamine (T1Am), 3-iodothyroacetic acid (T1Ac), 3,5-Diiodothyroacetic acid (Diac),
triiodothyroacetic acid (Triac) and tetraiodothyroacetic acid (Tetrac). LoDs as fmol on column for all

THs were 4.0 (T4), 0.79 (T3), 0.76 (rT3), 0.56 (T2), 0.60 (3,3'-T2), 0.74 (T1), 0.23 (T0), 0.05 (T1Am), 9.4 (T1Ac), 15 (Diac), 10 (Triac), 9.6 (Tetrac). Neat standard ten-point equimolar calibration curves (0.04 – 20.0 pmol/mL TH, n=2) were prepared in 5 % methanol and all vials contained a fixed amount IS and ICS (15.2 pmol/mL). Data analysis was conducted in MassHunter version 10.1 (Agilent Technologies). Detectable hormones in our samples were T4, T3, T2, T1Am and T0. Statistical analysis and visualization of the gene expression biomarkers and TH concentrations were done using R ver. 3.6.3 (40). The univariate statistical analysis was performed by Kruskal-Wallis test followed by pairwise comparison by Dunn's test (41). To allow for statistical comparisons, TH levels below the limit of detection (LoD) were replaced by ½ LoD. Data between the LoD and limit of quantification (LoQ) remained as they were. The data was visualized using ggplot2 (42). Principal component analyses (PCA) were done using prcomp and visualized by ggbiplot (43).

2.5   Untargeted metabolomics

The untargeted analysis was performed on a nanoflow UHPLC Orbitrap mass spectrometer system (ThermoFisher Scientific) with a preconcentration trap-column setup described elsewhere (33). The instrument was operated in data-dependent mode by automatically switching between MS and MS/MS fragmentation. Samples were analyzed in randomized order and a pooled sample was injected in between every five samples to correct for any instrumentational fluctuations and monitor any systematic errors. Quality and validity of the analysis was confirmed by principal component analysis (PCA) showing that the composite quality control (QC) samples were centrally located in the plot (Fig. S2). Compound Discoverer (CD) software version 3.3 (Thermo Scientific) was used for data processing and analysis as previously described (33).

A univariate analysis (one-way ANOVA with Tukey as post-hoc test and p-values adjusted by Benjamin-Hochberg algorithm) was performed in Compound Discoverer (CD) to filter out compounds of particular interest to differentiate between developmental stages. Using R ver. 3.6.3, abundance of selected compounds was visualized in a heatmap using the R-package pheatmap (44) with Euclidian clustering on rows and no clustering on columns. A multivariate analysis by sparse partial least squares discriminant analysis (sPLS-DA) was performed on the 421 selected compounds using the R-package mixOmics (45) in R ver. 4.3.1 (40). Putatively annotated compounds were based on comparisons of the experimental MS2-spectrum with in-house spectral libraries (annotation level 1) and the online spectral library, mzCloud, (level 2)(46). Putatively characterized compound classes (level 3) were based on assigned predicted composition, *in silico* fragmentation by SIRIUS4 (47), and plausible, tentative candidates to that composition. For instance, whether the compound was likely to naturally occur in tadpole serum. Compounds that did not meet these criteria were annotated to level 4 and not incorporated in the functional analysis. The functional analysis was based on the Kyoto Encyclopedia of Genes and Genomes (KEGG) database(48), BioCyc(49), and human metabolome database (HMDB)(50).

3   Results

3.1   Thyroid hormone profile and gene expression bioindicators

Of the targeted THs T4, T3, T2, T0 and T1Am were above the instrumental limit of detection. Metamorphosis-related concentration dynamics was observed for T4 (Fig. 1A), T3 (Fig. 1B) and T2 (Fig. 1C). The concentration of T0 and T1Am did not change significantly over the recorded

209 developmental window (Figure S1), except for a transient decrease in T0 around metamorphic climax
210 in stage 41-44 compared to premetamorphic stage 31-33 (Fig. S1A).
211
The metamorphosis related concentration pattern was similar for T4 and T3. Until Gosner stage 40,
T4 was below the limit of detection (LoD, 1.3 ng/mL), after which the concentration gradually
increased reaching a maximum of 28 ± 10 ng/mL at Gosner stage 44 (Table S2) resulting in a 20-fold
increase. In the froglets (Gosner 46), T4 was again below the detection limit. T3 was quantifiable in
all samples and, like T4, increased gradually during metamorphosis, reaching the highest
concentration at Gosner stage 44 (2.3 ± 0.5 ng/mL) before decreasing to premetamorphic levels. The
difference between premetamorphosis and metamorphic climax was about 5-fold. T2 concentrations
were generally lower with higher variation (Figure 1A). However, in contrast to T3 and T4, the
concentration of T2 was already significantly higher at stage 40 compared to the initial pre-
metamorphic measurement in Gosner stage 31-33 animals (Dunn's post hoc test p-value =0.002).
This higher level was maintained at about 5-fold through metamorphic climax at Gosner stage 44
before returning to basal levels in the froglet.
224
Back skin gene expression of the thyroid hormone receptor β (*thrb*, Fig. 1E) and TH-induced bZip
protein-encoding transcripts (*thibz*, Fig. 1F) also showed the expected correlation with Gosner stage.
Both transcripts increased quickly in abundance from Gosner stage 40, peaked at Gosner stage 41,
and then gradually decreased to basal levels in the frog (Gosner stage 46). The change was
particularly noticeable for *thibz;* abundance increased around 500-fold compared to Gosner stage 40.
In comparison, the change in *thrb* was only 3-fold. Thyroid hormone receptor a (*thra*, Fig. 1D)
showed a stable abundance in these back skin samples until stage 44, where it decreased to a
significantly lower level compared to the premetamorphic group (Gosner stage 31-33). The fold
change was, however, minor.
234

Figure 1. Box plots showing serum concentration (ng/mL) of (A) T4, (B) T3, and (C) T2 and back skin gene expression of (D) thra, (E) thrb, and (F) thibz for the grouped Gosner stages. The lower and upper hinges correspond to the first and third quartiles. The whiskers extend from the hinge to the largest/smallest value no further than 1.5 times the inter-quartile range. Circles show individual replicates. Asterisks indicate significant difference from the premetamorphic Gosner stage 31-33 (Dunn's post hoc test, p-value <0.05).

As animal-specific TH quantification and gene expression was obtained, a multivariate analysis by PCA was performed to observe whether the developmental stages would cluster across the two principal components of the data (Fig. 2). Driven mainly by the concentration of T3 and T4 as well as the *thibz* and *thrb* transcript abundance there is a division along component 1 (55% explained variation) that separates the pre- and early prometamorphic stages (Gosner 31-40) plus the juveniles (Gosner 46) from the animals at the late prometamorphic stage and metamorphic climax (Gosner 41-44). The direction of the vectors represents the correlation with the principal component and the length represents the contribution to the component. The vector of T4 is nearly parallel to the X axis, indicating that the difference of PC1 is mainly due to the contribution of this TH. Furthermore, it is mainly *thra* abundance that drives the separation on principal component 2.

We also investigated the individual Spearman's rank correlations between gene transcript abundance and TH concentrations and found a strong and highly significant correlation for T4 and T3 versus *thibz* ($\rho>0.6$, p-value < 0.0001) and *thrb* ($\rho>0.4$, p-value < 0.001). T2 was also weakly correlated with *thrb* and *thibz* ($\rho>0.3$, p-value = 0.01). For all correlations see supplementary Table S3.

Figure 2. Principal component analysis (PCA) biplot for T4, T3, and T2 concentrations and gene transcript abundance of thra, thrb and thibz. Dots represents samples from individuals and colors represents Gosner stages.

3.2 Untargeted metabolomics

After filtering out background peaks from more than 200,000 features, 6,062 endogenous metabolites remained. The measured metabolites were dominated by compounds active in lipid metabolism including fatty acids, bile acids, and steroid hormone derivatives. A good coverage of arachidonic acid metabolism and linoleic acid metabolism was observed.

269
270

Figure 3. Heatmap (A) of 421 compounds that differed significantly in relative concentrations (ANOVA, adj p-value <0.05) between Gosner stage 44 and either stages 31-33 (premetamorphosis) or stage 46 (juveniles). Boxplots (B-G) showing examples of metabolites with their relative concentration plotted against the grouped Gosner stages. The positions of metabolites appearing in the heatmap with corresponding boxplots are outlined by red boxes. The relative concentration is found by normalizing peak areas to the average peak area for

Gosner 31-33. Asterisks indicate significant difference from the premetamorphic Gosner stage 31-33 (Dunn's post hoc test, p-value <0.05).

Based on the observation that T4 and T3 reached a maximum concentration at Gosner stage 44, and that this was significantly different from the concentration in premetamorphosis and in juveniles, we wanted to look for metabolites that showed similar concentration dynamics. A differential analysis was performed comparing Gosner stage 44 to premetamorphic stages (Gosner 31-39) and juveniles (Gosner 46). This resulted in 421 significant metabolites (p-value$_{adj}$<0.05). As exemplified by Figure 3A, most of them followed the curve shape of T4 (Fig. 1A) and T3 (Fig. 1B), including a gradual increase up to a maximum reached at metamorphic climax, after which the level dropped down again at the froglet stage. Examples of compounds are given to the right including arachidonic acid (Fig. 3B), a prostaglandin (Fig. 3C) and a hydroxyeicosatetraenoic acid (Fig. 3D), which are all related to the arachidonic acid metabolic pathway. The latter two, especially, show a pattern similar to the THs and exhibit high fold changes. In contrast to previous observations (30), arachidonic acid does not follow this trend in the data set. Rather, there is considerable variation in measurements with a substantial reduction in the juvenile (Fig. 3A). Dodecenoylcarnitine (Fig. 3E) and palmitoyl glutamic acid (Fig. 3F) are other examples of compounds that follow the pattern of the THs. Carnitines are involved in the transport of long chain fatty acids into mitochondria for beta-oxidation and energy generation. Palmitoyl glutamic acid is a palmitic acid amide of glutamic acid and belongs to the compound class known as N-acylamides.

All 421 compounds may be efficient bioindicators of metamorphosis and possibly for its disruption. An sPLS-DA analysis was performed to discover compounds that best discriminate between Gosner stages (Figure S3A). The first component explained 45% of the variation and metabolites, which contributed to this first component, mainly separated the prometamorphosis and metamorphic climax, where T4, T3, and T2 levels were maximal. After tuning the model, 20 variables (i.e., metabolites) were selected on component 1 and all the selected variables had the highest abundance in Gosner stage 44 animals (Fig. S3B). The compounds included dodecenoylcarnitine (Fig 3E) and decanoylcarnitine both identified with a spectral match (level 1). Table 1 shows 12 of these compounds, for which a chemical class could be assigned, including their stability (i.e., the proportion of cross validation folds (across repeats) where it was selected to be used for component 1). The details for all 20 compounds can be found in Table S4.

Table 1. Annotated metabolites (n=12) which contributed to the first component in the sPLS-DA analysis. Chemical class, ID level (46) and name (tentative for all ID-level<1) is given as well as their sPLS-DA stability score (i.e., the proportion of cross validation folds across repeats where it was selected to be used for component 1). Further details are provided in Table S4.

Chemical class	ID level	Name	sPLS-DA Stability score	Formula
Acyl carnitines	1	Dodecenoylcarnitine	0.83	C19 H35 N O4
Acyl carnitines	3	Undecylcarnitine	0.67	C18 H35 N O4
Acyl carnitines	3	(5E)-tetradecenoyl-L-carnitine	0.57	C21 H39 N O4
Acyl carnitines	1	Decanoyl-L-carnitine	0.4	C17 H33 N O4
Prostaglandin	3	Prostaglandin C1	0.37	C20 H32 O4
Linoleic acids and derivatives – Fatty acyls	3	9-OxoOTrE	1	C18 H28 O3

Linoleic acids and derivatives – Fatty acyls	3	2R-HpOTrE	0.43	C18 H30 O5
Linoleic acids and derivatives	3		0.5	C18 H28 O4
Long-chain fatty acid	3		1	C18 H28 O3
Long-chain fatty acid	3		0.77	C18 H28 O3
Medium-chain hydroxy acids and derivatives	3		0.47	C19 H32 O5
Medium-chain hydroxy acids and derivatives	3		0.87	C19 H32 O5

4   Discussion

The present study brings forth essential new knowledge of baseline levels of THs and TH-responsive genes during metamorphosis. Currently, published data largely rely on pooled samples, limited replication, and sparse reporting of variation (Table 2). In contrast, the present study includes individual sampling of 5-10 animals per developmental stage and the methodology reflects state-of-the-art quantification of biomolecules. This also includes the quantification of TH metabolites other than T3 and T4, such as T2, which provides novel insight into the thyroid hormone system. Table 2 summarizes maximal levels of THs (ng/mL) measured during anuran metamorphosis in published studies. The average reported value for all species is 6.7±0.6 ng/mL and 2.0±0.6 ng/mL for T4 and T3, respectively. The studies of *R. catesbeiana* show slightly lower average values of 6.0±0.9 ng/mL and 1.1±0.9 ng/mL for T4 and T3, respectively. In contrast, the concentrations measured in the present study are higher. As shown in the final row of Table 2, we report maximum levels of 28±10 ng/mL and 2.3±0.5 ng/mL for T4 and T3, respectively. The reason for this dissimilarity could reflect differences the developmental stage that was sampled for quantification (see Gosner stage in Table 2) and the masking of variability by using pooled samples. Additionally, the extraction method can cause differences especially for T4. T4 is largely protein-bound (51), and to obtain a quantification of total T4, the extraction method in the present study included a protein denaturation by urea to release all protein-bound hormone. This differs from earlier applied methodologies and could explain the higher quantity measured in the present study. Notably in the only two previous studies examining T4 levels in individual tadpoles, it was noted that all stages had some individuals where T4 was not detected (20,21). In the present study, we found Gosner stage 44 animals all had detectable T4 concentrations but two individuals had T4 levels below the LoQ. The present study results are, therefore, consistent with these earlier findings in that T4 is highly variable, which contributes to the high variation in T4 levels during metamorphic climax.

In addition to updated information on circulating thyroid hormones and metabolites (T4, T3, T2, T0 and T1Am), this is also the first study to simultaneously analyse same-individual back skin gene expression of the thyroid hormone receptors α and β (*thra*, *thrb*) and TH-induced bZip protein (*thibz*) throughout metamorphosis. Efforts have been made earlier to combine published data on *X. laevis* to get this simultaneous view (8,52), but the same individual measurement, with high level of replication offers an unprecedented depiction of the endocrinological dynamics during anuran postembryonic development, and by approximation, vertebrate development as well. In previously published illustrations, the peak in T3 concentration was reported to occur simultaneously with the peak in *thrb* transcript abundance and precede the maximum in T4 (23,52). In the present study, however, we observed that the peak in T3 and T4 occurs simultaneously, and that *thrb* abundance preceded the increased concentration of both T3 and T4. This exemplifies the exquisite timing of events during the metamorphic process as the synthesis of the receptor occurs in anticipation of the peak in TH concentration.

The observed timewise correlation between THs and the abundance of *thibz* and *thrb* was confirmed
by correlation analyses. Furthermore, the PCA showed that T3 and T4 as well as *thrb* and *thibz*
transcript abundance were the main factors to separate the metamorphic climax from other
developmental stages. The abundance of *thra* transcripts did not, however, follow the same pattern;
its abundance remained stable until the end of metamorphic climax where it decreased significantly
and remained low in the juveniles. This is in good correspondence with an earlier study in *X. laevis*,
where a plateau in *thra* expression also was observed until a drop after metamorphic climax (53).
Relative THR expression is, however, known to be tissue specific, so it is important to note that our
results illustrate the transcript abundance in back skin, a tissue that remodels but is retained
throughout metamorphosis. Other tissues may exhibit different dynamics as exemplified by the
differences in tissue responsiveness to THs (29) and tissue specific expression profiles of THRs (54).
In particular, *thra* seems to have a very diverse expression pattern across tissues (27,29,54).
366

Table 2. Summary of maximal levels of THs (ng/mL) measured during metamorphosis in
published studies. All previous studies used pooled samples with two exceptions where
measured samples were from individuals (20, 21). The final row lists the values measured in the
present study. RIA, radioimmunoassay; SEM, standard error of the mean; UHPLC, ultrahigh-
performance liquid chromatography; UPLC, ultra-performance liquid chromatography; ID,
isotope dilution, MS/MS tandem mass spectrometry

Species	Gosner Stage	Tissue	Assay Used	T_4 (ng/mL) Mean±SEM	T_3 (ng/mL) Mean±SEM	T_2 (ng/mL) Mean±SEM	Reference
R. catesbeiana[a]	45	Plasma	RIA	5.0±0.3	0.8±0.1	-	(18)
R. catesbeiana[a]	45	Plasma	RIA	4.6±0.5	0.8±0.2	-	(18)
R. catesbeiana	43	Plasma	RIA	5.0±0.4	1.4±0.1	-	(13)
R. catesbeiana	43-44	Plasma[b]	RIA	6.0	1.5±0.3	-	(19)
R. catesbeiana	43-45	Serum	RIA	9.4±0.6	-	-	(20)
R. clamitans	45	Plasma	RIA	7.2±0.5	-	-	(21)
R. esculenta	44	Serum	RIA	2.6	0.1	-	(22)
X. laevis	44	Plasma	RIA	7.5	5.2	-	(23)
X. laevis	44	Serum	RIA	5.4	0.2	-	(22)
X. laevis	44	Plasma [c]	UPLC-ID-MS/MS	10.7±0.8	-	-	(14)
X. borealis	42-44	Plasma	RIA	8.7	4.5	-	(15)
X. clivii	42-44	Plasma	RIA	9.2	2.0	-	(15)
X. muelleri	42-44	Plasma	RIA	6.4	4.0	-	(15)
R. catesbeiana	44	Serum	UHPLC-MS/MS	28±10	2.3±0.5	0.9±0.3	The present study

373 [a]Two data sets were provided in Regard et al, 1978
374 [b]From ethanol extract
375 [c]From a single sample
376

To the best of our knowledge, this is the first study to quantify T2 throughout tadpole
metamorphosis. Interestingly, T2 increased earlier than T3 and T4 prior to the significant increase in
transcript abundance of *thrb* and *thibz* and remained high during metamorphosis. This indicates a
specific and distinct biological role of this biomolecule in anuran postembryonic development. There
is limited research on the action of T2 in different vertebrates and, to the best of our knowledge, there
are no studies about the biological role in amphibians. In humans and rodents it has, however, been
established that this iodinated molecule has biological activity distinct from that of T4 and T3 (55),

including for instance activation of fatty acid oxidation and down-regulation of lipogenesis in hepatic fatty acid metabolism (56,57). (58–61)In the freshwater fish, tilapia, T2 acts as a selective activator of a specific long isoform of THR-β named L-TRβ1 and modulates its expression *in vivo* (58). T2 also specifically regulates gene networks associated with cell signaling and transcriptional pathways in brain and liver, which are distinct from those affected by T3 (59,60). Furthermore, T2 affects myelination in cerebellum (59), which is essential for neurodevelopment and participates in growth processes via L-TRb1 activation (61).

Based on these findings and the fact that T2 concentration was significantly increased earlier than T3 and T4, it is tempting to speculate that T2 may modulate THR expression in amphibians. More research is clearly needed to investigate this hypothesis. Generally, TH-related research tends to focus on T4 and T3, of which the biological role is well-documented. However, deiodinated products like T1 and T2, as well as aminated metabolites (e.g., T1Am) and iodothyroacetic acids (e.g., 5-diiodothyroacetic acid, Diac), are increasingly being recognized as biologically active (57,62,63). These molecules will be an important focus in furthering our understanding of TH dynamics.

The untargeted analysis showed that the changes in THs during metamorphosis are accompanied by similar dynamics in many other biomolecules. Earlier untargeted studies in the serum of *R. catesbeiana* using ultraperformance liquid chromatography coupled with mass spectrometry (UPLC-MS) have found a high degree of remodelling in five core metabolic pathways during metamorphosis. This included arginine and purine/pyrimidine, cysteine/methionine, sphingolipid, and eicosanoid metabolism and the urea cycle (30). In addition, a major role for lipids during the postembryonic process was established and arachidonic acid metabolism also underwent remodeling during metamorphosis (30). In accordance with this, the current study also observed metamorphosis-related changes in several lipids including eicosanoids and arachidonic acid metabolites (Figure 3). The bioindicators that showed the highest predictive power for distinguishing metabolite dynamics like the THs, however, included prostaglandins, acyl-carnitines and fatty acyls (Table 1).

L-Carnitine is an endogenous metabolite found in all mammalian species (64). The most important role of L-carnitine is to facilitate the transport of fatty acids into the mitochondria for beta-oxidation (64). During this process, L-carnitine forms a high-energy bond with a fatty acid to give an acyl carnitine derivative that is transported into the mitochondrial matrix. Here it undergoes transesterification with coenzyme A (CoA) to form the corresponding acyl-CoA, which then enters the fatty acid b-oxidation (64). In this way, L-carnitine is also able to regulate the intracellular ratio of free CoA to acyl-CoA by forming acylcarnitines and eventually remove surplus of acyl groups from the organism by preferential renal excretion. This means that the intramitochondrial relationship between free CoA and acyl-CoA is reflected in the extracellular L-carnitine to acylcarnitine ratio, giving indirect evidence of altered mitochondrial metabolism (64). Considering this, our observation of altered carnitine levels during metamorphosis, which additionally correlates with the TH-dynamics, is consistent with the fact that the mitochondrial metabolism is modified during metamorphosis. A tissue specific metabolic reorganization during metamorphosis has already been observed for *Rana omeimontis* tadpoles (65). This study found that, by metamorphic climax, these tadpoles showed overall increased energy metabolism. In the liver, this was accompanied by upregulated amino acid catabolism and downregulated β-oxidation and glycolysis, suggesting a metabolic shift from carbohydrate and lipids to amino acid. A similar shift is likely occurring in *R. catesbeiana*. Therefore, the results in the current study show that the acyl-carnitines are excellent bioindicators for the metabolic shift during metamorphosis.

The ratio of L-carnitine to acyl-carnitines is currently applied in neonatal screening for inborn errors
of metabolism, so the potency of these bioindicators is already known from other vertebrates (66).
Acyl carnitine concentrations and THs can be biochemically linked as T3 is a known inducer of
carnitine palmitoyltransferase I (CPT-I) gene expression in the liver (67). This enzyme catalyzes the
transfer of long chain fatty acyl groups from CoA to carnitine for translocation across the
mitochondrial inner membrane. A long chain fatty acyl, which was tentatively annotated as 2R-
HpOTrE, and its oxidation product, 9-OxoOTrE, were actually also among the 20 compounds
selected on sPLS-DA component 1. The effect of THs on the transferases for short and -medium
chained fatty acyls to the best of our knowledge, not been investigated. However, the two metabolites
that were classified as medium-chain hydroxy acids and derivatives (Table 1) could potentially be
related to this.
443
We also observed modifications to the arachidonic acid metabolism pathway and eicosanoids during
metamorphosis (Figure 3). Furthermore, a prostaglandin was among the 20 compounds that best
discriminate between Gosner stages (Figure 3C, Table 1). Prostaglandins are synthesized by
sequential enzymatic modification of arachidonic acid (68). They are mainly recognized for their role
in inflammatory responses (69) but they also play a role in neurogenesis (70). A link to TH
homeostasis have been observed in rats with high iodide intake-induced hypothyroidism where
increased levels of prostaglandins were found (71). Therefore, prostaglandins and other eicosanoids
present an additional chemical class of important bioindicators of metamorphosis and vertebrate
developmental endocrinology that could be applied in future studies.
453
The steroid hormone, testosterone, was interestingly found to follow a similar Gosner stage profile as
T3, T4, carnitines and prostaglandins. Testosterone is the main sex hormone in vertebrate males.
However, because the animals in the study were of mixed sex, testosterone seems to serve a different
purpose in this developmental window. It should be noted that the observed increase in testosterone
occurred after anticipated sexual differentiation and is completely independent of sexual maturation.
Much empirical evidence exists that confirms crosstalk between the hypothalamic-pituitary-thyroid
(HPT) axis which involves TH synthesis and the hypothalamic-pituitary-gonadal axis which involves
synthesis of steroid hormone (7,72). As reviewed in Thambirajah et al. (7) this includes the effect of
androgens like testosterone on the HPT axis. However, the exact mechanisms and physiological
consequences are still unclear.
464
The vital crosstalk between the HPT axis and the hypothalamic-pituitary-adrenal axis, which signals
the production of corticosteroids, is also well established. This includes hormones like cortisol and
corticosterone. Corticosterone is maximal at metamorphic climax much like T3 and T4 and it can
accelerate TH-dependent metamorphosis. It has also recently been shown to also be required for
survival through metamorphosis in *X. tropicalis* (73). A metabolite tentatively annotated as
corticosterone was found among the compounds that displayed same Gosner stage profile as T3 and
T4, however it could not be annotated with a spectral match. These findings underline what many
others has called for before, namely that crosstalk between endocrine axes receives more attention in
future studies. It is one more piece in the puzzle that is essential for our understanding of vertebrate
developmental endocrinology. The non-target data provides a necessary context for interpreting TH-
mediated developmental processes and potential molecular crosstalk mechanisms. Furthermore, this
provides a solid foundation from which alternative bioindicators of metamorphosis may be chosen as
feasible endpoints in future risk assessment of chemicals and their potency for causing
developmental toxicity
479

5   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

6   Author Contributions

Rikke Poulsen: Conceptualization, methodology, investigation and formal analysis, writing – original draft. **Shireen H. Jackman:** Methodology, investigation and formal analysis, writing – review and editing. **Martin Hansen:** Conceptualization, methodology, writing – review and editing, funding acquisition and resources. **Caren C. Helbing:** Conceptualization, methodology, investigation and formal analysis, writing – review and editing, funding acquisition and resources.

7   Funding

The work was supported by Natural Sciences and Engineering Council of Canada (NSERC) Discovery grant (RGPIN-2018-03816) to CCH, European Union's Horizon 2020 research and innovation program, under grant agreement No. 825753 (ERGO) and the Carlsberg Foundation (grant no. CF20-0422). MH further acknowledge the financial support from Aarhus University Research Foundation (AUFF-T-2017- FLS-7-4).

8   Acknowledgments

We thank Sara Ohora and Emil Egede Frøkjær for technical assistance.

9   Data Availability Statement

The metabolomics dataset generated for this study can be found in the MassIVE database under the accession number MSV000090714

10   References

1.   Warner A, Mittag J. Thyroid hormone and the central control of homeostasis. J Mol Endocrinol. 2012;49(1):29–35.

2.   Flamant F, Koibuchi N, Bernal J. Editorial: "Thyroid Hormone in Brain and Brain Cells." Front Endocrinol (Lausanne) [Internet]. 2015 Jun 22;6(9):3713–24. Available from: http://journal.frontiersin.org/Article/10.3389/fendo.2015.00099/abstract

3.   Mullur R, Liu Y-Y, Brent GA. Thyroid Hormone Regulation of Metabolism. Physiol Rev [Internet]. 2014 Apr;94(2):355–82. Available from: https://www.physiology.org/doi/10.1152/physrev.00030.2013

4.   Bernal J. Thyroid Hormones in Brain Development and Function. In: KR F, B A, A B, et al., editors. Endotext [Internet]. South Dartmouth (MA): MDText.com, Inc.; Available from: https://www.ncbi.nlm.nih.gov/books/NBK285549/

5.   Hayes TB. Amphibian Metamorphosis: An Integrative Approach. Am Zool. 1997;37(2):121–
3.

6.   Weber R. Anuran Metamorphosis [Internet]. Metamorphosis: Postembryonic Reprogramming
of Gene Expression in Amphibian and Insect Cells. ACADEMIC PRESS, INC.; 1996. 567–
597 p. Available from: http://dx.doi.org/10.1016/B978-0-12-283245-1.50018-X

7.   Thambirajah AA, Wade MG, Verreault J, Buisine N, Alves VA, Langlois VS, et al. Disruption
by stealth - Interference of endocrine disrupting chemicals on hormonal crosstalk with thyroid
axis function in humans and other animals. Environ Res [Internet]. 2022 Jan;203(August
2021):111906. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0013935121012019

8.   Paul B, Sterner ZR, Buchholz DR, Shi Y-B, Sachs LM. Thyroid and Corticosteroid Signaling
in Amphibian Metamorphosis. Cells [Internet]. 2022 May 10;11(10):1595. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0012160607002217

9.   Thambirajah AA, Koide EM, Imbery JJ, Helbing CC. Contaminant and Environmental
Influences on Thyroid Hormone Action in Amphibian Metamorphosis. Front Endocrinol
(Lausanne) [Internet]. 2019 May 14;10(MAY). Available from:
https://www.frontiersin.org/article/10.3389/fendo.2019.00276/full

10.   Buchholz DR. More similar than you think: Frog metamorphosis as a model of human
perinatal endocrinology. Dev Biol [Internet]. 2015;408(2):188–95. Available from:
http://dx.doi.org/10.1016/j.ydbio.2015.02.018

11.   Sachs LM, Buchholz DR. Insufficiency of thyroid hormone in frog metamorphosis and the
role of glucocorticoids. Front Endocrinol (Lausanne). 2019;10(MAY):17–20.

12.   OECD. Amphibian Metamorphosis Assay (AMA) (OECD TG 231). In: Revised Guidance
Document 150 on Standardised Test Guidelines for Evaluating Chemicals for Endocrine
Disruption [Internet]. Paris, France: OECD Publishing; 2018. p. 211–20. Available from:
https://www.oecd-ilibrary.org/environment/revised-guidance-document-150-on-standardised-
test-guidelines-for-evaluating-chemicals-for-endocrine-disruption/amphibian-metamorphosis-
assay-ama-oecd-tg-231_9789264304741-8-en

13.   Suzuki S, Suzuki M. Changes in thyroidal and plasma Iodine compounds during and after
metamorphosis of the bullfrog, Rana catesbeiana. Gen Comp Endocrinol. 1981;45(1):74–81.

14.   Luna LG, Coady K, McFadden JR, Markham DA, Bartels MJ. Quantification of total
thyroxine in plasma from xenopus laevis. J Anal Toxicol. 2013;37(6):326–36.

15.   Buscaglia M, Leloup J. Thyroid hormone secretion during metamorphosis in Xenopus genus.
(Abstract). Gen Comp Endocrinol. 1980;40:369.

16.   Hansen M, Luong X, Sedlak DL, Helbing CC, Hayes T. Quantification of 11 thyroid
hormones and associated metabolites in blood using isotope-dilution liquid chromatography
tandem mass spectrometry. Anal Bioanal Chem [Internet]. 2016 Aug 23;408(20):5429–42.
Available from: http://dx.doi.org/10.1007/s00216-016-9614-9

17.   Simon R, Tietge JE, Michalke B, Degitz S, Schramm KW. Iodine species and the endocrine
555   system: Thyroid hormone levels in adult Danio rerio and developing Xenopus laevis.
556   Fresenius J Anal Chem. 2002;372(3):481–5.

557   18.   Regard E, Taurog A, Nakashima T. Plasma thyroxine and triiodothyronine levels in
558   spontaneously metamorphosing Rana catesbeiana tadpoles and in adult anuran amphibia.
559   Endocrinol 102. 1978;674–84.

560   19.   Miyauchi H, LaRochelle RJ, Suzuki M, Freeman M, Frieden E. Studies on thyroid hormones
561   and their binding in bullfrog tadpole plasma during metamorphosis. Gen Comp Endocrinol.
562   1977;33:254–66.

563   20.   Mondou P, Kaltenbach J. Thyroxine concentrations in blood serum and pericardial fluid of
564   metamorphosing tadpoles and adult frogs. Gen Comp Endocrinol. 1979;39:343–9.

565   21.   Weil M. Changes in plasma thyroxine levels during and after spontaneous metamoprhosis in a
566   natural population of the green frog, Rana clamitans. Gen Comp Endocrinol. 1986;62:8–12.

567   22.   Schultheiss H. T3 and T4 concentrations during metamorphosis of Xenopus laevis and Rana
568   esculenta and in the neotenic mexican axolotl. (Abstract). Gen Comp Endocrinol.
569   1980;40:372.

570   23.   Leloup J, Buscaglia M. La triiodothyronine: hormone de la metamorphose des amphibiens. C
571   R Acad Sci. 1977;284:2261–3.

572   24.   Spencer CA. Assay of Thyroid Hormones and Related Substances. In: KR Feingold, Anawalt
573   B, Boyce A, Al E, editors. Endotext [Internet]. South Darthmouth: MDText.com, Inc.;
574   Available from: http://www.ncbi.nlm.nih.gov/pubmed/25905337

575   25.   Davidsen N, Ramhøj L, Lykkebo CA, Kugathas I, Poulsen R, Rosenmai AK, et al. PFOS-
576   induced thyroid hormone system disrupted rats display organ-specific changes in their
577   transcriptomes. Environ Pollut. 2022;305(February).

578   26.   Flamant F, Cheng S, Hollenberg AN, Moeller LC, Samarut J, Wondisford FE, et al. Thyroid
579   Hormone Signaling Pathways: Time for a More Precise Nomenclature. Endocrinology
580   [Internet]. 2017 Jul 1;158(7):2052–7. Available from:
581   https://academic.oup.com/endo/article/158/7/2052/3784596

582   27.   Wen L, He C, Sifuentes CJ, Denver RJ. Thyroid hormone receptor alpha is required for
583   thyroid hormone-dependent neural cell proliferation during tadpole metamorphosis. Front
584   Endocrinol (Lausanne). 2019;10(JUN):1–13.

585   28.   Laslo M, Denver RJ, Hanken J. Evolutionary conservation of thyroid hormone receptor and
586   deiodinase expression dynamics in ovo in a direct-developing frog, eleutherodactylus coqui.
587   Front Endocrinol (Lausanne). 2019;10(MAY):1–14.

588   29.   Maher SK, Wojnarowicz P, Ichu TA, Veldhoen N, Lu L, Lesperance M, et al. Rethinking the
589   biological relationships of the thyroid hormones, L-thyroxine and 3,5,3'-triiodothyronine.
590   Comp Biochem Physiol - Part D Genomics Proteomics [Internet]. 2016;18:44–53. Available
591   from: http://dx.doi.org/10.1016/j.cbd.2016.04.002

30. Ichu TA, Han J, Borchers CH, Lesperance M, Helbing CC. Metabolomic insights into system-
wide coordination of vertebrate metamorphosis. BMC Dev Biol. 2014;14(1).

31. Poulsen R, De Fine Licht HH, Hansen M, Cedergreen N. Grandmother's pesticide exposure
revealed bi-generational effects in Daphnia magna. Aquat Toxicol [Internet]. 2021
Jul;236(January):105861. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0166445X2100120X

32. Ma NL, Hansen M, Roland Therkildsen O, Kjær Christensen T, Skjold Tjørnløv R, Garbus
SE, et al. Body mass, mercury exposure, biochemistry and untargeted metabolomics of
incubating common eiders (Somateria mollissima) in three Baltic colonies. Environ Int
[Internet]. 2020;142(January):105866. Available from:
https://doi.org/10.1016/j.envint.2020.105866

33. Marana MH, Poulsen R, Thormar EA, Clausen CG, Thit A, Mathiessen H, et al. Plastic
nanoparticles cause mild inflammation, disrupt metabolic pathways, change the gut microbiota
and affect reproduction in zebrafish: A full generation multi-omics study. J Hazard Mater
[Internet]. 2022 Feb;424(October):127705. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S030438942102673X

34. Rasmussen JA, Villumsen KR, Ernst M, Hansen M, Forberg T, Gopalakrishnan S, et al. A
multi-omics approach unravels metagenomic and metabolic alterations of a probiotic and
synbiotic additive in rainbow trout (Oncorhynchus mykiss). Microbiome [Internet].
2022;10(1):1–19. Available from: https://doi.org/10.1186/s40168-021-01221-8

35. Poulsen R, Zekri Y, Guyot R, Flamant F, Hansen M. Effect of in utero and lactational
exposure to a thyroid hormone system disrupting chemical on mouse plasma metabolome and
brain transcriptome. J Hazard Mater Lett. 2022;

36. Gosner KL. A Simplified Table for Staging Anuran Embryos and Larvae with Notes on
Identification. Herpetologica. 1960;16(3):183–90.

37. Corrie LM, Kempe MN, Blajkevitch O, Shang D, Helbing CC. Dioctyl Sodium Sulfosuccate
as a Potential Endocrine Disruptor of Thyroid Hormone Activity in American bullfrog, Rana
(Lithobates) catesbeiana, Tadpoles. Arch Environ Contam Toxicol [Internet]. 2021;80(4):726–
34. Available from: https://doi.org/10.1007/s00244-021-00835-1

38. Veldhoen N, Skirrow RC, Brown LLY, Van Aggelen G, Helbing CC. Effects of acute
exposure to the non-steroidal anti-inflammatory drug ibuprofen on the developing North
American bullfrog (Rana catesbeiana) tadpole. Environ Sci Technol. 2014;48(17):10439–47.

39. Davidsen N, Ramhøj L, Lykkebo CA, Kugathas I, Poulsen R, Rosenmai AK, et al. PFOS-
induced thyroid hormone system disrupted rats display organ-specific changes in their
transcriptomes. Environ Pollut. 2022;305(April):119340.

40. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna: R
Foundation for Statistical Computing; 2020. Available from: https://www.r-project.org/.

41. Alexis Dinno. dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums [Internet].
2017. Available from: https://cran.r-project.org/package=dunn.test

42.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-Verlag; 2016.

43.   Vu VQ. ggbiplot: A ggplot2 based biplot. [Internet]. 2011. Available from:
http://github.com/vqv/ggbiplot

44.   Kolde R. pheatmap: Pretty Heatmaps [Internet]. 2019. Available from: https://cran.r-
project.org/package=pheatmap

45.   Cao K-A Le, Florian Rohart, Ignacio Gonzalez, Sebastien Dejean with key contributors Benoit
Gautier F, Bartolo, contributions from Pierre Monget, Jeff Coquery FY and BL. mixOmics:
Omics Data Integration Project. [Internet]. 2016. Available from: https://cran.r-
project.org/package=mixOmics

46.   Viant MR, Ebbels TMD, Beger RD, Ekman DR, Epps DJT, Kamp H, et al. Use cases, best
practice and reporting standards for metabolomics in regulatory toxicology. Nat Commun
[Internet]. 2019;10(1). Available from: http://dx.doi.org/10.1038/s41467-019-10900-y

47.   Dührkop K, Fleischauer M, Ludwig M, Aksenov AA, Melnik A V., Meusel M, et al. SIRIUS
4: a rapid tool for turning tandem mass spectra into metabolite structure information. Nat
Methods [Internet]. 2019;16(4):299–302. Available from: http://dx.doi.org/10.1038/s41592-
019-0344-8

48.   Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. KEGG as a reference resource
for gene and protein annotation. Nucleic Acids Res [Internet]. 2016 Jan 4;44(D1):D457–62.
Available from: https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/gkv1070

49.   Karp PD, Billington R, Caspi R, Fulcher CA, Latendresse M, Kothari A, et al. The BioCyc
collection of microbial genomes and metabolic pathways. Brief Bioinform [Internet]. 2019 Jul
19;20(4):1085–93. Available from: https://academic.oup.com/bib/article/20/4/1085/4084231

50.   Wishart DS, Feunang YD, Marcu A, Guo AC, Liang K, Vázquez-Fresno R, et al. HMDB 4.0:
the human metabolome database for 2018. Nucleic Acids Res [Internet]. 2018 Jan
4;46(D1):D608–17. Available from: http://academic.oup.com/nar/article/46/D1/D608/4616873

51.   Refetoff S. Thyroid Hormone Serum Transport Proteins [Internet]. KR F, B A, A B, Al. E,
editors. Vol. 3, Endotext. 2000. 1–15 p. Available from:
http://www.ncbi.nlm.nih.gov/pubmed/25905421

52.   Shi Y, Wong J, Puzianowska-Kuznicka M, Stolow MA. Tadpole competence and tissue-
specific temporal regulation of amphibian metamorphosis: Roles of thyroid hormone and its
receptors. BioEssays [Internet]. 1996 May;18(5):391–9. Available from:
https://onlinelibrary.wiley.com/doi/10.1002/bies.950180509

53.   Yaoita Y, Brown DD. A correlation of thyroid hormone receptor gene expression with
amphibian metamorphosis. Genes Dev. 1990;4(11):1917–24.

54.   Wang X, Matsuda H, Shi YB. Developmental regulation and function of thyroid hormone
receptors and 9-cis retinoic acid receptors during Xenopus tropicalis metamorphosis.
Endocrinology. 2008;149(11):5610–8.

55.   Köhrle J. The Colorful Diversity of Thyroid Hormone Metabolites. Eur Thyroid J.
669   2019;8(3):115–29.

670   56.   Damiano F, Rochira A, Gnoni A, Siculella L. Action of thyroid hormones, T3 and T2, on
671   hepatic fatty acids: Differences in metabolic effects and molecular mechanisms. Int J Mol Sci.
672   2017;18(4).

673   57.   Bouazza A, Favier R, Fontaine E, Leverve X, Koceir E-A. Potential Applications of Thyroid
674   Hormone Derivatives in Obesity and Type 2 Diabetes: Focus on 3,5-Diiodothyronine (3,5-T2)
675   in Psammomys obesus (Fat Sand Rat) Model. Nutrients [Internet]. 2022 Jul 25;14(15):3044.
676   Available from: https://medium.com/@arifwicaksanaa/pengertian-use-case-a7e576e1b6bf

677   58.   Mendoza A, Navarrete-Ramírez P, Hernández-Puga G, Villalobos P, Holzer G, Renaud JP, et
678   al. 3,5-T2 Is an Alternative Ligand for the Thyroid Hormone Receptor β1. Endocrinology.
679   2013;154(8):2948–58.

680   59.   Hernández-Linares Y, Olvera A, Villalobos P, Lozano-Flores C, Varela-Echavarría A, Luna
681   M, et al. 3,5-T2 and 3,3′,5-T3 Regulate Cerebellar Thyroid Hormone Signalling and Myelin
682   Molecular Dynamics in Tilapia. Sci Rep. 2019;9(1):1–9.

683   60.   Olvera A, Martyniuk CJ, Buisine N, Jiménez-Jacinto V, Sanchez-Flores A, Sachs LM, et al.
684   Differential transcriptome regulation by 3,5-T2 and 3′,3,5-T3 in brain and liver uncovers novel
685   roles for thyroid hormones in tilapia. Sci Rep. 2017;7(1):1–12.

686   61.   Navarrete-Ramírez P, Luna M, Valverde-R C, Orozco A. 3,5-Di-Iodothyronine Stimulates
687   Tilapia Growth Through an Alternate Isoform of Thyroid Hormone Receptor B1. J Mol
688   Endocrinol. 2013;52(1):1–9.

689   62.   Köhrle J, Lehmphul I, Pietzner M, Renko K, Rijntjes E, Richards K, et al. 3,5-T2—A Janus-
690   Faced Thyroid Hormone Metabolite Exerts Both Canonical T3-Mimetic Endocrine and
691   Intracrine Hepatic Action. Front Endocrinol (Lausanne). 2020;10(January):1–15.

692   63.   Köhrle J, Biebermann H. 3-Iodothyronamine—A Thyroid Hormone Metabolite With Distinct
693   Target Profiles and Mode of Action. Endocr Rev. 2019;40(2):602–30.

694   64.   Reuter SE, Evans AM. Carnitine and acylcarnitines: Pharmacokinetic, pharmacological and
695   clinical aspects. Clin Pharmacokinet. 2012;51(9):553–72.

696   65.   Zhu W, Chang L, Zhao T, Wang B, Jiang J. Remarkable metabolic reorganization and altered
697   metabolic requirements in frog metamorphic climax. Front Zool. 2020;17(1):1–16.

698   66.   McCann MR, De la Rosa MVG, Rosania GR, Stringer KA. L-carnitine and acylcarnitines:
699   Mitochondrial biomarkers for precision medicine. Metabolites. 2021;11(1):1–21.

700   67.   Jackson-Hayes L, Song S, Lavrentyev EN, Jansen MS, Hillgartner FB, Tian L, et al. A thyroid
701   hormone response unit formed between the promoter and first intron of the carnitine
702   palmitoyltransferase-Iα gene mediates the liver-specific induction by thyroid hormone. J Biol
703   Chem [Internet]. 2003;278(10):7964–72. Available from:
704   http://dx.doi.org/10.1074/jbc.M211062200

705   68.   Hata AN, Breyer RM. Pharmacology and signaling of prostaglandin receptors: Multiple roles
706   in inflammation and immune modulation. Pharmacol Ther. 2004;103(2):147–66.

707   69.   Ricciotti E, Fitzgerald GA. Prostaglandins and inflammation. Arterioscler Thromb Vasc Biol.
708   2011;31(5):986–1000.

709   70.   Tamiji J, Crawford DA. Prostaglandin E2 and misoprostol induce neurite retraction in Neuro-
710   2a cells. Biochem Biophys Res Commun [Internet]. 2010;398(3):450–6. Available from:
711   http://dx.doi.org/10.1016/j.bbrc.2010.06.098

712   71.   Liu Q, Zhang Y, Zhao H, Yao X. Increased Epoxyeicosatrienoic Acids and
713   Hydroxyeicosatetraenoic Acids After Treatment of Iodide Intake Adjustment and 1,25-
714   Dihydroxy-Vitamin D3 Supplementation in High Iodide Intake–Induced Hypothyroid
715   Offspring Rats. Front Physiol. 2021;12(July):1–12.

716   72.   Duarte-Guterman P, Navarro-Martín L, Trudeau VL. Mechanisms of crosstalk between
717   endocrine systems: Regulation of sex steroid hormone synthesis and action by thyroid
718   hormones. Gen Comp Endocrinol [Internet]. 2014;203:69–85. Available from:
719   http://dx.doi.org/10.1016/j.ygcen.2014.03.015

720   73.   Shewade LH, Schoephoerster JA, Patmann MD, Kulkarni SS, Buchholz DR. Corticosterone Is
721   Essential for Survival through Frog Metamorphosis. Endocrinol (United States).
722   2020;161(12):1–15.