LXR-inducible host E3 ligase IDOL targets a human cytomegalovirus reactivation determinant to promote latency

Luwanika Mlera[1,2], Donna Collins-McMillen[1,2], Sebastian Zeltzer[1,2], Jason C. Buehler[1,2†], Melissa Moy[3†], Kristen Zarrella[1], Katie Caviness[1,2,4§], Louis Cicchini[2,5¶], David J. Tafoya[1,2], and Felicia Goodrum[1,2,3,5*]

[1]Department of Immunobiology, University of Arizona, Tucson, AZ 85721, USA

[2]BIO5 Institute, University of Arizona, Tucson, AZ 85721, USA

[3]Graduate Interdisciplinary Program in Cancer Biology, University of Arizona, Tucson, AZ 85721, USA

[4]Graduate Interdisciplinary Program in Genetics, University of Arizona, Tucson, AZ 85721, USA

[5]Department of Molecular and Cellular Biology, University of Arizona, Tucson, AZ 85721, USA

[†]Current address: Imanis Life Sciences, Rochester, MN 55901, USA

[§]Current address: National Biodefense Analysis and Countermeasures Center, Fort Detrick, MD 21702, USA

[¶]Current address: PPD, Inc., 929 N Front St. Wilmington, NC 28401, USA

*Correspondence: fgoodrum@arizona.edu

Summary

Liver X receptor (LXR) signaling broadly restricts virus replication; however, the mechanisms of restriction are poorly defined. Here, we demonstrate that the LXR-inducible cellular E3 ligase IDOL (inducible degrader of low-density lipoprotein receptor, LDLR) restricts replication of the beta-herpesvirus, human cytomegalovirus (HCMV), for the establishment of viral latency. IDOL is highly expressed in undifferentiated hematopoietic cells where HCMV establishes latency, but is sharply downregulated upon differentiation, a stimulus for reactivation. Importantly, IDOL restricts replication by driving the instability of a key viral determinant required for reactivation, the 33-kDa protein encoded by *UL136* (UL136p33). *UL136* encodes multiple proteins that differentially impact latency and reactivation. UL136p33 is targeted for rapid turnover by the proteasome and its stabilization by mutation of lysine residues to arginine results in a failure to quiet replication for latency. We show that IDOL interacts with and targets UL136p33 for turnover, but not the stabilized variant. While induction of IDOL prevents reactivation from latency, IDOL depletion increases viral gene expression in undifferentiated hematopoietic cells. This work establishes the UL136p33-IDOL interaction as a key regulator of the bistable switch between latency and reactivation. It further suggests a model whereby a key viral determinant of HCMV reactivation is regulated by a host E3 ligase and acts as a sensor at the tipping point between the decision to maintain the latent state or exit latency for reactivation.

Significance

Herpesviruses establish life-long latent infections, which pose an important risk for disease particularly in the immunocompromised. Our work is focused on the beta-herpesvirus, human cytomegalovirus (HCMV) that latently infects the majority of the population worldwide. Defining the mechanisms by which HCMV establish latency or reactivate from latency is important to controlling disease. Here, we demonstrate that the cellular inducible degrader of low-density lipoprotein receptor, IDOL, targets a HCMV determinant of reactivation for degradation. The instability of this determinant is important for the establishment of latency. This work defines a pivotal virus-host interaction that allows HCMV to sense changes in host biology to navigate decisions to establish latency or replicate.

Introduction

Viral latency, the ability of a virus to persist indefinitely in the infected host, is a hallmark of herpesviruses. Latency is defined as a non-replicative state punctuated by sporadic reactivation events where the virus replicates productively for transmission. Human cytomegalovirus (HCMV) is a beta-herpesvirus that establishes latent infection in hematopoietic progenitor cells (HPCs) and cells of the myeloid lineage (1). Reactivation of HCMV from latency in the immunocompromised can result in life-threatening disease and is a particularly important risk factor in the context of stem cell and solid organ transplantation. The molecular mechanisms by which HCMV senses changes in the host to make decisions to maintain latency or reactivate from latency is important in developing novel strategies to control viral reactivation and subsequent CMV disease.

Latency is a complex multifactorial phenomenon dependent on viral determinants, host

cell intrinsic, innate, and adaptive responses, as well as host cell signaling and chromatin

remodeling (2). Current paradigms hold that HCMV reactivation is tightly connected to

hematopoietic cell differentiation (3-7) or inflammatory signaling (3, 8-10). However, much

remains to be understood about the precise host contributions and key viral factors that

navigate the entry into or exit from latency. We and others have defined roles of *UL133-*

138 gene locus encoded within the UL*b'* region of the HCMV genome for latency and

reactivation. *UL138* is required for latency and its protein functions by regulating cell

surface receptor trafficking and signaling (11-14), innate immune responses (15) and also

contributes to repression of viral gene expression (16). *UL135*, by contrast, is required for

reactivation from latency, functioning in part by opposing the function of *UL138* (17-19).

UL136 is expressed as 5 protein isoforms that have differential effects on the

establishment of latency or reactivation. The largest, membrane-associated isoforms,

UL136p33 (UL136p33) and p26, are required for reactivation, whereas the small, soluble

isoforms are required for latency in both *in vitro* and *in vivo* experimental models (20, 21).

Importantly, UL136 proteins accumulate with delayed kinetics relative to UL135 and

UL138 and maximal accumulation of *UL136* gene products depend on the onset of viral

DNA synthesis and/or entry into late phase. Given its kinetics and dependence on viral

DNA synthesis for maximal expression, we postulate that UL136 may negotiate

checkpoints that function to maintain latency or commitment to reactivation from latency.

Identifying host pathways regulating these viral determinants is key to defining the

mechanistic basis by which HCMV senses and responds to changes in host biology to

maintain latency or reactivate.

The cellular environment the virus encounters upon infection undoubtedly plays important roles in the establishment of latency with a host of restriction factors prepared to silence the viral genome, as well as a transcriptional milieu that impacts viral gene expression. As an example, we have shown that the cellular transcription factor early growth response factor-1 (EGR-1) drives the expression of UL138 for the establishment of latency (19, 22). As EGR-1 is required to maintain stemness in the bone marrow niche and is highly expressed in undifferentiated hematopoietic cells (23), cells are primed to drive the expression of UL138 for latency upon infection. However, HCMV also regulates EGR-1 expression through the differential trafficking and turnover of the host receptor tyrosine kinase, epidermal growth factor receptor (EGFR) upstream of EGR-1 (11, 18, 19) and through direct targeting by microRNAs that promote reactivation (22). Thus, HCMV has evolved a multi-pronge strategy to manipulate and utilized host pathways to regulate its gene expression, hardwiring infection into host signaling pathways. With regards to reactivation, activation of the major immediate early (MIE) transcriptional unit, encoding the major transactivators for viral replication, is highly dependent on host transcription factors in myeloid cells (24). Host transcription factors CREB, FOXO3a and AP-1 are downstream of major signaling pathways, including Src, PI3K/Akt, and MEK/ERK, and are important for re-expression of MIE genes (25-27). Further, UL7 encodes a ligand that binds the cellular Fms-like tyrosine kinase 3 receptor (Flt-3R) that when secreted upon reactivation activates PI3K/AKT and MAPK signaling to prevent cell death and stimulate differentiation for reactivation (28, 29). Through its co-evolution with the human host, the biology of HCMV infection has become tightly intertwined with the biology its host, which

reflects the complex mechanisms the virus has evolved to "sense" and "respond" to

changes in the cellular environment to make "decisions" to maintain latency or reactivate

from latency.

Here we demonstrate that the largest isoform of UL136, UL136p33, is targeted for rapid

turnover relative to other UL136 protein isoforms and that its instability ($t_{1/2}$~1 h) is

important to the establishment of latency. We identified the LXR-inducible degrader of

low-density lipoprotein receptor (LDLR) IDOL (also known as myosin regulatory light

chain interacting protein or MYLIP), as the host E3 ligase that targets UL136p33 for

turnover. IDOL expression is induced by oxysterol activation of the liver X receptor (LXR)

(30) and is regulated through hematopoietic differentiation (31, 32). In addition to LDLR,

IDOL also targets VLDLR and ApoER-2 and, therefore, is an important homeostatic

regulator of intracellular cholesterol (30, 33-35). The targeting of UL136p33 by IDOL

suggests that the virus has evolved to be responsive to changes in cellular cholesterol in

its commitment to reactivation. A model emerges whereby high levels of IDOL maintains

low levels of UL136p33 for the establishment of latency, and the downregulation of IDOL

by differentiation results in the accumulation of UL136p33 to drive reactivation. Indeed,

induction of IDOL results in increased turnover of UL136p33 and prevents reactivation of

HCMV from latency, whereas depletion of IDOL increases viral gene expression in

hematopoietic cells. Our findings define UL133p33 is a key viral determinant of HCMV

reactivation that acts as a sensor at the tipping point between the decision to maintain the

latent state or exit latency for reactivation.

Results

UL136p33 is unstable and is targeted for proteasomal degradation

UL136 is expressed as at least 5 co-terminal protein isoforms (Fig. 1A). These isoforms are encoded by transcripts with unique 5' ends and have distinct roles in viral latency and reactivation (20, 21). The largest of these isoforms is a 33-kDa protein (UL136p33) that is required for replication in endothelial cells as well as reactivation from latency in CD34+ HPCs and in humanized mice (20). Conversely, the two small isoforms p23 and p19 suppress replication in endothelial cells and promote latency in CD34+ HPCs and humanized mice (21). The diametric nature of the UL136 system presupposes additional levels of control to coordinate and commit to one infection program over another, thereby allowing for either optimal replication/reactivation or robust latency. The UL136 isoforms, and particularly the full-length isoform UL136p33, are uniquely labile and sensitive to mechanical lysis while being processed (21). Based on this, we hypothesized that the concentration of some UL136 isoforms is regulated by increased proteolysis, and that the instability of UL136p33 may be important to establishing or maintaining latency.

To explore the possibility that UL136 isoforms have different stabilities, we analyzed the levels of UL136 isoforms in infected fibroblasts treated with cycloheximide (CHX) to halt protein synthesis. Cycloheximide treatment resulted in a striking loss of UL136p33, p25, and p23, but not p26 or p19 (Fig. 1B, left panel). These results indicated differential stability whereby UL136p33, p25, and p23 may be subject to more rapid turnover, while other UL136 isoforms are relatively stable. Proteasome inhibition robustly rescued UL136p33, p25, and p23 isoforms (Fig. 1B, right panel). Inhibition of the lysosome with

NH$_4$Cl had no effect on rescue of any of the UL136 isoforms; however, inhibition of both

the lysosome and proteasome showed a somewhat additive effect. Together, these

results indicate that the rapid turnover of UL136p33, p25, and p23 is regulated by the

proteasome.

To establish the kinetics of isoform turnover, we analyzed the decay of UL136 isoforms

in fibroblasts infected with TB40/E-*UL136*$_{myc}$ and treated with CHX at peak UL136 isoform

expression (Fig 1C). UL136p33 decayed with a half-life of 1 h ($t_{1/2}$=1 h) and p25 exhibited

a $t_{1/2}$=3 h, while all other isoforms were stable over the 6 h time course (Fig 1D). The p23

isoform accumulates to very low levels making the decay of this protein is difficult to

determine. The rapid decay of UL136p33 relative to the other isoforms, as well as its

significance in HCMV reactivation prompted us to focus on determining the factor(s)

contributing to its instability.

Lysine to arginine mutations stabilize UL136p33

UL136 has four lysine (K) residues (denoted by red lines in Fig. 1A), three of which lie

upstream of the transmembrane domain (TMD) and are unique to the p33 isoform. To

determine if the turnover of UL136p33 was regulated by ubiquitin-mediated proteolysis,

we substituted all lysine residues for arginine (R) through site-directed mutagenesis in an

expression vector encoding UL136p33 to generate *UL136p33*$_{myc}$*ΔK→R*. The p33$_{ΔK→R}$

mutant had a $t_{1/2}$=5 h, compared to ~1 h for UL136p33 following CHX treatment (Fig. 2A).

This observation, together with the stabilization of UL136p33 by proteasomal inhibition

(Fig. 1B), suggests that UL136p33 turnover is regulated by ubiquitin-mediated

proteasomal decay. Further, the rapid turnover of exogenously expressed UL136p33 (Fig.

2A, left panel) indicates that UL136p33's degradation is not dependent on virus or virus-

induced factors.

To determine the importance of UL136p33 instability in infection, we introduced the lysine

to arginine mutations into the $UL136_{myc}$ gene within the bacterial artificial chromosome

clone of TB40/E ($UL136_{myc}\Delta K \rightarrow R$). We measured the relative rates of decay in fibroblasts

infected with either the parental $UL136_{myc}$ or $UL136_{myc}\Delta K \rightarrow R$ virus and treated with CHX

and found that $p33_{\Delta K \rightarrow R}$ was stabilized during infection relative to wild type UL136p33 (Fig.

2B). Interestingly, we stabilization of UL136p33 diminished levels of UL136p26 and p25

isoforms by mechanisms that have yet to be determined.

We then performed multistep growth curves on fibroblasts infected with $UL136_{myc}$ or

$UL136_{myc}\Delta K \rightarrow R$ to analyze the impact of stabilizing UL136p33 on virus replication. No

change in the kinetics or yield of viral growth was observed during the productive infection

in fibroblasts when UL136p33 is stabilized relative to the WT infection (Fig. 2C). This was

not surprising since the UL136p33 isoform, while important for replication in endothelial

cells and CD34+ HPCs, is dispensable for replication in fibroblasts (20, 21). In summary,

UL136p33 instability is the result of lysine-dependent, proteasomal degradation and

mutating lysine residues to arginine in $UL136$ stabilizes the protein with no effect on virus

replication in fibroblasts.

Since UL136p33 is required for reactivation from latency in CD34+ HPCs (20), we

hypothesized that the instability of UL136p33 allows for a tunable concentration gradient

where low levels are important for the establishment of latency and accumulation past a

threshold is important for reactivation. To test this hypothesis, we performed a latency

assay in primary human CD34+ HPCs infected with the parental $UL136_{myc}$ or

$UL136_{myc}\Delta K\rightarrow R$ viruses. $UL136_{myc}$-infected CD34+ HPCs generated >2-fold frequencies

of infectious centers following reactivation compared to the pre-reactivation control (Fig.

2D). However, the pre- and post-reactivation frequencies of infectious centers were not

different in the $UL136_{myc}\Delta K\rightarrow R$ infection and were almost two-fold above $UL136_{myc}$ pre-

reactivation. These results indicate that HCMV replication in CD34+ HPCs is not quieted

when UL136p33 is stabilized, and that this virus does not establish a latent infection. As

is the case with many viral proteins, UL136 isoforms, even with a myc epitope tag, are

not expressed to detectable levels in hematopoietic cells, and we have been unsuccessful

in generating antibodies to UL136. Nevertheless, given the requirement for UL136p33 for

reactivation and replication in CD34+ HPCs, the $UL136_{myc}\Delta K\rightarrow R$ infection phenotype is

consistent with accumulation of UL136p33 beyond a threshold critical for commitment to

reactivation.

The host E3 ligase IDOL targets UL136p33 for proteasomal degradation

Ubiquitination of cargo by E3 ligases is an important step in targeting a protein for

proteasomal degradation (36, 37). Because UL136p33 is rapidly turned over by the

proteasome even in the absence of infection and lysine to arginine substitutions stabilized

UL136p33, we hypothesized that a host E3 ligase targets UL136p33 for turnover. We

identified UL136 interacting proteins by a yeast two-hybrid screen (Y2H) using the C-

terminal region of UL136 (amino acids 141-240) as bait. Intriguingly, we identified the host

E3 ligase IDOL (**i**nducible **d**egrader **o**f **l**ow-density lipoprotein receptor), also known as

MYLIP, as a candidate host interactor of UL136 (Supporting information, SI Table 1).

IDOL induces the degradation of the low-density lipoprotein receptor family members

LDLR, VLDLR, and ApoER-2 (34, 35). The degradation of these receptors is a

physiological response to increased cholesterol, which is modified to oxysterols that

activate the LXR transcription factor to induce *IDOL* expression (35, 38, 39).

To determine if IDOL targets UL136p33, we analyzed the ability of IDOL to interact with

and impact UL136p33 levels. We validated the interaction between IDOL and UL136p33

by immunoprecipitating (IP) UL136p33-IDOL complexes from cells transiently

overexpressing myc-tagged UL136p33 or HA-tagged IDOL. Transient overexpression is

necessary due to the lack of commercially available antibodies specific and sensitive

enough to detect or immunoprecipitate endogenous IDOL. Pulldown of $IDOL_{HA}$ or $p33_{myc}$

coprecipitated the alternant partner (Fig. 3A), confirming the interaction between

UL136p33 and IDOL. Further analysis indicated that co-expression of IDOL with

UL136p33 resulted in a ~60% reduction in UL136p33 levels compared to when UL136p33

was co-expressed with an empty vector (Fig. 3B top and quantified in Fig. 3C), consistent

with IDOL driving UL136p33 turnover. By contrast, $p33_{\Delta K \to R}$ levels were not affected by

IDOL co-expression (Fig. 3B bottom and quantified in Fig. 3C), in agreement with the

hypothesis that ubiquitination on lysine residues is required for UL136p33 turnover via

the proteasome. Taken together, IDOL functions as a host E3 ligase that targets

UL136p33 for proteasomal degradation.

We next induced endogenous IDOL expression with an LXR agonist GW3965 or LXR-

independent agonist HBX41-108 in cells transiently expressing UL136p33 or $p33_{\Delta K \to R}$.

Because endogenous IDOL cannot be detected by western blot using available

antibodies (33, 40, 41), the induction of IDOL was verified by the loss of its target ApoER-

2 (Fig. 3D). Induction of IDOL, by both GW3965 and HBX41-108 resulted in a reduction

of UL136p33 (Fig. 3D, top), but not $p33_{\Delta K \to R}$ (Fig. 3D, bottom). These results further

confirm that endogenous IDOL targets UL136p33 for ubiquitin-dependent degradation.

The Y2H bait was composed of the C-terminal region of UL136 spanning amino acids

141-240 downstream of the predicted transmembrane domain. These residues are

shared by all the UL136 isoforms (Fig. 1A). Further, our data indicated that p25 and p23

proteins are also labile following CHX treatment and these smaller isoforms could be

rescued, like UL136p33, by inhibiting the proteasome (Fig. 1B, right panel). The p25 and

p23 isoforms have a single lysine which is located downstream of the TMD in the full

length UL136 isoform (Fig. 1A). Transient co-expression of IDOL with either p25 or p23

did not lead to any notable turnover of these proteins (SI Fig. 1A-C). These data

collectively led us to conclude that IDOL specifically targets UL136p33 for turnover, and

it does not mediate the degradation of p25 and p23. The mechanism by which p25 and

p23 are turned over remains to be elucidated.

To further define the specificity of IDOL-mediated turnover of UL136p33, we co-

expressed UL136p33 with another E3 ligase, Cbl. Cbl targets tyrosine kinase receptors,

such as EGFR, for ubiquitination and subsequent degradation by the lysosome (42).

Levels of UL136p33 were unaffected by co-expression with Cbl (SI Fig. 1D-E), indicating

specificity of IDOL for proteasomal turnover of UL136p33.

IDOL impacts UL136p33 protein levels in HCMV-infected primary fibroblasts

In the context of a productive infection in fibroblasts, we found that infection with either

the parental $UL136_{myc}$ or $UL136_{myc}\Delta K{\rightarrow}R$ induced *IDOL* transcripts by at least 5-fold

compared to mock-infection by reverse transcriptase, quantitative PCR (RT-qPCR), (SI

Fig. 2A). *IDOL* transcript upregulation was sustained over time during infection and

peaked at close to 10-fold by 3 dpi. No changes in *IDOL* transcript levels were detected

in mock controls over the same time-course. Because *IDOL* induction is a consequence

of increased cellular cholesterol, we measured cholesterol content in infected cells to

correlate it to the upregulation of *IDOL* transcripts. Using this approach, we noted that

HCMV increased cholesterol concentration by at least 29% from 1 dpi onward (SI Fig.

2B), consistent with previous findings on HCMV-modulation of lipid biosynthesis (43-48).

Taken together, infection in fibroblasts upregulates expression levels of IDOL transcripts,

which is correlated with increased intracellular cholesterol levels.

To address the effect of HCMV-mediated induction of IDOL on UL136p33 levels in a

productive infection, we again induced IDOL expression with the LXR agonist GW3965

or HBX41-108 in TB40/E-$UL136_{myc}$- or -$UL136_{myc}\Delta K{\rightarrow}R$-infected fibroblasts. IDOL

induction resulted in a 50% reduction in the levels of UL136p33 but did not significantly

affect $p33_{\Delta K \to R}$ levels (SI Fig. 2C and D). Diminished levels of smaller UL136 isoforms are

again apparent in the *UL136*$_{myc}\Delta K \to R$ infection. We next disrupted *IDOL* expression in

fibroblasts to answer the question whether IDOL depletion would rescue UL136p33

turnover. CRISPR-mediated knockout (KO) of *IDOL* using guide RNAs to disrupt all the

isoforms of IDOL was validated by Sanger sequencing and increased ApoER-2 as a proxy

for *IDOL* KO (SI Fig. 2E). Further, CRISPR KO of *IDOL* had no effect on HDAC1, a

potential off-target hit when knocking out *IDOL*. In fibroblasts depleted of IDOL, UL136p33

levels increased at 2 and 3 dpi compared to luciferase KO controls, but did not affect

levels of $p33_{\Delta K \to R}$ (SI Fig. 2F and quantified in 2G). Further, IE2 levels were also

significantly increased in IDOL KO cells infected with *UL136*$_{myc}$, but not changed by IDOL

KO in the *UL136*$_{myc}$K$\to$R infection (SI Fig. 2H). Together, these results indicate that IDOL

regulates UL136p33 concentration, which indirectly impacts IE expression in the context

of infection.

IDOL restricts HCMV reactivation

HCMV latency is studied in a variety of models. Due to limitations unique to each model,

there is no perfect model and, therefore, we explore latency questions using multiple

complementary models. To explore the role of IDOL in regulating UL136p33 levels in

latency and reactivation, we will use two models of HCMV: (1) THP-1 cell line model and

(2) CD34+ primary HPC model. In the THP-1 monocytic cell line model, cells are infected

in an undifferentiated state for the establishment of latency. Viral gene expression is

detected early (1 dpi) following infection, but is silenced as the virus establishes a latent-

like state (49). Re-expression of viral genes for reactivation can be stimulated by differentiating the cells into macrophages with a phorbol ester, tetradecanoylphorbol acetate (TPA) (50-55). This is a strong model for looking at transcriptional changes associated with latency and reactivation; however, this model is limited in that THP-1 cells do not robustly or consistently support viral genome synthesis or production of viral progeny following TPA induction of reactivation. The gold-standard for HCMV latency is the ability of the virus to establish and reactivate from latency in primary human CD34+ HPCs (56). The primary CD34+ HPC model supports viral genome synthesis and the formation of infectious progeny in response to reactivaiton, the gold-standard for reactivation. However, the model is limited by substantial donor-variability of primary human cells, heterogeneity of the population, the availability of the cells in numbers to provide sufficient starting material for some assays, and the ability to manipulate these cells (e.g. gene knockdowns or overexpression) without affecting their differentiation, which impacts HCMV infection. Further, because reactivation of HCMV from CD34+ HPCs occurs at low frequency (1 in ~9,000 cells), analysis of molecular events associated with reactivation is challenging. The use of both models yields complementary insights into the patterns of gene expression upon infection or reactivation (THP-1), genome synthesis and virus production (CD34+ HPCs) to understand latency and reactivation.

Dynamic gene expression changes are associated with monocyte-to-macrophage differentiation (31, 57, 58). Because IDOL is highly expressed in bone marrow (59), we hypothesized that *IDOL* expression may be dictated by the differentiation state of the cell and would regulate UL136 levels in a differentiation-dependent manner. Indeed, we found

that *IDOL* was highly expressed in undifferentiated THP-1 monocytes prior to infection,

but was sharply downregulated upon THP-1 cell differentiation (Fig. 4A). Further, while

no significant differences in *IDOL* expression were observed between uninfected cells

and a $UL136_{myc}$ infection, $UL136_{myc}ΔK→R$ infection significantly downregulated *IDOL*

expression in the undifferentiated, latent phase (Fig. 4A). This result suggests that

UL136p33 levels may feedback to impact IDOL expression in a cell type-dependent

manner.

To determine how IDOL impacts HCMV infection in hematopoietic cells, we knocked out

IDOL in THP-1 cells using CRISPR. The knockdown was validated using an ELISA assay

for the detection of IDOL protein (Fig. 4B); IDOL protein was reduced in undifferentiated

cells to levels induced by differentiation. Further, *IDOL* KO had no off-target effects on

HDAC1 (Fig. 4C). Given the challenges of detecting UL136 isoforms, particularly in

hematopoietic cells, we examined IE1, IE2 and UL135 gene expression as proxies for

determining the effect of *IDOL* KO on infection in THP-1 cells. UL135 and IE1/2 proteins

are detected early during infection in THP-1 cells but diminish as the virus establishes

latency between 3 and 5 dpi (Fig. 4D-E) (50). *IDOL* KO resulted in an increased initial

burst in the expression of IE1, IE2, and UL135 at 1 dpi relative to control *Luc* KO cells.

Notably, the major HCMV transactivator, IE2, was only detected in the IDOL KO condition.

The increased viral gene expression in the IDOL KO is consistent with increased viral

activity and UL136p33 levels. However, gene expression was still relatively quieted by 5

dpi, suggesting not unexpectedly that other viral or cellular factors are required for

progression of rectivation. However, as we have never identified a condition or mutant

virus that is not ultimately silenced in THP-1 cells, it may be that the program for restriction

and silencing in this model is robust as indicated by the inability of these cells to support

viral DNA synthesis and progeny formation even with TPA treatment. Importantly, *IDOL*

KO had no effect on IE1, IE2, or UL135 expression in *UL136$_{myc}$ΔK→R* infection (Fig. 4D-

E), consistent with a role for IDOL in regulating the establishment of latency by targeting

UL136p33 for turnover. In contrast to *UL136*$_{myc}$ infection, IE2 is detected in all conditions

of the *UL136*$_{myc}$K→R infection and is not increased further by IDOL KO. While a potent

transactivator, IE2 feeds back to negatively regulate the major immediate early promoter

and its own expression (60-63). The differences in IE2 expression in IDOL KO in *UL136*$_{myc}$

and *UL136*$_{myc}$K→R infection may reflect complexities in the regulation of IE gene

expression and the possible dysregulation and loss of middle UL136 isoforms in

UL136$_{myc}$K→R infection (Fig. 2). These results indicate that IDOL negatively regulates

viral gene expression for the establishment of latency in undifferentiated THP-1 cells,

consistent with its role in maintaining low levels of UL136p33 and the role of UL136p33

in driving HCMV replication in hematopoietic cells (64). Further, these data indicate that

UL136p33 may negatively regulate IDOL expression.

To further explore the importance of IDOL in latency, we turned to the primary CD34+

HPC experimental model. We purified (GFP+) CD34+ HPCs infected with TB40/E-

UL136$_{myc}$ or -*UL136*$_{myc}$K→R by fluorescent activated cell sorting and seeded the cells into

long-term bone marrow cultures. IDOL was induced at 5 dpi after the establishment of

latency by GW396 treatment. IDOL induction was validated by the increased IDOL

transcripts in GW3965-treated cells compared to DMSO-treated cells (Fig. 5A). At 10 dpi,

the culture was split and half the cells were seeded by limiting dilution into co-culture with

fibroblasts and cytokine-rich media to stimulate reactivation from latency. The other half

were mechanically lysed and lystates were seeded by limiting dilution onto fibroblast

monolayers to determine the virus present prior to reactivation (pre-reactivation). Relative

to the DMSO control, GW3965 treatment restricted $UL136_{myc}$ reactivation (Fig. 5B),

consistent with a requirement for accumulation of UL136p33 for reactivation. Strikingly,

reactivation of $UL136_{myc}\Delta K\rightarrow R$ was not restricted by GW3965-induction of IDOL. Given

the limited availability of human CD34+ HPCs, data from two independent donors are

shown. These results indicate that IDOL restricts virus reactivation in a manner

dependent on its ability to turnover UL136p33. Taken together with the THP-1

experiments, these results support the hypothesis that UL136p33 stability is required for

reactivation from latency and that induction of IDOL at reactivation suppresses UL136p33

from accumulating to levels sufficient for reactivation (Fig. 6).

Discussion

Latent viruses require mechanisms to sense the cellular environment, filter noise and

respond to host cues to make solid decisions to maintain a quiescent latent state or to

reactivate replication (1). While the decision to enter or exit latency is often depicted as a

simple binary on/off switch, the reality is more likely that a number of thresholds must be

reached to clear checkpoints in the commitment to productive reactivation. Developing

strategies to control entry into or exit from CMV latency and ultimately control CMV

disease rests upon identifying the virus-host interactions that govern the decisions around

latency and reactivation. Here we demonstrate the existence of an unstable viral

determinant of reactivation, UL136p33. We demonstrate that its instability is dependent

upon a host E3 ligase, IDOL, which is regulated by hematopoietic differentiation. Our

results support a model whereby UL136p33 is maintained at low levels for the

establishment of latency because IDOL levels are high (Fig. 6). Upon differentiation, IDOL

levels fall, allowing UL136p33 levels to rise, which facilitates HCMV crossing at least one

threshold towards reactivation. UL136p33 accumulation is amplified and re-enforced by

the onset of viral DNA synthesis and entry into late phase (21) and high levels of

UL136p33 suppresses IDOL expression (Fig. 4A). These findings define novel virus-host

interactions that govern entry into and exit from latency. Moreover, these findings suggest

that by utilizing IDOL to regulate levels of a critical determinant for reactivation, the virus

evolved to hardwire this decision into LXR signaling and the the regulation of intracellular

cholesterol, which ultimately controls IDOL levels.

IDOL is transcriptionally upregulated by excess cholesterol derivatives (oxysterols) that

are sensed by LXR and in a cell type-dependent manner (35, 65-68). RNA sequencing to

classify the tissue-specific expression of genes across a representative set of all major

human organs and tissues revealed that IDOL is highly upregulated in the bone marrow,

where CD34+ HPCs and monocytes are generated (69). The level of IDOL expression in

the BM is only second to the placenta. High expression of IDOL in undifferentiated

progenitor cells suggests they are intolerant of high cholesterol levels. Consistent with

this, undifferentiated THP-1 monocytes express significantly higher levels of cholesterol

efflux genes, such as CES1, and express significantly lower levels of influx genes (70).

By contrast, THP-1 monocyte-derived macrophages upregulate cholesterol influx genes,

such as scavenger receptor A1 (SR-A1) and CD36 to become lipid-loaded macrophages

or foam cells. Further, statin treatment to lower cholesterol reduces CD34+ HPC

populations in the circulation and high cholesterol promotes CD34+ HPC proliferation,

myelopoiesis as well as monocyte and granulocytic differentiation (71-73). While carefully

designed studies are required to understand how statins specifically affect HCMV latency

and/or reactivation, these findings suggest the possibility that untreated

hypercholesterolemia might lead to increased hematopoietic differentiation and a higher

frequency of HCMV reactivation events, whereas statin treatments may restrict HCMV

reactivation and replication. Given the role demonstrated here for IDOL in the regulation

of cholesterol and UL136p33 expression, the high levels of IDOL expression make

progenitor cells an ideal environment for HCMV latency in which UL136p33 is suppressed

by rapid turnover.

HCMV infection remodels the cellular lipid metabolic landscape by upregulating *de novo*

cholesterol biosynthesis during a productive infection (44, 47, 48). It is, therefore,

expected that HCMV infection would induce IDOL expression. While IDOL and

cholesterol are induced by infection in fibroblasts (SI Fig. 2A-B), IDOL expression was

not induced by infection in hematopoietic cells. In fact, IDOL expression was

downregulated by *UL136*$_{myc}$K$\rightarrow$R infection (Fig. 4A). While cholesterol increases in

macrophage cells, their higher tolerance to cholesterol than HPCs or monocytes (74-76)

may not necessarily correspond to higher levels of IDOL, as we have shown. Accordingly,

despite increase cholesterol in macrophages, IDOL levels are strikingly downregulated in

THP-1 cells induced to differentiate with TPA (Fig. 4A).

IDOL recognizes a conserved motif WxxKNxxSI/MxF within the cytoplasmic tail of its targets (ApoER-2, LDLR and VLDLR), to mediate ubiquitination for lysosomal degradation (30, 34, 35, 77). Canonical IDOL targets reside at the plasma membrane and are typically internalized via the endocytosis and trafficked to the lysosome (77). By contrast, UL136p33 is predominantly Golgi-associated (21), lacks a WxxKNxxSI/MxF motif, and is degraded via the proteasome (Fig. 1B). Differences in localization and trafficking pathways between natural IDOL targets and UL136p33 may differentially direct the degradation pathway.

Activation of LXR in human foreskin fibroblasts with GW3965 prior to HCMV infection has been shown to depress expression levels of viral proteins, such as IE2, pp28, pp65 and gB (78). GW3965 treatment is also reported to induce IFN-γ (79), a cytokine which directly inhibits IE1/2 mRNA expression (80). These findings might suggest that possibility that the effects of LXR agonists on UL136p33 levels are due to a global repression of viral gene expression. However, our results argue against this scenario. First, we observed no effect on IE gene expression at 2 dpi in infected fibroblasts treated with GW3965 or HBX41-108 (Fig. S1C), consistent with the attenuation of IFN-γ at later times (81). Second, specific depletion of *IDOL* increases UL136p33, consistent with IDOL-specific effects on UL136p33 levels. Nevertheless, to avoid this potential pitfall, cells were treated with these agonists following the establishment of infection and IE gene expression in our studies.

The LXR axis is important for the biology of other herpesviruses. A recent report demonstrates that the alpha herpesvirus pseudorabies virus (PRV), etiologic agent of the economically important Aujeszky's disease, has evolved to inhibit LXR at both transcription and protein levels (82). Using an inverse LXR agonist SR9243, Wang and colleagues showed that inhibiting LXR increased PRV replication, whereas 3 different LXR agonists and the oxysterol 22R-hydroxycholesterol all inhibited PRV replication (82). However, PRV can be rescued from the suppressive effects of an LXR agonist by supplementing with cholesterol (82), consistent with a viral need for cholesterol to replicate. LXR also inhibits the murine gamma herpesvirus MHV68 as it replicates better in *LXR$^{-/-}$* macrophages and, contrary to PRV, MHV68 infection increases LXRα (83). Importantly, LXRα restricts reactivation of MHV86 in peritoneal cells, but not in splenocytes (84). The cell type-dependent role of LXR signaling for reactivation suggests the possibility that a LXR signaling might control a MHV68 determinant of reactivation in peritoneal cells that is dispensable in splenocytes. Similarly, disruption of or stabilization of UL136p33 did not impact replication in fibroblasts (Fig. 2C) (21), but did impact replication in HPCs and endothelial cells (Fig. 2D) (20). Viral factors encoded by MHV68 and regulated by LXR signaling have yet to be determined.

In addition to herpesviruses, LXR signaling suppresses the replication of other viruses, including Hepatitis C Virus (HCV) (85, 86), human immunodeficiency virus (HIV) (87-89),

and Newcastle disease virus (NDV) (90). Intriguingly, activation of LXR signaling

suppresses hepatitis C virus (HCV) replication in an IDOL-dependent manner. The

suppression of HCV is presumably due to the requirement of LDLR for an entry receptor

since once infection is established, IDOL induction does not seem to have an effect on

HCV replication (91). However, there are mixed reports on the requirement for LDLR for

HCV entry or its specific function in HCV replication (92-94).

While many host pathways are implicated in the regulation of herpesvirus latency, the

interplay between viral determinants and their regulation by host mechanisms to

terminate activity is less well appreciated, particularly for HCMV. The expression of HSV-

1 ICP0, a critical protein for reactivation and replication, is suppressed by the host

neuronal microRNA-138 to facilitate latency (95, 96). Moreover, ICP0 was recently shown

to be regulated by the host E3 ligase, TRIM23, which is in turn, countered by the HSV-1

late protein, $\gamma_1 34.5$ (97). While speculative, TRIM23 may moderate ICP0 protein levels to

suppress reactivation, where $\gamma_1 34.5$ ensures the commitment to replication by blocking

TRIM23-mediated turnover of ICP0. Further, the E3 ubiquitin ligase, MDM2, restricts

levels of the Kaposi's sarcoma-associated herpesvirus (KSHV) transactivator,

RTA/ORF50, acting as pro-viral factor for latency (98). RTA/ORF50 in turn induces

degradation of vFLIP through the host Itch E3 ubiquitin ligase to attenuate vFLIP-driven

NFκB signaling for reactivation (99, 100). Coopting and regulating host E3 ligases to

target viral proteins for latency allows infections to respond quickly to host cues (e.g., prior

to changes requiring transcription of genes and translation of proteins) for reactivation.

Further, it will be important to determine if UL136p33 feeds back to regulate *IDOL*

expression based on the observation that stabilization of UL136p33 represses *IDOL* expression in undifferentiated THP-1 cells (Fig. 5A). It will be important to further define the mechanisms by which UL136p33 interacts with host pathways to impact viral gene expression and promote context-dependent replication and reactivation.

The evolution of viruses to depend on host factor-mediated turnover of critical viral determinants is shared across highly divergent systems. Bacteriophages establish a dormant lysogenic state to cope with their variable and unpredictable environment (101). Bacteriophage λ lysogeny depends on the competition between repressors and activators of gene expression that serve as a gatekeeper for the lytic pathways (102, 103). As a primary example, competitive binding of the lysogeny determinant CI and the immediate early gene Cro at regulatory operator elements puts CI-Cro at the heart of a bistable switch controlling lysogenic-lytic decisions. Another determinant, CII, is expressed with later kinetics and is also critical to the commitment to lysogeny (104, 105). CII promotes expression of CI while it represses expression of the anti-terminator Q, that must accumulate to a threshold to drive lytic gene expression. Critical effectors in bacteriophage lysogeny, such as CI and CII, are susceptible to destruction by host proteases that are induced in response to stresses, such as DNA damage (106, 107). As such, the decision to establish lysogeny depends on the accumulation of lysogenic effectors and is responsive to host determinants regulated by environmental cues. Further, while CI is robust, the expression of CII, like UL136, is sensitive to genome copy number or other fluctuations in the system, providing additional checkpoints in regulating the decision.

With respect to the system emerging for HCMV, the *UL133-UL138* locus coordinates the expression of multiple genes impacting infection fate decisions. While UL135 and UL138 are not known to directly regulate gene expression, their opposing impact on host EGFR/PI3K signaling (11, 17, 19) has parallels to the Cro-CI bistable genetic switch. Further, UL136p33 is expressed with later kinetics than the UL135/UL138 determinants, and robust UL136p33 expression is dependent on crossing a threshold of commitment to viral DNA synthesis (21). In addition, UL136p33 is remarkably unstable compared to other *UL133-UL138* genes and other UL136 isoforms, and, as we have shown here, its instability is dependent upon a host E3 ligase regulated by differentiation-linked changes in IDOL expression. Although the UL136p33 protein is undetectable in THP-1 monocytes, KO of *IDOL* increases expression of genes critical for replication (e.g., IEs and UL135). While UL135 is critical for overcoming the repressive effects of UL138 for successful reactivation, we assert that the subsequent accumulation of UL136p33 beyond a threshold is also important for commitment to the replicative state (Fig. 6). It will be important to understand the relationship between UL135 and UL136p33 in the commitment to HCMV reactivation since as both are required for reactivation and how other UL136 isoforms, for example the latency-promoting UL136p23/19 isoforms (20), may contribute to the establishment of latency or counter the progression towards reactivation to maintain a latent infection. This is a particularly intriguing question as the stabilization of UL136p33 not only affects levels of IDOL mRNA, but also affects levels of the middle UL136 isoforms (Fig. 2B). These findings indicate that the LXR-IDOL axis may be a potential target for controlling HCMV reactivation clinically.

Methods

Cells and viruses

Human primary embryonic fibroblasts (MRC-5; ATCC) and human kidney embryonic

293T (HEK 293T; ATCC) cells were maintained in Dulbecco's Essential Media (DMEM).

The DMEM was supplemented with 10% fetal bovine serum (FBS), 10 mM HEPES, 1 mM

sodium pyruvate, 2 mM l-alanyl-glutamine, 0.1 mM nonessential amino acids, 100 U/ml

penicillin, and 100 µg/ml streptomycin. CD34+ hematopoietic progenitor cells (HPCs)

were isolated from de-identified medical waste following bone marrow isolations from

healthy donors for clinical procedures at the Banner-University Medical Center at the

University of Arizona (56). THP-1 cells were maintained in RPMI1640 (Gibco)

supplemented with 10% FBS, 1 mM sodium pyruvate, 0.1 mM β-mercapto-ethanol

(Sigma).

The wild type TB40E/05 strain expressing GFP as a marker for infection (108) was

engineered to fuse the myc epitope tag in frame at the 3' end of *UL136* (*UL136$_{myc}$* virus)

using BAC recombineering as described previously (21). The *UL136$_{myc}$ΔK→R* virus was

constructed following methods previously described (Caviness 2014). See SI for detailed

description of virus recombineering.

CRISPR/CAS9-mediated knock out of *IDOL*

Guide RNAs (gRNAs) used to knock out IDOL were designed using online gRNA design

tools at atum.bio and synthego.com. Sequences of the gRNAs are shown in SI Table 2.

The gRNAs were cloned into lentiCRISPR v2 (109). Lentiviruses to deliver the gRNAs

were generated by co-transfecting the lentiCRISPR v2 constructs with pDMG.2 and

psPAX2 in HEK 293T cells. MRC-5 and THP-1 cells were transduced at a MOI of 3 and

10, respectively. Transduced cells were selected with 2 µg/ml puromycin for 2 days.

Validation of knock-out was done by blotting for ApoER-2, Sanger sequencing or

quantifying levels of IDOL by ELISA. The Synthego gRNA design algorithm

(synthego.com) indicates potential off-target genes for the gRNAs and we selected

HDAC1 to evaluate potential off-target effects by western blot analyses.

IDOL ELISA

We used a sandwich ELISA assay (Cusabio) to quantify IDOL protein levels in THP-1

cells. Cells were washed in cold PBS and resuspended at 10^8 cells/mL followed by

freezing at -20 °C. Following 3 freeze-thaw cycles, IDOL was quantified according to the

manufacturer's protocol. Optical densities were determined using a Clariostar plate

reader (BMG Labtech) and sample values were interpolated from a standard curve in

GraphPad Prism v9.

Cholesterol assay

Fibroblasts were seeded at 3×10^5 cells per well in 6-well dishes and incubated overnight.

Infections were done at a MOI of 1 for 1 h and the inoculum was removed followed by

washing cells twice with PBS. Infected cells were incubated in serum-free media and 50

000 cells were harvested at appropriate time points. Samples were prepared using a

cholesterol Glo bioluminescent assay (Promega) according to the manufacturer's

instructions. Relative light units were measured using a Clariostar Plus plate reader (BMG

Labtech) and sample values were extrapolated from a standard curve in GraphPad Prism

v9.

Latency assays

Latency assays in CD34+ HPCs were carried out as described previously (56). IDOL expression

was induced prior to reactivation by treatment with 1 μM GW3965. Infections for latency in THP-

1 cells were done as described previously (50)

Please see SI for detailed description of materials and methods.

Acknowledgements

This study was supported by the National Institutes of Allergy and Infectious Diseases/

National Institutes of Health (NIAID/NIH) grants AI079059 and AI131598 to FG. We are

grateful for the support of the Flow Cytometry and Human Immune Monitoring Shared

Resource, supported through the Cancer Center Support Grant P30 CA023074 awarded

to University of Arizona. We appreciate technical assistance from Lincoln Gay. We thank

Drs. James Alwine (University of Pennsylvania), Lynn Enquist (Princeton University),

Jean Wilson (University of Arizona), and John Purdy (University of Arizona) for critical

discussions on this work.

References

1.   F. Goodrum, "The complex biology of human cytomegalovirus latency and reactivation" in
Advances in Virus Research, T. C. M. M. Kielian, M. J. Roossinck, Ed. (Academic Press, 2022), vol.
112.
2.   F. Goodrum, W. Britt, E. S. Mocarski, "Cytomegalovirus" in Fields Virology: DNA Viruses, P. M.
Howley, D. M. Knipe, J. L. Cohen, B. A. Damania, Eds. (Wolters Kluwer, 2022), vol. 2, chap. 12.

3.   G. Hahn, R. Jores, E. S. Mocarski, Cytomegalovirus remains latent in a common precursor of
dendritic and myeloid cells. *Proc Natl Acad Sci U S A* **95**, 3937-3942 (1998).

4.   C.-K. Min *et al.*, The Differentiation of Human Cytomegalovirus Infected-Monocytes Is Required
for Viral Replication. **10** (2020).

5.   M. B. Reeves, P. A. MacAry, P. J. Lehner, J. G. P. Sissons, J. H. Sinclair, Latency, chromatin
remodeling, and reactivation of human cytomegalovirus in the dendritic cells of healthy carriers.
Proceedings of the National Academy of Sciences of the United States of America **102**, 4140 (2005).

6.   M. B. Reeves, J. H. Sinclair, Circulating dendritic cells isolated from healthy seropositive donors
are sites of human cytomegalovirus reactivation in vivo. *J Virol* **87**, 10660-10667 (2013).

7.   J. Taylor-Wiedeman, J. G. P. Sissons, L. K. Borysiewicz, J. H. Sinclair, Monocytes are a major site of
persistence of human cytomegalovirus in peripheral blood mononuclear cells. **72**, 2059-2064
(1991).

8.   E. Forte *et al.*, Tumor Necrosis Factor Alpha Induces Reactivation of Human Cytomegalovirus
Independently of Myeloid Cell Differentiation following Posttranscriptional Establishment of
Latency. *mBio* **9**, e01560-01518 (2018).

9.   C. Söderberg-Nauclér, K. N. Fish, J. A. Nelson, Reactivation of Latent Human Cytomegalovirus by
Allogeneic Stimulation of Blood Cells from Healthy Donors. *Cell* **91**, 119-126 (1997).

10.   M. B. Reeves, T. Compton, Inhibition of inflammatory interleukin-6 activity via extracellular signal-
regulated kinase-mitogen-activated protein kinase signaling antagonizes human cytomegalovirus
reactivation from dendritic cells. *J Virol* **85**, 12750-12758 (2011).

11.   J. Buehler *et al.*, Opposing Regulation of the EGF Receptor: A Molecular Switch Controlling
Cytomegalovirus Latency and Replication. *PLOS Pathogens* **12**, e1005655 (2016).

12.   M. P. Weekes *et al.*, Latency-Associated Degradation of the MRP1 Drug Transporter During Latent
Human Cytomegalovirus Infection. *Science* **340**, 199 (2013).

13.   V. T. K. Le, M. Trilling, H. Hengel, The Cytomegaloviral Protein pUL138 Acts as Potentiator of Tumor
Necrosis Factor (TNF) Receptor 1 Surface Density To Enhance ULb'-
Encoded Modulation of TNF-α Signaling. *Journal of Virology* **85**, 13260 (2011).

14.   C. Montag *et al.*, The Latency-Associated UL138 Gene Product of Human Cytomegalovirus
Sensitizes Cells to Tumor Necrosis Factor Alpha (TNF-α) Signaling by Upregulating TNF-α Receptor
1 Cell Surface Expression. **85**, 11409-11421 (2011).

15.   E. R. Albright, C. K. Mickelson, R. F. Kalejta, Human Cytomegalovirus UL138 Protein Inhibits the
STING Pathway and Reduces Interferon Beta mRNA Accumulation during Lytic and Latent
Infections. *mBio* **12**, e0226721-e0226721 (2021).

16.   S. H. Lee, E. R. Albright, J.-H. Lee, D. Jacobs, R. F. Kalejta, Cellular defense against latent
colonization foiled by human cytomegalovirus UL138 protein. *Sci Adv* **1**, e1501164-e1501164
(2015).

17.   M. Umashankar *et al.*, Antagonistic Determinants Controlling Replicative and Latent States of
Human Cytomegalovirus Infection. *Journal of Virology* **88**, 5987 (2014).

18.   M. A. Rak *et al.*, Human Cytomegalovirus UL135 Interacts with Host Adaptor Proteins To Regulate
Epidermal Growth Factor Receptor and Reactivation from Latency. **92**, e00919-00918 (2018).

19.   J. Buehler *et al.*, Host signaling and EGR1 transcriptional control of human cytomegalovirus
replication and latency. *PLOS Pathogens* **15**, e1008037 (2019).

20.   K. Caviness *et al.*, Complex Interplay of the UL136 Isoforms Balances
Cytomegalovirus Replication and Latency. *mBio* **7**, e01986-01915 (2016).

21.   K. Caviness, L. Cicchini, M. Rak, M. Umashankar, F. Goodrum, Complex expression of the UL136
gene of human cytomegalovirus results in multiple protein isoforms with unique roles in
replication. *J Virol* **88**, 14412-14425 (2014).

22.   I. Mikell *et al.*, HCMV miR-US22 down-regulation of EGR-1 regulates CD34+ hematopoietic
702   progenitor cell proliferation and viral reactivation. *PLoS Pathog* **15**, e1007854 (2019).
703   23.   I. M. Min *et al.*, The Transcription Factor EGR1 Controls Both the Proliferation and Localization of
704   Hematopoietic Stem Cells. *Cell Stem Cell* **2**, 380-391 (2008).
705   24.   D. Collins-McMillen, J. Kamil, N. Moorman, F. Goodrum, Control of Immediate Early Gene
706   Expression for Human Cytomegalovirus Reactivation. *Frontiers in cellular and infection*
707   *microbiology* **10**, 476 (2020).
708   25.   B. A. Krishna, A. B. Wass, C. M. O'Connor, Activator protein-1 transactivation of the major
709   immediate early locus is a determinant of cytomegalovirus reactivation from latency. *Proc Natl*
710   *Acad Sci U S A* **117**, 20860-20867 (2020).
711   26.   A. E. Hale *et al.*, FOXO transcription factors activate alternative major immediate early promoters
712   to induce human cytomegalovirus reactivation. *Proc Natl Acad Sci U S A* **117**, 18764-18770 (2020).
713   27.   V. G. Kew, J. Yuan, J. Meier, M. B. Reeves, Mitogen and stress activated kinases act co-operatively
714   with CREB during the induction of human cytomegalovirus immediate-early gene expression from
715   latency. *PLoS Pathog* **10**, e1004195 (2014).
716   28.   L. B. Crawford *et al.*, Human Cytomegalovirus Encodes a Novel FLT3 Receptor Ligand Necessary
717   for Hematopoietic Cell Differentiation and Viral Reactivation. *MBio* **9** (2018).
718   29.   M. H. Hancock *et al.*, Human Cytomegalovirus UL7, miR-US5-1, and miR-UL112-3p Inactivation of
719   FOXO3a Protects CD34(+) Hematopoietic Progenitor Cells from Apoptosis. *mSphere* **6** (2021).
720   30.   E. Scotti *et al.*, Targeted disruption of the idol gene alters cellular regulation of the low-density
721   lipoprotein receptor by sterols and liver x receptor agonists. *Mol Cell Biol* **31**, 1885-1893 (2011).
722   31.   R. G. de Bruin *et al.*, Quaking promotes monocyte differentiation into pro-atherogenic
723   macrophages by controlling pre-mRNA splicing and gene expression. *Nat Commun* **7**, 10846
724   (2016).
725   32.   Y. Liu, M. Tingart, S. Lecouturier, J. Li, J. Eschweiler, Identification of co-expression network
726   correlated with different periods of adipogenic and osteogenic differentiation of BMSCs by
727   weighted gene co-expression network analysis (WGCNA). *BMC Genomics* **22**, 254 (2021).
728   33.   E. Scotti *et al.*, IDOL stimulates clathrin-independent endocytosis and multivesicular body-
729   mediated lysosomal degradation of the low-density lipoprotein receptor. *Mol Cell Biol* **33**, 1503-
730   1514 (2013).
731   34.   C. Hong *et al.*, The E3 ubiquitin ligase IDOL induces the degradation of the low density lipoprotein
732   receptor family members VLDLR and ApoER2. *J Biol Chem* **285**, 19720-19726 (2010).
733   35.   N. Zelcer, C. Hong, R. Boyadjian, P. Tontonoz, LXR regulates cholesterol uptake through Idol-
734   dependent ubiquitination of the LDL receptor. *Science* **325**, 100-104 (2009).
735   36.   M. H. Glickman, A. Ciechanover, The ubiquitin-proteasome proteolytic pathway: destruction for
736   the sake of construction. *Physiological reviews* **82**, 373-428 (2002).
737   37.   C. M. Pickart, Back to the future with ubiquitin. *Cell* **116**, 181-190 (2004).
738   38.   J. Zhang, Q. Liu, Cholesterol metabolism and homeostasis in the brain. *Protein Cell* **6**, 254-264
739   (2015).
740   39.   L. Zhang, K. Reue, L. G. Fong, S. G. Young, P. Tontonoz, Feedback regulation of cholesterol uptake
741   by the LXR-IDOL-LDLR axis. *Arterioscler Thromb Vasc Biol* **32**, 2541-2546 (2012).
742   40.   J. Gao *et al.*, The E3 ubiquitin ligase IDOL regulates synaptic ApoER2 levels and is important for
743   plasticity and learning. *Elife* **6**, e29178 (2017).
744   41.   J. K. Nelson *et al.*, Deubiquitylase Inhibition Reveals Liver X Receptor-independent Transcriptional
745   Regulation of the E3 Ubiquitin Ligase IDOL and Lipoprotein Uptake. *J Biol Chem* **291**, 4813-4825
746   (2016).
42.   Anonymous, Cytomegalovirus. *Am J Transplant* **4 Suppl 10**, 51-58 (2004).

43.   Y. Du, G. Zhang, Z. Liu, Human cytomegalovirus infection and coronary heart disease: a systematic
review. *Virol J* **15**, 31-31 (2018).
44.   N. Gudleski-O'Regan, T. M. Greco, I. M. Cristea, T. Shenk, Increased expression of LDL receptor-
related protein 1 during human cytomegalovirus infection reduces virion cholesterol and
infectivity. *Cell Host Microbe* **12**, 86-96 (2012).
45.   Y. Xi, S. Harwood, L. M. Wise, J. G. Purdy, Human Cytomegalovirus pUL37x1 Is Important for
Remodeling of Host Lipid Metabolism. *J Virol* **93** (2019).
46.   Y. Xi, L. Lindenmayer, I. Kline, J. von Einem, J. G. Purdy, Human Cytomegalovirus Uses a Host Stress
Response To Balance the Elongation of Saturated/Monounsaturated and Polyunsaturated Very-
Long-Chain Fatty Acids. *mBio* **12** (2021).
47.   Y. Yu, T. G. Maguire, J. C. Alwine, Human cytomegalovirus infection induces adipocyte-like
lipogenesis through activation of sterol regulatory element binding protein 1. *J Virol* **86**, 2942-
2949 (2012).
48.   Y. Yu, F. J. Pierciey, Jr., T. G. Maguire, J. C. Alwine, PKR-like endoplasmic reticulum kinase is
necessary for lipogenic activation during HCMV infection. *PLoS Pathog* **9**, e1003266 (2013).
49.   D. Collins-McMillen *et al.*, Alternative promoters drive human cytomegalovirus reactivation from
latency. *Proc Natl Acad Sci U S A* **116**, 17492-17497 (2019).
50.   D. Collins-McMillen *et al.*, Alternative promoters drive human cytomegalovirus reactivation from
latency. *Proceedings of the National Academy of Sciences* **116**, 17492 (2019).
51.   M.-C. Arcangeletti *et al.*, Human cytomegalovirus reactivation from latency: validation of a
"switch" model in vitro. *Virol J* **13**, 179-179 (2016).
52.   L.-F. Yee, P. L. Lin, M. F. Stinski, Ectopic expression of HCMV IE72 and IE86 proteins is sufficient to
induce early gene expression but not production of infectious virus in undifferentiated
promonocytic THP-1 cells. *Virology* **363**, 174-188 (2007).
53.   B. G. Weinshenker, S. Wilton, G. P. Rice, Phorbol ester-induced differentiation permits productive
human cytomegalovirus infection in a monocytic cell line. *J Immunol* **140**, 1625-1631 (1988).
54.   N. Wagenknecht *et al.*, Contribution of the Major ND10 Proteins PML, hDaxx and Sp100 to the
Regulation of Human Cytomegalovirus Latency and Lytic Replication in the Monocytic Cell Line
THP-1. *Viruses* **7**, 2884-2907 (2015).
55.   C. H. Lee *et al.*, Factors affecting human cytomegalovirus gene expression in human monocyte cell
lines. *Mol Cells* **9**, 37-44 (1999).
56.   M. Peppenelli, J. Buehler, F. Goodrum, Human Hematopoietic Long-Term Culture (hLTC) for
Human Cytomegalovirus Latency and Reactivation. *Methods Mol Biol* **2244**, 83-101 (2021).
57.   R. N. Ramirez *et al.*, Dynamic Gene Regulatory Networks of Human Myeloid Differentiation. *Cell
systems* **4**, 416-429.e413 (2017).
58.   F. O. Martinez, S. Gordon, M. Locati, A. Mantovani, Transcriptional Profiling of the Human
Monocyte-to-Macrophage Differentiation and Polarization: New Molecules and Patterns of Gene
Expression. *The Journal of Immunology* **177**, 7303 (2006).
59.   B. Fagerberg, J. Gnarpe, H. Gnarpe, S. Agewall, J. Wikstrand, Chlamydia pneumoniae but not
cytomegalovirus antibodies are associated with future risk of stroke and cardiovascular disease:
a prospective study in middle-aged to elderly men with treated hypertension. *Stroke* **30**, 299-305
(1999).
60.   M. C. Pizzorno, G. S. Hayward, The IE2 gene products of human cytomegalovirus specifically down-
regulate expression from the major immediate-early promoter through a target located near the
cap site. *J Virol* **64**, 6154-6165 (1990).
61.   J. M. Cherrington, E. L. Khoury, E. S. Mocarski, Human cytomegalovirus ie2 negatively regulates
alpha gene expression via a short target sequence near the transcription start site. *J Virol* **65**, 887-
896 (1991).

62.   B. Liu, T. W. Hermiston, M. F. Stinski, A cis-acting element in the major immediate early (IE)
promoter of human cytomegalovirus is required for negative regulation by IE2. *J Virol* **65**, 897-903
(1991).
63.   M. P. Macias, M. F. Stinski, An in vitro system for human cytomegalovirus immediate early 2
protein (IE2)-mediated site-dependent repression of transcription and direct binding of IE2 to the
major immediate early promoter. *Proc Natl Acad Sci U S A* **90**, 707-711 (1993).
64.   K. Caviness *et al.*, Complex Interplay of the UL136 Isoforms Balances Cytomegalovirus Replication
and Latency. *mBio* **7**, e01986 (2016).
65.   S. D. Lee, P. Tontonoz, Liver X receptors at the intersection of lipid metabolism and atherogenesis.
Atherosclerosis **242**, 29-36 (2015).
66.   B. A. Janowski *et al.*, Structural requirements of ligands for the oxysterol liver X receptors LXRalpha
and LXRbeta. *Proc Natl Acad Sci U S A* **96**, 266-271 (1999).
67.   B. A. Janowski, P. J. Willy, T. R. Devi, J. R. Falck, D. J. Mangelsdorf, An oxysterol signalling pathway
mediated by the nuclear receptor LXRα. *Nature* **383**, 728-731 (1996).
68.   D. J. Peet, B. A. Janowski, D. J. Mangelsdorf, The LXRs: a new class of oxysterol receptors. *Current
Opinion in Genetics & Development* **8**, 571-575 (1998).
69.   L. Fagerberg *et al.*, Analysis of the human tissue-specific expression by genome-wide integration
of transcriptomics and antibody-based proteomics. *Molecular & cellular proteomics : MCP* **13**,
397-406 (2014).
70.   I. Fernandez-Ruiz, P. Puchalska, C. A. Narasimhulu, B. Sengupta, S. Parthasarathy, Differential lipid
metabolism in monocytes and macrophages: influence of cholesterol loading[S]. *Journal of Lipid
Research* **57**, 574-586 (2016).
71.   T. R. Cimato, B. A. Palka, J. K. Lang, R. F. Young, LDL Cholesterol Modulates Human CD34+ HSPCs
through Effects on Proliferation and the IL-17 G-CSF Axis. *PLOS ONE* **8**, e73861 (2013).
72.   P. K. Morgan, L. Fang, G. I. Lancaster, A. J. Murphy, Hematopoiesis is regulated by cholesterol
efflux pathways and lipid rafts: connections with cardiovascular diseases: Thematic Review Series:
Biology of Lipid Rafts. *Journal of Lipid Research* **61**, 667-675 (2020).
73.   O. Soehnlein, F. K. Swirski, Hypercholesterolemia links hematopoiesis with atherosclerosis. *Trends
in endocrinology and metabolism: TEM* **24**, 129-136 (2013).
74.   J. M. Radley, S. Ellis, M. Palatsides, B. Williams, I. Bertoncello, Ultrastructure of primitive
hematopoietic stem cells isolated using probes of functional status. *Experimental hematology* **27**,
365-369 (1999).
75.   S. Shariatmadar *et al.*, Electronic volume of CD34 positive cells from peripheral blood apheresis
samples. *Cytometry. Part B, Clinical cytometry* **74**, 182-188 (2008).
76.   M. P. Vacanti, A. Roy, J. Cortiella, L. Bonassar, C. A. Vacanti, Identification and initial
characterization of spore-like cells in adult mammals. *Journal of cellular biochemistry* **80**, 455-460
(2001).
77.   A. C. Calkin *et al.*, FERM-dependent E3 ligase recognition is a conserved mechanism for targeted
degradation of lipoprotein receptors. **108**, 20107-20112 (2011).
78.   B. Liu *et al.*, Inhibition of Human Cytomegalovirus Particle Maturation by Activation of Liver X
Receptor. *Front Microbiol* **13**, 846386-846386 (2022).
79.   Q. Wang *et al.*, Identification of interferon-γ as a new molecular target of liver X receptor.
Biochemical Journal **459**, 345-354 (2014).
80.   B. Sainz, H. L. LaMarca, R. F. Garry, C. A. Morris, Synergistic inhibition of human cytomegalovirus
replication by interferon-alpha/beta and interferon-gamma. *Virol J* **2**, 14 (2005).
81.   K. A. Kropp *et al.*, Reversible inhibition of murine cytomegalovirus replication by gamma
interferon (IFN-γ) in primary macrophages involves a primed type I IFN-signaling subnetwork for

full establishment of an immediate-early antiviral state. *Journal of virology* **85**, 10286-10299
(2011).
82.   Y. Wang *et al.*, Pseudorabies Virus Inhibits Expression of Liver X Receptors to Assist Viral Infection.
Viruses **14**, 514 (2022).
83.   P. T. Lange, C. Schorl, D. Sahoo, V. L. Tarakanova, Liver X Receptors Suppress Activity of Cholesterol
and Fatty Acid Synthesis Pathways To Oppose Gammaherpesvirus Replication. *mBio* **9**, e01115-
01118 (2018).
84.   P. T. Lange, C. N. Jondle, E. J. Darrah, K. E. Johnson, V. L. Tarakanova, LXR Alpha Restricts
Gammaherpesvirus Reactivation from Latently Infected Peritoneal Cells. *Journal of virology* **93**,
e02071-02018 (2019).
85.   J. Zeng *et al.*, Liver X receptors agonists impede hepatitis C virus infection in an Idol-dependent
manner. *Antiviral Res* **95**, 245-256 (2012).
86.   S. Bocchetta *et al.*, Up-regulation of the ATP-binding cassette transporter A1 inhibits hepatitis C
virus infection. *PLoS One* **9**, e92140 (2014).
87.   H. Jiang *et al.*, Retinoic acid and liver X receptor agonist synergistically inhibit HIV infection in CD4+
T cells by up-regulating ABCA1-mediated cholesterol efflux. *Lipids Health Dis* **11**, 69 (2012).
88.   L. Dubrovsky *et al.*, Liver X receptor agonist inhibits HIV-1 replication and prevents HIV-induced
reduction of plasma HDL in humanized mouse model of HIV infection. *Biochem Biophys Res*
Commun **419**, 95-98 (2012).
89.   T. M. Hanley, G. A. Viglianti, Nuclear receptor signaling inhibits HIV-1 replication in macrophages
through multiple trans-repression mechanisms. *J Virol* **85**, 10834-10850 (2011).
90.   X. X. Sheng *et al.*, The LXR ligand GW3965 inhibits Newcastle disease virus infection by affecting
cholesterol homeostasis. *Arch Virol* **161**, 2491-2501 (2016).
91.   J. Zeng *et al.*, Liver X receptors agonists impede hepatitis C virus infection in an Idol-dependent
manner. *Antiviral Research* **95**, 245-256 (2012).
92.   G. H. Syed *et al.*, Hepatitis C virus stimulates low-density lipoprotein receptor expression to
facilitate viral propagation. *Journal of virology* **88**, 2519-2529 (2014).
93.   A. Albecka *et al.*, Role of low-density lipoprotein receptor in the hepatitis C virus life cycle.
Hepatology (Baltimore, Md.) **55**, 998-1007 (2012).
94.   S. Molina *et al.*, The low-density lipoprotein receptor plays a role in the infection of primary
human hepatocytes by hepatitis C virus. *Journal of hepatology* **46**, 411-419 (2007).
95.   B. Sun *et al.*, Regulation of host and virus genes by neuronal miR-138 favours herpes simplex virus
1 latency. *Nature microbiology* **6**, 682-696 (2021).
96.   D. Pan *et al.*, A neuron-specific host microRNA targets herpes simplex virus-1 ICP0 expression and
promotes latency. *Cell Host Microbe* **15**, 446-456 (2014).
97.   X. Liu *et al.*, Herpesvirus-mediated stabilization of ICP0 expression neutralizes restriction by
TRIM23. *Proc Natl Acad Sci U S A* **118** (2021).
98.   T. H. Chang *et al.*, Regulation of the Abundance of Kaposi's Sarcoma-Associated Herpesvirus
ORF50 Protein by Oncoprotein MDM2. *PLoS Pathog* **12**, e1005918 (2016).
99.   J. C. Chmura *et al.*, The Itch ubiquitin ligase is required for KSHV RTA induced vFLIP degradation.
Virology **501**, 119-126 (2017).
100.   E. S. Ehrlich, J. C. Chmura, J. C. Smith, N. N. Kalu, G. S. Hayward, KSHV RTA abolishes NFkappaB
responsive gene expression during lytic reactivation by targeting vFLIP for degradation via the
proteasome. *PLoS One* **9**, e91359 (2014).
101.   S. Gandon, Why Be Temperate: Lessons from Bacteriophage λ. *Trends in Microbiology* **24**, 356-
365 (2016).

102.   S. Atsumi, J. W. Little, Role of the lytic repressor in prophage induction of phage lambda as
analyzed by a module-replacement approach. *Proceedings of the National Academy of Sciences*
of the United States of America **103**, 4558-4563 (2006).
103.   R. A. Schubert, I. B. Dodd, J. B. Egan, K. E. Shearwin, Cro's role in the CI Cro bistable switch is critical
for {lambda}'s transition from lysogeny to lytic development. *Genes Dev* **21**, 2461-2472 (2007).
104.   M. G. Cortes, J. T. Trinh, L. Zeng, G. Balazsi, Late-Arriving Signals Contribute Less to Cell-Fate
Decisions. *Biophys J* **113**, 2110-2120 (2017).
105.   O. Kobiler, A. Rokney, A. B. Oppenheim, Phage lambda CIII: a protease inhibitor regulating the
lysis-lysogeny decision. *PLoS One* **2**, e363 (2007).
106.   J. W. Little, D. P. Shepley, D. W. J. T. E. j. Wert, Robustness of a gene regulatory circuit. **18**, 4299-
4307 (1999).
107.   J. D. Hernandez-Doria, V. Sperandio, Bacteriophage Transcription Factor Cro Regulates Virulence
Gene Expression in Enterohemorrhagic Escherichia coli. *Cell Host Microbe* **23**, 607-617.e606
(2018).
108.   M. Umashankar *et al.*, A Novel Human Cytomegalovirus Locus Modulates Cell Type-Specific
Outcomes of Infection. *PLOS Pathogens* **7**, e1002444 (2011).
109.   N. E. Sanjana, O. Shalem, F. Zhang, Improved vectors and genome-wide libraries for CRISPR
screening. *Nat Methods* **11**, 783-784 (2014).

Figure legends

Figure 1. UL136p33 isoform is rapidly turned over by the proteasome. (A) Schematic

depiction of the *UL133-UL138* polycistronic gene locus and UL136 protein isoforms. A C-

terminal myc tag is used to detect UL136 isoforms. Red lines represent lysine residues,

and the black box shows the putative transmembrane domain. (B-D) Fibroblasts were

infected with TB40/E-*UL136$_{myc}$* virus at a MOI of 1. (B) At 48 hpi, cells were treated with

50 µg/ml cycloheximide (CHX) for 6 h. Alternatively, mock- or TB40/E-infected cells were

treated with CHX in the context of proteasomal (MG132) or lysosomal (NH$_4$Cl) inhibition

(right panel). UL136$_{myc}$ protein isoforms or tubulin as a loading control were detected by

immunoblotting using by a monoclonal α-myc antibody or tubulin, respectively (left panel).

(C) Cells were untreated or treated with CHX at 48 hpi and UL136 decay was analyzed

over a 6-hour time course by immunoblotting. (D) Relative quantification of UL136 isoforms normalized to tubulin. HPT, hours post treatment

Figure 2. Stabilizing UL136p33 results in virus that fails to establish latency. (A) Fibroblasts were transduced with lentiviruses engineered to express $p33_{myc}$ or $p33_{K \to R}$ and protein decay was analyzed by immunoblotting following CHX treatment. (B) Fibroblasts were infected with TB40/E-$UL136_{myc}$ or $-UL136_{myc}\Delta K \to R$ viruses (MOI =1) and treated with CHX at 48 hpi or left untreated. Protein decay was analyzed as described in Figure 1C. (C) Fibroblasts were infected at a MOI of 0.02 and samples collected for $TCID_{50}$ at indicated times. All data points are an average of 3 biological replicates and error bars represent standard deviation. (D) CD34+ HPCs were infected with $UL136_{myc}$ or $UL136_{myc}\Delta K \to R$ viruses (MOI=2). Infected cells were purified by FACS at 1 dpi seeded into long term culture. At 10 days, half of the culture was reactivated by seeding intact cells by limiting dilution into co-culture with fibroblasts in a cytokine-rich media to promote myeloid differentiation (reactivation). The other half of the culture was mechanically lysed and seeded onto fibroblasts as the pre-reactivation control. The frequency of infectious centers was determined for each sample by extreme limiting dilution. (C-D) The data represents replicates with cells from 3 independent donors. The means were compared with two-way analysis of variance (ANOVA) with a Tukey post hoc test and the were no significant differences. **, $p<0.01$; ***, $p<0.001$.

Figure 3. The host E3 ligase IDOL specifically targets UL136p33 for turnover. (A) HEK cells were transiently transfected with plasmids expressing UL136p33, IDOL, empty vector, or the indicated combinations. UL136p33 or IDOL was immunoprecipitated from

protein lysates using α-myc or α-HA magnetic beads. Protein was detected by immunoblotting using the antibodies indicated. Lysates were blotted as a control for expression and GAPDH used as a loading control. (B) HEK cells were transiently transfected with plasmids expressing p33$_{\Delta K \to R}$, IDOL, or empty vector. Proteins were detected by immunoblotting using α-myc or α-HA monoclonal antibodies. (C) Protein densities of UL136p33 (from panel A) or p33$_{\Delta K \to R}$ (from panel B) relative to their loading control. (D) HEK cells were transiently transfected to express UL136p33 or p33$_{\Delta K \to R.}$ IDOL was induced 48 hours later with 2 μM of LXR-agonists, GW3965 or HBX41-108, or vehicle (DMSO) for 6 h. Detection of ApoER-2 was used as a proxy for validating the induction of IDOL. p33$_{myc}$, p33$_{K \to R}$, and tubulin as a loading control were detected by immunoblotting using α-myc or α-tubulin. All data points represent the mean of at least 3 independent replicates and the means were compared by two-way ANOVA with a Tukey post hoc test. Error bars represent standard deviation. ns, *p*>0.05. ****, p<0.0001.

Figure 4. IDOL impacts viral protein expression in a THP-1 cell latency model. (A) Total RNA was extracted from mock-, *UL136$_{myc}$*- or *UL136$_{myc}$ΔK➔R*-infected THP-1 cells at indicated time points. At 5 dpi, cells were treated with TPA to induce differentiation or the vehicle control (DMSO). cDNA was synthesized from the RNA and qPCR was used to determine the relative expression levels of *IDOL* transcripts compared to the low copy cellular housekeeping gene *H6PD* using the ΔΔCT method. (B) Validation of *IDOL* KO in undifferentiated THP-1 cells was confirmed by ELISA. Where indicated, cells were treated with TPA as a positive control for the loss of IDOL. (C) Immunoblot of HDAC1 in *Luc* control or *IDOL* KO THP-1 cells to rule out off-target CRISPR/CAS9 effects. (D) *IDOL* or

Luc KO THP-1 cells were infected with TB40/E-*UL136*$_{myc}$ (top panel) or *-UL136$_{myc}$K→R*

(bottom panel) at a MOI of 2 and cell lysates were collected at indicated time points. IE1/2

and UL135 accumulation was evaluated by immunoblotting protein lysates over a time

course using a monoclonal α-IE1/2 and polyclonal α-UL135. (E) IE1/2 and UL135 were

quantified, and levels normalized to Tubulin and levels present at 1 dpi in the Luc KO

control. All data points are representative of the mean 3 independent replicates and error

bars represent standard deviation. The means were compared by two-way ANOVA with

a Tukey post hoc test. ns, $p > 0.05$; *, $p < 0.05$; **, $p < 0.01$; ****, $p < 0.0001$.

Figure 5. Induction of IDOL in CD34+ HPCs inhibits reactivation. (A) CD34+ HPCs

were mock- or TB40/E-infected at a MOI of 2. At 1 dpi, cells were treated with 1 μM

GW3965 for 6 h and total RNA isolated for qPCR to measure *IDOL* transcript levels

relative to *H6PD*. The data points represent the mean from 3 independent experiments.

The means were analyzed by two-way ANOVA with a Tukey post-hoc test. ***$p < 0.001$;

****, $p < 0.0001$. (B) CD34+ HPCs were infected at a MOI of 2. At 24 hpi, infected CD34+

cells were isolated by FACS and seeded into long term culture. GW3965 (1 μM) at 24 hpi.

At 10 days, half of the culture was reactivated by seeding intact cells by limiting dilution

into co-culture with fibroblasts in a cytokine-rich media to promote myeloid differentiation

(reactivation). The other half of the culture was mechanically lysed and seeded onto

fibroblasts as the pre-reactivation control. The frequency of infectious centers was

determined for each sample by ELDA. The data points represent replicates from 2

independent donors and the range is indicated by squares (pre-reactivation) and dots

(post-reactivation).

Figure 6. Model for differentiation-dependent IDOL regulation to control UL136p33 levels. Progenitor cells are relatively small and are intolerant of high cholesterol, which they control by upregulating IDOL. We propose that high levels of IDOL maintain low levels of UL136p33 for the establishment of latency. Upon cell differentiation, cells downregulate IDOL, which would result in a corresponding increase in UL136p33 levels to drive infection towards reactivation. UL136p33 levels are further increased upon the commitment to viral DNA synthesis. Stabilization of UL136p33 results in corresponding downregulation of IDOL mRNAs, suggesting feedback to suppress of IDOL gene expression. This work implicates IDOL as a host factor that fine-tunes levels of UL136p33 to control decisions to enter and exit latency. These findings further indicate a key role for cholesterol homeostasis as a determinant of HCMV reactivation, whereby UL136p33 acts as a sensor at the tipping point between the decision to maintain the latent state or exit latency for reactivation. Cholesterol represented by beige droplets. Figure made with Biorender.com.

1006

1007   **Figure 1**

1008

Figure 2

1013   **Figure 3**

1014

1015

Figure 4

1019 **Figure 5**

A

TB40/E	−	−	+	+
GW3965	−	+	−	+

B

$U136_{myc}$	+	+	−	−
$UL136_{myc}\Delta K \rightarrow R$	−	−	+	+
DMSO	+	+	+	+
GW3965	−	+	−	+

1020

1021 **Figure 6**

1022

 SUPPORTING INFORMATION

1024

1025 **SI Figure 1**

1026

1027

1028

SI Figure 1. IDOL specifically targets the UL136p33 isoform. (A-C) HEK 293T cells were
transfected with plasmids expressing (A) IDOL, p25 or empty vector or (B) IDOL, p23 or empty
vector. Proteins detected by immunoblotting using antibodies to FLAG or HA to detect the
indicated proteins. Tubulin or GAPDH were used as loading controls for normalization. (C)
Relative p25 or p23 densities. (D) HEK 293T cells were co-transfected with plasmids expressing
UL136p33, the E3 ligase Cbl or empty vector. Proteins were detected by immunoblotting using
antibodies to the myc epitope tag or cbl. (E) Relative UL136p33 densities were measured and
quantified. All data points represent the mean of at least 3 independent replicates and the means
were compared by two-way ANOVA with a Tukey post hoc test. Error bars represent standard
deviation. ns, non-significant

1039

1040

SI Figure 2

SI Figure 2. IDOL expression in HCMV-infected fibroblasts impacts UL136p33 stability. (A) Fibroblasts were mock-infected or infected with *UL136$_{myc}$* or *UL136$_{myc}$K→R* viruses at a MOI of 1 and total RNA extracted at indicated time points. RT-qPCR was used to quantify relative *IDOL* transcripts using the ΔΔCT method. *IDOL* transcripts were normalized to GAPDH and compared to expression levels in mock samples at 1 dpi. (B) Fibroblasts were infected with TB40/E at a MOI of 1 followed by incubation in serum-free media. Cholesterol concentrations were measured and compared to mock-infected cells at 1 dpi. (C) Fibroblasts were mock-infected or infected with *UL136$_{myc}$* or *UL136$_{myc}$ΔK→R* viruses at a MOI of 1. At 2 dpi, IDOL expression was induced by treatment with 2 μM GW3965 or HBX41-108 for 6 h followed by western blot evaluation of UL136p33 levels (left). Relative UL136p33 levels are quantified in (D). (E) *IDOL* KO cell lysates were analyzed by western blot to evaluate ApoER-2 expression levels compared to a *Luc* KO control. To evaluate CRISPR/CAS9 off-target effects we measured HDAC1 in *Luc* or *IDOL* KO cell lysates. (F) *UL136$_{myc}$* or *UL136$_{myc}$ΔK→R* viruses were used to infect either *Luc* or *IDOL* KO cells at a MOI of 1. Cell lysates were collected at indicated times and evaluated for UL136p33 expression levels that are quantified in (G). (H) Relative quantification of IE2 protein densities in *Luc* or *IDOL* KO fibroblasts. Quantitative data points represent the mean of 3 independent replicates and error bars represent standard deviation. Data points are means of 3 independent replicates and error bars represent standard deviation. Differences in the means were compared using two-way ANOVA with a Tukey post hoc test. ns, $p > 0.05$; *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.

 SI Table 1. Candidate UL136 interactors from Y2H screen

Name		Known Functions	Accession Number
MYLIP	Myosin regulatory light chain interacting protein	E3 Ubiquitin Ligase that mediates ubiquitination and degradation of myosin regulatory light chain (MRLC)	XM_005249032.2
PIK3R4	Phosphoinositide-3-kinase, regulatory subunit 4	Also called VPS15; serine-threonine protein kinase found at the cytoplasmic face of late Golgi and may regulate membrane trafficking in the late endocytic pathway	NM_014202.2
AP3B1	Adapter-related protein complex 3, beta 1 subunit	Subunit of non-clathrin- and clathrin-associated adaptor protein complex 3 that functions in transmembrane protein sorting to lysosome or lysosomal-related organelles.	NM_003664.4
CTSL2	Cathespin L2	Member of peptidase C1 family; lysosomal cysteine proteinase	NM_001333.3
TNFR-SF11B	Tumor necrosis factor receptor superfamily, member 11B	TNFR superfamily member; may act as a decoy receptor for TRAIL thereby protecting against apoptosis	NM_002546.3
CTNNAL1	Catenin (cadherin-associated protein), alpha-like 1	May modulate Rho pathway and act as anchor for PKA	XM_011519161.1
MARK3	MAP/microtubule affinity-regulating kinase 3	Activated by phosphorylation, then phosphorylates MAP2, MAP4, and CDC25	XM_006720146.2
ZFP42	Zinc finger protein 42	Also called Rex1; marker of pluripotency, regulation is critical to maintaining pluripotent state in ES cells and may be involved in transcriptional regulation	NM_001304358.1

MYLIP	Myosin regulatory light chain interacting protein	E3 Ubiquitin Ligase that mediates ubiqutination and degradation of myosin regulatory light chain (MRLC)	XM_005249032.2
SCEL	Sciellin	Localizes to periphery of cells; is a precursor of the cornified envelope of terminally differentiated keratinocytes	XM_011535281.1
CHCHD4	Coiled-coil-helix-coiled-coil-helix domain containing 4	Mitochondrial chaperone to catalyze disulfide bonds. Required for the import and folding of small cysteine containing proteins into the mitochondrial intermembrane space	NM_001098502.1
ADSL	Adenylosuccinate lyase	De novo synthesis of purines; catalyzes reactions in synthesis of AMP	XM_011529976.1
BTF3	Basic transcription factor 3	Forms stable complex with RNA polymerase II; required for transcription initiation	NM_001037637.1
DARS	Aspartyl-tRNA synthetase	Part of multienzyme complex of aminoacyl-tRNA synthetases; charges its cognate tRNA with aspartate	NM_001349.3
TERF2IP	Telomeric repeat binding factor 2, interacting protein	Part of a complex involved in telomere length regulation	NM_018975.3
CCDC15	Coiled-coil domain containing 15	?	NM_025004.2
CPORF43	Chromosome 9, open reading frame 43	?	NM_152786.2

1080

1081

1082 **SI Table 2. gRNAs used to KO *IDOL* and *luciferase (Luc)* gene (from *Photinus pyralis*)**

Gene	Position	Forward sequence	Reverse sequence
IDOL	16,145,148	CACCG**GCGCCGTGATGATG CAGUAU**[a]	AAACAUACUGCAUCAUCACGG CGCC
IDOL	16,143,526	CACCG**CCUUUUUGUUUUAG GUGUGC**	AAACGUUUCUCAGGUUUAGCC AUAC
IDOL	16,143,613	CACCG**UAUGGCUAAACCUG AGAAAC**	AAACGUUUCUCAGGUUUAGCC AUAC
Luc	1355	CACCG**ACAACUUUACCGAC CGCGCC**	AAAC**GGCGCGGUCGGUAAAGU UGU**

1083 [a]gRNA sequence is in bold text, and the preceding sequence was used to facilitate cloning onto
1084 the lentiCRISPR v2 vector.

1085

1086 **SI Table 3. Antibodies used in this study**

Antibodies		
Rabbit monoclonal anti-ApoER-2	Abcam	Cat#ab108208
Rabbit polyclonal anti-FLAG tag	Sigma-Aldrich	Cat#F7425
Rabbit monoclonal anti-GAPDH (clone D16H11)	Cell Signaling Technology	Cat#5174
Rabbit polyclonal anti-HA tag	Sigma-Aldrich	Cat#H6908
Mouse HDAC1 (clone 10E2)	Santa Cruz Biotech	Cat#sc-81598
IE1/2 (3H4)	Princeton University/Dr. Tom Shenk	N/A
Rabbit monoclonal anti-Myc tag (clone 71D10)	Cell Signaling Technology	Cat#2278
Tubulin (clone DM1A)		Cat#T6199
UL135	Open Biosystems	Custom antibody
Goat anti-mouse secondary antibody (IR Dye 680)	LI-COR Biosciences	Cat#926-68070

| Goat anti-rabbit secondary antibody (IR Dye 800) | LI-COR Biosciences | Cat#926-32211 |

1087

1088

1089

1090

Materials and Methods

1092

Yeast two-hybrid screen

A yeast screen for human cellular interacting proteins of pUL136 was performed using the Matchmaker Gold Yeast Two-Hybrid System (Clontech) and the Mate and Plate Universal Human Library (Clontech). This yeast two-hybrid universal library was constructed from human cDNA that has been normalized to remove high copy number cDNAs (overrepresented transcripts) to facilitate the discovery of low copy number novel protein-protein interactions. All protocols were carried out according to the manufacturer's guidelines. Yeast dual transformation and yeast mating was performed according to established protocols from Clontech. Yeast prey vectors were isolated from Y187 yeast and subsequently transformed into DH10B *E. coli*. After selection for *E coli* containing yeast prey vectors, the plasmids were isolated and sequenced to identify interacting partners.

1104

Bacterial Artificial Chromosome Recombineering

For generation of *UL136*$_{myc}$$\Delta$K$\rightarrow$R, *UL136*$_{myc}$$\Delta$K$\rightarrow$R pGEM-T Easy construct was created by site-directed Phusion mutagenesis according to the manufacturer's instruction (New England Biolabs). Inverse PCR was used to mutagenize lysines 4, 20, 25, and 113 to arginine using the primers: 5'-GACGTTGGAAATAGATGGCGGCGTCGAAGGCCCTGAGTCGC-3' (forward); CAAGTCCCACGTCATTTCTGGCATCTCCACGCCCCTGACTGACAT-3' (reverse) for the first 3 lysine residues in the N-terminal region and 4th lysine was mutagenized to arginine with the forward primer 5'-ACGAGCGGCAGAGAAGCAGACGGC-3', and reverse primer GGTGCCGACGGCACCTCTCAGGATAATG-3'. Following verification by Sanger sequencing, a region of the *UL136K-Rmyc* pGEM-T Easy was PCR amplified from bp 854 of UL135 [UL135(bp854)] to UL138 (bp139). This PCR product was gel purified and recombined with a ΔUL136<GalK> BAC as previously described (21). BAC integrity was tested by enzyme digest fragment analysis and sequencing of the UL133/8 viral genomic region. All BAC genomes were maintained in Escherichia coli SW102, and viral stocks were propagated by transfecting 15 to 20 µg of each BAC genome, along with 2 µg of a plasmid encoding UL82 (pp71), into 5 × 106 MRC-5 fibroblasts and subsequently purified and stored as previously described (21).

1121

Immunoblot analyses

Cells were washed twice in PBS (Gibco) and lifted from cell culture vessels with 0.25X trypsin (Gibco). Cells were washed in cold PBS, pelleted at 1000 xg for 2 minutes followed by storage at

1125 -80 °C before lysis in RIPA buffer. Total protein was quantified using a BCA assay and 50 µg of
1126 each sample was prepared for loading onto 12% precast PAGE gels (GenScript). Wet transfers
1127 were performed onto 0.45 uM pore-sized PVDF membranes (Immobilon). Blocking was achieved
1128 by 5% skimmed milk in 1X TBST, or 5% BSA in 1X TBST. Primary and secondary antibodies
1129 used for western blot analyses are indicated in SI Table 3.

1130

1131 RT-qPCR assays

Cells were washed twice in cold PBS and lysed in 400 µl of RNA/DNA lysis buffer (Zymo
Research). Total cellular RNA was extracted according to the manufacturer's protocol. cDNA was
synthesized from 1 µg of DNAse-treated RNA using the Vilo cDNA synthesis kit (Invitrogen).
qPCR reactions were prepared with *IDOL* TaqMan assay (ID Hs00982312_m1; ThermoFisher
Scientific), Platinum™ Quantitative PCR SuperMix-UDG w/ROX (ThermoFisher Scientific) and 50
ng cDNA. qPCR reactions were run in a QuantStudio 3 real time PCR machine (Applied
Biosystems). Gene expression was normalized to GAPDH (Hs01922876_u1) or H6PD
(Hs00188728_m1) and the ΔΔCT method was used to calculate relative gene expression levels
(110, 111).

1141

1142 Co-immunoprecipitation

Cells were washed twice with PBS and trypsinized. The PBS-washed cell pellet was lysed on ice
for 30 minutes in a lysis buffer made of 100 mM Tris/HCl (pH 7.4), 100 mM NaCl, 50 mM NaFl,
and 0.5% NP-40. The complete protease inhibitor cocktail (Roche) was added to the lysis buffer
according to manufacturer recommendations. Lysates were clarified by centrifugation at 3000 x
g for 5 minutes at 4 °C. A milligram of the lysate was incubated with Pierce anti-myc or anti-HA
magnetic beads for 1 h at 4 °C. The beads were washed 3 times and protein eluted according to
manufacturer's protocol.

1150

1151

1152 References for Supporting Information

1153

1.   F. Goodrum, "The complex biology of human cytomegalovirus latency and reactivation"
in Advances in Virus Research, T. C. M. M. Kielian, M. J. Roossinck, Ed. (Academic Press, 2022), vol.
112.
2.   F. Goodrum, W. Britt, E. S. Mocarski, "Cytomegalovirus" in Fields Virology: DNA Viruses,
P. M. Howley, D. M. Knipe, J. L. Cohen, B. A. Damania, Eds. (Wolters Kluwer, 2022), vol. 2, chap.
12.
3.   G. Hahn, R. Jores, E. S. Mocarski, Cytomegalovirus remains latent in a common precursor
of dendritic and myeloid cells. *Proc Natl Acad Sci U S A* **95**, 3937-3942 (1998).
4.   C.-K. Min *et al.*, The Differentiation of Human Cytomegalovirus Infected-Monocytes Is
Required for Viral Replication. **10** (2020).

5.   M. B. Reeves, P. A. MacAry, P. J. Lehner, J. G. P. Sissons, J. H. Sinclair, Latency, chromatin
remodeling, and reactivation of human cytomegalovirus in the dendritic cells of healthy carriers.
Proceedings of the National Academy of Sciences of the United States of America **102**, 4140 (2005).
6.   M. B. Reeves, J. H. Sinclair, Circulating dendritic cells isolated from healthy seropositive
donors are sites of human cytomegalovirus reactivation in vivo. *J Virol* **87**, 10660-10667 (2013).
7.   J. Taylor-Wiedeman, J. G. P. Sissons, L. K. Borysiewicz, J. H. Sinclair, Monocytes are a major
site of persistence of human cytomegalovirus in peripheral blood mononuclear cells. **72**, 2059-
2064 (1991).
8.   E. Forte *et al.*, Tumor Necrosis Factor Alpha Induces Reactivation of Human
Cytomegalovirus Independently of Myeloid Cell Differentiation following Posttranscriptional
Establishment of Latency. *mBio* **9**, e01560-01518 (2018).
9.   C. Söderberg-Nauclér, K. N. Fish, J. A. Nelson, Reactivation of Latent Human
Cytomegalovirus by Allogeneic Stimulation of Blood Cells from Healthy Donors. *Cell* **91**, 119-126
(1997).
10.   M. B. Reeves, T. Compton, Inhibition of inflammatory interleukin-6 activity via
extracellular signal-regulated kinase-mitogen-activated protein kinase signaling antagonizes
human cytomegalovirus reactivation from dendritic cells. *J Virol* **85**, 12750-12758 (2011).
11.   J. Buehler *et al.*, Opposing Regulation of the EGF Receptor: A Molecular Switch Controlling
Cytomegalovirus Latency and Replication. *PLOS Pathogens* **12**, e1005655 (2016).
12.   M. P. Weekes *et al.*, Latency-Associated Degradation of the MRP1 Drug Transporter
During Latent Human Cytomegalovirus Infection. *Science* **340**, 199 (2013).
13.   V. T. K. Le, M. Trilling, H. Hengel, The Cytomegaloviral Protein pUL138 Acts as Potentiator
of Tumor Necrosis Factor (TNF) Receptor 1 Surface Density To Enhance
ULb'-Encoded Modulation of TNF-α Signaling. *Journal of Virology* **85**,
13260 (2011).
14.   C. Montag *et al.*, The Latency-Associated UL138 Gene Product of Human Cytomegalovirus
Sensitizes Cells to Tumor Necrosis Factor Alpha (TNF-α) Signaling by Upregulating TNF-α Receptor
1 Cell Surface Expression. **85**, 11409-11421 (2011).
15.   E. R. Albright, C. K. Mickelson, R. F. Kalejta, Human Cytomegalovirus UL138 Protein
Inhibits the STING Pathway and Reduces Interferon Beta mRNA Accumulation during Lytic and
Latent Infections. *mBio* **12**, e0226721-e0226721 (2021).
16.   S. H. Lee, E. R. Albright, J.-H. Lee, D. Jacobs, R. F. Kalejta, Cellular defense against latent
colonization foiled by human cytomegalovirus UL138 protein. *Sci Adv* **1**, e1501164-e1501164
(2015).
17.   M. Umashankar *et al.*, Antagonistic Determinants Controlling Replicative and Latent
States of Human Cytomegalovirus Infection. *Journal of Virology* **88**, 5987 (2014).
18.   M. A. Rak *et al.*, Human Cytomegalovirus UL135 Interacts with Host Adaptor Proteins To
Regulate Epidermal Growth Factor Receptor and Reactivation from Latency. **92**, e00919-00918
(2018).
19.   J. Buehler *et al.*, Host signaling and EGR1 transcriptional control of human
cytomegalovirus replication and latency. *PLOS Pathogens* **15**, e1008037 (2019).
20.   K. Caviness *et al.*, Complex Interplay of the UL136 Isoforms
Balances Cytomegalovirus Replication and Latency. *mBio* **7**, e01986-01915 (2016).
21.   K. Caviness, L. Cicchini, M. Rak, M. Umashankar, F. Goodrum, Complex expression of the
UL136 gene of human cytomegalovirus results in multiple protein isoforms with unique roles in
replication. *J Virol* **88**, 14412-14425 (2014).
22.   I. Mikell *et al.*, HCMV miR-US22 down-regulation of EGR-1 regulates CD34+ hematopoietic
progenitor cell proliferation and viral reactivation. *PLoS Pathog* **15**, e1007854 (2019).

23.   I. M. Min *et al.*, The Transcription Factor EGR1 Controls Both the Proliferation and
Localization of Hematopoietic Stem Cells. *Cell Stem Cell* **2**, 380-391 (2008).
24.   D. Collins-McMillen, J. Kamil, N. Moorman, F. Goodrum, Control of Immediate Early Gene
Expression for Human Cytomegalovirus Reactivation. *Frontiers in cellular and infection*
microbiology **10**, 476 (2020).
25.   B. A. Krishna, A. B. Wass, C. M. O'Connor, Activator protein-1 transactivation of the major
immediate early locus is a determinant of cytomegalovirus reactivation from latency. *Proc Natl*
Acad Sci U S A **117**, 20860-20867 (2020).
26.   A. E. Hale *et al.*, FOXO transcription factors activate alternative major immediate early
promoters to induce human cytomegalovirus reactivation. *Proc Natl Acad Sci U S A* **117**, 18764-
18770 (2020).
27.   V. G. Kew, J. Yuan, J. Meier, M. B. Reeves, Mitogen and stress activated kinases act co-
operatively with CREB during the induction of human cytomegalovirus immediate-early gene
expression from latency. *PLoS Pathog* **10**, e1004195 (2014).
28.   L. B. Crawford *et al.*, Human Cytomegalovirus Encodes a Novel FLT3 Receptor Ligand
Necessary for Hematopoietic Cell Differentiation and Viral Reactivation. *MBio* **9** (2018).
29.   M. H. Hancock *et al.*, Human Cytomegalovirus UL7, miR-US5-1, and miR-UL112-3p
Inactivation of FOXO3a Protects CD34(+) Hematopoietic Progenitor Cells from Apoptosis.
mSphere **6** (2021).
30.   E. Scotti *et al.*, Targeted disruption of the idol gene alters cellular regulation of the low-
density lipoprotein receptor by sterols and liver x receptor agonists. *Mol Cell Biol* **31**, 1885-1893
(2011).
31.   R. G. de Bruin *et al.*, Quaking promotes monocyte differentiation into pro-atherogenic
macrophages by controlling pre-mRNA splicing and gene expression. *Nat Commun* **7**, 10846
(2016).
32.   Y. Liu, M. Tingart, S. Lecouturier, J. Li, J. Eschweiler, Identification of co-expression
network correlated with different periods of adipogenic and osteogenic differentiation of BMSCs
by weighted gene co-expression network analysis (WGCNA). *BMC Genomics* **22**, 254 (2021).
33.   E. Scotti *et al.*, IDOL stimulates clathrin-independent endocytosis and multivesicular body-
mediated lysosomal degradation of the low-density lipoprotein receptor. *Mol Cell Biol* **33**, 1503-
1514 (2013).
34.   C. Hong *et al.*, The E3 ubiquitin ligase IDOL induces the degradation of the low density
lipoprotein receptor family members VLDLR and ApoER2. *J Biol Chem* **285**, 19720-19726 (2010).
35.   N. Zelcer, C. Hong, R. Boyadjian, P. Tontonoz, LXR regulates cholesterol uptake through
Idol-dependent ubiquitination of the LDL receptor. *Science* **325**, 100-104 (2009).
36.   M. H. Glickman, A. Ciechanover, The ubiquitin-proteasome proteolytic pathway:
destruction for the sake of construction. *Physiological reviews* **82**, 373-428 (2002).
37.   C. M. Pickart, Back to the future with ubiquitin. *Cell* **116**, 181-190 (2004).
38.   J. Zhang, Q. Liu, Cholesterol metabolism and homeostasis in the brain. *Protein Cell* **6**, 254-
264 (2015).
39.   L. Zhang, K. Reue, L. G. Fong, S. G. Young, P. Tontonoz, Feedback regulation of cholesterol
uptake by the LXR-IDOL-LDLR axis. *Arterioscler Thromb Vasc Biol* **32**, 2541-2546 (2012).
40.   J. Gao *et al.*, The E3 ubiquitin ligase IDOL regulates synaptic ApoER2 levels and is
important for plasticity and learning. *Elife* **6**, e29178 (2017).
41.   J. K. Nelson *et al.*, Deubiquitylase Inhibition Reveals Liver X Receptor-independent
Transcriptional Regulation of the E3 Ubiquitin Ligase IDOL and Lipoprotein Uptake. *J Biol Chem*
291, 4813-4825 (2016).
42.   Anonymous, Cytomegalovirus. *Am J Transplant* **4 Suppl 10**, 51-58 (2004).

43.   Y. Du, G. Zhang, Z. Liu, Human cytomegalovirus infection and coronary heart disease: a systematic review. *Virol J* **15**, 31-31 (2018).

44.   N. Gudleski-O'Regan, T. M. Greco, I. M. Cristea, T. Shenk, Increased expression of LDL receptor-related protein 1 during human cytomegalovirus infection reduces virion cholesterol and infectivity. *Cell Host Microbe* **12**, 86-96 (2012).

45.   Y. Xi, S. Harwood, L. M. Wise, J. G. Purdy, Human Cytomegalovirus pUL37x1 Is Important for Remodeling of Host Lipid Metabolism. *J Virol* **93** (2019).

46.   Y. Xi, L. Lindenmayer, I. Kline, J. von Einem, J. G. Purdy, Human Cytomegalovirus Uses a Host Stress Response To Balance the Elongation of Saturated/Monounsaturated and Polyunsaturated Very-Long-Chain Fatty Acids. *mBio* **12** (2021).

47.   Y. Yu, T. G. Maguire, J. C. Alwine, Human cytomegalovirus infection induces adipocyte-like lipogenesis through activation of sterol regulatory element binding protein 1. *J Virol* **86**, 2942-2949 (2012).

48.   Y. Yu, F. J. Pierciey, Jr., T. G. Maguire, J. C. Alwine, PKR-like endoplasmic reticulum kinase is necessary for lipogenic activation during HCMV infection. *PLoS Pathog* **9**, e1003266 (2013).

49.   D. Collins-McMillen *et al.*, Alternative promoters drive human cytomegalovirus reactivation from latency. *Proc Natl Acad Sci U S A* **116**, 17492-17497 (2019).

50.   D. Collins-McMillen *et al.*, Alternative promoters drive human cytomegalovirus reactivation from latency. *Proceedings of the National Academy of Sciences* **116**, 17492 (2019).

51.   M.-C. Arcangeletti *et al.*, Human cytomegalovirus reactivation from latency: validation of a "switch" model in vitro. *Virol J* **13**, 179-179 (2016).

52.   L.-F. Yee, P. L. Lin, M. F. Stinski, Ectopic expression of HCMV IE72 and IE86 proteins is sufficient to induce early gene expression but not production of infectious virus in undifferentiated promonocytic THP-1 cells. *Virology* **363**, 174-188 (2007).

53.   B. G. Weinshenker, S. Wilton, G. P. Rice, Phorbol ester-induced differentiation permits productive human cytomegalovirus infection in a monocytic cell line. *J Immunol* **140**, 1625-1631 (1988).

54.   N. Wagenknecht *et al.*, Contribution of the Major ND10 Proteins PML, hDaxx and Sp100 to the Regulation of Human Cytomegalovirus Latency and Lytic Replication in the Monocytic Cell Line THP-1. *Viruses* **7**, 2884-2907 (2015).

55.   C. H. Lee *et al.*, Factors affecting human cytomegalovirus gene expression in human monocyte cell lines. *Mol Cells* **9**, 37-44 (1999).

56.   M. Peppenelli, J. Buehler, F. Goodrum, Human Hematopoietic Long-Term Culture (hLTC) for Human Cytomegalovirus Latency and Reactivation. *Methods Mol Biol* **2244**, 83-101 (2021).

57.   R. N. Ramirez *et al.*, Dynamic Gene Regulatory Networks of Human Myeloid Differentiation. *Cell systems* **4**, 416-429.e413 (2017).

58.   F. O. Martinez, S. Gordon, M. Locati, A. Mantovani, Transcriptional Profiling of the Human Monocyte-to-Macrophage Differentiation and Polarization: New Molecules and Patterns of Gene Expression. *The Journal of Immunology* **177**, 7303 (2006).

59.   B. Fagerberg, J. Gnarpe, H. Gnarpe, S. Agewall, J. Wikstrand, Chlamydia pneumoniae but not cytomegalovirus antibodies are associated with future risk of stroke and cardiovascular disease: a prospective study in middle-aged to elderly men with treated hypertension. *Stroke* **30**, 299-305 (1999).

60.   M. C. Pizzorno, G. S. Hayward, The IE2 gene products of human cytomegalovirus specifically down-regulate expression from the major immediate-early promoter through a target located near the cap site. *J Virol* **64**, 6154-6165 (1990).

61.   J. M. Cherrington, E. L. Khoury, E. S. Mocarski, Human cytomegalovirus ie2 negatively
regulates alpha gene expression via a short target sequence near the transcription start site. *J*
Virol **65**, 887-896 (1991).
62.   B. Liu, T. W. Hermiston, M. F. Stinski, A cis-acting element in the major immediate early
(IE) promoter of human cytomegalovirus is required for negative regulation by IE2. *J Virol* **65**, 897-
903 (1991).
63.   M. P. Macias, M. F. Stinski, An in vitro system for human cytomegalovirus immediate early
2 protein (IE2)-mediated site-dependent repression of transcription and direct binding of IE2 to
the major immediate early promoter. *Proc Natl Acad Sci U S A* **90**, 707-711 (1993).
64.   K. Caviness *et al.*, Complex Interplay of the UL136 Isoforms Balances Cytomegalovirus
Replication and Latency. *mBio* **7**, e01986 (2016).
65.   S. D. Lee, P. Tontonoz, Liver X receptors at the intersection of lipid metabolism and
atherogenesis. *Atherosclerosis* **242**, 29-36 (2015).
66.   B. A. Janowski *et al.*, Structural requirements of ligands for the oxysterol liver X receptors
LXRalpha and LXRbeta. *Proc Natl Acad Sci U S A* **96**, 266-271 (1999).
67.   B. A. Janowski, P. J. Willy, T. R. Devi, J. R. Falck, D. J. Mangelsdorf, An oxysterol signalling
pathway mediated by the nuclear receptor LXRα. *Nature* **383**, 728-731 (1996).
68.   D. J. Peet, B. A. Janowski, D. J. Mangelsdorf, The LXRs: a new class of oxysterol receptors.
Current Opinion in Genetics & Development **8**, 571-575 (1998).
69.   L. Fagerberg *et al.*, Analysis of the human tissue-specific expression by genome-wide
integration of transcriptomics and antibody-based proteomics. *Molecular & cellular proteomics :*
MCP **13**, 397-406 (2014).
70.   I. Fernandez-Ruiz, P. Puchalska, C. A. Narasimhulu, B. Sengupta, S. Parthasarathy,
Differential lipid metabolism in monocytes and macrophages: influence of cholesterol loading[S].
Journal of Lipid Research **57**, 574-586 (2016).
71.   T. R. Cimato, B. A. Palka, J. K. Lang, R. F. Young, LDL Cholesterol Modulates Human CD34+
HSPCs through Effects on Proliferation and the IL-17 G-CSF Axis. *PLOS ONE* **8**, e73861 (2013).
72.   P. K. Morgan, L. Fang, G. I. Lancaster, A. J. Murphy, Hematopoiesis is regulated by
cholesterol efflux pathways and lipid rafts: connections with cardiovascular diseases: Thematic
Review Series: Biology of Lipid Rafts. *Journal of Lipid Research* **61**, 667-675 (2020).
73.   O. Soehnlein, F. K. Swirski, Hypercholesterolemia links hematopoiesis with
atherosclerosis. *Trends in endocrinology and metabolism: TEM* **24**, 129-136 (2013).
74.   J. M. Radley, S. Ellis, M. Palatsides, B. Williams, I. Bertoncello, Ultrastructure of primitive
hematopoietic stem cells isolated using probes of functional status. *Experimental hematology* **27**,
365-369 (1999).
75.   S. Shariatmadar *et al.*, Electronic volume of CD34 positive cells from peripheral blood
apheresis samples. *Cytometry. Part B, Clinical cytometry* **74**, 182-188 (2008).
76.   M. P. Vacanti, A. Roy, J. Cortiella, L. Bonassar, C. A. Vacanti, Identification and initial
characterization of spore-like cells in adult mammals. *Journal of cellular biochemistry* **80**, 455-460
(2001).
77.   A. C. Calkin *et al.*, FERM-dependent E3 ligase recognition is a conserved mechanism for
targeted degradation of lipoprotein receptors. **108**, 20107-20112 (2011).
78.   B. Liu *et al.*, Inhibition of Human Cytomegalovirus Particle Maturation by Activation of
Liver X Receptor. *Front Microbiol* **13**, 846386-846386 (2022).
79.   Q. Wang *et al.*, Identification of interferon-γ as a new molecular target of liver X receptor.
Biochemical Journal **459**, 345-354 (2014).
80.   B. Sainz, H. L. LaMarca, R. F. Garry, C. A. Morris, Synergistic inhibition of human
cytomegalovirus replication by interferon-alpha/beta and interferon-gamma. *Virol J* **2**, 14 (2005).

81. K. A. Kropp *et al.*, Reversible inhibition of murine cytomegalovirus replication by gamma
interferon (IFN-γ) in primary macrophages involves a primed type I IFN-signaling subnetwork for
full establishment of an immediate-early antiviral state. *Journal of virology* **85**, 10286-10299
(2011).
82. Y. Wang *et al.*, Pseudorabies Virus Inhibits Expression of Liver X Receptors to Assist Viral
Infection. *Viruses* **14**, 514 (2022).
83. P. T. Lange, C. Schorl, D. Sahoo, V. L. Tarakanova, Liver X Receptors Suppress Activity of
Cholesterol and Fatty Acid Synthesis Pathways To Oppose Gammaherpesvirus Replication. *mBio*
9, e01115-01118 (2018).
84. P. T. Lange, C. N. Jondle, E. J. Darrah, K. E. Johnson, V. L. Tarakanova, LXR Alpha Restricts
Gammaherpesvirus Reactivation from Latently Infected Peritoneal Cells. *Journal of virology* **93**,
e02071-02018 (2019).
85. J. Zeng *et al.*, Liver X receptors agonists impede hepatitis C virus infection in an Idol-
dependent manner. *Antiviral Res* **95**, 245-256 (2012).
86. S. Bocchetta *et al.*, Up-regulation of the ATP-binding cassette transporter A1 inhibits
hepatitis C virus infection. *PLoS One* **9**, e92140 (2014).
87. H. Jiang *et al.*, Retinoic acid and liver X receptor agonist synergistically inhibit HIV infection
in CD4+ T cells by up-regulating ABCA1-mediated cholesterol efflux. *Lipids Health Dis* **11**, 69
(2012).
88. L. Dubrovsky *et al.*, Liver X receptor agonist inhibits HIV-1 replication and prevents HIV-
induced reduction of plasma HDL in humanized mouse model of HIV infection. *Biochem Biophys
Res Commun* **419**, 95-98 (2012).
89. T. M. Hanley, G. A. Viglianti, Nuclear receptor signaling inhibits HIV-1 replication in
macrophages through multiple trans-repression mechanisms. *J Virol* **85**, 10834-10850 (2011).
90. X. X. Sheng *et al.*, The LXR ligand GW3965 inhibits Newcastle disease virus infection by
affecting cholesterol homeostasis. *Arch Virol* **161**, 2491-2501 (2016).
91. J. Zeng *et al.*, Liver X receptors agonists impede hepatitis C virus infection in an Idol-
dependent manner. *Antiviral Research* **95**, 245-256 (2012).
92. G. H. Syed *et al.*, Hepatitis C virus stimulates low-density lipoprotein receptor expression
to facilitate viral propagation. *Journal of virology* **88**, 2519-2529 (2014).
93. A. Albecka *et al.*, Role of low-density lipoprotein receptor in the hepatitis C virus life cycle.
Hepatology (Baltimore, Md.) **55**, 998-1007 (2012).
94. S. Molina *et al.*, The low-density lipoprotein receptor plays a role in the infection of
primary human hepatocytes by hepatitis C virus. *Journal of hepatology* **46**, 411-419 (2007).
95. B. Sun *et al.*, Regulation of host and virus genes by neuronal miR-138 favours herpes
simplex virus 1 latency. *Nature microbiology* **6**, 682-696 (2021).
96. D. Pan *et al.*, A neuron-specific host microRNA targets herpes simplex virus-1 ICP0
expression and promotes latency. *Cell Host Microbe* **15**, 446-456 (2014).
97. X. Liu *et al.*, Herpesvirus-mediated stabilization of ICP0 expression neutralizes restriction
by TRIM23. *Proc Natl Acad Sci U S A* **118** (2021).
98. T. H. Chang *et al.*, Regulation of the Abundance of Kaposi's Sarcoma-Associated
Herpesvirus ORF50 Protein by Oncoprotein MDM2. *PLoS Pathog* **12**, e1005918 (2016).
99. J. C. Chmura *et al.*, The Itch ubiquitin ligase is required for KSHV RTA induced vFLIP
degradation. *Virology* **501**, 119-126 (2017).
100. E. S. Ehrlich, J. C. Chmura, J. C. Smith, N. N. Kalu, G. S. Hayward, KSHV RTA abolishes
NFkappaB responsive gene expression during lytic reactivation by targeting vFLIP for degradation
via the proteasome. *PLoS One* **9**, e91359 (2014).

101.   S. Gandon, Why Be Temperate: Lessons from Bacteriophage λ. *Trends in Microbiology* **24**,
356-365 (2016).
102.   S. Atsumi, J. W. Little, Role of the lytic repressor in prophage induction of phage lambda
as analyzed by a module-replacement approach. *Proceedings of the National Academy of Sciences*
of the United States of America **103**, 4558-4563 (2006).
103.   R. A. Schubert, I. B. Dodd, J. B. Egan, K. E. Shearwin, Cro's role in the CI Cro bistable switch
is critical for {lambda}'s transition from lysogeny to lytic development. *Genes Dev* **21**, 2461-2472
(2007).
104.   M. G. Cortes, J. T. Trinh, L. Zeng, G. Balazsi, Late-Arriving Signals Contribute Less to Cell-
Fate Decisions. *Biophys J* **113**, 2110-2120 (2017).
105.   O. Kobiler, A. Rokney, A. B. Oppenheim, Phage lambda CIII: a protease inhibitor regulating
the lysis-lysogeny decision. *PLoS One* **2**, e363 (2007).
106.   J. W. Little, D. P. Shepley, D. W. J. T. E. j. Wert, Robustness of a gene regulatory circuit. **18**,
4299-4307 (1999).
107.   J. D. Hernandez-Doria, V. Sperandio, Bacteriophage Transcription Factor Cro Regulates
Virulence Gene Expression in Enterohemorrhagic Escherichia coli. *Cell Host Microbe* **23**, 607-
617.e606 (2018).
108.   M. Umashankar *et al.*, A Novel Human Cytomegalovirus Locus Modulates Cell Type-
Specific Outcomes of Infection. *PLOS Pathogens* **7**, e1002444 (2011).
109.   N. E. Sanjana, O. Shalem, F. Zhang, Improved vectors and genome-wide libraries for
CRISPR screening. *Nat Methods* **11**, 783-784 (2014).
110.   X. Rao, X. Huang, Z. Zhou, X. Lin, An improvement of the 2^(-delta delta CT) method for
quantitative real-time polymerase chain reaction data analysis. *Biostat Bioinforma Biomath* **3**, 71-
85 (2013).
111.   J. Winer, C. K. Jung, I. Shackel, P. M. Williams, Development and validation of real-time
quantitative reverse transcriptase-polymerase chain reaction for monitoring gene expression in
cardiac myocytes in vitro. *Anal Biochem* **270**, 41-49 (1999).

1428

1429

1430

1431

1432

1433