Identification of basic helix-loop-helix transcription factors that activate betulinic

acid biosynthesis by RNA-sequencing of hydroponically cultured *Lotus japonicus*

Hayato Suzuki[1,2], Hirokazu Takahashi[3], Ery Odette Fukushima[1,4], Mikio Nakazono[3],

Toshiya Muranaka[1,5], and Hikaru Seki[1,5*]

[1] Department of Biotechnology, Graduate School of Engineering, Osaka University, 2-1

Yamadaoka, Suita, Osaka 565-0871, Japan

[2] Bioproduction Research Institute, National Institute of Advanced Industrial Science

and Technology (AIST), Higashi 1-1-1, Tsukuba, Ibaraki 305-8562, Japan

[3] Graduate School of Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa,

Nagoya 464-8601, Japan

[4] Plant Translational Research Group, Universidad Regional Amazónica IKIAM,

Muyuna kilómetro 7 vía a Alto Tena 150101, Ecuador

[5] Industrial Biotechnology Initiative Division, Institute for Open and Transdisciplinary

Research Initiatives, Osaka University, 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan

* Corresponding author (hseki@bio.eng.osaka-u.ac.jp; Tel +81-6-6879-7423)

ORCIDs

Hayato Suzuki, https://orcid.org/0000-0001-6120-9471

Hirokazu Takahashi, https://orcid.org/0000-0001-6182-3236

Ery Odette Fukushima, https://orcid.org/0000-0003-4099-4292

Mikio Nakazono, https://orcid.org/0000-0001-7119-2052

Toshiya Muranaka, https://orcid.org/0000-0003-1058-2473

Hikaru Seki, https://orcid.org/0000-0002-6426-0368

Word counts: Summary (198), Main body in total (6293), Introduction (592), Materials

and Methods (2392), Results (1883), Discussion (1426)

Number of: Figures (7; Figures other than Fig. 3 will be coloured), Tables (2)

Type and number of supporting information: Figures (9); Tables (7)

Summary

● Although triterpenes are ubiquitous in plant kingdom, their biosynthetic regulatory

mechanisms are limitedly understood. Here, we found that hydroponic culture dramatically

activated betulinic acid (BA) biosynthesis in the model Fabaceae *Lotus japonicus*, and

investigated its transcriptional regulation.

● Fabaceae plants develop secondary aerenchyma (SA) on the surface of hypocotyls and

roots during flooding for root air diffusion. Hydroponic culture induced SA in *L. japonicus*

and simultaneously increased the accumulation of BA and the transcript levels of its

biosynthetic genes. RNA-sequencing of soil-grown and hydroponically cultured plant

tissues, including SA collected by laser microdissection, revealed that several transcription

factor genes were co-upregulated with BA biosynthetic genes.

● Overexpression of *LjbHLH32* and *LjbHLH50* in *L. japonicus* transgenic hairy roots

upregulated the expression of BA biosynthesis genes, resulting in enhanced BA

accumulation. However, transient luciferase reporter assays in *Arabidopsis* mesophyll cell

protoplasts showed that LjbHLH32 transactivated promoters of biosynthetic genes in the

soyasaponin pathway but not the BA pathway, like its homolog GubHLH3, a soyasaponin

biosynthesis regulator in *Glycyrrhiza uralensis*. This suggested the evolutionary origin and

complex regulatory mechanisms of BA biosynthesis in Fabaceae.

● This study sheds light on the unrevealed biosynthetic regulatory mechanisms of triterpenes

in Fabaceae plants.

Key words: betulinic acid, bHLH, hydroponics, laser microdissection (LMD), *Lotus*

japonicus, secondary aerenchyma, transcription factor, triterpene

Introduction

Triterpenes are plant specialized (secondary) metabolites with diverse structures and biological properties (Thimmappa *et al.*, 2014). The common precursor, 2,3-oxidosqualene, is composed of six isoprene units supplied by the mevalonate (MVA) pathway. Multiple oxidosqualene cyclases (OSCs), cytochrome P450 monooxygenases (P450s), and UDP-dependent glycosyltransferases (UGTs) catalyze conversion of the substrates to structurally diversified non-glycosylated triterpenes and triterpene glycosides (saponins; Seki *et al.*, 2015).

Fabaceae plants produce a large amount of triterpenes with marked chemodiversity. Soyasaponins are specific to Fabaceae plants such as *Glycine max*, *Medicago* spp., *Glycyrrhiza* spp., and *Lotus japonicus* (Fenwick, 1992; Hayashi *et al.*, 1998; Tsuno *et al.*, 2018). *Medicago truncatula* produces oleanolic acid (OA)-derived hemolytic saponins (Huhman *et al.*, 2005). Thickened roots of *Glycyrrhiza uralensis* accumulate glycyrrhizin, but more OA and betulinic acid (BA) were produced in tissue-cultured stolons (Kojoma *et al.*, 2010; Tamura *et al.*, 2017). *L. japonicus* accumulates C-28-oxidized triterpenes [ursolic acid (UA), OA, and BA] in a tissue-dependent manner (Suzuki *et al.*, 2019). Biosynthesis pathways of these Fabaceae triterpenes have been almost understood by the recent identification of CSyGT (cellulose-synthase-derived glycosyltransferase) enzymes catalyzing the glucuronosylation of triterpene aglycones (Jozwiak *et al.*, 2020; Chung *et al.*, 2020). However, their physiological functions and biosynthetic regulatory mechanisms are unclear.

Plant genomes possess a large number of basic helix-loop-helix (bHLH)-type transcription factors (TFs), which are classified into approximately 25 clades based on

sequence similarity in the bHLH domain and other conserved domains (Heim *et al.*,

2003). Fabaceae plants possess more clade IVa bHLH genes than related plant families,

which are subdivided into three subclades (IVa1-3) based on a phylogenetic tree of the

full-length proteins (Suzuki *et al.*, 2021). Some members of subclade IVa1 are

regulators of Fabaceae triterpene saponin biosynthesis. TSAR1 and TSAR2 are positive

regulators of soyasaponins and hemolytic saponins, respectively, in *M. truncatula*

(Mertens *et al.*, 2016). TSAR3, their paralog, is a regulator of hemolytic saponin

biosynthesis during seed development (Ribeiro *et al.*, 2020). GubHLH3, a TSAR

homolog, upregulated soyasaponin biosynthesis in *G. uralensis* (Tamura *et al.*, 2018).

TFs regulating BA biosynthesis have yet been identified in Fabaceae plants. On the

other hand, BpbHLH9 (clade Ia) was identified as an OA and BA biosynthesis regulator

in the Betulaceae plant *Betula platyphylla* (Yin *et al.*, 2017). The clade Ib bHLH TFs Bl

and Bt are regulators of cucurbitane-type triterpenes in leaves and fruits, respectively, of

cucumber (Cucurbitaceae; Shang *et al.*, 2014). Thus, identification of triterpene-related

bHLH TFs in *L. japonicus* provides insight into the evolutionary history of triterpene

biosynthesis regulators in the Fabaceae and other plant families.

We performed triterpene profiling to characterize triterpene production in *L.*

japonicus. A large amount of BA accumulated in secondary aerenchyma (SA) tissue in

hypocotyl and roots when cultured hydroponically. SA is a white, spongy tissue

developed from phellogen [secondary meristem (SM)], whose formation is an adaptive

trait of Fabaceae plants to survive flooding by promoting oxygen diffusion in roots

(Mochizuki *et al.*, 2000; Yamauchi *et al.*, 2013). Because the expression of BA

biosynthetic genes (*OSC3* and *CYP716A51*) was upregulated in whole roots of

hydroponically cultured plants and SA, we performed comparative RNA sequencing

(RNA-seq) analysis to identify candidate TFs that positively regulate BA biosynthesis.

From comparisons between soil-cultured and hydroponically cultured whole roots and

between SA and SM collected by laser microdissection (LMD), we identified 23 TFs

co-upregulated in hydroponic roots and SA. Among them, we focused on LjbHLH32

(subclade IVa2) and LjbHLH50 (clade Ia) based on their co-expression with BA

biosynthetic genes and phylogenetic analysis. The functions of these two bHLH-type

TFs in triterpene biosynthesis regulation were examined and are discussed.

Materials and Methods

Plant materials

Seeds of *L. japonicus* Gifu B-129 accession and *M. truncatula* cv. R108 and Jemalong

A17 were sterilized using a five-fold-diluted sodium hypochlorite solution containing

0.02% Tween-20 for 15 min with mild agitation. After washing three times in sterilized

water, they were soaked in water with mild agitation overnight. The seeds were

germinated on 0.8% agar with incubation for 4 days (*L. japonicus*) or 3 days (*M.*

truncatula) in the dark and for 2 days on a 16 h light/8 h dark cycle at 25°C. The

seedlings were transplanted to soil pots or hydroponic solutions and cultured in a plant

growth room maintained at 23°C until harvest. We used basal nutrient solution (BNS)

for all hydroponic solutions (Conn *et al.*, 2013). *L. japonicus* seedlings were cultured in

5 mL sampling tubes for 2 weeks and transferred to 50 mL sampling tubes with BNS. *M.*

truncatula seedlings were cultured in 50 mL tubes with BNS from transplant to harvest.

For induction of adventitious roots, the roots of 1-week-old seedlings were isolated and

cultured in 100 mL of half-strength Gamborg's B5 liquid medium supplemented with

3% sucrose and 0.2 mg/L of indole-3-butyric acid. The culture was maintained at 23°C

with shaking at 90 rpm and subcultured every 3 to 4 weeks. The *L. japonicus* Lj

suspension cell line (rpc00032) was maintained at 27°C with shaking at 120 rpm in

Murashige and Skoog (MS) medium (pH 5.7) supplemented with 1 mg/L of 2,4-D and

3% sucrose, and subcultured every 7 days. For metabolite analysis of cultured cells,

ethanol or 100 μM methyl jasmonate (MeJA) was added and the cells were cultured for

4 additional days.

Triterpene extraction and gas chromatography-mass spectrometry (GC-MS)

analysis

Triterpenes were extracted from 10 mg of dried plant powder and analyzed by GC-MS

following our previous report (Suzuki *et al.*, 2019).

RNA extraction from whole roots

Whole roots (including the hypocotyl) of 2.5-month-old *L. japonicus* cultured on soil or

in hydroponics were harvested and immediately frozen in liquid nitrogen. Total RNA

extraction, recombinant DNase I treatment, and column purification were performed as

described previously (Suzuki *et al.*, 2019).

RNA extraction from SA and SM

Sample fixation, paraffin-embedding, microtome sectioning, collection of SA and SM

cell layers by LMD, and RNA extraction were performed following a previous report

(Takahashi *et al.*, 2010) with some modifications.

Hypocotyls of *L. japonicus* seedlings cultured hydroponically for 7 weeks were

harvested. Segments of hypocotyls were obtained at 5 to 10 mm from the cotyledons.

The 5-mm-long hypocotyls were fixed in methanol by incubation for 30 min under a

vacuum of 0.09 MPa. This procedure was repeated three times with new methanol.

We employed a microwave-assisted method for paraffin-embedding using an

Energy Beam Sciences H2850 Microwave Tissue Processor (Agawan, MA). A vial

containing fixed tissue and pre-chilled methanol was placed on ice and microwaved

three times at 250 W at 37°C for 15 min. The methanol was replaced each time. The

temperature in the vial was < 20°C during the first and second microwave treatments;

however, 37°C was allowed in the third treatment. The solvent was gradually replaced

with   1-butanol,   using   methanol:1-butanol   (3:1),   methanol:1-butanol   (1:1),

methanol:1-butanol (1:3), and 1-butanol, with microwaving at each step at 300 W and

37°C for 1.5 min. 1-Butanol was gradually replaced with melted paraffin wax

(Surgipath Paraplast X-tra Tissue Infiltration/Embedding Media; Leica Biosystems,

Wetzlar, Germany) using 1-butanol:paraffin (3:1), 1-butanol:paraffin (1:1), and

1-butanol:paraffin (1:3) with microwaving at each step at 300 W and 58°C for 10 min.

Two treatments at 250 W at 58°C for 10 min and five treatments at 250 W at 58°C for

30 min were performed with new paraffin solution. The specimens were embedded in

paraffin wax and cooled to room temperature. The paraffin blocks were stored at 4°C.

RNAsecure reagent (Thermo Fisher Scientific, Waltham, MA) was diluted

25-fold with UltraPure Distilled Water (Thermo Fisher Scientific). The diluted

RNAsecure solution was placed on an Arcturus PEN Membrane Glass Slide (Thermo

Fisher Scientific) heated on a hot plate (57°C). Paraffin sections (12 μm thickness) were

floated on the surface of the solution and incubated for several tens of seconds. After

removing the solution, the sections were dried at 42°C for 1 h. The slides were

immersed in Histoclear II (National Diagnostics, Atlanta, GA) for 10 min twice to

remove the paraffin, and air-dried. LMD was performed using a PALM MicroBeam IV

Laser-capture Microdissection System (Zeiss, Jena, Germany) with a MicroBeam Laser

(cut energy: 55–58, cut focus: 80, LPC energy: 30, LPC focus: 2). AdhesiveCap 500

clear (Zeiss) was used to collect cell layers cut by the laser. RNA was purified using an

Arcturus PicoPure RNA Isolation Kit (Thermo Fisher Scientific).

Quantitative reverse transcription-PCR (qRT-PCR) analysis

First-strand cDNA was synthesized from purified total RNA using PrimeScript RT

Master Mix (Perfect Real Time; Takara Bio, Shiga, Japan) following the manufacturer's

instructions. The reaction time was 15 and 60 min, for total RNA of whole roots and

that of LMD samples, respectively. We performed qRT-PCR analysis following a

previous report (Suzuki *et al.*, 2019) using primer nos. 1–20 in Table S1.

RNA-seq

Transcriptome profiling of soil-grown and hydroponic roots was performed with three

biological replicates (Table S2). Complementary DNA library construction was

performed using a TruSeq Stranded mRNA Library Prep Kit (Illumina, San Diego, CA)

according to the manufacturer's instructions. The resulting cDNA libraries were

sequenced using HiSeq 2500 (Illumina) with 100 bp single-end (SE) reads. Adapter

sequences and low-quality reads were removed from raw RNA-seq reads by

Trimmomatic v. 0.38 (Bolger *et al.*, 2014). Quantification of transcripts was conducted

with Salmon v. 0.14.2 (Patro *et al.*, 2017). Reference *L. japonicus* coding sequences

(predicted from genome assembly build v. 1.2 of Gifu B-129 accession; Kamal *et al.*,

2020) were downloaded from *Lotus* Base (https://lotus.au.dk/). A mapping-based index

was built from the reference using an auxiliary k-mer hash over k-mers of length 31 as

recommended. Summarized gene counts were obtained from the transcript counts using

the R package tximport v. 1.16.1 (Soneson *et al.*, 2015). Differentially expressed genes

(DEGs) were identified using the graphical user interface for TCC (TCC-GUI) (Su *et al.*,

2019). Low count genes (0 mapped reads) were filtered out from TCC computation and

default settings were kept for the other parameters.

Transcriptome profiling of the LMD samples was performed with four

biological replicates. RNA-seq libraries were constructed and sequenced using a

Novaseq 6000 (Illumina) with 150-bp paired-end (PE) reads by Takara Bio. Because

two of four replicates of SA samples had smaller total read numbers, they were

combined into one FastQ file and analyzed with three biological replicates (Table S2).

Data processing was conducted as described above after removing unpaired reads.

Gene cloning

The primers used are listed in Table S1. PCR was performed using Prime STAR MAX

or Prime STAR GXL DNA Polymerase (Takara Bio). The plasmid pENTR1A (Thermo

Fisher Scientific) was linearized by PCR (primer nos. 21 and 22). The full coding

sequences of *LjbHLH32* and *LjbHLH50* were amplified from cDNA of hydroponically

cultured roots using primer nos. 24–27 and were integrated into pENTR1A using

NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs, Ipswich, MA) to

obtain gateway entry clones (pENTR1A-*LjbHLH32* and pENTR1A-*LjbHLH50*). *GFP*

was amplified from pUB-GWS-GFP (GenBank accession no. AB303066) using primer

nos. 28 and 29, and cloned into pENTR1A to obtain pENTR1A-*GFP*.

Construction of the gateway destination vector pSD11

The binary vector pUB-GW-GFP is used for overexpression in *L. japonicus* hairy roots (Maekawa *et al.*, 2008). After constructing pUB-GW-GFP, the *Arabidopsis thaliana* heat shock protein (HSP) 18.2 terminator, which is suitable for gene overexpression in plants, was identified (Nagaya *et al.*, 2010; Matsui *et al.*, 2014). Here, we constructed the binary vector pSD11 by replacing the terminators of pUB-GW-GFP with the HSP terminator (Fig. S1).

The primers used are listed in Table S1. The 875 bp native *AtHSP18.2* terminator (HSPter) was amplified from *A. thaliana* Col-0 genomic DNA using primer nos. 30 and 31, and cloned into pJET1.2 using a CloneJET PCR Cloning Kit (Thermo Fisher Scientific). The vector backbone was amplified from pUB-GWS-GFP using primer nos. 32 and 33. The HSP terminator was amplified from pJET1.2-HSPter using primer nos. 34 and 35. These two fragments were assembled by NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs) to construct pSD8.

The vector backbone, including the gateway cassette and GFP, and fragments, including the *LjUBQ1* promoter, were amplified from pUB-GWS-GFP using primer sets 33/36 and 38/39, respectively. These two fragments were assembled with the HSP terminator fragment (primer nos. 34 and 37) to produce pSD9.

The gateway destination vector pSD11 was constructed from PCR-amplified fragments of pSD8 (primer nos. 40 and 41) and pSD9 (primer nos. 42 and 43) by the golden gate cloning method using BsaI-HF v. 2 and T4 DNA ligase (both New England Biolabs).

Generation and analysis of TF-overexpressing hairy roots

The expression vectors (pSD11-*GFP*, *LjbHLH32*, and *LjbHLH50*) were generated by

LR reaction of pSD11 and entry clones (pENTR1A-*GFP*, *LjbHLH32*, and *LjbHLH50*)

using Gateway LR Clonase II Enzyme Mix (Thermo Fisher Scientific). These three

vectors were introduced into *Agrobacterium rhizogenes* ATCC15834 by electroporation.

Agrobacterium rhizogenes-mediated hairy root transformation was performed

following our previous study (Suzuki *et al.*, 2019), with some modifications. The

incubation temperature was maintained at 25°C in the germination, co-culture, and hairy

root induction steps. Isolated hairy roots were cultured at 23°C. For liquid culture of

hairy roots, the rotary shaker agitation speed was set to 90 rpm. The

germination/co-cultivation medium contained 1/2 B5 basal salts. Hairy root elongation

(HRE) medium is composed of 1/2 B5 basal salts, 1/4 Gamborg's B5 vitamins, and 300

mg/L of cefotaxime sodium. All media were adjusted to pH 5.5 before autoclaving and

freshly prepared. Seeds were germinated as described above. After infection and

co-cultivation (Suzuki *et al.*, 2019), seedlings were placed on HRE agar supplemented

with 1% sucrose and a filter paper strip was placed on the hypocotyls to prevent the

hairy roots from drying. Hairy roots were elongated under a 16 h light/8 h dark cycle for

3 weeks, followed by shoot removal. Isolated hypocotyls possessing hairy roots were

incubated on new HRE agar containing 3% sucrose without light for 10 days. Single

hairy roots were isolated and cultured in 5 mL of liquid HRE medium supplemented

with 3% sucrose using six-well cell culture plates. They were subcultured twice every

16 days. Hairy root lines emitting GFP fluorescence were selected and each was

subcultured into six wells. After 26 days of culture, hairy roots were harvested for

GC-MS, qRT-PCR, and RNA-seq analysis (150 bp PE reads, Novaseq 6000 platform,

by Takara Bio; Table S2) as described above.

Cloning of promoter sequences

Putative promoter sequences, including the 5'-untranslated region (2–2.5 kb upstream from the start codon), of triterpene biosynthetic genes were retrieved using the sequence retrieval tool in *Lotus* Base (https://lotus.au.dk/; Table S3). The primers used to amplify the promoter sequences are listed in Table S1 (primer nos. 44–67). The plasmid pENTR1A (Thermo Fisher Scientific) was linearized by PCR (primer nos. 21 and 23). The promoter sequences of *OSC1*, *OSC3*, *OSC9*, *OSC5*, *CYP716A51*, *LjCYP93E1*, *LjCYP72A61*, *LjCSyGT*, putative 2,3-dihydro-3,5-dihydroxy-6-methyl-4*H*-pyran-4-one (DDMP) transferase (*LjDDMPT*), HMG-CoA reductase 1 (*LjHMGR1*), and squalene epoxidase 1 (*LjSQE1*) were integrated into pENTR1A by NEBuilder HiFi DNA assembly cloning.

Vector construction for transient effector-reporter analysis

The *Sfi*I-digested PCR fragment, including the gateway cassette (primer nos. 68 and 69 in Table S1) and *Sfi*I-digested pGL4.1HSP [encoding firefly luciferase (fLUC); Nakata *et al.*, 2013], were ligated using T4 DNA ligase (New England Biolabs) to generate the pGL4.1HSP-GW vector. Promoter sequences were integrated into pGL4.1HSP-GW by the LR reaction to generate 11 reporter constructs (pGL4.1HSP-GW-*OSC1pro*, *OSC3pro*, *OSC9pro*, *OSC5pro*, *CYP716A51pro*, *LjCYP93E1pro*, *LjCYP72A61pro*, *LjCSyGTpro*, *LjDDMPTpro*, *LjHMGR1pro*, and *LjSQE1pro*). pENTR-D-*GuCYP93E3pro* (Tamura *et al.*, 2018) was used to generate pGL4.1HSP-GW- *GuCYP93E3pro*.

TF genes were transferred from entry clones [pENTR-D-*GubHLH3* (Tamura *et al.*, 2018), pENTR1A-*LjbHLH32*, and pENTR1A-*LjbHLH50*] to pDEST35SHSP

(Oshima *et al.*, 2013) to produce three effector constructs (pDEST35SHSP-*GubHLH3*,

LjbHLH32, and *LjbHLH50*). The plasmid pDEST35SHSP-*vesicle-associated membrane*

protein 722 (*Vamp722*; Yoshida *et al.*, 2013) was used as a negative control.

We employed phRLHSP [encoding a modified *Renilla* luciferase (RLUC);

Yoshida *et al.*, 2013] as an internal reference to normalize the results of our dual

luciferase reporter assays.

Transient effector-reporter analysis

A. thaliana Col-0 was grown under a 16 h light/8 h dark cycle at 22°C for 3–4 weeks.

Rosette leaves were used for protoplast isolation after peeling by the Tape-*Arabidopsis*

Sandwich method (Wu *et al.*, 2009). Protoplasts were isolated as described (Yoshida *et*

al., 2013) with minor modifications. Peeled leaves were incubated for 1 h in enzyme

solution [400 mM mannitol, 20 mM KCl, 10 mM $CaCl_2$, 10 g/L of cellulase onozuka

R10 (Yakult Pharmaceutical Industry Co., Ltd., Tokyo, Japan), 2.5 g/L of macerozyme

R10 (Yakult Pharmaceutical Industry Co., Ltd.), 10 mM 2-mercaptoethanol, and 20 mM

2-morpholinoethanesulfonic acid (MES), pH 5.7] in a shaker (22°C, 60 rpm). The

protoplasts were filtered through a 70 μm cell strainer and collected by centrifugation at

$100 \times g$ for 10 min. The supernatant was removed by decantation and the remaining

pellet was washed thrice with W5 buffer [150 mM NaCl, 125 mM $CaCl_2$, 5 mM KCl,

and 2 mM MES, pH 5.7] followed by centrifugation at $100 \times g$ for 5 min. The pellet

was washed with MMg solution [400 mM mannitol, 15 mM $MgCl_2$, and 4 mM MES,

pH 5.7] and resuspended in 5 mL of MMg solution (when using 24 leaves).

Reporter, effector, and reference plasmids were co-transfected into protoplasts

by the polyethylene glycol (PEG)-assisted method (Yoshida *et al.*, 2013). After

incubating transfected protoplasts at 22°C for 16–19 h, FLUC and RLUC activities were

measured using a Pikka Gene Dual Assay Kit (Toyo Ink, Tokyo, Japan).

Phylogenetic analysis

Sequences were collected as described previously (Suzuki *et al.*, 2021). The full-length

peptide sequence of CjbHLH1 (BAJ40865.1; Yamada *et al.*, 2011) was retrieved from

GenBank (https://www.ncbi.nlm.nih.gov/genbank/). The sequences of Bl

(Csa5g156220) and Bt (Csa5g157230; Shang *et al.*, 2014) were obtained from the

Cucurbit Genomics Database (http://cucurbitgenomics.org/). A phylogenetic tree was

constructed by the neighbor-joining method with 1000 replicates using MEGA6

software (Tamura *et al.*, 2013) following alignment by MUSCLE (Edgar, 2004).

A phylogenetic tree of lateral organ boundaries domain (LBD) proteins was

generated using Clustal Omega v. 1.2.3 (Sievers *et al.*, 2011), FastTree v. 2.1.10 (Price

et al., 2010), and MEGA X (Kumar *et al.*, 2018) following our previous report (Suzuki

et al., 2021).

Results

Hydroponic culture enhances BA accumulation in *L. japonicus* hypocotyl and roots

Previously, we reported triterpene profiling in the model Fabaceae *L. japonicus* grown

on soil (Fig. S2; Suzuki *et al.*, 2019). To investigate triterpene production characteristics

in this plant, we conducted triterpene sapogenin profiling of whole roots from

soil-grown and hydroponically cultured plants, and liquid-cultured adventitious roots

(Figs. 1a and S3). Because triterpenes and saponins were extracted and acid-hydrolyzed

to remove sugar moieties attached to aglycones, we detected soyasapogenols instead of

soyasaponins by GC-MS. BA (peak 6) and soyasapogenol B (SB; a major

soyasapogenol in this plant, peak 10) were detected in the extract of soil-grown roots.

Liquid culture of adventitious roots induced by indole-3-butyric acid accumulated BA

and SB with lower peak intensities. Hydroponic roots of *L. japonicus* accumulated a

larger amount of BA than soil-cultured roots and adventitious roots, which caused peak

leading in the GC-MS chromatogram. BA was detected in the extracts of roots without

acid hydrolysis, indicating that BA is not glycosylated *in planta*. MeJA induces

soyasaponin but not BA biosynthesis in Fabaceae plants (Hayashi *et al.*, 2003;

Broeckling *et al.*, 2005; Suzuki *et al.*, 2005). We analyzed Lj suspension cultured cells

with and without MeJA (Figs. 1b and S3). Neither BA nor SB was detected in the

extract of untreated cultured cells. Extracts of the MeJA-treated cells contained a small

amount of SB but no BA.

We also analyzed the model Fabaceae plant *M. truncatula* (Figs. 1c and S3).

Although soil-grown roots of *M. truncatula* (cv. R108) did not accumulate BA, BA was

detected in the liquid-cultured adventitious roots and roots of hydroponically cultured

plants. To our knowledge, this is the first report of BA accumulation in *Medicago* spp.

We also analyzed the adventitious roots of cv. Jemalong A17 and confirmed the

presence of BA. However, the amount of BA in *M. truncatula* was much lower than that

in *L. japonicus*. *L. japonicus* is more appropriate to study the regulatory mechanisms of

BA biosynthesis enhanced in hydroponic culture.

SA produces and accumulates BA

Next, we evaluated BA production in hydroponic roots. We analyzed triterpenes

extracted from whole roots of *L. japonicus* plants hydroponically cultured for 2–12

weeks and observed a time course-dependent increase in BA and enlargement of the

hypocotyl and main roots (Fig. S4a–c). Development of a white, spongy tissue was

confirmed by anatomical analysis (Fig. 2a), and we assumed that this tissue was

homologous to SA in *G. max* (soybean). Soybean develops SA from the SM on stem,

hypocotyl, main and lateral roots, and nodules as an adaptive trait to survive flooding by

promoting oxygen diffusion in roots (Shimamura *et al.*, 2010; Yamauchi *et al.*, 2013).

Hydroponic culture likely resulted in a lack of oxygen in roots, similar to flooding.

Hydroponic roots accumulated 10-fold more BA than soil-grown roots, and its quantity

reached 6.5 mg/100 mg-dry weight (Fig. 2b). Isolated SA accumulated a higher amount

of BA (> 12 mg/100 mg-dry weight) than stele and lateral roots (Fig. 2b). However, SB

did not increase in hydroponic roots and SA (Fig. 2c).

We analyzed the expression levels of biosynthetic genes for BA (*OSC3* and

CYP716A51) and SB (*OSC1* and *CYP93E1*) (Fig. 3) by qRT-PCR. When comparing

soil-grown and hydroponic roots, BA biosynthetic genes were upregulated in

hydroponic roots (Fig. 3). For metabolite analysis (Fig. 2), we harvested SA, including

cortical cells. To identify the tissue biosynthesizing BA, we isolated SA and SM cell

layers by LMD (Fig. S5) and performed RNA extraction and qRT-PCR. Because BA

biosynthetic gene expression was higher in SA (Fig. 3), more BA is likely synthesized

in SA. These results suggest that transcriptional regulation is responsible for the

increased BA level in SA of *L. japonicus* hydroponic roots.

Candidate regulators of BA biosynthesis as identified by RNA-seq

We performed an RNA-seq-based comparative transcriptome analysis for mining

potential transcriptional regulators of BA biosynthesis. RNA evaluated by qRT-PCR

(Fig. 3) was used for the analysis, and sequencing of the whole root samples and the

LMD samples was conducted in three and four replicates, respectively. Data processing

was performed in triplicate for all samples as mentioned in Materials and Methods. We

performed differential expression analyses independently in data set 1 [soil-grown roots

(So) and hydroponic roots (Hy)] (Figs. 4a and S6a) and data set 2 (SM and SA) (Figs.

4b and S6b). Based on the MA plot (Fig. S6), we identified 3152 and 1994 DEGs whose

expression was higher in So and Hy, respectively (Table S4). SM and SA had 2169 and

1647 DEGs with elevated expression levels (Table S4). There were 256 DEGs

overlapping between Hy and SA (Fig. 4c), including 23 putative TF genes (Fig. 4c and

Table S5), as well as BA biosynthetic genes (*OSC3* and *CYP716A51*) and genes for

rate-limiting enzymes for triterpene biosynthesis [3-hydroxy-3-methyl-glutaryl-CoA

reductase1 (HMGR1) and squalene epoxidase1 (SQE1)].

We assessed the correlation between the expression patterns of BA biosynthetic

genes and those of candidate TF genes using the gene expression atlas in *Lotus* Base

(https://lotus.au.dk/expat/; 81 conditions based on annotation and prediction in

Miyakojima MG-20 genome v. 3.0). Seven TF genes were co-expressed with *OSC3*,

CYP716A51, *HMGR1*, and *SQE1* (Pearson's correlation coefficient [PCC] threshold:

0.4; Table 1). Among the seven TFs, we selected LjbHLH32 (LotjaGi4g1v0043000) and

LjbHLH50 (LotjaGi1g1v0340900) for further analysis because bHLH TFs are triterpene

biosynthesis regulators in Fabaceae (Mertens *et al.*, 2016; Tamura *et al.*, 2018), *Panax*

spp. (Zhang *et al.*, 2017), cucumber (Shang *et al.*, 2014), and *B. platyphylla* (Yin *et al.*,

2017).

The bHLH TFs were classified into approximately 25 clades based on sequence

similarity of the bHLH domain and other conserved domains (Heim *et al.*, 2003). We

constructed a phylogenetic tree of full-length bHLH proteins of clades Ia, Ib, and IVa,

including representative bHLH proteins involved in specialized plant metabolism (Fig.

4d). Fabaceae clade IVa bHLH TFs were further subdivided into three subclades, and

subclade IVa1 included all Fabaceae triterpene saponin biosynthesis activators identified

in clade IVa (Suzuki *et al.*, 2021). Interestingly, LjbHLH32 was a member of subclade

IVa2 but not IVa1 (Fig. 4d). The orthologs in subclade IVa2 were highly conserved in

Fabaceae plants and were commonly expressed in roots and nodules of model Fabaceae

plants (Suzuki *et al.*, 2021). However, no subclade IVa2 bHLHs have been functionally

characterized to date. The closest homolog of another candidate LjbHLH50 in *M.*

truncatula is MtbHLH1 (Medtr3g099620), which is suggested to regulate nitrogen

transfer between nodule and root cells by controlling nodule vascular patterning

(Godiard *et al.*, 2011). LjbHLH50 and MtbHLH1 showed 60% amino acid identity

(99% coverage) and belonged to clade Ia (Fig. 4d), similar to BpbHLH9, an OA and BA

biosynthesis regulator in *B. platyphylla* (Yin *et al.*, 2017). These phylogenetic

relationships between our candidates and known triterpene biosynthesis regulators

suggest that LjbHLH32 and LjbHLH50 are involved in BA biosynthesis regulation in *L.*

japonicus.

Overexpression of LjbHLH32 or LjbHLH50 enhances BA production in transgenic

hairy roots

We generated GFP-expressing (control), LjbHLH32-overexpressing (LjbHLH32-OX),

and LjbHLH50-OX hairy root lines for functional identification of LjbHLH32 and

LjbHLH50 *in planta*. We chose three independent lines for each construct and analyzed

their triterpene content by GC-MS (Fig. 5a). Compared to the control lines,

LjbHLH32-OX and LjbHLH50-OX hairy roots accumulated 5.73- and 4.31-fold more

BA (Fig. 5a, right), but the amount of SB was weakly elevated (Fig. 5a, left). Next, we

assessed the expression of triterpene biosynthetic genes by qRT-PCR (Fig. 5b).

LjbHLHs-OX lines overexpressed the corresponding bHLH genes. LjbHLH32

overexpression increased the transcript levels of the BA biosynthetic genes *LjHMGR1*,

LjSQE1, *OSC3*, and *CYP716A51*, whereas branch pathway genes [cycloartenol synthase

gene (*OSC5*), *OSC1*, and *LjCYP93E1*] were only slightly affected. LjbHLH50-OX lines

showed 1.56–2.26-fold higher expression of *LjHMGR1*, *LjSQE1*, *OSC5*, *OSC1*, and

LjCYP93E1. However, the expression of *OSC3* and *CYP716A51* was enhanced to a

greater degree.

We performed an RNA-seq analysis of nine hairy root lines (Fig. 5) to test the

effect of LjbHLH32 and LjbHLH50 overexpression on a genome-wide scale. The

control lines and bHLH-OX lines were distinguished by principal component analysis

(PCA) of transcriptome data (Fig. 6a and b). DEGs were identified based on an MA plot

(Fig. S6c and d). Compared to the control lines, 527 and 1045 genes were significantly

upregulated in LjbHLH32-OX and LjbHLH50-OX, respectively (Table S4). Among the

DEGs upregulated in bHLH-OX hairy roots and 256 DEGs commonly upregulated in

Hy and SA samples (Fig. 4c), 6 overlapped (Fig. 6c). These six genes included the BA

biosynthetic genes *OSC3* and *CYP716A51* but no genes involved in other triterpene

biosynthesis pathways (Table S6). Some of the enzyme genes putatively involved in

precursor supply for BA biosynthesis were significantly upregulated by overexpression

of bHLHs (Table S7), which may contribute to increased BA production in hairy roots

(Fig. 5a).

The genes upregulated in the LjbHLH32- and LjbHLH50-OX lines (Fig. 6c)

are likely involved in the metabolism of primary metabolites and plant hormones,

transport of metabolites, signaling, and defense responses (Table S6). *LjHMGR1* and

LjSQE1 were significantly upregulated by LjbHLH32-overexpression (Table 2).

Interestingly, LjbHLH32-OX hairy roots showed significantly elevated expression of

genes (*AMY2*, *CYP88D4*, and *CYP88D5*) in the *AMY2* gene cluster (Tables 2 and S7;

Krokida *et al.*, 2013). The products of the *AMY2* gene cluster in *L. japonicus* are

unknown. AMY2 produced β-amyrin and lupeol in a yeast expression system

(Iturbe-Ormaetxe *et al.*, 2003), and β-amyrin and dihydro-lupeol by agro-infiltration

into tobacco leaves (Krokida *et al.*, 2013). CYP88D4 and CYP88D5, homologs of

CYP88D6 (involved in glycyrrhizin biosynthesis in *G. uralensis*), did not oxidize the

tested triterpenes (Seki *et al.*, 2008; Krokida *et al.*, 2013). CYP71D353 showed

dihydro-lupeol oxidation activity when transiently expressed in tobacco leaves (Krokida

et al., 2013); however, the expression level of *CYP71D353* was not affected by

overexpression of LjbHLH32 (Table 2).

LjbHLH50-overexpression resulted in significant upregulation of several of the

23 TF genes selected by RNA-seq (Table 1), but those TF genes were not upregulated

by LjbHLH32 overexpression (Table S6).

LjbHLH32 shows similar trends to GubHLH3 in a transient effector-reporter

analysis

Analyses of hairy roots indicated that overexpression of LjbHLH32 or LjbHLH50

upregulated BA biosynthesis directly or indirectly. To obtain insight into how they

regulate BA biosynthesis, we examined the transactivation abilities of LjbHLH32 and

LjbHLH50 against the promoter sequences of triterpene biosynthetic genes (Table S3)

by transient effector-reporter analysis. Effector (*35S::bHLH*), reporter

(*promoter*::*FLUC*), and reference (*35S*::*RLUC*) constructs were co-transfected into

Arabidopsis mesophyll cell protoplasts and transactivation activities were measured as

relative luciferase activities (FLUC/RLUC; Fig. 7). We used the combination of

GubHLH3 and *GuCYP93E3pro* as the positive control and confirmed a similar level of

transactivation (39-fold) as a previous transient effector-reporter analysis using tobacco

BY-2 cell protoplasts (Tamura *et al.*, 2018). Similarly, GubHLH3 transactivated

promoters of soyasaponin biosynthetic genes in *L. japonicus* except *OSC1pro*

(*LjCYP93E3pro*, *LjCYP72A61pro*, *LjCSyGTpro*, and *LjDDMPTpro*) and the genes

involved in precursor supply (*LjHMGR1pro* and *LjSQE1pro*). GubHLH3 weakly

transactivated the promoter of *OSC5* (*cycloartenol synthase*), consistent with a previous

study (Tamura *et al.*, 2018). LjbHLH32 and LjbHLH50 did not transactivate *OSC3pr*o

and *CYP716A51pro*. Unexpectedly, LjbHLH32 significantly transactivated the

promoters of genes in the soyasaponin, precursor, and sterol pathways. Although the

fold-changes induced by LjbHLH32 were generally lower than those by GubHLH3, the

transactivation pattern of LjbHLH32 was highly correlated to that of GubHLH3

(PCC = 0.99). LjbHLH50 did not positively affect any promoters and the results were

not correlated to those of GubHLH3 (PCC = 0.36).

Discussion

SA formation and BA biosynthesis

Plant specialized metabolites are often produced and accumulated in specific cell types

and organs, such as artemisinin in *Artemisia annua* (Duke & Paul, 1993; Duke *et al.*,

1994), nicotine in tobacco (Katoh *et al.*, 2005), glycyrrhizin in *Glycyrrhiza* spp.

(Kojoma *et al.*, 2010; Li *et al.*, 2014), and monoterpene indole alkaloid (MIA) in

Catharanthus roseus (Yamamoto *et al.*, 2019). In this study, we identified SA as a sink

tissue of BA in *L. japonicus*, which is formed on the surface of hypocotyls and roots

when cultured hydroponically (Figs. 2 and S4). BA negatively regulates nodulation by

controlling *ENOD40* (early nodulin gene) expression in *L. japonicus* (Delis *et al.*, 2011;

Suzuki *et al.*, 2019). More BA accumulated in SA than in roots of soil-grown plants

(Figs. 1a and 2), suggesting an unknown and SA-specific role of BA. The role of

aerenchyma, including SA, is to supply air to roots under flooded conditions (Colmer &

Pedersen, 2008; Shimamura *et al.*, 2010; Takahashi *et al.*, 2014). Our recent study

revealed that *G. max* also accumulated BA in SA under flooded conditions and

suggested that the accumulation of hydrophobic BA was essential to exclude water from

gas spaces and promote air diffusion (Takahashi et al. 2022).

Sucrose supply from leaves is required for SA formation (Striker & Colmer,

2017; Takahashi *et al.*, 2018). However, it remains elusive whether sucrose functions

only as a nutrient during cell differentiation from SM to SA, or also as a signaling

molecule (Tognetti *et al.*, 2013). Hydroponically cultured *L. japonicus* will be useful to

investigate such molecular mechanisms of SA formation.

Triterpene biosynthesis is downregulated in underground tissues of *Glycyrrhiza*

spp. lacking leaves during winter (Hayashi *et al.*, 1998; Ramilowski *et al.*, 2013), and

the total amount of triterpene decreased in aseptic liquid cultures of *G. uralensis* stolons

lacking aerial parts (Kojoma *et al.*, 2010; Tamura *et al.*, 2017). Liquid-cultured *L.*

japonicus adventitious roots also showed decreased triterpene contents compared to the

roots of soil-grown plants (Fig. 1a). Photosynthesis and sucrose supply from aerial parts

may affect not only SA formation but also triterpene biosynthesis in underground tissues

of Fabaceae plants.

Subclade IVa1 and IVa2 bHLHs are likely functionally diverged in Fabaceae

triterpene biosynthesis regulation

Clade IVa bHLHs are conserved factors regulating jasmonate-inducible terpene

metabolism in plants (Goossens *et al.*, 2017; da Silva Magedans *et al.*, 2020), including

that of MIA in *C. roseus* (Van Moerkercke *et al.*, 2015, 2016), and triterpene saponins

in *Chenopodium quinoa* (Jarvis *et al.*, 2017), *Panax notoginseng* (Zhang *et al.*, 2017),

and Fabaceae plants (Mertens *et al.*, 2016; Tamura *et al.*, 2018; Ribeiro *et al.*, 2020).

We showed that clade IVa bHLHs in Fabaceae are subdivided into three subclades; all

saponin biosynthesis activators identified to date in Fabaceae fell into subclade IVa1

(Suzuki *et al.*, 2021). BA biosynthesis is jasmonate-independent in Fabaceae plants

(Hayashi *et al.*, 2003; Broeckling *et al.*, 2005; Suzuki *et al.*, 2005), and LjbHLH32, a

member of subclade IVa2, activated BA biosynthesis in *L. japonicus* transgenic hairy

roots (Fig. 5). These results suggest that subclade IVa2 gained new target specificities

during the evolution from subclade IVa1 in Fabaceae plants.

Interestingly, similar to the soyasaponin biosynthesis activator GubHLH3,

LjbHLH32 transactivated the promoters of soyasaponin biosynthetic genes but not of

BA biosynthetic genes in transient effector-reporter assays using *A. thaliana* mesophyll

cell protoplasts (Fig. 7). The changes in *cis*-elements in promoters and DNA-binding

domains in TFs are likely important to rewire gene regulatory networks (Shoji, 2019).

In the case of MtTSARs, GubHLH3, and LjbHLH32, however, peptide sequences of the

basic domains are well conserved (Fig. S7). Differences in protein-protein interactions

among TFs may contribute to changes in target recognition by TFs. The bHLH-MYB

complex is implicated in the transcriptional regulation of anthocyanin biosynthesis. In

kiwifruit, *Actinidia chinensis*, the complex of AcbHLH42 and AcMYB123 regulated

tissue-specific anthocyanin biosynthesis (Wang *et al.*, 2019). Although single

overexpression of AcbHLH42 or AcMYB123 in *Arabidopsis* was not enough to

increase the anthocyanin content, their co-expression induced anthocyanin production in

seedlings (Wang *et al.*, 2019). LjbHLH32 may have missed its interacting partners in

Arabidopsis mesophyll cells (Fig. 7), unlike *L. japonicus* hairy roots (Fig. 5). To prove

this hypothesis, it is necessary to identify direct targets and interacting partners of

LjbHLH32.

Conserved roles of clade Ia bHLH TFs in BA biosynthesis regulation

Clade Ia bHLHs are subdivided into the SMF subclade (described in detail later)

regulating stomatal development and the others subclade (Ran *et al.*, 2013). BpbHLH9

in the others subclade is an OA and BA biosynthesis regulator in *B. platyphylla* (Yin *et*

al., 2017). LjbHLH50 is also of the others subclade and enhanced BA biosynthesis in

Lotus transgenic hairy roots (Fig. 5), suggesting that bHLHs in this subclade are

conserved regulators of BA biosynthesis in fabids.

LjbHLH50 did not transactivate the promoters tested (Fig. 7), suggesting that it

lacked its interacting partners in *Arabidopsis* mesophyll cells or indirectly regulated BA

biosynthesis via downstream TF cascades. Although overexpression of BpbHLH9

enhanced expression of some triterpene biosynthesis genes and accumulation of

triterpene compounds, the genes directly regulated by BpbHLH9 are unknown (Yin *et*

al., 2017). Our first transcriptome analysis of whole roots from soil-grown and

hydroponically cultured plants, and SA and SM cell layers isolated by LMD revealed 23

candidate TFs, including LjbHLH32 and LjbHLH50 (Fig. 4 and Table 1). We also

performed transcriptome profiling of transgenic hairy root lines (Fig. 6). We found

upregulation of 6 of 23 TF candidate genes by overexpression of LjbHLH50, which

were annotated as LBD, ERF, NAC, and SCREAM (SCRM)-like proteins (Table S6).

LjbHLH50 (and BpbHLH9) may indirectly regulate BA biosynthesis by upregulating

these TF genes.

587   In higher plants, SPEECHLESS (SPCH), MUTE, and FAMA in the SMF

subclade regulate stomatal development (Ohashi-Ito & Bergmann, 2006; MacAlister *et*

al., 2007; Pillitteri *et al.*, 2007). The clade IIIb bHLHs SCRM (ICE1) and SCRM2 are

essential for stomatal development because they formed heterodimers with SPCH,

MUTE, and FAMA (Kanaoka *et al.*, 2008). AtSCRM protein (At3g26744.1; 494 aa) has

bHLH (298–350 aa) and putative zipper (ZIP; 351–380 aa) domains (Chinnusamy *et al.*,

2003). Alignments between AtSCRM and AtSCRM-like (At2g40435.1) proteins

showed that SCRM-like proteins lacked a bHLH domain but possessed a partial ZIP

domain (Fig. S8). An analysis of the interaction between LjbHLH50 (BpbHLH9) and

SCRM-like proteins is interesting.

Sleeping BA biosynthesis ability in *M. truncatula*

The *M. truncatula* genome has putative lupeol synthase genes but lupeol derivatives

have not been detected in *Medicago* spp. (Naoumkina *et al.*, 2010). We found that

whole roots from hydroponically cultured plants and liquid-cultured adventitious roots

accumulated trace amounts of BA (Fig. 1c). Because homologs of LjbHLH32 and

LjbHLH50 were present in *M. truncatula* (Fig. 4d), expression of these TFs might limit

BA biosynthesis in this plant.

605   Silencing of *OSC3* expression and reduced accumulation of lupeol derivatives

results in upregulated *ENOD40* expression and enhanced nodulation in *L. japonicus*

(Delis *et al.*, 2011). Similarly, in *M. truncatula* roots, overexpression of chimeric

MtbHLH1 fused to the EAR repression domain (transgenic plants should show

dominant loss-of-function phenotypes; Hiratsu *et al.*, 2003) resulted in abnormal nodule

vascular bundle development and slightly increased numbers of nodules per plant

although the growth of untransformed aerial parts was significantly reduced in the

composite plants (Godiard *et al.*, 2011). Although it is unknown whether MtbHLH1

positively regulates BA biosynthesis similar to its ortholog LjbHLH50 and whether *M.*

truncatula nodules accumulate lupeol derivatives, Delis *et al.* (2011) and Godiard *et al.*

(2011) likely reported similar phenomena. Re-evaluation of these mutants will provide

rigid understanding for the role of lupeol derivatives and clade Ia bHLHs in nodule

formation.

Godiard *et al*. (2011) suggested that an LBD-type TF gene (Medtr7g007010) is

a potential target of MtbHLH1 by transcriptome analysis of transgenic roots

overexpressing MtbHLH1-EAR. An LBD gene was co-upregulated in hydroponic roots,

SA, and LjbHLH50-OX hairy roots (Lj5g3v2099760/LotjaGi5g1v0323400; Table S6)

and closely related to Medtr07g007010 in the phylogenetic tree (Fig. S9), suggesting

the involvement of LBD proteins in this yet-to-be-defined TF cascade.

Concluding remarks

We identified two bHLH-type TFs enhancing BA biosynthesis by transcriptome analysis

of hydroponically cultured *L. japonicus* and a functional analysis using transgenic hairy

roots. However, neither LjbHLH32 nor LjbHLH50 transactivated the promoters of BA

biosynthetic genes in transient effector-reporter assays using a heterologous host

(*Arabidopsis*), suggesting that these bHLHs interact with unknown TFs or indirectly

regulate BA biosynthesis via downstream TFs. The 23 putative TFs found in the first

screening are promising targets for further analysis (Table 1). This study is a first step

toward understanding the complicated transcriptional regulatory mechanisms of BA

biosynthesis in *L. japonicus*.

Data Availability

The nucleotide sequence reported in this paper has been submitted to the DNA Data

Bank of Japan (DDBJ) under the accession numbers of LC609881 (*LjbHLH32*) and

LC609882 (*LjbHLH50*). The raw RNA-seq reads obtained were deposited in the DDBJ

Sequence Read Archive (DRA) under the accession number DRA011635.

Acknowledgements

We thank Miyazaki University and National BioResource Project (NBRP) for providing

pUB-GWS-GFP plasmid and *L. japonicus* Gifu B-129 seeds, RIKEN BRC for

providing Lj suspension cell line (rpc00032) through NBRP, the Institut National de la

Recherche Agronomique for providing *M. truncatula R108* and Jemalong A17 seeds,

Prof. Shuhei Yasumoto (Osaka University) for providing genomic DNA of *A. thaliana*

Col-0, and Dr. Atsushi Toyoda (National Institute of Genetics) for construction and

sequencing of RNA-seq library of soil-grown and hydroponically-cultured whole roots.

We also thank Dr. Nobutaka Mitsuda, Dr. Shingo Sakamoto and Ms. Tomoko Niki

(AIST) for technical support on transient effector-reporter analysis. This work was

supported by the Ministry of Education, Culture, Sports, Science and Technology

(MEXT), Japan [JSPS KAKENHI grant Nos. 16H06279 (PAGS), 17K07754 and

JP20H02913 to HSe, JP19H02921 to TM and JP19J10245 to HSu, and the Frontier

Research Base for Global Young Researchers, Osaka University, to EOF].

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

Author Contribution

EOF, TM and HSe conceived and supervised the study; All authors designed the

experiments; HSu and HT performed the experiments; HSu wrote the manuscript; HT,

EOF, MN, TM and HSe made manuscript revisions. All authors read and approved the

final manuscript.

References

Bolger AM, Lohse M, Usadel B. **2014**. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**: 2114–2120.

Broeckling CD, Huhman DV, Farag MA, Smith JT, May GD, Mendes P, Dixon RA,
Sumner LW. **2005**. Metabolic profiling of *Medicago truncatula* cell cultures reveals
the effects of biotic and abiotic elicitors on metabolism. *Journal of Experimental Botany*
56: 323–336.

Chinnusamy V, Ohta M, Kanrar S, Lee B, Hong X, Agarwal M, Zhu J-K. **2003**.
ICE1: a regulator of cold-induced transcriptome and freezing tolerance in *Arabidopsis*.
Genes & Development **17**: 1043–1054.

Chung SY, Seki H, Fujisawa Y, Shimoda Y, Hiraga S, Nomura Y, Saito K,
Ishimoto M, Muranaka T. **2020**. A cellulose synthase-derived enzyme catalyses 3- *O*
-glucuronosylation in saponin biosynthesis. *Nature Communications* **11**: 5664.

Colmer TD, Pedersen O. **2008**. Oxygen dynamics in submerged rice (*Oryza sativa*).

New Phytologist **178**: 326–334.

Conn SJ, Hocking B, Dayod M, Xu B, Athman A, Henderson S, Aukett L, Conn V,
Shearer MK, Fuentes S, *et al.* **2013**. Protocol: optimising hydroponic growth systems
for nutritional and physiological analysis of *Arabidopsis thaliana* and other plants.
Plant Methods **9**: 4.

da Silva Magedans YV, Phillips MA, Fett-Neto AG. **2020**. Production of plant
bioactive triterpenoid saponins: from metabolites to genes and back. *Phytochemistry*
Reviews **20**: 461–482.

Delis C, Krokida A, Georgiou S, Peña-Rodríguez LM, Kavroulakis N, Ioannou E,
Roussis V, Osbourn AE, Papadopoulou KK. **2011**. Role of lupeol synthase in *Lotus*
japonicus nodule formation. *New Phytologist* **189**: 335–346.

Duke SO, Paul RN. **1993**. Development and fine structure of the glandular trichomes
of *Artemisia annua* L. *International Journal of Plant Sciences* **154**: 107–118.

Duke MV, Paul RN, Elsohly HN, Sturtz G, Duke SO. **1994**. Localization of
artemisinin and artemisitene in foliar tissues of glanded and glandless biotypes of
Artemisia annua L. *International Journal of Plant Sciences* **155**: 365–372.

Edgar RC. **2004**. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Research* **32**: 1792–1797.

Fenwick G. **1992**. Saponins. *Toxic substances in crop plants*: 285–327.

Godiard L, Lepage A, Moreau S, Laporte D, Verdenaud M, Timmers T, Gamas P.
2011. MtbHLH1, a bHLH transcription factor involved in *Medicago truncatula* nodule
vascular patterning and nodule to plant metabolic exchanges. *New Phytologist* **191**:
391–404.

Goossens J, Mertens J, Goossens A. **2017**. Role and functioning of bHLH
transcription factors in jasmonate signalling. *Journal of Experimental Botany* **68**: 1333–
1347.

Hayashi H, Hiraoka N, Ikeshiro Y, Yamamoto H, Yoshikawa T. **1998**. Seasonal
variation of glycyrrhizin and isoliquiritigenin glycosides in the root of *Glycyrrhiza*
glabra L. *Biological & Pharmaceutical Bulletin* **21**: 987–989.

Hayashi H, Huang P, Inoue K. **2003**. Up-regulation of soyasaponin biosynthesis by
methyl jasmonate in cultured cells of *Glycyrrhiza glabra*. *Plant and Cell Physiology* **44**:
404–411.

Heim MA, Jakoby M, Werber M, Martin C, Weisshaar B, Bailey PC. 2003. The
basic helix-loop-helix transcription factor family in plants: A genome-wide study of
protein structure and functional diversity. *Molecular Biology and Evolution* **20**: 735–
747.

Hiratsu K, Matsui K, Koyama T, Ohme-Takagi M. **2003**. Dominant repression of
target genes by chimeric repressors that include the EAR motif, a repression domain, in
Arabidopsis. *The Plant Journal* **34**: 733–739.

Horacio P, Martinez-Noel G, Tognetti JA. **2013**. Sucrose signaling in plants: A world
yet to be explored. *Plant Signaling & Behavior* **8**: e23316.

Huhman DV, Berhow MA, Sumner LW. **2005**. Quantification of saponins in aerial
and subterranean tissues of *Medicago truncatula*. *Journal of Agricultural and Food
Chemistry* **53**: 1914–1920.

Iturbe-Ormaetxe I, Haralampidis K, Papadopoulou K, Osbourn AE. **2003**.
Molecular cloning and characterization of triterpene synthases from *Medicago
truncatula* and *Lotus japonicus*. *Plant Molecular Biology* **51**: 731–743.

**Jarvis DE, Ho YS, Lightfoot DJ, Schmöckel SM, Li B, Borm TJA, Ohyanagi H,
Mineta K, Michell CT, Saber N, *et al.* 2017**. The genome of *Chenopodium quinoa*.
Nature **542**: 307–312.

**Jozwiak A, Sonawane PD, Panda S, Garagounis C, Papadopoulou KK, Abebie B,
Massalha H, Almekias-Siegl E, Scherf T, Aharoni A**. **2020**. Plant terpenoid
metabolism co-opts a component of the cell wall biosynthesis machinery. *Nature
Chemical Biology* **16**: 740–748.

**Kamal N, Mun T, Reid D, Lin JS, Akyol TY, Sandal N, Asp T, Hirakawa H,
Stougaard J, Mayer KFX, Sato S, Andersen SU. 2020.** Insights into the evolution of
symbiosis gene copy number and distribution from a chromosome-scale *Lotus
japonicus* Gifu genome sequence. *DNA Research* **27**:dsaa015.

Kanaoka MM, Pillitteri LJ, Fujii H, Yoshida Y, Bogenschutz NL, Takabayashi J,

Zhu J-K, Torii KU. **2008**. *SCREAM/ICE1* and *SCREAM2* specify three cell-state
transitional steps leading to *Arabidopsis* stomatal differentiation. *The Plant Cell* **20**:
1775–1785.

Katoh A, Ohki H, Inai K, Hashimoto T. **2005**. Molecular regulation of nicotine
biosynthesis. *Plant Biotechnology* **22**: 389–392.

Kojoma M, Ohyama K, Seki H, Hiraoka Y, Asazu SN, Sawa S, Sekizaki H,
Yoshida S, Muranaka T. **2010**. *In vitro* proliferation and triterpenoid characteristics of
licorice (*Glycyrrhiza uralensis* Fischer, Leguminosae) stolons. *Plant Biotechnology* **27**:
59–66.

Krokida A, Delis C, Geisler K, Garagounis C, Tsikou D, Peña-Rodríguez LM,
Katsarou D, Field B, Osbourn AE, Papadopoulou KK. **2013**. A metabolic gene
cluster in *Lotus japonicus* discloses novel enzyme functions and products in triterpene
biosynthesis. *New Phytologist* **200**: 675–690.

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. **2018**. MEGA X: Molecular
evolutionary genetics analysis across computing platforms. *Molecular Biology and*
Evolution **35**: 1547–1549.

Li B, Bhandari DR, Janfelt C, Römpp A, Spengler B. **2014**. Natural products in
Glycyrrhiza glabra (licorice) rhizome imaged at the cellular level by atmospheric
pressure matrix-assisted laser desorption/ionization tandem mass spectrometry imaging.
The Plant Journal **80**: 161–171.

MacAlister CA, Ohashi-Ito K, Bergmann DC. **2007**. Transcription factor control of
asymmetric cell divisions that establish the stomatal lineage. *Nature* **445**: 537–540.

Maekawa T, Kusakabe M, Shimoda Y, Sato S, Tabata S, Murooka Y, Hayashi M.
2008. Polyubiquitin promoter-based binary vectors for overexpression and gene
silencing in *Lotus japonicus*. *Molecular Plant-Microbe Interactions* **21**: 375–382.

Matsui T, Sawada K, Takita E, Kato K. **2014**. The longer version of *Arabidopsis*
thaliana heat shock protein 18.2 gene terminator contributes to higher expression of
stably integrated transgenes in cultured tobacco cells. *Plant Biotechnology* **31**: 191–194.

Mertens J, Pollier J, Bossche RV, Lopez-Vidriero I, Franco-Zorrilla JM, Goossens
A. **2016**. The bHLH transcription factors TSAR1 and TSAR2 regulate triterpene

saponin biosynthesis in *Medicago truncatula*. *Plant Physiology* **170**: 194–210.

Mochizuki T, Takahashi U, Shimamura S, Fukuyama M. **2000**. Secondary
aerenchyma formation in hypocotyl in summer leguminous crops. *Japanese Journal of*
Crop Science **69**: 69–73.

Nagaya S, Kawamura K, Shinmyo A, Kato K. **2010**. The HSP terminator of
Arabidopsis thaliana increases gene expression in plant cells. *Plant and Cell Physiology*
51: 328–332.

Naoumkina MA, Modolo LV, Huhman DV, Urbanczyk-Wochniak E, Tang Y,
Sumner LW, Dixon RA. **2010**. Genomic and coexpression analyses predict multiple
genes involved in triterpene saponin biosynthesis in *Medicago truncatula*. *The Plant*
Cell **22**: 850–866.

Ohashi-Ito K, Bergmann DC. **2006**. *Arabidopsis* FAMA controls the final
proliferation/differentiation switch during stomatal development. *The Plant Cell* **18**:
2493–2505.

Oshima Y, Shikata M, Koyama T, Ohtsubo N, Mitsuda N, Ohme-Takagi M. **2013**.
MIXTA-Like transcription factors and WAX INDUCER1/SHINE1 coordinately
regulate cuticle development in *Arabidopsis* and *Torenia fournieri*. *The Plant Cell* **25**:
1609–1624.

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. **2017**. Salmon provides fast
and bias-aware quantification of transcript expression. *Nature Methods* **14**: 417–419.

Pillitteri LJ, Sloan DB, Bogenschutz NL, Torii KU. **2007**. Termination of asymmetric
cell division and differentiation of stomata. *Nature* **445**: 501–505.

Price MN, Dehal PS, Arkin AP. **2010**. FastTree 2 – approximately
maximum-likelihood trees for large alignments. *PLoS ONE* **5**: e9490.

Ramilowski JA, Sawai S, Seki H, Mochida K, Yoshida T, Sakurai T, Muranaka T,
Saito K, Daub CO. **2013**. *Glycyrrhiza uralensis* transcriptome landscape and study of
phytochemicals. *Plant and Cell Physiology* **54**: 697–710.

Ran J-H, Shen T-T, Liu W-J, Wang X-Q. **2013**. Evolution of the bHLH genes
involved in stomatal development: implications for the expansion of developmental

complexity of stomata in land plants. *PLoS ONE* **8**: e78997.

Ribeiro B, Lacchini E, Bicalho KU, Mertens J, Arendt P, Bossche RV, Calegario G,
Gryffroy L, Ceulemans E, Buitink J, *et al.* **2020**. A seed-specific regulator of
triterpene saponin biosynthesis in *Medicago truncatula*. *The Plant Cell* **32**: 2020–2042.

Seki H, Ohyama K, Sawai S, Mizutani M, Ohnishi T, Sudo H, Akashi T, Aoki T,
Saito K, Muranaka T. **2008**. Licorice β-amyrin 11-oxidase, a cytochrome P450 with a
key role in the biosynthesis of the triterpene sweetener glycyrrhizin. *Proceedings of the*
National Academy of Sciences **105**: 14204–14209.

Seki H, Tamura K, Muranaka T. **2015**. P450s and UGTs: Key players in the
structural diversity of triterpenoid saponins. *Plant and Cell Physiology* **56**: 1463–1471.

Shang Y, Ma Y, Zhou Y, Zhang H, Duan L, Chen H, Zeng J, Zhou Q, Wang S, Gu
W, *et al.* **2014**. Biosynthesis, regulation, and domestication of bitterness in cucumber.
Science **346**: 1084–1088.

Shimamura S, Yamamoto R, Nakamura T, Shimada S, Komatsu S. **2010**. Stem
hypertrophic lenticels and secondary aerenchyma enable oxygen transport to roots of
soybean in flooded soil. *Annals of Botany* **106**: 277–284.

Shoji T. **2019**. The recruitment model of metabolic evolution: jasmonate-responsive
transcription factors and a conceptual model for the evolution of metabolic pathways.
Frontiers in Plant Science **10**: 560.

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,
Remmert M, Söding J, *et al.* **2011**. Fast, scalable generation of high-quality protein
multiple sequence alignments using Clustal Omega. *Molecular Systems Biology* **7**: 539.

Soneson C, Love MI, Robinson MD. **2015**. Differential analyses for RNA-seq:
transcript-level estimates improve gene-level inferences. *F1000Research* **4**: 1521.

Striker GG, Colmer TD. **2017**. Flooding tolerance of forage legumes. *Journal of*
Experimental Botany **68**: 1851–1872.

Su W, Sun J, Shimizu K, Kadota K. **2019**. TCC-GUI: a Shiny-based application for
differential expression analysis of RNA-Seq count data. *BMC Research Notes* **12**: 133.

Suzuki H, Fukushima EO, Shimizu Y, Seki H, Fujisawa Y, Ishimoto M, Osakabe K,

Osakabe Y, Muranaka T. **2019**. *Lotus japonicus* triterpenoid profile and
characterization of the *CYP716A51* and *LjCYP93E1* genes involved in their
biosynthesis in planta. *Plant and Cell Physiology* **60**: 2496–2509.

Suzuki H, Reddy MSS, Naoumkina M, Aziz N, May GD, Huhman DV, Sumner
LW, Blount JW, Mendes P, Dixon RA. **2005**. Methyl jasmonate and yeast elicitor
induce differential transcriptional and metabolic re-programming in cell suspension
cultures of the model legume *Medicago truncatula*. *Planta* **220**: 696–707.

Suzuki H, Seki H, Muranaka T. **2021**. Insights into the diversification of subclade IVa
bHLH transcription factors in Fabaceae. *BMC Plant Biology* **21**: 109.

Takahashi H, Abo C, Suzuki H, Romsuk J, Oi T, Yanagawa A, Gorai T, Tomisaki
Y, Jitsui M, Shimamura S *et al*. **2022.** Triterpenoids in aerenchymatous phellem
contribute to internal root aeration and waterlogging adaptability in soybean. *Research*
Square. https://doi.org/10.21203/rs.3.rs-2230730/v1

Takahashi H, Kamakura H, Sato Y, Shiono K, Abiko T, Tsutsumi N, Nagamura Y,
Nishizawa NK, Nakazono M. **2010**. A method for obtaining high quality RNA from
paraffin sections of plant tissues by laser microdissection. *Journal of Plant Research*
123: 807–813.

Takahashi H, Xiaohua Q, Shimamura S, Yanagawa A, Hiraga S, Nakazono M.
2018. Sucrose supply from leaves is required for aerenchymatous phellem formation in
hypocotyl of soybean under waterlogged conditions. *Annals of Botany* **121**: 723–732.

Takahashi H, Yamauchi T, Colmer TD, Nakazono M. **2014**. Aerenchyma formation
in plants. In: van Dongen JT, Licausi F, eds. Plant Cell Monographs. Low-Oxygen
Stress in Plants. Springer, Vienna, 247–265.

Tamura K, Seki H, Suzuki H, Kojoma M, Saito K, Muranaka T. **2017**.
CYP716A179 functions as a triterpene C-28 oxidase in tissue-cultured stolons of
Glycyrrhiza uralensis. *Plant Cell Reports* **36**: 437–445.

Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. **2013**. MEGA6: molecular
evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* **30**: 2725–
2729.

Tamura K, Yoshida K, Hiraoka Y, Sakaguchi D, Chikugo A, Mochida K, Kojoma

M, Mitsuda N, Saito K, Muranaka T, *et al.* 2018. The basic helix–loop–helix
transcription factor GubHLH3 positively regulates soyasaponin biosynthetic genes in
Glycyrrhiza uralensis. *Plant and Cell Physiology* **59**: 783–796.

Thimmappa R, Geisler K, Louveau T, O'Maille P, Osbourn A. **2014**. Triterpene
biosynthesis in plants. *Annual Review of Plant Biology* **65**: 225–257.

Tsuno Y, Fujimatsu T, Endo K, Sugiyama A, Yazaki K. **2018**. Soyasaponins: a new
class of root exudates in soybean (*Glycine max*). *Plant and Cell Physiology* **59**: 366–
375.

Van Moerkercke A, Steensma P, Gariboldi I, Espoz J, Purnama PC, Schweizer F,
Miettinen K, Vanden Bossche R, De Clercq R, Memelink J, *et al.* 2016. The basic
helix-loop-helix transcription factor BIS2 is essential for monoterpenoid indole alkaloid
production in the medicinal plant *Catharanthus roseus*. *The Plant Journal* **88**: 3–12.

Van Moerkercke A, Steensma P, Schweizer F, Pollier J, Gariboldi I, Payne R,
Vanden Bossche R, Miettinen K, Espoz J, Purnama PC, *et al.* 2015. The bHLH
transcription factor BIS1 controls the iridoid branch of the monoterpenoid indole
alkaloid pathway in *Catharanthus roseus*. *Proceedings of the National Academy of*
Sciences **112**: 8130–8135.

Wang L, Tang W, Hu Y, Zhang Y, Sun J, Guo X, Lu H, Yang Y, Fang C, Niu X, *et*
***al.* 2019**. A MYB/bHLH complex regulates tissue-specific anthocyanin biosynthesis in
the inner pericarp of red-centered kiwifruit *Actinidia chinensis* cv. Hongyang. *The Plant*
Journal **99**: 359–378.

Wu F-H, Shen S-C, Lee L-Y, Lee S-H, Chan M-T, Lin C-S. **2009**. Tape-*Arabidopsis*
Sandwich - a simpler *Arabidopsis* protoplast isolation method. *Plant Methods* **5**: 16.

Yamada Y, Kokabu Y, Chaki K, Yoshimoto T, Ohgaki M, Yoshida S, Kato N,
Koyama T, Sato F. **2011**. Isoquinoline alkaloid biosynthesis is regulated by a unique
bHLH-type transcription factor in *Coptis japonica*. *Plant and Cell Physiology* **52**:
1131–1141.

Yamamoto K, Takahashi K, Caputi L, Mizuno H, Rodriguez-Lopez CE, Iwasaki T,
Ishizaki K, Fukaki H, Ohnishi M, Yamazaki M, *et al.* 2019. The complexity of
intercellular localisation of alkaloids revealed by single-cell metabolomics. *New*
Phytologist **224**: 848–859.

Yamauchi T, Shimamura S, Nakazono M, Mochizuki T. **2013**. Aerenchyma
formation in crop species: A review. *Field Crops Research* **152**: 8–16.

Yin J, Li X, Zhan Y, Li Y, Qu Z, Sun L, Wang S, Yang J, Xiao J. **2017**. Cloning and
expression of *BpMYC4* and *BpbHLH9* genes and the role of *BpbHLH9* in triterpenoid
synthesis in birch. *BMC Plant Biology* **17**: 214.

Yoshida K, Sakamoto S, Kawai T, Kobayashi Y, Sato K, Ichinose Y, Yaoi K,
Akiyoshi-Endo M, Sato H, Takamizo T, *et al.* 2013. Engineering the *Oryza sativa* cell
wall with rice NAC transcription factors regulating secondary wall formation. *Frontiers*
in Plant Science **4**: 383.

Zhang X, Ge F, Deng B, Shah T, Huang Z, Liu D, Chen C. **2017**. Molecular Cloning
and Characterization of PnbHLH1 Transcription Factor in *Panax notoginseng*.
Molecules : A Journal of Synthetic Chemistry and Natural Product Chemistry **22**: 1268.

Tables

Table 1. Co-expression analysis of triterpene biosynthetic genes and 23 candidate

TF genes

Name	Gene ID		Pearson's correlation coefficient*			
	Gifu B-129 ver 1.2	Miyakojima MG-20 ver 3.0	OSC3	HMGR1	CYP716A51	SQE1
OSC3	LotjaGi2g1v0235400	Lj2g3v0914630	1.00	0.51	0.49	0.50
HMGR1	LotjaGi2g1v0352700	Lj2g3v1574250	0.51	1.00	0.66	0.68
CYP716A51	LotjaGi4g1v0438900	Lj4g3v3014670	0.49	0.66	1.00	0.63
SQE1	LotjaGi5g1v0037500	Lj5g3v0347610	0.50	0.68	0.63	1.00
NAC	LotjaGi1g1v0222300	Lj0g3v0171479	0.20	0.24	0.25	0.31
bHLH50 (Ia)	**LotjaGi1g1v0340900**	**Lj1g3v1525930**	**0.57**	**0.68**	**0.58**	**0.85**
MYB	LotjaGi1g1v0548000	-	-	-	-	-
MYB	LotjaGi1g1v0650400	Lj1g3v4012910	-0.54	-0.47	-0.54	-0.75
bHLH (Ib)	LotjaGi1g1v0667300	Lj0g3v0202379	0.23	0.14	0.07	0.48
WRKY	LotjaGi1g1v0694700	Lj1g3v4578840	-0.13	-0.03	-0.21	-0.25
bZIP	**LotjaGi1g1v0724300**	**Lj1g3v4315340**	**0.44**	**0.62**	**0.44**	**0.54**
WRKY	**LotjaGi1g1v0754400**	**Lj1g3v4830200**	**0.63**	**0.77**	**0.54**	**0.64**
Zinc finger	LotjaGi2g1v0209200	Lj2g3v0740620	-	-	-	-
LBD	**LotjaGi2g1v0242300**	**Lj2g3v0934070**	**0.58**	**0.79**	**0.81**	**0.82**
ERF	LotjaGi3g1v0009300	Lj0g3v0243159	-0.28	-0.48	-0.27	-0.39
LBD	LotjaGi3g1v0035700	Lj3g3v0397200	-0.01	0.30	0.67	0.26
SCREAM-like protein	LotjaGi3g1v0203400	Lj3g3v1404080	-	-	-	-
MYB_related	LotjaGi3g1v0380600	Lj0g3v0318769	-0.29	-0.24	-0.07	-0.11
NAC	LotjaGi4g1v0002900	Lj4g3v0050650	-0.49	-0.34	-0.31	-0.55
bHLH32 (IVa2)	**LotjaGi4g1v0043000**	**Lj0g3v0292969**	**0.74**	**0.79**	**0.85**	**0.67**
WRKY	**LotjaGi4g1v0379800**	**Lj4g3v2616130**	**0.67**	**0.70**	**0.47**	**0.69**
ERF	LotjaGi4g1v0422400	Lj4g3v2871880	0.39	0.50	0.34	0.56
Zinc finger	**LotjaGi5g1v0010400**	**Lj5g3v0165540**	**0.66**	**0.66**	**0.56**	**0.82**
NAC	LotjaGi5g1v0158000	Lj5g3v1093960	0.38	0.39	0.25	0.22
Zinc finger	LotjaGi5g1v0173100	-	-	-	-	-
LBD	LotjaGi5g1v0323400	Lj5g3v2099760	0.10	0.09	0.10	0.44
bHLH (IVc)	LotjaGi6g1v0344900	Lj6g3v2168640	-0.38	-0.43	-0.39	-0.52

906   *Pearson's correlation coefficient (PCC) values based on transcriptome data of MG-20 deposited in

907   *Lotus* Base. Bold indicates putative TF genes showing a > 0.4 of PCC to four BA biosynthetic genes.

908   **Table 2. Changes in the expression of triterpene biosynthetic genes**

Pathway	Name	Hy (vs So)	SA (vs SM)	LjbHLH32-OX (vs GFP)	LjbHLH50-OX (vs GFP)
TFs	*bHLH32*	**1.53**	**3.07**	**5.49**	1.00
	bHLH50	**1.68**	**2.01**	**-0.76**	**3.00**
Precursor	*HMGR1*	**1.53**	**4.03**	**2.88**	0.39
	SQE1	**1.22**	**3.02**	**1.30**	-0.11
Betulinic acid	*OSC3 (LUS)*	**4.24**	**2.73**	**4.18**	**2.42**
	CYP716A51	**3.92**	**4.02**	**3.18**	**1.41**
Soyasaponin /Oleanolic acid	*OSC1 (bAS)*	-0.21	**-1.17**	0.038	0.38
Soyasaponin	*LjCYP93E1*	-0.59	-0.59	-0.20	-0.023
	LjCYP72A61	-0.87	0.56	0.20	-0.11
	CSyGT	-0.47	0.28	0.11	0.27
	DDMPT	-0.87	0.79	0.20	-0.21
Ursolic acid	*OSC9 (aAS)*	-	-2.23	-	-
Sterol	*OSC5 (CAS)*	-0.2	-1.09	0.046	0.51
	OSC7 (LAS)	0.57	0.62	-0.30	0.43
AMY2 gene cluster	*AMY2 (mixAS)*	-0.95	-0.032	**1.56**	**-0.82**
	CYP71D353	1.03	-1.15	0.54	0.36
	CYP88D4	**2.14**	**4.34**	**1.43**	0.41
	CYP88D5	**1.13**	**2.25**	**1.48**	0.18

909   Log base 2 values of fold changes of expression levels are listed. Positive values mean upregulation

910   in BA-rich samples relative to control samples. Differentially expressed genes are indicated in bold.

911   Details are provided in Table S7.

 Figure Legends

Fig. 1. Triterpene sapogenin profiling by gas chromatography-mass spectrometry

(GC-MS). (a) Analysis of *Lotus japonicus* adventitious roots (AR) and whole roots of

soil-grown and hydroponically cultured plants. Increased betulinic acid (BA)

accumulation was observed in the extracts of hydroponic roots. A BA peak was

observed without acid hydrolysis during triterpene extraction. **(b)** Analysis of Lj

suspension cultured cells (LjCC) with and without methyl jasmonate (MeJA). Extracted

ion chromatograms (m/z = 189 and 306) are shown to assess the presence of BA and

soyasapogenol B (SB). **(c)** Extracted ion chromatograms (m/z = 189) of hydroponically

cultured and soil-grown roots and AR of *Medicago truncatula* cv. R108 and Jemalong

A17. The following authentic standards were analyzed (Suzuki *et al.*, 2019): β-amyrin

(1), α-amyrin (2), lupeol (3), 24-OH-β-amyrin (4), oleanolic acid (5), BA (6),

sophoradiol (7), ursolic acid (8), soyasapogenol E (9), SB (10), soyasapogenol A (11).

Fig. 2. Accumulation of betulinic acid (BA) in secondary aerenchyma (SA) tissue

induced by hydroponic culture. (a) Images of sections of hypocotyls. White, spongy

tissue (SA) had developed on the surface of hypocotyls and roots of *L. japonicus*

cultured hydroponically. Quantitation of BA **(b)** and soyasapogenol B **(SB; c)**.

Hydroponic roots and SA accumulated more BA than soil-grown roots (Soil) and stele,

respectively. Means ± standard errors (SEs, n = 3). LR, lateral roots.

Fig. 3. Quantitative reverse transcription-PCR (qRT-PCR) analysis of triterpene

biosynthetic genes. Relative expression levels of betulinic acid (BA) and soyasaponin

biosynthesis genes were measured in triplicate. The values were normalized to the

expression level of a reference gene (*LjUBQ1*). Means ± standard deviations (SDs). Soil,

soil-grown roots; hydroponics, hydroponically grown roots; SM, secondary meristem;

secondary aerenchyma, SA.

Fig. 4. Selection of candidate transcription factors (TFs) regulating betulinic acid

(BA) biosynthesis. Principal component analysis (PCA) of the transcriptomes of

soil-grown (So) and hydroponically cultured (Hy) roots **(a)**, and secondary meristem

(SM) and secondary aerenchyma (SA) tissues **(b)**. **(c)** Venn diagram of differentially

expressed genes (DEGs) upregulated in Hy and SA compared to So and SM,

respectively. **(d)** Phylogenetic relationships of basic helix-loop-helix proteins. TFs

known to regulate triterpene biosynthesis are highlighted by black circles. Blue and red

circles indicate LjbHLH32 and LjbHLH50, respectively. ER, endoplasmic reticulum;

OA, oleanolic acid; IQA, isoquinoline alkaloid.

Fig. 5. Metabolite and gene expression analysis of LjbHLH32- and

LjbHLH50-overexpressing (OX) transgenic hairy root lines. (a) Triterpenes were

extracted from three control (GFP-expressing) lines and a basic helix-loop-helix

(bHLH)-OX line. Accumulation of betulinic acid (BA) and soyasapogenol B (SB) in

hairy roots was analyzed by gas chromatography-mass spectrometry (GC-MS). Relative

amounts were calculated from the peak areas of BA and SB. **(b)** Relative expression

levels of bHLH and biosynthetic genes by quantitative reverse transcription-PCR

(qRT-PCR). Error bars indicate standard deviations (SD, n = 3).

Fig. 6. RNA-sequencing (RNA-seq) analysis of LjbHLH32- and

LjbHLH50-overexpressing (OX) transgenic hairy root lines. Principal component

analysis (PCA) of the transcriptomes of control and LjbHLH32-OX roots **(a)**, and

control and LjbHLH50-OX roots **(b)**. **(c)** Venn diagram of 256 differentially expressed

genes (DEGs) commonly upregulated in hydroponic whole roots (Hy) and secondary

aerenchyma (SA), and DEGs upregulated in bHLH-OX hairy root lines compared to

control (GFP-expressing) hairy roots.

Fig. 7. Transient effector-reporter analysis of LjbHLH32 and LjbHLH50 for the

promoters of triterpene biosynthetic genes. Effector, reporter, and reference plasmids

were co-transfected into *Arabidopsis* mesophyll cell protoplasts. Luciferase (LUC)

activity levels were measured after overnight incubation. Error bars indicate standard

errors (SE, n = 4). Asterisks indicate significant differences between the negative

control (NC) and transcription factors (TFs) by Dunnett's test (*$P < 0.05$, **$P < 0.01$,

***$P < 0.001$). PC, positive control; OA, oleanolic acid; BA, betulinic acid; UA, ursolic

acid.

Supporting Information

Fig. S1. Transfer DNA regions of gateway destination vectors for *L. japonicus*

transformation.

Fig. S2. Triterpene biosynthetic pathways in *L. japonicus*.

Fig. S3. Mass spectra of peaks 6 and 10 in the chromatograms in Fig. 1.

Fig. S4. Time-course-dependent increase in BA in hydroponic roots of *L. japonicus*.

Fig. S5. Magnified image of secondary aerenchyma from a hydroponic *L. japonicus*

plant.

Fig. S6. MA plots for differentially expressed gene (DEG) detection.

Fig. S7. Protein alignments of the basic domains of clade IVa bHLHs.

Fig. S8. Alignments of AtSCRM and AtSCRM-like proteins.

Fig. S9. Phylogenetic tree of LBD (LOB) proteins in *M. truncatula* and *L. japonicus*.

Table S1. List of primers.

Table S2. List of FastQ files obtained.

Table S3. Promoter sequences isolated in this study.

Table S4. List of differentially expressed genes (DEGs) identified in this study.

Table S5. Annotation of 256 differentially expressed genes (DEGs) commonly

upregulated in Hy and SA samples.

Table S6. Annotation of differentially expressed genes (DEGs) in Fig. 6c.

Table S7. Detailed results related to triterpene biosynthesis genes obtained by

TCC-GUI analysis.

The English in this document has been checked by at least two professional editors,

both native speakers of English. For a certificate, please see:

http://www.textcheck.com/certificate/3bPF0g

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig.6

Fig. 7