Title: Inferring the distributions of fitness effects and proportions of strongly

deleterious mutations

Running title: Inferring fraction of strongly deleterious mutations

Authors: Anders P. Charmouh[1], Greta Bocedi[1], Matthew Hartfield[2].

Author's affiliations: [1]School of Biological Sciences, University of Aberdeen, United

Kingdom. [2]Institute of Ecology and Evolution, The University of Edinburgh, EH9 3FL,

United Kingdom.

Corresponding author: Anders P. Charmouh (apc@birc.au.dk)

Abstract

The distribution of fitness effects is a key property in evolutionary genetics as it has implications for several evolutionary phenomena including the evolution of sex and mating systems, the rate of adaptive evolution, and the prevalence of deleterious mutations. Despite the distribution of fitness effects being extensively studied, the effects of strongly deleterious mutations are difficult to infer since such mutations are unlikely to be present in samples of haplotypes, so genetic data may contain very little information about them. Recent work has attempted to correct for this issue by expanding the classic gamma-distributed model to explicitly account for strongly deleterious mutations. Here, we use simulations to investigate one such method, adding a parameter (p_{lth}) to capture the proportion of strongly deleterious mutations. We show that p_{lth} can improve the model fit when applied to individual species but can underestimate the true proportion of strongly deleterious mutations. The parameter can also artificially maximize the likelihood when used to jointly infer a distribution of fitness effects from multiple species. As p_{lth} and related parameters are used in current inference algorithms, our results are relevant with respect to avoiding model artifacts and improving future tools for inferring the distribution of fitness effects.

Introduction

The distribution of fitness effects (DFE) among new mutations has long been recognized as having a fundamental importance in evolutionary genetics since the shape of this distribution affects many evolutionary phenomena. For example, the DFE governs the rate of adaptive evolution (Fisher 1930; Schultz and Lynch 1997; Spigler et al. 2017), the frequency (and hence prevalence) of mutations with different effect sizes (Haldane 1937; Agrawal and Whitlock 2012) and has important implication for phenomena such as inbreeding avoidance, the evolution of sex and mating system evolution (Kondrashov 1985; Hartfield and Keightley 2012; Hedrick and Garcia-Dorado 2016). Because of this, the DFE has been extensively studied over several decades (for a review, see Eyre-Walker and Keightley 2007). Many important results on the DFE have emerged from this large body of literature; the DFE differs among species (Charlesworth and Eyre-Walker 2006; Castellano et al. 2018) and the shape depends on how well-adapted the population is and its effective population size N_e (Eyre-Walker 2002; Lynch and Conery 2003; Goldstein 2013; Huber et al. 2017). While the full DFE is probably complex and multi-modal, parts of the DFE of deleterious mutations can be modeled as a leptokurtic gamma distribution (Loewe and Charlesworth 2006; Silander et al. 2007) or by a series of discrete bins showing the proportions of mutations with different effects (Keightley and Eyre-Walker 2010).

Most of these results on the DFE have been obtained through two experimental techniques: mutation-inducing experiments and mutation accumulation experiments. In the former, mutations are induced and their effects are evaluated by comparing different strains of a model organism using some trait, such as fertility or growth rate, as a fitness proxy (Sanjuán et al. 2004). In the latter and more common approach, an ancestral strain is split into small, separate populations which are kept for many generations under favourable conditions that impose very little selection (Ohnishi 1977; Halligan and Keightley 2009; Böndel et al.

2019). The effects of accumulated mutations can then be evaluated by comparing the fitness

proxy of the population strains with that of the ancestral strain. The DFE can also be

estimated by sampling and comparing haplotypes from individuals in the wild (Dobzhansky

and Wright 1941; Crow and Temin 1964; Eyre-Walker et al. 2006; Castellano et al. 2018).

Data from the sampled haplotypes are often analysed in the form of a site frequency spectrum

(SFS) which describes the frequency with which different alleles segregates in the sampled

population (Fisher 1931; Wright 1938; Evans et al. 2007).

One major challenge in DFE inference is measuring the occurrence of strongly

deleterious mutations. Assuming deleterious alleles are not completely recessive, their

mutation-selection frequency is generally proportional to $1/s$, where s is the mutation's

selection coefficient (Wright 1937; Crow and Kimura 1970), and mutations are less likely to

be present in a sample of haplotypes as they become more strongly deleterious due to the

large selective disadvantage they entail. In other words, the probability distribution of

mutational effects in segregating mutations only will almost certainly be different from the

probability distribution of all mutations, since strongly deleterious mutations (with $s \gg$

$1/2N_e$) have a tiny probability of showing up in a population sample (Crow and Kimura

1970; Reich and Lander 2001; Rice et al. 2015). This emphasises the importance of

distinguishing between an *observed* DFE, consisting of alleles segregating in the gene pool of

a population at a given timepoint, and an *input* DFE that represents the actual probability

distribution of mutational effects as they arise by mutation and are added to the gene pool.

Much work has been done on inferring what proportion of the DFE consists of

strongly deleterious mutations; a study on human polymorphism data puts the estimate at

<15% (Eyre-Walker et al. 2006). However, this proportion differs greatly among different

branches of the tree of life. The proportion of lethal mutations was found to be approximately

40% in the vesicular stomatitis virus (Sanjuán et al. 2004), and even above 70% in

Drosophila (Keightley and Eyre-Walker 2007). However, two mutations which are both

described as strongly deleterious can still be orders of magnitude different in effect as

mutations with, for example, $s = 10^2/N_e$ and $s = 10^5/N_e$ both satisfy $s \gg 1/2N_e$. As a

result, modern DFE inference algorithms such as DFE-alpha (Eyre-Walker and Keightley

2007; Keightley et al. 2016) and polyDFE (Tataru et al. 2017; Tataru and Bataillon 2020)

which all work by analysing a SFS, may struggle to describe the extent to which many

different strongly deleterious mutations arise. This is partly because very strongly deleterious

mutations are unlikely to show up in a genome sample. However, the number of non-

segregating sites in a genome sample (some of which are presumed to be under strong

purifying selection) may provide information about the rate at which strongly deleterious

mutations occur. In any case, it is hard to differentiate the effect sizes of mutations once the

effect size is large, since such a mutation will have essentially no probability of appearing in

a genome sample (Galtier and Rousselle 2020).

A few studies have attempted to account for these strongly deleterious mutations by

fitting a DFE under the assumption that a certain fraction of mutations are so strongly

deleterious that they will remain absent from a haplotype sample, regardless of N_e (Eyre-

Walker et al. 2006; Boyko et al. 2008; Elyashiv et al. 2010; Kim et al. 2017; Galtier and

Rousselle 2020). In particular, Galtier and Rousselle (2020) used DFE methods to infer the

mean $N_e s$ of mutations among species, thereby obtaining estimate of the difference in N_e

between different species under the assumption that species shared a common, gamma-

distributed DFE. The field of DFE research has often parameterized gamma distributions

through the usage of a shape parameter, often denoted β, and a mean value (Keightley and

Eyre-Walker 2007). The optimal shape parameter of the gamma-distributed DFE varied

greatly among species, but a good fit for similar shape parameters was obtained when a

fraction of strongly selected mutations, with an effect size making them deleterious to the

point of being "essentially independent of N_e" was included in the SFSs. Here, N_e-

independent mutations were defined as those deleterious enough so that their probability of

showing up in a haplotype sample is essentially zero regardless of N_e. Galtier and Rousselle

(2020) implemented this approach was implemented by assuming that a proportion of

$0 \leq p_{lth} \leq 1$ mutations were either lethal or so strongly selected that they never appeared in

a SFS; each entry in the selected SFS was subsequently scaled by $1 - p_{lth}$. The method

yielded a high model likelihood but produced very surprising results such that estimates of

the proportion of strongly deleterious mutations (including lethal and nearly lethal mutations)

in *Drosophila* being over 50%, which is well above an order of magnitude larger than

previous estimates (Lethal or very strongly deleterious mutations are estimated to occur at a

rate of approx. 10% of the rate of mildly deleterious mutations; Simmons and Crow 1977;

Crow and Simmons 1983). While this method provides a way of accounting for strongly

deleterious mutations that will be absent from genomic data, the properties and accuracies of

this approach has yet to be fully investigated.

Here, we investigate to what extent explicitly accounting for strongly deleterious

mutations in a gamma distributed DFE model improves the accuracy of inference when

inferring an input DFE from simulated and genomic data. We do this by formulating a

genetically explicit individual-based Wright-Fisher model to simulate steady-state observed

DFEs under different strengths of selective effects, then infer the DFEs using the state-of-the-

art DFE inference software polyDFE (Tataru et al. 2017). We show that explicitly accounting

for a proportion of strongly deleterious mutations can increase the accuracy of inference

when inferring the DFE of a single population. However, we also demonstrate that including

this parameter artificially increases the inferred proportion of strongly deleterious mutations

when considering SFSs from multiple different populations. We further show that the p_{lth}

resulting in the best model fit is not equivalent to the true proportion of lethal or strongly

deleterious mutations in both cases.

Methods

To simulate the observed DFE of a population, we model a diploid, sexually reproducing

Wright-Fisher population with discrete generations and a default population size of $N = 500$.

Individuals have two different chromosomes, each with two homologs. Both chromosomes

have a map length of $[0, R]$ where we used $R = 10$ such that any real number r with $0 \leq r \leq$

R denotes a unique site on the chromosome; the potential number of sites is therefore

effectively infinite (e.g. Roze and Rousset 2009). In the first chromosome, mutations are non-

neutral, although mutations that are effectively neutral can occur, while in the second

chromosome, mutations are fully neutral. The sites on the neutral and non-neutral

chromosome are used to calculate a neutral and non-neutral site frequency spectrum,

respectively. These are then used as input datasets for the DFE inference software (see

below).

Individuals go through a life cycle similar to that of SFS_CODE (Hernandez 2008).

Specifically, individuals are sampled with a probability relative to their fitness standardized

by the mean fitness of their sex. The fitness of the ith individual, w_i is defined as

$$w_i = \prod_{x=1}^{Nhet} (1 - s_x/2) \prod_{y=1}^{Nhom} (1 - s_y) \tag{1}$$

Thus fitness is the product of the effects of all $Nhet$ heterozygous mutations and all

$Nhom$ homozygous mutations, thereby using a model of selection where individuals effects

are additive within a locus (i.e., $h = 1/2$ for all mutations) and multiplicative between loci.

Selection coefficients s are sampled from a gamma distribution (see parameterization below).

Parents each provide one of their two homologs to their offspring after a number of

recombination events, sampled from a Poisson distribution with mean $R = 10$, has occurred.

Mutations then occur in offspring and the number of mutations is sampled from a Poisson

distribution with a mean of $U = 1.0$ mutations per chromosome per generation, yielding a per

genome mutation rate similar to that of some eukaryotes (Haag-Liautard et al. 2007). The

number of neutral and non-neutral mutations are sampled independently with the same mean.

The selection coefficients of new mutations are sampled from a gamma distribution with a

mean of $S = 2N_e s$ and a shape parameter of β. Sampled values of S are re-scaled to s by

calculating $s = S/2N_e$ using the estimated N_e for that parameter set (see below). Offspring

are created by randomly sampling a mother and a father for each offspring, until N viable

($w > 0$) offspring have been produced with a constant 1:1 sex ratio. Once N viable offspring

have been formed, these will constitute the new parent generation.

To estimate N_e we added a single, neutral, linked locus to the centre of the non-

neutral chromosome and mutated the locus at a Poisson-distributed rate with a mean of 1 by

adding a random value sampled from $N(0,1)$ to the allelic value, thus having the locus satisfy

$V_m = 1$, which is the new genetic variance from mutation appearing per generation (Lynch

and Hill 1986; Barton and Turelli 1989; Keightley and Otto 2006). The N_e of the population

was defined as the mean (over 50 replicates) steady-state variance at the neutral linked locus

after a burn-in period of $10N$ generations. Without selection, the expected variance at this

locus is $NV_M = N_e V_M$ (Lynch and Hill 1986) and this was confirmed by simulation

(Supporting Information; S1). Mutational effects were sampled as $2N_e s$ (Hernandez 2008),

then re-scaled by dividing the sampled value with $2N_e$ before applying it in Eq. 1. To

accomplish this, N_e was estimated before the Wright-Fisher simulation by plotting a standard

curve of estimated steady-state N_e against an assumed N_e (the value used to re-scale $2N_e s$ to

s) for a given gamma distribution of selective effects. This allowed us to know the

approximate steady-state N_e which would emerge in our Wright-Fisher simulation before we

used these to simulate data for SFS construction (Supporting information; S2).

Data for calculating SFSs were outputted from the simulation after a burn-in period of

$10N$ generations, after which the full neutral and selected genomes of 10 random individuals

(i.e. 20 haplotypes) where outputted and used to calculated 1 neutral and 1 selected unfolded

SFS. Unfolded SFSs were used throughout.

To infer the mean scaled selection coefficients of mutations $S = 2N_e s$ and the shape

parameter β, we used polyDFE v1.11 (Tataru et al. 2017; Tataru and Bataillon 2020). To

determine whether these parameters could be obtained using our simulation setup, we

estimated them from SFSs calculated from simulated populations with the same S and β

combinations that Tataru et al. (2017) used to test the inference of purely deleterious DFEs

(see their Fig. 4). We found that polyDFE could accurately infer them from our simulations

(Supporting Information; S3) For each single-species polyDFE run, an input file with 50

neutral and 50 selected SFS was used, wherein each pair of neutral and selected SFS was

calculated from the basis of a single replicate of a Wright-Fisher simulation (hence 50

Wright-Fisher simulations per polyDFE run set for single-species DFE). The assumed

sequence lengths were 20,000 base pairs. For the multi-species polyDFE runs, an input files

with 20 neutral and 20 selected SFSs from 3 different species was used (60 SFS). The

polyDFE command lines and init-files used are available in Supporting Information; S4.

The concept of p_{lth} was previously used when fitting a single DFE model to a

collection of SFSs obtained from multiple different species (Galtier & Rousselle 2020). To

examine the efficacy and validity of p_{lth} for fitting multi-species DFEs, we simulated various

gamma-distributed DFEs in different populations, calculated output SFSs for each

population, and inferred DFEs for datasets containing SFSs from multiple different

populations. We repeated this analysis with DFEs inferred from genomic data from different

species (Chen et al. 2017).

For single species inferences, we first used the model described to simulate

populations with $S = 2N_e s$ in the range of $[1000, 10000]$ with increments of 1000 and

$\beta = 0.4$. Each simulation was replicated 50 times (500 replicates in total). For each

simulation set, we then inferred S and β using polyDFE under 10 different p_{lth} values in the

range $[0.0, 0.9]$ with increments of 0.1. The parameter p_{lth} was implemented, following

Galtier and Rousselle (2020), by setting an initial p_{lth} value and scaling each entry of the

selected SFS by $1 - p_{lth}$. For each p_{lth} value, 10 polyDFE runs were performed (1000

polyDFE runs in total) since the performance of polyDFE is affected by the initial S value

used. For each combination of p_{lth} value and simulation set, half of the polyDFE runs were

run using randomly sampled initial values of the estimated parameters. These values were

sampled from a uniform real distribution $U(1000, 10000)$ for S and $U(0.2, 0.8)$ for β. For

the other half of the polyDFE runs, the true values of S and β were used as the initial value of

the estimated parameters (following Tataru et al. 2017, Supplemental file S1, page 3). To

thoroughly test how p_{lth} affects the overall DFE inference under weaker selection, we also

studied the effect of p_{lth} on S and β inference given different combinations of $S =$

$50, 250, 2500$ and $\beta = 0.15, 0.40, 0.65$. For the simulation sets with $S = 2N_e s$ in the range

of $[1000, 10000]$ with increments of 1000 and $\beta = 0.4$, we used standard results from

probability theory to calculate the true proportion of lethal mutations resulting from these

distributions (Appendix A) and compared it to the value of p_{lth} under which the highest

accuracy of inference was found.

We also studied the effects of p_{lth} on inference of multi-species DFEs from more than

one species, and in particular if β is fixed between species. This was studied with both

arbitrarily defined weakly deleterious DFEs (which were used because they contained hardly

any strongly deleterious mutations), and DFEs inferred from genomic data (Chen et al. 2017,

2020). First, we simulated 3 Wright-Fisher populations with different DFEs with the shape

parameters β and means S of $\beta = 0.15, S_D = 5, \beta = 0.40, S_D = 20$ and $\beta = 0.65, S_D = 40$.

We then constructed polyDFE input files using 20 neutral and selected SFSs from each

population. Using polyDFE, S and β were estimated from this input file under four different

p_{lth} values using similar procedure of replication and definition of initial values as for the

single-species case. Second, using DFEs estimated from genomic data, we ran the same

analysis by simulating the DFEs estimated for populations of *Arabidopsis lyrata*, *Capsella*

grandiflora and *Zea mays* using the data of Chen et al. (2017). We compared the likelihood

of the estimated parameters under 4 different p_{lth} values, so that they cover cases when the

initial values were correctly defined and when these were purposely highly inaccurate.

Following Galtier & Rousselle (2020), we also studied the effect of p_{lth} on S inference from

these DFEs given different fixed values of β (as opposed to jointly estimated). Confidence

intervals of estimates shown in plots were calculated as $\bar{x} \pm Z \cdot \sigma/\sqrt{n}$ with $\bar{x}$ being the mean

value, n being the sample size, and Z being the 97.5 percentile value of the *t*-distribution

given $n - 1$ degrees of freedom.

Results

Analytical results

To illustrate the utility of including a p_{lth} parameter in DFE inference and determine

where it might be most useful, we first present analytical results demonstrating which

deleterious alleles are unlikely to be present in a population sample. Assume that mutations

are introduced at a rate $\theta = 4N_e\mu L$ for N_e the effective population size, μ the per-site

mutation probability per generation, and L the number of sites where selected sites are likely

to arise (e.g., nonsynonymous sites;(Galtier 2016)). A focal mutation has a scaled selection

coefficient $S = 2N_e s$ as a heterozygote, and is in the population at a frequency x. If we sample

n haploid genomes, then the expected probability that this mutation is present in i of n

samples is (Tataru et al. 2017, Equation 1):

$$\mathbb{E}(P = i|\theta, S) = \theta \int_0^1 B(i, n, x) H(S, x) \, dx \tag{2}$$

251 Where B indicates the binomial distribution of sampling i selected alleles from n

252 genomes, and H the expected time that the allele spends between frequencies x and $x + \delta x$

253 (Wright 1938):

$$H = \frac{1 - e^{S(1-x)}}{x(1 - x)(1 - e^S)} \tag{3}$$

Note there are a few differences between Eq. 3 and the version in Tataru *et al.* (2017).

Eq. 2 scales s by $2N_e$ whereas Tataru *et al.* (2017) use $4N_e$ scaling, due to the different

manner in which heterozygote fitness is calculated. That is, we consider mutations to be

additive with a dominance coefficient of $h = 1/2$, such that the fitness effect of a single

heterozygous and homozygous mutation is $1 - sh$ and $1 - s$, respectively. S is also

multiplied by -1 in Eq. 3, so a positive value denotes a deleterious mutation with that

selection magnitude.

Using Poisson Random Field theory (Sawyer and Hartl 1992; Sethupathy and

Hannenhalli 2008), the probability of observing a certain number of sites with i selected

alleles is Poisson distributed with mean $\mathbb{E}(P = i)$ (Eq. 2). Hence, the probability that no sites

will have i selected alleles is $e^{-\mathbb{E}(P=i)}$, and the probability that no selected alleles are present

at all equals $\prod_{i=1}^{n-1} e^{-\mathbb{E}(P=i)} = e^{-\sum_{i=1}^{n-1}\mathbb{E}(P=i)}$. The repository with the code used for all

simulations also contains a *Mathematica* notebook to perform these calculations (see Data

Availability).

Fig. 1 outlines the probability that no selected allele will be present in a sample of

genomes, for different sample sizes and mutation rates. We see that if the mutation rate is low

($\theta = 0.1$) then even weakly-deleterious mutations ($S > 10$) are unlikely to be present. For $\theta =$

1, while a small sample is also unlikely to contain strongly deleterious mutations ($S > 100$), a

large sample size of 100 is likely to capture some strongly deleterious mutations with $100 < S$

< 1000. A very large sample can capture even more strongly deleterious mutations if the

mutation rate is high ($\theta = 10$). However, under all cases considered, it becomes unlikely to

sample extremely deleterious mutations (S approaching 5000), even with large sample sizes

and mutation rates. Overall, for realistic sample sizes and mutation rates, once S exceeds 100

it becomes difficult to capture strongly deleterious mutations in a population sample, and it is

in this parameter space that p_{lth} might be useful in capturing the proportion of these

deleterious mutations. A p_{lth}-type parameter might also be necessary to calculate the fraction

of all deleterious variants if the population mutation rate is very low.

281
Figure 1: The probability that a deleterious mutation with scaled selection coefficient S will

not be present in a sample of n genomes. Results are plotted for different n, θ values as

denoted by the legend. Note that we do not plot $S < 10$ since calculations do not assume

neutrality and break down as S approaches 0. Strongly deleterious mutations are less likely to

end up in a sample of haplotypes, making it is difficult to differentiate the effects sizes by

analysing site frequency spectra.

Model-fitting for single-species DFEs

In simulations with a high S, the accuracy of inference of S and β was affected by p_{lth} (Fig.

2A-B) and the accuracy of inference in 4/10 of all cases was highest under a non-zero value

of p_{lth} (Fig. 2C-D). In these cases, the accuracy of inference (measured as the difference

between $\log_2(estimate/simulation)$ and 0) decreased by a value between 0.13 and 0.99. In

terms of the actual estimates of S, this means that in the four cases where p_{lth} improved the

accuracy of inference, S was estimated 3%-49% (depending on simulation set) more

accurately than in the corresponding set where $p_{lth} = 0.0$. In general, S inference becomes

more inaccurate as p_{lth} increases, however the accuracy of inference of β is less affected

unless p_{lth} is very high (Fig 2). At low mutation rates, high accuracy of inference can be

achieved under a high p_{lth}, although this comes at the expense of the accuracy of inference of

β (see Supporting Information; S7).

300
301

Figure 2: Mean accuracy of inference for S (A) and β (B) under the assumption of different

values of p_{lth} for different Wright-Fisher simulation sets (differing in the simulation S,

ranging from 1000 to 10000). Each point shows the mean value of 10 polyDFE replicates.

Bars denote 95% confidence intervals (if they are not visible then they lie within the point).

For each simulation set (differing in S; color), bottom panels show the subset that had the

highest accuracy of inference of S. For these subsets, the accuracy of inference for S (C) and

β (D) are shown. A value of 0 is equivalent to perfect accuracy of inference. The results show

that for each simulation set (these differ in their value of S), the best accuracy of inference is

found when p_{lth} is either 0 or very low ($p_{lth} = 0.1, p_{lth} = 0.2$) when a DFE model is fitted

to SFS data from a single species.

Under low values of S and varying values of β, non-zero values of p_{lth} generally

decreased the accuracy of inference, highlighting that p_{lth} only improves the accuracy of

inference when S is high (Fig. 3).

Figure 3: As Figure 2, but instead using 9 simulation sets with different combinations of S

and β ($S = 50,\ 250,\ 2500$ and $\beta = 0.15, 0.40, 0.65$).

By calculating the true proportion of lethals (Appendix A), which can be used as a

lower bound for the proportion of strongly deleterious mutations, we found that the p_{lth} value

resulting in the highest accuracy of inference (Fig. 2, 3) were not equivalent to the true

proportion of lethals (Fig. 4), with the true proportions being much higher than those inferred

using p_{lth}.

Figure 4: The true proportion of strongly deleterious mutations as a function of S (same

values as used in Fig. 2) given $\beta = 0.40$ and $N_e = 500$ (Appendix A). Red lines show the

interval containing p_{lth} value which resulted in the highest accuracy of inference (Fig. 2, 3).

The results show that the lower bounds for true proportion of lethals for each simulation set

(points) is higher than the p_{lth} which results in the best accuracy of inference (0.0-0.2, i.e.

between the red lines).

Model-fitting for multi-species DFEs

We simulated weakly deleterious DFEs for multiple species that had essentially zero

probability of yielding lethal mutations (Supporting Information; S5). Despite this setup, we

found that the likelihood of the estimated parameters could be artificially maximized when

assuming a high p_{lth} value, when inferring a joint set of DFE parameters from SFSs across

different species (Fig. 5A). When a DFE model is fitted to SFSs from multiple species, the

model likelihood is artificially maximized because a non-zero value of p_{lth} pushes the entries

of the different SFSs closer together by making them numerically similar (Fig. 6). That is,

species with very different SFSs will seem to have more similar SFSs under a high value of

p_{lth}, because p_{lth} will result in the entries of these different SFS all being closer to 0.

Because of this, fitting a single DFE model to joint SFS data yields a better model likelihood

once p_{lth} is applied. Thus, we found that p_{lth} can erroneously result in a high likelihood of a

model despite essentially no strongly deleterious mutations being present.

344
Figure 5: Mean likelihood with 95% confidence intervals returned by polyDFE for a multi-

species DFE model against 10 p_{lth} values. β was inferred and S was assumed fixed at a

randomly sampled value. polyDFE runs were conducted on 20 neutral and selected SFSs

from 3 different simulated DFEs. **(A)** Model likelihood for inference based on simulated

weakly deleterious DFEs with the shape parameters β and means S of $\beta = 0.15, S = 5$,

$\beta = 0.40, S = 20$ and $\beta = 0.65, S = 40$ combined into one data input file. **(B)** Model

likelihood for inference based on the DFE of three populations simulated parameterized with

data from Chen et al. 2017 using DFEs inferred for *A. lyrata*, *C. grandiflora* and *Z. mays*. The

results show that (A) model likelihood can be maximized under a high value of p_{lth} despite

little or no strongly deleterious mutations being present in the DFE because p_{lth} makes

different SFSs more similar and (B) this effect can also occur for DFE estimated from natural

populations.

358
Figure 6: Value of the 20 entries of the SFSs of 3 different simulated DFEs (black, red, blue)

under four different p_{lth} values: (A) $p_{lth} = 0.0$, (B) $p_{lth} = 0.5$, (C) $p_{lth} = 0.6$, (D) $p_{lth} =$

0.7.Confidence intervals fall within points as shown. The results show that increasing the p_{lth}

value pushes the entries of different SFS together by making them more numerically similar.

p_{lth} can also artificially maximise the likelihood of DFE models when simulating

DFEs estimated from genomic data (Fig. 5B). Whether the likelihood of the DFE model

levels off or keeps increasing with p_{lth} depends on the sampled initial values for the polyDFE

runs. For high p_{lth} values, the initial values sampled for each polyDFE run becomes

increasingly important. This seems to happen as the likelihood of the inferred DFE model

drops substantially if the initial S fed into polyDFE was low, and the SFSs are modified by a

high p_{lth}value resulting in a seemingly very deleterious selected SFS. Because of this effect,

95% confidence intervals on the mean likelihood increase as p_{lth} increases (Fig. 5). The

parameter p_{lth} can also artificially maximize the likelihood of a DFE model in cases where β

is fixed and S is inferred, similar to the approach used by Galtier & Rousselle (2020) (Fig. 7).

374
Figure 7: Mean likelihood returned by polyDFE for estimates of S given combinations of

fixed β and p_{lth}. A multi-species DFE where 20 neutral and selected SFSs from DFEs based

on those estimated for *A. lyrata*, *C. grandiflora* and *Z. mays* by Chen et al. (2017) were

combined was used a data input for polyDFE. The dashed line show the mean β estimated for

A. lyrata, *C. grandiflora* and *Z. mays*. The results again shows that model likelihood can be

artificially maximized under the assumption of a high value of p_{lth} and the extent to which

this happens depends on the initial values supplies to the DFE inference software.

 Exploring wider parameter space

        The effect of p_{lth} on the accuracy of inference was also tested under larger population

 sizes and a larger number of sampled haplotypes (Supporting information; S6). While p_{lth}

 was less useful for correcting the underestimation of the mean S when a larger population

 size was used, the results remained qualitatively similar such that p_{lth} had a more positive

 effect on the accuracy of inference as S was increased. The ability of p_{lth} to artificially

 maximize the likelihood of a model fitted to SFS data from multiple species was also tested

 under larger simulated population and sample size and this yielded a result that was

 qualitatively the same (Supplementary Information; S6). That is, p_{lth} will also artificially

 maximize the likelihood of a DFE model based on multiple species when these species have

 larger populations. This makes intuitive sense given the mechanism through which p_{lth}

 seems to cause the artificial maximization of likelihood; that is, the entries of different SFSs

 will become numerically similar as a higher p_{lth} is used, regardless of the sample size.

        We also investigated inference in a single species using a five-fold lower mutation

 rate. Here, higher values of p_{lth} can result in the best accuracy of inference for S, however,

 this happens at the expense of poor accuracy of inference of β (Supporting information S7).

 Discussion

        Improving the accuracy with which we can infer the DFE is important since this is a

 fundamental parameter of evolutionary genetics with implications for many branches of

 biology. Constructing an accurate description of the DFE boils down to constructing a model

 which can describe a wide range of effect sizes of mutations at a broad range of frequencies,

 and hence can be applied to many different species. This is not a trivial problem, because

 while several models consisting of parametric distributions have been suggested, including

gamma, exponential and lognormal distributions, the evidence shows the DFE of deleterious

mutations follows some complex and multi-modal distribution rather than a simple

parametric one (Nielsen and Yang 2003; Sawyer et al. 2003; Eyre-Walker and Keightley

2007; Kousathanas and Keightley 2013). Thus, fitting a single parametric distribution model

to a deleterious DFE can result in a biased estimate, and this bias is especially prone to affect

the inferred proportion of strongly deleterious mutations (Kim et al. 2017). Our results

illustrate that once strongly deleterious mutations are present, expanding the typical gamma

distribution model of a DFE to one which accounts for a proportion of strongly deleterious

mutations (Galtier and Rousselle 2020) can in principle improve the accuracy of inference,

although it can artificially increase the likelihood of the inferred parameters when a single

DFE is fitted to data from different species or populations.

We show that when inferring a DFE from a single species, a model under which

$p_{lth} > 0$, can sometimes yield the highest likelihood even when the DFE does not contain a

fraction of p_{lth} strongly deleterious mutations. This result highlights that p_{lth} is not the true

proportion of lethals and should not be viewed as such. Future developments might produce a

more useful correcting factor that accounts for underrepresentation of strongly deleterious

mutations while being compatible with classic parametric distributions such as gamma

distributions. This would likely involve first calculating how much underestimating might

occur, assuming some gamma distributed DFE, and part of answering this would be to

calculate the mean effect size of mutations with $2N_e s > 2N_e$.

Further, the classic notion of a model likelihood is not well-defined when using p_{lth},

since p_{lth} is used to multiply all entries of the SFSs by $1 - p_{lth}$. This rescaling results in the

data itself being modified and consequently two DFE models with differing values of p_{lth}

will be models on different datasets. This means that the likelihood of two different DFE

models with different p_{lth} values cannot be compared in the same way. This explanation does

not mean that there is a problem with the concept of model likelihood in general; rather, it

instead means that model likelihood cannot be used for model selection in the same way

when p_{lth} is used, since different p_{lth} models are effectively attempting to explain different

datasets.

Given SFS data from a single species, our results illustrate that DFE inference

performs better on strongly deleterious DFEs when entries in the input SFS are modified by

some value p_{lth} to reflect the proportion of strongly deleterious mutations that will not end up

in a haplotype sample. This is in line with the results of Fig. 4A in Tataru et al. (2017)

showing that mutational effects are underestimated when the input DFE is strongly

deleterious, indicating that strongly selected mutations are not fully detected or represented in

the haplotype sample. This is one parameter space where including p_{lth} may help with

improving the accuracy of inference, albeit the resulting p_{lth} values are unlikely to reflect the

actual proportion of strongly deleterious mutations. This means that although including some

value p_{lth} may improve the model likelihood, p_{lth} cannot reliably be used to infer the

proportion of strongly deleterious mutations. A recent study wherein a DFE model was

derived on completely different principles (by considering gene regulatory networks and

metabolic pathways) also concluded that no parametric distribution suffices to describe the

DFE of strongly deleterious mutations, and that a satisfactory model must involve some extra

class of strongly deleterious mutation akin to p_{lth} (Brajesh et al. 2019). These findings

suggest our conclusion has broad generality beyond the Wright-Fisher model. In practice,

information about the optimal value of p_{lth} to use when fitting a DFE model to a particular

species may eventually be determined *a priori* based on the species in question. For example,

an application of Fisher's geometric model (Fisher 1930) to DFE theory yields the prediction

that as the complexity of an organism increases, a larger proportion of the DFE should be

strongly deleterious (Lourenço et al. 2011; Tenaillon 2014). This has been tested several

times and found to be consistent with data by using the level of pleiotropy as a proxy for the

complexity of an organism (Martin and Lenormand 2006; Huber et al. 2017).

It is well-known that the likelihood of an individual DFE model drops when fitted to

multiple species (Huber et al. 2017; Galtier and Rousselle 2020), which highlights that the

DFE differs among species (Eyre-Walker 2002; Lynch and Conery 2003). In this study, we

also show that p_{lth} can artificially improve the likelihood of parameter estimates when a

single DFE is fitted to multiple species. This result has several important implications. First,

it illustrates the point that while the data strongly suggests the DFE of deleterious mutations

usually follows a gamma and lethal model, it can still result in a misleading description of the

DFE when fitted to data from multiple species. Second, it shows that since p_{lth} consists of a

modification of the data, we should bear this in mind when assessing the likelihood of a

model; we find the likelihood of a DFE model is maximized under a high p_{lth} in cases where

the true DFE is known to be very weakly deleterious and known to not contain a proportion

of p_{lth} strongly deleterious mutations. Because of this, we can conclude that if the likelihood

of a given DFE model for multiple species is maximized under the assumption of some

$p_{lth} > 0$, this should not be considered evidence that the DFEs being modeled does indeed

contain a proportion of p_{lth} strongly deleterious mutations. Third, our results offer an

explanation of the mechanism by which this effect occurs, namely that p_{lth} makes entries of

different SFSs more numerically similar, thereby inflating the likelihood of the resulting DFE

model. Several studies model DFEs as following a gamma and lethal model, or gamma and

"point mass" model (Eyre-Walker et al. 2006; Elyashiv et al. 2010), but when such models

involve applying a correction factor to SFSs, they seem unsuitable for fitting a single DFE to

data from a group of species.

While our study is limited to purely deleterious DFEs, more work on improving the

accuracy of inference for DFEs with beneficial mutations is also needed, since recent work

shows that the occurrence of strongly beneficial mutations (much like strongly deleterious

mutations) can make DFE inference inaccurate (Booker 2020). Even in the study of purely

deleterious DFEs, attention has almost exclusively been focused on the effects of SNPs

(resulting from point mutations), but the DFE of INDELs (insertions and deletions) is

understudied and accounting for such mutations might require a slightly different modelling

approach (Barton and Zeng 2018). As in our study, mutational effects are most often assumed

to be additive, however the average dominance coefficient of new mutations appears to be

substantially lower than $h = 1/2$ (Mukai et al. 1972; Simmons and Crow 1977; Lynch et al.

1999; Fernández et al. 2004; Spigler et al. 2017). While it is implicitly assumed that current

DFE inference algorithms remains accurate when the assumption of additive dominance is

violated, very little work has been done to test this assumption (but see Wade et al. 2022).

Similarly, other "gamma + lethal" DFE models exist and since these differ slightly from p_{lth}

implemented in Rousselle & Galtier (2020), testing their ability compared to the p_{lth}

implementation of a "gamma + lethal" DFE model is a topic worthy of further research

(Boyko et al. 2008; Kim et al. 2017).

Experimental evidence suggest that the DFE of deleterious mutations follow a

bimodal or perhaps even multi-modal distribution, meaning that a good model fit may not be

possible under classic parametric distributions such as exponential, gamma, or lognormal

(Nielsen and Yang 2003; Sawyer et al. 2003; Eyre-Walker and Keightley 2007; Kousathanas

and Keightley 2013). Because of this, it would be worthwhile for future research to explore

whether DFE model fitting under different distribution could result in an improve model fit.

In a recent study, the DFE was represented by using a non-parametric distribution in the form

of several non-overlaping uniform distributions (Johri et al. 2020). Some of the current DFE

inference software can also be set to infer a DFE where selection coefficients take discrete

values rather than necessarily conforming to a single continuous distribution (Tataru et al.

2017). Expanding the set of distributions typically used for DFE model could prove to be

fertile grounds for new research and result in more accurate models.

Conclusion

While the parameter p_{lth} can in principle improve the accuracy of inference, obtaining a good

model fit under some non-zero p_{lth} value should not be viewed as evidence for a proportion

of p_{lth} mutations segregating in the population in question. This is because it can be shown

that p_{lth} is not equivalent to the true proportion of lethals. When inferring a single DFE for a

group of species or populations, the usage of p_{lth} is also problematic since it modifies the

data, resulting in different SFSs becoming more alike and artificially increases the likelihood

of the model inferred. Thus, comparing the likelihood of two models with different p_{lth}

values cannot be done in a standard way, since these two models will effectively have

different data. We have presented a detailed study of some of the problems with p_{lth} as a

concept which will be useful to anyone modelling the DFE, especially with regards to

avoiding model artifacts.

Data availability

522 The simulation software was implemented in C++ and the full source code is available at

523 https://github.com/r02ap19/DFE_Wright-Fisher01.

Acknowledgements

We thank Adam Eyre-Walker, Brian Charlesworth, Thomas Bataillon, Jane M. Reid, Roslyn

Henry, and Max Tschol for many helpful ideas, comments, and suggestions. We also thank

Nicolas Galtier for reading an earlier draft and providing many useful comments and much

valuable feedback. Anders P. Charmouh was supported by the University of Aberdeen. Greta

Bocedi was supported by a Royal Society University Research Fellowship (UF160614).

Matthew Hartfield is supported by a NERC Independent Research Fellowship

(NE/R015686/1) and a UKRI Frontier Research Guarantee Grant (EP/X027570/1).

References

Agrawal, A. F., and M. C. Whitlock. 2012. Mutation load: The fitness of individuals in

populations where deleterious alleles are abundant. Annu. Rev. Ecol. Evol. Syst.

43:115–135.

Barton, H. J., and K. Zeng. 2018. New methods for inferring the distribution of fitness effects

for INDELs and SNPs. Mol. Biol. Evol. 35:1536–1546.

Barton, N. H., and M. Turelli. 1989. Evolutionary quantitative genetics: How little do we

know? Annu. Rev. Genet. 23:337–370.

Böndel, K. B., S. A. Kraemer, T. Samuels, D. McClean, J. Lachapelle, R. W. Ness, N.

Colegrave, and P. D. Keightley. 2019. Inferring the distribution of fitness effects of

spontaneous mutations in Chlamydomonas reinhardtii. PLoS Biol. 17:1–24.

Booker, T. R. 2020. Inferring parameters of the distribution of fitness effects of new

mutations when beneficial mutations are strongly advantageous and rare. G3 Genes,

Genomes, Genet. 10:2317–2326.

Boyko, A. R., S. H. Williamson, A. R. Indap, J. D. Degenhardt, R. D. Hernandez, K. E.

Lohmueller, M. D. Adams, S. Schmidt, J. J. Sninsky, S. R. Sunyaev, T. J. White, R.

Nielsen, A. G. Clark, and C. D. Bustamante. 2008. Assessing the evolutionary impact of

amino acid mutations in the human genome. PLoS Genet. 4.

Brajesh, R. G., D. Dutta, and S. Saini. 2019. Distribution of fitness effects of mutations
obtained from a simple genetic regulatory network model. Sci. Rep. 9:1–11.

Castellano, D., J. James, and A. Eyre-Walker. 2018. Nearly Neutral Evolution across the
Drosophila melanogaster Genome. Mol. Biol. Evol. 35:2685–2694.

Charlesworth, J., and A. Eyre-Walker. 2006. The rate of adaptive evolution in enteric
bacteria. Mol. Biol. Evol. 23:1348–1356.

Chen, J., S. Glémin, and M. Lascoux. 2020. From drift to draft: How much do beneficial
mutations actually contribute to predictions of Ohta's slightly deleterious model of
molecular evolution? Genetics 214:1005–1018.

Chen, J., S. Glémin, and M. Lascoux. 2017. Genetic diversity and the efficacy of purifying
selection across plant and animal species. Mol. Biol. Evol. 34:1417–1428.

Crow, J. F., and J. M. Simmons. 1983. The Genetics and Biology of Drossophila. Academic
Press, New York.

Crow, J. F., and R. G. Temin. 1964. Evidence for the Partial Dominance of Recessive Lethal
Genes in Natural Populations of Drosophila. Am. Nat. 98:21–33.

Crow, J., and M. Kimura. 1970. An Introduction to Population Genetics Theory. The
Blackburn Press.

Dobzhansky, T., and S. Wright. 1941. Genetics of Natural Populations. V. Relations Between
Mutation Rate and Accumulation of Lethals in Populations of Drosophila
Pseudoobscura. Genetics 26:23–51.

Elyashiv, E., K. Bullaughey, S. Sattath, Y. Rinott, M. Przeworski, and G. Sella. 2010. Shifts
in the intensity of purifying selection: An analysis of genome-wide polymorphism data
from two closely related yeast species. Genome Res. 20:1558–1573.

Evans, S. N., Y. Shvets, and M. Slatkin. 2007. Non-equilibrium theory of the allele frequency

spectrum. Theor. Popul. Biol. 71:109–119.

Eyre-Walker, A. 2002. Changing effective population size and the McDonald-Kreitman test.

Genetics 162:2017–2024.

Eyre-Walker, A., and P. D. Keightley. 2007. The distribution of fitness effects of new

mutations. Nat. Rev. Genet. 8:610–618.

Eyre-Walker, A., M. Woolfit, and T. Phelps. 2006. The distribution of fitness effects of new

deleterious amino acid mutations in humans. Genetics 173:891–900.

Fernández, B., A. García-Dorado, and A. Caballero. 2004. Analysis of the estimators of the

average coefficient of dominance of deleterious mutations. Genetics 168:1053–1069.

Fisher, R. 1930. The Genetical Theory of Natural Selection. Oxford University Press, Oxford.

Fisher, R. A. 1931. XVII.—The Distribution of Gene Ratios for Rare Mutations. Proc. R.

Soc. Edinburgh 50:204–219.

Galtier, N. 2016. Adaptive Protein Evolution in Animals and the Effective Population Size

Hypothesis. PLoS Genet. 12:1–23.

Galtier, N., and M. Rousselle. 2020. How much does Ne vary among species? Genetics

216:559–572.

Goldstein, R. A. 2013. Population size dependence of fitness effect distribution and

substitution rate probed by biophysical model of protein thermostability. Genome Biol.

Evol. 5:1584–1593.

Haag-Liautard, C., M. Dorris, X. Maside, S. Macaskill, D. L. Halligan, B. Charlesworth, and

P. D. Keightley. 2007. Direct estimation of per nucleotide and genomic deleterious

mutation rates in Drosophila. Nature 445:82–85.

Haldane, J. B. S. 1937. The Effect of Variation of Fitness. Am. Nat. 71:337–349. Science

Press.

Halligan, D. L., and P. D. Keightley. 2009. Spontaneous mutation accumulation studies in

evolutionary genetics. Annu. Rev. Ecol. Evol. Syst. 40:151–172.

Hartfield, M., and P. D. Keightley. 2012. Current hypotheses for the evolution of sex and

recombination. Integr. Zool. 7:192–209.

Hedrick, P. W., and A. Garcia-Dorado. 2016. Understanding Inbreeding Depression, Purging,

and Genetic Rescue. Trends Ecol. Evol. 31:940–952. Elsevier Ltd.

Hernandez, R. D. 2008. A flexible forward simulator for populations subject to selection and

demography. Bioinformatics 24:2786–2787.

Huber, C. D., B. Y. Kim, C. D. Marsden, and K. E. Lohmueller. 2017. Determining the

factors driving selective effects of new nonsynonymous mutations. Proc. Natl. Acad.

Sci. U. S. A. 114:4465–4470.

Johri, P., B. Charlesworth, and J. D. Jensen. 2020. Toward an Evolutionarily Appropriate

Null Model⬚: Genetics 215:173–192.

Keightley, P. D., J. L. Campos, T. Booker, and B. Charlesworth. 2016. Inferring the

frequency spectrum of derived variants to quantify adaptive molecular evolution in

protein-coding genes of Drosophila melanogaster. Genetics 203:975–984.

Keightley, P. D., and A. Eyre-Walker. 2007. Joint inference of the distribution of fitness

effects of deleterious mutations and population demography based on nucleotide

polymorphism frequencies. Genetics 177:2251–2261.

Keightley, P. D., and A. Eyre-Walker. 2010. What can we learn about the distribution of

fitness effects of new mutations from DNA sequence data? Philos. Trans. R. Soc. B

Biol. Sci. 365.

Keightley, P. D., and S. P. Otto. 2006. Interference among deleterious mutations favours sex

and recombination in finite populations. Nature 443:89–92.

Kim, B. Y., C. D. Huber, and K. E. Lohmueller. 2017. Inference of the distribution of

selection coefficients for new nonsynonymous mutations using large samples. Genetics

206:345–361.

Kondrashov, A. S. 1985. Deleterious mutations as an evolutionary factor. II. Facultative

apomixis and selfing. Genetics 111:635–653.

Kousathanas, A., and P. D. Keightley. 2013. A comparison of models to infer the distribution

of fitness effects of new mutations. Genetics 193:1197–1208.

Loewe, L., and B. Charlesworth. 2006. Inferring the distribution of mutational effects on

fitness in Drosophila. Biol. Lett. 2:426–430.

Lourenço, J., N. Galtier, and S. Glémin. 2011. Complexity, pleiotropy, and the fitness effect

of mutations. Evolution (N. Y). 65:1559–1571.

Lynch, M., J. Blanchard, D. Houle, T. Kibota, S. Schultz, L. Vassilieva, and J. Willis. 1999.

Perspective: Spontaneous deleterious mutation. Evolution (N. Y). 53:645–663.

Lynch, M., and J. S. Conery. 2003. The Origins of Genome Complexity. Science (80-.).

302:1401–1404.

Lynch, M., and W. G. Hill. 1986. Phenotypic Evolution by Neutral Mutation. Evolution (N.

Y). 40:915.

Martin, G., and T. Lenormand. 2006. a General Multivariate Extension of Fisher'S

Geometrical Model and the Distribution of Mutation Fitness Effects Across Species.

Evolution (N. Y). 60:893.

Mukai, T., S. I. Chigusa, L. E. Mettler, and J. F. Crow. 1972. Mutation rate and dominance of

genes affecting viability in Drosophila melanogaster. Genetics 72:335–355.

Nielsen, R., and Z. Yang. 2003. Estimating the distribution of selection coefficients from

phylogenetic data with applications to mitochondrial and viral DNA. Mol. Biol. Evol.

20:1231–1239.

Ohnishi, O. 1977. Spontaneous And Ethyl Methanesulfonate-Induced Mutations Controlling

Viability In Dosophila Melanogaster II. Homozygous Effect of Polygenic Mutations.

Genetics 87:529–545.

Reich, D. E., and E. S. Lander. 2001. On the allelic spectrum of human disease. Trends

Genet. 17:502–510.

Rice, D. P., B. H. Good, and M. M. Desai. 2015. The evolutionarily stable distribution of

fitness effects. Genetics 200:321–329.

Roze, D., and F. Rousset. 2009. Strong effects of heterosis on the evolution of dispersal rates.

J. Evol. Biol. 22:1221–1233.

Sanjuán, R., A. Moya, and S. F. Elena. 2004. The distribution of fitness effects caused by

single-nucleotide substitutions in an RNA virus. Proc. Natl. Acad. Sci. U. S. A.

101:8396–8401.

Sawyer, S. A., R. J. Kulathinal, C. D. Bustamante, and D. L. Hartl. 2003. Bayesian Analysis

Suggests that Most Amino Acid Replacements in Drosophila Are Driven by Positive

Selection. J. Mol. Evol. 57:154–164.

Schultz, S. T., and M. Lynch. 1997. Mutation and extinction: The role of variable mutational

effects, synergistic epistasis, beneficial mutations, and degree of outcrossing. Evolution

(N. Y). 51:1363–1371.

Silander, O. K., O. Tenaillon, and L. Chao. 2007. Understanding the evolutionary fate of

finite populations: The dynamics of mutational effects. PLoS Biol. 5:922–931.

Simmons, M. J., and J. F. Crow. 1977. Mutations affecting fitness in Drosophila populations.

Annu. Rev. Genet. 11:49–78.

Spigler, R. B., K. Theodorou, and S. M. Chang. 2017. Inbreeding depression and drift load in

small populations at demographic disequilibrium. Evolution (N. Y). 71:81–94.

Tataru, P., and T. Bataillon. 2020. polyDFE: Inferring the Distribution of Fitness Effects and

Properties of Beneficial Mutations from Polymorphism Data. Pp. 125–146 *in* J. Y.

Dutheil, ed. Statistical Population Genomics. Springer US, New York, NY.

Tataru, P., M. Mollion, S. Glémin, and T. Bataillon. 2017. Inference of Distribution of

Fitness Effects and. Genetics 207:1103–1119.

Tenaillon, O. 2014. The utility of Fisher's geometric model in evolutionary genetics. Annu.

Rev. Ecol. Evol. Syst. 45:179–201.

Wade, E. E., C. C. Kyriazis, M. I. A. Cavassim, and K. E. Lohmueller. 2022. Quantifying the

fraction of new mutations that are recessive lethal. bioRxiv 2022.04.22.489225.

Wright, S. 1937. The distribution of gene frequencies in populations. Science (80-.). 85:504.

Wright, S. 1938. The Distribution of Gene Frequencies Under Irreversible Mutation. Proc.

Natl. Acad. Sci. 24:253–259.

Appendix A

Here, we show how to calculate the fraction of mutations with $s > 1$ given some N_e and

some gamma distributed input distribution (Fig. 4). The calculations can in principle be

performed for any input distribution where the probability density function and its parameters

are known. As an example, we will use DelHSD from Tataru et al. (2017): a gamma

distribution with a shape parameter of $\beta = 0.4$ and $S = 10000$.

For any gamma distribution with shape parameter β and scale parameter θ, the probability

density function is

$$f(x) = \frac{1}{(\beta - 1)!\,\theta^{\beta}} x^{\beta - 1} e^{-\frac{x}{\theta}}$$

696 which we can use to find the fraction of mutations with $s > 1$ by integrating f over $[0, 4N_e]$

$$\int_0^{4Ne} f(x)\,dx$$

697 Since $E[f(x)] = \beta\theta$, the probability density function for DelHSD becomes

$$f(x) = \frac{1}{(0.4 - 1)!\left(\dfrac{10000}{0.4}\right)^{0.4}} x^{0.4 - 1} e^{-\frac{x}{\frac{10000}{0.4}}}$$

698 And with $N_e = 500$, we get

$$1 - \int_0^{4Ne} f(x)\,dx = 1 - \int_0^{2000} f(x)\,dx = 1 - 0.4012 \approx 0.5988 = 59.88\%.$$