1

Pioneer factor Pax7 initiates two-step cell-cycle dependent chromatin opening

Arthur Gouhier[1,2], Justine Dumoulin-Gagnon[1], Vincent Lapointe-Roberge[1,2], Aurelio Balsalobre[1]

and Jacques Drouin[1,2,3] *

[1] Laboratoire de génétique moléculaire, Institut de recherches cliniques de Montréal (IRCM)

Montréal, Québec, H2W 1R7 Canada

[2] Department of Biochemistry, McGill University, Montreal, Quebec, Canada.

[3] Département de Biochimie, Université de Montréal, Montréal, Québec, Canada.

Word count: 2618 / 2500

*Corresponding author: Jacques Drouin

16   Laboratoire de génétique moléculaire

17   Institut de recherches cliniques de Montréal (IRCM)

18   110, avenue des Pins Ouest

19   Montréal QC H2W 1R7 Canada

20   Tel.: (514) 987-5579; Fax: (514) 987-5575

21   jacques.drouin@ircm.qc.ca

Summary

Pioneer transcription factors direct cell differentiation by deploying new enhancer repertoires through their unique ability to target and initiate remodelling of closed chromatin. The initial steps of their action remain undefined although pioneers were shown to interact with nucleosomal target DNA and with some chromatin remodelling complexes. We now define the sequence of events that provide pioneers with their unique abilities. Chromatin condensation exerted by linker histone H1 is the first constraint on pioneer recruitment, and this establishes the initial speed of chromatin remodelling. The first step of pioneer action involves recruitment of the LSD1 H3K9me2 demethylase for removal of this repressive mark, as well as recruitment of the MLL complex for deposition of the H3K4me1 mark. Further progression of pioneer action requires passage through cell division, and this involves dissociation of pioneer targets from perinuclear lamins. Only then, the SWI/SNF remodeling complex and the coactivator p300 are recruited, leading to nucleosome displacement and enhancer activation. Thus, the unique features of pioneer actions are those occurring in the lamin-associated compartment of the nucleus. This model is consistent with much prior work that showed a dependence on cell division for establishment of new cell fates.

Introduction

Cell fates are established and maintained through the action of specific combinations of transcription factors, including cell-restricted factors that define unique cell identities. The implementation of new cell fates relies on activation of new enhancer repertoires and this is achieved by pioneer transcription factors through their unique ability to access target sites in closed chromatin and trigger chromatin opening[1]. Many aspects of this general scheme remain undefined, notably pioneer interaction with closed chromatin and the initial events of chromatin remodelling[1].

Investigation of DNA sequences targeted for pioneer action did not show preferential occurrence

of sequence subsets, with some pioneers exhibiting degenerate recruitment sites[2] while others

have stronger recruitment at pioneered sites[3,4]. Closed chromatin access may include the ability

to bind target DNA sequences on nucleosomes, one of the earliest features ascribed to many,

but not all[5], pioneer factors[6]. Structural evidence suggests that pioneer binding may weaken the

interaction of DNA with nucleosomes and hence contribute to initial chromatin alteration[7-9]. This

is compatible with the appearance of so-called "accessible nucleosome conformation" caused by

FOXA recruitment[10] or by subtle chromatin rearrangements that precede frank chromatin opening

following the action of the pioneer Pax7 (REF.[3]). However, nucleosomal organisation cannot on

its own explain barriers to pioneer action, and the data rather suggest that chromatin environment

dictates whether chromatin is permissive or not for pioneer action. For example, the highly

compacted constitutive heterochromatin prevents pioneer access[11,12]. The nature of crucial

events initiating pioneer action remain elusive[1].

It is likely that chromatin remodelling complexes identified for their role in development or in

transcription are also involved in chromatin opening by pioneer factors. For example, the MLL

complex is recruited to FOXA-pioneered sites that are targets of the oestrogen receptor in breast

cancer cells[13]. Another pioneer, Pax7, interacts with the methyltransferase CARM1 and the

MLL1/2 proteins[14]. The pioneers OCT4, GATA3 and ISL1, interact with Brg1, the ATPase of the

SWI/SNF chromatin-remodelling complex[15-17]. But it is presently unclear how and when these

interactions may be relevant to pioneer-dependent chromatin opening.

In the present study, we used an inducible system to define the time course of chromatin opening

at Pax7-pioneered enhancers[3]. Amongst its different roles[18], Pax7 specifies the intermediate

pituitary melanotrope cell fate by opening a repertoire of >2000 enhancers[3,19]. While Pax7 is

necessary to trigger chromatin opening, it also requires cooperation with the nonpioneer

transcription factor Tpit to fully open and activate enhancer chromatin[20]. We now show that Pax7

pioneer action occurs in two steps, a first step that is limited by Pax7 recruitment strength and

inversely correlated with the level of linker histone H1. And a second step that requires passage

through replication and cell division, for recruitment of the SWI/SNF complex, nucleosome

displacement and enhancer activation. This sequential scheme of pioneer action defines a

conceptual framework to further probe and control the pioneering process.

Results

Pax7 primes enhancers for activation

We previously provided an exhaustive analysis of chromatin opening by the pioneer factor Pax7

at two enhancer repertoires, one where Pax7 action results in complete enhancer activation and

another where enhancers only reach the primed state[3]. To study the kinetics of enhancer opening

by Pax7, we used a tamoxifen-inducible ER-Pax7 chimera system (Fig. 1a). The present study

compared subsets of enhancers that are either activated or primed in response to Pax7 with a set

of constitutively active enhancers where Pax7 binding does not alter chromatin organization (Fig.

1b and Supplementary Fig. 1a, b). Having previously observed rapid Pax7 binding to pioneered

sites in closed chromatin but relatively slower remodelling[3], we first assessed whether the primed

state is indeed a transitory state towards complete enhancer activation. The status of pioneered

enhancers was assessed by ChIP-Seq for the chromatin mark histone monomethylated histone

H3 Lys4 (H3K4me1) at different times following activation of ER-Pax7 (Fig. 1c). Thus, pioneered

enhancers are observed first (12h) in the primed state followed by complete activation (Fig. 1c).

The primed state was previously documented as presenting with a weak single peak of H3K4me1,

whereas fully activated enhancers present with a stronger bimodal distribution of H3K4me1 that

reflects nucleosome displacement at the center of pioneered enhancers[1].

Pioneered sites are heterogeneous

ChIP-Seq analyses of pioneered sites show significant heterogeneity with regards to Pax7

recruitment and deposition of chromatin marks (Fig. 1b). To assess the variability in

implementation of Pax7 chromatin opening, we selected a panel of 14 pioneered sites

representative of the Pax7 recruitment distribution (Fig. 1d). We validated that pioneered

enhancers are first remodelled into a primed state by H3K4me1 ChIP-qPCR using two sets of

primers, one centered on the Pax7 binding site and another shifted to one of the H3K4me1

bimodal deposition summit: indeed, H3K4me1 central deposition (average half-time of 24h)

occurs before lateral (average half-time of 32h) accumulation (Fig. 1e,f). Prior data had

suggested interaction between Pax7 and the MLL complex that has H3K4me1 methyltransferase

activity[14]. We therefore assessed MLL complex recruitment at Pax7 pioneered enhancers by

ChIP-qPCR for the MLL3/4 and Ash2 components of the MLL complex: both MLL4 and Ash2 are

recruited with a half-time (average of the 14 sites) of ~12 hours following Pax7 activation (Fig. 1f

and Supplementary Fig. 1c). This half-time corresponds to a time when enhancers are in a primed

state (Fig. 1c) and precedes bulk H3K4me1 deposition (Fig.1f). This delay taken together with the

prior demonstration of a delay between chromatin opening measured by ATAC-Seq and Pax7

recruitment[3] suggests that chromatin opening initiated by Pax7 may involve sequential steps.

Chromatin opening occurs in two steps

To assess the sequence of events leading to chromatin opening, we performed ChIP-qPCR time

courses for various chromatin state markers and chromatin remodellers (Fig. 2a-d). Pax7

recruitment itself has an average half-time of ~12 hours and we found a similar time course for

recruitment of Tpit, a cooperating transcription factor that is required for chromatin opening at the

Pax7 pioneered enhancers[20] (Fig. 2a). Recruitment of these factors parallels MLL4 and Ash2

recruitment: they all precede deposition of H3K4me1 at the center of enhancers (Fig. 2a). We

next assessed chromatin opening by ATAC-qPCR: the half-time of chromatin opening of ~24

hours is similar to H3K4me1 deposition (Fig. 2b). These longer time courses are similar to those

of nucleosome displacement at these enhancers as revealed by ChIP-qPCR for histone H3 and

for recruitment of Brg1 (Fig. 2b). The recruitment of the SWI/SFN complex (Brg1) occurs in parallel

with recruitment of the general coactivator and histone acetylase p300 and of the cohesin complex

protein SMC1 (Fig. 2b). These precede deposition of the active enhancer mark H3K27ac (Fig.

2c). These time courses clearly identify two steps in Pax7-dependent chromatin opening: an initial

step marked by Pax7 recruitment and a delayed second step that involves nucleosome

displacement (Fig. 2a-c and Supplementary Fig. 2a, b).

We previously showed depletion of the repressive histone mark H3K9me2 that is typical of

facultative heterochromatin at the Pax7 pioneered enhancers[3]. We next assessed the time course

of this depletion and observed a half-time of ~12 hours (Fig. 2d). Since facultative heterochromatin

is compacted, we assessed the levels of linker histone H1 at these sites and observed a time

course of H1 depletion similar to that of H3K9me2 depletion (Fig. 2d). The initiation of Pax7

pioneering thus correlates with an initial perturbation of the chromatin environment represented

by the loss of linker histone H1 and the loss of the facultative heterochromatin mark H3K9me2,

together with implementation of the primed state reflected by a weak deposition of H3K4me1 (Fig.

1c).

Pioneering kinetics depend on Pax7 recruitment strength

While the average time response curves defined two steps in the pioneering process, examination

of parameters for the individual 14 pioneer sites revealed heterogeneity in the kinetics of pioneer

action (Fig. 2e). Color-coded individual time course curves for Pax7 recruitment, histone H3

depletion, ATAC signal, H3K4me1 deposition, Brg1 recruitment and H3 depletion indicate that

the stronger Pax7 recruitment sites (Fig. 1d, red) are remodelled quicker than the weaker sites

(blue). The correlation between recruitment strength and half-time of remodelling is observed for

all parameters (Fig. 2f). Similar correlations are observed for all pioneered enhancers genome-

wide (Supplementary Fig. 3a). Computation of time course curves for the five strongest and five

weakest sites indicates an average delay of ~6 hours between the two subgroups (Supplementary

Fig. 3b). These data clearly support the conclusion that recruitment strength determines the onset

of chromatin remodelling at pioneered sites.

To assess whether these correlations relate to long-term or initial levels of Pax7 recruitment, we

performed similar comparisons at 1 hour after Pax7 activation. These revealed that recruitment

strength at 1 and 48 hours are directly correlated (Fig. 2g and Supplementary Fig. 4) and that

initial recruitment is inversely correlated with the time course of chromatin remodelling (Fig. 2h).

We next assessed the impact of initial chromatin organization on the strength of Pax7 recruitment.

Whereas nucleosome content reflected by total histone H3 ChIP levels and the level of the

repressive mark H3K9me2 (Supplementary Fig. 5a) do not correlate with Pax7 recruitment,

chromatin compaction dependent on linker histone H1 determines the initial (1h) strength of Pax7

recruitment (Fig. 2i). Similar observations were made at 48h after Pax7 activation (Supplementary

Fig. 5b) and confirmed genome-wide by UMAP representation of Pax7 recruitment compared to

histone H1 levels (Fig. 2j). Hence, the higher the level of histone H1, the weaker the initial Pax7

recruitment and the longer it takes to remodel these sites (Fig. 2i). In summary, chromatin

condensation reflected by the level of histone H1 constrains Pax7 recruitment and determines the

time course of chromatin remodelling.

Activation of Pax7 primed enhancers requires cell division

The half-time of the second step of the pioneering process (~24 hours) corresponds roughly to

the half-time for AtT-20 cell division. Thus, the second step of the pioneering process may require

DNA replication or passage through cell division. We assessed this by blocking AtT-20 cell

replication for 12 hours with mimosine (Fig. 3a, b). Mimosine-blocked AtT-20 cells are viable and

can re-enter cell cycle upon release (Supplementary Fig. 6a). We assessed each step of the

pioneering process by comparison of mimosine-blocked with normal cycling cells. These

experiments indicate that Pax7 recruitment and H3K9me2 depletion, two parameters of the first

pioneering step, occur independently of cell replication and division (Fig. 3c). In marked contrast,

all marks of the second step of the pioneering process are blocked by mimosine treatment,

including chromatin opening assessed by ATAC-qPCR, Brg1 and p300 recruitment, and

deposition of H3K27ac (Fig. 3d and Supplementary Fig. 6b). Since prior experiments showed that

the first step of the pioneering process yields enhancers in the primed state, we assessed the

pattern of H3K4me1 in mimosine-blocked cells that fail to undergo the second step of the

pioneering process. The ChIP-Seq average profiles of pioneered enhancers in mimosine-blocked

cells remain in the primed state (Fig. 3e) in contrast to normal cycling cells that are in the activated

state at a similar time following Pax7 activation (Fig. 1c).

Cell division is required for dissociation from nuclear lamins

Since the first change observed in parallel with Pax7 recruitment is the depletion of the repressive

mark H3K9me2, we queried a putative mechanism for this depletion. The analysis of proteins

associated with Pax7 in RIME experiments[21] provided two candidate demethylases, KDM1A

(LSD1) and KDM3A (Harris et al. In preparation). Hence, we assessed recruitment of these

demethylases at Pax7 pioneered enhancers and found that KDM1A is recruited with a half-time

corresponding to step 1 of the process (Fig. 4a) whereas KDM3A is recruited at step 2 of the

process (Supplementary Fig. 7a). The early loss of H3K9me2 may be a mechanism to alter the

localization of pioneered sites in the nucleoplasm. Indeed, H3K9me2-marked loci are found in the

nuclear periphery B compartment and their localization is altered by removal of H3K9me2

(REF.[22]). We thus assessed the association of the Pax7 pioneered sites with lamin B1, a marker

of the nuclear B compartment (Fig. 4b) and found that high levels of lamin B1 correlated with high

levels of H3K9me2 at pioneered sites (Fig. 4c). Despite the loss of H3K9me2 at step 1, we

observed dissociation of pioneered sites from lamin B with a half-time corresponding to the cell

cycle-dependent second step (Fig. 4a). If the requirement for cell division observed for step two

of the pioneering process depends on nuclear compartment switching, the blockade of cell

division should prevent dissociation from lamin B, and this is indeed what was observed in

mimosine-blocked cells (Fig. 4d). In this context, we would expect that enhancers that are de

novo primed by Pax7 would not show significant H3K9me2 depletion as is indeed observed (Fig.

4e). We surmise that the slight decrease of H3K9me2 at primed sites is not sufficient for

compartment switching and this is supported by the similar levels of H3K9me2 at sites that are

were already in the primed state before Pax7 activation and that get transcriptionally activated by

Pax7 (Activated enhancers, Fig. 4e). Further, we observe recruitment of the SWI/SNF complex

only after full activation but not with the primed state as revealed by Brg1 ChIP-Seq (Fig. 4e), in

agreement with restricted nuclear localisation of Brg1 in the central nucleoplasm[23]. We found

similar features for recruitment of RNA Pol II and components (MED1 and MED12) of the Mediator

complex (Supplementary Fig. 7b). Why do some enhancers fail to complete the process of

activation? A failure to recruit an essential cooperating transcription factor could be responsible

and indeed, we observe no recruitment of Tpit, a nonpionner factor required for full activation of

melanotrope enhancers[20], at de novo primed sites (Fig. 4e).

In conclusion, and taken collectively, the present data suggest a model for pioneer factor action

(Fig. 4f, g) in which the first steps are compatible with actions within the perinuclear compartment

B: these include removal of repressor marks together with deposition of activation marks

corresponding to the primed state. The ensuing steps for enhancer opening require passage

through cell division that allows for lamin dissociation, and presumably switching to the A

compartment, where the loci become accessible to the SWI/SNF complex for nucleosome

displacement and transcriptional activation.

Discussion

The present work shows that chromatin compaction elicited by histone H1 determines the strength

of pioneer Pax7 recruitment and speed of its actions. Beyond this limiting step, the different

parameters of chromatin opening follow a similar time course while the next limiting step is

imposed by cell division and its importance for lamin, and presumably compartment B,

dissociation (Fig. 4g). The association of SWI/SNF (Brg1) recruitment and nucleosome

displacement is consistent with the nuclear sublocalization of BRG1 and Brm in compartment A,

but not in the perinuclear compartment B, of the nucleoplasm[23].

As the ability of loci to switch from compartments B to A depends on the level of H3K9me2 (REF.[22]

[24]), it follows that enhancers that are primed (but not activated) by Pax7 do not show significant

H3K9me2 depletion (Fig. 4e). In contrast, these primed enhancers have deposition of H3K4me1

presenting as a single weak peak (Fig. 4f). Consistent with an early recruitment of the MLL

complex (Fig. 2a), the data indicate that targeted enhancers can be primed in the lamin-

associated compartment including deposition of H3K4me1. The limiting factor for complete

enhancer activation is thus controlled by demethylation of H3K9me2 and this appears to require

the cooperating nonpioneer Tpit (Fig. 4e). The variable effectiveness of Pax7 to open chromatin

at different sites may thus be related to chromatin constraints such as dictated by histone H1,

and/or to the relative affinity for DNA binding sites and/or to nonpioneer interaction. In models that

may involve different cooperating nonpioneers, a single pioneer factor may thus implement

various subsets of primed enhancers ready for activation in response to an array of developmental

or signaling cues mediated through the activities of different nonpioneers.

References

1   Balsalobre, A. & Drouin, J. Pioneer factors as master regulators of the epigenome and cell
fate. *Nat Rev Mol Cell Biol* **23**, 449-464 (2022). https://doi.org:10.1038/s41580-022-
00464-z
2   Soufi, A. *et al.* Pioneer transcription factors target partial DNA motifs on nucleosomes to
initiate   reprogramming.   *Cell*   **161**,   555-568   (2015).
https://doi.org:10.1016/j.cell.2015.03.017
3   Mayran, A. *et al.* Pioneer factor Pax7 deploys a stable enhancer repertoire for specification
of cell fate. *Nat Genet* **50**, 259–269 (2018). https://doi.org:10.1038/s41588-017-0035-2
4   Pelletier, A., Mayran, A., Omichinski, J. & Drouin, J. Pax7 Pioneer action requires both
paired and homeo DNA binding domains. *Nucleic Acids Research*, 1–13 (2021).
https://doi.org:https://doi.org/10.1093/nar/gkab561
5   Minderjahn, J. *et al.* Mechanisms governing the pioneering and redistribution capabilities
of   the   non-classical   pioneer   PU.1.   *Nat   Commun*   **11**,   402   (2020).
https://doi.org:10.1038/s41467-019-13960-2
6   Cirillo, L. A. *et al.* Binding of the winged-helix transcription factor HNF3 to a linker histone
site   on   the   nucleosome.   *EMBO   J*   **17**,   244-254   (1998).
https://doi.org:10.1093/emboj/17.1.244
7   Dodonova, S. O., Zhu, F., Dienemann, C., Taipale, J. & Cramer, P. Nucleosome-bound
SOX2 and SOX11 structures elucidate pioneer factor function. *Nature* **580**, 669-672
(2020). https://doi.org:10.1038/s41586-020-2195-y
8   Michael, A. K. *et al.* Mechanisms of OCT4-SOX2 motif readout on nucleosomes. *Science*
368, 1460-1465 (2020). https://doi.org:10.1126/science.abb0074
9   Tanaka, H. *et al.* Interaction of the pioneer transcription factor GATA3 with nucleosomes.
Nat Commun **11**, 4136 (2020). https://doi.org:10.1038/s41467-020-17959-y
10   Iwafuchi-Doi, M. *et al.* The Pioneer Transcription Factor FoxA Maintains an Accessible
Nucleosome Configuration at Enhancers for Tissue-Specific Gene Activation. *Mol Cell* **62**,
79-91 (2016). https://doi.org:10.1016/j.molcel.2016.03.001
11   Soufi, A., Donahue, G. & Zaret, K. S. Facilitators and impediments of the pluripotency
reprogramming factors' initial engagement with the genome. *Cell* **151**, 994-1004 (2012).
12   Nicetto, D. *et al.* H3K9me3-heterochromatin loss at protein-coding genes enables
developmental   lineage   specification.   *Science*   **363**,   294-297   (2019).
https://doi.org:10.1126/science.aau0583
13   Jozwik, K. M., Chernukhin, I., Serandour, A. A., Nagarajan, S. & Carroll, J. S. FOXA1
Directs H3K4 Monomethylation at Enhancers via Recruitment of the Methyltransferase
MLL3. *Cell Rep* **17**, 2715-2723 (2016). https://doi.org:10.1016/j.celrep.2016.11.028
14   Kawabe, Y., Wang, Y. X., McKinnell, I. W., Bedford, M. T. & Rudnicki, M. A. Carm1
regulates Pax7 transcriptional activity through MLL1/2 recruitment during asymmetric
satellite   stem   cell   divisions.   *Cell   Stem   Cell*   **11**,   333-345   (2012).
https://doi.org:10.1016/j.stem.2012.07.001
15   King, H. W. & Klose, R. J. The pioneer factor OCT4 requires the chromatin remodeller
BRG1 to support gene regulatory element function in mouse embryonic stem cells. *Elife*
6 (2017). https://doi.org:10.7554/eLife.22631
16   Takaku, M. *et al.* GATA3-dependent cellular reprogramming requires activation-domain
dependent   recruitment   of   a   chromatin   remodeler.   *Genome Biol* **17**, 36 (2016).
https://doi.org:10.1186/s13059-016-0897-0

17 Gao, R. *et al.* Pioneering function of Isl1 in the epigenetic control of cardiomyocyte cell
fate. *Cell Res* **29**, 486-501 (2019). https://doi.org:10.1038/s41422-019-0168-1
18 Mayran, A., Pelletier, A. & Drouin, J. Pax factors in transcription and epigenetic
remodelling.   *Semin   Cell   Dev   Biol*   **44**,   135-144   (2015).
https://doi.org:10.1016/j.semcdb.2015.07.007
19 Budry, L. *et al.* The selector gene Pax7 dictates alternate pituitary cell fates through its
pioneer action on chromatin remodeling. *Genes & Development* **26**, 2299-2310 (2012).
https://doi.org:10.1101/gad.200436.112
20 Mayran, A. *et al.* Pioneer and nonpioneer factor cooperation drives lineage specific
chromatin   opening.   *Nature   Communications*   **10**,   3807   (2019).
https://doi.org:10.1038/s41467-019-11791-9
21 Mohammed, H. *et al.* Rapid immunoprecipitation mass spectrometry of endogenous
proteins (RIME) for analysis of chromatin complexes. *Nat Protoc* **11**, 316-326 (2016).
https://doi.org:10.1038/nprot.2016.020
22 Poleshko, A. *et al.* H3K9me2 orchestrates inheritance of spatial positioning of peripheral
heterochromatin through mitosis. *Elife* **8** (2019). https://doi.org:10.7554/eLife.49278
23 Reyes, J. C., Muchardt, C. & Yaniv, M. Components of the human SWI/SNF complex are
enriched in active chromatin and are associated with the nuclear matrix. *J Cell Biol* **137**,
263-274 (1997). https://doi.org:10.1083/jcb.137.2.263
24 Smith, C. L., Poleshko, A. & Epstein, J. A. The nuclear periphery is a scaffold for tissue-
specific   enhancers.   *Nucleic   Acids   Res*   **49**,   6181-6195   (2021).
https://doi.org:10.1093/nar/gkab392
25 Langlais, D., Couture, C. & Drouin, J. The Stat3/GR interaction code: predictive value of
direct/indirect DNA recruitment for transcription outcome. *Molecular Cell* **47**, 38-49 (2012).
311

ACKNOWLEDGMENTS

We are very grateful to Dr. Ali Shilatifard for the MLL3 and MLL4 antibodies, to Odile Neyret and Sarah Boisset for NGS analyses, and to Valérie Magoon for her expert secretarial assistance. Data analyses were possible thanks to the support of Compute Canada. This work was supported by Foundation grant FDN-154297 to J.D. from the Canadian Institutes of Health Research.

AUTHOR CONTRIBUTIONS

A.G. and J.D. conceived the study, A.G., A.B. and J.D. conceived and designed the experiments, A.G., J.D.-G. and V.L.-R. performed ChIP experiments, A.G. and J.D. wrote the manuscript.

COMPETING INTERESTS

The authors declare no competing financial interests.

Legends to Figures

Fig. 1 | Pax7 primes enhancers for activation. a) The role of Pax7 in specification of the

intermediate pituitary melanotrope cell fate is illustrated in contrast to the corticotrope cell fate.

These two hormone-secreting lineages express the same hormone precursor gene,

proopiomelanocortin (POMC), and their fates are determined by the Tbox transcription factor Tpit.

The mouse AtT-20 cells are a model of the corticotrope lineage and Pax7 expression reprograms

these cells towards a melanotrope-like fate. **b**) Heat map representation of ChIP-Seq and ATAC-

Seq data for three subsets of Pax7-targeted enhancers. Data are shown before (0h) and after

48h induction of Pax7 activity in AtT-20 cells harbouring a tamoxifen-inducible ER-Pax7 chimera[3].

Pioneered enhancers have no marks before Pax7 activation and present with hallmarks of

transcriptionally active enhancers after, namely appearance of an ATAC-Seq signal together with

a bimodal distribution of H3K4me1 deposition. Pax7 primed enhancers also show appearance of

weaker ATAC-Seq and H3K4me1 signals, the latter presenting as a single weak peak. Finally, a

set of constitutively active enhancers serve as control for comparisons. These enhancers bind

Pax7, but this binding is not associated with any change in chromatin signature. These enhancer

subsets were previously described[3,4]. **c**) Average profiles of H3K4me1 deposition at subsets of

Pax7-remodelled enhancers at indicated times following tamoxifen (Tam) activation of ER-Pax7.

d) Selection of 14 Pax7-pioneered sites used for detailed investigation by qPCR and color-coded

from red to blue according to strength of Pax7 recruitment. Pax7 and H3K4me1 ChIP-Seq, and

ATAC-Seq, profiles are shown for these individual peaks at 48h following Pax7 activation. **e**) Time

course of H3K4me1 deposition determined by ChIP-qPCR at the 14 pioneered sites illustrated

above. For each site, a red curve illustrates H3K4me1 accumulation under the Pax7 peak

(centered), whereas a blue curve represents H3K4me1 deposition over a lateral peak of

deposition (shifted) that characterizes the bimodal H3K4me1 distribution of fully active enhancers.

f) Average (of 14 pioneered sites) time course of H3K4me1 accumulation (centered and shifted)

is compared with ChIP-qPCR data for recruitment of two protein components of the MLL

remodelling complex, MLL4 and Ash2. Data represent the means ± sem of duplicate biological

samples.

Fig. 2 | Pax7-dependent chromatin opening occurs in two steps. a) Average (14 pioneered

sites of Fig.1d represented as means ± sem of duplicate biological samples, $n \geq 2$) profiles of

ChIP-qPCR data for Pax7, Tpit, MLL4 and Ash2 that are recruited with a similar time course (T ½

~ 12 hrs) compared to accumulation of H3K4me1 (center). **b**) Compared to Pax7 recruitment,

ATAC-Seq signals and recruitment of Brg1, p300 and SMC1, together with depletion of Histone

H3 occur with an average T ½ of ~ 24h ($n \geq 2$). **c**) Deposition of activation mark H3K27ac and

bimodal H3K4me1 (shifted) are slightly delayed compared to ATAC signal ($n \geq 2$). **d**) Depletion of

the repressive mark H3K9me2 and of linker Histone H1 occur with a T ½ of ~ 12h ($n \geq 2$). **e**)

Individual site ChIP-qPCR time course data for recruitment of Pax7, H3K4me1 and Brg1, together

with time courses of ATAC signals and depletion histones H1 and H3 shown for the 14 pioneered

sites color-coded (in Fig.1d) from red to blue to reflect the strength of Pax7 recruitment. **f**)

Correlations between Pax7 recruitment strength and various markers depicted in (e). **g**)

Correlation between rapid (1h) and long-term (48h) Pax7 recruitment strength. **h**) Inverse

correlation between initial (1h) Pax7 recruitment strength and the average half-times of all

markers of chromatin remodelling at the 14 pioneered sites. **i**) Correlations between the initial

levels of histone H1 at the 14 pioneered sites and the strength of Pax7 recruitment and with the

half-time of chromatin remodelling markers. **j**) UMAP representation of Pax7 recruitment strengths

at the indicated times after Tam addition compared to initial histone H1 levels measured by ChIP-

Seq at pioneered enhancers depicted in Fig. 1b.

Fig. 3 | Activation of Pax7 pioneered enhancers requires cell division. a) Experimental

design for assessment of chromatin remodelling in cells with mimosine-blocked cell cycle

compared to normally cycling cells. **b**) FACS profiles of normal (no mimosine) cells compared to

cells incubated for 12h with mimosine that exhibit cell-cycle arrest in G1. Representative of n=2

is shown. **c**) Both recruitment of Pax7 and depletion of H3K9me2 measured by ChIP-qPCR at the

14 pioneered sites are not affected by blocking AtT-20 cell cycle with mimosine (n=2). **d**) In

contrast, chromatin opening (ATAC) recruitments of Brg1, p300 and H3K27ac measured by ChIP-

qPCR are all severely curtailed in mimosine-blocked cells (n=2). **e**) Average profiles of H3K4me1

accumulation at Pax7-remodelled enhancer subsets measured by ChIP-Seq at different times

after Tam activation of ER-Pax7.

Fig. 4 | Pax7-dependent chromatin remodelling involves H3K9me2 demethylation and

dissociation from lamins. a) Average (14 pioneered sites of Fig.1d represented as means ±

sem of duplicate biological samples) time courses for depletion of H3K9me2 and dissociation

from lamin B1 measured by ChIP-qPCR compared to the time courses of recruitment for the

H3K9me2 demethylase, KDM1A (LSD1). **b**) Pax7 pioneered sites have high levels of H3K9me2

and lamin B1 before Pax7 action. **c**) Correlation between initial levels of H3K9me2 and lamin B1

at 14 pioneered sites before Pax7 action (n $\geq$ 2). **d**) Lamin depletion is prevented by cell cycle

blockade with mimosine (n=2). **e**) Heatmaps of ChIP-Seq profiles for Pax7, H3K4me1, H3K9me2,

Tpit and Brg1 at subsets of Pax7 remodelled enhancers. Data for other markers presented in

Supplementary Fig. 7. **f**) Schematic representation of proteins shown in the present work to be

associated with either primed or active enhancer states. **g**) Sequential model of Pax7 pioneer

action. Pax7 pioneer targets are enriched in H3K9me2 and associated with lamin B1, which is

typical of chromatin that is present in the perinuclear B compartment. Step 1 of the pioneering

process involves recruitment of the H3K9me2 demethylase KDM1A (LSD1), together with the

MLL complex. These lead to demethylation of H3K9me2 and H3K4me1 deposition, respectively.

Cell cycle blockade with mimosine prevents Pax7-dependent remodelling from proceeding

further. At mitosis, masking of the H3K9me2 mark by H3S10 phosphorylation accompanies

dissociation of A and B compartments of the nucleus[22]. Following mitosis and nuclear reassembly,

Pax7-pioneered targets that have undergone H3K9me2 demethylation will remain in the central

nuclear A compartment where the Brg1 and Brm ATPase's of the SWI/SNF complex are

localised[23] and recruited for nucleosome displacement and complete enhancer activation. Pax7

primed enhancers do not recruit the cooperating nonpioneer factor Tpit and hence, the

cooperation between Tpit and Pax7 appears required for significant H3K9me2 demethylation and

continuation of the Pax7-dependent remodelling process to the cell cycle-dependent step 2.

Fig. 1 | Pax7 primes enhancers for activation.

Fig. 2 | Pax7-dependent chromatin opening occurs in two steps.

Fig. 3 | Activation of Pax7 pioneered enhancers requires cell division.

Fig. 4 | Pax7-dependent chromatin remodelling involves H3K9me2 demethylation and dissociation from lamins.