1

Pioneer factor Pax7 initiates two-step cell-cycle dependent chromatin opening

Arthur Gouhier[1,2], Justine Dumoulin-Gagnon[1], Vincent Lapointe-Roberge[1,2], Juliette Harris[1],

Aurelio Balsalobre[1] and Jacques Drouin[1,2,3] *

[1] Laboratoire de génétique moléculaire, Institut de recherches cliniques de Montréal (IRCM)

Montréal, Québec, H2W 1R7 Canada

[2] Department of Biochemistry, McGill University, Montreal, Quebec, Canada.

[3] Département de Biochimie, Université de Montréal, Montréal, Québec, Canada.

Word count: 3915

*Corresponding author: Jacques Drouin
16   Laboratoire de génétique moléculaire
17   Institut de recherches cliniques de Montréal (IRCM)
18   110, avenue des Pins Ouest
19   Montréal QC H2W 1R7 Canada
20   Tel.: (514) 987-5579; Fax: (514) 987-5575
21   jacques.drouin@ircm.qc.ca

Summary

Pioneer transcription factors direct cell differentiation by deploying new enhancer repertoires through their unique ability to target and initiate remodelling of closed chromatin. The initial steps of their action remain undefined although pioneers were shown to interact with nucleosomal target DNA and with some chromatin remodelling complexes. We now define the sequence of events that enable the pioneer Pax7 with its unique abilities. Chromatin condensation exerted by linker histone H1 is the first constraint on Pax7 recruitment, and this establishes the initial speed of chromatin remodelling. The first step of pioneer action involves recruitment of the KDM1A (LSD1) H3K9me2 demethylase for removal of this repressive mark, as well as recruitment of the MLL complex for deposition of the activating H3K4me1 mark. Further progression of pioneer action requires passage through cell division, and this involves dissociation of pioneer targets from perinuclear lamin B. Only then, the SWI/SNF remodeling complex and the coactivator p300 are recruited, leading to nucleosome displacement and enhancer activation. Thus, the unique features of pioneer actions are those occurring in the lamin-associated compartment of the nucleus. This model is consistent with prior work that showed a dependence on cell division for establishment of new cell fates.

Introduction

Cell fates are established and maintained through the action of specific combinations of transcription factors, including cell-restricted factors that define unique cell identities. The implementation of new cell fates relies on activation of new enhancer repertoires and this is achieved by pioneer transcription factors through their unique ability to access target sites in closed chromatin and trigger chromatin opening[1]. Many aspects of this general scheme remain

undefined, notably pioneer interaction with closed chromatin and the initial events of chromatin

remodelling[1].

Investigation of DNA sequences targeted for pioneer action did not show preferential occurrence

of sequence subsets, with some pioneers exhibiting degenerate recruitment sites[2] while others

have stronger recruitment at pioneered sites[3,4]. Closed chromatin access may include the ability

to bind target DNA sequences on nucleosomes, one of the earliest features ascribed to many,

but not all[5], pioneer factors[6]. Structural evidence suggests that pioneer binding may weaken the

interaction of DNA with nucleosomes and hence contribute to initial chromatin alteration[7-9]. This

is compatible with the appearance of so-called "accessible nucleosome conformation" caused by

FOXA recruitment[10] or by subtle chromatin rearrangements that precede frank chromatin opening

following the action of the pioneer Pax7 (REF.[3]). However, nucleosomal organisation cannot on

its own explain barriers to pioneer action, and the data rather suggest that chromatin environment

dictates whether chromatin is permissive or not for pioneer action. For example, the highly

compacted constitutive heterochromatin prevents pioneer access[11,12]. The nature of crucial

events initiating pioneer action remain elusive[1].

It is likely that chromatin remodelling complexes identified for their role in development or in

transcription are also involved in chromatin opening by pioneer factors. For example, the MLL

complex is recruited to FOXA-pioneered sites that are targets of the oestrogen receptor in breast

cancer cells[13]. The pioneer, Pax7, interacts with the methyltransferase CARM1 and the MLL1/2

proteins[14]. The pioneers OCT4, GATA3 and ISL1, interact with Brg1, the ATPase of the SWI/SNF

chromatin-remodelling complex[15-17]. But it is presently unclear how and when these interactions

may be relevant to pioneer-dependent chromatin opening.

In the present study, we used an inducible system to define the time course of chromatin opening

at Pax7-pioneered enhancers[3]. Amongst its different roles[18], Pax7 specifies the intermediate

pituitary melanotrope cell fate by opening a repertoire of >2000 enhancers[3,19]. While Pax7 is

necessary to trigger chromatin opening, it also requires cooperation with the nonpioneer

transcription factor Tpit to fully open chromatin and activate enhancers [20]. We now show that Pax7

pioneer action occurs in two steps, a first step that is limited by Pax7 recruitment strength and

inversely correlated with the level of linker histone H1, and a second step that requires passage

through replication and cell division for dissociation of large domains from lamin B, recruitment of

the SWI/SNF complex, nucleosome displacement and enhancer activation. This sequential

scheme of pioneer action defines a conceptual framework to further probe and control the

pioneering process.

Results

Pax7 primes enhancers for activation

We previously provided an exhaustive analysis of chromatin opening by the pioneer factor Pax7

at two enhancer repertoires, one where Pax7 action results in complete enhancer activation and

another where enhancers only reach the primed state[3]. To study the kinetics of enhancer opening

by Pax7, we used a tamoxifen-inducible ER-Pax7 chimera system (Fig. 1a) that relies on

expression of ER-Pax7 in AtT-20 cells at levels that are similar to those of mouse pituitary

melanotrope cells (Extended Data Fig. 1a). This re-programs AtT-20 cells into melanotrope-like

cells[3,19,20]. The present study compared subsets of enhancers that are either activated or primed

in response to Pax7 with a set of constitutively active enhancers where Pax7 binding does not

alter chromatin organization (Fig. 1b and Extended Data Fig. 1b, c). Having previously observed

rapid Pax7 binding to pioneered sites in closed chromatin but relatively slower remodelling[3], we

first assessed whether the primed state is indeed a transitory state towards complete enhancer

activation. The status of pioneered enhancers was assessed by ChIP-Seq for the activating

chromatin mark histone monomethylated histone H3 Lys4 (H3K4me1) at different times following

activation of ER-Pax7 (Fig. 1c). Thus, pioneered enhancers are observed first (12h) in the primed

state followed by complete activation (Fig. 1c). The primed state was previously documented as

presenting with a weak single peak of H3K4me1, whereas fully activated enhancers present with

a stronger bimodal distribution of H3K4me1 that reflects nucleosome displacement at the center

of pioneered enhancers[1].

Pioneered sites are heterogeneous

ChIP-Seq analyses of pioneered sites show significant heterogeneity with regards to Pax7

recruitment and deposition of chromatin marks (Fig. 1b). To assess the variability in

implementation of Pax7 chromatin opening, we selected a panel of 14 pioneered sites chosen

randomly and representative of the Pax7 recruitment distribution (Fig. 1d and Extended Data Fig.

1d). We validated that pioneered enhancers are first remodelled into a primed state by H3K4me1

ChIP-qPCR using two sets of primers, one centered on the Pax7 binding site and another shifted

to one of the H3K4me1 bimodal deposition summit: indeed, H3K4me1 central deposition (average

half-time of 24h) occurs before lateral (average half-time of 32h) accumulation (Fig. 1e,f and

Extended Data Fig. 1e). Prior data had suggested interaction between Pax7 and the MLL

complex that has H3K4me1 methyltransferase activity[14]. We therefore assessed MLL complex

recruitment at Pax7 pioneered enhancers by ChIP-qPCR for the MLL3, MLL4 and Ash2

components of the MLL complex: both MLL4 and Ash2 are recruited with a half-time (average of

the 14 sites) of ~12 hours following Pax7 activation (Fig. 1g and Extended Data Fig. 2). This half-

time corresponds to a time when enhancers are in a primed state (Fig. 1c) and precedes bulk

H3K4me1 deposition (Fig.1f). This delay taken together with the prior demonstration of a delay

between chromatin opening measured by ATAC-Seq and Pax7 recruitment[3] suggests that

chromatin opening initiated by Pax7 may involve sequential steps.

Chromatin opening occurs in two steps

To assess the sequence of events leading to chromatin opening, we performed ChIP-qPCR time courses for various chromatin state markers and chromatin remodellers (Fig. 2a-d). Pax7 recruitment itself has an average half-time of ~12 hours and we found a similar time course for recruitment of the cooperating transcription factor Tpit (Fig. 2a). Recruitment of these factors parallels MLL4 and Ash2 recruitment, and they all precede deposition of H3K4me1 at the center of enhancers (Fig. 2a). We next assessed chromatin opening by ATAC-qPCR: the half-time of chromatin opening of ~24 hours is similar to H3K4me1 deposition (Fig. 2b). These longer time courses are similar to those of nucleosome displacement at these enhancers as revealed by ChIP-qPCR for histone H3 and for recruitment of Brg1 (Fig. 2b). The recruitment of the SWI/SNF complex (Brg1) occurs in parallel with recruitment of the general coactivator and histone acetylase p300 and of the cohesin complex protein SMC1 (Fig. 2b). These precede deposition of the active enhancer mark histone H3 acetylated Lys27, H3K27ac (Fig. 2c). These time courses clearly identify two steps in Pax7-dependent chromatin opening (Fig. 2a-c): an initial step marked by Pax7 recruitment and a delayed second step that involves nucleosome displacement (Fig. 2e, f).

We previously showed depletion of the repressive histone mark histone H3 dimethylated Lys9 (H3K9me2) that is typical of facultative heterochromatin at the Pax7 pioneered enhancers[3]. We assessed the time course of this depletion and observed a half-time of ~12 hours (Fig. 2d). Since linker histone H1 contributes to chromatin compaction, we measured its levels at the pioneered sites and observed a time course of H1 depletion similar to that of H3K9me2 (Fig. 2d). The initiation of Pax7 pioneering thus correlates with an initial perturbation of the chromatin environment represented by the loss of linker histone H1 and the loss of the repressive mark H3K9me2, together with implementation of the primed state reflected by a weak deposition of H3K4me1 (Fig. 1c).

Pioneering kinetics depend on Pax7 recruitment strength

While the average time response curves defined two steps in the pioneering process, examination of parameters for the individual 14 pioneer sites revealed heterogeneity in the kinetics of pioneer action (Fig. 3a). Color-coded individual time course curves for Pax7 recruitment, histone H1 depletion, ATAC signal, H3K4me1 deposition, Brg1 recruitment and H3 depletion indicate that the stronger Pax7 recruitment sites (Fig. 1d, red) are remodelled quicker than the weaker sites (blue). The correlation between recruitment strength and half-time of remodelling is observed for all parameters (Fig. 3b). Similar correlations are observed for all pioneered enhancers genome-wide (Fig. 3c and Extended Data Fig. 3c). Computation of time course curves for the five strongest and five weakest sites indicates an average delay of ~6 hours between the two subgroups (Extended Data Fig. 3a, b). These data clearly support the conclusion that recruitment strength determines the onset of chromatin remodelling at pioneered sites.

To assess whether these correlations relate to long-term or initial levels of Pax7 recruitment, we performed similar comparisons at 1 hour after Pax7 activation. These revealed that recruitment strength at 1 and 48 hours are directly correlated (Fig. 3d and Extended Data Fig. 4) and that initial recruitment is inversely correlated with the time course of chromatin remodelling (Fig. 3e).

We next assessed the impact on the strength of Pax7 recruitment of initial chromatin compaction as reflected by the levels of linker histone H1 that are higher at pioneered compared to constitutive sites (Fig. 3f). Whereas nucleosome content reflected by total histone H3 ChIP levels and the level of the repressive mark H3K9me2 (Extended Data Fig. 3d) do not correlate with Pax7 recruitment, chromatin compaction dependent on linker histone H1 inversely correlates with the initial (1h) strength of Pax7 recruitment (Fig. 3g). Similar observations were made at 48h after Pax7 activation (Extended Data Fig. 3c) and confirmed genome-wide by UMAP representation of

Pax7 recruitment compared to histone H1 levels (Fig. 3h). Hence, the higher the level of histone

H1, the weaker the initial Pax7 recruitment and the longer it takes to remodel these sites (Fig.

3h). It is noteworthy that long-term Pax7 recruitment and chromatin opening are not directly

correlated with initial recruitment (Extended Data Fig. 4). In summary, chromatin condensation

reflected by the level of histone H1 appear to constrain initial Pax7 recruitment and determine the

time course of chromatin remodelling without affecting final pioneered enhancer activation.

171 Activation of Pax7 primed enhancers requires cell division

The half-time of the second step of the pioneering process (~24 hours) corresponds roughly to

the half-time for AtT-20 cell division. Thus, the second step of the pioneering process may require

DNA replication or passage through cell division. We assessed this by blocking AtT-20 cell

replication for 12 hours with mimosine (Fig. 4a, b). Mimosine-blocked AtT-20 cells are viable and

can re-enter cell cycle upon release (Extended Data Fig. 5a). We assessed each step of the

pioneering process by comparison of mimosine-blocked with normal cycling cells. These

experiments indicate that Pax7 and Tpit recruitment, as well as H3K9me2 depletion, three

parameters of the first pioneering step, occur independently of cell replication and division (Fig.

4c). In marked contrast, all marks of the second step of the pioneering process are blocked by

mimosine treatment, including chromatin opening assessed by ATAC-qPCR, Brg1 and p300

recruitment, and deposition of H3K27ac (Fig. 4c and Extended Data Fig. 6). The mimosine

treatment did not affect the levels of Brg1 and p300 but led to a ~2-fold decrease of Tpit (Extended

Data Fig. 5b); however, this did not change Tpit recruitment in the first phase of Pax7 action (Fig.

4c). Since prior experiments showed that the first step of the pioneering process yields enhancers

in the primed state, we assessed the pattern of H3K4me1 in mimosine-blocked cells that fail to

undergo the second step of pioneering. The ChIP-Seq average profiles of pioneered enhancers

in mimosine-blocked cells show that they remain in the primed state (Fig. 4d) in contrast to normal

cycling cells in which they are in the activated state at a similar time following Pax7 activation

(Fig. 1c). Also, mimosine-blocked cells only show minimal gains of ATAC-Seq signals (Fig. 4e)

consistent with their primed status.

Cell division is required for dissociation from nuclear lamins

Since the first change observed in parallel with Pax7 recruitment is the depletion of the repressive

mark H3K9me2, we queried a putative mechanism for this depletion. The analysis of proteins

associated with Pax7 in RIME experiments[21] provided two candidate demethylases, KDM1A

(LSD1) and KDM3A (Harris et al. In preparation). Hence, we assessed recruitment of these

demethylases at Pax7 pioneered enhancers and found that KDM1A is recruited with a half-time

corresponding to step 1 of the process (Fig. 5a) whereas KDM3A is recruited at step 2 (Extended

Data Fig. 7a). The early loss of H3K9me2 may be a mechanism to alter the localization of

pioneered sites in the nucleoplasm. Indeed, H3K9me2-marked loci are found in the nuclear

periphery B compartment and their localization is altered by removal of H3K9me2 (REF.[22,23]). We

thus assessed the association of the Pax7 pioneered sites with lamin B1, a marker of the nuclear

B compartment (Fig. 5b) and found that high levels of lamin B1 correlated with high levels of

H3K9me2 at pioneered sites (Fig. 5c, d). Despite the loss of H3K9me2 at step 1, we observed

dissociation of pioneered sites from lamin B with a half-time corresponding to the cell cycle-

dependent second step (Fig. 5a). If the requirement for cell division observed for step 2 of the

pioneering process depends on nuclear compartment switching, the blockade of cell division

should prevent dissociation from lamin B, and this is indeed what was observed in mimosine-

blocked cells (Fig. 5e). In this context, we would expect that enhancers that are de novo primed

by Pax7 would not show significant H3K9me2 depletion as is observed (Fig. 5f). We surmise that

the slight decrease of H3K9me2 at primed sites is not sufficient for lamin B dissociation and this

is supported by the similar levels of H3K9me2 at sites that were already in the primed state before

Pax7 activation and that get transcriptionally activated by Pax7 (Activated enhancers, Fig. 5f).

Further, the recruitment of the SWI/SNF complex is observed only after full activation but not with

the primed state as revealed by Brg1 ChIP-Seq (Fig. 5f), in agreement with restricted nuclear

localisation of Brg1 in the central nucleoplasm[24]. We found similar features for recruitment of RNA

Pol II and components (MED1 and MED12) of the Mediator complex (Extended Data Fig. 7b).

Why do some enhancers fail to complete the process of activation? A failure to recruit an essential

cooperating transcription factor could be responsible and indeed, we observe no recruitment of

Tpit, a nonpionner factor required for full activation of melanotrope enhancers[20], at de novo primed

sites (Fig. 5f).

In summary, the first step of pioneer action involves two biochemical alterations of critical histone

marks: deposition of H3K4me1 presumably implemented by the MLL complex and removal of the

repressive mark H3K9me2 by KDM1A. Both alterations occur independently of cell replication

and division (Fig. 4c, d) but primed enhancers are characterized by the presence of the activation

mark H3K4me1 without significant depletion of the repressive H3K9me2. We used an inhibitor of

KDM1A to test whether its enzyme activity is essential for H3K9me2 demethylation and

downstream chromatin changes. Whereas treatment with the KDM1A inhibitor OG-L002 does not

prevent Pax7 recruitment (Fig. 5g), it decreases H3K9me2 demethylation (Fig. 5h), lamin B

dissociation (Fig. 5i) and recruitment of p300 (Fig. 5j). These results support that KDM1A-

dependent removal of the repressive H3K9me2 mark is needed for lamin B dissociation and

complete enhancer activation. Further, the pioneer-dependent priming of enhancers only

depends on H3K4me1 deposition independently of H3K9me2 depletion.

TAD size lamin B dissociation initiated by Pax7

We next assessed the extent of epigenetic changes associated with pioneer-dependent chromatin

opening using qPCR ChIP measurements of histones H3 and H1, together with levels of lamin B

association at three pioneered loci (Fig. 6). These loci are enriched for H3K9me2 and lamin B

(Fig. 6a, b) and ATAC-Seq profiles performed on wild-type and Pax7 mutant mouse pituitary

tissues defined TAD size domains (Fig. 6a) that show Pax7-dependent opening (Fig. 6b). In

contrast to the highly localized nucleosome depletion revealed by histone H3 levels at the center

of Pax7 pioneered enhancers, displacement of histone H1 is more variable, appears locus-

specific and may extend over several Kb's (Fig. 6c, d). The dissociation from lamin B is more

striking as it extends the full length of TADs containing the pioneered sites (Fig. 6). This is

consistent with our prior observation of TAD size chromatin opening at loci containing

melanotrope-specific genes such as *Pcsk2* and *Drd2* [20].

Whereas the three loci have different numbers of Pax7 recruitment sites and/or Pax7-dependent

changes in ATAC-Seq profiles, the depletion of histones H3 and H1 is localized around the Pax7

peaks in contrast with lamin B dissociation. These three loci have only one (Pio-14 and -6) or two

(Pio7) pioneered enhancers; the other Pax7 recruitment sites were in the primed state before

Pax7. By contrast to Pio7 that shows many Pax7 recruitment sites, loci Pio14 and Pio6 only have

one other significant Pax7 sites that may contribute to lamin B dissociation. It thus appears that a

single pioneered site may be sufficient to drive domain-scale lamin B dissociation.

In conclusion, and taken collectively, the present data suggest a model for pioneer factor action

(Fig. 7) in which the first steps are compatible with actions within the perinuclear compartment B:

these include removal of repressor marks together with deposition of activation marks

corresponding to the primed state. The ensuing steps for enhancer opening require passage

through cell division that allows for lamin dissociation, and presumably switching to the A

compartment, where the loci become accessible to the SWI/SNF complex for nucleosome

displacement and transcriptional activation.

Discussion

The present work shows that chromatin compaction elicited by histone H1 determines the initial

strength of pioneer Pax7 recruitment and speed of its actions. Beyond this limiting step, the

different parameters of chromatin opening follow a similar time course while the next limiting step

is imposed by cell division and its importance for lamin B dissociation (Fig. 7). Lamin B association

has been related to the levels of H3K9 methylation and the strength of this association correlated

with localization within perinuclear compartment B[22,23] or with subdomains of this compartment[25].

Whereas the extent of lamin B association is correlated with the level of H3K9me2 at various loci

(Fig. 5d), demethylation of H3K9me2 occurs earlier (step1) than lamin dissociation (step 2) with

the latter coinciding with cell division (Fig. 5a). Although pioneered loci were not visualized within

the nucleus in the present work, their association with lamin B suggest that the Pax7 pioneer

target enhancers are localized within the perinucleoplasmic compartment B. The association of

large LAD domains with the nuclear lamina and their dissociation following the loss of the

H3K9me2 mark[22,23] is consistent with our qPCR-ChIP data on lamin B association of pioneered

loci and the TAD-size decrease of lamin B association following Pax7-dependent remodeling (Fig.

6). This large scale lamin B dissociation contrasts with the more localized depletion of histone H1

around Pax7 pioneered enhancers that also exhibit nucleosome displacement. The relationship

between localized Pax7-dependent enhancer chromatin opening and the TAD size

rearrangements remain to be elucidated. Whether the decreased lamin B association of

pioneered enhancers reflects compartment (or sub-compartment[25]) switching or not, the apparent

dissociation reflects a significant change of organization as it is correlated with the ability for full

activation including recruitment of the SWI/SNF ATPase Brg1 that is only found in nucleoplasmic

compartment A [24].

Both our previous analyses[3] and recent papers[26,27] showed that pioneered sites exhibit strong

pioneer recruitment whereas weaker sites are more frequently associated with the primed states

although there is significant overlap between these subsets. These observations led some

authors to suggest that recruitment strength is critical for pioneer action[26,27]. We found that Pax7

isoforms or mutants that have decreased affinity fail to open subsets of targets but these analyses

also revealed intrinsic target site differences that presumably reflects context-dependent

chromatin heterogeneity[4,28]. Irrespective of these relations, the present work suggests a model

where permissive sites of Pax7 pioneer action are in various states of compaction as reflected by

the level of histone H1. While this compaction appears to hamper initiation of Pax7 action, it does

not prevent it as long-term levels of Pax7 recruitment and action are not directly related to the 1h

and 48h levels (Extended Data Fig. 4). Hence, the impact of histone H1 compaction is limited to

initial Pax7 recruitment.

In parallel with histone H1 displacement, Pax7 recruits the MLL complex that has histone

H3K4me1 methylase activity: enhancers that are primed after Pax7 recruitment have this

signature (Fig. 7c). The priming of enhancers is thus possible without dissociation from lamin B

(and possibly compartment B). Complete enhancer opening by Pax7 involves recruitment of the

cooperating factor Tpit and of the H3K9me2 demethylase KDM1A (Fig. 7b): the activity of this

enzyme is, at least partly, responsible for H3K9me2 demethylation and for subsequent steps of

enhancer activation, including dissociation from lamin B (Fig.5g-j). It will be interesting to compare

the stepwise process described here for Pax7-dependent pioneer action with other pioneers.

While there is as yet no similar temporal description of pioneer action, a two-tier process that is

consistent with our model was proposed for Zelda[29]. The targets of many pioneers, or the pioneers

themselves, were shown to recruit the MLL[13,14] and SWI/SNF[5,15,30] complexes; however, their

sequence of action or their relative involvement at the naïve or primed states are not documented.

Similarly, KDM1A was associated with the actions of pioneers, but also many TFs[31,32].

The realization that Pax7 pioneering targets are enriched in H3K9me2 and that removal of this

mark by KDM1A is required for activation raises the question of the nature of permissive sites for

other pioneers. While it is quite possible that different pioneers may target different "closed"

chromatin, it is difficult to assess the H3K9me2 status of chromatin sites targeted by other

pioneers as this mark was either not investigated or the naïve versus primed status of the targets

not sorted out[31] . Irrespective of pioneer involvement, the association of differentiation genes with

LAD switching from compartments B to A is however documented in many systems[25,33].

Similar limitations make it difficult to compare the role of replication or cell division in pioneer

action. This was investigated for FOXA-mediated chromatin opening by *Donaghey et al.[34]* who

showed that blockade of replication prevents FOXA-dependent DNA demethylation without

affecting bulk chromatin opening. Meaningful comparisons between different studies would

require clear distinction of pioneer target subsets as either naïve or primed because this impacts

dependence on replication for enhancer activation. Most cell reprogramming systems[35,36] typically

take place over a few days and would involve cell divisions. As far as we can tell, the link between

pioneering and replication is indirect as cell division appears required for nuclear compartment

switching[22] rather than for the unique aspects of pioneer action, namely the priming that results

in H3K4me1 deposition (Fig. 7). The recruitment of KDM1A to the pioneer Pax7 may thus

represent a mechanism for chromatin opening at enhancers that are within H3K9me2-associated

LADs. Whether recruitment of the MLL complex and of KDM1A are due to direct interactions with

Pax7 or Tpit respectively, or whether they are interdependent, remains to be established. But the

independence of the two critical changes of chromatin marks at pioneered enhancers supports

models where a single pioneer factor may implement various subsets of primed enhancers ready

for activation in response to an array of developmental (such as Tpit in our system) or signaling

(such as Stat3 and GR[3]) cues mediated through the activities of different nonpioneers.

References

1   Balsalobre, A. & Drouin, J. Pioneer factors as master regulators of the epigenome and cell fate.
Nature reviews. Molecular cell biology **23**, 449-464, doi:10.1038/s41580-022-00464-z (2022).
2   Soufi, A. *et al.* Pioneer transcription factors target partial DNA motifs on nucleosomes to initiate
reprogramming. *Cell* **161**, 555-568, doi:10.1016/j.cell.2015.03.017 (2015).
3   Mayran, A. *et al.* Pioneer factor Pax7 deploys a stable enhancer repertoire for specification of cell
fate. *Nat Genet* **50**, 259–269 doi:10.1038/s41588-017-0035-2 (2018).
4   Pelletier, A. *et al.* Pax7 pioneer factor action requires both paired and homeo DNA binding
domains. *Nucleic Acids Res* **49**, 7424-7436, doi:10.1093/nar/gkab561 (2021).
5   Minderjahn, J. *et al.* Mechanisms governing the pioneering and redistribution capabilities of the
non-classical pioneer PU.1. *Nat Commun* **11**, 402, doi:10.1038/s41467-019-13960-2 (2020).
6   Cirillo, L. A. *et al.* Binding of the winged-helix transcription factor HNF3 to a linker histone site on
the nucleosome. *EMBO J* **17**, 244-254, doi:10.1093/emboj/17.1.244 (1998).
7   Dodonova, S. O., Zhu, F., Dienemann, C., Taipale, J. & Cramer, P. Nucleosome-bound SOX2 and
SOX11 structures elucidate pioneer factor function. *Nature* **580**, 669-672, doi:10.1038/s41586-
020-2195-y (2020).
8   Michael, A. K. *et al.* Mechanisms of OCT4-SOX2 motif readout on nucleosomes. *Science* **368**, 1460-
1465, doi:10.1126/science.abb0074 (2020).
9   Tanaka, H. *et al.* Interaction of the pioneer transcription factor GATA3 with nucleosomes. *Nat
Commun* **11**, 4136, doi:10.1038/s41467-020-17959-y (2020).
10   Iwafuchi-Doi, M. *et al.* The Pioneer Transcription Factor FoxA Maintains an Accessible Nucleosome
Configuration at Enhancers for Tissue-Specific Gene Activation. *Mol Cell* **62**, 79-91,
doi:10.1016/j.molcel.2016.03.001 (2016).
11   Soufi, A., Donahue, G. & Zaret, K. S. Facilitators and impediments of the pluripotency
reprogramming factors' initial engagement with the genome. *Cell* **151**, 994-1004 (2012).
12   Nicetto, D. *et al.* H3K9me3-heterochromatin loss at protein-coding genes enables developmental
lineage specification. *Science* **363**, 294-297, doi:10.1126/science.aau0583 (2019).
13   Jozwik, K. M., Chernukhin, I., Serandour, A. A., Nagarajan, S. & Carroll, J. S. FOXA1 Directs H3K4
Monomethylation at Enhancers via Recruitment of the Methyltransferase MLL3. *Cell Rep* **17**,
2715-2723, doi:10.1016/j.celrep.2016.11.028 (2016).
14   Kawabe, Y., Wang, Y. X., McKinnell, I. W., Bedford, M. T. & Rudnicki, M. A. Carm1 regulates Pax7
transcriptional activity through MLL1/2 recruitment during asymmetric satellite stem cell
divisions. *Cell Stem Cell* **11**, 333-345, doi:10.1016/j.stem.2012.07.001 (2012).
15   King, H. W. & Klose, R. J. The pioneer factor OCT4 requires the chromatin remodeller BRG1 to
support gene regulatory element function in mouse embryonic stem cells. *eLife* **6**,
doi:10.7554/eLife.22631 (2017).
16   Takaku, M. *et al.* GATA3-dependent cellular reprogramming requires activation-domain
dependent recruitment of a chromatin remodeler. *Genome Biol* **17**, 36, doi:10.1186/s13059-016-
0897-0 (2016).
17   Gao, R. *et al.* Pioneering function of Isl1 in the epigenetic control of cardiomyocyte cell fate. *Cell
Res* **29**, 486-501, doi:10.1038/s41422-019-0168-1 (2019).
18   Mayran, A., Pelletier, A. & Drouin, J. Pax factors in transcription and epigenetic remodelling. *Semin
Cell Dev Biol* **44**, 135-144, doi:10.1016/j.semcdb.2015.07.007 (2015).

19   Budry, L. *et al.* The selector gene Pax7 dictates alternate pituitary cell fates through its pioneer
action on chromatin remodeling. *Genes & Development* **26**, 2299-2310,
doi:10.1101/gad.200436.112 (2012).
20   Mayran, A. *et al.* Pioneer and nonpioneer factor cooperation drives lineage specific chromatin
opening. *Nature Communications* **10**, 3807, doi:10.1038/s41467-019-11791-9 (2019).
21   Mohammed, H. *et al.* Rapid immunoprecipitation mass spectrometry of endogenous proteins
(RIME) for analysis of chromatin complexes. *Nat Protoc* **11**, 316-326, doi:10.1038/nprot.2016.020
(2016).
22   Poleshko, A. *et al.* H3K9me2 orchestrates inheritance of spatial positioning of peripheral
heterochromatin through mitosis. *eLife* **8**, doi:10.7554/eLife.49278 (2019).
23   Kind, J. *et al.* Single-cell dynamics of genome-nuclear lamina interactions. *Cell* **153**, 178-192,
doi:10.1016/j.cell.2013.02.028 (2013).
24   Reyes, J. C., Muchardt, C. & Yaniv, M. Components of the human SWI/SNF complex are enriched
in active chromatin and are associated with the nuclear matrix. *J Cell Biol* **137**, 263-274,
doi:10.1083/jcb.137.2.263 (1997).
25   Shah, P. P. *et al.* An atlas of lamina-associated chromatin across twelve human cell types reveals
an intermediate chromatin subtype. *Genome Biol* **24**, 16, doi:10.1186/s13059-023-02849-5
(2023).
26   Hansen, J. L., Loell, K. J. & Cohen, B. A. A test of the pioneer factor hypothesis using ectopic liver
gene activation. *eLife* **11**, doi:10.7554/eLife.73358 (2022).
27   Blassberg, R. *et al.* Sox2 levels regulate the chromatin occupancy of WNT mediators in epiblast
progenitors responsible for vertebrate body formation. *Nat Cell Biol* **24**, 633-644,
doi:10.1038/s41556-022-00910-2 (2022).
28   Bascunana, V. *et al.* Chromatin opening ability of pioneer factor Pax7 depends on unique isoform
and C-terminal domain. *Nucleic Acids Res*, 2023.2003. 2001.530660, doi:10.1093/nar/gkad520
(2023).
29   Brennan, K. J. *et al.* Chromatin accessibility is a two-tier process regulated by transcription factor
pioneering and enhancer activation. 2022.2012. 2020.520743 (2022).
30   Frederick, M. A. *et al.* A pioneer factor locally opens compacted chromatin to enable targeted
ATP-dependent nucleosome remodeling. *Nature structural & molecular biology* **30**, 31-37,
doi:10.1038/s41594-022-00886-5 (2023).
31   Heslop, J. A., Pournasr, B., Liu, J. T. & Duncan, S. A. GATA6 defines endoderm fate by controlling
chromatin accessibility during differentiation of human-induced pluripotent stem cells. *Cell Rep*
35, 109145, doi:10.1016/j.celrep.2021.109145 (2021).
32   Ismail, T. *et al.* KDM1A microenvironment, its oncogenic potential, and therapeutic significance.
Epigenetics & chromatin **11**, 33, doi:10.1186/s13072-018-0203-3 (2018).
33   Poleshko, A. *et al.* Genome-Nuclear Lamina Interactions Regulate Cardiac Stem Cell Lineage
Restriction. *Cell* **171**, 573-587.e514, doi:10.1016/j.cell.2017.09.018 (2017).
34   Donaghey, J. *et al.* Genetic determinants and epigenetic effects of pioneer-factor occupancy. *Nat
Genet* **50**, 250-258, doi:10.1038/s41588-017-0034-3 (2018).
35   Sardina, J. L. *et al.* Transcription Factors Drive Tet2-Mediated Enhancer Demethylation to
Reprogram Cell Fate. *Cell Stem Cell* **23**, 727-741.e729, doi:10.1016/j.stem.2018.08.016 (2018).
36   Stadhouders, R. *et al.* Transcription factors orchestrate dynamic interplay between genome
topology and gene regulation during cell reprogramming. *Nat Genet* **50**, 238-249,
doi:10.1038/s41588-017-0030-7 (2018).
37   Jiang, Y. *et al.* The methyltransferase SETDB1 regulates a large neuron-specific topological
chromatin domain. *Nat Genet* **49**, 1239-1250, doi:10.1038/ng.3906 (2017).

38   Langlais, D., Couture, C. & Drouin, J. The Stat3/GR interaction code: predictive value of
direct/indirect DNA recruitment for transcription outcome. *Molecular Cell* **47**, 38-49 (2012).
39   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. *Genome Biol* **10**, R25 (2009).
40   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079,
doi:10.1093/bioinformatics/btp352 (2009).
41   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data analysis.
Nucleic Acids Res **44**, W160-165, doi:10.1093/nar/gkw257 (2016).
42   Zhang, Y. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137, doi:10.1186/gb-
2008-9-9-r137 (2008).
43   Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-
regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-589 (2010).

ACKNOWLEDGMENTS

We are very grateful to Dr. Ali Shilatifard for the MLL3 and MLL4 antibodies, to Odile Neyret and

Sarah Boisset for NGS analyses, and to Valérie Magoon for her expert secretarial assistance.

Data analyses were possible thanks to the support of Compute Canada. This work was supported

by Foundation grant FDN-154297 to J.D. from the Canadian Institutes of Health Research.

AUTHOR CONTRIBUTIONS

A.G. and J.D. conceived the study, A.G., A.B. and J.D. conceived and designed the experiments,

A.G., J.D.-G., V.L.-R. and J. H. performed experiments, A.G., V.L.-R. and J.D. wrote the

manuscript.

COMPETING INTERESTS

The authors declare no competing financial interests.

Legends to Figures

Fig. 1 | Pax7 primes enhancers for activation. a, The role of Pax7 in specification of the

intermediate pituitary melanotrope cell fate is illustrated in contrast to the corticotrope cell fate.

These two hormone-secreting lineages express the same hormone precursor gene,

proopiomelanocortin (POMC), and their fates are determined by the Tbox transcription factor Tpit.

The mouse AtT-20 cells are a model of the corticotrope lineage and Pax7 expression reprograms

these cells towards a melanotrope-like fate. **b,** Heatmap representation of ChIP-Seq and ATAC-

Seq data for three subsets of Pax7-targeted enhancers determined in Extended Fig. 1b before (0

h) and after 48 h of Pax7 activation (tamoxifen treatment) in AtT20-ERPax7 cells. Pioneered

enhancers have no marks before Pax7 activation and present with hallmarks of transcriptionally

active enhancers after, namely appearance of an ATAC-Seq signal together with a bimodal

distribution of H3K4me1 deposition. Pax7 primed enhancers also show appearance of weaker

ATAC-Seq and H3K4me1 signals, the latter presenting as a single weak peak. Finally, a set of

constitutively active enhancers serve as control for comparisons. These enhancers bind Pax7,

but this binding is not associated with any change in chromatin signature. These enhancer

subsets were previously described[3,4]. **c,)** Mean profiles of H3K4me1 deposition at subsets of

Pax7-remodelled enhancers at indicated times following tamoxifen (Tam) activation of ER-Pax7.

d, Pax7 and H3K4me1 ChIP-Seq and ATAC-Seq signals at 14 randomly selected (Extended Data

Fig. 1d) enhancers pioneered by Pax7 after 48 h of tamoxifen activation. Sites are sorted (red to

blue) by the strength of Pax7 recruitment. **e,** Time course of H3K4me1 deposition determined by

ChIP-qPCR at the 14 pioneered sites illustrated above. For each site, a red curve illustrates

H3K4me1 accumulation under the Pax7 peak (centered), whereas a blue curve represents

H3K4me1 deposition over a lateral peak of deposition (shifted) that characterizes the bimodal

H3K4me1 distribution of fully active enhancers. **f,** Distribution of H3K4me1 deposition $t_{1/2}$ from

(e). P-value from two-tailed t-test (*** ≤ 0.001, ** ≤ 0.01, * ≤ 0.5, – otherwise). **g,** Mean (± standard

error of the mean, n ≥ 2) time course at 14 pioneered sites of H3K4me1 accumulation (centered

and shifted) is compared with ChIP-qPCR data for recruitment of two protein components of the

MLL remodelling complex, MLL4 and Ash2.

Fig. 2 | Pax7-dependent chromatin opening occurs in two steps. a, Average (14 pioneered

sites of Fig.1d represented as means ± sem of duplicate biological samples, n ≥ 2) profiles of

ChIP-qPCR data for Pax7, Tpit, MLL4 and Ash2 that are recruited with a similar time course (T ½

~ 12 hrs) compared to accumulation of H3K4me1 (center). **b,** Compared to Pax7 recruitment,

ATAC-Seq signals and recruitment of Brg1, p300 and SMC1, together with depletion of Histone

H3 occur with an average T ½ of ~ 24h (n ≥ 2). **c,** Deposition of activation mark H3K27ac and

bimodal H3K4me1 (shifted) are slightly delayed compared to ATAC signal (n ≥ 2). **d,** Depletion of

the repressive mark H3K9me2 and of linker Histone H1 occur with a T ½ of ~ 12h (n ≥ 2). **e,**

Median (± median absolute deviation) half-times of change for the indicated parameter at the 14

pioneer sites. **f,** Median (± median absolute deviation) half-times for all parameters altered by

Pax7 pioneering regrouped for step 1 (blue) and step 2 (pink) of the pioneering process. Data are

shown for each of 14 pioneered enhancers (n ≥ 2). All panels: p-values from two-tailed t-tests (***

≤ 0.001, ** ≤ 0.01, * ≤ 0.5, – otherwise).

Fig. 3 | Initiation of Pax7 recruitment is delayed by histone H1-dependent chromatin

compaction. a, Individual site ChIP-qPCR time course data (means ± standard error of the mean,

n ≥ 2) for recruitment of Pax7, H3K4me1 and Brg1, together with time courses of ATAC signals

and depletion histones H1 and H3 shown for the 14 pioneered sites color-coded (in Fig.1d) from

red to blue to reflect the strength of Pax7 recruitment. **b,** Correlations between Pax7 recruitment

strength and various markers depicted in (a). **c,** Pax7 ChIP-Seq and ATAC-Seq time courses at

all genome-wide pioneered enhancers. Sites color-coded according to Pax7 recruitment at 1 h

and 48 h. **d,** Correlation at 14 pioneered enhancers between Pax7 recruitment signals at 1 h and

48 h. **e,** Inverse correlation between initial (1h) Pax7 recruitment strength and the average half-

times of all markers of chromatin remodelling at the 14 pioneered sites. **f,** Distribution of initial

histone H1 levels at constitutively active enhancers (con.), pioneered enhancers (pio.) and

random sites (rnd) sites determined either genome-wide or from the qPCR panel (Extended Data

Fig. 2). **g,** Correlations between initial histone H1 levels at the 14 pioneered enhancers and Pax7

recruitment at 1 h as well as with the mean $t_{1/2}$ of all markers of the pioneering process. **h,** UMAP

representation of Pax7 recruitment strengths at the indicated times after Tam addition compared

to initial histone H1 levels measured by ChIP-Seq at pioneered enhancers depicted in Fig. 1b. All

panels: p-values from two-tailed t-tests (*** ≤ 0.001, ** ≤ 0.01, * ≤ 0.5, – otherwise).

Fig. 4 | Activation of Pax7 pioneered enhancers requires cell division. a, Experimental design

for assessment of chromatin remodelling in cells with mimosine-blocked cell cycle compared to

normally cycling cells. **b,** FACS profiles of untreated cells compared to cells incubated for 12h

with mimosine that exhibit cell-cycle arrest in G1. Representative of n=2 is shown. **c,** The

recruitment of both Pax7 and Tpit, as well as depletion of H3K9me2, measured by ChIP-qPCR at

the 14 pioneered sites are not affected by blocking AtT-20 cell cycle with mimosine (n=2). In

contrast, DNA accessibility (ATAC-qPCR) as well as recruitments of Brg1, p300 and H3K27ac

measured by ChIP-qPCR are all severely curtailed in mimosine-arrested cells (n≥2). P-values

from two-tailed t-tests (*** ≤ 0.001, ** ≤ 0.01, * ≤ 0.5, – otherwise). **d, e,** Average profiles of

H3K4me1 (**d**) and ATAC-Seq signals (**e**) accumulation at Pax7-remodelled enhancer subsets

measured by ChIP-Seq at different times after Tam activation of ER-Pax7 in dividing or mimosine-

arrested cells as indicated.

Fig. 5 | Pax7-dependent H3K9me2 demethylation and dissociation from lamin B. a, Mean

(14 pioneered sites of Fig.1d represented as means ± sem of duplicate biological samples) time

courses for depletion of H3K9me2 and dissociation from lamin B1 measured by ChIP-qPCR

compared to the time courses of recruitment for the H3K9me2 demethylase, KDM1A (LSD1). **b,**

c, Pax7 pioneered sites have high levels (ChIP-Seq) of lamin B1 (**b**) and H3K9me2 (**c**) before

Pax7 action. **d,** Correlation between initial levels (ChIP-qPCR, n ≥ 2) of H3K9me2 and lamin B1

at 14 pioneered sites. **e,** Ratios of Lamin B1 levels (ChIP-qPCR) at 48 over 0 h of Pax7

activation at constitutively active enhancers (con.), pioneered enhancers (pio.) and random loci

(rnd) in dividing compared to mimosine-arrested AtT20 cells (n ≥ 2). **f,** Heatmaps of ChIP-Seq

profiles for Pax7, H3K4me1, H3K9me2, Tpit, KDM1A(LSD1) and Brg1 at subsets of Pax7

remodelled enhancers. Data for other markers presented in Extended Data Fig. 7. **g-j,** ChIP-

qPCR variation of Pax7 (g) and p300 (j) recruitment as well as H3K9me2 (h) and Lamin B1 (i)

levels between no and 48 h of Pax7 activation at constitutively active enhancers (con.), the 14

pioneered enhancers (pio.) and random loci (rnd) in normal and cells treated (using highest

concentration devoid of cytotoxicity) with the KDM1A inhibitor OG-L002 (n ≥ 2). Experimental

design similar to mimosine treatment depicted in Fig. 4a. All panels: p-values from two-tailed t-

tests (*** ≤ 0.001, ** ≤ 0.01, * ≤ 0.5, – otherwise).

Fig. 6 | Pax7 initiates domain-wide lamin B dissociation. a, HiC contact scores in neural

cells[37] and annotated genes at three TADs harboring a Pax7-pioneered enhancer (including the

hallmark Psck2 enhancer, Pio-14) and their flanking regions. **b,** Stable Pax7 ChIP-seq

recruitment in AtT20-Pax7 cells. H3K9me2 and Lamin B1 ChIP-seq in AtT20 cells. ATAC-seq in

WT and Pax7-KO intermediate lobe (IL) mouse pituitary cells depicting the dependance on

Pax7 for activation of these TADs. Units: reads per million. **c,** Mean changes (± standard error

of the mean, n ≥ 3) in histones H3, H1 and Lamin B1 levels following stable Pax7 expression in

AtT20 cells. Data acquired by ChIP-qPCR at numerous sites across target TADs and flanking

regions. By contrast to the local histone H3 and H1 depletions, Lamin B1 dissociation spans the

entirety of the TADs. Flanking regions are not affected by Pax7. **d,** Same as **(c)** but zoomed on

Pax7-targeted enhancer. **e,** Local Pax7 recruitment by ChIP-Seq at the pioneered enhancers. **f,**

Distribution of Pax7 recruitment, histone H3 and H1 and Lamin B1 changes depicted in **(c)** with

measured sites clustered as: flanking TADs, within target TAD excluding Pax7 target ± 5 kb and

within Pax7 target ± 5 kb. P-values from two-tailed t-tests (*** ≤ 0.001, ** ≤ 0.01, * ≤ 0.5, –

otherwise).

Fig. 7 | Model of Pax7-initiated chromatin opening and enhancer activation. a, Schematic

representation of proteins shown in the present work to be associated with either primed or active

enhancer states. **b,** Sequential model of Pax7 pioneer action. Pax7 pioneer targets are enriched

in H3K9me2 and associated with lamin B1, which is typical of chromatin that is present in the

perinuclear B compartment. Step 1 of the pioneering process involves recruitment of Pax7 and

Tpit together with the H3K9me2 demethylase KDM1A (LSD1) and the MLL complex. These lead

to demethylation of H3K9me2 and H3K4me1 deposition, respectively. Cell cycle blockade with

mimosine prevents Pax7-dependent remodelling from proceeding further. At mitosis, masking of

the H3K9me2 mark by H3S10 phosphorylation accompanies dissociation of A and B

compartments of the nucleus[22]. Following mitosis and nuclear reassembly, Pax7-pioneered

targets that have undergone H3K9me2 demethylation will remain in the central nuclear A

compartment where the Brg1 and Brm ATPase's of the SWI/SNF complex are localised[24] and

recruited for nucleosome displacement and complete enhancer activation. **c,** Pax7 primed

enhancers do not recruit the cooperating nonpioneer factor Tpit and hence, the cooperation

between Tpit and Pax7 appears required for significant H3K9me2 demethylation and continuation

of the Pax7-dependent remodelling process to the cell cycle-dependent step 2.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

a

Priming $t_{1/2}$ = 12 h
Varies with chromatin condensation

Pax7
Tpit   MLL   KDM1A

Activation $t_{1/2}$ = 24 h
Requires H3K9me2 loss and cell division

ACTIVE ENHANCER

Pax7
Tpit   Brg1   p300

b

B
A

| Pax7 | Tpit |
| MLL | KDM1A |

+ H3K4me1 − H3K9me2

B
A

Lamins dissociation

REPLICATION & MITOSIS

H3K9me2-mediated reassociation

Brg1 ↓ p300

B
A

c

| Pax7 | ~~Tpit~~ |
| MLL | ~~KDM1A~~ |

+ H3K4me1

Stable primed state

✗

○ H3K4me1   ○ H3K9me2   ⬭ Cohesin   ✕✕ Nuclear lamins
○ H3K27ac   ○ H3S10ph   [H1] Histone H1   **A****B** Compartments

Fig. 7