Novel regulatory logic in the antibiotic resistance response of *Enterococcus faecalis* against cell envelope targeting antibiotics

Sali M. Morris[1], Georg Fritz[2], Tim Rogers[3], Susanne Gebhard[1*]

[1]Life Sciences Department, Milner Centre for Evolution, University of Bath, Claverton Down, Bath, BA2 7AY, United Kingdom; [2]School of Molecular Sciences, The University of Western Australia, WA 6009 Crawley, Australia; [3]Department of Mathematical Sciences, University of Bath, Claverton Down, Bath, BA2 7AY, United Kingdom

*Corresponding author. Phone: +44 1225 386421; E-mail: s.gebhard@bath.ac.uk;

Running title: Antibiotic resistance control in *E. faecalis*

Data accessibility statement: The data that support the findings of this study are available in the supplementary material.

Funding statement: This work was supported in part by grant MR/N0137941/1 for the GW4 BIOMED MRC DTP, awarded to the Universities of Bath, Bristol, Cardiff and Exeter from the Medical Research Council (MRC)/UKRI.

Conflicts of interest: The authors declare no conflicts of interest.

Summary

Enterococcal infections frequently show high levels of antibiotic resistance, including to cell envelope-acting antibiotics like daptomycin (DAP). While we have a good understanding of the resistance mechanisms, less is known about the control of such resistance genes in enterococci. Previous work unveiled a bacitracin resistance network, comprised of the sensory ABC transporter SapAB, the two-component system (TCS) SapRS and the resistance ABC transporter RapAB. Interestingly, components of this system have recently been implicated in DAP resistance, a role usually regulated by the TCS LiaFSR. We here explored the interplay between these two regulatory pathways. Our results show the regulation by SapR of an additional resistance operon, *dltXABCD,* and show that LiaFSR regulates the expression of *sapRS*, placing SapRS target genes under dual control: *dltXABCD* expression depends on both antibiotic-induced cellular damage (LiaFSR) *and* the presence of a substrate drug for the sensory transporter (SapAB). We further show that this network protects *E. faecalis* from antimicrobials produced by potential competitor bacteria, providing a potential rationale for the evolution of this regulatory strategy. The network structure described here can explain why clinical DAP resistance often emerges via mutations in regulatory components, which may ultimately lead to the discovery of new therapeutic targets.

Keywords: antimicrobial peptides, cell envelope stress, antimicrobial resistance, two-component system, signalling

Introduction

The rise of antibiotic resistant bacteria is one of the greatest current threats to public health, resulting in 670,000 infections a year and 33,000 deaths in Europe alone[1]. Of these infections, the "ESKAPE" organisms[2] (*Enterococci* spp., *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, *Enterobacter* spp.) have driven the rising number of nosocomial and antibiotic-resistant infections in the past decade. Of these bacteria, the enterococci are the second most common causative agent of nosocomial infections in the US; including bacteraemia, endocarditis and urinary tract infections[3–5]. The two species most frequently isolated, *Enterococcus faecalis* and *Enterococcus faecium*, remain a major infection-control challenge, particularly in healthcare settings.

Enterococci became recognised as important nosocomial pathogens due to their high level of intrinsic resistance to several antimicrobials[6] (e.g. penicillin, ampicillin and cephalosporins) and their capacity to acquire further resistance determinants. One such acquired resistance is to the glycopeptide antibiotic vancomycin, which occurs through plasmid acquisition and was first reported in the 1988[7,8], 30 years after vancomycin was introduced for clinical use[9]. Despite the molecular mechanisms of vancomycin resistance in enterococci being well understood today[10], infections by vancomycin resistant enterococci (VRE) still result in serious health and economic impacts[11] and are an increasing problem worldwide.

One of the last-resort antibiotics used to treat these VRE infections is the lipopeptide antibiotic daptomycin (DAP)[12]. Disappointingly, within 2 years of clinical introduction of the drug in 2003, DAP-resistant enterococcal isolates were reported[13], and in contrast to vancomycin resistance, this occurred through a subtle chromosomal change based on the mutations in genes *liaF*, *cls* and *gdpD*[14–17]. LiaF is a transmembrane protein involved in monitoring the integrity of the cell envelope and responding to damage[18], whereas Cls (cardiolipin synthase) and GdpD (glycerol-phosphodiester phosphodiesterase) are both involved in phospholipid metabolism[14]. Of these, mutation of *liaF* has been proposed to be the first pivotal event towards daptomycin resistance[15,19]. However, the exact role of the *liaF* mutations in DAP resistance so far remains unclear.

LiaF is part of a three-component regulatory system, LiaFSR, which is important amongst the Firmicutes for coordinating the cell envelope stress response (CESR) against antimicrobial-induced damage[20–23]. The system is comprised of LiaF and a conventional two-component system (TCS): the sensor kinase LiaS and the response regulator LiaR[20,24,25]. LiaF is an inhibitor of LiaS, maintaining the sensor kinase in an inactive conformation[20]. Rather than detecting a specific antimicrobial compound, LiaFSR responds to cell envelope damage, although the exact stimulus is unknown[26]. Upon sensing this damage, LiaF releases LiaS, which is then able to phosphorylate LiaR to induce the expression of the system's target operon, *liaXYZ*[19,27]. The *liaXYZ* cluster is involved in sensing and binding antimicrobials at the cell surface to provide resistance[27].

To add to this complexity, the Lia system does not exist in isolation, but is just one of many TCSs involved in monitoring cell envelope integrity, each responding to their own individual stimuli and activating a unique set of target genes. The CroRS system, unique to the enterococci, is the main regulator required for cephalosporin resistance[28], whereas the VicKR (YycFG) system is essential across the low-CG Gram positives and is involved in regulating cell division, lipid biosynthesis, biofilm and virulence[29]. An additional element of the network is the serine/threonine kinase IreK and the phosphatase IreP, involved in maintaining cell wall integrity by potentially regulating peptidoglycan biosynthesis and metabolism[30–32].

A further TCS involved in monitoring the cell envelope is EF0926/27, which we have now renamed SapRS (<u>S</u>ensor of <u>A</u>ntimicrobial <u>P</u>eptides), identified in our previous work[33]. SapRS is part of a bacitracin resistance network comprised of the histidine kinase SapS, the response regulator SapR and the Bce-like ABC transporters: SapAB (EF2752/51) and RapAB (<u>R</u>esistance against <u>A</u>ntimicrobial <u>P</u>eptides) (EF2050/49)[33]. The use of the sensory transporter SapAB to control the activity of SapRS implements a 'flux-sensing' mechanism to regulate bacitracin resistance, as was shown for the homologous system in *Bacillus subtilis*[34]. In brief, upon exposure, bacitracin binds to its membrane-associated target molecule, undecaprenol-pyrophosphate (UPP), blocking the dephosphorylation and recycling of UPP in the lipid II cycle of cell wall biosynthesis[35,36]. The sensory ABC transporter SapAB, based on biochemical evidence from the *B. subtilis* system, forms a sensory complex with SapRS[37]. In

the absence of bacitracin, the transporter maintains the histidine kinase, SapS, in an 'OFF' state[38]. Upon

detection of bacitracin-UPP (BAC-UPP) complexes, SapAB switches its role from repressor to activator

of SapS, resulting in kinase autophosphorylation[38]. SapS can then phosphorylate SapR, which in turn

induces the production of the resistance transporter RapAB. RapAB, again based on evidence from its

B. subtilis homologue, frees UPP from the inhibitory grip of bacitracin using a target protection

mechanism[39], allowing dephosphorylation of UPP and continuation of cell wall synthesis, rendering the

cell resistant to bacitracin. The equivalent system in *B. subtilis* is a self-contained module, involving

only a single transporter (BceAB) and TCS (BceRS)[40] with no known further genes involved in its

signalling or resistance mechanism. However, in *E. faecalis*, we observed that the TCS SapRS is itself

transcriptionally induced by bacitracin, however we had not identified the regulatory system that

controls this expression[33].

Interestingly, recent work by the Arias lab demonstrated that when experimentally evolving *E. faecalis*

for DAP resistance in a Δ*liaFSR* background, mutations were observed in the sensory transporter

SapAB[41,42], suggesting a functional link between the Sap and Lia systems. In *B. subtilis*, the Lia system

is known to be one of the main components of bacitracin resistance[43,44], but there is currently no

evidence for a role of LiaFSR in response to bacitracin in *E. faecalis*. Neither is there any indication of

a role for the Sap system in responding to DAP exposure. However, this recent evidence suggested an

interplay between both the Sap and Lia systems and that, potentially, both systems may be contributing

to resistance against bacitracin and daptomycin.

In accordance with the need to deepen our knowledge of the CESR in *E. faecalis,* in this study we

sought to examine the potential interplay between the Sap and Lia systems and investigate the

involvement of further genes associated with this regulatory network. Utilising mutagenesis and

analysis of promoter activity, we provide evidence of the activation of LiaFSR signalling in response

to bacitracin and describe a direct functional link between the Lia and Sap systems by demonstrating

the regulation of *sapR* by LiaR in response to antibiotic exposure. We also show the contribution of an

additional resistance operon involved in response to bacitracin and unravel a differential response of

the network between bacitracin and daptomycin treatment. Our data show that interplay between SapRS

and LiaFSR signalling effectively implements a logic 'AND' gate, whereby expression of SapR target

genes is under dual control and requires two separate signalling inputs. These observations also explain

the occurrence of mutations in both regulatory systems during clinical and experimental emergence of

DAP resistance in enterococci found previously, thus providing a compelling model for the complex

CESR network in *E. faecalis*.

Results

LiaFSR controls expression of SapRS. As mentioned above, from previous work we had identified

increased expression of *sapRS* under exposure to bacitracin or mersacidin[33]. Interestingly, we also

showed that *sapRS* was not autoregulated, in contrast to other two component systems such as CroRS

and LiaFSR[20,28], and regulation did not depend on either SapAB or RapAB[33]. We therefore here first

aimed to identify the regulator of *sapRS*. For possible candidates, we considered potential regulators

that are common amongst the Firmicutes and are known to respond to bacitracin. A potential candidate

was LiaFSR, deemed the 'master regulator' of the CESR in *Bacillus subtilis*[43] and recently shown to

have a possible functional link to the Sap system[41,42]. Because of this evidence, we first aimed to

examine the relationship between these two systems.

Firstly, we had to test if LiaFSR was able to respond to bacitracin treatment. It is well established that

liaX is under the control of LiaFSR[27], and therefore we exposed *E. faecalis* harbouring a P$_{liaX}$-*lacZ*

transcriptional fusion to increasing levels of bacitracin, as a readout for LiaFSR activity. The results

showed a ~10-fold increase in *liaX* expression at 32 μg ml^{-1} bacitracin compared with untreated cells

(Fig. 1A, black line), indicating that LiaFSR can indeed respond to bacitracin exposure in *E. faecalis*.

Figure 1. The LiaFSR operon responds to bacitracin exposure and induces the expression of both *liaXYZ* and *sapRS*, with the absence of *sapRS* resulting in the hypersensitivity of the *liaXYZ* promoter. Cells of *E. faecalis* JH2-2 wild-type or isogenic deletion strains harbouring a P_{liaX}-*lacZ* (panel A) or P_{sapRS}-*lacZ* (panel B) transcriptional fusion were grown to exponential phase and challenged with increasing concentrations of bacitracin. β-galactosidase activity, expressed as Miller units (MU), was assayed following 1h incubation. Results are means and standard deviations for three biological repeats. The significance of induction relative to untreated cells was calculated for each strain by a one-tailed, pairwise *t*-test analysis. Significance is indicated by a filled symbol ($p < 0.05$); unfilled symbols represent no significant increases compared to uninduced conditions.

Following this, we tested if LiaR can regulate the expression of *sapRS*. To do this, we monitored the response of a transcriptional P_{sapRS}-*lacZ* fusion to increasing bacitracin. The results showed that in wild-type *E. faecalis* carrying the transcriptional *sapRS-lacZ* fusion, bacitracin concentrations of 16 µg ml⁻¹ or higher led to significant induction of the reporter, with a three-fold increase compared to uninduced cells at 32 µg ml⁻¹ (Fig. 1B, black line). Deletion of *liaR* in the reporter strain resulted in a complete loss of *sapRS* induction, with expression remaining at basal levels. This indicated that LiaR indeed regulates the expression of *sapRS* in response to bacitracin exposure, presenting first evidence of a direct functional link between the Lia and Sap regulatory systems.

A surprising feature of this LiaR regulation was the high concentration required to induce a response, as *liaX* expression only significantly increased at 32 µg ml⁻¹. One possible explanation for this might be that the response was masked by other components of the bacitracin stress response in *E. faecalis*, for example RapAB, which plays an active role in the removal of bacitracin from UPP[33], protecting the

cell from damage. In *B. subtilis*, it was observed that the RapAB equivalent, BceAB, was the primary

response to protect the cell from bacitracin exposure and masked the response of the Lia system[45]. To

prevent the induction of the main bacitracin resistance genes of *E. faecalis*, controlled by SapRS[33] and

thus remove potential interference with Lia signalling, we therefore introduced the P*liaX*-*lacZ*

transcriptional fusion into the Δ*sapRS* background. Compared to the wild type, the expression of the

liaX promoter in response to bacitracin challenge of the deletion strain showed increased sensitivity,

significantly inducing expression from 4 μg ml^{-1} bacitracin and reaching overall higher activities (Fig.

1A, blue line). This showed that the weak response of Lia signalling to bacitracin in the wild type was

indeed due to masking effects of the resistance genes controlled by SapRS, implying the presence of

multiple layers of protection, similar to those observed in *B. subtilis*[45], as well as physiological links

between the SapRS- and LiaFSR-dependent components of the CESR in *E. faecalis*.

SapRS controls the expression of the *dltABCD* operon. To expand our understanding of the Lia/Sap

regulatory network, we next considered if other genes may also be under its control. Currently, the only

known gene under SapRS regulation is the resistance ABC-transporter operon *rapAB* [33]. In *B. subtilis*,

both the Lia and Bce systems act as self-contained modules, controlling the regulation of only a single

resistance operon each, *liaIH* and *bceAB* respectively, each encoded adjacently to its regulatory operon

on the chromosome [24,40,43]. However, the regulatory setup in *E. faecalis* seems to be much more

complex. This is supported by evidence from the literature, with the indication that the Lia system has

a larger operon than just itself and *liaXYZ* [27]. There is also a proposal that Lia contributes to DAP

resistance through the regulation of *dltXABCD* (*dlt*), although there is no evidence to date that Lia is a

direct regulator of this operon. Dlt is responsible for the D-alanylation of teichoic acids (TA) on the

bacterial cell surface, resulting in a decrease in the negative charge of the cell envelope[46], a mechanism

commonly involved in DAP resistance amongst the low-CG bacteria[47].

Interestingly, recent evidence has suggested a regulatory link between SapRS and *dlt*. When evolving

E. faecium for DAP resistance in the absence of LiaFSR, mutations occurred in the *sapAB* homologue,

ycvRS. These mutations correlated with an increase in *dlt* transcription when measured by qPCR and

resulted in an increase in cell surface charge[41,42]. In addition, the *dltABCD* operon is located directly

downstream from *sapAB* on the chromosome, suggesting a possible functional link between the genes. Moreover, the promoter region of *dlt* contains a putative SapR binding site, similar to that of the *rapAB* promoter (Fig. 2). This body of evidence therefore led us to investigate the contribution of LiaFSR and SapRS to *dlt* regulation.

Figure 2. Schematic of the promoter regions of both *rapAB* and *dltABCD*. The sequence 150 base pairs upstream of the ATG start codons are shown. The proposed binding site for SapR on both the *rapAB* and *dltABCD* promoters is in bold and capitalised, and the likely -10 and -35 elements are underlined. The experimentally confirmed transcriptional start sites are capitalised and underlined (+1) and the translational start site is shown in italicised capitals[33,84].

To this end, we first constructed a transcriptional P$_{dltX}$-*lacZ* fusion to test *dlt* induction by bacitracin. The results showed that in wild-type JH2-2 carrying the fusion, increasing bacitracin concentrations led to a dose-dependent increase in *dlt* expression, resulting in ~4-fold higher activity at 32 μg ml^{-1} compared to untreated conditions (Fig. 3A, black line), showing that *dlt* expression is indeed induced in response to bacitracin in *E. faecalis*. When we tested the response of the reporter in strains carrying deletions of either *sapRS* or *sapAB,* the results showed a decrease in basal activity in both strains and a complete loss of the promoter's bacitracin response in Δ*sapR* (Fig. 3A, blue and magenta lines). Loss of SapAB still resulted in induction compared with uninduced cells, but overall activities were considerably lower than in the wild-type strain. This indicated that SapRS was essential for *dlt*

199   expression in response to bacitracin, but a residual amount of *dlt* induction remained in the *sapAB*

200   deletion. This is consistent with the less direct role of SapAB in signalling, i.e. via controlling SapRS

201   activity and not the target genes directly. The residual bacitracin response observed in the *sapAB* deleted

202   strain can likely be explained by increased production of SapRS due to LiaFSR signalling, which would

203   be expected to lead to increased basal activity of SapRS and thus the *dlt* promoter.

Figure 3. Induction of the resistance operon *dltXABCD* by bacitracin. Panel A indicates the experimental induction of *dltX*. Cells of *E. faecalis* harbouring a *PdltX-lacZ* transcriptional fusion were grown to exponential phase and challenged with increasing concentrations of bacitracin. Beta-galactosidase activity, expressed as Miller units (MU), was assayed following 1h incubation in wild type (WT) and deletion strain backgrounds. Results are means and standard deviations for three biological repeats The significance of induction relative to untreated cells was calculated for each strain by a one-tailed, pairwise *t*-test analysis. Significance is indicated by a filled symbol ($p < 0.05$), unfilled symbols represent no significant increases compared to uninduced conditions. Panel B indicates the mathematical model of *dltX* induction in the *E. faecalis* strains indicated in panel A.

As we had shown above that LiaR regulates *sapRS* expression, we next aimed to find out if deletion of

liaR also influenced *dlt* expression. When we tested the response of the *dltX* promoter fusion to

bacitracin in a strain carrying deletion of *liaR*, we observed that deletion of *liaR* resulted in a noticeably

weaker amplitude of *dlt* expression in response to bacitracin, being ~2-fold lower than the wild-type

strain at 32 μg ml^{-1} (Fig. 3A, yellow line). In addition, the sensitivity of the *dlt* promoter response was

much lower in the *liaR* deletion compared with the wild-type response, with significant activation over

baseline occurring at 16 μg ml^{-1} rather than at 2 μg ml^{-1} in the wild type. However, there was no effect

on basal activity of *dlt*, in contrast to deletion of *sapR* and *sapAB*. These data are consistent with LiaR

regulating *sapRS* expression, and SapRS being the actual regulator of *dlt* expression in response to

bacitracin.

As we had observed before with the *liaX* and *sapRS* promoters, we also wanted to investigate if the

protection provided by the SapRS-target RapAB was dampening the expression of *dlt* in response to

bacitracin. To examine this, we tested the response of *dlt* expression in the absence of *rapAB*. In a

ΔrapAB background, the *dlt* promoter demonstrated markedly increased sensitivity, and a much

stronger response to bacitracin, resulting in a ~3-fold increase in expression at 2 μg ml^{-1} when compared

to the wild-type (Fig. 3A, orange line). This response demonstrated the presence of a layered protection

with RapAB activity moderating *dlt* expression.

Overall, these findings were rather surprising, as the signalling pathway appeared remarkably complex

to result in a relatively simple outcome, i.e. inducing the expression of two resistance genes in response

to an antibiotic. To test if our understanding of the regulatory pathway was plausible, we therefore

developed a representative mathematical model to see if this would reproduce the behaviour we had

observed in the experiments. At the core, this model was based on a simplified form of the flux-sensing

mechanism described previously for the *B. subtilis* BceRS-BceAB system[34], which was then expanded

upon to reproduce the hypothesised network structure investigated here. In brief, the model considered

bacitracin binding to UPP to form UPP-bacitracin complexes with the on-rate dependent on the

bacitracin concentration. These complexes then drive transport activity of, in this case, SapAB,

according to a soft-switch type functional response (see Methods), which the model translates into

activation of SapRS and thus *dlt* expression. Importantly, SapRS activity also drives expression of the

resistance transporter operon, in this case *rapAB*, and production of RapAB leads to a reduction in

formation of UPP-bacitracin complexes due to the target protection activity of the transporter[34,39]. This

creates the negative feedback loop that is characteristic of the flux-sensing mechanism and leads to the

gradual response behaviour of the output promoters (Fig. 3B, black symbols). To adapt this model to

the Lia-Sap regulatory pathway of *E. faecalis*, we additionally considered the activity of the Lia system.

This was modelled to also respond to UPP-bacitracin complexes in a switch like manner representing

the generation of cellular damage caused by these complexes. The model then linked the Lia and Sap

components by modulating the SapRS signalling output (i.e. *dlt* and *rapAB* expression) according to

Lia activity, driven by bacitracin. The mathematical details of the model are explained in the Methods

section. Fitting to experimental data was performed by matching differential activity of SapRS against

both *rapAB* and *dlt* recorded at different bacitracin concentrations.

This model accurately reflected the behaviour of our experimental strains, depicting the same gradual

response to bacitracin of the *dltX* promoter in wild-type *E. faecalis* (Fig. 3B, black symbols), as well as

the hypersensitive response in the *rapAB* deletion strain, where the negative feedback from RapAB-

driven removal of bacitracin was missing (orange symbols). Importantly, the model gave the same

complete loss of *dltX* activity when *sapRS* was deleted (blue symbols) as observed experimentally, as

well as the normal basal level activity but loss of bacitracin-dependent induction when *liaR* was deleted

(yellow symbols). A *sapAB* deletion strain was not specifically considered as the model did not

differentiate between SapAB and SapRS activities and thus both strains would have been

mathematically identical. This close agreement between theoretical and experimental data strongly

suggested that our reconstruction of the regulatory pathway and connection between the Lia and Sap

systems was theoretically plausible and that no further major players were needed to explain the

behaviour of the *dltX* target promoter in the various mutant backgrounds analysed here.

Daptomycin induces the expression of *sapRS* but not *dltABCD*. We had now established a sequential

order of *dlt* regulatory control; in response to bacitracin, LiaFSR induces the expression of *sapRS*, and

in turn SapRS, activated by its sensory transporter SapAB, induces the expression of *dlt*. However,

treatment with bacitracin made it difficult to separate out the individual contributions to regulation by

LiaFSR and SapRS, as both signalling systems respond to this antibiotic. Bacitracin forms a complex

with UPP, which acts as the input for SapRS signalling via SapAB[39]; at the same time, bacitracin

induces cell envelope damage, triggering LiaFSR activation[26]. To allow us to differentiate between the

contribution made by each regulator to the network, we instead required an antibiotic that would only

trigger signalling by one of the systems. To this end, we turned to daptomycin (DAP), which activates

the Lia system by causing damage to the cell envelope[48,49]. However, it does not bind to UPP or the

sugar-pyrophosphate moiety of Lipid II, which is the common feature of substrates for BceAB-type

transporters[39]. Indeed, the tripartite complex of DAP with Lipid II and phosphatidyl glycerol in the

membrane[50] is unlikely to be recognised by SapAB, and thus DAP exposure should not trigger SapRS

activation, allowing us to study the specific impact of LiaFSR signalling in the regulatory pathway.

Figure 4. Exposure to daptomycin results in the induction of *sapRS* but does not induce the expression of *dltXABCD*. Cells of *E. faecalis* harbouring a P_{sapRS}-*lacZ* (panel A) or a P_{dltX}-*lacZ* (panel B) transcriptional fusion were grown to exponential phase and challenged with increasing concentrations of daptomycin. Beta-galactosidase activity, expressed as Miller units (MU), was assayed following 30 minutes incubation in wild type (WT) and deletion strain backgrounds indicated. Results are means and standard deviations for two biological repeats. The significance of induction relative to untreated cells was calculated for each strain by a one-tailed, pairwise *t*-test analysis. Significance is indicated by a filled symbol (P < 0.05), unfilled symbols represent no significant increases compared to uninduced conditions.

To investigate the effect of DAP exposure on LiaFSR and SapRS signalling and target gene expression,

we again utilised transcriptional promoter fusions. The P_{sapRS}-*lacZ* transcriptional fusion (LiaFSR

target) in the wild-type background showed a ~2-fold induction upon DAP treatment when compared

to unexposed levels (Fig. 4A). As we had seen for bacitracin-induction, deletion of *liaR* again

completely abolished *sapRS* activation, showing the induction of *sapRS* by DAP was occurring through

LiaR regulation. This showed that DAP exposure indeed triggered LiaFSR signalling.

To test how DAP induction was propagated further down the regulatory pathway, we next exposed the

P_{dltX}-*lacZ* transcriptional fusion (SapRS target) to increasing DAP concentrations. Surprisingly, in wild-

type *E. faecalis*, we found daptomycin treatment did not result in any induction of *dlt*, with expression

remaining at basal levels across all tested concentrations (Fig. 4B). As seen for the bacitracin challenge,

deletion of *sapRS* resulted in a decrease in basal expression, from 20 MU to 7.5 MU, and again, there

was no induction of the *dlt* promoter in response to daptomycin exposure. This indicated that although

sapRS transcription was induced in response to daptomycin treatment, remarkably, expression of its

target gene *dlt* remained at basal levels. This discovery suggested a differential response in the

signalling cascade dependent on the inducer, indicating that bacitracin and daptomycin might be

differentially sensed by SapRS, as further explained in the discussion.

The network components vary in importance during interactions with antimicrobial producer

strains. As stated above, we were surprised by the complexity of the regulatory pathway controlling

what in other Firmicutes bacteria is a fairly straightforward response to antibiotic challenge, where each

regulatory system controls its own resistance genes. To shed some light on the reasons for the

complexity of signalling in enterococci, we considered the environments these bacteria can be found

in. *E. faecalis* is a common member of many natural microbial communities, such as in soil and water

or the gastrointestinal tract of humans and animals. There, the enterococci reside within the small and

large intestine and represent up to 1% of the faecal flora[51–54]. In such environments, *E. faecalis* interacts

with other microbes and must defend itself against antimicrobial producing bacteria. Therefore, we next

wanted to assess the role of the individual components of the resistance network in protecting *E. faecalis*

from antimicrobial activity produced by potential competitor bacteria. To do this, we utilised deferred

antagonism assays to simulate relevant environmental pressures from other microbes the enterococci

may encounter. We used four antimicrobial producing strains: the subtilin producer *Bacillus subtilis*

ATCC6633[55], bacitracin producer *Bacillus lichenformis* ATCC10716[56], nisin-A producer *Lactococcus*

lactis NZ9000[57] and nisin-P producer *Streptococcus gallolyticus* AB39[58]. Each producer was spotted

onto a plate, and the antimicrobial was allowed to accumulate over multiple days. This was followed

by the addition of an overlay containing the different *E. faecalis* strains to assess their sensitivity against

the produced compounds based on the size of the zone of inhibition (Fig. 5).

Firstly, we tested the *E. faecalis* strains against the antimicrobial subtilin, produced by *B. subtilis*

ATCC6633. Of the components under investigation in this study, the LiaFSR system appeared to

contribute most strongly to resistance against subtilin, as the Δ*liaR* mutant displayed the largest increase

in zone of inhibition compared to the wild type. The deletions of *sapR*, *sapAB* and *rapAB* also displayed

an increased zone of inhibition compared to the wild type, but to a lesser extent than Δ*liaR*. This

therefore suggests they play a lesser role in subtilin resistance.

In contrast, when testing the *E. faecalis* strains against the bacitracin producer *B. lichenformis*

ATCC10716, deletion of *sapR, sapAB* and *rapAB* showed similar increased sensitivity, consistent with

their contribution to the bacitracin resistance network that we have examined in this study and

previously[33]. Although also presenting increased sensitivity compared with the wild type, deletion of

liaR resulted in a marginally smaller zone of inhibition compared with the other deletion strains. This

is in line with our data presented above, showing the Lia system has a more indirect role in controlling

the bacitracin response by regulating *sapRS* expression.

Figure 5. _E. faecalis_ deletion strains show differential zones of inhibition in deferred antagonism tests against antimicrobials produced by other firmicute bacteria. The antimicrobial producer strains indicated at the top were grown overnight and then adjusted to an OD_{600} of 0.5 with fresh media. Aliquots (5ul) from each producer strain was then spotted and incubated at 25°C for 36-45 hours to allow antimicrobial accumulation. Overnight cultures of the _E. faecalis_ strain indicated on the left were then added as a soft agar overlay. Plates were incubated at 25°C for 24 hours and zones of inhibition in the _E. faecalis_ lawn were used to assess susceptibility to antimicrobials produced by the central strain.

We next tested the _E. faecalis_ mutants against the nisin producers, _L. lactis_ NZ9000 and _S. gallolyticus_

AB39. Although there was some inhibition of wild type _E. faecalis_ by the nisin-A producer _L. lactis_,

there was no difference in sensitivity between the wild type and deletion strains. This suggests either

that the genes of our regulatory pathway do not contribute to resistance against this antibiotic, or that

under the chosen conditions the inhibitory activity of *L. lactis* NZ9000 is primarily due to a different

antimicrobial than nisin-A. In contrast, the nisin-P producer *S. gallolyticus AB39*, despite having no

inhibitory effect on wild type *E. faecalis*, was able to strongly inhibit growth of the *liaR* deletion strain.

Deletions of both *sapR* and *sapAB* also resulted in increased sensitivity but to a lesser degree than *liaR*

deletion. Deletion of *rapAB* had little to no effect compared to the wild type, suggesting that the other

target genes of the regulatory pathway but not the RapAB transporter are responsible for nisin P

resistance.

These findings indicate that in response to different antimicrobials, the various members of this

resistance network have differing roles to play and also differ in their relative importance. In response

to subtilin and nisin-P, LiaR clearly plays a very important role in resistance, but it is a less important

component in response to bacitracin exposure. This reflects LiaR's role as a key regulator in the cell

envelope stress response more widely, but with mostly a moderating role in response to bacitracin. The

sapR, *sapAB* and *rapAB* deletions all present with very similar effects in response to the antimicrobials,

reflecting the interdependent functional relationship between the three components. The notable

exception to this is RapAB in the context of nisin-P, where the transporter does not appear to contribute

to protection of the cell.

Discussion

In this study, we aimed to expand our understanding of the cell envelope stress response of *E. faecalis*

using the previously described bacitracin resistance network as our foundation. Our findings revealed

the regulation of the BceRS-type TCS SapRS via the damage sensor, LiaFSR. In addition, we also

demonstrated the regulation of the resistance operon *dltXABCD* via SapRS. This work describes a direct

functional link between the Lia and Sap signalling systems in *E. faecalis*, which can explain previous

observations of suppressor mutations in *sapAB* when a LiaR-deficient strain was evolved for DAP

resistance[41,42]. Moreover, through creation of deletion strains and analysing promoter fusions, we have

revealed the presence of a differential network response through the damage-sensing LiaFSR and drug-

sensing SapAB-SapRS systems, leading to *dlt* expression in response to bacitracin, but not daptomycin.

This discovery implies there is distinct control on subsections of the network depending on the inducer.

To understand the mechanisms of this differential control, the mode-of-actions of both antibiotics must

be considered. Bacitracin is a peptide antibiotic, which forms a complex with undecaprenyl

pyrophosphate (UPP), an intermediate of the Lipid II cycle[35,36]. The formation of this complex prevents

the dephosphorylation of UPP and slows cell wall formation[59,60]. In contrast, daptomycin is a

lipopeptide antibiotic which forms a tripartite complex between lipid II and the membrane phospholipid

phosphatidyl glycerol[50]. The formation of this complex leads to a loss of membrane potential and cell

death. The similarity between both antibiotics is the formation of cell envelope damage, the known

stimulus for the activation of the "damage sensor" LiaFSR. However, they differ in their mechanism of

action with bacitracin preventing cell wall synthesis and daptomycin directly affecting cell membrane

integrity.

Since both antibiotics cause cell envelope damage, both lead to induction of LiaR target genes (Figures

1&3), including *sapRS*. SapAB, the sensory transporter that controls SapRS activity, however, so far

has been shown to be activated by only two antibiotics, bacitracin and mersacidin [33], both of which bind

to lipid II cycle intermediates[35,61,62]. This is consistent with the current working model for BceAB-type

transporters, to which SapAB belongs, recognising the complex between a peptide antibiotic and a lipid

II cycle intermediate as their substrate[39]. Importantly, the type of antibiotics to which BceAB-type

transporters respond all interact with the lipid II cycle intermediates via at least the pyrophosphate group

(e.g., bacitracin) or the sugar-pyrophosphate moiety of Lipid II (e.g. the lantibiotics, including

mersacidin)[62,63]. In the case of bacitracin, it is therefore the presence of bacitracin-UPP complexes that

triggers SapAB activity, leading to SapRS activation via the transport flux-sensing mechanism

demonstrated for the *B. subtilis* system[34,38]. Thus, the presence of bacitracin triggers both Lia and Sap

signalling simultaneously, inducing the expression of *dlt*.

In contrast, DAP interacts with its cellular target in a fundamentally different way, forming a tripartite

complex between Lipid II and phosphatidyl glycerol (PG)[50]. It therefore would not be expected to form

the type of complexes that could act as substrate for SapAB[39], and Sap signalling is not activated.

BceAB-type transporters have been shown to repress histidine kinase activity in the absence of

substrate[38]. Thus, the lack of a SapAB substrate likely leads to the transporter maintaining SapS in an

'OFF' conformation, preventing activation of SapR target genes, i.e. *dlt*. Such a mechanism can explain

why in the presence of DAP, when the LiaFSR system activates expression of the *sapRS* operon, no

induction of *dlt* expression was observed, because SapAB – unable to detect the tripartite DAP/lipid

II/PG complex - would have prevented SapRS signalling.

Taken together, these findings indicate the presence of a logic 'AND' gate, whereby two inputs (the

damage sensing input provided by LiaFSR and the drug-sensing input provided by SapAB) are needed

to obtain the output, which is the induction of *dlt* expression (Figure 6). This mode of regulation offers

an explanation for the mutations observed previously in *sapAB* when evolving for DAP resistance in

the absence of *liaR*[41,42]. A gain-of-function mutation in *sapAB* would allow the activation of Sap

signalling in the absence of a substrate and induce *dlt* expression, essentially overriding the 'AND' gate

and providing DAP resistance. This working model also explains the low but statistically significant

induction of *dlt* expression by bacitracin we observed in the absence of *sapAB* (Fig. 3). The mutant still

possesses SapRS and thus bacitracin-challenge leads to activation of *sapRS* expression via LiaR. This

in turn is likely to cause an increase in basal activity of SapRS-signalling and thus moderate *dlt*

activation, even if the full activation via SapAB is missing. A similar effect has been described

previously for the second known SapRS target, *rapAB*, where deletion of *sapAB* resulted in low

promoter activity of *rapAB*, but with a small level of residual bacitracin response remaining[33].

The regulatory wiring we have uncovered here between the LiaFSR and the Bce-type SapAB-SapRS

systems, resulting in the control of the *dltXABCD* and *rapAB* resistance operons, is remarkably complex

compared with other Firmicutes bacteria. For example, in *B. subtilis* LiaFSR controls its own regulon,

comprised solely of *liaIH(G)* and its own encoding genes [24]. In *S. aureus*, the LiaFSR homologue VraRS

has been shown to regulate the expression of multiple genes involved in cell wall biosynthesis[64], but

has not been observed to regulate a BceRS-type TCS. Bce-type systems in most Firmicutes bacteria

either solely control their own transporter, as found with BceAB-RS in *B. subtilis*[40], or can be involved

with other Bce-type systems, such as the BraRS-VraDE[65] and GraRS-VraFG[66] systems in *S. aureus*. In

some cases, Bce-type systems can also control additional genes, for example the GraRS TCS has also been shown to regulate the *dlt* operon[67–69]. However, to our knowledge there have been no prior reports of direct regulatory relationships that hardwire the Lia and Bce-type pathways together.

Figure 6. Proposed model of the antibiotic resistance network. Schematic illustrating the genes and proteins involved in antibiotic detection and resistance, with transcriptional regulation and translation demonstrated by black and red arrows respectively. Exposure to bacitracin (BAC) causes damage to the cell envelope, which acts as the stimulus for the removal of repression by LiaF, triggering the activation of LiaS. LiaS activity leads to phosphorylation of LiaR, which in turn induces the expression of its target promoters: its own, *liaXYZ* and *sapRS*, indicated by the black arrows. Simultaneously, bacitracin is also detected by SapAB. Communication between SapAB and SapS is indicated by a faded arrow, resulting in the activation of SapS and phosphorylation of SapR. SapR then induces the expression of the resistance genes, *rapAB* and *dlt,* which remove bacitracin from UPP and reduce the positive charge of the cell envelope, respectively. Conversely, exposure to daptomycin (DAP) again results in cell envelope damage, which triggers the phosphorylation of LiaR and induces the expression of *sapRS*. However, daptomycin is not recognised by SapAB, therefore SapS is kept in an 'OFF' conformation and SapR is not phosphorylated. SapR is therefore unable to induce the expression of *rapAB* and *dlt*.

Our findings reported here show that although the enterococci possess the same regulators as *S. aureus* and *B. subtilis*, evolution has taken these components down different routes. A potential reasoning behind *E. faecalis* maintaining this intricate setup could be that having *dlt* and *rapAB* under Sap control, and *sapRS* itself under Lia regulation, may provide the cell with a layered response. Under this premise,

when the cell is experiencing a mild bacitracin attack, signalling in the Lia-Sap pathway is likely to be

dominated by the Sap system, which responds to the presence of UPP-bacitracin complexes, even if

those do not yet cause cellular damage. Sap signalling then activates both *rapAB* and *dltXABCD*. Of

these, RapAB provides the main target protection mechanism, especially at lower antibiotic

concentrations, whereas expression of *dlt* is likely of minor importance, based on evidence from *B.*

subtilis[39,45]. This is consistent with our observations in *E. faecalis,* where the strength of *rapAB*

induction is much higher than that of *dlt*: at 4μg/mL bacitracin, *rapAB* expression reaches >100 MU [33],

compared to *dlt*, which reaches a maximum of 25 MU (Fig. 3). However, when the antibiotic threat

begins to exceed the protection level RapAB can provide, damage to the cell occurs and the Lia system

is activated. LiaFSR is then able to boost the signalling cascade by increasing *sapRS* expression. We

see this in the absence of *rapAB*, which mimics an extreme failure of target protection, leading to a

drastic increase in *dlt* expression (Fig. 3). Thus, the regulatory structure between Lia and Sap signalling

in *E. faecalis* allows a second line of defence to be mounted if the primary resistance system is no longer

sufficient to fully protect the cell.

The genes discussed here are all part of the core genome of *E. faecalis* and thus contribute to intrinsic,

rather than acquired drug resistance in this bacterium. To better understand the evolutionary context

that appears to have given rise to the unusually complex regulation of the resistance genes, we

considered the broader scope of this network - was it able to protect the enterococci from antimicrobial

challenge from potential competitors in their natural environments? The importance of the role played

by LiaFSR in the cell envelope stress response was well demonstrated by the antagonism assays, in

which deletion of *liaR* resulted in increased sensitivity against all tested antimicrobial producers, except

for *L. lactis* NZ9000 (nisin-A). In contrast, components of the Sap system only contributed to protection

from the bacitracin producer *B. lichenformis* ATCC10716 and the subtilin producer *B. subtilis*

ATCC6633. The only antimicrobial producer against which neither of the genes appeared to give a

protective effect was the nisin-A producer *L. lactis* NZ9000. However, we cannot currently differentiate

whether this was because *L. lactis* did not produce sufficient nisin-A under the chosen growth conditions

to detect differences, or because *E. faecalis* uses alternative resistance mechanisms against nisin-A not

controlled by either the Lia or Sap systems. Overall, it appears that the regulatory network studied here

plays a key role in protecting *E. faecalis* from antimicrobial peptide producing bacteria that it would

realistically encounter in its natural habitats, offering an explanation for the existence of such a complex

regulatory strategy.

One of our most striking results was that daptomycin challenge, although able to activate Lia signalling,

was unable to activate the Sap system, potential mechanistic reasons for which are explained above.

This was a surprising finding, as *dlt* is known to respond to daptomycin in other Firmicutes, e.g. *S.*

aureus [70–74]. The lack of *dlt* expression in response to DAP may, however, explain the sensitivity to this

antibiotic observed throughout the *Enterococcus* genus and its effectiveness when treating VRE

infections[75]. This is despite the bacteria being in possession of an effective resistance mechanism, except

this is not induced upon DAP challenge. It is then plausible to see why clinical DAP resistance could

result from overriding of the molecular controls investigated here to trigger high *dlt* operon expression,

specifically via mutations in genes encoding the components of the LiaFSR system[19], or experimentally

in *sapAB* as suppressor mutations of a LiaR-deficient strain[41,42]. DAP was introduced for treatment of

enterococci infections in 2003[12] and the first resistant isolates were identified already in 2005[13]. It may

be that the rapid emergence of DAP resistance is in part derived from a dysregulation of *dlt* expression

by such mutations of regulatory components.

Overall, our findings contribute to an increasing understanding of the regulatory network protecting *E.*

faecalis from cell envelope attack and provide insights into the regulatory cascade that is involved in

controlling resistance gene expression in response to cell envelope targeting antibiotics. Our findings

also provide valuable context in which to better understand the emergence of DAP resistance in clinical

environments. Ultimately, a system-wide understanding of antimicrobial resistance regulation may lead

to identification of an 'Achilles' heel' within the network and identify new therapeutic targets.

Experimental Procedures

Bacterial strains and growth conditions. All bacterial strains and plasmids used in this study are listed in Table S1 in the supplementary material. *E. coli* MC1061 was used for cloning with pTCVlac, and strain DH5α was used for all other cloning. *E. coli*, *Bacillus licheniformis* and *Bacillus subtilis* were routinely grown in lysogeny broth (LB) at 37°C with agitation (200 rpm). *Lactococcus lactis* was grown routinely in M17 supplemented with 0.5% lactose at 30°C without agitation. *Enterococcus faecalis and Streptococcus gallolyticus were* grown routinely in brain heart infusion (BHI) at 37°C without agitation, with media for the latter being supplemented with 5% (v/v) foetal bovine calf serum. Solid media contained 15 g l^{-1} agar. Selective media contained chloramphenicol (10 μg ml^{-1} for *E. coli*; 15 μg ml^{-1} for *E. faecalis*), kanamycin (50 μg ml^{-1} for *E. coli*; 1000 μg ml^{-1} for *E. faecalis*), spectinomycin (100 μg ml^{-1} for *E. coli*; 500 μg ml^{-1} for *E. faecalis*). For blue-white screening, 5-bromo-4-chloro-3-indolyl-β-d-galactopyranoside (X-Gal) was used at 120 μg/ml. Bacitracin was supplied as the Zn^{2+} salt. All media for experiments with daptomycin were supplemented with 50 μM CaCl$_2$.

E. faecalis was transformed by electroporation as previously described[76]. *E. coli* was transformed by heat-shock of CaCl$_2$ competent cells, followed by 1 hour recovery time[77]. Growth was measured as optical density at 600 nm (OD$_{600}$) on a the Biochrom™ Novaspec Pro Spectrophotometer using cuvettes with 1 cm light path length or in 96-well plates with 100 μL culture volumes on a Spark®Microplate reader (Tecan).

Construction of plasmids and genetic techniques. All primer sequences used for cloning are listed in Table S2 in the supplementary material. Transcriptional promoter fusions to *lacZ* in *E. faecalis* were constructed in the vector pTCVlac[78]. All fragments were cloned via the EcoRI and BamHI sites of the vector. Constructs for unmarked deletions in *E. faecalis* were cloned into pLT06[79]. For each gene or operon to be deleted, 700- to 1000-bp located immediately before the start codon of the first gene ("up" fragment) and after the stop codon of the last gene ("down" fragment) were amplified. The primers were designed to create a 17- to 20-bp overlap between the PCR products (Table S2), facilitating the fusion of the fragments by PCR overlap extension[80] and were subsequently cloned into the NcoI and BamHI site of the vector pLT06. Gene deletions were performed as previously described[79]. Briefly, following transformation of the parent strain with the temperature-sensitive vector pLT06, overnight

cultures were grown at 30° containing chloramphenicol and reinoculated 1:1000 into 10 mL BHI the

next morning. Cells were then grown at 30° for 2.5 hours, followed by increasing to 42° for a further

2.5 hours to force single-site integration. Cells were then serially diluted onto BHI agar containing

chloramphenicol and X-Gal and incubated at 42°. Blue colonies growing at 42°C were screened for the

targeted integration using PCR with primers flanking the site of integration. Positive clones were then

serially passaged for two days from overnight culture in BHI medium with no selection at 30° to allow

a second site recombination event. Cultures were then serially diluted on to MM9-YEG agar[81]

containing 10 mM *p*-chloro-phenylalanine for counter-selection and X-Gal at 37°C. The resulting white

colonies were screened for the deletion of the target genes by PCR. All cloned constructs were checked

for PCR fidelity by Sanger sequencing, and all created strains were verified by PCR using the primers

given in Table S2.

Antimicrobial susceptibility assays. For antibiotic susceptibility assays, minimum inhibitory

concentrations (MICs) were determined by broth dilution assays in BHI medium in 96-well plates,

containing serial two-fold dilutions of antibiotic. Colonies of *E. faecalis* were suspended in sterile

Phosphate Buffered Saline (PBS) to 0.5 McFarland standard turbidity and diluted 1:1,000 in a total

volume of 100μL. After 24 h incubation at 37°C, growth was determined by measuring optical density

(OD_{600}) on a Spark®Microplate reader (Tecan). The MIC was scored as the lowest antibiotic

concentration where no growth was observed following subtraction of the OD_{600} of a well containing

sterile medium.

β-Galactosidase assays. For quantitatively assessing induction of *lacZ* reporter constructs in *E.*

faecalis, exponentially growing cells (OD_{600} = 0.4-0.5) inoculated 1:250 from overnight cultures in BHI

medium were exposed to different concentrations of bacitracin for 1 h or daptomycin for 30 minutes.

Cells were harvested via centrifugation and stored at -20°C. β-Galactosidase activities were assayed in

permeabilised cells and expressed in Miller units (MU)[82]. For this, cells were resuspended in 1 ml Z-

buffer (8.04 g $Na_2HPO_4*7H_2O$, 2.76 g $NaH_2PO_4*H_2O$, 0.123 g $MgSO_4*7H_2O$ and 5 mL 1M KCl in 495

mL dH_2O, pH 7). The samples were adjusted to OD_{600} = 0.5 in a 1 ml volume of Z-buffer and from this,

two volumes were taken: 200 μl and 400 μl cells made up to 1 mL each with Z-buffer. This volume

corresponds to the 'volume of cells' in the Miller Unit (MU) equation below. Following this, 20 μl

0.1% (w/v) SDS and 40 μl chloroform were added and vortexed for 5 seconds, then rested for 5-10

minutes. Reactions were started by adding 200 μl *o*-nitrophenyl-β-D-galactopyranoside (ONPG) (4 mg

mL^{-1} in Z-buffer) and incubated at room temperature until yellow colouration was observed. If no colour

change was visible, the reaction was incubated for 20 minutes. Reactions were stopped by adding 500

μl 1M Na$_2$CO$_3$, and the time recorded, which corresponds to the 'time' in the Miller Unit (MU) equation

below. Absorbance at 420 nm (A$_{420}$) was then read. MUs were calculated using the following equation:

533
$$Miller\ Units\ (MU) = \frac{A420 * 1000}{Time\ (minutes) * volume\ of\ cells\ (in\ ml) * OD600}$$

Computational modelling. Rather than directly modelling the temporal dynamics of the regulatory

network as done for the *B. subtilis* Bce system[34,83], we chose to focus on the (meta) stable state reached

in response to challenge with a given bacitracin concentration. The quantities modelled are the

concentrations of bacitracin [*bac*], UPP [*UPP*], UPP-bound bacitracin [*UPPbac*], and the effective

activities of SapRS [*SapRS*], RapAB [*RapAB*], LiaX [*Lia*], and DltXABCD [*Dlt*]. For the modelling,

we made repeated use of a soft-switch sigmoid type function

540
$$f_y(x; y_{base}, y_{max}, x_0, k) = f_{max} \left(1 + \left(\frac{y_{max}}{y_{base}} - 1 \right) e^{-k(x-x_0)} \right)^{-1},$$

541 where x is the input quantity and y the output which varies between y_{base} and y_{max}, with x_0 controlling

542 the switching threshold and k setting the sharpness of the transition. The equations of state for the model

543 are then simply

544
$$k_{on}([Dlt])([bac] - [UPPbac])[UPP] - k_{off}([RapAB])[UPPbac] = 0$$

545
$$[Lia] = f_{Lia}([UPPbac])$$

546
$$[SapRS] = f_{SapRS}([Lia])$$

547
$$[Dlt] = f_{dlt}([SapRS])$$

548
$$[RapAB] = f_{RapAB}([SapRS])$$

We modelled the binding and off rates for the UPP / bacitracin interaction as being linearly proportional

to [*Dlt*] and [*RapAB*], respectively, with Dlt reducing bacitracin binding and RapAB increasing the off-

rate. The biological reasoning for this is explained in the Results section.

For [*bac*], actual values from the experimental work were used, and the switching parameters for [*Dlt*]

and [*RapAB*] as functions of [*SapRS*] were determined by fitting actual activity levels observed

experimentally of these quantities for different bacitracin levels. The remaining parameters were fitted

to achieve a description of experimental results for the wild-type strain by the model output. The values

for each parameter are given in Table S3. Solving the equations of state for a given input bacitracin

level [*bac*] yielded predictions for [*UPP*] and [*UPPbac*], which in turn drive the response curves plotted

in Figure 3B.

Antagonism assays on solid media. A plate-based assay was utilised to measure growth inhibition

between antibiotic producer strains and *E. faecalis* strains. For each species, cultures of the

antimicrobial-producing bacteria were grown overnight in the respective growth medium and

temperatures stated above and adjusted to $OD_{600} = 0.5$. Aliquots (5 µl) from each producer culture were

spotted onto the centre of a BHI agar plate and incubated at room temperature (20-25°C) for 3-5 days

to allow accumulation of antimicrobial products. Overnight cultures of *E. faecalis* strains were grown

at 37°C in BHI medium, inoculated 1:100 into fresh medium and grown to OD = 0.5. Next, 3 mL of

liquid BHI soft agar (7.5 g/L, 50°C) were inoculated with 30 µL culture. The soft agar was then poured

onto the plate containing the pre-grown antimicrobial-producing strains and left to dry. The plate was

then incubated overnight at room temperature to allow *E. faecalis* to grow to visualise the zone of

inhibition. Results were recorded photographically using a PowerShot G camera attached to a lightbox.

Data accessibility. The numerical data underpinning the results shown in Figures 1, 3 and 4 are

available in Table S4. The *Mathematica* file for the model is provided in the supplementary material.

Acknowledgements

This work was supported in part by grant MR/N0137941/1 for the GW4 BIOMED MRC DTP, awarded

to the Universities of Bath, Bristol, Cardiff and Exeter from the Medical Research Council

(MRC)/UKRI. The authors gratefully acknowledge the Technical Staff within the Life Sciences

Department at the University of Bath for technical support and assistance in this work. We would also

like to thank Mathew Upton for the gift of *Streptococcus gallolyticus*.

Author contributions

581 SMM, SG, TR and GF contributed to the conception of the study; SMM and SG designed the

582 experimental work; SMM carried out all data acquisition; SMM, TR and SG carried out data analysis

583 and interpretation; SMM, SG, TR and GF wrote the manuscript.

Graphical abstract

Abbreviated summary

588 This study explored a regulatory network controlling resistance against daptomycin and bacitracin in

589 *E. faecalis*. We show that the *dltXABCD* operon, which can protect against both antibiotics, is

590 controlled by two regulatory systems, LiaFSR and SapRS. Strikingly, this strategy allows *dtlXABCD*

591 expression only in response to bacitracin, but not daptomycin, potentially explaining the natural DAP-

592 sensitivity of *E. faecalis* and why clinical resistance can emerge via regulatory mutations.

References

1.   Organisation for Economic Co-operation and Development. Antimicrobial Resistance.

Tackling the Burden in the European Union. *Eur Cent Dis Prev Control*. 2020:1-20.

https://www.oecd.org/health/health-systems/AMR-Tackling-the-Burden-in-the-EU-OECD-

ECDC-Briefing-Note-2019.pdf.

2.   Zhen X, Lundborg CS, Sun X, Hu X, Dong H. Economic burden of antibiotic resistance in

ESKAPE organisms: a systematic review. *Antimicrob Resist Infect Control*. 2019;8:137.

doi:10.1186/s13756-019-0590-7

3.   Miller WR, Murray BE, Rice LB, Arias CA. Vancomycin-Resistant Enterococci: Therapeutic

Challenges in the 21st Century. *Infect Dis Clin North Am*. 2016;30(2):415-439.

doi:10.1016/j.idc.2016.02.006

4.   Chiang H-Y, Perencevich EN, Nair R, *et al*. Incidence and Outcomes Associated With

Infections Caused by Vancomycin-Resistant Enterococci in the United States: Systematic

Literature Review and Meta-Analysis. *Infect Control Hosp Epidemiol*. 2017;38(2):203-215.

doi:10.1017/ice.2016.254

5.   Said MS, Tirthani E, Lesho E. *Enterococcus Infections*. Rochester General Hospital: StatPearls

Publishing, Treasure Island (FL); 2021. http://europepmc.org/abstract/MED/33620836.

6.   Arias CA, Murray BE. The rise of the Enterococcus: Beyond vancomycin resistance. *Nat Rev*

Microbiol. 2012;10(4):266-278. doi:10.1038/nrmicro2761

7.   Leclercq R, Derlot E, Duval J, Courvalin P. Plasmid-Mediated Resistance to Vancomycin and

Teicoplanin in *Enterococcus faecium*. *N Engl J Med*. 1988;319(3):157-161.

doi:10.1056/NEJM198807213190307

8.   Uttley AHC, George RC, Naidoo J, *et al*. High-level vancomycin-resistant enterococci causing

hospital infections. *Epidemiol Infect*. 1989;103(1):173-181. doi:DOI:

10.1017/S0950268800030478

9.   Levine DP. Vancomycin: A History. *Clin Infect Dis*. 2006;42(Supplement_1):S5-S12.

doi:10.1086/491709

10.   Courvalin P. Vancomycin Resistance in Gram-Positive Cocci. *Clin Infect Dis*.

2006;42(Supplement_1):S25-S34. doi:10.1086/491711

11.   Puchter L, Chaberny IF, Schwab F, Vonberg R-P, Bange F-C, Ebadi E. Economic burden of

nosocomial infections caused by vancomycin-resistant enterococci. *Antimicrob Resist Infect*

Control. 2018;7(1):1. doi:10.1186/s13756-017-0291-z

12.   Sauermann R, Rothenburger M, Graninger W, Joukhadar C. Daptomycin: A Review 4 Years

after First Approval. *Pharmacology*. 2008;81(2):79-91. doi:10.1159/000109868

13.   Munoz-Price LS, Lolans K, Quinn JP. Emergence of Resistance to Daptomycin during

Treatment of Vancomycin-Resistant *Enterococcus faecalis* Infection. *Clin Infect Dis*.

2005;41(4):565-566. doi:10.1086/432121

14.   Arias CA, Panesso D, McGrath DM, *et al*. Genetic Basis for In Vivo Daptomycin Resistance

in Enterococci. *N Engl J Med*. 2011;365(10):892-900. doi:10.1056/NEJMoa1011138

15.   Tran T, Panesso D, Mishra Nagendra N, *et al*. Daptomycin-Resistant *Enterococcus faecalis*

Diverts the Antibiotic Molecule from the Division Septum and Remodels Cell Membrane

Phospholipids. *MBio*. 2013;4(4):e00281-13. doi:10.1128/mBio.00281-13

16.   Reyes J, Panesso D, Tran TT, *et al*. A *liaR* Deletion Restores Susceptibility to Daptomycin and

Antimicrobial Peptides in Multidrug-Resistant *Enterococcus faecalis*. *J Infect Dis*.

2015;211(8):1317-1325. doi:10.1093/infdis/jiu602

17.   Palmer K, Daniel A, Hardy C, Silverman J, Gilmore M. Genetic Basis for Daptomycin

Resistance in Enterococci. *Antimicrob Agents Chemother*. 2011;55(7):3345-3356.

doi:10.1128/AAC.00207-11

18.   Tran TT, Miller WR, Shamoo Y, Arias CA. Targeting cell membrane adaptation as a novel

antimicrobial strategy. *Curr Opin Microbiol*. 2016;33:91-96.

doi:https://doi.org/10.1016/j.mib.2016.07.002

19.   Miller C, Kong J, Tran TT, Arias CA, Saxer G, Shamoo Y. Adaptation of *Enterococcus*

faecalis to daptomycin reveals an ordered progression to resistance. *Antimicrob Agents*

Chemother. 2013;57(11):5373-5383. doi:10.1128/AAC.01473-13

20.   Jordan S, Junker A, Helmann J, Mascher T. Regulation of LiaRS-Dependent Gene Expression

in *Bacillus subtilis*: Identification of Inhibitor Proteins, Regulator Binding Sites, and Target

Genes of a Conserved Cell Envelope Stress-Sensing Two-Component System. *J Bacteriol*.

2006;188(14):5153-5166. doi:10.1128/JB.00310-06

21.   Kuroda M, Kuroda H, Oshima T, Takeuchi F, Mori H, Hiramatsu K. Two-component system

VraSR positively modulates the regulation of cell-wall biosynthesis pathway in

Staphylococcus aureus. *Mol Microbiol*. 2003;49(3):807-821.

doi:https://doi.org/10.1046/j.1365-2958.2003.03599.x

22.   Martínez B, Zomer AL, Rodríguez A, Kok J, Kuipers OP. Cell envelope stress induced by the

bacteriocin Lcn972 is sensed by the lactococcal two-component system CesSR. *Mol*

Microbiol. 2007;64(2):473-486. doi:https://doi.org/10.1111/j.1365-2958.2007.05668.x

23.   Suntharalingam P, Senadheera MD, Mair RW, Levesque CM, Cvitkovitch DG. The LiaFSR

system regulates the cell envelope stress response in *Streptococcus mutans*. *J Bacteriol*.

2009;191(9):2973-2984. doi:10.1128/JB.01563-08

24.   Wolf D, Kalamorz F, Wecke T, *et al*. In-Depth Profiling of the LiaR Response of *Bacillus*

subtilis. *J Bacteriol*. 2010;192(18):4680-4693. doi:10.1128/JB.00543-10

25.   Schrecke K, Jordan S, Mascher T. Stoichiometry and perturbation studies of the LiaFSR

system of *Bacillus subtilis*. *Mol Microbiol*. 2013;87(4):769-788.

doi:https://doi.org/10.1111/mmi.12130

26.   Wolf D, Domínguez-Cuevas P, Daniel R, Mascher T. Cell Envelope Stress Response in Cell

Wall-Deficient L-Forms of *Bacillus subtilis*. *Antimicrob Agents Chemother*.

2012;56(11):5907-5915. doi:10.1128/AAC.00770-12

27. Khan A, Davlieva M, Panesso D, *et al*. Antimicrobial sensing coupled with cell membrane

remodeling mediates antibiotic resistance and virulence in *Enterococcus faecalis*. *Proc Natl*

Acad Sci. 2019;116(52):26925-26932. doi:10.1073/pnas.1916037116

28. Comenge Y, Quintiliani R, Li L, *et al*. The CroRS Two-Component Regulatory System Is

Required for Intrinsic β-Lactam Resistance in *Enterococcus faecalis*. *J Bacteriol*.

2003;185(24):7184-7192. doi:10.1128/JB.185.24.7184-7192.2003

29. Winkler ME, Hoch JA. Essentiality, bypass, and targeting of the YycFG (VicRK) two-

component regulatory system in gram-positive bacteria. *J Bacteriol*. 2008;190(8):2645-2648.

doi:10.1128/JB.01682-07

30. Kristich C, Wells C, Dunny G. A eukaryotic-type Ser/Thr kinase in *Enterococcus faecalis*

mediates antimicrobial resistance and intestinal persistence. *Proc Natl Acad Sci*.

2007;104(9):3508-3513. doi:10.1073/pnas.0608742104

31. Kristich C, Little J, Hall C, Hoff J, Low D. Reciprocal Regulation of Cephalosporin Resistance

in *Enterococcus faecalis*. *MBio*. 2011;2(6):e00199-11. doi:10.1128/mBio.00199-11

32. Iannetta AA, Minton NE, Uitenbroek AA, *et al*. IreK-Mediated, Cell Wall-Protective

Phosphorylation in *Enterococcus faecalis*. *J Proteome Res*. 2021;20(11):5131-5144.

doi:10.1021/acs.jproteome.1c00635

33. Gebhard S, Fang C, Shaaly A, *et al*. Identification and characterization of a bacitracin

resistance network in *Enterococcus faecalis*. *Antimicrob Agents Chemother*. 2014.

doi:10.1128/AAC.02111-13

34. Fritz G, Dintner S, Treichel SN, *et al*. A New Way of Sensing: Need-Based Activation of

Antibiotic Resistance by a Flux-Sensing Mechanism. *MBio*. 2015;6(4):e00975-15.

doi:10.1128/mBio.00975-15

35. Storm DR. Mechanism of Bacitracin Action: A Specific Lipid-Peptide Interaction. *Ann N Y*

Acad Sci. 1974;235(1):387-398. doi:https://doi.org/10.1111/j.1749-6632.1974.tb43278.x

36.   Storm DR, Strominger JL. Binding of bacitracin to cells and protoplasts of Micrococcus

lysodeikticus. *J Biol Chem*. 1974;249(6):1823-1827. doi:10.1016/s0021-9258(19)42861-5

37.   Dintner S, Heermann R, Fang C, Jung K, Gebhard S. A sensory complex consisting of an

ATP-binding cassette transporter and a two-component regulatory system controls bacitracin

resistance in *Bacillus subtilis*. *J Biol Chem*. 2014;289(40):27899-27910.

doi:10.1074/jbc.M114.596221

38.   Koh A, Gibbon MJ, Van der Kamp MW, Pudney CR, Gebhard S. Conformation control of the

histidine kinase BceS of *Bacillus subtilis* by its cognate ABC-transporter facilitates need-based

activation of antibiotic resistance. *Mol Microbiol*. 2021;115(1):157-174.

doi:https://doi.org/10.1111/mmi.14607

39.   Kobras C, Piepenbreier H, Emenegger J, Sim A, Fritz G, Gebhard S. BceAB-Type Antibiotic

Resistance Transporters Appear To Act by Target Protection of Cell Wall Synthesis.

Antimicrob Agents Chemother. 2020;64(3):e02241-19. doi:10.1128/AAC.02241-19

40.   Ohki R, Giyanto, Tateno K, *et al*. The BceRS two-component regulatory system induces

expression of the bacitracin transporter, BceAB, in *Bacillus subtilis*. *Mol Microbiol*.

2003;49(4):1135-1144. doi:https://doi.org/10.1046/j.1365-2958.2003.03653.x

41.   Prater A, Mehta H, Kosgei A, *et al*. Environment Shapes the Accessible Daptomycin

Resistance Mechanisms in *Enterococcus faecium*. *Antimicrob Agents Chemother*.

2019;63(10):e00790-19. doi:10.1128/AAC.00790-19

42.   Prater A, Mehta H, Beabout K, *et al*. Daptomycin Resistance in *Enterococcus faecium* Can Be

Delayed by Disruption of the LiaFSR Stress Response Pathway. *Antimicrob Agents*

Chemother. 2021;65(4):e01317-20. doi:10.1128/AAC.01317-20

43.   Mascher T, Margulis NG, Wang T, Ye RW, Helmann JD. Cell wall stress responses in

Bacillus subtilis: the regulatory network of the bacitracin stimulon. *Mol Microbiol*.

718   2003;50(5):1591-1604. doi:https://doi.org/10.1046/j.1365-2958.2003.03786.x

719 44. Rietkötter E, Hoyer D, Mascher T. Bacitracin sensing in *Bacillus subtilis*. *Mol Microbiol*.

720   2008;68(3):768-785. doi:https://doi.org/10.1111/j.1365-2958.2008.06194.x

721 45. Radeck J, Gebhard S, Orchard PS, *et al*. Anatomy of the bacitracin resistance network in

722   *Bacillus subtilis*. *Mol Microbiol*. 2016;100(4):607-620. doi:10.1111/mmi.13336

723 46. Peschel A, Otto M, Jack RW, Kalbacher H, Jung G, Götz F. Inactivation of the *dlt* Operon in

724   *Staphylococcus aureus* Confers Sensitivity to Defensins, Protegrins, and Other Antimicrobial

725   Peptides. *J Biol Chem*. 1999;274(13):8405-8410. doi:10.1074/jbc.274.13.8405

726 47. Tran T, Munita J, Arias C. Mechanisms of drug resistance: daptomycin resistance. *Ann N Y*

727   *Acad Sci*. 2015;1354(1):32-53. doi:https://doi.org/10.1111/nyas.12948

728 48. Wecke T, Zühlke D, Mäder U, *et al*. Daptomycin versus Friulimicin B: In-Depth Profiling of

729   *Bacillus subtilis* Cell Envelope Stress Responses. *Antimicrob Agents Chemother*.

730   2009;53(4):1619-1623. doi:10.1128/AAC.01046-08

731 49. Hachmann A., Angert E, Helmann J. Genetic Analysis of Factors Affecting Susceptibility of

732   *Bacillus subtilis* to Daptomycin. *Antimicrob Agents Chemother*. 2009;53(4):1598-1609.

733   doi:10.1128/AAC.01329-08

734 50. Grein F, Müller A, Scherer KM, *et al*. Ca^{2+}-Daptomycin targets cell wall biosynthesis by

735   forming a tripartite complex with undecaprenyl-coupled intermediates and membrane lipids.

736   *Nat Commun*. 2020;11(1):1455. doi:10.1038/s41467-020-15257-1

737 51. Tendolkar PM, Baghdayan AS, Shankar N. Pathogenic enterococci: new developments in the

738   21st century. *Cell Mol Life Sci C*. 2003;60(12):2622-2636. doi:10.1007/s00018-003-3138-0

739 52. Eckburg P, Bik E, Charles B, *et al*. Diversity of the Human Intestinal Microbial Flora. *Science*

740   *(80-)*. 2005;308(5728):1635-1638. doi:10.1126/science.1110591

741 53. Sghir A, Gramet G, Suau A, Rochet V, Pochart P, Dore J. Quantification of Bacterial Groups

within Human Fecal Flora by Oligonucleotide Probe Hybridization. *Appl Environ Microbiol*.

2000;66(5):2263-2266. doi:10.1128/AEM.66.5.2263-2266.2000

54.   Lorian V. Colonization resistance. *Antimicrob Agents Chemother*. 1994;38(7):1693.

doi:10.1128/AAC.38.7.1693

55.   Banerjee S, Hansen JN. Structure and expression of a gene encoding the precursor of subtilin,

a small protein antibiotic. *J Biol Chem*. 1988;263(19):9508-9514.

doi:https://doi.org/10.1016/S0021-9258(19)76571-5

56.   Johnson B, Anker H, Meleney F. Bacitracin: A New Antibiotic Produced by a Member of the

B. subtilis Group. *Science (80-)*. 1945;102(2650):376-377. doi:10.1126/science.102.2650.376

57.   Kuipers OP, de Ruyter PGGA, Kleerebezem M, de Vos WM. Quorum sensing-controlled gene

expression in lactic acid bacteria. *J Biotechnol*. 1998;64(1):15-21.

doi:https://doi.org/10.1016/S0168-1656(98)00100-X

58.   Aldarhami A, Felek A, Sharma V, Upton M. Purification and characterization of nisin P

produced by a strain of *Streptococcus gallolyticus*. *J Med Microbiol*. 2020;69(4):605-616.

doi:https://doi.org/10.1099/jmm.0.001170

59.   Storm DR, Strominger JL. Complex Formation between Bacitracin Peptides and Isoprenyl

Pyrophosphates. *J Biol Chem*. 1973;248(11):3940-3945. doi:10.1016/s0021-9258(19)43823-4

60.   Economou N, Cocklin S, Loll P. High-resolution crystal structure reveals molecular details of

target recognition by bacitracin. *Proc Natl Acad Sci*. 2013;110(35):14207-14212.

doi:10.1073/pnas.1308268110

61.   Cudic P, Kranz J, Behenna D, *et al*. Complexation of peptidoglycan intermediates by the

lipoglycodepsipeptide antibiotic ramoplanin: Minimal structural requirements for

intermolecular complexation and fibril formation. *Proc Natl Acad Sci*. 2002;99(11):7384-

7389. doi:10.1073/pnas.102192099

62.   Staroń A, Finkeisen DE, Mascher T. Peptide antibiotic sensing and detoxification modules of

767  *Bacillus subtilis. Antimicrob Agents Chemother*. 2011;55(2):515-525.

768  doi:10.1128/AAC.00352-10

769 63. Gebhard S, Mascher T. Antimicrobial peptide sensing and detoxification modules: Unravelling

770  the regulatory circuitry of *Staphylococcus aureus. Mol Microbiol*. 2011;81(3):581-587.

771  doi:10.1111/j.1365-2958.2011.07747.x

772 64. Wu S, Lin K, Liu Y, Zhang H, Lei L. Two-component signaling pathways modulate drug

773  resistance of *Staphylococcus aureus* (Review). *Biomed Rep*. 2020;13(2):5.

774  doi:10.3892/br.2020.1312

775 65. Arii K, Kawada-Matsuo M, Oogai Y, Noguchi K, Komatsuzawa H. Single mutations in BraRS

776  confer high resistance against nisin A in *Staphylococcus aureus. Microbiologyopen*.

777  2019;8(11):e791. doi:https://doi.org/10.1002/mbo3.791

778 66. Meehl M, Herbert S, Götz F, Cheun A. Interaction of the GraRS Two-Component System with

779  the VraFG ABC Transporter To Support Vancomycin-Intermediate Resistance in

780  *Staphylococcus aureus. Antimicrob Agents Chemother*. 2007;51(8):2679-2689.

781  doi:10.1128/AAC.00209-07

782 67. Ledger EVK, Mesnage S, Edwards AM. Human serum triggers antibiotic tolerance in

783  *Staphylococcus aureus. Nat Commun*. 2022;13(1):2041. doi:10.1038/s41467-022-29717-3

784 68. Cheung AL, Bayer AS, Yeaman MR, *et al*. Site-specific mutation of the sensor kinase *graS* in

785  *Staphylococcus aureus* alters the adaptive response to distinct cationic antimicrobial peptides.

786  *Infect Immun*. 2014;82(12):5336-5345. doi:10.1128/IAI.02480-14

787 69. Kraus D, Herbert S, Kristian SA, *et al*. The GraRS regulatory system controls *Staphylococcus*

788  *aureus* susceptibility to antimicrobial host defenses. *BMC Microbiol*. 2008;8(1):85.

789  doi:10.1186/1471-2180-8-85

790 70. Bayer AS, Mishra NN, Cheung AL, Rubio A, Yang S-J. Dysregulation of *mprF* and *dltABCD*

791  expression among daptomycin-non-susceptible MRSA clinical isolates. *J Antimicrob*

Chemother. 2016;71(8):2100-2104. doi:10.1093/jac/dkw142

71.   Yang S, Kreiswirth BN, Sakoulas G, *et al*. Enhanced Expression of *dltABCD* Is Associated

with the Development of Daptomycin Nonsusceptibility in a Clinical Endocarditis Isolate of

Staphylococcus aureus. *J Infect Dis*. 2009;200(12):1916-1920. doi:10.1086/648473

72.   Cafiso V, Bertuccio T, Purrello S, *et al*. *dltA* overexpression: A strain-independent keystone of

daptomycin resistance in methicillin-resistant *Staphylococcus aureus*. *Int J Antimicrob Agents*.

2014;43(1):26-31. doi:https://doi.org/10.1016/j.ijantimicag.2013.10.001

73.   Mishra NN, Bayer AS, Weidenmaier C, *et al*. Phenotypic and Genotypic Characterization of

Daptomycin-Resistant Methicillin-Resistant *Staphylococcus aureus* Strains: Relative Roles of

mprF and *dlt* Operons. *PLoS One*. 2014;9(9):e107426.

https://doi.org/10.1371/journal.pone.0107426.

74.   Bertsche U, Weidenmaier C, Kuehner D, *et al*. Correlation of Daptomycin Resistance in a

Clinical *Staphylococcus aureus* Strain with Increased Cell Wall Teichoic Acid Production and

d-Alanylation. *Antimicrob Agents Chemother*. 2011;55(8):3922-3928.

doi:10.1128/AAC.01226-10

75.   Turnidge J, Kahlmeter G, Cantón R, MacGowan A, Giske CG. Daptomycin in the treatment of

enterococcal bloodstream infections and endocarditis: a EUCAST position paper. *Clin*

Microbiol Infect. 2020;26(8):1039-1043. doi:10.1016/j.cmi.2020.04.027

76.   Cruz-Rodz AL, Gilmore MS. High efficiency introduction of plasmid DNA into glycine

treated *Enterococcus faecalis* by electroporation. *Mol Gen Genet MGG*. 1990;224(1):152-154.

doi:10.1007/BF00259462

77.   Hanahan D, Glover DM. DNA cloning: a practical approach. *DNA cloning a Pract approach*.

1985;1:109-135.

78.   Poyart C, Trieu-Cuot P. A broad-host-range mobilizable shuttle vector for the construction of

transcriptional fusions to β-galactosidase in Gram-positive bacteria. *FEMS Microbiol Lett*.

1997;156(2):193-198. doi:10.1111/j.1574-6968.1997.tb12726.x

79.   Thurlow L, Chittezham TV, Hancock L. Capsular Polysaccharide Production in *Enterococcus*

faecalis and Contribution of CpsF to Capsule Serospecificity. *J Bacteriol*. 2009;191(20):6203-

6210. doi:10.1128/JB.00592-09

80.   Ho SN, Hunt HD, Horton RM, Pullen JK, Pease LR. Site-directed mutagenesis by overlap

extension using the polymerase chain reaction. *Gene*. 1989;77(1):51-59.

doi:https://doi.org/10.1016/0378-1119(89)90358-2

81.   Kristich CJ, Chandler JR, Dunny GM. Development of a host-genotype-independent

counterselectable marker and a high-frequency conjugative delivery system and their use in

genetic analysis of *Enterococcus faecalis*. *Plasmid*. 2007;57(2):131-144.

doi:https://doi.org/10.1016/j.plasmid.2006.08.003

82.   Gebhard S, Tran SL, Cook GM. The Phn system of Mycobacterium smegmatis: a second high-

affinity ABC-transporter for phosphate. *Microbiology*. 2006;152(11):3453-3465.

doi:https://doi.org/10.1099/mic.0.29201-0

83.   Piepenbreier H, Sim A, Kobras CM, *et al*. From Modules to Networks: a Systems-Level

Analysis of the Bacitracin Stress Response in *Bacillus subtilis*. *mSystems*. 2020;5(1):e00687-

19. doi:10.1128/mSystems.00687-19

84.   Allen D. Transcriptional regulation of the *dlt* operon in *Enterococcus faecalis* and further

characterization of a *dltA* mutant. 2008.

Supplementary Material

838   **Table S1.** Vectors, plasmids and strains used in this study.

839   **Table S2.** Primers used in this study.

840   **Table S3.** Parameter values used in the mathematical model.

841   **Table S4.** Experimental numerical data for Figures 1, 3 and 4.

842 **File S1.** Mathematical model file for the application *Mathematica*