1 **Root Walker: an automated pipeline for large scale quantification of early root growth responses at**
2 **high spatial and temporal resolution**
3

Platre Matthieu Pierre[1,♩,*], Halvorson Zachary[1,*], Mehta Preyanka[1], Brent Lukas[1], Gleason F. Matias[1], Faizi
Kian[1], Busch Wolfgang[1,♩].
[*]author contributed equally
[♩]Correspondence: mplatre@salk.edu, wbusch@salk.edu

[1] Salk Institute for Biological Studies, Plant Molecular And Cellular Biology Laboratory, 10010 N Torrey Pines Rd, La
Jolla, CA 92037, USA.

10 Summary

Plants are sessile organisms that constantly need to adapt to their changing environment. The root is
exposed to numerous environmental signals ranging from nutrients and water to microbial molecular
patterns. These signals can trigger distinct responses including the rapid increase or decrease of root
growth. Consequently, using root growth as a read out for signal perception allows for deciphering which
cues root senses and how they integrate it. To date, studies measuring root growth responses using large
numbers of roots are often limited either by the lack of high-throughput image acquisition, scalable
analysis methods, and by low spatiotemporal resolution. We developed the "Root Walker" pipeline, which
utilizes automated microscopes to acquire images of many roots exposed to controlled treatments with
high-spatiotemporal-resolution in conjunction with a fast and automated image analysis software. We
demonstrate the power of this pipeline by quantifying fast root growth rate responses in the order of
minutes upon treatments with natural auxin, and by quantifying slower root growth rate responses in the
order of hours upon treatment with two mitogen-associated protein kinase cascade inhibitors. We find a
concentration dependent root growth response to auxin and expose the specificity of one MAPK inhibitor
for affecting root growth. Overall, the "Root Walker" toolkit is a relevant tool for the plant community to
conduct large-scale screening of root growth changes for a variety of purposes including genetic screens
for root sensing and response mechanisms.

28
29

30 Introduction

The root is an organ that is involved in a plethora of functions such as anchoring the plant into the soil and
nutrient and water uptake. However, resources are not evenly distributed in the soil. Moreover, plants
interact with beneficial and pathogenic microorganisms which can promote, limit resources, or pose a
threat. It is therefore not surprising that roots can sense and respond to many different cues, ranging from
physical cues such as gravity and touch to chemical cues such as water, nutrients or microbially associated
molecules (Morris *et al.*, 2017). Root responses to the environment are almost always resulting in a change
of root growth. These responses are key for plant adaptation and over time shape the root system
architecture (RSA). During the last decade, the root growth has been successfully used to novel genes and

molecular mechanisms that underlie root responses to the environment (Satbhai *et al.*, 2017; Li *et al.*,
2019; Platre *et al.*, 2022). However, these studies have utilized screens on relatively late read outs of root
growth change, corresponding to days after exposure to distinct growth environments. Such approaches
might not be very powerful to identify sensory mechanisms that are still not yet identified for many
environmental cues such as nutrients (Jia, Giehl and von Wirén, 2022). We hypothesized that it might be
more powerful to measure responses dynamically during the first hours after transfer to a treatment
condition. However, tools to precisely measure such responses on the order of hundreds or thousands of
roots with high spatiotemporal resolution are spars .
To accurately measure even small changes in root growth, spatial resolution is of high importance. High-
resolution CCD scanner images that are otherwise suited for high-throughput studies (Slovak *et al.*, 2014)
won't yield the required resolution, as, even with a high scanner resolution, only ~21 µm/pixel was
achieved. Camera systems are frequently used to achieve medium resolution for root imaging and can
detect subtle root growth changes (Durham Brooks, Miller and Spalding, 2010). Microscopes in
conjunction with growth chambers or microfluidics chambers have been used to study root growth
changes with high time resolution while inducing a stress (Grossmann *et al.*, 2011; Busch *et al.*, 2012; de
Luis Balaguer *et al.*, 2016; von Wangenheim *et al.*, 2017; Stanley *et al.*, 2018). However, this method is
particularly efficient to catch the extremely fast response this remains an expensive solution for tracking
only ~8 roots at a time preventing any high throughput image acquisition. Moreover, production of
growth chambers, the use of confocal microscopy and dedicated software to automate these microscopes
frequently pose hurdles towards their adoptability (Grossmann *et al.*, 2011; Busch *et al.*, 2012; de Luis
Balaguer *et al.*, 2016; von Wangenheim *et al.*, 2017; Stanley *et al.*, 2018).
Another important element to study root growth in a high throughput manner is the ability to
automatically measure the roots at different timepoints. Many efforts have been made to develop root
image analysis tools that allows for this. For instance, the software MyRoot can calculate root length in a
high throughput manner using scanner images (Betegón-Putze *et al.*, 2019), but requires human
interaction and thus limits the processing of the time series at high time resolution. Tools allow for
assessing root growth variation with high time resolution using time series images such as RootTrace and
RootTip softwares (French *et al.*, 2009; Geng *et al.*, 2013) require the use of a camera which provides
sufficient resolution to detect root growth variation at time-scales from 3 minutes to 15 minutes,
respectively. However, the use of a camera limits high through put phenotyping e.g. from 1 for RootTrace
to 6 plates for RootTip at a time containing 24 seedlings. Moreover, the downstream time series image
analysis is semi-automated method which becomes very time consuming with increasing sample sizes.
Lastly, a vertical confocal microscope and microfluidic system are often not provided with an automated
software to calculate the root growth making hard to automate root growth analysis (Grossmann *et al.*,
2011; Busch *et al.*, 2012; de Luis Balaguer *et al.*, 2016; von Wangenheim *et al.*, 2017; Stanley *et al.*, 2018).
Taken together, tools that can be used to acquire and automatically analyze root growth in a high
throughput manner with high spatiotemporal resolution are currently not available.
Here, we present an analytic phenotypic platform that allows for the high throughput acquisition and
quantification of early root growth response at high temporal and spatial resolution named "Root
Walker". This method allows to acquire bright-field roots pictures using a microscope within minutes after
imposing treatments and for up to 12 hours. The acquired images are then processed by a MATLAB® script
in which roots are segmented and their growth after transfer are automatically quantified. To illustrate

the accuracy and power of this method, we used *Arabidopsis thaliana* treated with different
concentrations of the auxin analog, indole-3-acetic acid (IAA), as well as another treatment with two
mitogen-associated protein kinase (MAPK) inhibitors (U0126 and PD98059). We show that the Root
Walker pipeline allows for the determination of root length in a high throughput manner with high time
and spatial resolution. We propose that this platform is highly suited to conduct large scale targeted or
exploratory quantitative genetic screens to identify genes controlling early root growth responses to
stresses, nutrients or other internal or external cues.

Results

A simple growth chamber in conjunction with automated microscopy allows for root imaging at high
temporal and spatial resolution

We set out to develop an automated platform for capturing root growth responses at a large-scale. For
this, we wanted to image at a resolution that could capture even small changes of root growth. We also
used an automated platform to capture numerous roots at a high time resolution. For this, we used an
inverted BZ-X810 KEYENCE® microscope that allowed to automatically capture images with a resolution
of ~3µm/pixel for using a 2X objective (NA/0.1) or a resolution of ~1.5 µm/pixel using a 4X objective
(NA/0.2). This level of resolution allows for the detection of even subtle root growth rate changes. For
capturing many roots, used an agar filled chamber in which we could place ~20 seedlings (Figure S1a). To
allow the accurate quantification of root growth responses, plants were first germinated under standard
growth condition on plates and then transferred to imaging chambers containing the treatment medium
(Figure S1). Per chamber, either 9 images or 17 images are acquired depending on the objective used. Up
to three of these chambers can be mounted for imaging (Figure S1b-c), resulting in an imaging capacity of
~60 seedlings per experimental round. As we used 4 microscopes, we were able to measure ~240 roots
at a time. The described imaging setup allows for taking images at a highly frequent rate which varies
according to the objective used and the number of imaging chambers loaded on the microscope stage.
For example, these values are ranging from 18 seconds loading one chamber using the 2X objective to 59s
loading 3 chambers with the 4X objective. Here, we used two different setups, acquiring images every
minute for 60 minutes with the 4X objective and every 5 minutes for 12 hours with the 2X objectives.
Taken together, this microscope set up provides a platform to acquire images of roots at high time and
spatial resolution in a high-throughput manner.

A user-friendly software-tool allows to calculate the root length in an automated and high-throughput
manner

After successful image acquisition, each image is processed by a user-friendly MATLAB® script that only
requires three user inputs: i) select a parent folder containing the images; ii) select a folder to store the
output files, and iii) supply a descriptive file-prefix to be used for the output files. This script automatically
quantifies root length in the time series images acquired with the Keyence® BZ-X810 microscope of

Arabidopsis thaliana roots by performing several steps: (1) convert raw images to a large stitched image
(2) transform into a binarized stitched image, (3) convert binary stitched images into arrays of root center
midpoints, (4) "walk" along the center root array to virtually reconstruct the root, (5) calculate root length
and remove outliers, (6) allow for quality control (QC) through the generation of videos and graphical plot
output files (7) export of raw data in .csv files (8) further data analysis and graphs can be continued using
Excel® (Figure 1a). The script takes only a ~190 ms per single image (if 9 images is contained in one
stitched image the time is ~1.710 seconds per stitches) using a desktop computer with Intel® Xeon® W-
2135 CPU @3.70 GHz and 64GB of random-access memory. Note that we developed two different scripts
adjusted to quantify root growth when acquiring images every minute for 60 minutes using the 4X
objectives and every 5 minutes for 12 hours using the 2X objective as mentioned in the previous
paragraph. In the subsequent sections, we describe the steps of the software in detail.
132

Root segmentation

The first step (1) of the algorithm consists of stitching all images that were acquired during a single time
point together (Figure 1b). The next step (2) is then to accurately segment all roots that are contained in
the stitched image enabling a precise measurement of the root length (Figure 1c). For this, a binarized
image is generated using Otsu's thresholding method, which minimizes the intraclass variance of the
binary pixels (Figure 1c). Then a locally adaptive thresholding, and darker foreground than background
methods are applied to increase the thresholding accuracy (Figure 1c). Finally, background noise and air
bubbles coming from the camera sensor itself or from the plant transfer procedure to the imaging
chamber are eliminated, respectively. This step removes small objects (< 8000 pixels) and very large
objects (> 150 000 pixels) to obtain images with only roots as the remaining objects (Figure 1c). Note that
we particularly focused on annotating carefully the MATLAB® script to offer the possibility for the users
to fine tune these parameters depending on the users' needs and deliver a versatile tool. The resulting
image contains only segmented roots. However, these segmented roots have very jagged, fractal-like
edges, which when used to track the center line of each root leads to an overestimation of the root length
(analogous to the coastline paradox). To reduce this error, the image is convolved with a smoothing kernel
to smooth out the jagged root edges. The borders are then cleared of any black pixels created during the
smoothing process (Figure 1c). The binarized stitched images are now saved for the subsequent
generation of diagnostic videos and data analysis.
151

Determination of the root length using the Root Walker

Once, the images are segmented, the root center line is determined based on the center array of each
root object to calculate the root length at each time point (step 3, Figure 1d). Each stitched image is then
processed to create a set of arrays that corresponds to the center point of each root for each row of every
image. This process consists of several distinct steps: (1) for each row of each stitched image, the indices
of the black pixels (Figure 1d), corresponding to an object are recorded. This set of indices is saved as a
vector for each row of each stitched image and is of the same height in dimension as the image (Figure
6a, eg. 1920 pixels). (2) The entire set of black pixel indices is separated into vectors of connected sets of
black pixel indices, separating each object that is not touching or connected in the image. The objects are

now considered as distinct which then allows to process them individually in the next steps and iterative
functions. (3) The mean of each vector of the black pixel indices is taken, replacing the vector with a single
value corresponding to the center point index for each object for each row in the stitched image. As a
result, the data set is the same height as the input stitched image but is now much smaller in width which
reduces the amount of data to process and speeds up the analysis (Figure 1e). This step is performed for
each row of each stitched image that was previously created.
The center point index array does not reflect the root length after transfer yet as each root object is stored
in a data set which is the same height but for which the width is variable as indicated in Figure 1e.
Therefore, further processing needs to be conducted by "walking" along each root pixel array previously
generated to artificially reconstruct the root (step 4, Figure 1f). The value of the center point index array
is used to determine which objects correspond to roots based on its horizontal distance in the next row.
The method selects the first center point index array as a start and loops through each row of the array,
stepping down each row from the top to the bottom, and determines if there is an index in the next row
that is within a certain horizontal distance (default of 30 for the 60 minutes timelapse script or 20 pixels
for 12 hours timelapse script) of the root's center point in the previous row. Note that we particularly
focused on annotating carefully the MATLAB® script to offer the possibility for the users to fine tune these
parameters depending on the users' needs and deliver a versatile tool. This process is repeated for the
entire center index array, saving each center point within that set criteria as the next value in that "root"
vector (Figure 1f). The method repeats this process for each timestep, for each stitched image, for each
starting point or object in the center point index array, and for each row of the stitched image. While this
is a repetitive process, it is computationally extremely fast due to the simplicity of the calculation being
performed. After the method has carried out this step, the dataset now includes position vectors for each
center point for each row of each object in each stitched image.
To increase the robustness of our root growth rate measurements, the software next integrates different
timesteps and conducts further processing. This aims to address curvature due to circumnutation (Taylor
et al., 2021) that may create imprecision in measuring the root growth, as well as potential distortions
due to our focus on pixel root center array (we need three width pixels but a thresholded root tip may
contain only one or two pixels in some timesteps). To overcome these issues, only new growth is being
measured at each time point and this is achieved by overwriting each timestep with the previous timestep
ensuring that only new growth relative to the previous root tip location is measured.
Next, the position vectors for each root are utilized to perform a center line distance calculation to
determine the length of each root at each timestep (Figure 1g and Figure S2). The center line positions for
each row are used to calculate the distance between each row, summing each small change in length
between each row of the image to calculate the entire root length. This process is completed for all roots
in the current root center array.
196
197 **Outlier removal**
198
The method then filters out outlier growth rates that might correspond to image processing errors or the
failure to remove a non-root object from analysis earlier in the method (step 5, Figure 1h). This method
utilizes the root length data that was previously generated and removes any elements that do not change
in length for 20 or 45 timesteps, for short or long-time lapses, respectively. Note that we particularly

focused on annotating carefully the MATLAB® script to offer the possibility for the users to fine tune these
parameters depending on the users' needs and deliver a versatile. These outliers might be roots that are
not growing due to damage while transferring them to the imaging chamber referenced as "stuck" roots
or a non-root object (such as a scratch or a bubble) that was included in the binarized image and remains
in the same location. The method also removes any roots that "jump", having an increase in length greater
than 45 pixels in a one timestep for our long time-lapse set up and 20 pixels for the short time-lapse set
up representing approximately in both cases a jump of ∼28μm per minute, being more than 7 times faster
than the root growth rate that we observe in healthy roots (∼3.9μm/min, Figure 1h). Note that we
particularly focused on annotating carefully the MATLAB® script to offer the possibility for the users to
fine tune these parameters depending on the users' needs and deliver a versatile. Depending on the
experimental setup, the percentage of "stuck" and 'jump" events detected, as well as undetected, false
negatives led to a loss of about 8% for the 60 minutes script (124 roots detected and 114 roots for which
the root length after transfer has been calculated over 7 independent experiments) and about 25% for 12
hours script (171 roots detected and 129 roots for which the root length after transfer has been
determined over 6 independent experiments) in our experimental set-up. Note that we did not detect any
false positives in this set up. After outliers and artifacts removal, the root length for each timestep is used
to determine the amount of new growth of each root per timestep by subtracting the previous timestep's
length from the current timestep as previously described. This is performed iteratively for each timestep
to construct a dataset containing the change in root length for each root in each timestep.
222
223　**Generation of output files**
224
To allow for a quality control step of the analysis, a timelapse video with annotations for manual
inspection is generated (step 6, Figure 1i and Video S1). For this, the raw and processed stitched images
are displayed as a video with the calculated root center lines plotted on the associated figure (Video S1).
The video includes the raw image with center lines for all detected roots, including "Jump" and "Stuck"
roots, as well as the binarized and thresholded image that contains the root objects that are remaining
after the artifacts and outliers removal. This step takes approximately half of the full image analysis time
(∼95 ms per single image (if 9 images is contained in one stitched image the time is 855ms per stitches)
using a processor Intel® Xeon® W-2135 CPU @3.70 GHz and 64GB of random-access memory. If diagnostic
videos are not needed, we provided an alternative script – thus, one script including this step and one in
which this step is omitted.
To provide a succinct analysis of the experiment, the dataset is plotted for visual inspection (step6) by
generating two plots. The first plot contains the root lengths versus time, showing each individual root
throughout the entire experiment. The second plot groups the roots by genotype and condition and plots
the average root lengths for each group versus time. Both plots are saved and exported as .jpeg. Note that
these plots are being intended for visual inspection and can be used to detect potential outliers that were
not detected by the automated analysis or to help to adapt in a fast manner every tunable parameters in
the script.
The processed root datasets are exported as .csv (using pixels as units) to allow for a comprehensive data
analysis (step 7, Figure 1j). To facilitate high throughput data organization in the .csv output file, the date,
the number of the replicate, the genotype and the growing conditions are directly extracted from the

parent folder. The data organization is made to be compatible with the "Pivot Table" function in Excel®
allowing to easily process large amounts of data (Figure 1j). Using this Excel® function the remaining
outliers, which might have passed the previous thresholding steps to filter out objects that "Jump" and
are "Stuck" can be removed. Moreover, the pixels units can be converted into physical measurements and
any root length derived traits, such as root growth rate and other parameters can be calculated (step 8).
250

251 **Validation of the platform studying the early root growth response to the auxin analog indole-3-acetic**
252 **acid (IAA) for 20 minutes**
253

Before submitting roots to a treatment condition, we evaluated the accuracy of our method by comparing
root growth data measured by a human observer versus the root walker. This experiment revealed that
our tool is highly accurate since no significant differences were observed when comparing the
quantification methods (Figure 2a-d). Moreover, scatter plot with linear regression and Pearson
correlation matrix analysis based on the root growth rate confirm our finding further validating the
reliability of our quantification tool (Figure 1e-f). The root growth rate in our system was ~2.3 +/-0.52
µm/min thus being in the same range as previously reported (Figure 2; Serre et al., 2021 Nat Plants;
Fendrych et al., 2018 Nat Plants, Prigge et al., 2020, eLife). To test the utility of the Root Walker platform,
we studied the response of Arabidopsis root growth rate upon treatment with indole-3-acetic acid (IAA).
To our knowledge this is the only treatment which has been reported to impact the root growth within
minutes when applied in nanomolar range (Serre et al., 2021 Nat Plants; Fendrych et al., 2018 Nat Plants,
Prigge et al., 2020, eLife). We applied IAA at 1nM (IAA[1]), 10nM (IAA[10]) and 100nM (IAA[100]) and
followed the root growth taking images every minute for an hour. In standard condition, the root growth
rate was still in the range of our previous results ~3.9+/-1.37 µm/min (Figure 3). Furthermore, in line with
the literature, we observed a decrease of the root growth rate in a concentration dependent manner in
the order of µm per minute (Figure 3 and Video S2, Serre et al. 2021). A significant decrease of the root
length was already observed 1 minute after transfer upon treatment with 10nM IAA and 100nM IAA
(Figure 3b). At the lower concentration of 1nM IAA, the first significant difference of root growth rate was
observed after 11 minutes timepoint, highlighting a concentration-dependent root growth response
(Figure 3b). Altogether, these results show that the Root Walker platform can capture extremely fast root
growth changes through its capability to acquire images with high time and spatial resolution and with
high throughput, making it a powerful tool for studying root growth responses.
276

277 **Investigation of two MAPK inhibitors on the root growth for 12 hours**
278

As we had shown that our platform is very well suited to study very fast responses as observed for IAA,
we also wanted to test whether it was capable to accurately capture growth responses that only occur
after several hours. For this, we investigated the impact of two well-known mitogen associated protein
kinase homolog 2 (MPK2) inhibitors on root growth. The ubiquitous MPK2 is part of the MAPK
phosphorylation cascade which is central in the regulation of eukaryotic developmental signaling events
(Ge, Fu and Meadows, 2002; Hotokezaka *et al.*, 2002; Bjornson *et al.*, 2014). We used two inhibitors of
MPK2 that have different modes of action: PD98059 inhibits the MAPK cascade by binding the inactive
form of MPK2 while U0126 directly inhibits MPK2 kinase activity (Bjornson *et al.*, 2014). To study the root

growth response to these MPK2 inhibitors, we monitored root length for 12 hours taking images every 5
minutes. We noticed that the root growth rate (mm/hour) was significantly lower when plants were
exposed to U0126, but no differences were apparent for PD98059 compared to the mock treatment
(Figure 4 Video S3). Moreover, a statistical different root length between U0126 and mock conditions was
already apparent after 2 hour and 25 minutes (Figure 4b). We concluded that the inhibition of MPK2
kinase activity rather than the sequestration of the inactive form is critical in regulating root growth. This
experiment showed that the Root Walker platform can shed light on new early root growth regulatory
mechanisms.
295
296 **DISCUSSION**
297
Here, we have described a new analytic high-throughput platform with high spatial and temporal
resolution allowing to accurately quantify early root growth responses to different external stimuli.
Minutes after being transferred to imaging chambers containing the treatment of about 20 roots per
imaging chamber can be imaged at the same time. By using an automated digital microscope, we were
able to mount 3 chambers per microscope and by using 4 microscopes in parallel, resulting in 240 roots
being imaged at a time. Using such a system, a time resolution of more than 30 seconds after the transfer
could be achieved at a spatial resolution of ~3μm/pixel for 2X objective and ~1.5 μm/pixel for 4X
objective. To process the data, we developed a MATLAB® pipeline to automatically quantify the root
lengths from these experiments. This is step is particularly user friendly since the initialization required
only three user inputs and works without any image preprocessing or human intervention. The platform
presented here is highly versatile and can be then adoptable by a broader set of scientist; 1) the Keyence®
microscope is affordable compared to confocal microscopes, 2) the imaging chambers used are
commercially available and can be used on any microscope types, 3) the script is tediously annotated
offering to the user to fine tune several parameters (thresholding value, outliers object size, stitching
overlap, distance of the index, stuck and jump) without requiring any skills in script coding, 4) the script
can work on any type of stitched images, 5) the root growth tracking time is wide, ranging from minutes
to 12 hours.
To demonstrate the utility and the power of this approach to capture early root growth responses, we
utilized the exogenous application of IAA for which, the root growth has been described to be inhibited
minutes after being exposed to this hormone at nanomolar ranges. Using our platform, we observed that
this response was concentration-dependent and occurred minutes after transfer confirming the
sensitivity of the Root Walker platform. To show the versatility of the platform at a different timescale,
we investigated longer root growth response, 12 hours after transfer using two MAPK inhibitors known
to interfere with the MAPK signaling cascade. This revealed that only U0126 triggers a significant root
growth decrease, suggesting that MPK2 kinase activity is required to regulate root growth. In the
literature, discrepancy between these inhibitors has been already reported while studying the plant
general stress response in shoots (Bjornson *et al.*, 2014). They found that the MAPK signaling cascade is
critical in mediating the rapid stress response element through the CALMODULIN-BINDING
TRANSCRIPTIONAL ACTIVATOR 3 (Bjornson *et al.*, 2014). However, opposite to our findings, in shoots, only
PD98059 was able to delay the response while no changes were observed in presence of U0126
highlighting that the sequestration of MPK2 rather than its catalytic activity is critical in triggering this

rapid response in shoots. Taken together, this might suggest that there are tissue and time scale
dependent MPK2 regulatory mechanisms in plants.
To conclude, the Root Walker platform is an excellent tool to study early root growth responses ranging
from minutes to 12 hours after transfer to the treatment conditions. Compared to other publicly available
platforms or software, the Root Walker provides a simple but comprehensive all-in-1 large-scale pipeline
to study early root growth rate (Slovak *et al.*, 2014; Durham Brooks, Miller and Spalding, 2010; Betegón-
Putze *et al.*, 2019; French *et al.*, 2009; Geng *et al.*, 2013; von Wangenheim *et al.*, 2017). It not only allows
to acquire images with high spatiotemporal resolution in a high throughput manner but also to analyze
root growth automatically. We do think that this comprehensive platform is a prime choice to conduct
large scale screening or early root growth response to various stimuli which might enable the
identification of genes involved in sensing external stimuli.
340
341 **Experimental procedures**
342
Plant materials and growth conditions. For surface sterilization, *Arabidopsis thaliana* seeds
that had been produced under uniform growth conditions and were placed for 1 h in opened 1.5-
mL Eppendorf tubes in a sealed box containing chlorine gas generated from 130 mL of 10%
sodium hypochlorite and 3.5 mL of 37% hydrochloric acid. For stratification, seeds were imbibed
in water and stratified in the dark at 4 °C for 2-3 days. Seeds were then put on the surface of
media described in Gruber et al., 2013, using 12-cm x 12-cm square plates (Gruber *et al.*, 2013).
The Gruber et al., 2013 media contains, 750 µM of $MgSO_4$-7H2O, 625 µM of KH_2PO_4, 1000 µM
of NH_4NO_3, 9400 µM of KNO_3, 1500 µM of $CaCL_2$-$2H_2O$, 0.055 µM of $CoCL_2$-$6H_2O$, 0.053 µM of
$CuCl_2$-$2H_2O$, 50 µM of H_3BO_3, 2.5 µM of KI, 50 µM of $MnCl_2$-$4H_2O$, 0.52 µM of Na_2MoO_4-$2H_2O$,
15 µM of $ZnCl_2$, 75 µM of Na-Fe-EDTA, 1000 µM of MES adjusted to pH 5.5 with KOH and 0.5%
sucrose and 1% BactoAgar®. Plants were grown in long day conditions (16/8h) in walk in growth
chambers at 21°C, 50uM light intensity, 60% humidity. During nighttime, temperature was
decreased to 15°C.
356
Phenotyping of the root growth responses. Wild type seeds were sowed in media described
in Gruber et al., 2013 and stratified for 2-3 days at 4°C (Gruber *et al.*, 2013). Five days after
planting, about 20 seedlings were transferred using forceps to an imaging chamber (Lab-Tek,
Chambers #1.0 Borosilicate Coverglass System, catalog number: 155361) filled with medium
described in Gruber et al., 2013. Note that the transfer took about 45-60 seconds. Images were
acquired every minute or 5 minutes for 1 hours or 12 hours, respectively, in brightfield conditions
using a Keyence® microscope model BZ-X810 with a BZ NIKON Objective Lens (4X or 2X) CFI
Plan-Apo Lambda.
365
Treatment with IAA, U0126 and PD98059. The imaging chamber was filled up with the media
described in Gruber et al., 2013 in which IAA dissolved in DMSO (stock solution 1000X compared
to the final solution) at 1nM, 10nM or 100nM and DMSO was added (Gruber *et al.*, 2013).
Moreover, U0126 and PD98059 dissolved in DMSO (stock solution 20mM or 10mM, respectively)

at a concentration of 10µM and DMSO were added to the gruber media when poured to the
imaging chamber

Manual measurement of the root growth. Time lapse images were used to create videos with
the Keyence software®. The videos were opened in Fiji (Schindelin *et al.*, 2012) and the root tip
displacement was measured using the segmented line tool at each time point.

Statistical analysis. Each experiment has been repeated independently at least twice, as in
every case the same trend has been recorded for independent experiments, the data have been
pooled for further statistical analysis. Each sample were subjected to four different normality tests
(Jarque-Bera, Lilliefors, Anderson-Darling and Shapiro-Wilk), sample were considered as a
Gaussian distribution when at least one test was significant ($p < 0.05$) using Xlstat. As a normal
distribution was observed a two-ways ANOVA coupled with post hoc Tukey test was performed
($p < 0.05$) using Xlstat. As a normal distribution was not observed at two-ways Kruskal-Wallis
coupled with post hoc Steel-Dwass-Critchlow-Fligner procedure was performed ($p < 0.05$) using
Xlstat. The type of test performed in indicated in the figure legend.

Acknowledgements: We thank all the Busch lab members for critical discussions. This study was funded
by the National Institute of General Medical Sciences of the National Institutes of Health (grant number
R01GM127759 to W. Busch), and start-up funds from the Salk Institute for Biological Studies (W. Busch).
M.P. Platre was supported by a long-term postdoctoral fellowship (LT000340/2019 L) by the Human
Frontier Science Program Organization.

Declaration of interest: The authors declare no conflict of interest.

Code availability: Matlab® script can be found here, https://github.com/mplatre/Root-Walker-Script.git

REFERENCES

Betegón-Putze, I. *et al.* (2019) 'MyROOT: a method and software for the semiautomatic measurement of
primary root length in Arabidopsis seedlings', *The Plant Journal*, 98(6), pp. 1145–1156. Available at:
https://doi.org/10.1111/tpj.14297.

Bjornson, M. *et al.* (2014) 'Distinct Roles for Mitogen-Activated Protein Kinase Signaling and
CALMODULIN-BINDING TRANSCRIPTIONAL ACTIVATOR3 in Regulating the Peak Time and Amplitude of
the Plant General Stress Response1[W][OPEN]', *Plant Physiology*, 166(2), pp. 988–996. Available at:
https://doi.org/10.1104/pp.114.245944.

Busch, W. *et al.* (2012) 'A microfluidic device and computational platform for high-throughput live imaging
of gene expression', *Nature Methods*, 9(11), pp. 1101–1106. Available at:
https://doi.org/10.1038/nmeth.2185.

Durham Brooks, T.L., Miller, N.D. and Spalding, E.P. (2010) 'Plasticity of Arabidopsis Root Gravitropism
throughout a Multidimensional Condition Space Quantified by Automated Image Analysis', *Plant
Physiology*, 152(1), pp. 206–216. Available at: https://doi.org/10.1104/pp.109.145292.

French, A. *et al.* (2009) 'High-Throughput Quantification of Root Growth Using a Novel Image-Analysis
Tool', *Plant Physiology*, 150(4), pp. 1784–1795. Available at: https://doi.org/10.1104/pp.109.140558.

Ge, X., Fu, Y.M. and Meadows, G.G. (2002) 'U0126, a mitogen-activated protein kinase kinase inhibitor,
inhibits the invasion of human A375 melanoma cells', *Cancer Letters*, 179(2), pp. 133–140. Available at:
https://doi.org/10.1016/s0304-3835(02)00004-6.

Geng, Y. *et al.* (2013) 'A Spatio-Temporal Understanding of Growth Regulation during the Salt Stress
Response   in   Arabidopsis',   *The   Plant   Cell*,   25(6),   pp.   2132–2154.   Available   at:
https://doi.org/10.1105/tpc.113.112896.

Grossmann, G. *et al.* (2011) 'The RootChip: An Integrated Microfluidic Chip for Plant Science', *THE PLANT
CELL ONLINE*, 23(12), pp. 4234–4240. Available at: https://doi.org/10.1105/tpc.111.092577.

Gruber, B.D. *et al.* (2013) 'Plasticity of the Arabidopsis Root System under Nutrient Deficiencies', *Plant
Physiology*, 163(1), pp. 161–179. Available at: https://doi.org/10.1104/pp.113.218453.

Hotokezaka, H. *et al.* (2002) 'U0126 and PD98059, specific inhibitors of MEK, accelerate differentiation of
RAW264.7 cells into osteoclast-like cells', *The Journal of Biological Chemistry*, 277(49), pp. 47366–47372.
Available at: https://doi.org/10.1074/jbc.M208284200.

Jia, Z., Giehl, R.F.H. and von Wirén, N. (2022) 'Nutrient–hormone relations: Driving root plasticity in
plants', *Molecular Plant*, 15(1), pp. 86–103. Available at: https://doi.org/10.1016/j.molp.2021.12.004.

Li, B. *et al.* (2019) 'GSNOR provides plant tolerance to iron toxicity via preventing iron-dependent
nitrosative and oxidative cytotoxicity', *Nature Communications*, 10, p. 13.

de Luis Balaguer, M.A. *et al.* (2016) 'Multi-sample Arabidopsis Growth and Imaging Chamber (MAGIC) for
long term imaging in the ZEISS Lightsheet Z.1', *Developmental Biology*, 419(1), pp. 19–25. Available at:
https://doi.org/10.1016/j.ydbio.2016.05.029.

Morris, E.C. *et al.* (2017) 'Shaping 3D Root System Architecture', *Current Biology*, 27(17), pp. R919–R930.
Available at: https://doi.org/10.1016/j.cub.2017.06.043.

Platre, M.P. *et al.* (2022) 'The receptor kinase SRF3 coordinates iron-level and flagellin dependent defense
and   growth   responses   in   plants',   *Nature   Communications*,   13(1),   p.   4445.   Available   at:
https://doi.org/10.1038/s41467-022-32167-6.

Satbhai, S.B. *et al.* (2017) 'Natural allelic variation of FRO2 modulates Arabidopsis root growth under iron
deficiency', *Nature Communications*, 8, p. 15603. Available at: https://doi.org/10.1038/ncomms15603.

Schindelin, J. *et al.* (2012) 'Fiji: an open-source platform for biological-image analysis', *Nature Methods*,
9(7), pp. 676–682. Available at: https://doi.org/10.1038/nmeth.2019.

Slovak, R. *et al.* (2014) 'A Scalable Open-Source Pipeline for Large-Scale Root Phenotyping of Arabidopsis',
The Plant Cell, 26(6), pp. 2390–2403. Available at: https://doi.org/10.1105/tpc.114.124032.

Stanley, C.E. *et al.* (2018) 'Dual-flow-RootChip reveals local adaptations of roots towards environmental
asymmetry at the physiological and genetic levels', *The New Phytologist*, 217(3), pp. 1357–1369. Available
at: https://doi.org/10.1111/nph.14887.

Taylor, I. *et al.* (2021) 'Mechanism and function of root circumnutation', *Proceedings of the National*
Academy of Sciences, 118(8), p. e2018940118. Available at: https://doi.org/10.1073/pnas.2018940118.

von Wangenheim, D. *et al.* (2017) 'Live tracking of moving samples in confocal microscopy for vertically
grown   roots',   *eLife*.   Edited   by   C.S.   Hardtke,   6,   p.   e26792.   Available   at:
https://doi.org/10.7554/eLife.26792.

Figure 1. Overview of the Root Walker analysis platform. (a) Step diagram of the root walker analysis platform. b) Pictures that are acquired overlap by 30% . (**b**) Stitched picture (collage). (**c**) Upper panel, binary image obtained by using three steps: thresholding, removing background and smoothing. Lower panel, binary matrix (subset from red box) obtained from binary image, 0 represents a black pixel and 1 a white pixel. (**d**) Determination of the Root Center Array from the binary image using connected sets of black pixels and extraction of the root center values (blue text). (**e**) Root center value array stored as a data sheet. The algorithm walks down each row in the matrix to determine if there is a connected object in the next row progressing vertically down the image. The connected objects are represented by the same color and the number represent the root center array value. (**f**) The connected objects are stored separately as arrays in a data sheet. (**g**) Distances are then calculated for each root array in the matrix to calculate the root length. (**h**) removal of improbable root growth values. (**i**) Graphical output file representing the root length (in pixels) of each root, left, or in for the same genotype in one condition according to time. (**j**) Example of the data stored in the .csv file when opened in Excel®.

Figure 2. Comparison of root growth rate between Root Walker analysis script and human observer. (**a-c**) Root growth of individual 7-day old seedlings after transfer to the imaging chamber containing standard medium. Root length increase of individual roots over 60 minutes manually quantified (**a**), by the Root Walker script (**b**), or averaged (**c**). Error bars represent the standard deviation. (**d**) Root growth rate of data presented in a,b. [two-way student t-test; ($p<0.05$)]. n.s., non-significant. Red crosses represent the average and the red bars the median. Black dots individual measurements. (**e**) Scatter plot of the manually measured values in function of the root walker measured values. (**f**) From top to bottom, descriptive data of the root growth rate values manually measured, or root walker measured, correlation matric (Pearson), p-values (Pearson) and coefficient of determination (Pearson).

Figure 3. Exogenous application of Indole-3-acetic acid for an hour restricts root growth in a concentration-dependent manner. (**a**) From left to right, representative images of roots of 5-day old seedlings treated with DMSO, IAA at 1nM, 10nM and 100nM for over the course of one hour. Scale bar, 100 μm. (**b**) Graphical representation of the root length increases over time after transfer to treatment medium. [two-way student t-test; (p<0.05)]. Asterisks indicate statistical differences compared to the DMSO and the color corresponds to the curve. Error bars represent the standard deviation. (**c**) Quantification of root growth rate expressed in μm per minute of seedlings treated with DMSO, IAA at 1nM, 10nM and 100nM for an hour. [two-way ANOVA followed by a Fisher LSD test; letters indicate statistical differences (p<0.05)]. Red crosses represent the average and the red bars the median. Black dots individual measurements

Figure 4. Exogenous application of the MPK cascade inhibitor U0126 but not PD98059 restricts root growth. (**a**) From left to right, representative images of roots of 5-day old seedlings treated with DMSO, PD98059 and U0126 at 10µM over the course of 12 hours. Scale bar, 200 µm. (**b**) Graphical representation of the root length increases after transfer to DMSO, PD98059 and U0126 at 10µM for 12 hours. [two-way student t-test; (p<0.05)]. Asterisks indicate statistical differences compared to DMSO and the color corresponds to the curve. Error bars represent the standard deviation. (**c**) Quantification of the root growth rate expressed in mm per hour of seedlings treated with DMSO, PD98059 and U0126 at 10µM for 12 hours. [two-way ANOVA followed by a post hoc Tukey HSD test; letters indicate statistical differences (p<0.05)]. Red crosses represent the average and the red bars the median. Black dots individual measurements.