Initiation of fibronectin fibrillogenesis is an enzyme-dependent process

Shay Melamed[1], Shelly Zaffryar-Eilot[1], Elisabeth Nadjar-Boger[1], Rohtem Aviram[1], Huaning Zhao[2], Wesal Yaseen-Badarne[1], Rotem Kalev-Altman[3], Dalit Sela-Donenfeld[3], Oded Lewinson[4], Sophie Astrof[2], Peleg Hasson[1,*], Haguy Wolfenson[1,*]

[1]Department of Genetics and Developmental Biology, The Rappaport Faculty of Medicine and Research Institute, Technion – Israel Institute of Technology, Haifa 31096, Israel

[2]Department of Cell Biology and Molecular Medicine, New Jersey Medical School, Rutgers Biomedical and Health Sciences, Newark, NJ 07103, USA.

[3]Koret School of Veterinary Medicine, Faculty of Agriculture, Food and Environmental Sciences, The Hebrew University, Rehovot, Israel.

[4]Department of Molecular Microbiology, The Rappaport Faculty of Medicine and Research Institute, Technion – Israel Institute of Technology, Haifa 31096, Israel

*correspondence: phasson@technion.ac.il, haguyw@technion.ac.il

Summary:

Fibronectin fibrillogenesis and mechanosensing both depend on integrin-mediated force transmission to the extracellular-matrix. However, force transmission is in itself dependent on fibrillogenesis, and fibronectin fibrils are found in soft embryos where sufficient force cannot be applied, demonstrating that force cannot be the sole initiator of fibrillogenesis. Here we identify a novel nucleation step prior to force generation, driven by fibronectin oxidation mediated by lysyl-oxidase enzyme family members. This oxidation induces fibronectin clustering that promotes early adhesion, alters cellular response to soft matrices, and enhances force transmission to the matrix. In contrast, absence of fibronectin oxidation abrogates fibrillogenesis, perturbs cell-matrix adhesion, and compromises mechanosensation. Moreover, FN oxidation promotes cancer cells colony formation in soft agar as well as collective and single-cell migration. These results reveal a yet unidentified, force-independent enzyme-dependent mechanism that initiates fibronectin fibrillogenesis, establishing a critical step in cell adhesion and mechanosensing.

Introduction:

Cell adhesion to the extracellular matrix (ECM) is a fundamental property that is essential for multiple cellular processes including migration, proliferation, and differentiation. As a result, cellular functions often depend on alterations within the ECM, including changes in its composition, structure, and mechanical properties[1,2]. Fibronectin (FN) is an essential ECM protein that promotes cell adhesion to the matrix[3,4], and is also an important regulator of ECM alterations owing to its binding to numerous ECM proteins, including collagen[5], fibrillin[6], fibrinogen[7], and Tenascin-C[8]. FN is secreted as an inactive dimer and it promotes matrix assembly and integrin-mediated cell adhesion to the matrix in its active multimeric fibrillar form[9,10]. The dimeric to pro-fibrillar transition, which activates the initiation of FN fibrillogenesis, is considered to be driven by intracellular forces generated and transmitted via the actomyosin cytoskeleton to the extracellular FN molecules through transmembrane integrins[11–13]. When generated, the forces lead to conformational changes within FN, exposing cryptic self-association binding sites which allow adjacent dimers to interact, thus gradually promoting the assembly of FN fibrils[14,15]. This process is associated with growth of nascent integrin adhesions into mature focal- or fibrillar-adhesions[16,17]. Indeed, in situations where high forces cannot be generated, as in cultured cells grown on soft matrices, FN fibrillogenesis is inhibited, resulting in reduced cell adhesion[14]. In developing embryos, fibril assembly depends on cadherin-dependent enhanced contractility that leads to integrin-FN translocation[18]. However, in very early stages of embryonic development, the embryo rigidity is so low[19,20] such that it is not expected to provide sufficient resistance to support transmission of high cellular forces to the matrix. Thus, the observed FN fibrils at such early embryonic stages[21,22], which are essential for facilitating cell adhesion and migration, appear to depend on additional mechanisms which can overcome and compensate for the low rigidity and the resulting inability to exert force. Here we set to dissect this longstanding conundrum. We found a novel force-independent mechanism that is essential for the initiation of FN fibrillogenesis.

Results:

Lysyl oxidases promote FN fibrillogenesis and cell spreading on soft matrices

The presence of FN fibrils in early, soft, embryos whose stiffness cannot support transmission of high forces thar are involved in integrin-dependent FN conformational changes, suggests that an additional mechanism is involved in FN fibrillogenesis. Our previous finding that members of the Lysyl oxidase enzyme family bind FN, that LOX-Like 3 (LOXL3) oxidizes it, and that its enzymatic activity induces FN-dependent integrin activation[23], raised the possibility of an enzyme-dependent *initiation of fibrillogenesis*, a force-independent mechanism which relies on enzymes and their substrates' availability. In line with this notion, we found that in cultured cells expressing GFP-tagged FN[24], FN fibrils are observed under β1-integrin adhesions and LoxL3 is distributed at nascent adhesion sites (Figure 1A). Notably, LoxL3 is expressed in neural crest cells (Figure S1A) and following its deletion (*LoxL3$^{\Delta/\Delta}$*) the mutant embryos display a cleft palate phenotype and reduced neural crest cell numbers at branchial arches (Figure S1B-H), both of which are FN-associated phenotypes[25,26]. Further, abnormal FN-fibril formation along the somitic boundaries is also observed in *LoxL3$^{\Delta/\Delta}$* mutant embryos (Figure S1I-K). Collectively, these findings suggest a role for Lysyl oxidases in FN fibrillogenesis.

To test the relationship between substrate rigidity, FN fibrillogenesis, and the Lysyl oxidases, we took advantage of primary human aortic smooth muscle cells (HASMCs), which a) normally reside on a rigid artery wall which, depending on stretching, is 15-88 kPa[27]; and b) primarily express the lysyl oxidase family member, LOX[28]. Following infection with short hairpin RNA against *LOX* (*shLOX*) or scrambled RNA (*shCtrl*) (Figure S1L), we cultured the cells for 24 hrs on 1 kPa or 40 kPa FN-coated silicone surfaces and monitored FN-fibrils by immunostaining. Whereas *shCtrl* cells displayed significant FN fibril formation, cells devoid of LOX, despite secreting FN to the same extent (Figure S1M), did not form any FN fibrils on both rigidities (Figure 1B-G).

We next wished to carry out the converse experiment and test whether pre-treatment of FN with lysyl oxidases overrides rigidity sensing and promotes early cell adhesion on soft matrices. Towards that end, we used recombinant human LOXL3 or LOXL2, and treated FN-coated 1.0 and 0.35 kPa surfaces with either enzyme. Mouse embryonic fibroblasts (MEF) stably expressing Paxillin-GFP (Pax-GFP) were then added onto the plates and

fixed after 60 and 120 minutes on the sham-, LOXL3- or LOXL2-treated FN (Figure 1H-W and Figure S1N-ZC). FN fibrils were undetectable at these early times of cell spreading and therefore we monitored cell areas as readouts of cellular interaction with FN. This analysis revealed a significant increase in the areas of cells plated on the LOXL3- or LOXL2-treated FN in comparison to the sham-treated FN or to those treated with the enzyme mixed with the pan LOX inhibitor beta-aminopropionitrile (βAPN)[29].

Previous work has noted significant differences in mechanosensing whether the ECM proteins were physically absorbed or chemically crosslinked to the substrate as cell-generated forces can pull-off ECM molecules from the substrates when they are physisorbed[30]. Although we used soft silicone gels (CY 52-276) on which this phenotype does not occur[31], to verify that the above observations are not due to crosslinking of FN to the underlying gel via the LOX family members but rather due to a possible change in FN, we crosslinked LOXL3-treated or untreated FN to 0.35 kPa gels and monitored cell spreading after 2 hours. We found that in a similar manner to that observed above, also in this case, the treatment of FN with a LOX family member enhanced cell spreading compared to the untreated control (Figure S1ZD), demonstrating that the crosslinking step failed to significantly enhance the link between FN and the substrate beyond the physisorbed FN. Altogether, these results suggest a general role for the enzymatic activity of lysyl oxidases that promotes FN fibrillogenesis and cell adhesion.

Figure 1. Lysyl oxidases regulate FN fibrillogenesis on soft matrices. MEF expressing FN-GFP stained for β1-integrin and LoxL3 **(A)**. FN immunostaining, color-coded for intensity (blue – low, yellow – high), of shLOX and shControl HASMC on 1 kpa **(B-C)** and 40 kpa **(D-E)**. Magnification of representative regions in **(D and E)**. Quantification of FN fibrils show significant reduction following LOX knockdown on both 1 kpa and 40 kpa **(F-G, n>40)**. Pax-GFP on sham- and LOXL3-treated FN with or without βAPN, seeded on 1 kPa **(H-M)** or 0.25 kPa **(P-U)** gels and fixed after 1 hour **(H-J, P-R)** and 2 hours **(K-M, S-U)**. Quantification of cell area show significant differences between cells seeded on treated FN regardless of rigidities **(N-O, V-W, n>28)**. 1**p<0.0001, **p<0.001, *p<0.05, n.s. = non-significant.

LOX-treated FN promotes nascent adhesion formation

The above results suggest that LOX family members modulate the cells' response to the external rigidity by initiating FN fibrillogenesis and enhancing adhesion formation. We therefore wished to test whether the direct treatment of FN with a lysyl oxidase affects the mechanism by which cells perform mechanosensing of ECM rigidity. We postulated that the enzymatic activity of lysyl oxidases promotes a structural change in FN that allows creating more stable nascent adhesions through which early mechanosensing of matrix rigidity occurs[32,33]. To test this, we plated cells on arrays of deformable PDMS pillars, with 0.5 μm diameter and 2.3 μm height (bending stiffness 1.5 pN/nm[32,34], equivalent to ~2 kPa[35]), and performed live-cell time-lapse imaging of early attachment and spreading of the cells (Figure 2A-C). To test the hypothesis that LOX-induced local structural change in FN enhances nascent adhesion formation, we coated the pillars with FN at a density which was less than saturation level (1 μg/ml, equivalent to ~25% coverage of the pillar tops). Analyses of early times of cell edge interactions with individual pillars revealed repeated catch-and-release events which were 3 standard deviations above the mean noise level (Figure 2A). We analyzed the strength of the linkage in each of these events by measuring the pillar displacement level at the time of release, translating it to force. Comparison of the histograms of release forces between sham- and LOXL3-treated FN showed that the latter displayed significantly more events per unit time at 2 pN (Figure 2B), consistent with the formation of integrin-mediated FN-actin 2 pN bonds in the presence of trimers of FN's integrin-binding domain FNIII7-10, but not with monomers[36]. These results are suggestive of formation of more closely packed FN clusters following LOXL3 treatment. Further, analyses of the longer-term pillar displacements that succeeded the initial catch-and-release events showed a significant shift toward higher

values of the maximal displacement histogram in LOXL3-treated compared to control FN (Figure 2C). Together, these results suggest that the ability to form nascent adhesions on soft surfaces is enhanced upon LOX treatment, which assists in overcoming the low rigidity of the matrix (Figure 1).

We next considered the fact that nascent adhesion formation even on very stiff matrices requires at least four RGD (Arg-Gly-Asp) ligands that are localized within ~60 nm from each other[37,38]. Thus, if LOX treatment led to an adhesion promoting structural change in FN, one should expect to see a difference in early adhesion even on stiff surfaces such as glass whose stiffness is at the GPa range. Towards that end, following enzymatic treatment, FN was coated on glass coverslip and Pax-GFP cells were added and shortly after (15 or 30 minutes) fixed and analyzed. Quantifications of adhesion sizes at 15 minutes showed a significant increase in the relative count of adhesions larger than 0.1 μm^2 on LOXL3-or LOXL2- treated FN and to a concomitant increase in cell spreading area (Figure 2D-F, J, L and Figure S2A-B, E, G).

Importantly, the increases in adhesion sizes and cell areas were blocked when βAPN was added to the FN-LOXL3 mix throughout the reaction (Figure 2J, L, I). A similar trend was observed at 30 minutes, albeit to a lesser extent (Figure 2G-L, K, M and Figure S2C-D, F, H), altogether suggesting that the primary effect of the enzymatic treatment occurred at the very early stages of adhesion assembly. A comparable effect was observed with two other fibroblast cells (Figure S2I-P), indicating that the effect of lysyl oxidases on adhesion to FN was not specific to a particular cell line. Notably, since α5β1 integrin can interact with LOX enzymes[39], we wished to verify that the observed effect on adhesion occurred through changes in FN and not through direct integrin activation by residual enzyme that remained attached to FN (Figure S2U). Indeed, when βAPN was added following the FN treatment, no difference was observed in cell spreading (Figure S2Q-T, V), demonstrating that the enzymatic activity of LOXL3 directly affects FN and that the presence of the enzyme by itself (when inactive) has no effect. Taken together, the similar results observed following LOXL2 and LOXL3 treatment (Figure 2 & Figure S2), and the observations on LOX effects on FN fibrillogenesis (Figure 1 & Figure S1), reinforce the idea of a general enzymatic role among LOX family members in affecting cell adhesion through initiation of FN fibrillogenesis.

Figure 2. LOX family stimulate adhesion initiation. Representative image of a cell on pillars and a graph of pillar displacement; the red arrowheads point to catch-and-release events. **(A)**. Histograms of the pillar release force during catch-and-release events by cells seeded on sham or LOXL3-treated FN, demonstrating enhanced formation of 2 pN bonds upon LOXL3 treatment **(B, n>243)**. Histogram of the maximal displacement of pillars by cells seeded on LOXL3-treated FN **(C, n>243)**. Pax-GFP on sham- and LOXL3-treated FN with or without βAPN, seeded on glass and fixed after 15 **(D-F)** and 30 minutes **(G-I)**. Quantifications of cell area show significant enhanced spreading of cells seeded on treated FN compared to controls **(J-K, n>39)**. Histogram

of focal adhesion area of cells seeded on sham- or LOXL3-treated FN demonstrate that LOXL3 treatment promotes larger adhesions **(L-M)**. ***p<0.0001, **p<0.001, *p<0.05, n.s. = non-significant.

LOX induces FN clustering

We next wished to directly test whether alteration to FN organization occurs following LOX treatments. To this end, we deposited treated and untreated FN samples, as well as fresh FN, on glass coverslips at low density (100 ng/ml). After immunostaining the samples, we imaged regions completely devoid of cells using direct stochastic optical reconstruction microscopy (dSTORM). We then used particle analysis to identify FN clusters in the dSTORM images (Figure 3A-B). The histograms of the detected FN clusters indicate that LOXL3 treatment led to significant increase in the number of clusters larger than 0.009 µm^2 compared to fresh or sham-treated FN (Figure 3C). Collectively, these results indicate that LOX treatment enhances FN clustering, which results in cells perceiving the matrix as stiffer.

Since adhesion stabilization plays a major role in maintenance of a leading edge in cell migration[40], we postulated that the larger FN clusters following treatment would lead to improved migration. We therefore plated Pax-GFP cells on sham- and LOXL3-treated FN and performed live-cell imaging to track their migratory patterns and quantify the overall displacement length, migration distance, and speed (Figure 3D-H). All three parameters were significantly increased following LOXL3 FN-treatment, demonstrating that not only were the cells able to rapidly generate adhesions on the treated FN but importantly, that the cells' migration pattern was more persistent (evident by the increased displacement length), further reinforcing the notion that the adhesions that form on the LOXL3-treated FN are more stable.

Figure 3. LOX induces FN clustering. Representative dSTORM images of sham- and LOXL3-treated FN nanodomains; the curves below are the intensity values along the white lines shown in each image **(A-B)**. dSTORM analysis of FN cluster area shows a significant increase following LOXL3 treatment **(C)**. Representative color-coded images generated from time-lapse videos tracking fluorescence-labeled nuclei on sham- and LOXL3-treated FN **(D-E)**. Quantifications of migration parameters show significant increases in the measured parameters of cells seeded on LOXL3-treated FN **(F-H, n>426)**. LoxL3 and LoxL2 immunostaining of non-polar cells **(I-L)** and polar cells **(M-P)** showing that the Lox family members are localized to lamellipodial regions. 3D dSTORM image and zoom-in x-z representation of the cell edge demonstrating LoxL3 secretion **(Q)**. Quantification of the percent of LoxL3 molecules localized below the cell leading edge **(R, n=17)**. ***p<0.0001, **p<0.001.

The overall results thus far suggest that exogenous FN treatment with a member of the lysyl oxidase family promotes a faster and more robust cell adhesion. As members of the family are secreted enzymes, we next wished to test whether their secretion is localized to cellular regions where such robust integrin-dependent adhesions are mostly required. To that end, Pax-GFP cells were immunostained for LoxL2 or LoxL3 following 60 minutes of spreading. We find that in polarized cells, LoxL2 and LoxL3 expression is typically localized to the most prominent lamellipodial protrusion of the cell (Figure 3I-L). Surprisingly, even in non-polarized cells, narrow stripes in lamellipodial regions were also observed (Figure 3M-P). Further, 3-dimensional (3D) dSTORM analysis demonstrates that these matrix modifying enzymes are specifically secreted from the leading edge (Figure 3Q-R). Taken together, these results support a model whereby localized secretion of LOX enzymes at the cell edge enhances FN clustering, thereby supporting nascent adhesion formation and persistent cell migration.

FN oxidation is essential for FN fibrillogenesis

Since a) Lysyl oxidases' enzymatic activity is required for FN-mediated cell adhesion and migration; b) treatment of FN with LOXL3 promotes its clustering; and c) LOXL3 oxidizes FN[23], prompted us to further explore the oxidation and dissect its necessity. Liquid chromatography–mass spectrometry (LC-MS/MS) analysis identified three lysine residues (K116, K2391, K2401) that were specifically oxidized by both LOXL2 and LOXL3 (Figure 4A). Since no active recombinant LOX enzyme is available, we cultured HASMC and submitted the FN they secrete to the LC-MS/MS analysis. We find that in a similar manner to the two LOX-like proteins, LOX also specifically oxidizes lysine 116 (Figure 4A

and Figure S3A), altogether demonstrating the specificity of the reaction and reinforcing the above observations of the shared FN-dependent activities the LOX enzymes carry. Analysis of the oxidized lysine residues demonstrates that they are all localized to FN type 1 (FN1) repeats (FN1 repeat #2 and FN repeat #12), one close to the FN N-terminus and the other in its C-terminal region. Both regions encompassing these lysine residues are highly conserved in vertebrate FN (Figure 4B) suggesting they play an important role in FN activity.

To directly test the role that these three lysine residues play in cell adhesion and FN fibrillogenesis, we generated a lentiviral construct expressing a mutant FN form where we mutated them to Alanines, rendering them incompetent for oxidation (FN$^{\Delta Oxi}$). As a control, we mutated the α5β1 and αv integrins binding site RGD to RGE (FN$^{\Delta RGD}$) thus blocking their binding through this site[41]. Western blot analysis from cells overexpressing these two constructs demonstrated that they are expressed to similar levels as that of cells over-expressing the full length wild-type FN protein (FN^{WT}) (Figure 4C-D).

Considering that LOX treatment enhances FN clustering (Figure 2), we took into account that under classical adhesion assays with excess FN adsorbed to the surface (typically 10-20 µg/ml) stochastic clustering readily occurs. Thus, to avoid the possible inference of pre-adsorbed FN, we performed early adhesion assays of the cells expressing the FN variants directly on glass coverslips with no FN coating. We cultured the cells for 1 hour and quantified their areas to test whether the non-oxidizable FN variant inhibits initiation of cell adhesion. We find that the over-expression of FN$^{\Delta RGD}$ led to a significant reduction in cell area and adhesion sizes, demonstrating the inhibition of the FN-integrin interactions. Notably, the over-expression of FN$^{\Delta Oxi}$ led to a similar reduction in cell area and adhesion sizes as that observed with the over-expression of FN$^{\Delta RGD}$, demonstrating the role that these oxidation sites play in the initiation of FN-dependent integrin activation (Figure 4E-I).

To explore whether these mutations affect also FN fibrillogenesis, we first tested whether cells fibrilize their own secreted FN. To that end, we coated coverslips with Alexa 647-labelled FN and cultured MEF FN-GFP (in which GFP is knocked-in to both alleles of the FN gene[24]) for 48 hours, followed by imaging of the FN fibrils. All FN fibrils we observed were GFP labelled, demonstrating that cells use their own secreted FN in forming the

initial fibrils (Figure S3B-C). Next, we cultured the cells expressing the FN variants for 24 hours, immunostained them for FN, and monitored FN fibrils. Cells over-expressing the $FN^{\Delta RGD}$ demonstrated a significant 27% reduction in the ratio between fibrillar FN vs. total FN; thus, even though integrin binding was perturbed in the lack of the RGD site, alternative interactions with αvβ3 integrin receptors could promote FN fibrillogenesis[41], but to a lesser extent. Strikingly, cells over-expressing $FN^{\Delta Oxi}$ displayed significantly less fibrillar FN (>62% reduction), despite containing the wild-type RGD site, demonstrating that the oxidation is a prerequisite for adhesion initiation (Figure 4J-M). As these cells still express endogenous wild-type FN, these results strongly demonstrate the critical and essential role played by members of the LOX-family enzymes in regulating FN fibrillogenesis, establishing that initiation of the process is an enzyme- rather than force-dependent process, in line with their ability to generate force-bearing stress fibers (Figure 4N). Similarly, when we used HASMC infected with shCtrl lentiviral particles we found that they nicely formed FN fibrils when cultured directly on glass, but shLOX HASMC did not form any FN fibrils, even though they were cultured on a very stiff matrix and even though they too formed stress fibers (Figure 4O-R), further demonstrating the necessity for LOX in regulating FN fibrillogenesis, integrin activation and cell adhesion.

Figure 4. LOX is essential for FN fibrillogenesis. Cartoon of a FN monomer containing the oxidized lysines (marked by red asterisks) **(A)**. Conservation of the three oxidized lysines among vertebrates **(B)**. Cartoon representing the two FN constructs that were generated **(C)**. Western blot against FN from lysates of cells expressing FN^wt, FN^ΔRGD and FN^ΔOxi **(D)**. Phalloidin staining of FN^WT, FN^ΔRGD and FN^ΔOxi cells seeded on glass without external FN coating and fixed after 1

hour **(E-G)**. Quantification of cell area shows a significant decrease in FN$^{\Delta Oxi}$ cells **(H, n>39)**. Quantification of mean fluorescence intensity of β1-integrin staining per cell shows a significant decrease in FN$^{\Delta Oxi}$ cells **(I, n>20)**. FN immunostaining, color-coded for intensity (blue – low, yellow – high), of cells expressing FN^{WT}, FN$^{\Delta RGD}$ and FN$^{\Delta Oxi}$ after 24 hours of seeding on glass without external FN coating; magnification of representative regions shown below **(J-L)**. Quantification of FN fibrils show a significant decrease in FN$^{\Delta Oxi}$ cells, revealing a critical contribution of LOX family to FN fibrillogenesis **(M, n>71)**. Representative images of cells expressing FN^{WT} and FN$^{\Delta Oxi}$ after 24 hours of seeding on glass without external FN coating, stained for F-actin and FN (color-coded for intensity), showing the lack of FN fibrils even when stress fibers are assembled in FN$^{\Delta Oxi}$ cells **(N)**. Actin and FN immunostaining (color-coded for intensity) of shControl **(O-P)** and shLOX **(Q-R)** HASMC seeded on glass without external FN coating and fixed after 24 hours, demonstrating the inability of shLOX cells to form FN fibrils even when actin stress fibers are assembled. Images on the right are zoom-ins of the orange boxes in each image. ***p<0.0001, n.s. = non-significant.

FN oxidation promotes cancer cells colony formation in soft agar and migration

ECM organization and deposition play key roles in tumorigenesis. Since FN fibrillogenesis underlies these ECM properties and breast cancer progression is highly dependent on FN, we set to test the dependence of breast cancer cell migration and their ability to form colonies on FN oxidation. Towards that end, we stably expressed 67NR non-metastatic breast cancer cells with FN^{WT}, FN$^{\Delta RGD}$ or FN$^{\Delta Oxi}$ (Figure S4) and monitored the above properties. The soft agar colony formation assay serves as one of the hallmarks of carcinogenesis[42] and has been shown to be dependent on FN[43]. Hence, we monitored the ability of these breast cancer cells to form colonies. We find that the colonies over-expressing either FN$^{\Delta RGD}$ or FN$^{\Delta Oxi}$ were significantly smaller than those over-expressing FN^{WT} (Figure 5A-D).

We next set to monitor the effects of these FN-derivatives on cell migration. Quantification of cell migration in a scratch assay over 48 hours (Figure 5E-H) as well as in a single cell migration assay over 6 hours (Figure 5I-N) demonstrated that, as above, the cells over-expressing either of the two FN mutant forms exhibited a reduced cell migration capacity. Altogether, these results strongly demonstrate that the oxidation of FN on the 3 three lysine residues is essential for the cells ability to induce FN fibrillogenesis and as a result to modulate cell behavior.

Figure 5. Mutations in LOX-specific oxidation sites within FN significantly impairs cancer cell characteristics. Representative images of the FN^{WT}, FN$^{\Delta RGD}$ and FN$^{\Delta Oxi}$ colonies formed in soft agar **(A-C)**. Quantification of colonies' diameter show a significant decrease in FN$^{\Delta Oxi}$ cells **(D, n>55)**. Representative images of the FN^{WT}, FN$^{\Delta RGD}$ and FN$^{\Delta Oxi}$ 67NR cell monolayers during wound healing assays **(E-G)**. Cumulative XY graph shows faster wound closure by FN^{WT} cells compared to FN$^{\Delta Oxi}$ and FN$^{\Delta RGD}$ cells, demonstrating the significance of LOX-specific oxidation on cancer cell migration **(H, n>20)**. Representative time color-coded images generated from time-lapse videos tracking fluorescence-labeled nuclei FN^{WT}, FN$^{\Delta RGD}$ and FN$^{\Delta Oxi}$ 67NR cells **(I-K)**. Quantifications of single cell migration parameters show significant decrease in the measured parameters of cells infected with the FN mutants **(L-N, n>330)**. ***p<0.0001, **p<0.001, *p<0.01, n.s. = non-significant.

Discussion:

Our results reveal a novel regulatory step essential for the formation of FN fibrils. We show that the enzymatic activity of lysyl oxidases leads to oxidation of specific lysine residues within FN, resulting in a structural change in the FN dimers thereby promoting early adhesion. We propose that through this structural change, a local increase in RGD density occurs, thus serving as a nucleator of FN fibrillogenesis. Following this nucleation step, integrin-mediated cell adhesion is favored, leading to increased force transmission to the matrix and the subsequent formation of bona-fide FN fibrils (Figure 6). This process could particularly be important for cells to overcome the soft embryonic environment through local secretion of LOX enzymes along migration routes, e.g. in the case of neural crest cells. Thus, similar to nucleation of cytoskeletal fibers' polymerization (e.g. F-actin), which is regulated locally through the activity of nucleators (e.g. formins), initiation of FN fibrillogenesis also requires nucleation. As this nucleation step is an enzyme-dependent process, rather than a force-dependent one, these observations could bridge the observed discrepancies between the *in vivo* observations of FN fibrils in early embryos and in soft tissues, and the cell culture-based models which suggest that fibrillogenesis occurs primarily on stiff matrices.

Our results suggest that in the lack of LOX activity, cells perceive a soft environment even on rigid matrices. The association between LOX activity and rigidity perception highlight an alternative explanation to the observed association of lysyl oxidase upregulation and metastases[44]. Thus, cells initiate migration not only in response to general tissue stiffness but rather due to their local immediate environment, or the perception of their environment, even on soft matrices. As the majority of adhesion assays are carried out on FN coated matrices, we suggest that such conditions mask the LOX-dependent nucleation step and alter the perception of the immediate outside environment. Our results therefore imply that to identify and study the early steps of cell adhesion, cells must be cultured on matrices with either no FN or limiting FN levels. The ubiquitous expression of FN, and the ability of the various widely expressed members of the LOX family to oxidize it, suggest that the process that we have identified is general and is not specific to a distinct tissue or cell type. This process is thus expected to have important

implications for our understanding of various matrix-dependent cellular functions in different tissues and contexts.

Figure 6. Model depicting the stages **(1-3)** required for generating FN clusters that act as nucleators for the force-induced fibrillogenesis step **(4)**. Created with BioRender.com.

Methods:

Mice maintenance and genotyping: All experiments involving mice were performed according to the relevant regulatory standards (Technion IACUC and national animal welfare laws, guidelines, and policies). All mice are housed in IVC's (Techniplast) according to space requirements defined by the NRC. All rooms are set to have 22 ± 2 °C and humidity of 30–70%. HVAC parameters and light cycle are set by the computerized central system to full light of 10 hours, half-light of 2 hours and complete darkness of 12 hours. All mice were bred on a C57Bl/6 background purchased from Envigo (https://www.envigo.com). Embryonic day was staged according to Kaufmann, whereby the morning of the day in which a vaginal plug was observed was marked as E0.5. Mice were genotyped using PCR and the following primers: LoxL3: 5'-GCCAGGGTGAAGTGAAAGAC-3'; 3'-GATCTGGGATGCTGAAGACC-5'; 3'-GAACTTCGGAATAGGAACTTCG-5'. 300bp and 100bp represent wild-type and mutant PCR products, respectively.

Embryos' dissection, imaging and analysis: E9.5 and E18.5 embryos were dissected and underwent over-night fixation after somites were counted. Afterwards, whole mount staining was performed for the E9.5 embryos using FN and ap2α antibodies. Neural crest nuclei in the second branchial arch were then counted from confocal images. For analyses of FN organization in the somite borders, z-stack images were taken. We then used the trainable Weka segmentation plugin[45] of ImageJ (NIH) software to identify FN fibrils in each stack. The area of the quantified fibrils was divided by the total somite area in each stack and the ratios were then averaged. For detection of cleft palate, E18.5 embryos underwent alcian blue and alizarin red staining.

Wholemount embryo staining: Embryos were incubated in 600 μl of blocking buffer (PBS, 0.01% Triton X-100 and 10% non-immune donkey serum (Sigma cat # D9663-10ml) overnight at 4 °C, then with 600 μl of blocking buffer containing primary antibodies for 4 days at 4 °C, with gentle rocking. The following primary antibodies were used: anti-Fn1 (Abcam cat #199056, 1:500), anti-TFAP2α (Developmental Studies Hybridoma Bank, DSHB-3B5, 1:200). After the incubation with primary antibodies, embryos were rinsed and washed with PBST (PBS with 0.05% Triton X-100) for 2 days at 4 °C, with gentle rocking. Embryos were then incubated with 600 μl of blocking buffer containing DAPI

(ThermoFisher, cat #D3571, 5 mg/ml stock diluted 1:1000) and secondary antibodies diluted 1:300 for 4 days at 4 °C. Alexa-labeled secondary antibodies were purchased from Invitrogen (donkey anti-mouse Alexa-647 cat # A31571 and donkey anti-rabbit Alexa-488 cat #A21206). After staining with DAPI and secondary antibodies, embryos were washed with PBST. Prior to imaging, embryos were dehydrated gradually with 25%, 50%, 75% of methanol diluted in dH2O for 10 minutes for each step, and then incubated with 100% methanol twice for 10 min each, and cleared in 50% BABB (BABB and 100% methanol at 1:1 (v/v)), and 100% BABB. BABB was generated by mixing benzyl alcohol (Sigma, #B1042) and benzyl benzoate (Sigma, #B6630) at 1:2 (v/v). Cleared embryos were placed between two coverslips (VWR cat # 16004-312) separated by FastWells spacers (Grace Bio-Labs, cat # 664113). Images were acquired using Nikon A1-HD25 inverted microscope equipped with a 20x water immersion objective (cat # MRD77200), numerical aperture 0.95, working distance 0.95 mm, and the NIS-Elements AR 5.11.01 64-bit software. Optical sections were collected at 0.55 µm intervals.

Cell culture: MEF FN-GFP were described previously[24]. MEF Paxillin-GFP cells were kindly provided by B. Geiger (Weizmann institute of Science, Israel). Dermal fibroblasts were kindly provided by R. Shalom-Feuerstein (Technion, Israel). The three cell lines were cultured in DMEM (Sigma) with 10% FBS, 1% glutamine and 1% pen-strep (Biological Industries). HASMC (Sigma, 354-05A) were cultured with Smooth Muscle Cell Growth Medium 2 (PromoCell, C-22082).

Generation of mutant FN constructs: Point mutations were generated on a human fibronectin pMAX vector plasmid (Addgene, #120402) using the Q5 site-directed mutagenesis kit (BioLabs, #E05545). The mutagenesis procedure was performed using the following primers: Oxidation site 1: 5'-TGAGCGTCCTGCAGACTCCATG-3', 3'-TAAGTGTCACCCACTCGG-5'; RGD: 5'- GCCGTGGAGAAAGCCCCGCAA-3', 3'-CAGTGACAGCATACACAGTGATGGTATAATCAAC-5'; Oxidation site 2: 5'-TGATGATGGGGCGACATACCACG-3', 3'-TAACACGTTGCCTCATGAG-5'; Oxidation site 3: 5'- ACAGTGGCAGGCGGAATATCTCG-3', 3'- TCTCCTACGTGGTATGTC-5'. Next, mutated FN constructs were PCR amplified and ligated into BamH1+Xho1 cut NSPI vector using NEBuilder® HiFi DNA Assembly kit (BioLabs, #E5520).

Lenti viral infection: Plasmids were transfected into HEK 293-FT cells using CalFectin™ Mammalian Cell Transfection Reagent (SignaGen, SL100478). After 48 hours, conditioned medium was collected and used to infect MEF or HASMC cells. Plasmids used: Fibronectin mutant constructs, shLOX (Sigma, TRCN0000045991), shControl (Sigma, SHC016).

Spreading and soft gel experiments: Ibidi 8-well plates (Ibidi, 190814/1) were coated with human plasma FN (hFN, Sigma-Aldrich, FC010-10MG, 10 µg/ml) for 1 hour at 37ºC. Afterwards, the samples were treated with human recombinant LOXL3 (RhLOXL3, R&D systems, 6069-AO) or LOXL2 (RhLOXL2, R&D systems, 2639-AO) for 5 hours (or sodium borate buffer 50 mM as control), with or without 200 mM βAPN (Sigma, A3134-25G) and then washed. Then, cells were seeded on the FN matrix and fixed at varying time points after spreading, followed by immunostaining. For soft gel experiments, Ibidi 8-well plates were covered with silicone gels (DOWSIL, CY 52-276 A&B), prepared at different rigidities that were previously calibrated and described[46] previous to FN coating. Images were acquired using an LSM800 confocal microscope (Zeiss) with Airyscan function and 20X or 63X objectives.

Adhesion, cell area and fibronectin analysis: All of the adhesion and fibronectin fibrils analyses shown in this paper were performed using the trainable Weka segmentation plugin[45] of ImageJ (NIH) software.

Immunofluorescence: Cells were fixed with 4% paraformaldehyde solution (36% stock, Sigma-Aldrich, 50-00-0, diluted with PBS) and permeabilized with 0.1% Triton X-100 in PBS (PBT 0.1%). Primary antibodies were incubated with the fixed cells over night at 4ºC. After, secondary antibodies and phalloidin were incubated for 1 hour at room temperature. For dSTORM imaging, a second 10 minutes fixation was performed at the end of the immunostaining protocol.

Antibodies that were used: anti-total Paxillin (Abcam, ab32084, 1:200); anti-FN (Abcam, ab2413, 1:500); anti-LoxL3 and anti-LoxL2 (produced and kindly gifted by G. Neufeld, Technion). Phalloidin (Abcam, ab176757, 1:1000) and DAPI (Biolegend, 422801, 1:10,000) were also used.

<u>Single cell migration assay:</u> FN was treated with RhLOXL3, with or without βAPN as described above. Then, 24-well cell imaging plates (Eppendorf, EP-0030741005) were coated with the treated or untreated FN. Cells were incubated with 200 nM Sir-DNA (SpiroChrome, SC007) for 6 hours, and were then seeded. Two hours after seeding, a series of 64 images was taken (4 images/h) using the ImageXpress Micro Confocal High-Content Imaging system (Molecular Devices). Nuclei tracking analysis was performed using IMARIS software (Oxford instruments).

<u>dSTORM imaging and analysis:</u> Prior to imaging, FN was treated with RhLOXL3 as described above (molar ratio of 25:1) for 5 hours and was then diluted to 1 ng/ml. Then, #1.5 coverslips were coated with the diluted FN (treated, untreated or fresh FN without any incubation). To detect FN, αFN (Abcam, ab2413) and Alexa-647 conjugated secondary antibodies were used. Fresh dSTORM buffer was prepared as previously described[47]. 2D dSTORM images were taken using a SAFe360 module (Abbelight Ltd, Cachan, France) coupled to Olympus IX83 inverted microscope using a 100× oil-immersion TIRF objective (NA 1.49) and 640 nm laser. The system is equipped with two sCMOS cameras PCO.panda4.2. A total of 15,000 frames at 50 ms exposure time were acquired and used for single-molecule detections to reconstruct a dSTORM image. Resulting coordinate tables and images were processed and analyzed using SAFe NEO software (Abbelight Ltd, Cachan, France).

3D dSTORM images were acquired in a similar manner with the following changes: LoxL3 was detected using anti-LoxL3 antibody. F-actin was detected using Phalloidin-647 (Abcam ab176759). For identification of the Z axis position of each single molecule an astigmatic lens was place in front of the cameras. 40 nm fluorescent beads were added to the sample prior to cell seeding in order to correct the x-y-z drifts.

<u>Pillar array preparation and imaging:</u> Pillar fabrication was performed by pouring Polydimethylsiloxane (PDMS) (mixed at 10:1 with its curing agent, Sylgard 184; Dow Corning) into silicon molds (fabricated as previously described[34]) with holes at fixed depths and distances (center-to-center distance = 1 μm). The molds were then placed, face down, onto glass-bottom 35 mm dishes (#0 coverslip, Cellvis) which were incubated at 65°C for 12h to cure the PDMS. The molds were next peeled off while immersed in ethanol to prevent pillar collapse. On the day of the experiment, FN was treated with

RhLOXL3 for 5 hours (or sodium borate buffer 50 mM as control). Then, the pillars were coated with the treated or untreated FN for 1 hour at 37°c. Next, the buffer was replaced to HBSS + HEPES 20 mM (Biological industries), pH 7.2.

Pillar bending stiffness (k) was calculated by Euler–Bernoulli beam theory as described:

$$k = \frac{3\pi}{64} \frac{ED^4}{L^3}$$

where D and L are the diameter and length of the pillar, respectively, and E is the Young's modulus of the PDMS (= 2 MPa).

Cells were resuspended with the HBSS/HEPES buffer and then spread on the FN-coated pillars. Time-lapse imaging of cells spreading on the pillars was performed using an inverted microscope (Leica DMIRE2) at 37°C using a 63x 1.4 NA oil immersion objective. Brightfield images were recorded every 2 seconds with a Retiga EXi Fast 1394 CCD camera (QImaging). The microscope and camera were controlled by Micromanager software[48]. For each cell, a movie of 10-20 minutes was recorded. To minimize photo-damage to the cells, a 600 nm longpass filter was inserted into the illumination path.

Pillar displacement analysis: Pillar tracking was performed using the Nanotracking plugin of ImageJ, as described previously[32]. In short, the cross-correlation between the pillar image in every frame of the movie and an image of the same pillar from the first frame of the movie was calculated, and the relative x- and y-position of the pillar in every frame of the movie was obtained. To consider only movements of pillar from their zero-position, we only analyzed pillars that at the start of the movie were not in contact with the cell and that during the movie the cell edge reached to them. Drift correction was performed using data from pillars far from any cell in each movie. For each pillar, the displacement curve was generated by Matlab (MathWorks). Analyses of the catch-and-release events was performed by the 'findpeaks' function in Matlab, considering all the peaks above noise level with a minimum width of 3 frames.

Western Blot: Protein lysates were harvested from the cell lines using a lysis buffer (Tris 10 mM pH7, 2 nM EDTA, 1% NP-40, 0.1% DOC, 0.2 mM AEBSF). The lysates were loaded onto SDS-PAGE. Gels were transferred to nitrocellulose membrane that was blocked for 1 hour (5% BSA, 0.1% Tween in TBS). The membranes were then probed

with the primary antibody over night at 4 ºC. Secondary antibody was added for 2 hours at room temperature. Antibodies used for this method: anti-Lox (58135, Cell Signaling); anti-P97 (kindly provided by A. Stanhill, Technion); anti-Fibronectin (Abcam, ab23750).

Neural crest explants: Neural tube explants were carried out essentially as previously described[49]. Briefly, hindbrain regions of E8-8.5 embryos were dissected and the neural primordia consisting of the pre-migratory neural crest cells was isolated from surrounding tissues using fine dissecting pins. The neural tubes were then cultured on FN (50 µg/ml) coated dishes and incubated for 16 hours at 37ºC in DMEM supplemented with 10% fetal bovine serum, 1% 1M HEPES, 1% PenStrep, and 1% L-glutamate (Gibco, USA). Explants were then fixed with 4% PFA and immunofluorescently stained.

Soft agar assay: Protocol was done as previously described[50]. Briefly, the lower layer of 0.5% agarose was prepared using pre-melted 1% agarose and 2x concentrated culture medium and distributed in a 6-well dish. The dish was then incubated for 30 minutes at 4ºC to coagulate the agar. After, the upper layer of 0.3% agarose was prepared using pre-melted 0.6% agarose and 2x concentrated culture medium. 67NR cells were suspended in the upper layer (4000 cells/well) and were distributed above the 0.5% agarose layer that was previously prepared. The 6-well plate was incubated again to coagulate the agar. Then, 300ul of culture medium was added on top of the agar every 3 days. Cells were fixed after 17 days using 10% ethanol in PBS and stained with crystal violet. 4x images of the colonies were taken using light microscope and colonies' diameter was measured.

Wound healing assay: Scratch assay were performed and cell migration was monitored by the automated Incucyte ZOOM (Sartorius, Gottingen, Germany). 67NR cells infected with the FN mutant plasmids were seeded in an Essen Imagelock 96-well plate at the density of 1×60^4 cells/well and were maintained over night to reach 100% confluency. A wound was made in the cell monolayer using a 96-well wound-maker tool with PTFE pin tips (Sartorius) according to the manufacturer's instructions. Afterwards, the plate was inserted into the Incucyte ZOOM and incubated in 5% CO2 at 37°C for 53 hours. Images were captured at 1-hour intervals, and data were analyzed using the IncuCyte Zoom software (Sartorius). Relative wound density (RWD), which measures the relative cell

density in the wounded and non-wounded areas at each time point, was used to assess the rate of cell migration.

<u>Statistical analysis and plotting:</u> Matlab was used for statistical analysis and graph plotting. For all of the boxplots shown in the paper, the red central mark represents the median, the bottom and the top edges of the box represents the 25th and 75th percentiles, respectively, and the black top and bottom represent the minimal and maximal values that are not outliers, respectively. Outliers are represented as red crosses.

Acknowledgments:

We would like to thank Mike Sheetz, Tom Schultheiss and Ella Preger-Ben Noon for critical reading and helpful discussions. H.W. is an incumbent of the David and Inez Myers Career Advancement Chair in Life Sciences. This research was supported by Israel Science Foundation grant 1738/17 (H.W.), Israel Science Foundation grant 111/18 (P.H.), Rappaport Family Foundation (H.W. and P.H.), National Heart, Lung, and Blood Institute grants R01 HL103920, R01 HL134935, R01 HL158049 (S.A.)

Author contributions:

S.M. designed the experiments and methodology, performed the experiments, analyzed the data, and wrote the manuscript. S.Z.E., E.N.B, R.A., H.Z., W.Y.B & R.K.A performed the experiments. H.W. & P.H. were responsible for conceptualization, supervision, funding acquisition, reviewing, writing and editing the manuscript.

Declaration of interests:

Authors declare that they have no competing interests.

Inclusion and diversity:

We support inclusive, diverse, and equitable conduct of research

References:

1. Bonnans, C., Chou, J., and Werb, Z. (2014). Remodelling the extracellular matrix in development and disease. Nat. Rev. Mol. Cell Biol. *15*, 786–801. 10.1038/nrm3904.

2. Humphrey, J.D., Dufresne, E.R., and Schwartz, M.A. (2014). Mechanotransduction and extracellular matrix homeostasis. Nat. Rev. Mol. Cell Biol. *15*, 802–812. 10.1038/nrm3896.

3. Schwarzbauer, J.E., and DeSimone, D.W. (2011). Fibronectins, their fibrillogenesis, and in vivo functions. Cold Spring Harb. Perspect. Biol. *3*, 1–19. 10.1101/cshperspect.a005041.

4. Pankov, R., and Yamada, K.M. (2002). Fibronectin at a glance. J. Cell Sci. *115*, 3861–3863. 10.1242/jcs.00059.

5. Gold, L.I., and Pearlstein, E. (1980). Fibronectin–collagen binding and requirement during cellular adhesion. Biochem. J. *186*, 551–559. 10.1042/BJ1860551.

6. Sabatier, L., Chen, D., Fagotto-Kaufmann, C., Hubmacher, D., McKee, M.D., Annis, D.S., Osher, D.F., and Reinhardt, D.P. (2009). Fibrillin assembly requires fibronectin. Mol. Biol. Cell *20*, 846–858. 10.1091/MBC.E08-08-0830/ASSET/IMAGES/LARGE/ZMK0030989040010.JPEG.

7. Pereira, M., Rybarczyk, B.J., Odrljin, T.M., Hocking, D.C., Sottile, J., and Simpson-Haidaris, P.J. (2002). The incorporation of fibrinogen into extracellular matrix is dependent on active assembly of a fibronectin matrix. J. Cell Sci. *115*, 609–617. 10.1242/JCS.115.3.609.

8. Chung, C.Y., and Erickson, H.P. (1997). Glycosaminoglycans modulate fibronectin matrix assembly and are essential for matrix incorporation of tenascin-C. J. Cell Sci. *110*, 1413–1419. 10.1242/JCS.110.12.1413.

9. Singh, P., Carraher, C., and Schwarzbauer, J.E. (2010). Assembly of Fibronectin Extracellular Matrix. http://dx.doi.org/10.1146/annurev-cellbio-100109-104020 *26*,

397–419. 10.1146/ANNUREV-CELLBIO-100109-104020.

10. Früh, S.M., Schoen, I., Ries, J., and Vogel, V. (2015). Molecular architecture of native fibronectin fibrils. Nat. Commun. *6*. 10.1038/ncomms8275.

11. Zhong, C., Chrzanowska-Wodnicka, M., Brown, J., Shaub, A., Belkin, A.M., and Burridge, K. (1998). Rho-mediated contractility exposes a cryptic site in fibronectin and induces fibronectin matrix assembly. J. Cell Biol. *141*, 539–551. 10.1083/jcb.141.2.539.

12. Antia, M., Baneyx, G., Kubow, K.E., and Vogel, V. (2008). Fibronectin in aging extracellular matrix fibrils is progressively unfolded by cells and elicits an enhanced rigidity response. Faraday Discuss. *139*, 229–249. 10.1039/b718714a.

13. Cao, L., Nicosia, J., Larouche, J., Zhang, Y., Bachman, H., Brown, A.C., Holmgren, L., and Barker, T.H. (2017). Detection of an Integrin-Binding Mechanoswitch within Fibronectin during Tissue Formation and Fibrosis. ACS Nano *11*, 7110. 10.1021/ACSNANO.7B02755.

14. Carraher, C.L., and Schwarzbauer, J.E. (2013). Regulation of matrix assembly through rigidity-dependent fibronectin conformational changes. J. Biol. Chem. *288*, 14805–14814. 10.1074/jbc.M112.435271.

15. Baneyx, G., Baugh, L., and Vogel, V. (2002). Fibronectin extension and unfolding within cell matrix fibrils controlled by cytoskeletal tension. Proc. Natl. Acad. Sci. U. S. A. *99*, 5139–5143. 10.1073/PNAS.072650799/ASSET/5A4D4546-878F-46A5-BC7C-00ACD42B6CE5/ASSETS/GRAPHIC/PQ0726507003.JPEG.

16. Zamir, E., Katz, M., Posen, Y., Erez, N., Yamada, K.M., Katz, B.-Z., Lin, S., Lin, D.C., Bershadsky, A., Kam, Z., et al. (2000). Dynamics and segregation of cell-matrix adhesions in cultured fibroblasts. Nat. Cell Biol. *2*.

17. Wolfenson, H., Lavelin, I., and Geiger, B. (2013). Dynamic Regulation of the Structure and Functions of Integrin Adhesions. Dev. Cell *24*, 447–458.

18. Dzamba, B.J., Jakab, K.R., Marsden, M., Schwartz, M.A., and DeSimone, D.W. (2009). Cadherin Adhesion, Tissue Tension, and Noncanonical Wnt Signaling Regulate Fibronectin Matrix Organization. Dev. Cell *16*, 421–432.

10.1016/J.DEVCEL.2009.01.008.

19. Agero, U., Glazier, J.A., and Hosek, M. (2010). Bulk elastic properties of chicken embryos during somitogenesis. Biomed. Eng. Online *9*, 1–16. 10.1186/1475-925X-9-19.

20. Marrese, M., Antonovaite, N., Nelemans, B.K.A., Smit, T.H., and Iannuzzi, D. (2019). Micro-indentation and optical coherence tomography for the mechanical characterization of embryos: Experimental setup and measurements on chicken embryos. Acta Biomater. *97*, 524–534. 10.1016/J.ACTBIO.2019.07.056.

21. Rifes, P., and Thorsteinsdóttir, S. (2012). Extracellular matrix assembly and 3D organization during paraxial mesoderm development in the chick embryo. Dev. Biol. *368*, 370–381. 10.1016/j.ydbio.2012.06.003.

22. de Almeida, P.G., Pinheiro, G.G., Nunes, A.M., Gonçalves, A.B., and Thorsteinsdóttir, S. (2016). Fibronectin assembly during early embryo development: A versatile communication system between cells and tissues. Dev. Dyn. *245*, 520–535. 10.1002/dvdy.24391.

23. Kraft-Sheleg, O., Zaffryar-Eilot, S., Genin, O., Yaseen, W., Soueid-Baumgarten, S., Kessler, O., Smolkin, T., Akiri, G., Neufeld, G., Cinnamon, Y., et al. (2016). Localized LoxL3-Dependent Fibronectin Oxidation Regulates Myofiber Stretch and Integrin-Mediated Adhesion. Dev. Cell *36*, 550–561. 10.1016/j.devcel.2016.02.009.

24. Tomer, D., Arriagada, C., Munshi, S., Alexander, B.E., French, B., Vedula, P., Caorsi, V., House, A., Guvendiren, M., Kashina, A., et al. (2022). A new mechanism of fibronectin fibril assembly revealed by live imaging and super-resolution microscopy. bioRxiv, 2020.09.09.290130. 10.1101/2020.09.09.290130.

25. Wang, X., and Astrof, S. (2016). Neural crest cell-autonomous roles of fibronectin in cardiovascular development. Dev. *143*, 88–100. 10.1242/DEV.125286/256986/AM/NEURAL-CREST-CELL-AUTONOMOUS-ROLES-OF-FIBRONECTIN.

26. Mittal, A., Pulina, M., Hou, S.Y., and Astrof, S. (2010). Fibronectin and integrin alpha 5 play essential roles in the development of the cardiac neural crest. Mech.

Dev. *127*, 472–484. 10.1016/J.MOD.2010.08.005.

27. Matsumoto, T., Sugita, S., and Nagayama, K. (2016). Tensile Properties of Smooth Muscle Cells, Elastin, and Collagen Fibers BT - Vascular Engineering: New Prospects of Vascular Medicine and Biology with a Multidiscipline Approach. In, K. Tanishita and K. Yamamoto, eds. (Springer Japan), pp. 127–140. 10.1007/978-4-431-54801-0_7.

28. S., L.V., M., H.C., P., H.E., Nikkola, C., Ignaty, L., G., L.C., J., B.A., Dana, V., null, null, P., M.R., et al. (2016). Loss of function mutation in LOX causes thoracic aortic aneurysm and dissection in humans. Proc. Natl. Acad. Sci. *113*, 8759–8764. 10.1073/pnas.1601442113.

29. Sampath Narayanan, A., Siegel, R.C., and Martin, G.R. (1972). On the inhibition of lysyl oxidase by β-aminopropionitrile. Biochem. Biophys. Res. Commun. *46*, 745–751. https://doi.org/10.1016/S0006-291X(72)80203-1.

30. Trappmann, B., Gautrot, J.E., Connelly, J.T., Strange, D.G., Li, Y., Oyen, M.L., Cohen Stuart, M.A., Boehm, H., Li, B., Vogel, V., et al. (2012). Extracellular-matrix tethering regulates stem-cell fate. Nat Mater *11*, 742. 10.1038/nmat3339.

31. Missirlis, D., Haraszti, T., Heckmann, L., and Spatz, J.P. (2020). Substrate Resistance to Traction Forces Controls Fibroblast Polarization. Biophys. J. *119*, 2558–2572. 10.1016/j.bpj.2020.10.043.

32. Wolfenson, H., Meacci, G., Liu, S., Stachowiak, M.R.M.R., Iskratsch, T., Ghassemi, S., Roca-Cusachs, P., O'Shaughnessy, B., Hone, J., and Sheetz, M.P.M.P. (2016). Tropomyosin controls sarcomere-like contractions for rigidity sensing and suppressing growth on soft matrices. Nat. Cell Biol. *18*, 33–42. 10.1038/ncb3277.

33. Feld, L., Kellerman, L., Mukherjee, A., Livne, A., Bouchbinder, E., and Wolfenson, H. (2020). Cellular contractile forces are nonmechanosensitive. Sci. Adv. *6*. 10.1126/SCIADV.AAZ6997.

34. Ghassemi, S., Meacci, G., Liu, S., Gondarenko, A.A., Mathur, A., Roca-Cusachs, P., Sheetz, M.P., and Hone, J. (2012). Cells test substrate rigidity by local contractions on submicrometer pillars. Proc. Natl. Acad. Sci. U. S. A. *109*, 5328–

5333. 1119886109 [pii] 10.1073/pnas.1119886109.

35.   Ghibaudo, M., Saez, A., Trichet, L., Xayaphoummine, A., Browaeys, J., Silberzan,
       P., Buguin, A., and Ladoux, B. (2008). Traction forces and rigidity sensing
       regulate cell functions. Soft Matter *4*, 1836. 10.1039/b804103b.

36.   Jiang, G., Giannone, G., Critchley, D.R., Fukumoto, E., and Sheetz, M.P. (2003).
       Two-piconewton slip bond between fibronectin and the cytoskeleton depends on
       talin. Nature *424*, 334–337. 10.1038/nature01805 nature01805 [pii].

37.   Schvartzman, M., Palma, M., Sable, J., Abramson, J., Hu, X., Sheetz, M.P., and
       Wind, S.J. (2011). Nanolithographic control of the spatial organization of cellular
       adhesion receptors at the single-molecule level. Nano Lett *11*, 1306–1312.
       10.1021/nl104378f.

38.   Cavalcanti-Adam, E.A., Volberg, T., Micoulet, A., Kessler, H., Geiger, B., and
       Spatz, J.P. (2007). Cell spreading and focal adhesion dynamics are regulated by
       spacing of integrin ligands. Biophys. J. *92*, 2964–2974.

39.   Vallet, S.D., Berthollier, C., Salza, R., Muller, L., and Ricard-Blum, S. (2020). The
       Interactome of Cancer-Related Lysyl Oxidase and Lysyl Oxidase-Like Proteins.
       Cancers 2021, Vol. 13, Page 71 *13*, 71. 10.3390/CANCERS13010071.

40.   Parsons, J.T., Horwitz, A.R., and Schwartz, M.A. (2010). Cell adhesion:
       integrating cytoskeletal dynamics and cellular tension. Nat. Rev. Mol. Cell Biol.
       11, 633–643. 10.1038/nrm2957.

41.   Takahashi, S., Leiss, M., Moser, M., Ohashi, T., Kitao, T., Heckmann, D., Pfeifer,
       A., Kessler, H., Takagi, J., Erickson, H.P., et al. (2007). The RGD motif in
       fibronectin is essential for development but dispensable for fibril assembly. J. Cell
       Biol. *178*, 167–178. 10.1083/JCB.200703021.

42.   Du, F., Zhao, X., and Fan, D. (2017). Soft Agar Colony Formation Assay as a
       Hallmark of Carcinogenesis. Bio-protocol *7*, e2351. 10.21769/BioProtoc.2351.

43.   Gupta, R.K., and Johansson, S. (2013). Fibronectin Assembly in the Crypts of
       Cytokinesis-Blocked Multilobular Cells Promotes Anchorage-Independent Growth.
       PLoS One *8*, e72933. 10.1371/JOURNAL.PONE.0072933.

44. Amendola, P.G., Reuten, R., and Erler, J.T. (2019). Interplay Between LOX Enzymes and Integrins in the Tumor Microenvironment. Cancers (Basel). *11*. 10.3390/CANCERS11050729.

45. Chermnykh, E., Kalabusheva, E., and Vorotelyak, E. (2018). Extracellular matrix as a regulator of epidermal stem cell fate. Int. J. Mol. Sci. *19*. 10.3390/ijms19041003.

46. Gutierrez, E., and Groisman, A. (2011). Measurements of elastic moduli of silicone gel substrates with a microfluidic device. PLoS One *6*. 10.1371/journal.pone.0025534.

47. Nahidiazar, L., Agronskaia, A. V., Broertjes, J., Van Broek, B. Den, and Jalink, K. (2016). Optimizing imaging conditions for demanding multi-color super resolution localization microscopy. PLoS One *11*, 1–18. 10.1371/journal.pone.0158884.

48. Edelstein, A., Amodaj, N., Hoover, K., Vale, R., and Stuurman, N. (2010). Computer control of microscopes using microManager. Curr Protoc Mol Biol 92, 14.20.1–14.20.17. 10.1002/0471142727.mb1420s92.

49. Kalev-Altman, R., Hanael, E., Zelinger, E., Blum, M., Monsonego-Ornan, E., and Sela-Donenfeld, D. (2020). Conserved role of matrix metalloproteases 2 and 9 in promoting the migration of neural crest cells in avian and mammalian embryos. FASEB J. *34*, 5240–5261. https://doi.org/10.1096/fj.201901217RR.

50. Borowicz, S., Van Scoyk, M., Avasarala, S., Karuppusamy Rathinam, M.K., Tauler, J., Bikkavilli, R.K., and Winn, R.A. (2014). The soft agar colony formation assay. J. Vis. Exp., e51998. 10.3791/51998.