Dietary fibre controls blood pressure and cardiovascular risk by lowering large intestinal pH and activating the proton-sensing receptor GPR65

Running title: Large intestinal pH, GPR65 and blood pressure

Liang Xie[1, 2], Dakota Rhys-Jones[1], Rikeish R. Muralitharan[1,3], Evany Dinakis[1],

Michael Nakai[1], Madeleine Paterson[1], Alex Peh[1], Hamdi Jama[1], Ekaterina

Salimova[4], Dovile Anderson[5], Caroline Ang[6], Md Jahangir Alam[2], Yu-Anne Yap[7],

Darren Creek[5], Remy Robert[7], CK Yao[8], Daniel So[8], Geoffrey A. Head[9], Peter R.

Gibson[8], Jane Muir[8], Joanne A. O'Donnell[1], Charles R. Mackay[2, 10], Francine Z.

Marques[1, 11*]

Affiliations

[1]Hypertension Research Laboratory, School of Biological Sciences, Monash

University, Clayton, VIC 3800, Australia; [2]Department of Microbiology, Biomedicine

Discovery Institute, Monash University, Clayton, VIC 3800, Australia; [3]Institute for

Medical Research, Ministry of Health Malaysia, Kuala Lumpur, Malaysia; [4]Monash

Biomedical Imaging, Monash University, Clayton, VIC 3168, Australia; [5]Monash

Proteomics and Metabolomics Facility, Monash Institute of Pharmaceutical Sciences,

Monash University, Parkville, VIC 3052, Australia; [6]Department of Biochemistry and

Molecular Biology, Biomedicine Discovery Institute, Monash University, Clayton,

VIC 3800, Australia; [7]Department of Physiology, Biomedicine Discovery Institute,

Monash University, Clayton, VIC 3800, Australia; [8]Department of Gastroenterology,

Central Clinical School, Monash University, Melbourne, 3000, Victoria, Australia;

[9]Neuropharmacology Laboratory, Baker Heart and Diabetes Institute, Melbourne,

3000, Victoria, Australia; [10]School of Pharmaceutical Sciences, Shandong Analysis

and Test Center, Qilu University of Technology (Shandong Academy of Sciences),

Jinan, 250014, China; [11]Heart Failure Research Group Baker Heart and Diabetes

Institute, Melbourne, VIC 3000, Australia.

*Corresponding author: A/Prof Francine Z. Marques, PhD, Hypertension Research

Laboratory, School of Biological Sciences, Faculty of Science, Monash University,

Melbourne, Australia, Phone: +61-03-9905 6958. E-mail:

francine.marques@monash.edu

Abstract

Dietary fibre regulates blood pressure (BP) through gut microbial production of acidic

metabolites known as short-chain fatty acids (SCFAs). The specific mechanisms of

how SCFAs regulate BP are still emerging. We hypothesised that acidic metabolites

that are abundant in the large intestine may activate proton-sensing G-protein coupled

receptors, such as GPR65, thus conferring BP regulating effects. Using mouse models,

we found that dietary fibre levels determined the luminal pH in the large intestine,

through production of SCFAs by the gut microbiota. We then investigated a new

mouse model lacking GPR65, which spontaneously developed higher BP, cardiac and

renal hypertrophy and fibrosis. We identified that low pH, acting via GPR65

signalling, increased cAMP production and phosphorylation of CREB, and regulated

inflammatory cytokine production involved in hypertension. We showed that the

benefits of diets high in fibre, which usually prevent hypertension and its associated

phenotypes, were decreased in mice lacking GPR65. Finally, we provided proof-of-

concept evidence that the luminal pH profile in the colon of hypertensive participants

is higher than that of normotensive participants. Colonic pH was further associated

with dietary fibre, particularly in the colonic regions where fibre is fermented by the

gut microbiota. Together, we show that pH sensing by GPR65 underlies at least some

of the cardiovascular benefits of dietary fibre.

Keywords: hypertension, heart, metabolites, microbiome, pH, G-protein coupled

receptor   65

Introduction

High blood pressure (BP), known as hypertension, accounted for 19.2% of all deaths

(10.8 million deaths) in 2019, primarily due to cardiovascular disease (CVD), being

the leading global disease burden[1]. Insufficient intake of foods high in fibre, such as

vegetables, fruits and wholegrain foods, contributes to mortality and morbidity of

non-communicable diseases (NCDs)[2] resulting from higher BP and CVD[3]. Gut

microbial metabolites, short-chain fatty acids (SCFAs), produced from dietary fibre

fermentation, underlie the cardiovascular benefits conferred by dietary fibre[4,5]. As a

consequence, dietary interventions that increase colonic SCFAs (either via direct

administration or a high fibre diet) have been tested in animal studies[4,5] and, more

recently, in a randomised clinical trial as a novel and safe therapy against

hypertension[6]. SCFAs significantly lowered blood pressure not only in animal

hypertension models[4,5], but in human non-treated essential hypertensive patients as

well[7]. However, the underlying mechanism of how SCFAs lower BP remains to be

fully elucidated. Thus far, studies have focused on the direct role of SCFAs, binding

to GPCRs such as GPR43[4,5]. An important observation that has not been explored yet

is that the large production of SCFAs from bacterial fermentation of dietary fibre in

the colon reduces the luminal pH from ~8 in the distal small intestine to ~6.5[8-10].

Acidification of the colonic pH benefits the host's health through facilitating the

expansion of SCFA-producers and suppressing some gastrointestinal pathogens[11].

Nevertheless, the direct effects of the acidic luminal pH in the colon to the host's

physiology remain largely unknown. The acidic intestinal pH ~6.5 created by SCFAs

may activate proton-sensing receptors, including the G-protein coupled receptor

GPR65[12,13]. A set of genetic variants (e.g., rs3742704, rs8005161) in the human

GPR65 gene are associated with inflammatory bowel disease (IBD)[14-19]. Indeed,

GPR65-deficient mice exhibited exacerbated inflammation in experimental colitis

models[19-21]. More recently, other inflammatory diseases, were associated with the

GPR65 minor T allele of rs8005161 and the minor C allele of rs3742704 in phenome-

wide association studies[18,22]. These genotypes are associated with reduced GPR65

signalling, by ~50% measured by intracellular cAMP accumulation[19] and so provide

an insight into the relevance of GPR65 in human disease. High BP was initially

associated with rs8005161 (P=0.0029) but this was no longer significant after

adjustment by FDR (q=0.300)[18]. A potential explanation for this may be that a large

proportion of hypertensive patients (41% of female patients and 51% of male patients)

are not aware of their elevated BP[23]. Another explanation may be that activation of

GPR65 is not due to intrinsic parameters, such as genotype, and depends on pH.

Independently, GPR65 is highly expressed by various leukocyte subsets[24-26] and

regulates inflammatory responses that are relevant to hypertension. For example,

GPR65 inhibits the secretion of the pro-inflammatory cytokine, TNFα, which has

been implicated in hypertension pathogenesis[27], from $CD4^+$ and $CD8^+$ T cells,

particularly under acidic pH (pH 6.0)[18]. Thus, proton sensing by GPR65 may be an

instrumental link between fibre, colonic luminal pH and cardiovascular health.

Here we aimed to establish the role of colonic pH and the proton-sensing receptor

GPR65 in hypertension and its associated risk factors for CVD, particularly when

fermentable fibre intake is high. We designed a series of studies in mice that provide

evidence for a role of GPR65 in the cardiovascular protection conferred by dietary

fibre and described the mechanism involved. Furthermore, we provided proof-of-

concept that hypertensive patients have higher colonic luminal pH, compared to

normotensives, in the colonic region where dietary fibre is fermented. Collectively,

we provide translational evidence that dietary fibre confers cardiovascular protection

by lowering colonic pH, and activating downstream signalling of the proton-sensor

GPR65 that lowers inflammatory cytokines and inflammation. This represents a novel

way of gut microbiota-to-host communication and may form the basis for new

therapeutic strategies for hypertension.

Results

Dietary fibre lowers large intestinal pH, and this is dependent on the gut

microbiota

The large intestine, including caecum and colon, provides an anaerobic environment

for the microbiota, where fermentation of dietary fibre by the microbiota releases a

large amount of SCFAs, which reduce the local pH[8,9,28]. Therefore, colonic luminal

pH is a quantitative marker for carbohydrate, mostly dietary fibre, fermentation by the

microbiota, particularly bacteria, in the large intestine[29,30]. To determine the impact of

dietary fibre on the colonic luminal pH in the large intestine, we fed C57BL/6J (wild-

type, WT) mice a control diet (AIN93G) and diets high (SF11-025) and low (SF09-

028) in resistant starches (hereby referred to as 'fibre') for 7 days (Fig. 1a). As

expected, these dietary interventions significantly altered the composition of the gut

microbiome (Extended Data Fig. 1a-f), with SCFA producers, such as

Lachnospiraceae-NK4A136-group, being enriched with the high fibre diet (Extended

Data Fig. 1g). Low fibre diet significantly enriched *Alistipes* spp., which is associated

with intestinal inflammation and hypertension[31,32]. The luminal pH of the caecum and

colon was ~7.2 when mice were fed with the control diet (Fig. 1b). Notably, the

caecal and colonic pH were reduced by high fibre diet to 6-6.5 – this was

accompanied by a significant increase in the levels of acetate and total SCFAs in the

colonic contents (Fig. 1c). On the contrary, low fibre diet increased the luminal pH in

the caecum and colon to ~7.5 (Fig. 1b) and reduced the levels of acetate, butyrate,

propionate, which together consist of more than 95% SCFAs produced in the large

intestine[33], and total SCFAs in both the colonic contents and portal vein plasma

relatively to the high fibre intervention (Fig. 1c). However, these short-term dietary

interventions (7 days) did not alter the levels of SCFAs in the peripheral circulation

(Extended Data Fig. 1h), suggesting it is the colon where fiber intake produces

marked effects.

We then aimed to determine whether the pH lowering effect of dietary fibre is

dependent on the gut microbiota. To achieve this, we pre-treated mice with an

antibiotic cocktail to reduce the bacteria load before we started the same dietary

interventions (Fig. 1e). Indeed, the antibiotic cocktail effectively reduced the bacterial

load (Fig. 1f). This reduction in bacteria in the intestine resulted in a higher luminal

pH in the caecum and colon, independent of the level of fibre in the diet (Fig 1g-j).

These data suggest that acidic large intestinal pH produced when fibre is consumed is

dependent on the gut microbiota, particularly bacteria.

Deficiency of GPR65 results in spontaneous cardiovascular dysfunction

The colonic luminal pH produced by high fibre intake, pH 6.5, activates the family of

proton-sensing GPCRs, while that produced by low fibre intake, pH 7.5, silences

them[12,13]. Within this family, GPR65 has functional evidence supporting its protective

roles in gut homeostasis[14-20,22]. Thus, we examined the impact of GPR65 deficiency

within the gut. Using CRISPR/Cas9, we developed a new mouse strain that lacks

GPR65 (*Gpr65$^{-/-}$*). Under normal physiological conditions, *Gpr65$^{-/-}$* mice exhibited

similar gut microbiome composition (Extended Data Fig. 2a-c), immune cell

infiltration (Extended Data Fig. 2d) and tissue histology in the caecum (Extended

Data Fig. 2e-h) and small intestine (data not shown) compared to WT mice.

The role of GPR65 in the cardiovascular system and BP regulation remains

largely unknown. Therefore, we characterized multiple key cardiovascular phenotypes

of *Gpr65$^{-/-}$* mice. Compared to WT mice, male *Gpr65$^{-/-}$* mice had no difference in

body weight (Extended Data Fig. 3a), body composition (Extended Data Fig. 3b-d),

food and water intake (Extended Data Fig. 3e, f), or faeces and urine excretion

(Extended Data Fig. 3g, h). We discovered, however, that *Gpr65$^{-/-}$* mice spontaneously

developed significantly higher systolic BP, diastolic BP and mean arterial pressure

compared to WT mice (Fig. 2a-c). Consistently, *Gpr65$^{-/-}$* mice had significantly

enlarged heart, kidney and spleen (all adjusted to tibia length) compared to WT mice

(Fig. 2d-f). This was accompanied by significantly higher levels of cardiac fibrosis

(Fig. 2g, h), cardiac left ventricular posterior wall in the end diastole (LVPW;d, Fig.

2i) and cardiac output (Fig. 2j). Similarly, renal fibrosis was also significantly

elevated in *Gpr65*[-/-] mice (Fig. 2k, l). To investigate the influence of GPR65

deficiency on renal function, we performed a saline challenge as previously

described[5,34]. Compared to WT mice, *Gpr65*[-/-] mice excreted less water and sodium

over a 5-hour period (Fig. 2m, n), which suggests impaired diuretic and natriuretic

functions. Aligned with the significant functional phenotypes, urine creatinine levels

were significantly increased in *Gpr65*[-/-] mice (Fig. 2o).

We observed similar cardiovascular phenotypes in female *Gpr65*[-/-] mice,

including higher BP (Extended Data Fig. 4a-c), cardiorenal and splenic hypertrophy

(Extended Data Fig. 4d-f), thicker LVPW;d (Extended Data Fig. 4g), and elevated

cardiorenal fibrosis (Extended Data Fig. 4h-k) compared to female WT mice.

Collectively, these results clearly demonstrated that GPR65 deficiency leads to the

spontaneous development of high BP and associated end-organ damage which leads

to CVD.

Acidic pH regulates GPR65 signalling and inhibits pro-inflammatory pathways

A large body of evidence supports that a pro-inflammatory immune system

contributes to the elevation of BP[35,36]. Considering GPR65 expression is particularly

enriched in the immune compartment[24-26], we quantified the number of immune cells

in the spleen, a major lymphoid organ. Using a T cell panel (Extended Data Fig. 5a),

we found that *Gpr65*[-/-] mice had significantly higher numbers of CD45$^+$ immune cells

(Extended Data Fig. 5b), and major T cell subsets, including CD8$^+$ T cells (Fig. 3a),

CD4$^+$ T cells (Extended Data Fig. 5c), γδ T cells in the spleen (Extended Data Fig. 5e,

f). Furthermore, we observed significantly more CCR6$^+$ γδ T cells, which is a

deleterious subset in CVD[37], in the spleen (Extended Data Fig. 5g-i). However, using

a myeloid cell panel (Extended data Fig. 5j), we found that neutrophils and

macrophages/monocytes, and dendritic cells were not enriched in the spleens of

Gpr65[-/-] mice (Extended Data Fig. 5k-o). We then investigated the number of immune

cells in kidney, a critical organ for BP regulation. Flow cytometry with the T cell

panel (Extended Fig. 6a) demonstrated significantly elevated CD45$^+$ immune cells

(Extended Data Fig. 6b), CD8 T cells (Fig. 3b), and other major T cell subsets

(Extended Data Fig. 6c-i) in the kidneys of *Gpr65*[-/-] mice. Expression of an early

activation marker CD69 on T cells in kidney was not influenced by GPR65 deficiency

(Fig. 4b, Extended data Fig. 6d), except on γδ T cells whose expression of CD69 was

promoted in *Gpr65*[-/-] mice (Extended data Fig. 6h). Using the myeloid cell panel

(Extended data Fig. 6j), we found that neutrophils were not significantly increased in

the kidneys of *Gpr65*[-/-] mice, but macrophages/monocytes, particularly Ly6C[-]

macrophages/monocytes and dendritic cells, were elevated in the kidney (Extended

Data Fig. 6l-q) of *Gpr65*[-/-] mice. Thus, these data suggest *Gpr65*[-/-] mice spontaneously

develop a pro-inflammatory profile that is consistent with that observed in

hypertension.

We next aimed to determine whether pH modulates inflammatory responses via

GPR65 signalling. GPR65 is a Gαs-coupled receptor whose activation induces

intracellular cAMP accumulation[38]. cAMP production in response to low pH was

impaired in macrophages of GPR65 deficient mice[39]. We discovered that low (GPR65

activating) pH (6.5) increased the level of intracellular cAMP in splenic cells in a

GPR65 dependent manner (Fig. 3c). The increased intracellular cAMP can lead to the

phosphorylation of cAMP response element-binding protein (CREB). Phosphorylated

CREB (pCREB) has various roles in the immune responses, including suppressing the

production of pro-inflammatory cytokines, IFNγ and TNFα[40], which underlie the

pathogenesis of hypertension[35,36]. T cells, particularly CD8[+] T cells, have a causal role

in hypertension[41]. Thus, we investigated whether pH and GPR65 signalling regulated

the phosphorylation of CREB in splenic T cells *in vitro*. Indeed, compared to pH 7.5,

pH 6.5 significantly elevated pCREB levels in WT splenic CD8[+] T cells activated by

para-Methoxyamphetamine (PMA) and ionomycin (Fig. 3d). The elevation of pCREB

by acidic pH was impaired in *Gpr65[-/-]* splenic CD8[+] T cells (Fig. 3d). In CD4[+] T cells,

acidic pH also elevated pCREB levels in WT mice but the regulation of pCREB by

GPR65 was not observed (Extended Data Fig. 7a). To further characterise the impact

of pH and GPR65 on the effector functions of T cells, we examined their influence on

the production of IFNγ and TNFα by CD8[+] T cells. pH 6.5 inhibited the production of

IFNγ and TNFα by stimulated splenic CD8[+] T cells (Fig. 3e-h) and TNFα but not

IFNγ production by CD4[+] T cells (Extended Data Fig. 7a-d) compared to pH 7.5.

Splenic CD8[+] T cells from *Gpr65[-/-]* mice produced more IFNγ and TNFα after

stimulation, particularly at pH 6.5 (Fig. 3e-h), but we observed no difference in CD4[+]

T cells (Extended Data Fig. 7b-e), consistent with pCREB results. In summary, pH

change, which resembles the regulation of colonic luminal pH by dietary fibre,

regulates GPR65 signalling. Low pH, via GPR65 activation, suppresses the pro-

inflammatory cytokine production by CD8$^+$ T cells, which may reduce BP (Fig. 3i).

Depletion of GPR65 decreased the protection of dietary fibre against high blood

pressure

We next investigated whether pH-sensing by GPR65 conferred the cardiovascular

protection by dietary fibre. To achieve this, we challenged WT and *Gpr65$^{-/-}$* mice with

angiotensin-II (Ang-II, 0.5mg/kg body/weight/day) and fed the mice either high or

low fibre diets (Fig. 4a). As previously reported by us, high fibre diet suppressed the

elevation of systolic BP (Fig. 4b), diastolic BP (Fig. 4c) and mean arterial pressure

(Fig. 4d) induced by Ang-II infusion in WT mice[5]. Such suppression was significantly

lower in *Gpr65$^{-/-}$* mice (Fig. 4b-d, Extended Data Fig. 8a, b). GPR65 deficiency

significantly increased BP with hypertensive stimuli when mice were fed with high

fibre diet, while the impact of GPR65 genotype was negligible under low fibre diet

(Fig. 4b-d, Extended Data Fig. 8a, b). GPR65 deficient mice treated with high fibre

had higher cardiac weight to tibia length ratio (Fig. 4e), thicker LVPW;d (Fig. 4f),

reduced ejection fraction (Fig. 4g) and fractional shortening (Fig. 4h), and higher

cardiac fibrosis (Fig. 4i, j). Similarly, kidney weight to tibia length ratio (Fig. 5a) and

renal fibrosis (Fig. 5b, c) were increased, but not natriuretic or diuretic function (Fig.

5d, e, Extended Data Fig. 8c, d). We quantified the number of immune cells in the

kidneys and observed that high fibre fed $Gpr65^{-/-}$ mice had significantly more CD45^{+}

immune cells compared to the high fibre fed WT mice (Fig. 5f). Of note, the

infiltration of CD8^{+} T cells, whose responses weas regulated by acidic pH and GPR65

signalling, was significantly higher in the kidney of $Gpr65^{-/-}$ mice (Fig. 5g). In

addition, γδ T cells were also increased (Fig. 5h), but there was no change in CCR6^{+}

γδ   T   cells   (Fig.   5i).   Macrophages/monocytes,   particularly   Ly6C^{-}

macrophages/monocytes, were increased in the kidney (Fig. 5j, k), while Ly6C^{+}

macrophages/monocytes, CD4^{+} T cells, Foxp3^{+} regulatory T cells, neutrophils and

dendritic cells were not (Fig. 5l, Extended data 8e-h). However, when mice were fed

with a low fibre diet, which copies a pH where GPR65 is not active, there was no

difference on cardiac and renal parameters besides fibrosis, which was further

exacerbated in $Gpr65^{-/-}$ mice (Fig. 4e-j, 5a-e, Extended Data Fig. 8a-d). Consistent

with this, no difference was observed in immune cell infiltration in the kidney under a

low fibre diet (Fig. 5f-l, Extended Data Fig. 8e-h).

231   We then examined the intestinal tissue of mice fed low and high fibre diets. High

fibre intake thickened the muscularis propria layer and increased the length of

intestinal villi in caecum, irrespective of the genotype for *Gpr65* (Extended Data Fig.

9a-c), but did not influence the width of the fibrotic layer and the number of goblet

cells (Extended Data Fig. 9d, e). However, low fibre or GPR65 deficiency reduced the

number of goblet cells in the duodenum (Extended Data Fig. 9f, j). We also

characterized the gut microbiome composition by analysing 16S rRNA gene. The

main factor driving the differences in gut microbiota was diet rather than the genotype

for *Gpr65* (Extended Data Fig. 10a-e). This was also reflected in the abundance of

bacterial taxa, shown at the genus level (Extended Data Fig. 10f). *Lachnospiraceae-*

NK4A136-group, which was enriched by 7-day high fibre intervention (Extended

Data Fig. 1g), was also more abundant in mice fed high fibre diet in this study

(Extended Data Fig. 10h). The top leading taxonomic change explaining the

differences in gut microbiome was the enrichment of *Bacteroides* in the groups fed

high fibre diet (Extended Data Fig. 10g, h). *Bacteroides* species are the most

predominant anaerobes in the gut[42] with significant cardiovascular benefits[43-46]. Low

fibre diet expanded the genera *Bilophila* and *Alistipes* (Extended Data Fig. 10g, h),

associated with inflammation and higher risk of CVD.[31,32,47,48] Interestingly,

Bacteroides spp. are the dominant carbohydrate fermenters at pH 6.0 to 7.0, the

normal luminal pH within the caecum and colon.[49-51] The major products of

carbohydrate fermentation by *Bacteroides* spp. are propionate and acetate, rather than

butyrate[52,53]. Indeed, compared to low fibre, high fibre diet profoundly elevated

acetate and propionate levels in the caecal contents, independent of the genotype of

the mice, while no significant difference in caecal butyrate levels were observed

(Extended Data Fig. 10i).

The intestinal luminal pH profile in human hypertension: A pilot study

To translate our experimental findings, we then examined colonic luminal pH using a

wireless real-time capsule (SmartPill[TM]) in a pilot cohort study, the pH of Intestines

and Blood-pressure Regulation (pHibre) study (Fig. 6a, Extended Data Table 1).

Although relative to normotensive participants, those with hypertensive had a similar

total dietary fibre intake (Fig. 6b, Extended Data Table 2), dietary fibre intake of the

whole cohort was negatively associated with colonic minimum pH (Fig. 6c), but not

colonic median (Fig. 6d) or maximum pH (Fig. 6e). This is important as GPR65 is

activated by lower pH[13]. We discovered hypertensive patients had higher colonic

minimum pH (Fig. 6f and Extended Data Table 3), without changes in maximum (Fig.

6g) or median pH (Fig. 6h). These were independent of the transit time across the

small and large intestine, which were similar between groups (Extended Data Fig.

11a-c). Thus, we determined that fibre intake was associated with lower colonic

minimum pH, which was higher in hypertensive compared to normotensive

participants.

Most of the dietary fibre fermentation takes place in proximal colonic regions,

such as the ascending and transverse colon[29]. Due to gradual reductions of dietary

fibre availability, bacterial load and activity, the levels of SCFAs in the colon reduce

gradually from the ascending to the descending colon; therefore, the luminal pH

increases (Fig. 6i)[9,54,55]. Such pH elevation is technically difficult to be observed in

small rodents, like mice, due to their small size, but can be investigated using the

SmartPill[TM] in humans[9,56]. Thus, to study changes in pH in the different colonic

regions, we divided the colon into four equal quartiles, as previously reported[9,56] –

these quartiles are thought to approximately correspond to the main regions of the

large intestine. Our data indeed fitted with the expected pattern of pH distribution

along the colon (Fig. 6j). Colon minimum pH was significantly correlated with the

luminal pH in the first and second quartiles (Fig. 6k), consistent with higher fibre

fermentation and production of SCFAs in these proximal colonic regions. We

observed a non-significant increase in pH in hypertensive participants in the second

quartile (P=0.157 after adjustment for age, sex and BMI). This quartile may be

representative of the transverse colon, where slowly fermented fibres, such as

resistant starches, are fermented[29]. These types of fibres, including resistant starch

which was used in our animal studies, are a major source for SCFA-production and

were shown to reduce BP in animal models[4,5] and, more recently, a human clinical

trial[7].

Discussion

Dietary fibre and SCFAs are potent BP-lowering agents. SCFAs, particularly acetate,

propionate and butyrate, are well-known for their immunomodulatory effects through

activation of metabolite-sensing G-protein coupled receptors (GPCRs), such as

GPR41, GPR43 and GPR109A[57]. Some evidence suggests these receptors may confer

cardiovascular protection[5]. Here, we used a translational approach to identify a novel

mechanism of how dietary fibre regulates BP and cardiovascular health via colonic

luminal pH and GPR65. Using animal models, we determined high fibre diet lowers

large intestinal pH by increasing the production of SCFAs by bacteria. This change in

large intestinal pH activates the proton-sensing receptor GPR65 and its downstream

anti-inflammatory signalling cascade. We then used wireless real-time pH monitoring

in normotensive and hypertensive participants diagnosed using ambulatory blood

pressure monitoring. Consistently with our animal findings, we identified an

association between human hypertension and higher colonic minimum pH. Together,

our findings support that large intestinal pH-sensing by GPR65, at least partly,

explains the systemic cardiovascular protection by dietary fibre.

The colon is one of the rare sites in the body where pH is naturally acidic. This

largely results from the microbial production of acidic metabolites, particularly

SCFAs[8-10]. Acidic pH in the colon is essential in maintaining gut homeostasis through

suppressing the growth of pathogens and facilitating the expansion of beneficial

microorganisms, including SCFA producers[11]. Indeed in mice, we used dietary

interventions to show that fibre determined the luminal pH in the colon, and that it

was inversely associated with the levels of SCFAs in both the large intestine and

plasma. Such regulation was dependent on the existence of bacteria in the colon.

Moreover, we validated that total dietary fibre intake was negatively correlated with

the luminal pH in the first regions of the colon in participants of the pHibre cohort.

Among the host's pH sensors, GPR65 has critical roles in gastrointestinal

homeostasis[14-20,22]. In this study, we generated a new systemic GPR65 knockout

mouse model ($Gpr65^{-/-}$) using CRISPR/Cas9 technology. Aligned with a previous

report [21], no clear difference was observed in the gut homeostasis between naïve and

otherwise healthy $Gpr65^{-/-}$ and WT mice. However, characterisation of its

cardiovascular phenotype demonstrated that lack of GPR65 led to the spontaneous

development of cardiovascular disorders featured by higher BP, impaired cardiorenal

function, and cardiorenal remodelling. These results demonstrate GPR65 has a

protective role for cardiovascular health.

GPR65 is particularly enriched in immune cells and lymphoid tissues[24-26].

Evidence built in the last decade strongly support the contribution of inflammation in

the development of hypertension[35,36], making hypertension an unconventional

inflammatory disease. Our study suggested GPR65 deficiency increased immune cell

counts in both spleen and kidneys. T cells, particularly CD8$^+$ T cells, were more

abundant in *Gpr65*[-/-] mice. CD8$^+$ T cells are a well-known immune cell subset

contributing to the pathogenesis of hypertension[41]. Our data demonstrated that low pH,

similar as the large intestinal pH post high fibre intervention, suppresses pro-

inflammatory cytokine production by CD8$^+$ T cells, but not CD4$^+$ T cells, via

activating GPR65 signalling. This is a plausible mechanism linking dietary fibre,

large intestinal pH, GPR65 and cardiovascular protection. CD8$^+$ T cells express

GPR65 at higher levels compared to CD4$^+$ T cells[18,24], which may suggest this

receptor has a more significant role in CD8$^+$ T cells rather than CD4$^+$ T cells. CCR6$^+$

γδ T cells, which are a major initial source of the pro-inflammatory cytokine IL-17A[37],

were significantly more abundant in the spleens and kidneys of *Gpr65*[-/-] mice. GPR65

protects against myocardial infarction in mice, likely through suppressing IL-17A

producing CCR6$^+$ γδ T cells[58]. This may represent another mechanism involved in the

BP regulation by GPR65, which requires further investigation. Thus, we propose that

pH-sensing by GPR65 mediates the cardiovascular protection by dietary fibre via

immunosuppressive mechanisms.

Our study demonstrates that the cardiovascular protection conferred by dietary

fibre is, at least in part, mediated by GPR65. When challenged with Ang-II, *Gpr65*$^{-/-}$

mice had higher BP and cardiorenal hypertrophy compared to WT controls fed high

fibre, and similar BP when fed a low fibre diet, which would also abort GPR65

signalling in the large intestine. Indeed, we observed that high fibre fed *Gpr65*$^{-/-}$ mice

had more CD8$^+$ T cells in the kidney compared to high fibre fed WT mice. Analysis of

the microbiome using 16S rRNA sequencing demonstrated that diet, not *Gpr65*

genotype, drove the difference in the gut microbiota composition. High fibre diet

promoted *Bacteroides* spp., a genus with known cardiovascular benefits[43-46], and

increased the levels of acetate and propionate, but not butyrate, in the intestine.

Moreover, high fibre also inhibited certain taxa associated with inflammation and

cardiovascular disorders, including *Bilophila* and *Aliptipes*[31,32,47,48]. These data

suggested that GPR65 signalling does not regulate gut microbiota composition

directly and its effects in cardiovascular health are dependent on downstream

pathways triggered by GPR65, such as immunomodulation as discussed above.

We acknowledge that since we used a systemic knockout model, our animal

studies cannot exclude the effect of pH-sensing in other tissues. However, the

interventions with low and high fibre diets show robust differences in the phenotype

between WT and *Gpr65*[-/-] knockout mice that support that the importance of GPR65

signalling in hypertension starts in the colon. The *in vitro* experiments used the

luminal colonic pH we determined *in vivo*; however, we acknowledge these may not

correspond to colonic tissue pH. Moreover, the colonic quartiles used in the pHibre

cohort analysis were based on transit time and not geography, and thus, they may not

represent the exact colonic region we attempted to characterise.

In conclusion, this is the first study to report a role of luminal pH in the colon on

the host's cardiovascular health in both humans and mice. We provided new insights

on how dietary fibre and gut microbiota-derived metabolites regulate host health from

the perspective of pH. We established a role for GPR65 as a metabolite-sensing

GPCR by sensing pH changes caused by microbial fermentations. Since pH

alternation is happening universally, this mechanism through proton-sensor GPR65

may extend to broader applications. Agents reducing colonic luminal pH, or agonists

activating GPR65 may be potential tools for the global control of CVDs.

Conflict of Interest

None to declare.

Acknowledgements

This study was supported by a National Heart Foundation Vanguard Grant (102927)

and a National Health & Medical Research Council of Australia (NHMRC) Ideas

Grant (GTN2017382). L.X., M.J.A are supported by Monash Graduate Scholarships.

R.R.M is supported by a scholarship from the Faculty of Science, Monash University.

J.A.O. is supported by a NHMRC Fellowship.   F.Z.M is supported by a Senior

Medical Research Fellowship from the Sylvia and Charles Viertel Charitable

Foundation Fellowship and a National Heart Foundation Future Leader Fellowship

(105663). The Baker Heart & Diabetes Institute is supported in part by the Victorian

Government's Operational Infrastructure Support Program. The authors thank

Department of Gastroenterology, The Alfred, Monash Animal Research Platform,

Monash Genomic Modification Platform, Monash Histology Platform, FlowCore,

Monash Biomedical Imaging, Monash Proteomics and Metabolomics Facility,

Department of Pathology, Monash Health, Australian Genome Research Facility for

their technical supports.

Author contributions

L.X. planned and performed most of the animal in vivo and in vitro experiments, data

analyses, provided intellectual inputs and wrote the manuscript. D.R.J. coordinated

the pHibre and performed human data summarization, SmartPill^{TM} data analyses.

R.R.M., E.D. provided major contribution to animal in vivo experiments. J.A.O.

provided major contribution to pCREB detection. M.P., A.P. performed gut histology

analyses. H.J. contributed to animal in vivo experiments and rodent cardiac

ultrasound data analyses, heart and kidney histology analyses. E.S. performed rodent

cardiac ultrasound scanning and supervised the data analyses performed by L.X.. D.A.

performed SCFA measurement by LC-MS. C.A., M.J.A., Y.A.Y. contributed to animal

in vivo experiments, mouse strain maintenance and cAMP assay. D.C. supervised

SCFA measurement by LC-MS. R.R. contributed to animal in vivo experiments,

mouse strain maintenance and provided intellectual inputs. C.K.Y contributed pHibre

study and supervised SmartPill^{TM} data analyses performed by D.R.J.. D.S., G.A.H.,

P.R.G., J.M. contributed to pHibre study and provided intellectual inputs. C.R.M. and

F.Z.M. conceived the study and supervised the research. F.Z.M. secured the funding

supporting this study. All authors approved of and contributed to the final version of

the   manuscript.

References

1   Murray, C. J. L. *et al.* Global burden of 87 risk factors in 204 countries and

territories, 1990-2019: a systematic analysis for the Global Burden of Disease

Study 2019. *The Lancet* **396**, 1223-1249, doi:10.1016/S0140-6736(20)30752-

2 (2020).

2   Afshin, A. *et al.* Health effects of dietary risks in 195 countries, 1990–2017: a

systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*

393, 1958-1972, doi:10.1016/S0140-6736(19)30041-8 (2019).

3   Reynolds, A. *et al.* Carbohydrate quality and human health: a series of

systematic   reviews   and   meta-analyses.   *Lancet*   **393**,   434-445,

doi:10.1016/S0140-6736(18)31809-9 (2019).

4   Marques, F. Z. *et al.* High-Fiber Diet and Acetate Supplementation Change the

Gut Microbiota and Prevent the Development of Hypertension and Heart

Failure   in   Hypertensive   Mice.   *Circulation*   **135**,   964-977,

doi:10.1161/CIRCULATIONAHA.116.024545 (2017).

5   Kaye, D. M. *et al.* Deficiency of Prebiotic Fiber and Insufficient Signaling

Through Gut Metabolite-Sensing Receptors Leads to Cardiovascular Disease.

Circulation **141**, 1393-1403, doi:10.1161/CIRCULATIONAHA.119.043081

(2020).

6   Rhys-Jones, D. *et al.* Microbial Interventions to Control and Reduce Blood

Pressure in Australia (MICRoBIA): rationale and design of a double-blinded

randomised cross-over placebo controlled trial. *Trials* **22**, 496,

doi:10.1186/s13063-021-05468-2 (2021).

7   Jama, H. A. *et al.* Gut microbial metabolites lower 24-hour systolic blood

pressure in untreated essential hypertensive patients. *medRxiv*,

2022.2006.2020.22276673, doi:10.1101/2022.06.20.22276673 (2022).

8   Campbell, J. M., Fahey, G. C., Jr. & Wolf, B. W. Selected indigestible

oligosaccharides affect large bowel mass, cecal and fecal short-chain fatty

acids, pH and microflora in rats. *J Nutr* **127**, 130-136,

doi:10.1093/jn/127.1.130 (1997).

9   Cummings, J. H., Pomare, E. W., Branch, W. J., Naylor, C. P. & Macfarlane, G.

T. Short chain fatty acids in human large intestine, portal, hepatic and venous

447   blood. *Gut* **28**, 1221, doi:10.1136/gut.28.10.1221 (1987).

448   10   den Besten, G. *et al.* The role of short-chain fatty acids in the interplay

449   between diet, gut microbiota, and host energy metabolism. *Journal of Lipid*

450   *Research* **54**, 2325-2340, doi:10.1194/jlr.R036012 (2013).

451   11   Duncan, S. H., Louis, P., Thomson, J. M. & Flint, H. J. The role of pH in

452   determining the species composition of the human colonic microbiota.

453   *Environ Microbiol* **11**, 2112-2122, doi:10.1111/j.1462-2920.2009.01931.x

454   (2009).

455   12   Ludwig, M. G. *et al.* Proton-sensing G-protein-coupled receptors. *Nature* **425**,

456   93-98, doi:10.1038/nature01905 (2003).

457   13   Wang, J. Q. *et al.* TDAG8 is a proton-sensing and psychosine-sensitive G-

458   protein-coupled receptor. *J Biol Chem* **279**, 45626-45633,

459   doi:10.1074/jbc.M406966200 (2004).

460   14   Franke, A. *et al.* Genome-wide meta-analysis increases to 71 the number of

461   confirmed Crohn's disease susceptibility loci. *Nat Genet* **42**, 1118-1125,

462   doi:10.1038/ng.717 (2010).

15   Ke, X. Presence of multiple independent effects in risk loci of common

complex human diseases. *Am J Hum Genet* **91**, 185-192,

doi:10.1016/j.ajhg.2012.05.020 (2012).

16   Ballester, V. *et al.* Association of NOD2 and IL23R with inflammatory bowel

disease in Puerto Rico. *PLoS One* **9**, e108204,

doi:10.1371/journal.pone.0108204 (2014).

17   Liu, J. Z. *et al.* Association analyses identify 38 susceptibility loci for

inflammatory bowel disease and highlight shared genetic risk across

populations. *Nature Genetics* **47**, 979-986, doi:10.1038/ng.3359 (2015).

18   Xie, L. *et al.* pH and Proton Sensor GPR65 Determine Susceptibility to Atopic

Dermatitis. *The Journal of Immunology*, ji2001363,

doi:10.4049/jimmunol.2001363 (2021).

19   Lassen, K. G. *et al.* Genetic Coding Variant in GPR65 Alters Lysosomal pH

and Links Lysosomal Dysfunction with Colitis Risk. *Immunity* **44**, 1392-1405,

doi:10.1016/j.immuni.2016.05.007 (2016).

20   Tcymbarevich, I. *et al.* Lack of the pH-sensing Receptor TDAG8 [GPR65] in

Macrophages Plays a Detrimental Role in Murine Models of Inflammatory

Bowel Disease. *J Crohns Colitis*, doi:10.1093/ecco-jcc/jjy152 (2018).

21   Chen, X. *et al.* pH sensing controls tissue inflammation by modulating cellular

metabolism and endo-lysosomal function of immune cells. *Nature*

Immunology, doi:10.1038/s41590-022-01231-0 (2022).

22   Diogo, D. *et al.* Phenome-wide association studies across large population

cohorts support drug target validation. *Nature Communications* **9**, 4285,

doi:10.1038/s41467-018-06540-3 (2018).

23   Zhou, B. *et al.* Worldwide trends in hypertension prevalence and progress in

treatment and control from 1990 to 2019: a pooled analysis of 1201

population-representative studies with 104 million participants. *The Lancet*

398, 957-980, doi:10.1016/S0140-6736(21)01330-1 (2021).

24   Heng, T. S. P. *et al.* The Immunological Genome Project: networks of gene

expression in immune cells. *Nature Immunology* **9**, 1091-1094,

doi:10.1038/ni1008-1091 (2008).

25   Uhlen, M. *et al.* Proteomics. Tissue-based map of the human proteome.

Science **347**, 1260419, doi:10.1126/science.1260419 (2015).

26   Thul Peter, J. *et al.* A subcellular map of the human proteome. *Science* **356**,

eaal3321, doi:10.1126/science.aal3321 (2017).

27   Sriramula, S., Haque, M., Majid Dewan, S. A. & Francis, J. Involvement of

Tumor Necrosis Factor-α in Angiotensin II–Mediated Effects on Salt Appetite,

Hypertension, and Cardiac Hypertrophy. *Hypertension* **51**, 1345-1351,

doi:10.1161/HYPERTENSIONAHA.107.102152 (2008).

28   Rubinstein, R., Howard, A. V. & Wrong, O. M. In vivo dialysis of faeces as a

method of stool analysis. IV. The organic anion component. *Clin Sci* **37**, 549-

564 (1969).

29   So, D., Gibson, P. R., Muir, J. G. & Yao, C. K. Dietary fibres and IBS:

translating functional characteristics to clinical value in the era of personalised

medicine. *Gut* **70**, 2383, doi:10.1136/gutjnl-2021-324891 (2021).

30   Farmer, A. D., Mohammed, S. D., Dukes, G. E., Scott, S. M. & Hobson, A. R.

Caecal pH is a biomarker of excessive colonic fermentation. *World J*

Gastroenterol **20**, 5000-5007, doi:10.3748/wjg.v20.i17.5000 (2014).

31   David, L. A. *et al.* Diet rapidly and reproducibly alters the human gut

microbiome. *Nature* **505**, 559, doi:10.1038/nature12820

https://www.nature.com/articles/nature12820#supplementary-information (2013).

32   Kim, S. *et al.* Imbalance of gut microbiome and intestinal epithelial barrier

dysfunction in patients with high blood pressure. *Clinical Science* **132**, 701-

718, doi:10.1042/CS20180087 (2018).

33   Cook & Sellin. Review article: short chain fatty acids in health and disease.

Alimentary Pharmacology & Therapeutics **12**, 499-507,

doi:https://doi.org/10.1046/j.1365-2036.1998.00337.x (1998).

34   Kamat, N. V. *et al.* Renal transporter activation during angiotensin-II

hypertension is blunted in interferon-gamma-/- and interleukin-17A-/- mice.

Hypertension **65**, 569-576, doi:10.1161/HYPERTENSIONAHA.114.04975

(2015).

35   Drummond, G. R., Vinh, A., Guzik, T. J. & Sobey, C. G. Immune mechanisms

of hypertension. *Nature Reviews Immunology*, doi:10.1038/s41577-019-0160-

5 (2019).

36 Norlander, A. E., Madhur, M. S. & Harrison, D. G. The immunology of

hypertension. *The Journal of Experimental Medicine* **215**, 21,

doi:10.1084/jem.20171773 (2018).

37 Yan, X. *et al.* Deleterious Effect of the IL-23/IL-17A Axis and γδT Cells on

Left Ventricular Remodeling After Myocardial Infarction. *Journal of the*

American Heart Association **1**, e004408, doi:10.1161/JAHA.112.004408

(2012).

38 Robert, R. & Mackay, C. R. Gαs-coupled GPCRs GPR65 and GPR174.

Downers for immune responses. *Immunol Cell Biol* **96**, 341-343,

doi:10.1111/imcb.12027 (2018).

39 Mogi, C. *et al.* Involvement of proton-sensing TDAG8 in extracellular

acidification-induced inhibition of proinflammatory cytokine production in

peritoneal macrophages. *J Immunol* **182**, 3243-3251,

doi:10.4049/jimmunol.0803466 (2009).

40 Wen, A. Y., Sakamoto, K. M. & Miller, L. S. The role of the transcription

factor CREB in immune function. *J Immunol* **185**, 6413-6419,

doi:10.4049/jimmunol.1001829 (2010).

41 Trott, D. W. *et al.* Oligoclonal CD8+ T cells play a critical role in the

development of hypertension. *Hypertension* **64**, 1108-1115,

doi:10.1161/HYPERTENSIONAHA.114.04147 (2014).

42 Salyers, A. A. Bacteroides of the human lower intestinal tract. *Annual Review*

of Microbiology **38**, 293-313, doi:10.1146/annurev.mi.38.100184.001453

(1984).

43 Yoshida, N. *et al.* Bacteroides vulgatus and Bacteroides dorei Reduce Gut

Microbial Lipopolysaccharide Production and Inhibit Atherosclerosis.

Circulation **138**, 2486-2498, doi:10.1161/CIRCULATIONAHA.118.033714

(2018).

44 Emoto, T. *et al.* Analysis of Gut Microbiota in Coronary Artery Disease

Patients: a Possible Link between Gut Microbiota and Coronary Artery

Disease. *Journal of Atherosclerosis and Thrombosis* **advpub**,

doi:10.5551/jat.32672 (2016).

45 Emoto, T. *et al.* Characterization of gut microbiota profiles in coronary artery

disease patients using data mining analysis of terminal restriction fragment

length polymorphism: gut microbiota could be a diagnostic marker of

coronary artery disease. *Heart and Vessels* **32**, 39-46, doi:10.1007/s00380-

016-0841-y (2017).

46   Karlsson, F. H. *et al.* Symptomatic atherosclerosis is associated with an altered

gut metagenome. *Nature Communications* **3**, 1245, doi:10.1038/ncomms2266

(2012).

47   Qin, J. *et al.* A metagenome-wide association study of gut microbiota in type 2

diabetes. *Nature* **490**, 55-60, doi:10.1038/nature11450 (2012).

48   Devkota, S. *et al.* Dietary-fat-induced taurocholic acid promotes pathobiont

expansion and colitis in Il10–/– mice. *Nature* **487**, 104-108,

doi:10.1038/nature11225 (2012).

49   Walker Alan, W., Duncan Sylvia, H., McWilliam Leitch, E. C., Child Matthew,

W. & Flint Harry, J. pH and Peptide Supply Can Radically Alter Bacterial

Populations and Short-Chain Fatty Acid Ratios within Microbial Communities

from the Human Colon. *Applied and Environmental Microbiology* **71**, 3692-

3700, doi:10.1128/AEM.71.7.3692-3700.2005 (2005).

50   Ilhan, Z. E. *et al.* pH-Mediated Microbial and Metabolic Interactions in Fecal

Enrichment   Cultures.   *mSphere*   **2**,   e00047-00017,

doi:doi:10.1128/mSphere.00047-17 (2017).

51   Raba, G., Adamberg, S. & Adamberg, K. Acidic pH enhances butyrate

production from pectin by faecal microbiota. *FEMS Microbiology Letters* **368**,

fnab042, doi:10.1093/femsle/fnab042 (2021).

52   Salonen, A. *et al.* Impact of diet and individual variation on intestinal

microbiota composition and fermentation products in obese men. *The ISME*

Journal **8**, 2218-2230, doi:10.1038/ismej.2014.63 (2014).

53   Aguirre, M. *et al.* Diet drives quick changes in the metabolic activity and

composition of human gut microbiota in a validated in vitro gut model.

Research   in   Microbiology   **167**,   114-125,

doi:https://doi.org/10.1016/j.resmic.2015.09.006 (2016).

54   Evans, D. F. *et al.* Measurement of gastrointestinal pH profiles in normal

ambulant human subjects. *Gut* **29**, 1035-1041 (1988).

55 Macfarlane, G. T., Gibson, G. R. & Cummings, J. H. Comparison of

fermentation reactions in different regions of the human colon. *J Appl*

Bacteriol **72**, 57-64, doi:10.1111/j.1365-2672.1992.tb04882.x (1992).

56 Ringel-Kulka, T. *et al.* Altered Colonic Bacterial Fermentation as a Potential

Pathophysiological Factor in Irritable Bowel Syndrome. *Official journal of the*

American College of Gastroenterology | ACG **110** (2015).

57 Tan, J. *et al.* The role of short-chain fatty acids in health and disease. *Adv*

Immunol **121**, 91-119, doi:10.1016/B978-0-12-800100-4.00003-9 (2014).

58 Nagasaka, A. *et al.* The proton-sensing G protein-coupled receptor T-cell

death-associated gene 8 (TDAG8) shows cardioprotective effects against

myocardial infarction. *Scientific Reports* **7**, 7812, doi:10.1038/s41598-017-

07573-2 (2017).

Figures and Figure legends

Fig. 1. Dietary fibre lowers colonic pH through the gut microbiota.

a, The experiment design to characterize the impact of dietary fibre on luminal pH in caecum and colon. **b**, Caecal and colonic luminal pH at the endpoint post dietary interventions. **c**, Levels of acetic, propionic, butyric acids and total short-chain fatty acids (SCFAs) in caecal contents. **d**, Levels of acetic, propionic, butyric acids and total SCFAs in plasma from the portal vein. **e**, The experiment design to characterize the impact of dietary fibre on luminal pH in caecum and colon with antibiotic treatment. Time points with asterisks indicates when faecal samples were collected for bacterial load measurements. **f**, Bacterial load. **g**, Caecal and colonic luminal pH at the endpoint post dietary interventions with antibiotics treatments. **h-j**, Caecal and colonic luminal pH at the endpoint post dietary interventions with/without antibiotics treatments. Each data point represents an individual mouse, and n=6 mice in each group; All data represented as means ± SEM; **c**, **d**, **g**, One-way ANOVA; **h-j**, Student's t test; *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Fig. 2. GPR65 deficiency spontaneously increases tissue hypertrophy and blood

pressure in male mice.

Blood pressure (BP) was measured non-invasively by tail cuff at the age of 10 weeks.

a, Systolic BP. **b**, Diastolic BP. **c**, Mean arterial pressure. Tissues of male mice were

collected and weighed at the age of 12 weeks. **d**, Heart weight to tibia length index. **e**,

Kidney weight to tibia length index. **f**, Spleen weight to tibia length index. Heart

sections were stained with Masson's trichrome. **g**, Representative heart sections; For

upper panels, magnification = ×10, scale bar = 1mm; For lower left panels

(interstitial), magnification = ×200, scale bar = 50μm; For lower right panels

(perivascular), magnification = ×100, scale bar = 100μm. **h**, Percentage of fibrotic

area in the cardiac tissues. Cardiac ultrasound screening was performed at the age of

11 weeks. **i**, Left ventricle posterior wall in the end diastole. **j**, Cardiac outputs.

Kidney sections were stained with Masson's trichrome. **k**, Representative kidney

sections; For left panels, magnification= ×5, scale bar = 2mm; For right panels,

magnification = ×200, scale bar = 50μm. **l**, Percentage of fibrotic area in the renal

fibrosis. Saline challenge was performed at the age of 11 weeks. Mice were

intraperitoneally injected with 10% of their body weight of 37 °C 0.9% saline solution.

m, Renal diuretic functions measured by accumulated water excretion over 5 hours

post injection. **n**, Renal natriuretic functions measured accumulated sodium excretion

over 5 hours post injection. Urine samples were collected from 24h metabolic cage. **o**,

The level of creatinine in urine. For the left figures of panels **m** and **n**, n=10-14 mice

in each group; All data represented as means ± SEM; Two-way ANOVA. For other

panels, each data point represents an individual mouse, and n=6-26 mice in each

group; All data represented as means ± SEM; **a-c**, **h**, **j**, **l** and **o**, Student's t test; **d-f**, **i**,

Mann-Whitney test. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Fig. 3. pH and GPR65 signalling modulates immune responses related to

hypertension pathogenesis.

Single cell suspensions were prepared from spleens or kidneys of male mice at the

age of 12 weeks. **a**, CD8$^+$ T cell count in the spleen. **b**, CD8$^+$ T cell count and

percentage of CD69$^+$ cells from CD8$^+$ T cells in the kidney. Splenocytes were isolated

from mice at the age of 8-10 weeks and incubated under either pH 6.5 or pH 7.5 for

30 min. **c**, Levels of cAMP in the cell lysate extracts. 1×10^6 splenocytes were

incubated with 100ng/ml PMA and 1µg/ml ionomycin for 2 hours. **d**, Representative

histogram of pCREB (Ser133) levels of CD8$^+$ T cells and bar plot of pCREB (Ser133)

levels of CD8$^+$ T cells. 1×10^6 splenocytes were incubated with 100ng/ml PMA and

1µg/ml ionomycin for 4 hours. 1×10^6 splenocytes were incubated with 100ng/ml PMA

and 1µg/ml ionomycin for 4 hours. **e**, Representative pseudocolor plots for IFNγ

producing WT and *Gpr65$^{-/-}$* CD8$^+$ T cells at pH 6.5 and 7.5 (Gated from CD8$^+$ T cells).

f, Percentage of IFNγ producing CD8$^+$ T cells at pH 6.5 and 7.5 among WT and

Gpr65$^{-/-}$ CD8$^+$ T cells. **g**, Representative pseudocolor plots for TNFα producing WT

and *Gpr65$^{-/-}$* CD8$^+$ T cells at pH 6.5 and 7.5 (Gated from CD8$^+$ T cells). **h**, Percentage

of TNFα producing CD8$^+$ T cells at pH 6.5 and 7.5 among WT and *Gpr65$^{-/-}$* CD8$^+$ T

cells. **i**, Graphic summary of how pH regulates inflammatory responses contributing

to hypertension in CD8$^+$ T cells. Created with BioRender.com. Each data point

represents an individual mouse, and n=4-9 mice in each group; All data represented as

means ± SEM. **a**, **b**, Student's t test. **c**, **f**, **h**, Two-way ANOVA. *P<0.05, ***P<0.001,

***P<0.0001.

Fig. 4. Deficiency of GPR65 decreased the cardiac protective effects of dietary

fibre.

6-week-old male WT and *Gpr65*[-/-] were infused with 0.5mg/kg body weight/day

angiotensin-II for 28 days and fed with either HF or LF diet. **a**, The experiment

design. **b**, Systolic BP profile over 4 weeks of infusion; **c**, Diastolic BP profile over 4

weeks of infusion; **d**, Mean arterial pressure profile over 4 weeks of infusion; **e**, Heart

weight to tibia length index; Cardiac ultrasound scanning was performed at the week

4 of infusion. **f**, Left ventricle posterior wall in the diastole; **g**, Cardiac ejection

fraction; **h**, Cardiac fractional shortening; Heart sections were stained with Masson's

trichrome. **i**, Percentage of fibrotic area in the cardiac tissues. **j**, Representative heart

sections; For upper panels, magnification = ×10, scale bar = 1mm; For lower left

panels (interstitial), magnification = ×200, scale bar = 50μm; For lower right panels

(perivascular), magnification = ×100, scale bar = 100μm. Each data point represents

an individual mouse, and n=8-11 mice in each group; All data represented as means ±

SEM. Two-way ANOVA. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Figure 5. Deficiency of GPR65 impaired renal function and promoted renal inflammation post-hypertensive stimuli in mice fed with high fibre diet.

Kidneys were collected at the endpoint of the study. **a**, Kidney weight to tibia length index; Kidney sections were stained with Masson's trichrome. **b**, Percentage of fibrotic area in the renal tissues. **c**, Representative kidney sections; For left panels, magnification= ×5, scale bar = 2mm; For right panels, magnification = ×200, scale bar = 50μm. Mice were intraperitoneally injected with 10% of their body weight of

37 °C 0.9% saline solution at the 3^{rd} week post minipump implantation. **d**, Renal

diuretic functions measured by accumulated water excretion at the 5^{th} hour post

injection. **e**, Renal natriuretic functions measured accumulated sodium excretion at

the 5^{th} hour post injection. Single cell suspensions were prepared from kidneys of the

mice at the endpoint of the study. **f**, $CD45^+$ immune cell count; **g**, $CD8^+$ T cell count;

h, γδ T cell count; **i**, $CCR6^+$ γδ T cell count; **j**, Macrophage/Monocyte count; **k**, $Ly6C^-$

Macrophage/Monocyte count; **l**, $Ly6C^+$ Macrophage/Monocyte count. Each data point

represents an individual mouse, and n=6-11 mice in each group; All data represented

as means ± SEM. Two-way ANOVA. *$P<0.05$, **$P<0.01$, ***$P<0.001$.

Fig. 6. The intraluminal pH profile within the colon of normotensive and hypertensive participants.

a, Recruitment summary. Out of 91 expressions of interest, 55 participants have finished the trial so far. GI, gastrointestine; COVID, coronavirus

disease; BMI, body mass index; BP, blood pressure. The demographic summary and the relevant clinical information collected at the initial

screening is in Extended Data Table 1. **b**, Daily total dietary fibre intake. **c-e**, Correlations of daily total dietary fibre intake with colon minimum,

median and maximum pH generated by the SmartPill^{TM}, showing that fibre intake affected the colonic minimum pH the most. **f**, Colon minimum

pH, **g**, colon median pH, and **h**, colon maximum pH in normotensive (n=15) versus hypertensive participants (n=40). **i,** Graphic scheme of

regional pH profile in colon and respective bacterial activity, short-chain fatty acids (SCFAs) concentration. Adapted from [9,55] (Created with

BioRender.com). **j**, Regional pH profile in the colon of the pHibre cohort generated by the SmartPill^{TM}. **k**, Correlations of colon minimum pH

with luminal pH in different quartiles of colon. Each data point represents an individual participant. All data represented as mean ± SEM; P-

values from correlations, one-way ANOVA, or regression analysis adjusted for sex, age and BMI (n=55).