Dietary fiber controls blood pressure and cardiovascular risk by lowering large intestinal pH and activating the proton-sensing receptor GPR65

Running title: Colonic pH, GPR65 and blood pressure

Liang Xie[1,2,3], Rikeish R. Muralitharan[1], Evany Dinakis[1], Simona Antonacci[1], Kwan

Charmaine Leung[1], Zoe McArdle[4], Katrina Mirabito Colafella[4], Michael Nakai[1],

Madeleine Paterson[1], Alex Peh[1], Hamdi Jama[1], Ekaterina Salimova[5], Dovile Anderson[6],

Caroline Ang[7], Md Jahangir Alam[2,7], Yu-Anne Yap[7], Darren Creek[6], Remy Robert[7],

Joanne A. O'Donnell[1], Charles R. Mackay[8], Francine Z. Marques[1,9*]

Affiliations

[1]Hypertension Research Laboratory, Victorian Heart Institute and Department of

Pharmacology, Biomedical Discovery Institute, Faculty of Medicine, Nursing and Health

Sciences, Monash University, Clayton VIC 3800, Australia; [2]Department of Microbiology,

Biomedicine Discovery Institute, Monash University, Clayton, VIC 3800, Australia;

[3]Precision Medicine Translational Research Programme, Department of Obstetrics &

Gynaecology, Yong Loo Lin School of Medicine, National University of Singapore,

Singapore; [4]Cardiovascular Disease Program, Biomedicine Discovery Institute and

Department of Physiology, Faculty of Medicine, Nursing and Health Sciences, Monash

University, Clayton VIC 3800, Australia; [5]Bioimaging Platform, La Trobe University,

Melbourne Australia, Monash Biomedical Imaging, Monash University, Clayton, VIC 3168,

Australia; [6]Monash Proteomics and Metabolomics Facility, Monash Institute of

Pharmaceutical Sciences, Monash University, Parkville, VIC 3052, Australia; [7]Department of

Physiology, Biomedicine Discovery Institute, Monash University, Clayton, VIC 3800,

Australia; [8]School of Pharmaceutical Sciences, Shandong Analysis and Test Center, Qilu

University of Technology (Shandong Academy of Sciences), Jinan, 250014, China; [9]Baker

Heart and Diabetes Institute, Melbourne, VIC 3000, Australia.

***Corresponding author**: Prof Francine Marques, Hypertension Research Laboratory,

Victorian Heart Institute, Level 2, Victorian Heart Hospital, 631 Blackburn Road,

Clayton, VIC 3168 Monash University, Melbourne, Australia, Phone: +61-03-9905 6958. E-

mail: francine.marques@monash.edu

Abstract

High blood pressure (BP) is the most common cause of death globally, due to

increasing the risk of cardiovascular diseases. Dietary fiber regulates BP through gut

microbial production of acidic metabolites known as short-chain fatty acids (SCFAs).

The specific mechanisms of how SCFAs regulate BP are still emerging. In a

phenome-wide association study, we identified that the proton-sensing G-protein-

coupled receptor *GPR65* gene is associated with hypertension and its associated end-

organ damage phenotypes. We hypothesized that acidic metabolites produced from

the gut microbiota may activate GPR65, thus conferring BP regulating effects. We

found that dietary fiber levels determined the luminal and interstitial tissue pH in the

large intestine through production of SCFAs by the gut microbiota. We identified that

low pH produced by high fiber intake, acting via GPR65 signaling, increased cAMP

production and phosphorylation of CREB, and restricted the production of

hypertension-promoting inflammatory cytokines by CD8$^+$ T cells. *Gpr65$^{-/-}$* mice

spontaneously developed higher BP, cardiac and renal hypertrophy and fibrosis. We

showed that the benefits of a diet high in fiber, which prevented hypertension and

associated end-organ damage, were decreased in *Gpr65$^{-/-}$* mice. Finally, adoptive

transfers revealed that GPR65 deficiency in CD8$^+$ T cells causally explained this

phenotype. In conclusion, we showed that pH sensing by GPR65 in CD8+ T cells

mediates much of the cardiovascular benefits of dietary fiber. pH sensing represents a

novel gene-by-environment interaction of gut microbiota-to-host biological effects

and may form the basis for new therapeutic strategies for hypertension.

Keywords: hypertension, metabolites, microbiome, pH, G-protein coupled receptor

Introduction

Hypertension accounted for 19.2% of all deaths (10.8 million deaths) worldwide in 2019, primarily due to cardiovascular disease (CVD).[1] Insufficient intake of foods high in dietary fiber, such as vegetables, fruits and wholegrain foods, contributes to mortality and morbidity of non-communicable diseases[2] resulting from higher blood pressure (BP) and CVD.[3] Gut microbial metabolites, called short-chain fatty acids (SCFAs), produced upon dietary fiber fermentation, underlie the cardiorenal benefits conferred by dietary fiber.[4-7] As a consequence, dietary interventions that increase colonic SCFAs (either via direct administration of SCFAs or a high fiber diet) have been reported to reduce BP significantly not only in experimental hypertensive animals[4,5] but, more recently, in untreated essential hypertensive patients in a phase II randomized clinical trial.[8] However, the underlying mechanism of how SCFAs lower BP remains to be fully elucidated. Thus far, studies have focused on the direct role of SCFAs, via binding to and activating G-protein coupled receptors (GPCRs) such as GPR41 and GPR43.[4,5,9,10]

An important observation that has not been explored yet is that the production of SCFAs from the bacterial fermentation of dietary fiber in the colon reduces the luminal pH in the distal small intestine from ~8 to ~6.5.[11-13] Such acidic intestinal pH ~6.5 created by SCFAs may activate proton-sensing receptors, including the G-protein-coupled receptor GPR65.[14-17] GPR65 is critical for gut homeostasis. A set of genetic variants (e.g., rs3742704, rs8005161) in the human *GPR65* gene are strongly associated with inflammatory bowel disease (IBD),[18-23] and GPR65-deficient mice

exhibited exacerbated inflammation in experimental colitis models.[23-25] In addition,

the *GPR65* minor T allele of rs8005161 was associated with high BP in a phenome-

wide association study (PheWAS).[22] These genotypes are associated with ~50%

reduced GPR65 signaling,[23] and provide an insight into the relevance of GPR65 in

human disease. We validated these results in an independent PheWAS using the Atlas

of GWAS Summary Statistics data, showing that *GPR65* is associated with

hypertension, cardiac hypertrophy and function (e.g., ejection fraction, fractional

shortening), and heart failure (Table S1).[26] Moreover, GPR65 is highly expressed by

various leukocyte subsets[27-29] and regulates inflammatory responses relevant to

hypertension, such as inhibiting inflammatory cytokine production.[22] Therefore,

proton sensing by GPR65 may be an instrumental link between fiber, colonic luminal

pH, and cardiovascular health.

Here we aimed to establish the role of colonic pH and the proton-sensing receptor

GPR65 in hypertension and its associated end-organ damage that are risk factors for

CVD, particularly when fermentable fiber intake is high. We conducted a series of

studies in mice that provided evidence for a novel role of GPR65 in the

cardiovascular protection conferred by dietary fiber. We demonstrate that dietary fiber

confers cardiovascular protection by lowering colonic pH and activating downstream

signaling of the proton-sensor GPR65 that lowers inflammation via inhibiting

inflammatory cytokines produced by CD8+ T cells.

Methods

For detailed description of Methods, please see the Online Supplemental Files.

Animal studies

All animal care and experimental procedures used in this study were approved by the Animal Ethics Committee of Monash University (17465, 27929, 37720). Wild-type (WT) C57BL/6J mice were obtained from the Monash Animal Research Platform, Monash University, housed in the same facility under the same conditions. $Gpr65^{-/-}$ mice on a C57BL/6J background were generated by using CRISPR/Cas9-based protocol at the Monash Genome Modification Platform (MGMP) at Monash University. $Gpr65^{gfp/gfp}$ mice were purchased from The Jackson Laboratory (stock number 008577). $Gpr65^{gfp/gfp}$ mice have 90% of exon 2 coding sequences replaced by promoterless IRES-EGFP sequences to disrupt GPR65 function.[30] $Gpr65^{gfp/+}$ mice were generated by crossing $Gpr65^{gfp/gfp}$ mice with WT C57BL/6J mice. $Rag1^{-/-}$ mice on a C57BL/6J background were obtained from the Clive and Vera Ramaciotti Laboratories, Walter and Eliza Hall Institute of Medical Research. All mice were maintained under specific pathogen-free and controlled environmental conditions.

     Murine diets used in this study include: i) Normal chow diet (Barastoc rat and mouse pellets; Ridley); ii) AIN93G (Specialty Feeds); iii) a diet lacking resistant starches (low fiber, LF, SF09-028; Specialty Feeds); iv) a diet rich in resistant starches (high fiber with all carbohydrates replaced with resistant starches, HF, SF11-025; Specialty Feeds) (Table S2). To eliminate the gut microbiota, mice were treated with a combination of enrofloxacin (Baytril, 10 mg/kg body weight/ day) and amoxicillin

with clavulanic acid (50 mg/kg body weight/ day) in drinking water for a week before

the experimental procedures and maintained on this antibiotic cocktail until the

endpoint.

Tissue Collection

Upon euthanasia by CO_2 asphyxiation, blood and various tissues (heart, kidney, liver,

gut, cecal content, spleen) were collected for further analyses. Samples were either

snap-frozen using liquid nitrogen and stored at -80 °C for subsequent DNA/RNA

extractions, stored in formalin before paraffin embedding for histopathology studies,

or kept in cold phosphate buffer saline (PBS) to be used fresh for flow cytometry

studies.

Large intestinal intraluminal and interstitial fluid pH measurement in mouse

The pH was measured in samples collected from the cecum and colon using a pre-

calibrated Thermo-Scientific Orion Star A211 pH meter as previously described.[31,32]

Intestinal interstitial fluid pH was measured using a protocol generously shared by

Professor Dominik Muller (Max Delbrück Center for Molecular Medicine in the

Helmholtz Association, Berlin, Germany), adapted from [33]. The same Orion PerpHec

ROSS Combination pH Micro Electrode (Thermo-scientific, diameter 3mm) was used

to measure the pH of the intestinal interstitial fluid. We ensured the sample covered

the probe tip and acquired a stable reading.

Blood pressure measurement

BP was measured using radiotelemetry (probe TA11PA-C10, Data Sciences

International, using the Ponemah software, Data Sciences International) and tail-cuff

in a CODA non-invasive blood pressure system (Kent Scientific Corporation), as

previously decribed.[10,34]

CD8$^+$ T cells adoptive transfer

To isolate CD8$^+$ T cells, single-cell suspensions of splenocytes were subject to

Dynabeads™ Untouched™ Mouse CD8 Cells Kit (Thermo Fisher Scientific) for the

isolation of CD8$^+$ leukocytes according to the manufacturer's manuals. $5x10^6$ of either

WT or *Gpr65$^{-/-}$* CD8$^+$ T cells were resuspended in 200μL PBS and intravenously

injected into *Rag1$^{-/-}$* mice.

Angiotensin-II induced hypertension model

Six-week-old male WT and *Gpr65$^{-/-}$* mice fed low or high-fiber diets received a

subcutaneous minipump (Alzet model 2004) containing angiotensin II (Ang II, 0.5

mg/kg body weight/day; Auspep) placed during surgery under anesthesia with

isoflurane. Data were collected for 4 weeks after the implant. Littermate mice were

randomized into either group using an Excel randomization tool.

For the adoptive transfer model, one day after the adoptive transfer of the CD8$^+$ T

cells, ten-week-old male *Rag1$^{-/-}$* recipient mice underwent a subcutaneous minipump

(Alzet model 1002) implantation containing Ang II at 1.44 mg/kg body weight/day

(Auspep) for 2 weeks as described above. Mice were randomized into either group

using an Excel randomization tool.

Metabolic characterization

Mice were individually housed in metabolic cages for 24 hours. Body weight, food

intake, water intake, urine excretion and feces excretion were measured. Mice body

composition was analyzed by an EchoMRI 3-in-1 machine (Houston, TX).

Saline challenge studies

135 To examine if GPR65 deficiency affected diuretic and natriuretic responses, we

136 performed a saline challenge test according to the protocol kindly shared by Prof.

137 Alicia McDonough (Keck School of Medicine of the University of Southern

138 California).[35,36]

Cardiac ultrasound

140 The day before the endpoint, mice were anesthetized using isoflurane, and B- and M-

141 mode echocardiography was performed to image the left ventricle using a Vevo

142 2100 Ultra High Frequency ultrasound system and an MS550D transducer by an

143 experienced user at the Monash Biomedical Imaging Centre (E.S.).

Histological analysis

145 Mouse heart, kidney, and gut tissues were fixed in 10% neutral buffered formalin for

146 24-48 hours, processed, paraffin-embedded and sectioned at 4 μm. Masson's

147 trichrome staining was performed to analyze the collagen present in heart, kidney, and

148 intestinal samples. Periodic acid-Schiff alcian blue (PAS/AB) staining was performed

149 to analyze goblet cells in the gut.

Flow cytometry

Single cell suspensions were isolated from mouse spleen, kidney, and colon (See

Online Supplemental files for details). Cells were resuspended with FACS buffer

(PBS containing 2% FCS and 4 mM EDTA). Cell viability was evaluated using the

fixable viability stain 620 (BD Horizon). After blocking Fc receptors with mouse FcR

blocking reagents (Miltenyi Biotech) for 15 mins at room temperature, single-cell

suspensions were surface-stained at 4°C for 30 min. Cells were then fixed and

permeabilized at 4°C using a Foxp3/Transcription Factor Staining Buffer Set

(eBioscience). Intracellular staining was then performed. Cell counts were determined

using flow cytometry counting beads (CountBright Absolute; Life Technologies)

following manufacturer instructions. Flow cytometry was performed to quantify

tissue-specific immune cells and assess pCREB levels and cytokine production (See

Online Supplemental Files for details). Sample data were acquired using a five-laser

BD LSRFortessa X-20 flow cytometer and BD FACSDiva software (BD Biosciences)

and analyzed using FlowJo software (Tree Star).

In vitro **cell stimulation**

Mouse splenic or colon lamina propria cells were cultured in RPMI1640 medium

supplemented with 10% FBS and 4 mM L-glutamine, which was altered to pH 6.5 or

pH 7.5 using HCl and NaOH. Cells were stimulated with 100 ng/ml PMA (Sigma-

Aldrich) and 1 mg/ml ionomycin (Sigma-Aldrich) for 2-h for pCREB detection, and

for 4-h for intracellular cytokine analyses.

cAMP assay

1×10^6 splenic cells were resuspended in 200 µL RPMI1640 supplemented with 10%

FBS and 4 mM L-glutamine at pH 6.5 or pH 7.5. After incubating at 37°C for 30 mins,

cells were washed with PBS at pH 6.5 or pH 7.5, respectively. Intracellular cAMP

levels were measured by Invitrogen cAMP Competitive ELISA kit (ThermoFisher,

#Cat: EMSCAMPL), following the manufacturer's instructions.

DNA extraction from cecal contents and feces

DNA extraction of cecal/fecal contents was performed using the DNeasy PowerSoil

DNA isolation kit (Qiagen, Germany). DNA samples were quantified using Nanodrop

(Thermo Fisher Scientific).

Mouse fecal bacterial load measurement

Absolute bacterial load was measured using quantitative real-time PCR. Briefly, we

used Fast SYBRgreen (Thermo Scientific) and 16S primers 1114F/1221R in a

QuantStudio 7 qPCR Instrument (Thermo Scientific). Samples were loaded at

10ng/well, amplified in triplicates and compared to a standard containing 10^2 to 10^{10}

bacterial copies.

Short-chain fatty acids quantification by liquid chromatography-mass

spectrometry (LC-MS)

Quantification of SCFAs including acetate, propionate, butyrate, isobutyrate,

isovalerate, valerate and caproate was performed using LC-MS method, as previously

described (See Online Supplemental Files for details).[37]

Animal statistical analyses

All graphical representation of data was done using Prism 8.2.1 (GraphPad Software,

La Jolla, CA). The normality of the distribution of continuous data was assessed by

the Shapiro-Wilk test. Comparison between multiple groups of different treatments

and different mouse strains was analyzed using a two-way analysis of variance (not

for repetitive measures for 4 groups or for repetitive measures for weekly BP) with

adjustment using Tukey's multiple comparisons test. Data is presented as mean ±

standard error of the mean. *P*<0.05 was considered significant. All data was collected

and analyzed blindly, where possible.

16S ribosomal RNA sequencing

V4 region of the 16S ribosomal RNA (rRNA) was targeted and amplified using

primers previously published.[38] 240ng of the amplicon product per sample were

pooled and purified. Sequencing was performed at the Australian Genomic Research

Facility (AGRF, Melbourne, Australia) on MiSeq instrument (Illumina) to generate

300bp paired-end reads. Sequencing followed our protocol published elsewhere.[39]

Results

Dietary fiber lowers large intestinal pH, and this is dependent on the gut

microbiota

Mice intestinal pH profiles have mostly been reported under normal conditions

without dietary interventions[32] and have not been well characterized, particularly for

the large intestine. To determine the impact of dietary fiber on the large intestinal

luminal and interstitial tissue pH, we fed WT mice a control diet and diets high and

low in fermentable fiber for 7 days (Figure 1A). Besides dietary fiber, the nutritional

components of these diets are almost identical (Table S2). The cecum and colon

luminal pH was ~7.2 when mice were fed the control diet (Figure 1B, Figure S1A).

Notably, the cecal and colonic pH were reduced by high fiber diet to 6 and 6.5,

respectively (Figure 1B). This change in pH was mirrored by changes in colonic

interstitial fluid pH (Figure 1B). This was further accompanied by a significant

increase in the acetate and total SCFA levels in the colonic contents (Figure 1C). On

the contrary, relative to a high fiber diet, a low fiber diet increased the luminal pH in

the colon to ~7.5 (Figure 1B) and reduced the levels of SCFAs in both the colonic

contents and portal vein plasma (Figure 1C and 1D). However, these short-term

dietary interventions (7 days) did not alter the levels of SCFAs in the peripheral

circulation (Figure S1B), suggesting it is the colon where fiber intake produces

marked effects. Aligned with previous reports,[5,11,40] these dietary interventions

significantly altered the composition of the gut microbiome in mice, as indicated by

both β-diversity (Figure S1C) and α-diversity (Figure S1D-G), with SCFA producers,

such as Lachnospiraceae-NK4A136-group, being enriched with the high fiber diet

(Figure S1H-I). Low fiber diet significantly enriched *Alistipes* spp. (Figure S1I),

which is associated with intestinal inflammation and hypertension.[41,42]

We then aimed to determine whether the pH-lowering effect of dietary fiber is

dependent on the gut microbiota. To achieve this, we pre-treated mice with an

antibiotic cocktail to reduce the bacterial load before starting the same dietary

interventions (Figure 1E). Indeed, the antibiotic cocktail effectively reduced the

bacterial load (Figure 1F). This reduction in bacteria in the intestine resulted in a

higher luminal pH in the cecum and colon, independent of the level of fiber in the diet

(Figure 1G-J). This is consistent with previously reported data showing lacking gut

microbiota leads to large intestinal alkalization.[32] Our data further support the theory

that acidic large intestinal pH produced when fiber is consumed is dependent on the

gut microbiota, particularly bacteria.

243 **Acidic pH regulates GPR65 signaling and inhibits pro-inflammatory pathways in**

244 **both colon and spleen derived CD8[+] T cells**

245 The colonic luminal pH produced by high fiber intake, pH 6.5, activates a family of

246 proton-sensing GPCRs, while the pH produced by low fiber intake, pH 7.5, silences

247 them.[14,15]{Howard, 2024 #6584} Within this family, we chose to focus on GPR65 due

248 to its previously known association with inflammatory diseases related to gut

249 homeostasis,[18-23] as well as its more recent association with hypertension in PheWAS

250 studies (Table S1).[22] GPR65 expression is particularly enriched in the immune

251 compartment.[27-29] There are no robust antibodies against GPR65, so we leveraged a

252 mouse model where GFP replaced one copy of the GPR65 gene to detect where

253 GPR65 was being expressed. Using $Gpr65^{gfp/+}$ mice,[22,43] we found that CD8[+] T cells

254 highly expressed GPR65 protein compared to CD4[+] T cells (Figure 2A). This is

255 consistent with human GPR65 expression in a public database.[28] Interestingly, CD8[+]

256 T cells causally contribute to the elevation of BP.[36] Moreover, CD8[+] T cells extracted

257 from colon lamina propria had a higher level of GPR65 expression compared to

258 splenic CD8[+] T cells, while CD4[+] T cells extracted from colon lamina propria and

259 spleen did not differ in GPR65 expression (Figure 2A). In addition, we found that

260 $Gpr65^{-/-}$ mice had significantly larger spleens (Figure 2B) with significantly more

261 leukocytes, including CD4[+], CD8[+] and γδ T cells (Figure 2C, Figure S2A and S2B),

262 but not Foxp3[+] regulatory T cells (T_{reg}) (Figure S2C).

263 GPR65 is a Gαs-coupled receptor whose activation induces intracellular cAMP

264 accumulation.[44] cAMP production in response to low pH was impaired in

macrophages of GPR65 deficient mice.[45] We discovered that low (GPR65 activating)

pH (6.5) increased the level of intracellular cAMP in splenic cells in a GPR65-

dependent manner (Figure 2D). The increased intracellular cAMP can lead to the

phosphorylation of cAMP response element-binding protein (CREB). Phosphorylated

CREB (pCREB) has various roles in immune responses, including suppressing the

production of pro-inflammatory cytokines, tumor necrosis factor alpha (TNFα) and

interferon gamma (IFNγ),[46] which underlie the pathogenesis of hypertension.[47,48] Thus,

we investigated whether pH and GPR65 signaling regulated the phosphorylation of

CREB in T cells extracted from the spleen and colon lamina propria *in vitro*.

Compared to pH 7.5, pH 6.5 significantly elevated pCREB levels in WT splenic CD8$^+$

and CD4$^+$ T cells activated by PMA and ionomycin (Figure 2E and Figure S2D). The

elevation of pCREB by acidic pH was impaired in *Gpr65$^{-/-}$* splenic CD8$^+$ T cells

(Figure 2E), but not splenic CD4$^+$ T cells (Figure S2D). While we did not observe a

significant difference in pCREB levels in WT colonic T cells activated by PMA and

ionomycin at pH 6.5 and pH 7.5, GPR65 deficiency significantly reduced the pCREB

levels of colonic CD4$^+$ and CD8$^+$ T cells under pH 6.5 but not pH 7.5 (Figure 2F and

Figure S2E). Downstream of the GPR65-pCREB axis, acidic pH 6.5 indeed

suppressed TNFα and IFNγ production from splenic and colonic CD8$^+$ T cells in a

GPR65-dependent manner (Figure 2G and S2F). However, the impact of acidic pH

and GPR65 signaling was less profound on TNFα and IFNγ production from splenic

and colonic CD4$^+$ T cells (Figure S2G and S2H).

In summary, pH change, which resembles the regulation of colonic luminal pH by

dietary fiber, regulates GPR65 signaling. Low pH, via GPR65 activation, suppresses

the pro-inflammatory cytokine production by $CD8^+$ T cells both locally from the

colon and systemically from the spleen (Figure 2H). We hypothesized this may reduce

BP and protect against CVD.

Deficiency of GPR65 does not impair gut homeostasis

We then examined the impact of GPR65 deficiency on gut homeostasis. *Gpr65*$^{-/-}$ and

WT mice exhibited similar gut microbiome compositions (Figure S3A and S3B), and

immune cell composition in the colon, including $CD4^+$, $CD8^+$, $\gamma\delta$ T cells, and T_{reg}

cells (Figure S3C and S3D). However, *Gpr65*$^{-/-}$ mice had altered gut physiology

(Figure S3E), demonstrated in fibrosis (Figure S3F), muscularis propria thickness

(Figure S3G) and goblet cell frequency (Figure S3H) compared to WT mice.

Deficiency of GPR65 results in spontaneous cardiovascular dysfunction

The role of GPR65 in the cardiovascular system and BP regulation remains largely

unknown. Therefore, we characterized multiple key cardiovascular phenotypes of

Gpr65$^{-/-}$ mice. Compared to WT mice, male *Gpr65*$^{-/-}$ mice had no difference in body

weight (Figure S4A), body composition (Figure S4B through S4D), food and water

intake (Figure S4E and S4F), or feces and urine excretion (Figure S4G and S4H). We

discovered, however, that *Gpr65*$^{-/-}$ mice spontaneously developed higher diastolic BP

compared to WT mice (Figure 3A). Consistently, *Gpr65*$^{-/-}$ mice had significantly

enlarged heart and kidney compared to WT mice (Figure 3B and 3C). This was

accompanied by significantly higher levels of cardiac fibrosis (Figure 3D and 3E), left

ventricular posterior wall in the end diastole (LVPW;d, Figure 3F), and cardiac output

(Figure 3G). Similarly, renal fibrosis was significantly elevated in *Gpr65$^{-/-}$* mice

(Figure 3H and 3I). We performed a saline challenge to investigate the influence of

GPR65 deficiency on renal function. Compared to WT mice, *Gpr65$^{-/-}$* mice excreted

less water and sodium over a 5-hour period (Figure 3J and 3K), which suggests

impaired diuretic and natriuretic functions. Aligned with the significant functional

phenotypes, urine creatinine levels were significantly increased in *Gpr65$^{-/-}$* mice

(Figure 3L). We then investigated the number of immune cells in the kidney (Figure

S4I and S4J). We observed significantly elevated CD45^{+} immune cells (Figure 3M),

particularly CD8^{+} T cells (Figure 3N). CD4^{+} T cells (Figure 3O), γδ T cells (Figure

3P), macrophages/monocytes (Figure 3Q), and dendritic cells (Figure S4K) were also

more abundant in the kidneys of *Gpr65$^{-/-}$* mice, while no difference in T$_{reg}$ cells

(Figure S4L) and neutrophils (Figure S4M) were observed.

Importantly, we observed similar cardiovascular phenotypes in female *Gpr65$^{-/-}$*

mice, including higher BP (Figure S5A), cardiorenal hypertrophy (Figure S5B and

S5C), thicker LVPW;d (Figure S5D), and elevated cardiorenal fibrosis (Figure S4E

through S4H) compared to female WT mice. Collectively, these results clearly

demonstrated that GPR65 deficiency leads to the spontaneous development of high

BP and associated end-organ damage which leads to CVD.

Deletion of GPR65 decreased the protection of dietary fiber against high blood

pressure

We next investigated whether pH-sensing by GPR65 conferred cardiovascular

protection by dietary fiber. To achieve this, we challenged WT and *Gpr65$^{-/-}$* mice with

Ang II (0.5mg/kg body/weight/day) and fed the mice either high or low fiber diets

(Figure 4A). As previously reported by us, high fiber diet suppressed the elevation of

systolic BP (Figure 4B) induced by Ang II infusion in WT mice.[5] Such suppression

was significantly smaller in *Gpr65*[-/-] mice (Figure 4B). GPR65 deficiency

significantly increased BP with hypertensive stimuli when mice were fed with high

fiber diet, while the impact of GPR65 genotype was negligible under low fiber diet

(Figure 4B). GPR65 deficient mice fed high fiber had higher cardiac weight (Figure

4C), thicker LVPW;d (Figure 4D), reduced ejection fraction (Figure 4E) and

fractional shortening (Figure 4F), and higher cardiac fibrosis relative to WT mice

(Figure 4G and 4H). Similarly, kidney weight (Figure 4I) and renal fibrosis (Figure

S6A and S6B) were increased. Renal natriuretic and diuretic function were also

impaired by the GPR65 deficiency (Figure 4J and 4K). We quantified the number of

immune cells in the kidneys and observed that high fiber fed *Gpr65*[-/-] mice had

significantly more CD45[+] immune cells than the high fiber fed WT mice (Figure S6C).

Notably, the infiltration of CD8[+] T cells, whose responses were regulated by acidic

pH and GPR65 signaling, was significantly higher in the kidney of *Gpr65*[-/-] mice

(Figure 4L). In addition, $\gamma\delta$ T cells and macrophages/monocytes were also increased

(Figure S6D and S6E), while CD4[+] T cells, T$_{reg}$ cells, neutrophils and dendritic cells

were not (Figure S6F through S6I). However, when mice were fed with a low fiber

diet, which mimics an environment where GPR65 is not activated, there was no

differences in cardiac or renal parameters besides fibrosis, which was further

exacerbated in *Gpr65*[-/-] mice (Figure 4B through 4K, Figure S6A and S6B).

Consistent with this, no difference was observed in immune cell infiltration in the

kidney under a low fiber diet (Figure 4L, Figure S6C through S6I).

We then examined the large intestinal tissue of mice fed low and high fiber diets

challenged with Ang II. High fiber intake thickened the muscularis propria layer and

increased the length of intestinal villi in the cecum, irrespective of *Gpr65* genotype

(Figure S7A though S7C) but did not influence the width of the fibrotic layer and the

number of goblet cells (Figure S7D and S7E). We also characterized the gut

microbiome composition by analyzing the 16S rRNA gene. The main factor driving

the differences in gut microbiota was diet rather than the genotype for *Gpr65* (Figure

S7F and S7G). The top leading taxonomic change explaining the differences in gut

microbiome was the enrichment of *Bacteroides* in the groups fed high fiber diet

(Figure S7H and S7I). *Bacteroides* species are the most predominant anaerobes in the

gut[49] with significant cardiovascular benefits.[50-53] Interestingly, *Bacteroides* spp. are

the dominant carbohydrate fermenters at pH 6.0 to 7.0, the normal luminal pH within

the cecum and colon.[54-56] The major products of carbohydrate fermentation by

Bacteroides spp. are propionate and acetate, rather than butyrate.[57,58] Indeed,

compared to low fiber, high fiber diet profoundly elevated acetate and propionate

levels in the cecal contents, independent of the genotype of the mice, while no

significant difference in cecal butyrate levels was observed (Figure S7J).

Lachnospiraceae-NK4A136-group, enriched following the 7-day high fiber

intervention (Figure S1D), was also more abundant in mice fed high fiber diet in this

study (Figure S7I). Low fiber diet expanded the genera *Bilophila* and *Alistipes* (Figure

S7H and S7I), which are associated with inflammation and higher risk of

CVD.[41,42,59,60]

GPR65 absence in CD8$^+$ T cells elevated blood pressure under high fiber diet

We then investigated whether immune regulation by acidic pH and GPR65 in CD8+ T

cells causally contributed to elevated BP under a high-fiber diet. *Rag1$^{-/-}$* mice lack

both T and B cells and have blunted BP elevation in the Ang II-induced hypertension

model.[61] We adoptively transferred splenic CD8$^+$ T cells from WT or *Gpr65$^{-/-}$* mice

fed a high fiber diet (Figure S8A) into *Rag1$^{-/-}$* mice which were also fed high fiber

diet, then infused with Ang II for 14 days (Figure 5A). We indeed observed that *Rag1$^{-/-}$*

$^{/-}$ mice that received *Gpr65$^{-/-}$* CD8$^+$ T cells developed higher BP than *Rag1$^{-/-}$* mice that

received WT CD8$^+$ T cells (Figure 5B). *Gpr65$^{-/-}$* CD8$^+$ T cells also led to cardiac

hypertrophy in Ang II-infused *Rag1$^{-/-}$* mice but had no significant impact on kidney

size (Figure 5C). Using flow cytometry, we confirmed the presence of adoptive

transferred CD8$^+$ T cells in the recipient mice's kidneys, blood, and colon lamina

propria (Figure S8A). We also observed a trend towards more abundant CD8$^+$ T cells

in the kidneys of *Rag1$^{-/-}$* mice that received *Gpr65$^{-/-}$* CD8$^+$ T cells (Figure 5D) but not

in blood and colon lamina propria (Figure S8B and S8C). Moreover, *Gpr65$^{-/-}$* CD8$^+$ T

cells resulted in significantly higher CD45$^+$ leukocyte infiltration, particularly

macrophage/monocytes (Figure 5E). We did not observe these changes in blood and

colon lamina propria (Figure S8D and S8E). Together, these results demonstrated that

GPR65 signaling in CD8$^+$ T cells causally regulates BP and improves the

accompanied cardiovascular complications under a high fiber diet.

Discussion

Dietary fiber and SCFAs are potent BP-lowering agents. Promoting dietary fiber or SCFAs intake represents a novel, safe, and affordable therapy against hypertension. SCFAs, particularly acetate, propionate, and butyrate, are well-known for their immunomodulatory effects by activating metabolite-sensing GPCRs, such as GPR41, GPR43 and GPR109A.[5,9,10] Some evidence suggests these receptors may confer cardiovascular protection.[5,9,10] Here, we identified a novel mechanism of how dietary fiber regulates BP and cardiovascular health via colonic luminal pH and GPR65. We show that a high-fiber diet lowers large intestinal pH by increasing the production of SCFAs by bacteria. This reduction in large intestinal pH activates the proton-sensing receptor GPR65, particularly in CD8$^+$ T cells, and its downstream anti-inflammatory signaling cascade, which lowered BP elevation induced by Ang II. Together, our findings support that large intestinal pH-sensing by GPR65, at least partly, explains the systemic cardiovascular protection by dietary fiber.

Colonic epithelial cells are naturally acidic. This largely results from the microbial production of acidic metabolites, particularly SCFAs.[11-13] Acidic pH in the colon is essential in maintaining gut homeostasis by suppressing the growth of pathogens and facilitating the expansion of beneficial microorganisms, including SCFA producers.[62] Indeed, we used dietary interventions to show that fiber determined the luminal pH in the colon and that it was inversely associated with the levels of SCFAs in both the large intestine and portal vein plasma. Such regulation was dependent on the presence of bacteria in the colon.

Among the host's pH sensors, GPR65 has critical roles in gastrointestinal

homeostasis.[18-24,63] GPR65 is particularly enriched in immune cells and lymphoid

tissues.[27-29] Evidence built in the last decade strongly supports the contribution of

inflammation in the development of hypertension,[47,48] making hypertension an

unconventional inflammatory disease. $CD8^+$ T cells are a well-known immune cell

subset contributing to the pathogenesis of hypertension.[36] We found GPR65 is more

highly expressed by $CD8^+$ T cells than $CD4^+$ T cells, and the expression was

particularly enriched in colon lamina propria. This expression profile suggests this

receptor may have a more significant role in $CD8^+$ T cells than $CD4^+$ T cells,

particularly in the colon. Furthermore, our study suggested GPR65 deficiency

increased immune cell counts in both the spleen and kidneys. Our data demonstrated

that low pH, similar to the large intestinal pH post-high-fiber intervention, suppresses

pro-inflammatory cytokine production by CD8+ T cells extracted from the spleen and

colon lamina propria via activating GPR65 signaling. This is a plausible mechanism

linking dietary fiber, low intestinal pH in the large intestine, GPR65 and

cardiovascular protection.

Characterization of its cardiovascular phenotype demonstrated that lack of

GPR65 led to the spontaneous development of mild but significant cardiovascular

disorders featured by impaired cardiorenal function, and cardiorenal remodeling.

These results demonstrate GPR65 has a protective role for cardiovascular health. This

phenotype was exacerbated when mice lacking GPR65 were challenged with Ang II,

particularly in a low-fibre diet. This shows a novel gene-by-environment interaction.

We propose that pH-sensing by GPR65 mediates cardiovascular protection through

dietary fiber via anti-inflammatory mechanisms. Our study demonstrates that the

cardiovascular protection conferred by dietary fiber is, at least in part, mediated by

GPR65. When challenged with Ang II, *Gpr65*[-/-] mice had higher BP and cardiorenal

hypertrophy compared to WT controls fed high fiber, and similar BP when fed a low

fiber diet, which would also silence GPR65 signaling in the large intestine. Indeed,

we observed that high fiber-fed *Gpr65*[-/-] mice had more CD8[+] T cells in the kidney

compared to high fiber fed WT mice. Analysis of the microbiome using 16S rRNA

sequencing demonstrated that diet, not *Gpr65* genotype, drove the difference in the

gut microbiota composition. High fiber diet promoted *Bacteroides* spp., a genus with

known cardiovascular benefits,[50-53] and increased the levels of acetate and propionate,

but not butyrate, in the intestine. Moreover, high fiber also inhibited certain taxa

associated with inflammation and cardiovascular disorders, including *Bilophila* and

Aliptipes.[41,42,59,60] These data suggested that GPR65 signaling does not regulate gut

microbiota composition directly and its effects on cardiovascular health are dependent

on downstream pathways triggered by GPR65, such as immunomodulation as

discussed above. Indeed, the adoptive transfer model suggested that GPR65

deficiency in CD8[+] T cells could causally explain the BP elevation under high fiber

diet.

The interventions with low and high fiber diets show robust differences in the

phenotype between WT and *Gpr65*[-/-] knockout mice, supporting the notion that

GPR65 signaling in hypertension starts in the colon. We acknowledge that since we

used a systemic knockout model, our animal studies cannot exclude the effect of pH-

sensing in other tissues. However, the *in vitro* experiments used the luminal colonic

pH we determined *in vivo*; however, we acknowledge these may not correspond to

colonic tissue pH.

In conclusion, this is the first study to report the role of colonic luminal and tissue

pH on the host's cardiovascular health. We provided new insights into how dietary

fiber and gut microbiota-derived metabolites regulate host health from the perspective

of pH. We established a role for GPR65 as a metabolite-sensing GPCR by sensing pH

changes caused by microbial fiber fermentation, showing a novel gene-by-

environment interaction. Since pH alterations happen universally, this mechanism

may extend to broader applications. Agents reducing colonic luminal pH or agonists

activating GPR65 may be potential tools for the global control of CVDs.

Conflict of Interest

None to declare.

Acknowledgements

This study was supported by a National Heart Foundation Vanguard Grant (102927)

and a National Health & Medical Research Council of Australia (NHMRC) Ideas

Grant (GTN2017382). L.X., M.J.A were supported by Monash Graduate Scholarships.

R.R.M was supported by a scholarship from the Faculty of Science, Monash

University. J.A.O. is supported by an NHMRC Fellowship (GTN1124288). F.Z.M is

supported by a Senior Medical Research Fellowship from the Sylvia and Charles

Viertel Charitable Foundation Fellowship, a National Heart Foundation Future Leader

Fellowship (105663) and an NHMRC Emerging Leader Fellowship (GNT2017382).

The Baker Heart & Diabetes Institute is supported in part by the Victorian

Government's Operational Infrastructure Support Program. The authors thank Monash

Animal Research Platform, Monash Genomic Modification Platform, Monash

Histology Platform, Monash FlowCore, Monash Biomedical Imaging, Monash

Proteomics and Metabolomics Facility, Monash Bioinformatics, and the Australian

Genome Research Facility for their technical supports. Summary figures were created

with Biorender. We thank Prof Peter Gibson, Jane Muir, Dakota Rhys-Jones, Daniel

So, and Chu Yao, for their feedback on the manuscript.

497   Author contributions

L.X. planned and performed most of the animal in vivo and in vitro experiments, data

analyses, provided intellectual inputs and wrote the manuscript. R.R.M., E.D., S.A.,

K.C.L., Z.M., K.M.C. contributed to animal in vivo experiments. J.A.O. contributed

to pCREB detection. M.P., A.P. performed gut histology analyses. H.J. contributed to

animal in vivo experiments and rodent cardiac ultrasound data analyses, heart and

kidney histology analyses. E.S. performed rodent cardiac ultrasound scanning and

supervised the data analyses performed by L.X.. D.A. performed SCFA measurement

by LC-MS. C.A., M.J.A., Y.A.Y. contributed to animal in vivo experiments, mouse

strain maintenance and cAMP assay. D.C. supervised SCFA measurement by LC-MS.

R.R. contributed to animal in vivo experiments, mouse strain maintenance and

provided intellectual inputs. C.R.M. and F.Z.M. conceived the study and supervised

the research. F.Z.M. secured the funding to support this study. All authors approved of

and contributed to the final version of the manuscript.

References

1.   Murray, C. J. L. *et al.* Global burden of 87 risk factors in 204 countries and

territories, 1990-2019: a systematic analysis for the Global Burden of Disease

Study 2019. *The Lancet* **396**, 1223-1249 (2020).

2.   Afshin, A. *et al.* Health effects of dietary risks in 195 countries, 1990–2017: a

systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*

393, 1958-1972 (2019).

3.   Reynolds, A. *et al.* Carbohydrate quality and human health: a series of

systematic reviews and meta-analyses. *Lancet* **393**, 434-445 (2019).

4.   Marques, F. Z. *et al.* High-Fiber Diet and Acetate Supplementation Change the

Gut Microbiota and Prevent the Development of Hypertension and Heart

Failure in Hypertensive Mice. *Circulation* **135**, 964-977 (2017).

5.   Kaye, D. M. *et al.* Deficiency of Prebiotic Fiber and Insufficient Signaling

Through Gut Metabolite-Sensing Receptors Leads to Cardiovascular Disease.

Circulation **141**, 1393-1403 (2020).

6.   O'Donnell, J. A., Zheng, T., Meric, G. & Marques, F. Z. The gut microbiome

and hypertension. *Nat Rev Nephrol* **19**, 153-167 (2023).

7.   Jama, H. A. *et al.* Prebiotic intervention with HAMSAB in untreated essential

hypertensive patients assessed in a phase II randomized trial. *Nature*

Cardiovascular Research **2**, 35-43 (2023).

8.   Jama, H. A. *et al.* Prebiotic intervention with HAMSAB in untreated essential

hypertensive patients assessed in a phase II randomized trial. *Nature*

Cardiovascular Research (2023).

9.   Natarajan, N. *et al.* Microbial short chain fatty acid metabolites lower blood

pressure via endothelial G-protein coupled receptor 41. *Physiol Genomics* **48**,

826-834 (2016).

10.   Muralitharan, R. R. *et al.* GPR41/43 regulates blood pressure by improving

gut epithelial barrier integrity to prevent TLR4 activation and renal

inflammation. *bioRxiv*, 2023.2003.2020.533376 (2023).

11.   Campbell, J. M., Fahey, G. C., Jr. & Wolf, B. W. Selected indigestible

oligosaccharides affect large bowel mass, cecal and fecal short-chain fatty

acids, pH and microflora in rats. *J Nutr* **127**, 130-136 (1997).

12.   Cummings, J. H., Pomare, E. W., Branch, W. J., Naylor, C. P. & Macfarlane, G.

T. Short chain fatty acids in human large intestine, portal, hepatic and venous

blood. *Gut* **28**, 1221 (1987).

13.   den Besten, G. *et al.* The role of short-chain fatty acids in the interplay

between diet, gut microbiota, and host energy metabolism. *Journal of Lipid*

Research **54**, 2325-2340 (2013).

14.   Ludwig, M. G. *et al.* Proton-sensing G-protein-coupled receptors. *Nature* **425**,

93-98 (2003).

15.   Wang, J. Q. *et al.* TDAG8 is a proton-sensing and psychosine-sensitive G-

protein-coupled receptor. *J Biol Chem* **279**, 45626-45633 (2004).

16.   Xie, L., Alam, M. J., Marques, F. Z. & Mackay, C. R. A major mechanism for

immunomodulation: Dietary fibres and acid metabolites. *Seminars in*

Immunology **66**, 101737 (2023).

17.   Hajjar, S. & Zhou, X. pH sensing at the intersection of tissue homeostasis and

inflammation. *Trends in Immunology* **44**, 807-825 (2023).

18.   Franke, A. *et al.* Genome-wide meta-analysis increases to 71 the number of

confirmed Crohn's disease susceptibility loci. *Nat Genet* **42**, 1118-1125 (2010).

19.   Ke, X. Presence of multiple independent effects in risk loci of common

complex human diseases. *Am J Hum Genet* **91**, 185-192 (2012).

20.   Ballester, V. *et al.* Association of NOD2 and IL23R with inflammatory bowel

disease in Puerto Rico. *PLoS One* **9**, e108204 (2014).

21.   Liu, J. Z. *et al.* Association analyses identify 38 susceptibility loci for

inflammatory bowel disease and highlight shared genetic risk across

populations. *Nature Genetics* **47**, 979-986 (2015).

22.   Xie, L. *et al.* pH and Proton Sensor GPR65 Determine Susceptibility to Atopic

Dermatitis. *The Journal of Immunology*, ji2001363 (2021).

23.   Lassen, K. G. *et al.* Genetic Coding Variant in GPR65 Alters Lysosomal pH

and Links Lysosomal Dysfunction with Colitis Risk. *Immunity* **44**, 1392-1405

(2016).

24.   Tcymbarevich, I. *et al.* Lack of the pH-sensing Receptor TDAG8 [GPR65] in

Macrophages Plays a Detrimental Role in Murine Models of Inflammatory

Bowel Disease. *J Crohns Colitis* (2018).

25.   Chen, X. *et al.* pH sensing controls tissue inflammation by modulating cellular

metabolism and endo-lysosomal function of immune cells. *Nature*

Immunology (2022).

26.   Watanabe, K. *et al.* A global overview of pleiotropy and genetic architecture in

complex traits. *Nat Genet* **51**, 1339-1348 (2019).

27.   Heng, T. S. P. *et al.* The Immunological Genome Project: networks of gene

expression in immune cells. *Nature Immunology* **9**, 1091-1094 (2008).

28.   Uhlen, M. *et al.* Proteomics. Tissue-based map of the human proteome.

Science **347**, 1260419 (2015).

29.   Thul Peter, J. *et al.* A subcellular map of the human proteome. *Science* **356**,

eaal3321 (2017).

30.   Radu, C. G., Nijagal, A., McLaughlin, J., Wang, L. & Witte, O. N. Differential

proton sensitivity of related G protein-coupled receptors T cell death-

associated gene 8 and G2A expressed in immune cells. *Proc Natl Acad Sci U S*

A **102**, 1632-1637 (2005).

31.   McConnell, E. L., Basit, A. W. & Murdan, S. Measurements of rat and mouse

gastrointestinal pH, fluid and lymphoid tissue, and implications for in-vivo

experiments. *J Pharm Pharmacol* **60**, 63-70 (2008).

32.   Shimizu, K., Seiki, I., Goto, Y. & Murata, T. Measurement of the Intestinal pH

in Mice under Various Conditions Reveals Alkalization Induced by Antibiotics.

Antibiotics **10** (2021).

33.   Avery, E. G. *et al.* Intestinal interstitial fluid isolation provides novel insight

into the human host-microbiome interface. *Cardiovasc Res* (2025).

34.   Brown, R. D. *et al.* Influence of the adenosine A1 receptor on blood pressure

regulation and renin release. *American Journal of Physiology-Regulatory,*

Integrative and Comparative Physiology **290**, R1324-R1329 (2006).

35. Kamat, N. V. *et al.* Renal transporter activation during angiotensin-II

hypertension is blunted in interferon-gamma-/- and interleukin-17A-/- mice.

Hypertension **65**, 569-576 (2015).

36. Trott, D. W. *et al.* Oligoclonal CD8+ T cells play a critical role in the

development of hypertension. *Hypertension* **64**, 1108-1115 (2014).

37. Jama, H. *et al.* Maternal diet and gut microbiota influence predisposition to

cardiovascular disease in the offspring. *bioRxiv*, 2022.2003.2012.480450

(2022).

38. Caporaso, J. G. *et al.* Global patterns of 16S rRNA diversity at a depth of

millions of sequences per sample. *Proceedings of the National Academy of*

Sciences **108**, 4516-4522 (2011).

39. Nakai, M. *et al.* Essential Hypertension Is Associated With Changes in Gut

Microbial Metabolic Pathways: A Multisite Analysis of Ambulatory Blood

Pressure. *Hypertension* **78**, 804-815 (2021).

40. Marques, F. Z. *et al.* High-Fiber Diet and Acetate Supplementation Change the

Gut Microbiota and Prevent the Development of Hypertension and Heart

Failure in Hypertensive Mice. *Circulation* **135**, 964-977 (2017).

41. David, L. A. *et al.* Diet rapidly and reproducibly alters the human gut

microbiome. *Nature* **505**, 559 (2013).

42. Kim, S. *et al.* Imbalance of gut microbiome and intestinal epithelial barrier

dysfunction in patients with high blood pressure. *Clinical Science* **132**, 701-

718 (2018).

43. Wirasinha, R. C. *et al.* GPR65 inhibits experimental autoimmune

encephalomyelitis through CD4(+) T cell independent mechanisms that

include effects on iNKT cells. *Immunol Cell Biol* **96**, 128-136 (2018).

44. Robert, R. & Mackay, C. R. Gαs-coupled GPCRs GPR65 and GPR174.

Downers for immune responses. *Immunol Cell Biol* **96**, 341-343 (2018).

45. Mogi, C. *et al.* Involvement of proton-sensing TDAG8 in extracellular

acidification-induced inhibition of proinflammatory cytokine production in

peritoneal macrophages. *J Immunol* **182**, 3243-3251 (2009).

46. Wen, A. Y., Sakamoto, K. M. & Miller, L. S. The role of the transcription

factor CREB in immune function. *J Immunol* **185**, 6413-6419 (2010).

47. Drummond, G. R., Vinh, A., Guzik, T. J. & Sobey, C. G. Immune mechanisms

of hypertension. *Nature Reviews Immunology* (2019).

48. Norlander, A. E., Madhur, M. S. & Harrison, D. G. The immunology of

hypertension. *The Journal of Experimental Medicine* **215**, 21 (2018).

49. Salyers, A. A. Bacteroides of the human lower intestinal tract. *Annual Review*

of Microbiology **38**, 293-313 (1984).

50. Yoshida, N. *et al.* Bacteroides vulgatus and Bacteroides dorei Reduce Gut

Microbial Lipopolysaccharide Production and Inhibit Atherosclerosis.

Circulation **138**, 2486-2498 (2018).

51. Emoto, T. *et al.* Analysis of Gut Microbiota in Coronary Artery Disease

Patients: a Possible Link between Gut Microbiota and Coronary Artery

Disease. *Journal of Atherosclerosis and Thrombosis* **advpub** (2016).

52. Emoto, T. *et al.* Characterization of gut microbiota profiles in coronary artery

disease patients using data mining analysis of terminal restriction fragment

length polymorphism: gut microbiota could be a diagnostic marker of

coronary artery disease. *Heart and Vessels* **32**, 39-46 (2017).

53. Karlsson, F. H. *et al.* Symptomatic atherosclerosis is associated with an altered

gut metagenome. *Nature Communications* **3**, 1245 (2012).

54. Walker Alan, W., Duncan Sylvia, H., McWilliam Leitch, E. C., Child Matthew,

W. & Flint Harry, J. pH and Peptide Supply Can Radically Alter Bacterial

Populations and Short-Chain Fatty Acid Ratios within Microbial Communities

from the Human Colon. *Applied and Environmental Microbiology* **71**, 3692-

3700 (2005).

55. Ilhan, Z. E. *et al.* pH-Mediated Microbial and Metabolic Interactions in Fecal

Enrichment Cultures. *mSphere* **2**, e00047-00017 (2017).

56. Raba, G., Adamberg, S. & Adamberg, K. Acidic pH enhances butyrate

production from pectin by faecal microbiota. *FEMS Microbiology Letters* **368**,

fnab042 (2021).

57. Salonen, A. *et al.* Impact of diet and individual variation on intestinal

microbiota composition and fermentation products in obese men. *The ISME*

Journal **8**, 2218-2230 (2014).

58. Aguirre, M. *et al.* Diet drives quick changes in the metabolic activity and

composition of human gut microbiota in a validated in vitro gut model.

Research in Microbiology **167**, 114-125 (2016).

59.   Qin, J. *et al.* A metagenome-wide association study of gut microbiota in type 2

diabetes. *Nature* **490**, 55-60 (2012).

60.   Devkota, S. *et al.* Dietary-fat-induced taurocholic acid promotes pathobiont

expansion and colitis in Il10–/– mice. *Nature* **487**, 104-108 (2012).

61.   Guzik, T. J. *et al.* Role of the T cell in the genesis of angiotensin II induced

hypertension and vascular dysfunction. *J Exp Med* **204**, 2449-2460 (2007).

62.   Duncan, S. H., Louis, P., Thomson, J. M. & Flint, H. J. The role of pH in

determining the species composition of the human colonic microbiota.

Environ Microbiol **11**, 2112-2122 (2009).

63.   Diogo, D. *et al.* Phenome-wide association studies across large population

cohorts support drug target validation. *Nature Communications* **9**, 4285 (2018).

Figure. 1. Dietary fiber lowers colonic pH through the gut microbiota.

A, The experiment design to characterize the impact of dietary fiber on luminal pH in

cecum and colon. **B**, Colonic luminal and interstitial fluid pH at the endpoint, post

dietary interventions. **C**, Levels of acetic, propionic, butyric acids and total short-

chain fatty acids (SCFAs) in cecal contents and **D**, in plasma from the portal vein. **E**,

The experiment design to characterize the impact of dietary fiber on luminal pH in the

cecum and colon with antibiotic treatment. Time points with asterisks indicate when

fecal samples were collected for bacterial load measurements. **F**, Bacterial load. **G**,

Cecal and colonic luminal pH at the endpoint, post dietary interventions with

antibiotics treatments. **H**, Each data point represents an individual mouse, and n=6

mice in each group for all panels besides colonic interstitial fluid pH (n=8 for control

diet, n=9 low and high fiber diets); all data represented as means ± SEM; **C**, **D**, **G**,

One-way ANOVA with adjustment for multiple comparisons; **H**, Student's t test;

*P<0.05, **P<0.01, ***P<0.001, ****P<0.0001.

Figure 2. pH and GPR65 signaling modulate immune responses related to

hypertension pathogenesis.

Single cell suspensions were prepared from the spleens or colon lamina propria of

male *Gpr65*^{gfp/+} mice at the age of 12 weeks. **A**, GPR65 expression in CD4⁺ and

CD8⁺ T cells from spleen and colon lamina propria indicated as mean fluorescence

intensity (MFI) of GFP. Single cell suspensions were prepared from the colon lamina

propria of male wild-type C57BL/6J (WT) mice or *Gpr65*$^{-/-}$ mice at the age of 12

weeks. **B**, CD4$^+$ and CD8$^+$ T cell count in the colon lamina propria. Colon lamina

propria cells were incubated with PMA and ionomycin for 2 hours. **C**, pCREB

(Ser133) levels of CD8$^+$ T cells extracted from colon lamina propria. Colon lamina

propria cells were incubated with PMA and ionomycin for 4 hours. **D**, Percentage of

TNFα producing CD8$^+$ T cells at pH 6.5 and 7.5 among WT and *Gpr65*$^{-/-}$ CD8$^+$ T

cells derived from colon lamina propria. **E**, Percentage of IFNγ producing CD8$^+$ T

cells at pH 6.5 and 7.5 among WT and *Gpr65*$^{-/-}$ CD8$^+$ T cells derived from colon

lamina propria. **B**, Spleen weight to tibia length index from male mice at the age of 12

weeks. **C**, CD45$^+$ immune cell count, CD4$^+$ and CD8$^+$ T cell count in the spleen.

Splenocytes were isolated from mice at the age of 8-10 weeks and incubated under

either pH 6.5 or pH 7.5 for 30 min. **D**, Levels of cAMP in the cell lysate extracts. **E**,

pCREB (Ser133) levels of splenic CD8$^+$ T cells in splenocytes incubated with PMA

and ionomycin for 2 hours. **F**, pCREB (Ser133) levels of CD8$^+$ T cells extracted from

colon lamina propria incubated with PMA and ionomycin for 2 hours. **G**, Percentage

of TNFα and IFNγ producing CD8$^+$ T cells at pH 6.5 and 7.5 among WT and *Gpr65*$^{-/-}$

splenocytes were incubated with PMA and ionomycin for 4 hours. **H**, Graphic

summary of how pH regulates inflammatory responses contributing to hypertension in

CD8$^+$ T cells. Created with BioRender.com. Each data point represents an individual

mouse, and n=4-26 mice in each group; All data represented as means ± SEM. **A, D-**

G, Two-way ANOVA with adjustment for multiple comparisons. **B**, **C**, Student's t test.

*P<0.05, **P<0.01, ***P<0.001, ***P<0.0001.

Figure 3. GPR65 deficiency spontaneously increases blood pressure, tissue

hypertrophy, and renal inflammation in male mice.

A, Systolic blood pressure (BP) of male mice measured non-invasively by radio-

telemetry at the age of 10 weeks. Tissues of male mice were collected and weighed at

the age of 12 weeks. **B**, Heart weight to tibia length index. **C**, Kidney weight to tibia

length index. **D**, Representative heart sections stained with Masson's trichrome; For

upper panels, magnification = ×10, scale bar = 1mm; For lower left panels

(interstitial), magnification = ×200, scale bar = 50μm; For lower right panels

(perivascular), magnification = ×100, scale bar = 100μm. **E**, Percentage of fibrotic

area in the cardiac tissues. **F**, Left ventricle posterior wall in the end diastole and **G**,

cardiac outputs from ultrasound screening was performed at 11 weeks of age. **H**,

Representative kidney sections stained with Masson's trichrome; For left panels,

magnification= ×5, scale bar = 2mm; For right panels, magnification = ×200, scale

bar = 50μm. **I**, Percentage of fibrotic area in the renal fibrosis. Saline challenge was

performed at the age of 11 weeks. **J**, Renal diuretic functions measured by

accumulated water excretion over 5 hours post injection. **K**, Renal natriuretic

functions measured accumulated sodium excretion over 5 hours post injection. **L**, The

level of creatinine in urine from urine were collected from 24h metabolic caging. **M**,

$CD45^+$ immune cell count, **N**, $CD8^+$ T cell count, **O**, $CD4^+$ T cell count, **P**, γδ T cell

count and **Q** Macrophage/Monocyte count from kidneys of male mice at 12 weeks of

age. For the left figures of panels **J** and **K**, n=10-14 mice in each group; All data

represented as means ± SEM; Two-way ANOVA. For other panels, each data point

represents an individual mouse, and n=6-26 mice in each group; All data represented

as means ± SEM; **B, C, F**, Mann-Whitney test; All other panels, Student's t test.

*$P<0.05$, **$P<0.01$, ***$P<0.001$, ****$P<0.0001$.

Figure 4. Deficiency of GPR65 decreased the cardiorenal protective effects of

dietary fiber.

6-week-old male WT and *Gpr65*^{-/-} were infused with 0.5 mg/kg body weight/day

angiotensin-II for 28 days and fed with either high fiber (HF) or low fiber (LF) diet.

A, The experiment design. **B**, Systolic BP profile over 4 weeks of infusion; **C**, Heart

weight to tibia length index; Cardiac ultrasound scanning was performed at week 4 of

infusion. **D**, Left ventricle posterior wall in the diastole; **E**, Cardiac ejection fraction;

F, Cardiac fractional shortening; **G**, Percentage of fibrotic area in the cardiac tissues

stained with Masson's trichrome. **H**, Representative heart sections; For upper panels,

magnification = ×10, scale bar = 1mm; For lower left panels (interstitial),

magnification = ×200, scale bar = 50μm; For lower right panels (perivascular),

magnification = ×100, scale bar = 100μm. **I**, Kidney weight to tibia length index.

Mice were intraperitoneally injected with 10% of their body weight of 37 °C 0.9%

saline solution at the 3rd week post minipump implantation. **J**, Renal diuretic and **K,**

natriuretic functions measured by accumulated water excretion over 5 hours post

injection. **L**, CD8^{+} T cell count from kidneys of the mice at the endpoint of the study.

Each data point represents an individual mouse, and n=8-11 mice in each group; All

data represented as means ± SEM. Two-way ANOVA with adjustment for multiple

comparisons. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

Figure 5. Deficiency of GPR65 in CD8+ T cells causes BP elevation in mice under the high fiber diet.

10-week-old male *Rag1-/-* mice were adoptively transferred with either WT and *Gpr65-/-* CD8+ T cells, and then infused with 1.44 mg/kg body

weight/day angiotensin-II for 14 days. They were fed with HF diet. **A**, The experimental design; **B**, Systolic BP profile over 2 weeks of infusion;

C, Heart weight to tibia length and kidney weight to tibia length indexes. **D**, CD8+ T cell count, **E**, CD45+ immune cell count and

Macrophage/Monocyte count from kidneys of the mice at the endpoint of the study. Each data point represents an individual mouse, and n=5-7

mice in each group; All data represented as means ± SEM. **B**, Two-way ANOVA with adjustment for multiple comparisons. All other panels,

Student's t test. *P<0.05, ***P<0.001, ****P<0.0001.