Spatial localization ability of planarians identified through a light maze paradigm

Renzhi Qian[1], Yuan yan[1], Yu Pei[1], Yixuan Zhang[1], Yuanwei Chi[1], Yuxuan Chen[1], Kun Hao[1],

Zhen Xu[1], Guang Yang[1], Zilun Shao[1], Yuhao Wang[1], Xinran Li[1], Chenxu Lu[1], Xuan Zhang[1],

Kehan Chen[2], Wenqiang Zhang[2], Baoqing Wang[1], Zhengxin Ying[1] and Kaiyuan Huang[1]

[1]College of Biological Science, China Agricultural University; Beijing, China.

[2]College of Engineering, China Agricultural University; Beijing, China.

* Kaiyuan Huang

Email: cfhuangkaiyuan@gmail.com

Author Contributions: Renzhi Qian, Yuan Yan and Yu Pei contributed equally to this work.

Kaiyuan Huang, Renzhi Qian and Yuan yan designed research. Renzhi Qian, Yuan yan,

Yixuan Zhang, Yuanwei Chi , Yuxuan Chen, Kun Hao, Zhen Xu performed research. Guang

Yang, Zilun Shao, Yuhao Wang, Xinran Li, Chenxu Lu, Xuan Zhang and Yu Pei analyzed

data. Kehan Chen draw blueprint of the maze. Wenqiang Zhang, Baoqing Wang and

Zhengxin Ying gave suggestions to the research. Kaiyuan Huang and Yu Pei wrote the

manuscript. Wenqiang Zhang, Baoqing Wang and Zhengxin Ying revised the manuscript.

Abstract

Spatial localization ability is crucial for free-living animals to fit the environment. As shown by previous studies, planarians can be conditioned to discriminate directions. However, due to their simplicity and primitiveness, they had never been considered to have true spatial localization ability to retrieve locations of objects and places in the environment. Here, we introduce a light maze training paradigm to demonstrate that a planarian worm can navigate to a former recognized place from the start point, even if the worm is transferred into a newly produced maze. This finding identifies the spatial localization ability of planarians for the first time, which provides clues for the evolution of spatial learning. Since the planarians have a primitive brain with simple structures, this paradigm can also provide a simplified model for a detailed investigation of spatial learning.

Introduction

Spatial localization, the ability to learn and remember locations and routes, is crucial to survival for most animals. The ability can be used to forage for food, return to sites of storage or safety, and avoid sites of danger, thus helping animals survive and fit the environment. Thus this ability is considered to have evolved in almost all species.[1]

Planarians are free-living flatworms that usually live and stick under rocks, debris and water plants in streams, ponds, and springs[2]. They are considered one of the first class of animals to have a centralized brain structure, and they had evolved various sensory abilities, including

sensitivity to light[3], temperature[4], water currents[5], chemical gradients[6], vibration[7],

magnetic fields[8] and electric fields[9]. For freely living in the environment, it is reasonable to

hypothesize that planarians have evolved spatial localization ability. The study of planarian

behavior mainly focused on simple classical conditioning and proved that planarians can learn

simple associative tasks. Moreover, planarians are concluded to have abilities to discriminate

directions in a Y or T maze[10]. With such ability, they can turn their moving directions in

response to a certain stimulus. However, due to the primitiveness and simplicity of planarians,

they are not considered to have the ability of true, complex spatial localization ability[11],

which is used to localize a certain place in a familiar environment.

To prove the hypothesis that planarians have evolved spatial localization ability, we

designed a light maze paradigm inspired by the Morris water maze paradigm[12]. The Morris

water maze, designed by Morris in 1981, is widely used to evaluate the rats' spatial

localization ability through distal localization (when the goal object is invisible, inaudible and

cannot be detected by smell) rather than proximal localization (when the goal object is visible,

audible or detectable by smell)[12]. The Morris water maze uses water as the aversive stimuli

to prompt the rats to swim to a platform. Since planarians have innate behavior to avoid

light[3], similar to the Morris water maze, we chose light as the aversive stimuli to prompt

planarians to move to the dark chamber in the maze. The planarians' primitive eyes can only

sense the light rather than imaging; thus, they cannot visualize the dark chamber. To avoid

any chemical cues that a worm might leave in the dark chamber, in the test phase, worms

were transferred into newly manufactured mazes from their original mazes. Thus, the

information that a worm use to navigate to the dark chamber can only be distal. In the light

maze, the distal cues provided to the worm could be such as the topography of the maze, the

light intensity gradient, or the geomagnetic field which the goal object did not provide direct

information to the worm. The proximal cues could be such as chemicals from the goal object

which can be directly smelled by the worms. In the text, we use the word "escape" to show

that the worms solved the maze and successfully entered the dark chamber. We also use the

escape latency to measure the worms' performance, which is most commonly used in the

Morris water maze, defined as the time for the animal to find the dark chamber and escape

the maze. Our light maze paradigm is also inspired by the Barnes maze, which uses open

spaces as aversion stimuli to prompt rodents to seek shelter.

Here we demonstrated that planarians were able to use distal information to localize a

dark chamber in the light maze. To evaluate the learning effect of the worms, we measured

the escape latencies and drew typical routes of worms which showed learning effects.

These are the first findings suggesting that planarian worms have spatial localization abilities,

indicating that planarians might have higher cognition ability that had never been considered.

As a primitive forerunner of more evolved animals, the study of planarian cognition can be

evolutionarily instructive to the study of more evolved animals. Also, since the planarians'

nervous system has simple structures, it can provide a model for spatial learning studies.

77   Materials and Methods

78   Planarians

A laboratory strain of *D. japonica*, originating from wild collected *D. japonica* (identified by

cytochrome c oxidase subunit 1 gene) from the Cherry-Valley in Beijing Botanical Garden,

Haidian district, Beijing, China in 2019. Worms were maintained in Montjuic Water(Simulated

components of fresh water from Montjuic) in the dark and fed with chicken liver twice a week

on Monday night and Thursday night. The procedure in planarian rearing used protocol from

M. Shane Merryman et al. 2018 for reference [13]. The length of planarians used in the

experiment varied from 1 cm to 1.5 cm.

Experimental Setup

To investigate the spatial localization ability of the planarians, we designed a light maze. It

is a square maze with a dark chamber near one corner (Fig1.A). The maze consists of a

square pool with a side length of 65 mm and a depth of 6mm, the corners of the maze are

rounded to make it easier for planarians to crawl. A square dark room with inner side length of

14 mm and outer side length of 24 mm is located in one of the quadrants as the destination of

the maze. The starting position is located in the corner of the opposite quadrant of the dark

chamber. The dark chamber has only one entrance. 15 mL Montjuic water is added to the

maze before starting a trial.

The mazes were 3D-printed. The material used for 3D printing is photosensitive resin and

PLA (Polylactic acid), the main body of the mazes was printed in white using photosensitive

resin, and the dark chamber lid was printed in black using PLA (The dark chamber lid can be

detached from the main body of maze). In the SLA printing method, a photocurable and

photosensitive polymeric material is placed on a surface apparatus and is subsequently

opened to a UV light to cure the resin to form a initial layer. When initial layers cure,

the repetition of this process fabricates the 3D printing product[9]. The toxicity released by

the stereolithography (STL) printing materials (photosensitive resin in this experiment) is fatal

to worms, which causes the worms to die and disintegrate in hours, even in the mazes

handled by ultraviolet to have reduced toxicity. Although the fused deposition modeling (FDM)

materials like PLA is non-toxic, it is of lower printing precision, and its water leaking problem

may cause worms to escape from the maze while training. We found that a biocompatible

encapsulation material called Parylene to totally block the toxicity from the STL printed parts,

using which can make worms safely live in the maze without health problems for over 2

weeks.   The inner side of the mazes was coated with Parylene film (thickness of 12 μm) to

block the toxicity from the resin[14]. The structure and parameters of the maze are shown in

(Fig1.C).

During training, the mazes were put in a line on a black acrylic board to avoid light

reflections, and each maze was filled with 15 mL of Montjuic Water. A LED lighting tube was

set above the entrance of the dark chamber (Fig1.B) to make the entrance lighted and avoid

direct guidance to the entrance by the light. The length of the LED tube was 1.2 m, and the

power of the LED tube was 13 W. The distance between the bottom of the LED lighting tube

to the acrylic board was 125 mm. The LED lighting tube was set right above the dark

chamber. Light intensity distribution of the maze is shown in Fig1.D. Worms were carefully

manipulated through a smooth woolen brush.

Training Procedure

We totally used 36 worms (2 worms were excluded due to self-fission in the training

process) and they were divided into 6 groups. We used 6 mazes to train each group of worms

in the first 3 trials. In trial 4, all mazes used to train any of the worms were changed to newly

manufactured (36 new mazes). The training procedure started at 8:00 p.m. and ended at 9:00

p.m. each day for a total of 6 trials for each worm in 6 days. 1 day before the training, worms

were taken out from the home well to a petri dish. Diagram of the training procedure is shown

in Fig1.E. To control worms' satiety, they were fed at 10:00 a.m. on day 1 and day 4 in a 12-

well plate, each worm in a single well. Before training, worms were taken to a 12-well plate,

then the water in the mazes is changed and the mazes were set up as described in the last

section. The worm is put 5 mm away from wall a and wall b (the starting point might be a little

different in each trial because the worm need to sink to the bottom). The room temperature

during training was controlled at 20 ± 1 ℃. After the training, the worms were not taken out,

and the maze along with the acrylic board was put in an incubator to control the environment

temperature.

Fig 1. Experimental setup and training procedure. A. The conceptual drawing of the light

maze. B. The experimental setup of the light maze paradigm. C. The blueprint of the light

maze. The upper right shows the lateral view of section A. Length unit: mm. D. Light intensity

distribution of the maze. The light intensity distribution is measured according to vertical

distance from Wall a. Light intensity variance along Wall a is less than 5 Lux, which can be

neglected. E. Diagram of the training procedure.

Statistical Analysis

36 worms were involved in training and 2 of them were excluded due to self-fission.

However, possibly because the worms were gradually getting habituated to the light and the

environment, about one third of the worms in each trial quickly got to rest and stop moving in

each trial except trial 1. Hence, our data excluded worms that rest for more than 20 mins in 40

mins in each trial. So, from trial 1 to trial 6, n=34, 24, 24, 24, 23, 22. The latency of other

worms which did not enter the dark chamber (either rested or kept on moving) were counted

as 40 mins. Also, worms that went into the dark chamber but got out within 30 mins did not

count as a successful escape.

For statistical analysis, because the escape latencies in trial 1 and trial 4 were not normally

distributed (Kolmogorov–Smirnov test), we used the nonparametric Mann–Whitney U-test to

evaluate statistical significance of latencies in trial 1 and trial 4. One way ANOVA was applied

to determine whether there is a statistical significance between escape latencies of trial 2, 3,

4, 5 and 6. One way ANOVA and subsequent Turkey's multiple comparison test was applied

to determine whether worm speed in trial one is significantly higher than other trials. All data

were analyzed using PRISM (GraphPad Prism 9.0.0(121)). Routes are drawn by hand with

Photoleap (2.13).

Ethics statement

164   The overall study and animal experiments of this manuscript conformed to the guidelines

165   and regulatory standards of the Institutional Animal Care and Use Committee of China

166   Agricultural University and approved by the Institutional Animal Care and Use Committee of

167   China Agricultural University.

Results

We tested for recall of the location of the dark chamber of the worms in trial 4 when the

mazes were changed to newly manufactured. In trial 4, because the mazes were new, worms

cannot use local cues such as chemicals for navigation. The latency result shows that worms

in trial 4 displayed a significantly shorter time to escape compared with trial 1 (two-tailed U-

test, $P<0.01$; Fig.2A). To test whether this behavior is stable after changing the maze, we

continued trial 5 and trial 6. Comparing escape latencies in trial 4 to trial 2, 3, 5 and 6, there is

no significant difference, which means that changing to newly manufactured maze did not

affect worm's performance in the maze, showing that the worms might not use chemical cues

in this case (One way ANOVA, $P>0.1$; Fig.2A). We also counted the worms that underwent

successful escapes. From trial 3 to trial 6, there were 10-12 worms that successfully escaped

the maze (Fig2.C). However, from trial 2 to trial 6, a worm might successfully escape the

maze in some trials but not in other trials. From trial 2 to trial 6, the number of worms that

successfully escaped 0, 1, 2, 3, 4 and 5 times is 3, 2, 17, 8, 3 and 3, and the number of

worms that consecutively could not escape the maze for 2, 3, 4, and 5 times is 12, 8 ,1 and 3.

To exclude the effect of worm speed on latency, we counted the worm speed of a total

moving distance of 241 mm (the perimeter of the maze) for worms which moved for more

than 241 mm, otherwise, we used its actual moving distance to count its speed. The result

shows that worm speed in trial 1 are significantly higher than other trials (One way ANOVA,

P<0.001; Fig.2B), which means the worm speed did not shorten the latencies from trial 2 to

trial 6.

To further understand the worms' learning effect, we showed representative routes of trial 1

and trial 4 of 2 worms which successfully escaped the maze in trial 4 (Fig2.C-F). In trial 4, the

mazes are newly manufactured, and no chemicals or mucus can affect the navigation of the

worms. As the worms did not know the location of the dark chamber prior to the training, the

routes of trial 1 were mainly distributed on the walls of the maze. In trial 4, the worms that

showed learning effects quickly secede from the wall of the maze and find the entrance of the

dark chamber, which showed significantly shorter routes than routes in trial 1, demonstrating

that worms can use distal information to navigate to their destinations.

Fig 2. Results of the light maze experiment. A. Latency of each trial. Data are the average± SEM of worm's latency for worms moved for more than 20 mins in 40 mins. From trial 1 to 6, n=34, 24, 24, 24, 23, 22. B. Worm speed in each trial. C. Worms successfully escaped in each trial. C-G. Representative routes of learned worms. C, E show routes of trial 1. D, F show routes of trial 4. C and D are routes of 1 worm and E, F are routes of 1 worm. The green spots show the start point and the red spots show the end point.

Discussion

In our findings, we designed a light maze to demonstrate that planarians were able to use distal information to localize a dark chamber in the light maze. After training, the worms can use shorter time or shorter distances to navigate to the dark chamber. Below, I'll discuss details and issues in this paradigm.

From the 1950s to the 1960s, numerous studies had been done to understand planarian learning and behavior[15]. The most commonly used procedure is a classical conditioning protocol to make worms associate a light stimulus with an electrical stimulus[16]. Other studies showed that worms can learn more complicated tasks such as discriminating directions in a Y or T maze[10, 17]. However, due to the manual practical difficulties and the control of variables, a large number of research results failed to be reproduced in lots of cases[15]. Consequently, this field became largely abandoned, and even the ability of planarians to form long-term memory was questioned[18].

Some classical conditioning paradigms of the planarians can be attributed to

pseudoconditioning or sensitization instead of true learning and memory encoded by the

brain[19]. Since spatial learning is a more complex behavior that requires the integration of

different neural circuits in higher animals[20], spatial learning might also be a great point to

investigate the memory and learning of planarians, and can be regarded as true learning and

memory.

The control of variables and manipulations of the experimenters can also largely affect the

worms' behavior. In previous studies, light intensity[21], water temperature[22], water

existence[10], time of day[22], time of year[23], chemical components of water[23], chemical

components of food[24], worm's appetite level[25], slime trails[26], worm fatigue state[23],

magnetic fields[27], training conditions and manipulation of the experimenter[23] were

considered to affect the worms' behavior. Our light maze paradigm was designed to decrease

the number of variables as much as possible to make the experiment easy to reproduce. In

our paradigm, the light intensity, water components, water temperature, appetite level and

training time of day are strictly controlled.

The fabrication of mazes using 3D printing techniques is a great advantage in this

research. Designing such a number of mazes of different shapes requests a much higher

cost than 3D printing techniques. With the assistance of the 3D printing techniques, further

investigation of the worm's spatial learning will be much more convenient.

The results demonstrated that worms showed learning effects in each trial. However, a

worm might successfully escape the maze in some trials but not in other trials, which makes it

a little hard to judge the learning effect of the worm. We believe that this phenomenon might

be caused by multiple factors including the habituation phenomenon. I When the worms

habituate to the environment and light stimulation, it might ignore the light and choose to rest

under the light. Or this is because the navigation ability of planarians might be primitive, and

cannot accurately solve every task. So, many worms cannot solve the task in some trials.

Although we endeavored to exclude proximal cues that worms might use to navigate, we

still do not understand which kind of distal cues the worms might have utilized. We speculate

that the worms used the light intensity gradient combined with the geometric information of

the maze to navigate to the dark chamber. The worms might learn to get to brighter place to

find the dark chamber. Also, the worms can learn to secede the wall to find the dark chamber.

To further investigate this question, we think that the position of the dark chamber can be

changed to test whether learned worms can still get into the dark chamber. Also, the electric

and magnetic gradients from the led, earth and/or other lab equipment could also be used as

distal cues for worms to navigate.

In conclusion, we identified the spatial localization ability of planarians by presenting a

paradigm using the light maze we designed. The paradigm can also be a new tool for the

analysis of spatial learning in planarians. In this work, planarian worms were found to use

distal cues instead of proximal cues to navigate to the dark chamber. Since planarians are

now the most primitive animal found to have spatial learning ability, identifying the spatial

localization ability of them might provide insights into the evolution of spatial learning.

References

1.   Vorhees CV, Williams MT. Assessing Spatial Learning and Memory in Rodents. ILAR
Journal. 2014;55(2):310-32. doi: 10.1093/ilar/ilu013 %J ILAR Journal.

2.   Vila-Farré M, J CR. The Ecology of Freshwater Planarians. Methods in molecular
biology (Clifton, NJ). 2018;1774:173-205. Epub 2018/06/20. doi: 10.1007/978-1-4939-
7802-1_3. PubMed PMID: 29916156.
3.   Shettigar N, Joshi A, Dalmeida R, Gopalkrishna R, Chakravarthy A, Patnaik S, et al.
Hierarchies in light sensing and dynamic interactions between ocular and extraocular
sensory networks in a flatworm. Science advances. 2017;3(7):e1603025. Epub
2017/08/07. doi: 10.1126/sciadv.1603025. PubMed PMID: 28782018; PubMed Central
PMCID: PMCPMC5533540.
4.   Inoue T, Hoshino H, Yamashita T, Shimoyama S, Agata K. Planarian shows decision-
making behavior in response to multiple stimuli by integrative brain function. Zoological
Letters. 2015;1(1):7. doi: 10.1186/s40851-014-0010-z.
5.   Allen GD. Reversibility of the Reactions of Planaria Dorotocephala to a Current of
Water. Biological Bulletin. 1915;29(2):111-28. doi: 10.2307/1536302.
6.   Mason PR. Chemo-klino-kinesis in planarian food location. Animal behaviour.
1975;23(2):460-9. Epub 1975/05/01. doi: 10.1016/0003-3472(75)90095-0. PubMed
PMID: 1163855.
7.   !!! INVALID CITATION !!! {}.
8.   Brown FA, Chow CS. Differentiation between Clockwise and Counterclockwise
Magnetic Rotation by the Planarian, Dugesia dorotacephala. Physiological Zoology.
1975;48(2):168-76.
9.   Brown HM, Ogden TE. The electrical response of the planarian ocellus. The Journal
of general physiology. 1968;51(2):237-53. Epub 1968/02/01. doi: 10.1085/jgp.51.2.237.
PubMed PMID: 5641637; PubMed Central PMCID: PMCPMC2201127.
10.   Corning WC. Evidence of Right-Left Discrimination in Planarians. The Journal of
Psychology. 1964;58(1):131-9. doi: 10.1080/00223980.1964.9916729.
11.   Perry CJ, Barron AB, Cheng K. Invertebrate learning and cognition: relating
phenomena to neural substrate. 2013;4(5):561-82. doi:
https://doi.org/10.1002/wcs.1248.
12.   Morris RGM. Spatial localization does not require the presence of local cues.
Learning and Motivation. 1981;12(2):239-60. doi: https://doi.org/10.1016/0023-
9690(81)90020-5.
13.   Merryman S, Sánchez Alvarado A, Jenkin J. Culturing Planarians in the Laboratory.
17742018. p. 241-58.
14.   O'Grady BJ, Geuy MD, Kim H, Balotin KM, Allchin ER, Florian DC, et al. Rapid
prototyping of cell culture microdevices using parylene-coated 3D prints. Lab on a chip.
2021;21(24):4814-22. Epub 2021/11/18. doi: 10.1039/d1lc00744k. PubMed PMID:
34787148; PubMed Central PMCID: PMCPMC8717820.
15.   McConnell JV. Comparative Physiology: Learning in Invertebrates. 1966;28(1):107-
36. doi: 10.1146/annurev.ph.28.030166.000543. PubMed PMID: 5322973.
16.   Thompson R, Mc CJ. Classical conditioning in the planarian, Dugesia dorotocephala.
Journal of comparative and physiological psychology. 1955;48(1):65-8. Epub
1955/02/01. doi: 10.1037/h0041147. PubMed PMID: 14354075.

17. Best JB, Rubinstein I. Maze learning and associated behavior in planaria. Journal of
comparative and physiological psychology. 1962;55:560-6. Epub 1962/08/01. doi:
10.1037/h0044394. PubMed PMID: 13868584.
18. Travis GDL. Replicating Replication? Aspects of the Social Construction of Learning
in Planarian Worms. Social Studies of Science. 1981;11(1):11-32.
19. Kartry AL, Keith-Lee P, Morton WD. PLANARIA: MEMORY TRANSFER THROUGH
CANNIBALISM REEXAMINED. Science (New York, NY). 1964;146(3641):274-5. Epub
1964/10/09. doi: 10.1126/science.146.3641.274. PubMed PMID: 14185328.
20. Lin X, Amalraj M, Blanton C, Avila B, Holmes TC, Nitz DA, et al. Noncanonical
projections to the hippocampal CA3 regulate spatial learning and memory by
modulating the feedforward hippocampal trisynaptic pathway. PLoS biology.
2021;19(12):e3001127. Epub 2021/12/21. doi: 10.1371/journal.pbio.3001127. PubMed
PMID: 34928938; PubMed Central PMCID: PMCPMC8741299.
21. Harden Jones FR. The response of the planarian Dendrocoelum lacteum to an
increase in light intensity. Animal behaviour. 1971;19(2):269-76. doi:
https://doi.org/10.1016/S0003-3472(71)80007-6.
22. Vandeventer JM, Ratner SC. VARIABLES AFFECTING THE FREQUENCY OF
RESPONSE OF PLANARIA TO LIGHT. Journal of comparative and physiological
psychology. 1964;57:407-11. Epub 1964/06/01. doi: 10.1037/h0042776. PubMed PMID:
14155379.
23. McConnell JV. Specific Factors Influencing Planarian Behavior. In: Corning WC,
Ratner SC, editors. Chemistry of Learning: Invertebrate Research. Boston, MA: Springer
US; 1967. p. 217-33.
24. Mori M, Narahashi M, Hayashi T, Ishida M, Kumagai N, Sato Y, et al. Calcium ions in
the aquatic environment drive planarians to food. Zoological Letters. 2019;5(1):31. doi:
10.1186/s40851-019-0147-x.
25. Shomrat T, Levin M. An automated training paradigm reveals long-term memory in
planarians and its persistence through head regeneration. Journal of Experimental
Biology. 2013;216(20):3799-810. doi: 10.1242/jeb.087809 %J Journal of Experimental
Biology.
26. Riccio D, Corning WC. Slime and Planarian Behavior. The Psychological Record.
1969;19(3):507-13. doi: 10.1007/BF03393880.
27. Miller, Kellman J. The influence of geomagnetism on the maze behavior of Dugesia
Tigrina. 2022.

349