Genome Report: A blue mussel chromosome-scale genome assembly for aquaculture, marine
ecology and evolution

Tim Regan[1], Tiago S. Hori[2], Tim P. Bean[1]

[1] The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, UK.

[2] Atlantic Aqua Farms Ltd., Charlottetown, PE, Canada.

Keywords: Aquaculture, evolution, bivalve, mussel, *Mytilus edulis*

Abstract

The blue mussel, *Mytilus edulis* is part of the *Mytilus edulis* species complex, encompassing at least

three putative species: *M. edulis, M. galloprovincialis* and *M. trossulus*. These three species occur on

both sides of the Atlantic and hybridize in nature, and both *M. edulis* and *M. galloprovincialis* are

important aquaculture species. They are also invasive species in many parts of the world. Here, we

present a chromosome-level assembly of *Mytilus edulis* . We used a combination of PacBio sequencing

and Dovetail's Omni-C technology to generate an assembly with 14 long scaffolds containing 94% of

the predicted length of the *M. edulis* genome (1.6 out of 1.7 Gb). Assembly statistics were total length

1.65 Gb, N50 = 116 Mb, L50 = 7 and, L90 = 13. BUSCO analysis showed 92.55% eukaryote BUSCOs

identified. AB-*Initio* annotation using RNA-seq from mantle, gills, muscle and foot predicted 47,128

genes. These gene models were combined with Isoseq validation resulting in 65,505 gene models and

129,708 isoforms. Using GBS and shotgun sequencing, we also sequenced 3 North American

populations of *Mytilus* to characterize single-nucleotide as well as structural variance. This high-quality

genome for *M. edulis* provides a platform to develop tools that can be used in breeding, molecular

ecology and evolution to address questions of both commercial and environmental perspectives.

Introduction

The blue mussel (*Mytilus edulis*) is common to the North Atlantic from Arctic to Mediterranean

regions, with habitat ranging from upper shore to the shallow subtidal (Hayward and Ryland 2017).

M. edulis is known to hybridise with *M. trossulus* in North America and with *M. trossulus* and *M.*

galloprovincialis in Europe. Together, these three species form the *M. edulis* species complex (**Fig. 1**)

(McDonald et al. 1991).

This reef-building bivalve is an ecosystem engineer. Blue mussels dominate fouling communities in

shallow and substrata providing important secondary habitat (Norling and Kautsky 2007). Offshore

wind energy structure surveys found that they can cover the structures with up to 3.4 kg of biomass

m^{-2} (Krone et al. 2013). Through filter-feeding, eutrophication is reduced which can alter ecosystems

(Broszeit et al. 2016). This nutrient cycling ability has been harnessed by using *M. edulis* to study the

fate of persistent organic and metal pollutants (Chase et al. 2001; McEneff et al. 2014), for the

bioremediation of waste (Broszeit et al. 2016) and reduce environmental effects from salmon farms

(MacDonald et al. 2011).

Mussels, a key bivalve production species (FAO 2020), face decreasing wild spat availability for

aquaculture in the UK and elsewhere (Regan et al. 2021). These losses are attributed to multiple

stressors including warmer seas (Seuront et al. 2019) causing a poleward range contraction (Jones et

al. 2010). Additionally, warmer oceans elevate dissolved CO_2 leading to Ocean Acidification (OA)

impacting mussel viability (Asplund et al. 2014) and disease resistance (Ellis et al. 2015). This makes

mussels more susceptible to bacterial pathogenesis (Eggermont et al. 2017; Ripabelli et al. 1999), with

emerging pathogens posing a constant threat (Charles et al. 2020; Cano et al. 2022). Infectious disease

such as disseminated Neoplasia (DN) of *M. trossulus* origin is associated with reduced fitness in *Mytilus*

spp. (Burioli et al. 2021). Furthermore, the effects of hybridisation between the three species of the

Mytilus edulis complex are yet uncertain with suggested negative effect of *M. trossulus* hybridisation

and potential adaptive introgression in the case of *M. galloprovincialis* hybridisation (Fraïsse et al.

2014; Kenchington et al. 2020; Michalek et al. 2021).

To protect the aquaculture industry from these threats, hatchery efforts have been launched in the

UK (Regan et al. 2021), elsewhere in Europe (Kamermans et al. 2013) and in Canada (Gurney-Smith et

al. 2017). However, these efforts have not been straightforward and a better understanding of

fundamental biology is required to achieve commercial success. Despite their importance in

aquaculture and the valuable ecosystem services they provide, no chromosomal assembly existed for

any species within the *Mytilus edulis* species complex prior to this study. Improved genomic tools are

required to address fundamental biological questions such as inheritance patterns and adaptations.

Like many bivalves, the mussel genome is highly heterozygous (3.5%) with an estimated 43% repeat

content. The linear plot of k-mer abundance analysis clearly shows a heterozygous peak in addition to

the homozygous peak and estimates a shorter haploid genome length of 1.18 Gb compared to flow-

cytometry data. The estimated repeat content of the *M. edulis* genome is ~43%. These characteristics

make assembly of these genomes challenging. However, recently genomes for the American oyster,

Pacific Oyster, *M. corruscus* and *M. califoniaus* have been assembled to chromosome-level using short

and long-read sequencing technologies as well as Hi-C-based scaffolding (Yang et al. 2021; Paggeot et

al. 2022a; Penaloza et al. 2021; Gomez-Chiarri et al. 2015). We used a similar approach in this project

to produce a highly contiguous assembly. Practical application of this assembly is demonstrated in

cross-species synteny analyses and in population structure of *Mytilus* individuals sampled from

different regions of the Canadian Atlantic.

70
71 **Fig. 1. TimeTree for *Mytilus edulis***
72 Mytilidae diverged ~387 MYA. Generated using TimeTree (Kumar et al. 2022).

Methods

DNA extraction, library preparation for genome assembly

One naïve blue mussel sample (Anne) was selected from samples collected by the Provincial Department of Communities and Fisheries in the estuary Foxley river in PE. Foxley River was selected as a sampling site because there is no grow-out aquaculture there (i.e. seed from other bays are not transferred to the area). Due to potential introgression of other species of the *Mytlius* species complex (e.g. *M. trossulus*), this sample was genotyped using 12 SNPs described by (Wilson et al. 2018). We sampled the gill, mantle and muscle of sample "Anne" aseptically, flash froze fresh tissues in liquid nitrogen and preserved them at -80°C. Tissues were shipped to Dovetail Genomics in Scott's Valley, CA, in excess dry ice. Dovetail extracted high molecular weight (HMW) DNA using an in-house modified CTAB method. Dovetail prepared and sequenced PacBIO SMRTbell libraries to a depth of 196X using the Sequel II sequencer. Processed PacBio data (from Dovetail) can be found on SRA under accession number SRX11246493.

Raw contig assembly, scaffold formation and polishing

Dovetail generated a primary contig-level assembly using wtdbg2 (Ruan and Li 2020). The contig-level assembly was filtered of putative duplicated haplotypes and contaminants using Purge Haplotigs (Roach et al. 2018) and Blobtools2 (Challis et al. 2020), respectively. Scaffolding was performed using Omni-C libraries and the HiRise assembler (v1.0). The same DNA sample used by Dovetail was shipped to UWM (University of Wisconsin – Madison) and sequenced to a 100X using the NovaSeq sequencer. These data were trimmed using Trimmomatic (v) and used for polishing with racon (v1.4.3) (Vaser et al. 2017). Completeness was evaluated using compleasm v0.2.5 (Huang and Li 2023) (eukaryota_odb10: 255 BUSCOS, metazoa_odb10: 954 BUSCOS and mollusca_odb10: 5295 BUSCOS) (Manni et al. 2021) and Merqury (Rhie et al. 2020). Merqury analysis was carried out with the same read set used for polishing as the original PacBio CLR reads were not suitable for this analysis. General

length metrics were obtained using QUAST (v5.0.1) (Gurevich et al. 2013; Mikheenko et al. 2018).

Synteny mapping between the *M. edulis* and the *M. coruscus* (GCA_017311375.1) (Yang et al. 2021)

was done using the MCScanX.h function of MCScanX (v2) (Wang et al. 2012). Putative orthologous

groups were identified with Orthofinder (v.2.5.4) using predicted gene structures for *M. edulis* (this

work) and *M. coruscus* (Yang et al. 2021). Dot plots and circle plots were generated using MCScanX

(v.2).

RNA preparation and Isoseq analysis

Isoseq3 analysis of CSS data from muscle, gill, hemolymph, and foot (2.9 million reads -188,165 Mb)

identified 216,343 high-quality putative full-length (FL) transcripts (from 2.8 million reads containing

poly-A tails). We shipped flash frozen gill and adductor muscle tissue from sample "Anne" to the

Biotechnology Centre Core facility at the University of Wisconsin, Madison (UWM). Samples were

homogenized using a Qiagen Tissuelyser (2 min @ 20 Hz). RNA was extracted using the RNeasy Mini

Kit (Qiagen) with on-column DNAse treatment. UWM performed RNA QC with a nanodrop and

bioanalyzer and prepared libraries using the Iso-seq Library SMRTbell express template prep kit.

Libraries were sequenced using one Sequel II SMRT cell in CSS mode (i.e. HiFi reads). UWM provided

de-multiplexed processed HiFi reads. RNA samples from foot, mantle and gut were sent to the

Genome Excellence Centre (Genome Quebec) in Montreal. HiFi sequencing was carried out as above.

Putative   full-length   transcripts   were   identified   using   the   IsoSeq3   pipeline

(https://github.com/ylipacbio/IsoSeq3). Putative open read frames (ORF) were identified using

TransDecoder (v5.5.0) (https://github.com/TransDecoder/TransDecoder/wiki).

Annotation

Repeat modeller (4.1.0) (Flynn et al. 2020) was used to predict repeat motifs for *M. edulis* and Repeat

Masker (2.0.2a) (Smit et al. 2015) was used to mask the final assembly. Ab-initio annotation was done

using Augustus (v3.4.0) (Stanke et al. 2006) trained with genes from *Mytilus galloprovincialis*, *Mytilus*

coruscus and *Crassostrea virginica*. Additional hints were generated using short-read RNA-seq (data

not shown) and Isoseq data generated herein. The ab-initio annotation was updated using PASA (v2.5)

with alignments of a de-novo transcriptome produced using Trinity (v2.8.15) (Grabherr et al. 2011) in

Genome-Guided mode. Two runs through PASA were used to update the *ab-initio* annotation. Full-

length (FL) transcripts from Isoseq3 pipeline were mapped to the genome using pbmm2 with pre-sets

for Isoseq and filtered based on quality Isoseq3 collapse (minimum alignment identity/coverage:

0.90/0.90) and Isoseq3 refine. We used the resulting gff annotation to run SQANTI3 (v4.3.0). We used

sqanti_qc.py to generate quality data for sqanti_filter.py. Filtering to remove artifacts was carried out

using the default parameters. Lastly, ab-initio predications and filtered Isoseq FLs were merged with

AGAT (v.1.2.0). Using agat_sp_merge.pl, we removed duplicate gene models/isoform and assigned

orphan isoforms from the Isoseq data when possible. Amino acid sequences were translated from the

CDS using agat_sp_extract_sequences using options -t "CDS" -p --cfs --acs –asc.

Sample collection for SNP discovery and population structure

We collected gill samples for DNA extractions using standard molecular biology techniques and

preserved them in non-denatured 100% ethanol. For population genetics analysis, we sampled

mussels in sets of 96 from west to east PEI in Foxley River (FOX) (wild population), Malpeque Bay (MB)

(North Shore, Prince County); French River (FR), Stanley Bridge (ST), Whetley River (WR), Tracadie

(TRC) (North Shore, Queen county); Orwell (OR) (South Shore, Queen's county; Morell (MRL) (North

Shore – King's county) and; Murray River (MR) (South Shore, King county) (**Fig. 2**). Seed deployed on

these sites were originally collected in St. Peter's bay, Brudenell River, and Malpeque bay. We also

collected samples in the Bras D'or Lake in Cape Breton (Nova Scotia), the Magdalene Island (Québec)

and Notre Dame bay in Newfoundland.

SNP markers, Admixture analysis, population structures.

Samples for SNP discovery and population genetics were shipped to LGC genomics in Berlin.

Restriction-associated DNA libraries (RAD) libraries were prepared by LGC with using MslI, normalized

and sequenced using a NextSeq sequencer. The resulting reads were trimmed and checked for the

restriction site by LGC. This final read set was used to identify SNPs and call individual genotypes using

Tassel5 (v.5.2.4). SNPs were filtered using MAF (<0.01) and percent of individuals with genotypes. For

the SNP discovery, the samples from Cape Breton were excluded from filtering analysis. These samples

were shown to be a pure *M. trossulus* population and had missing calls for a significant number of

sites. For population genetics analysis, a second set of SNPs that were successfully called across all

populations was used. All samples were also genotyped for the 12 species discrimination SNPs from

(Wilson et al. 2018). All population genetics analyses were performed using the dpca function of the

R package Adegenet (v.2.1.7) and STRUCTURE (v.2.3.4) or fastSTRUCTURE (v1.0) (K 3 to K10, 5000

repetitions, 1000 burn in). Analysis of species discrimination SNPs included the genotypes published

by (Corrochano-Fraile et al. 2022) as outgroups.

Figure 2: Sampling Locations in PEI

Results and Discussion

Genome Assembly and Annotation

The chromosome-level assembly presented herein was produced in two stages. First, ~15 million

PacBio CLR reads (~340 Gb) were produced, representing coverage of 196X for an estimated genome

size of 1.7 Gb (Hinegardner 1974; Rodríguez-Juíz et al. 1996). These reads were assembled into contigs

using wtdbg2, which uses uncorrected reads (Ruan and Li 2020). The primary assembly was 1.96 Gb

long in 17,825 contigs and a N50 of 443 Kb. After haplotype purging and contaminant removal, the

final contig assembly had 10,111 contigs, for a total of 1,65 Gb and a N50 of 518 Kb. Following

scaffolding using Omni-C libraries and the HiRise assembler, we generated a primary chromosome-

level assembly made of 2,117 contigs. We removed putative contaminants using Blobtools by

eliminating sequences coming from non-molluscan organisms. This assembly was further filtered to

contain only sequences > 5,000 bp. The resulting draft is deposited on NCBI assembly under accession

number GCA_019925275.1. The final assembly is made of 1,119 contigs and has an N50 of 116 Mb.

The 14 putative *M. edulis* chromosomes are deposited under accession numbers CM034349.1 to

CM34362.1. Detailed statistics for the assemblies can be found in **Table 1**.

	Anne (This study)	Corrochano-Fraile *et al.* (2022)	xbMytEdul2.1, Darwin Tree of Life (2024)
Length	1,659,567,081 bp	1,827,085,763	1,374,471,240
# of scaffolds	1,119	3,339	2,563
# of contigs	9,866	5,966	3,754
N50	116,503,180	1,097,279	1,734,586
Coverage	196.0x	152.0x	30.0x
Completeness (Merqury)	76.71	NA	NA
QV	32.39	NA	NA

Table 1. Summary statistics for *M. edulis* assemblies

Compared to the assembly published in Corrochano-Fraile et al. (2022), the assembly presented

herein has better contiguity (**Table 1**). Our assembly is also shorter than the assembly from

Corrochano-Fraile et al. (2022). The two assemblies' length falls close to the estimated size of the

genome based on c-value (1.7) (Rodríguez-Juíz et al. 1996) and is significantly longer than what is

estimated by k-mer abundance analysis with GenomeScope (1.18 Gb).

Despite the total assembly length being close to that estimated using c-values, k-mer-based

completeness analysis recovers only 76% in a set of Illumina reads origination from sample "Anne".

When the putative purged haplotigs were added back to the assembly, recovery was ~83%. This

apparent low k-mer recovery is probably a combination of the consensus being different from either

haploid genomes, the error rate in the original PacBio data, the fact that the reads used for polishing

were not used from the primary assembly, the contigs removed based on length or contaminant

status, and the gaps arising from the Omni-C scaffolding. It is also possible that the high heterozygosity

affects the accuracy of k-mer abundance analysis, as shown by the large discrepancy between genome

size estimates.

Completeness analysis in Merqury resulted in 76.71% recovery of k-mers present in the polishing

Illumina data from the final version of the assembly. We also evaluated the completeness of the

assembly when combined with purged haplotypes, which was 83.44%. QV value for the primary

assembly was 32.39, while the combined draft had a QV of 30.73. Compleasm BUSCO analysis showed

a recovery of 92.55%, 91.95%, and 88.5% complete BUSCOS against the eukaryote, metazoan and

molluscan databases, respectively. The 14 putative chromosomes represent ~96% of the assembly,

with lengths varying from 140 Mbp to 90 Mbp. These data and the N50 metric show that this assembly

has high contiguity and that this assembly and its annotation will be highly useful for aquaculture,

evolution and molecular ecology studies. Herein, we illustrate the possible applications of this

assembly by performing population and synteny analyses.

SNP discovery and population structure

Due to the close relationship between the members of the *Mytilus* species complex, we wanted to

verify that the individual sampled (Anne) was pure *M. edulis*. Population Structure analysis conducted

using 12 SNPs (Wilson et al. 2018) clearly separated *M. edulis* populations in PEI from other regions of

Canada. We generated two sets of SNPs: the first set totalling 71,231 SNPs using only samples from

PEI and made a polymorphic collection of SNPs in *M. edulis.* The second set, with ~6,000 markers, is a

polymorphic set of SNPs in both *M. edulis* and *M. trossulus.* Population structure and putative

admixture are shown in **Figure 4**. In the DPCA and PCA, untrained clustering clearly separated both CB

from PEI/NL/MAG and also the putative populations in the Gulf of St. Lawrence. In green, are the

majority of samples from NL and MAG, while blue represents individuals from PEI.

We used *M. trossulus*, *M. galloprovincialis* and European *M. edulis* genotypes as outgroups. The later

inclusion of 96 individuals from Cape Breton (NS) showed no significant evidence of *M. trossulus*

introgression in PEI samples. Given that Cape Breton has long been considered a pure *M. trossulus*

population (Wilson et al., 2018), we are confident that the sample Anne represents an *M. edulis*

individual. We also genotyped over 500 PEI individuals and 96 samples from Magdalene Island and 96

individuals from Newfoundland with the same 12 SNP panel. As before no significant introgression of

M. trossulus was detected in PEI. However, we only genotyped 96 samples collected in an area with

no grow-out leases (Foxley River). Although unlikely, we cannot rule out the possibility of a sampling

bias in aquaculture sites favouring *M. edulis.* DPCA and Structure analysis indicate that there is low

population stratification between different regions of PEI, while the Magdalene islands and NL are

distinct populations. The populations from NL and the Magdalene Islands are more similar to each

other than from the populations from PEI. For animals from one sampling event at S.t Peter's bay, we

found evidence of shared genetics between PEI and the population to the northeast of the island.

Figure 4 Population structure in the North American Atlantic. (A) DPCA with 6,000 SNPs, (B) PCA
with 6,000 SNPs, (C) fastStructure with 6,000 SNPs. Samples from Cape Breton NS (CB), FR, TRC, MB,
FOX, MRL, U, WR, MR, OR (PEI), Magdalene Islands QC (MAG), Notre Dame Bay NL (NFL)

 Annotation and synteny analysis

We identified 196,111 putative open reading frames from the 216,343 FL transcripts using Isoseq3

analysis of CSS data from muscle and gill (2.9 million reads -188,165 Mb). BLASTp analysis against the

uniref90 database returned informative hits for ~80% (164,969) of these translated transcripts. *Ab-*

initio gene prediction in Augustus detected 46,604 gene models that produced 46,604 transcripts after

filtering based on evidence support. After two rounds of PASA updates, the final number of gene

models was 47,128 and the number of transcripts was 55,138. After Isoseq refine and collapse 85,099

isoforms survived and following SQANTI3 filtering 70,592 isoforms remained. They were assigned to

31,211 unique genes. Finally, the combined annotation has 65,505 gene models and 129,708 isoforms.

Proteins were translated from the CDS ensuring only complete CDS were translated and that isoforms

were not incorrectly fused together. This resulted in 45,379 amino acid sequences. Compleasm BUSCO

analysis of these 45,379 proteins showed a recovery of 78.43% and 76.83% complete BUSCOS against

the eukaryote and metazoan databases, respectively.

Synteny analysis showed a high degree of collinearity between putative chromosomes of *M. edulis*

and *M. coruscus* (**Fig. 3**). However, putative inversions, transposition and deletion can be observed in

almost all chromosomes. Gene order in chromosomes represented by sequences CM034349.1 (*M.*

edulis chromosome 1) - CM029595.1 (*M. coruscus* LG1) and CM034343.1 (*M. edulis* chromosome 4) –

CM029599.1 (*M. coruscus* LG5) showed the highest conservation. Putative orthologous relationships

between *M. edulis* chromosomes and *M. coruscus* linkage groups (LG) are shown in **Table 2**.

The taxonomic status of the "species" in the *Mytilus* species complex remains in debate. Chromosome-

level assemblies allow the study of macroevolution of the genome by looking at synteny across

species. Herein, we present the synteny analysis between the 14 putative chromosomes of *M. edulis*

and *M. corruscus* (Yang et al. 2021) to exemplify how chromosome-level assemblies may allow us to

better understand the phylogenetic relationships within the genus *Mytilus*. This analysis shows that

some of putative orthologous chromosomes of the 2 species maintain high-levels of collinearity (e.g.

chromosomes 1 and 4 from *M. edulis* with LG1 and LG5 from *M. coruscus* respectively) while others

present significant levels of re-arrangements (e.g. chromosomes 7 and 11 from *M. edulis* with LG2 and

LG9 from *M. coruscus* respectively). An in-depth analysis of chromosome synteny will shed light on

the level of collinearity between multiple members of the genus *Mytilus*. Homology between

chromosomes is a key element of the viability of hybrids. Reproductive isolation tends to increase

during speciation, and these resources will permit further studies on the reproductive compatibility

of the species in genus *Mytilus* at the chromosome level.

Figure 3. Synteny between the *M. edulis* and *M. coruscus* putative chromosomes

Table 2: Putative synteny between *M. edulis* and *M. coruscus*. Ids are NCBI Assembly database Molecule name and accession number.

M. edulis	*M. coruscus*
Chromosome 1 (CM034349.1)	LG01 (CM029595.1)
Chromosome 2 (CM034350.1)	LG04 (CM029598.1)
Chromosome 3 (CM034351.1)	LG03 (CM029597.1)
Chromosome 4 (CM034352.1)	LG05 (CM029599.1)
Chromosome 5 (CM034353.1)	LG10 (CM029604.1)
Chromosome 6 (CM034354.1)	LG06 (CM029600.1)
Chromosome 7 (CM034355.1)	LG02 (CM029596.1)
Chromosome 8 (CM034356.1)	LG11 (CM029605.1)
Chromosome 9 (CM034357.1)	LG07 (CM029601.1)
Chromosome 10 (CM034358.1)	LG08 (CM029602.1)
Chromosome 11 (CM034359.1)	LG09 (CM029603.1)
Chromosome 12 (CM034360.1)	LG14 (CM029608.1)
Chromosome 13 (CM034361.1)	LG12 (CM029606.1)
Chromosome 14 (CM034362.1)	LG13 (CM029607.1)

Here, we present a highly contiguous chromosome assembly for *Mytilus edulis* confirming species-

level individual purity through resequencing. To date, our resource has been applied in multiple

studies analysing *Mytilus* genome assemblies (Paggeot et al. 2022b; Gallardo-Escarate et al. 2023) and

cross-species gene orthology analyses (Saco et al. 2023). The gene annotations produced in this study

were generated using Augustus gene model predictions integrating full transcript Isoseq data and

applying stringent filtering parameters. This comprehensive approach provides a robust foundation

for future cross-species analyses and biological studies on gene function within the *Mytilus* species

complex.

Data Availability:

The resulting draft is deposited on NCBI assembly under accession number GCA_019925275.1.

Funding Statement:

"Breeding Better Mussels (Mytilus edulis): developing genomic tools for the implementation of a
modern and sustainable mussel Breeding Program" (Grant ID GA-PE-RP3, Genome Canada)

The authors would like to acknowledge funding from BBSRC Institute Strategic Programme grants
(BBS/E/RL/230001A, BBS/E/RL/230001B and BBS/E/RL/230002B).

Acknowledgements

Industry Partners: Mussel King (Scott Dockendorff), PEI Aqua Farms (John Paquet), Atlantic Aqua
Farms (Terry Ennis).

PEIMSO: Angela MacDonald, Neil McNair, Aaron Ramsey, Jessica Fry

Genome Atlantic: Kristin Tweel, Charmaine Gaudet, Nil d'Entremont

Service Providers: LGC Berlin and UK, Genome Quebec, Dovetail Genomics

Asplund, M.E., S.P. Baden, S. Russ, R.P. Ellis, N. Gong *et al.*, 2014 Ocean acidification and host-
pathogen interactions: blue mussels, Mytilus edulis, encountering Vibrio tubiashii. *Environ*
Microbiol 16 (4):1029-1039.
Broszeit, S., C. Hattam, and N. Beaumont, 2016 Bioremediation of waste under ocean acidification:
Reviewing the role of Mytilus edulis. *Marine Pollution Bulletin* 103 (1):5-14.
Burioli, E.A.V., M. Hammel, N. Bierne, F. Thomas, M. Houssin *et al.*, 2021 Traits of a mussel
transmissible cancer are reminiscent of a parasitic life style. *Scientific Reports* 11 (1):24110.
Cano, I., A. Parker, G.M. Ward, M. Green, S. Ross *et al.*, 2022 First Detection of Francisella halioticida
Infecting a Wild Population of Blue Mussels Mytilus edulis in the United Kingdom. *Pathogens*
11 (3).
Challis, R., E. Richards, J. Rajan, G. Cochrane, and M. Blaxter, 2020 BlobToolKit – Interactive Quality
Assessment of Genome Assemblies. *G3 Genes|Genomes|Genetics* 10 (4):1361-1374.
Charles, M., A. Villalba, G. Meyer, S. Trancart, C. Lagy *et al.*, 2020 First detection of Francisella
halioticida in mussels Mytilus spp. experiencing mortalities in France. *Dis Aquat Organ*
140:203-208.
Chase, M., S.H. Jones, P. Hennigar, J. Sowles, G. Harding *et al.*, 2001 Gulfwatch: Monitoring spatial
and temporal patterns of trace metal and organic contaminants in the Gulf of Maine (1991–
1997) with the blue mussel, Mytilus edulis L. *Marine Pollution Bulletin* 42 (6):490-504.
Corrochano-Fraile, A., A. Davie, S. Carboni, and M. Bekaert, 2022 Evidence of multiple genome
duplication events in Mytilus evolution. *BMC Genomics* 23 (1):340.
Eggermont, M., P. Bossier, G.S.J. Pande, V. Delahaut, A.M. Rayhan *et al.*, 2017 Isolation of
Vibrionaceae from wild blue mussel (Mytilus edulis) adults and their impact on blue mussel
larviculture. *FEMS Microbiology Ecology* 93 (4).
Ellis, R.P., S. Widdicombe, H. Parry, T.H. Hutchinson, and J.I. Spicer, 2015 Pathogenic challenge
reveals immune trade-off in mussels exposed to reduced seawater pH and increased
temperature. *Journal of Experimental Marine Biology and Ecology* 462:83-89.

FAO, 2020 The State of World Fisheries and Aquaculture 2020. Sustainability in action. , pp. #244 p.
in *The State of World Fisheries and Aquaculture (SOFIA)*. FAO, Rome, Italy.
Flynn, J.M., R. Hubley, C. Goubert, J. Rosen, A.G. Clark *et al.*, 2020 RepeatModeler2 for automated
genomic discovery of transposable element families. *Proceedings of the National Academy*
of Sciences 117 (17):9451-9457.
Fraïsse, C., C. Roux, J.J. Welch, and N. Bierne, 2014 Gene-flow in a mosaic hybrid zone: is local
introgression adaptive? *Genetics* 197 (3):939-951.
Gallardo-Escarate, C., V. Valenzuela-Munoz, G. Nunez-Acuna, D. Valenzuela-Miranda, F.J. Tapia *et al.*,
2023 Chromosome-Level Genome Assembly of the Blue Mussel Mytilus chilensis Reveals
Molecular Signatures Facing the Marine Environment. *Genes (Basel)* 14 (4).
Gomez-Chiarri, M., W.C. Warren, X. Guo, and D. Proestou, 2015 Developing tools for the study of
molluscan immunity: The sequencing of the genome of the eastern oyster, Crassostrea
virginica. *Fish Shellfish Immunol* 46 (1):2-4.
Grabherr, M.G., B.J. Haas, M. Yassour, J.Z. Levin, D.A. Thompson *et al.*, 2011 Full-length
transcriptome assembly from RNA-Seq data without a reference genome. *Nature*
Biotechnology 29 (7):644-652.
Gurevich, A., V. Saveliev, N. Vyahhi, and G. Tesler, 2013 QUAST: quality assessment tool for genome
assemblies. *Bioinformatics* 29 (8):1072-1075.
Gurney-Smith, H.J., A.J. Wade, and C.L. Abbott, 2017 Species composition and genetic diversity of
farmed mussels in British Columbia, Canada. *Aquaculture* 466:33-40.
Hayward, P.J., and J.S. Ryland, 2017 *Handbook of the marine fauna of North-West Europe*: Oxford
University Press.
Hinegardner, R., 1974 Cellular DNA content of the Mollusca. *Comp Biochem Physiol A Comp Physiol*
47 (2):447-460.
Huang, N., and H. Li, 2023 compleasm: a faster and more accurate reimplementation of BUSCO.
Bioinformatics 39 (10).
Jones, S.J., F.P. Lima, and D.S. Wethey, 2010 Rising environmental temperatures and biogeography:
poleward range contraction of the blue mussel, Mytilus edulis L., in the western Atlantic.
Journal of Biogeography 37 (12):2243-2259.
Kamermans, P., T. Galley, P. Boudry, J. Fuentes, H. McCombie *et al.*, 2013 11 - Blue mussel hatchery
technology in Europe, pp. 339-373 in *Advances in Aquaculture Hatchery Technology*, edited
by G. Allan and G. Burnell. Woodhead Publishing.
Kenchington, E.L., B.W. MacDonald, A. Cogswell, L.C. Hamilton, and A.P. Diz, 2020 Sex-specific
effects of hybridization on reproductive fitness in Mytilus. *Journal of Zoological Systematics*
and Evolutionary Research 58 (2):581-597.
Krone, R., L. Gutow, T.J. Joschko, and A. Schröder, 2013 Epifauna dynamics at an offshore foundation
– Implications of future wind power farming in the North Sea. *Marine Environmental*
Research 85:1-12.
Kumar, S., M. Suleski, J.M. Craig, A.E. Kasprowicz, M. Sanderford *et al.*, 2022 TimeTree 5: An
Expanded Resource for Species Divergence Times. *Mol Biol Evol*.
MacDonald, B.A., S.M.C. Robinson, and K.A. Barrington, 2011 Feeding activity of mussels (Mytilus
edulis) held in the field at an integrated multi-trophic aquaculture (IMTA) site (Salmo salar)
and exposed to fish food in the laboratory. *Aquaculture* 314 (1):244-251.
Manni, M., M.R. Berkeley, M. Seppey, and E.M. Zdobnov, 2021 BUSCO: Assessing Genomic Data
Quality and Beyond. *Current Protocols* 1 (12):e323.
McDonald, J., R. Seed, and R. Koehn, 1991 Allozymes and morphometric characters of three species
ofMytilus in the Northern and Southern Hemispheres. *Marine Biology* 111 (3):323-333.
McEneff, G., L. Barron, B. Kelleher, B. Paull, and B. Quinn, 2014 A year-long study of the spatial
occurrence and relative distribution of pharmaceutical residues in sewage effluent, receiving
marine waters and marine bivalves. *Science of the Total Environment* 476:317-326.

Michalek, K., D.L.J. Vendrami, M. Bekaert, D.H. Green, K.S. Last *et al.*, 2021 Mytilus trossulus
introgression and consequences for shell traits in longline cultivated mussels. *Evolutionary*
applications 14 (7):1830-1843.
Mikheenko, A., A. Prjibelski, V. Saveliev, D. Antipov, and A. Gurevich, 2018 Versatile genome
assembly evaluation with QUAST-LG. *Bioinformatics* 34 (13):i142-i150.
Norling, P., and N. Kautsky, 2007 Structural and functional effects of Mytilus edulis on diversity of
associated species and ecosystem functioning. *Marine Ecology Progress Series* 351:163-175.
Paggeot, L.X., M.B. DeBiasse, M. Escalona, C. Fairbairn, M.P.A. Marimuthu *et al.*, 2022a Reference
genome for the California ribbed mussel, Mytilus californianus, an ecosystem engineer. *J*
Hered.
Paggeot, L.X., M.B. DeBiasse, M. Escalona, C. Fairbairn, M.P.A. Marimuthu *et al.*, 2022b Reference
genome for the California ribbed mussel, Mytilus californianus, an ecosystem engineer. *J*
Hered 113 (6):681-688.
Penaloza, C., A.P. Gutierrez, L. Eory, S. Wang, X. Guo *et al.*, 2021 A chromosome-level genome
assembly for the Pacific oyster Crassostrea gigas. *Gigascience* 10 (3).
Regan, T., T.P. Bean, T. Ellis, A. Davie, S. Carboni *et al.*, 2021 Genetic improvement technologies to
support the sustainable growth of UK aquaculture. *Reviews in Aquaculture* 13 (4):1958-1985.
Rhie, A., B.P. Walenz, S. Koren, and A.M. Phillippy, 2020 Merqury: reference-free quality,
completeness, and phasing assessment for genome assemblies. *Genome Biology* 21 (1):245.
Ripabelli, G., M.L. Sammarco, G.M. Grasso, I. Fanelli, A. Caprioli *et al.*, 1999 Occurrence of Vibrio and
other pathogenic bacteria in Mytilus galloprovincialis (mussels) harvested from Adriatic Sea,
Italy. *International Journal of Food Microbiology* 49 (1):43-48.
Roach, M.J., S.A. Schmidt, and A.R. Borneman, 2018 Purge Haplotigs: allelic contig reassignment for
third-gen diploid genome assemblies. *BMC Bioinformatics* 19 (1):460.
Rodríguez-Juíz, A.M., M. Torrado, and J. Méndez, 1996 Genome-size variation in bivalve molluscs
determined by flow cytometry. *Marine Biology* 126:489-497.
Ruan, J., and H. Li, 2020 Fast and accurate long-read assembly with wtdbg2. *Nature Methods* 17
(2):155-158.
Saco, A., B. Novoa, S. Greco, M. Gerdol, and A. Figueras, 2023 Bivalves Present the Largest and Most
Diversified Repertoire of Toll-Like Receptors in the Animal Kingdom, Suggesting Broad-
Spectrum Pathogen Recognition in Marine Waters. *Molecular Biology and Evolution* 40 (6).
Seuront, L., K.R. Nicastro, G.I. Zardi, and E. Goberville, 2019 Decreased thermal tolerance under
recurrent heat stress conditions explains summer mass mortality of the blue mussel Mytilus
edulis. *Scientific Reports* 9 (1):17498.
Smit, A., R. Hubley, and P. Green, 2015 RepeatMasker Open-4.0. *http://www.repeatmasker.org*.
Stanke, M., O. Keller, I. Gunduz, A. Hayes, S. Waack *et al.*, 2006 AUGUSTUS: ab initio prediction of
alternative transcripts. *Nucleic acids research* 34 (suppl_2):W435-W439.
Vaser, R., I. Sović, N. Nagarajan, and M. Šikić, 2017 Fast and accurate de novo genome assembly
from long uncorrected reads. *Genome Res* 27 (5):737-746.
Wang, Y., H. Tang, J.D. Debarry, X. Tan, J. Li *et al.*, 2012 MCScanX: a toolkit for detection and
evolutionary analysis of gene synteny and collinearity. *Nucleic acids research* 40 (7):e49.
Wilson, J., I. Matejusova, R.E. McIntosh, S. Carboni, and M. Bekaert, 2018 New diagnostic SNP
molecular markers for the Mytilus species complex. *PLoS One* 13 (7):e0200654.
Yang, J.L., D.D. Feng, J. Liu, J.K. Xu, K. Chen *et al.*, 2021 Chromosome-level genome assembly of the
hard-shelled mussel Mytilus coruscus, a widely distributed species from the temperate areas
of East Asia. *Gigascience* 10 (4).

422