Semaphorin7A patterns neural circuitry in the lateral line of the zebrafish

Agnik Dasgupta[1,*], Caleb C. Reagor[1,2], Sang P. Paik[1,3], Lauren M. Snow[1,4], Adrian Jacobo[1,5], and A. J. Hudspeth[1]

[1] Howard Hughes Medical Institute and Laboratory of Sensory Neuroscience, The Rockefeller University, New York, NY, USA

*Corresponding author: Dr. Agnik Dasgupta

Laboratory of Sensory Neuroscience

The Rockefeller University

New York, NY 10065

Telephone: 212-327-7353

E-mail: adasgupta@rockefeller.edu

[2] Tri-Institutional PhD Program in Computational Biology and Medicine, New York, NY, USA

[3] Present address: Department of Functional Neuroanatomy, Institute for Anatomy and Cell Biology, Heidelberg University, Heidelberg, Germany

[4] Present address: Brigham and Women's Hospital, Boston, MA, USA

[5] Chan Zuckerberg Biohub San Francisco, San Francisco, CA, USA

Abstract

In a developing nervous system, axonal arbors often undergo complex rearrangements before

neural circuits attain their final innervation topology. In the lateral line sensory system of the

zebrafish, developing sensory axons reorganize their terminal arborization patterns to establish

precise neural microcircuits around the mechanosensory hair cells. However, a quantitative

understanding of the changes in the sensory arbor morphology and the regulators behind the

microcircuit assembly remain enigmatic. Here, we report that Semaphorin7A (Sema7A) acts

as an important mediator of these processes. Utilizing a semi-automated three-dimensional

neurite tracing methodology and computational techniques, we have identified and

quantitatively analyzed distinct topological features that shape the network in wild-type and

Sema7A loss-of-function mutants. In contrast to those of wild-type animals, the sensory axons

in Sema7A mutants display aberrant arborizations with disorganized network topology and

diminished contacts to hair cells. Moreover, ectopic expression of a secreted form of Sema7A

by non-hair cells induces chemotropic guidance of sensory axons. Our findings propose that

Sema7A likely functions both as a juxtracrine and as a secreted cue to pattern neural circuitry

during sensory organ development.

Introduction

Pathfinding axons are directed to their appropriate synaptic targets by a variety of guidance

cues. While approaching or traversing the target field, the growth cones of migrating axons

encounter secreted or cell surface-attached ligands and respond by steering toward or away

from their sources (1–4). Fine-grained control of these factors is critical for the establishment

of proper neural circuitry.

The zebrafish's lateral line provides a tractable *in vivo* system for exploring the

mechanisms that guide the assembly of neural circuits in a peripheral nervous system. In

particular, it is possible to detect individual hair cells and the sensory axons that innervate them

throughout the first week of development (5). The primary posterior lateral line on each side of

the zebrafish's tail consists of about seven neuromasts, which contain mechanoreceptive hair

cells of opposing polarities—half sensitive to headward (rostrad) water motion and the

complementary half to tailward (caudad) water motion—that cluster at their centers (6). The

sensory axons of the lateral-line nerve branch, arborize, and consolidate around the basolateral

surfaces of the hair cells (Figure 1A). To aid in an animal's swimming behavior, these hair cells

sense water currents and relay signals to the brain through the sensory axons (7). The well-

defined structure of the ramifications and synaptic contacts of the sensory axons highlight the

need to determine the molecular signals behind the assembly of such precise microcircuits (8).

We adopted a candidate-gene strategy to seek factors that direct the growth of afferent

growth cones in the zebrafish's lateral line. Analysis of single-cell RNA sequencing data

identified the *semaphorin7a* (*sema7a*) gene to be highly expressed in hair cells of that organ

(9). Semaphorins are important regulators of axonal growth and target finding during the

patterning of diverse neural circuits (10). The semaphorin family includes proteins that are

secreted, transmembrane, and cell surface-attached (11). Among these, Semaphorin7A

(Sema7A) is the only molecule that is anchored to the outer leaflet of the lipid bilayer of the cell

membrane by a glycosylphosphatidylinositol (GPI) linker (11). Unlike many other semaphorins,

which act as repulsive cues (12,13), Sema7A is involved in promoting axon growth (4) and

imparting directional signals by interacting with the integrin and plexin families of receptors

residing on the pathfinding axons (14–16). Sema7A likely occurs *in vivo* in both GPI-anchored

and soluble forms (17–19), and studies of neuronal explants confirm that both forms can induce

directed axonal outgrowth (4). Furthermore, it has been proposed that the GPI anchor is

cleaved by membrane-resident GPI-specific phospholipases (GPI-PLs) or matrix

metalloproteases to release the Sema7A into the extracellular environment and thus regulate

dynamic cellular processes (20,21). Because these observations indicate that Sema7A might

influence neuronal development both as a juxtracrine and as a diffusive signal, we investigated

the role of Sema7A in sculpting a vertebrate peripheral sensory organ.

Results

Sema7A expression and localization in hair cells. The zebrafish genome contains a single

sema7a gene that produces two transcripts. One transcript encodes Sema7A-GPI, a full-

length, GPI-linked, cell surface-attached protein. The second transcript yields Sema7A^{sec}, a

protein with a truncated C-terminus, which we conjecture is secreted into the local environment.

Each transcript encodes an N-terminal signal sequence and a single copy of the conserved

Sema domain (11). To detect the presence of both *sema7a* transcripts in the hair cells of the

lateral-line neuromasts, we used fluorescence-activated cell sorting (FACS) to capture the

labeled hair cells (22) and isolated total RNA. Using primers that flank the regions encoding

the conserved Sema domain and the distinct C-termini of the two transcripts, we performed

reverse transcription and polymerase chain reactions (RT-PCR) to identify both the membrane-

anchored and the secreted transcript in developing larvae (Figure 1B-D; Figure 1—figure

supplement 1A-D). Single-cell RNA sequencing data also have shown that the GPI-linked

sema7a transcript is particularly enriched in the immature and mature hair cells of the

neuromast during early larval development (9), a period when sensory axons of the posterior

lateral line contact hair cells and form synapses with them (5,23). Indeed, we have observed

Sema7A to be specifically localized in both mature and immature hair cells (Figure 1E; Figure

1—figure supplement 2A-D).

As neuromasts mature, the contacts between sensory axons and hair cells increase in

number, become stabilized, and establish synapses (5,23). Because Sema7A can modulate

axon guidance (4,14) and synapse formation (15), we wondered whether the level of Sema7A

also changes during neuromast development. Upon quantifying the average intensity of

Sema7A, we found similar expression in rostrally and caudally polarized hair cells of

neuromasts at 1.5, 2, 3, and 4 days post fertilization (dpf). The average Sema7A intensity

increased significantly over this period in both rostrally and caudally polarized hair cells. At

1.5 dpf, the neuromasts harbor primarily immature hair cells that make minimal contacts with

sensory axons and do not form stable association with axonal terminals (23). At this stage the

average Sema7A intensity in hair cells remained low. By 2-4 dpf the hair cells mature to form

stable contacts and well-defined synapses with the sensory axons (5,24). The average

Sema7A intensity in hair cells at each of these stages rose to levels that significantly exceeded

those of 1.5 dpf neuromasts (Figure 1F). This rise in the amount of Sema7A during neuromast

maturation supports a role for Sema7A in the guidance of sensory axons and their interaction

with hair cells.

We additionally demonstrated an anisotropic distribution of Sema7A along the

apicobasal axis of each hair cell. As neuromasts develop through 2, 3, and 4 dpf, Sema7A

remained highly enriched in the subapical region of the hair cell, the site of the Golgi network

(Figure 1E; Figure 1—figure supplement 2E). This localization suggests that Sema7A—like

other proteins of the semaphorin family (25)—is directed to the plasmalemma through Golgi-

mediated vesicular trafficking. Using the transgenic line *Tg(cldnb:LY-mScarlet)* that expresses

membrane-tethered mScarlet in all cells of the neuromast (26), we detected Sema7A at the

plasma membrane and potentially in vesicles (Figure 1—figure supplement 2F-I). While

Sema7A maintained a high level in the subapical region, the protein increasingly accumulated

at the base of the hair cell: the Sema7A level at the hair cell base was low at 2 dpf, but

increased sharply by 3 dpf and 4 dpf (Figure 1G; Figure 1—figure supplement 2J). During this

period the hair cells increase in number and mature, whereas the associated sensory axons

arborize to contact multiple hair cells and form robust synaptic boutons at their bases (5,24).

Does progressive basal accumulation of Sema7A facilitate association of sensory axon

terminals to the hair cell? To address this question, we utilized doubly labeled transgenic fish

Tg(myo6b:actb1-EGFP;neurod1:tdTomato) that marked the hair cells and the sensory axons

of the posterior lateral line with distinct fluorophores (Figure 1A) (27). We quantified the average

Sema7A intensity and the accumulation of sensory arbors, measured by td-Tomato intensity,

at the bases of the hair cells. Across developmental stages, the basal association of the

sensory arbors showed a positive correlation with Sema7A accumulation (Figure 1H,I;

Figure 1—figure supplement 3A-E, Video 1). The progressive enrichment of Sema7A at the

hair cell base with the sensory axons suggests that Sema7A acts as a mediator of contacts

between sensory axons and hair cells.

Sema7A restricts sensory axons around clustered hair cells. If Sema7A guides and

restricts the sensory arbors at and around the hair cells clustered in a neuromast, then

inactivating Sema7A function should disrupt this process. To test this hypothesis, we obtained

a *sema7a* mutant allele (*sema7a^{sa24691}*) with a point mutation that introduces a premature stop

codon in the conserved sema domain (Figure 2—figure supplement 1A). The homozygous

mutant, hereafter designated *sema7a^{-/-}*, should lack both the secreted and the membrane-

attached forms of Sema7A. The average intensity of Sema7A immunolabeling in hair cells

diminished by 61% in *sema7a^{-/-}* larvae (0.34 ± 0.01 arbitrary units) in comparison to controls

(0.87 ± 0.01 arbitrary units). In a few cases we observed minute accumulations of Sema7A at

the subapical region, but little or no protein at the base of the mutant hair cells (Figure 2A-E).

We quantified the number of hair cells in the neuromasts of the control and the *sema7a^{-/-}*

larvae. We did not observe significant changes in the hair cell number between the genotypes

across developmental stages (Figure 2—figure supplement 1B). Furthermore, we tested the

mechanotransduction ability of *sema7a^{-/-}* mutant hair cells by labeling with the styryl dye FM

4-64 (5). The hair cells of the 4 dpf *sema7a^{-/-}* larvae incorporated the FM 4-64 as effectively as

the age-matched control animals. We quantified the FM 4-64 intensities from 60 hair cells and

six neuromasts from three control larvae, and from 60 hair cells and six neuromasts from three

sema7a^{-/-} mutant larvae. The average intensities of FM 4-64 labeling between the genotypes

did not show a significant difference, which suggests that the mutation does not impact

essential hair cell functions (Figure 2—figure supplement 1C,D; Videos 2,3).

To characterize the impact of the *sema7a^{-/-}* mutation on arbor patterning, we utilized

doubly labeled transgenic fish *Tg(myo6b:actb1-EGFP;neurod1:tdTomato)* that distinctly

labeled the hair cells and the sensory arbors (27). In control neuromasts, the sensory axons

approached, arborized, and consolidated around the bases of clustered hair cells (Figure 2F).

Although the sensory axons approached the hair cells of *sema7a^{-/-}* neuromasts, they displayed

aberrant arborization patterns with wayward projections that extended transversely to the organ

(Figure 2G). To quantitatively analyze the arborization morphology, we traced the sensory

arbors in three dimensions to generate skeletonized network traces that depict–as

pseudocolored trajectories–the increase in arbor length from the point of arborization

(Figure 2H, I).

To visualize the hair cell clusters and associated arborization networks from multiple

neuromasts, we aligned images of hair cell clusters, registered them at their hair bundles, from

neuromasts at 2 dpf, 3 dpf, and 4 dpf (Figure 2—figure supplement 1E,F). Throughout

development, the arborization networks of control neuromasts largely remained within the

contours of the hair cell clusters; the few that reached farther nonetheless lingered nearby

(Figure 2J; Figure 2—figure supplement 1G,I). This result suggests that an attractive cue

retained the axons near the sensory organ. In *sema7a*$^{-/-}$ neuromasts, however the neuronal

arbors failed to consolidate within the boundaries of hair cell clusters and extended far beyond

them (Figure 2K; Figure 2—figure supplement 1H,J; Video 4).

We quantified the distribution of the sensory arbors around the center of the combined

hair cell clusters. For both the control and the *sema7a*$^{-/-}$ neuromasts, the arbor densities peaked

proximal to the boundaries of the hair cell clusters. Beyond the cluster boundaries, in control

neuromasts at 2 dpf, 3 dpf, and 4 dpf the arbor densities fell sharply at respectively 31.7 *µ*m,

36.6 *µ*m, and 32.7 *µ*m from the center. In *sema7a*$^{-/-}$ neuromasts of the same ages, the sensory

arbors extended respectively 38.5 *µ*m, 51.4 *µ*m, and 56.6 *µ*m from the center (Figure 2L;

Figure 2—figure supplement 1K,L; Figure 2—figure supplement 2A). Furthermore, we

quantified the degree of contact of the sensory arbors to their hair cell clusters in individual

neuromasts from both control and *sema7a*$^{-/-}$ mutants. In control neuromasts, the degree of

contact was 83 ± 1 % (mean ± SEM) at 2 dpf and 84 ± 1 % at 3 dpf, but significantly increased

to 90 ± 1 % at 4 dpf (Figure 2M; Figure 2—figure supplement 2B). This observation indicates

that the sensory arbors reinforce their association to the hair cell clusters as neuromasts

develop. However, in *sema7a*$^{-/-}$ neuromasts such contact was significantly reduced. Only

69 ± 4 % of sensory arbors at 2 dpf were closely associated with the corresponding hair cell

clusters, a value that remained at 67 ± 2 % at both 3 dpf and 4 dpf (Figure 2M). These findings

signify that, with simultaneous disruption of both signaling modalities of Sema7A, the organ

fails to restrict the localization of sensory arbors and has significantly diminished contact with

them.

Sema7A patterns the arborization network's topology. Accurate spatial arrangement of the

neuronal projections around hair cells is essential for proper retrieval and transmission of

sensory information. Mutations that perturb proper assembly of sensory neuronal microcircuits

lead to hearing impairment (28). Quantitative insight into the wiring pattern is therefore essential

to identify systematic aberrations in the microcircuit arising from genetic deficits. The three-

dimensional skeletonized network traces allowed us to accurately delineate the microcircuit

topology across individuals, developmental stages, and genotypes from living animals (Figure

2D, E). To quantitatively analyze the arbor network, we defined five topological attributes: (1)

nodes, denoted as junctions where arbors branch, converge, or cross; (2) loops, comprising

minimal arbor cycles forming topological holes; (3) bare terminals, defined as arbors with free

ends; (4) the node degree, calculated as the number of first-neighbor nodes incident to a node;

and (5) the arbor curvature, measured as the total curvature of a bare terminal (Figure 3A). We

quantified these attributes to accurately determine how the circuit topology is established

during maturation of the sensory organ in control larvae and how it is impacted by *sema7a*

deficit.

To determine how internode connectivity is distributed across networks, we quantified

the distribution of node degrees along the relative height of the hair cell cluster across

individuals, developmental stages, and genotypes. As for most other real-world networks (29),

we observed a highly skewed node-degree distribution: the majority of nodes had low node

degrees (three and four) and a small number of nodes—often acting as hubs, like the

arborization initiation point—had high node degrees (five to seven). The node degree varied

from three to seven in control larvae and three to six in *sema7a*$^{-/-}$ larvae. Across

developmental stages and genotypes, the high node degrees primarily occurred at the base

(region from 0 % to 10 % of the cluster height) of the hair cell cluster and acted as hubs for

the majority of branching events. Low node degrees primarily occurred along the basolateral

region (region from 10 % to 60 % of the cluster height) of the hair cell cluster. The apical region

(region from 60 % to 100 % of the cluster height) was sparsely populated with low node

degrees. The network's hierarchical organization along the organ's axis suggests that arbors

receive sensory inputs at multiple levels, allowing the network to carry out robust neural

computations. Even though nodes of low degree were most prevalent between genotypes

across developmental stages, their distribution along the relative organ axis became

significantly different at 4 dpf, with nodes in control larvae localized higher along the axis than

in $sema7a^{-/-}$ larvae (Figure 3A; Figure 3—figure supplement 1A-C; Table 1). These

observations indicate that Sema7A regulates both the spatial organization and node-degree

complexity of the sensory arbor network.

An average of 9.3 sensory axonal branches approach the hair cell cluster of a neuromast

(8). Pathfinding of sensory axons around the hair cell cluster results in the formation of axonal

fascicles that form numerous loops and bare terminals, which determine the microcircuit

topology (Figure 2B, D) (8). To optimally surround and contact the hair cell cluster, do the

sensory arbors prefer to adopt one topological feature over the other? To address this

question, we measured the correlation between the number of bare terminals and the number

of loops across individuals, developmental stages, and genotypes. In both 2 dpf and 3 dpf

larvae, the distributions remained highly scattered across individuals and genotypes.

However, by 4 dpf in the control larvae the arbor network contained either a high number of

loops with a low number of bare terminals or a high number of bare terminals with a low

number of loops. This result suggests that the stereotypical circuit topology observed in the

mature organ may emerge through transition of individual arbors from forming bare terminals

to forming closed loops encircling topological holes. That we did not observe a similar

transition in $sema7a^{-/-}$ neuromasts of the same age suggests that Sema7A may direct the

switch between these two distinct topological features (Figure 3B; Figure 3—figure

supplement 1D, E; Table 1).

Measurement of the contour length of both bare terminals and loops additionally

revealed that across developmental stages in control larvae the sensory arbors followed a

three-step process to choose the appropriate topological feature: arbors shorter than 10 μm

always formed bare terminals; arbors between 10 μm and 30 μm began to switch between

features, with arbors longer than 20 μm strongly preferring to form loops; and arbors longer

than 30 μm always formed loops. These results imply that the switch between bare terminals

and loops is partly dependent on the length of the arbors. However, this characteristic

topological preference was significantly perturbed in the *sema7a$^{-/-}$* larvae across

developmental stages, as many of the sensory arbors longer than 30 μm remained as bare

terminals (Figure 3C; Figure 3—figure supplement 1F,G; Table 1). Furthermore, in 4 dpf

control neuromasts the majority of the loops contained four nodes and utilized 30-40 μm of

arbors to form closed cycles. These conserved topological attributes were significantly

perturbed in the *sema7a$^{-/-}$* larvae of same age (Figure 3D, E; Figure 3—figure supplement

2A-D; Table 1). These findings indicate that Sema7A tightly monitors the sensory arbor loop

formation, including the preferred loop length and number of nodes per loop, to properly

surround and contact the hair cell cluster.

As observed previously, bare terminals in control neuromasts project transverse to the

organ and linger nearby, often curving back to form new loops and contact the hair cell cluster

(Figure 2D,F). To analyze the propensity of the bare terminals to curl backward, we measured

their total curvatures, defined as the sum of the curvature at each point along the length of an

arbor. Henceforth, we address the total curvature as the curvature of a bare terminal. We

classified the bare terminals in two categories: high-curvature arbors with curled morphology

and curvature greater than π/6 (30 degrees); and low-curvature arbors with linear morphology

and curvature less than π /6. In control neuromasts 44.8 % of arbors had low curvature

compared to 56.4 % in *sema7a$^{-/-}$* neuromasts at 4 dpf. Concomitantly, 55.2 % of arbors had

high curvature in control neuromasts compared to 43.6 % in *sema7a$^{-/-}$* neuromasts at 4 dpf

(Figure 3F; Figure 3—figure supplement 2E, F; Table 1). These observations further confirm

our hypothesis that a Sema7A-mediated attractive cue is involved in orienting bare neural

arbors toward the hair cell cluster to potentially form loops.

In summary, we propose that Sema7A mediates four key aspects of the topology of the

arborization network: (1) the node degree complexity along the organ's axis; (2) the transition

of arbors from bare terminals to loops; (3) the preference for forming loops containing four

nodes with 30-40 μm of arbors; and (4) the preference of bare terminals to adopt higher

curvature.

Sema7A^{sec} is a sufficient chemoattractive cue for sensory axon guidance. Because

genetic inactivation of the *sema7a* gene disrupts both signaling modalities of the cognate

protein, we could not assess the specific activity of the Sema7A^{sec} diffusive cue in guiding

lateral-line sensory axons. To independently verify the potential role of that isoform, we

therefore expressed the secreted variant ectopically and investigated the resultant arborization

patterns.

Analysis of the microcircuit connectivity of the neuromasts has demonstrated bare

sensory-axonal terminals in the perisynaptic compartments, where they do not contact the hair

cell membrane (8). We speculate that these bare terminals are attracted by a Sema7A^{sec}

diffusive cue. To test this hypothesis, we ectopically expressed Sema7A^{sec} protein tagged with

the fluorophore mKate2, then analyzed its effect on the morphology of sensory arbors.

Fertilized one-cell embryos were injected with the *hsp70:sema7a^{sec}-mKate2* plasmid and

raised to 3 dpf. Larvae expressing the transgenesis marker—the lens-specific red fluorescent

protein mCherry—were heat-shocked, incubated to allow expression, and subsequently

mounted for live imaging. In these larvae, a random mosaic of cells—often embryonic muscle

progenitors in the dermomyotome or mature myofibers—expressed the exogenous

Sema7A^{sec}-mKate2 protein. We selected only those neuromasts in which a single mosaic

ectopic integration had occurred near the network of sensory arbors (Figure 4A,B). In all 22

such cases, we observed robust axonal projections from the sensory arbor network toward the

ectopically expressing Sema7A^{sec} cells. When the exogenous *sema7a^{sec}* integration occurred

in embryonic muscle progenitor cells, which reside in a superficial layer external to the muscle

fibers and adjacent to the larval skin (30), the projections were able to form direct contacts with

them (Figure 4C; Videos 5,6). But more often, when mature myofibers expressed the

exogenous Sema7A^{sec} deep in the myotome, the axonal projections approached those targets

but failed to contact them (Figure 4D, Videos 7,8). This behavior likely arose because other

components in the myotomal niche—including the fibrous epimysium, other myofibers,

myoblasts, and immature myotubes whose surroundings are permeable to diverse diffusive

cues (31)—physically obstructed direct interaction of the extended neurites with the ectopically

expressing myofibers. All 18 of the injected but not heat-shocked control larvae did not express

ectopic Sema7A^{sec}, and we did not observe aberrant projections from the sensory arbor

network (Figure 4E).

To measure the accuracy of the extended axonal projections in finding the ectopic

Sema7A^{sec} source, we defined three parameters: (1) the source path, denoted as the linear

distance from the point of arborization to the boundary of an ectopically expressing Sema7A^{sec}

cell; (2) the projection path, representing the linear distance from the point of arborization to

the terminus of the extended projection; and (3) the projection proximity, calculated as the

linear distance from the terminus of the extended axonal projection to the nearest boundary of

the ectopically expressing Sema7A^{sec} cell (Figure 4B). We quantified these parameters from

18 of the 22 mosaic integration events; in the remainder the extended axonal projections–

although following the ectopic source–reentered the posterior lateral-line nerve so that

identification of the axon terminals was not possible (Figure 4—figure supplement 1A-A";

Video 9). The projection paths of the extended axons reached from 27.1 μm to 142.5 μm while

following the ectopic Sema7A^{sec} cues. When we plotted the projection path length against the

source-path length, we observed a nearly perfect correlation (Figure 4F). This result signifies

that irrespective of the location of the ectopic source–whether proximal or distal to the sensory

arbor network–the Sema7A^{sec} cue sufficed to attract axonal terminals from the sensory arbor.

Furthermore, the low average projection-proximity length of $3.5 \pm 0.7 \mu$m (mean ± SEM)

indicated that the extended projection terminals remained either in contact (8 of 18) or in the

vicinity (10 of 18) of the ectopic Sema7A^{sec} sources (Figure 4G). These observations together

demonstrate that the Sema7A^{sec} diffusive cue is sufficient to provide neural guidance *in vivo*.

As the posterior lateral-line ganglion matures, newly formed neurons extend their axonal

growth cones along the posterior lateral-line nerve to reach the neuromasts (32). We wondered

whether the axonal projections that had yet to be associated with a neuromast could also

respond to an ectopic source of Sema7A^{sec}. Indeed, on seven occasions in which *sema7a^{sec}*

integration had occurred between two neuromasts, we detected neurite extensions from the

posterior lateral-line nerve to the ectopic Sema7A^{sec} sources (Figure 4H; Figure 4—figure

supplement 1B-B", Videos 10-12). Live time-lapse video microscopy further revealed dynamic

association between the extending neurites and the ectopic Sema7A^{sec} sources. Long and thin

along their entire extents, the neurites were highly motile, extending and retracting rapidly near

the Sema7A^{sec} expressing cells. Many neurites maintained their contact with the ectopic

sources of Sema7A^{sec} from one hour to almost eight hours (Figure 4I,J; Figure 4—figure

supplement 2A, Videos 13-16). The responsiveness of the posterior lateral-line axons to

Sema7A^{sec} might therefore be an intrinsic property of the neurons that does not require the

neuromast microenvironment.

Sema7A-GPI fails to impart sensory axon guidance from a distance. We speculate that

the GPI-anchored transcript variant of the *sema7a* gene can function only through contact-

dependent pathways. To test this hypothesis, we expressed the Sema7A-GPI ectopically and

investigated the resultant arborization patterns of the sensory axons. Fertilized one-cell

embryos were injected with the *hsp70:sema7a-GPI-2A-mCherry* plasmid and raised to 3 dpf.

Larvae expressing the transgenesis marker—the lens-specific green fluorescent protein

(GFP)—were heat-shocked, incubated to allow expression, and subsequently mounted for live

imaging (Figure 5A). In these larvae, a few mature myofibers expressed the exogenous

Sema7A-GPI protein. We selected only those neuromasts in which an ectopic integration had

occurred near the network of sensory arbors. In all 15 such cases, no sensory axon left the

arbor network to approach the ectopic Sema7A-GPI sources. In all 17 of the uninjected and

heat-shocked control larvae, the sensory arbor network did not show any perturbation

(Figure 5B-C, Videos 17-19). These data suggest that the Sema7A-GPI, unlike its secreted

counterpart, functions at the surfaces of the expressing cells.

Sema7A deficiency impacts synaptic assembly. Contact stabilization between an axon and

its target cell is essential for synapse formation (33). Neural GPI-anchored proteins can act as

regulators of various synaptic-adhesion pathways (34). In the mouse olfactory system, GPI-

anchored Sema7A is enriched in olfactory sensory axons and mediates sustained interaction

with the dendrites of mitral and tufted cells in the olfactory bulb to establish synapses (15). In

the developing mouse brain–particularly in Purkinje cells–GPI-anchored Sema7A instead

regulates the elimination of synapses onto climbing fibers (35). The role of anchored Sema7A

in regulating synaptic architecture can thus vary according to the cell type and developmental

stage. Our demonstration that Sema7A is necessary to consolidate the contact between

sensory axons and hair cells during neuromast development suggests a possible role in

regulating synaptic structure.

Proper synaptogenesis requires the correct spatial organization of the presynaptic and

postsynaptic apparatus at the apposition of two cells. At the interface between hair cells and

sensory axons, the synaptic network involves two scaffolding proteins in particular: ribeye, a

major constituent of the presynaptic ribbons (36), and membrane-associated guanylate kinase

(MAGUK), a conserved group of proteins that organize postsynaptic densities (37). Utilizing

these scaffolding proteins as markers, we investigated the impact of the $sema7a^{-/-}$ mutation on

synapse formation. To analyze the distribution of presynaptic densities–immunolabeled with

an antibody against the ribeye-associated presynaptic constituent C-terminal binding protein

(CTBP) (36)–we counted the number and measured the area of CTBP punctae from multiple

neuromasts at several stages of neuromast development (Figure 6A,B). The CTBP punctae in

each hair cell ranged from zero to seven across developmental stages. In the zebrafish's

posterior lateral line, the formation of presynaptic ribbons is an intrinsic property of the hair cells

that does not require contact with lateral-line sensory axons. However, sustained contact with

the sensory axon influences the maintenance and stability of ribbons (38). As neuromasts

matured from 2 dpf to 4 dpf, we identified similar distribution patterns with a characteristic

increase in the number of CTBP punctae in both control and $sema7a^{-/-}$ larvae (Figure 6C,D).

This result implies that the formation of new presynaptic ribbons is not perturbed in the mutants,

even though there are on average fewer CTBP punctae (Figure 6—figure supplement 1A-C).

Measurement of the areas of CTBP punctae also showed similar distributions in both

genotypes (Figure 6E,F) but the average area of presynaptic densities was reduced in the

mutants (Figure 6—figure supplement 1D-F, Figure 6—figure supplement 2). Because GPI-

anchored Sema7A lacks a cytosolic domain, it is unlikely that Sema7A signaling directly

induces the formation of presynaptic ribbons. We propose that the decrease in average number

and area of CTBP punctae likely reflects decreased stability of the presynaptic ribbons owing

to lack of contact between the sensory axons and the hair cells.

Our analysis of the skeletonized network traces of the sensory arbor network identified

distinct topological features that are resultant of collective assemblies of each sensory axon.

Because postsynaptic aggregates are organized at the sensory axons, we wondered whether

they are associated with a specific topological feature of the network. To identify the distribution

of the postsynaptic aggregates along the arbor network, we utilized the transgenic animal

TgBAC(neurod1:EGFP) that labeled the sensory axons of the posterior lateral line in green

fluorescence protein (GFP), and marked the postsynaptic aggregates through immunostaining

with a pan-MAGUK antibody (Figure 6G). We generated skeletonized network traces of the

arbor network and determined the three-dimensional spatial locations of the MAGUK punctae

(Figure 6—figure supplement 2). Comparison of the topological features of the network with

the spatial locations of the MAGUK punctae revealed that the postsynaptic densities occur

more extensively on the loops than on the bare terminals (Figure 6G). We measured 567

MAGUK punctae from 16 sensory arbor networks, where 531 punctae were located on the

loops and only 36 punctae on the bare terminals (Figure 6H). Furthermore, the MAGUK

punctae predominantly aggregate near the node and their density sharply diminishes moving

farther from the node (Figure 6I, Figure 6—figure supplement 2). These observations suggest

that the distribution of the postsynaptic aggregates is closely associated with the topological

attributes of the sensory circuit. Because in the *sema7a^{-/-}* mutants the arbor networks have

significantly lower number of loops and nodes, and more bare terminals, we hypothesize that

in the mutants the occurrence of postsynaptic aggregates is perturbed. Moreover, Sema7A-

GPI-mediated juxtracrine signaling onto the apposed axonal membrane is essential to induce

postsynaptic assembly (15).

To characterize the impact of *sema7a* mutation on the formation of postsynaptic
aggregates, we counted the number and measured the area of MAGUK punctae from multiple
neuromasts at several stages of neuromast development (Figure 6J,K). At 4 dpf, 68.3 % of the
hair cells in the control neuromasts were associated with at least one MAGUK punctum. In the
sema7a$^{-/-}$ mutant this value fell to 37.5 % (Figure 6L,M). The average number of MAGUK
punctae also declined in the mutants (Figure 6—figure supplement 1G-I). As control
neuromasts matured from 2 dpf to 4 dpf, the MAGUK punctae showed a shift toward smaller
sizes, suggesting that the postsynaptic structure fragmented into smaller entities as observed
in other excitatory synapses (40). At 4 dpf, 77.5 % of the hair cells in the control neuromasts
had postsynaptic densities with areas between 0.10 μm^2 and 0.40 μm^2. In the *sema7a$^{-/-}$* mutant
the corresponding value was only 42.5 % (Figure 6N,O). The average area of MAGUK punctae
was also reduced in the mutants compared to the controls (Figure 6—figure supplement 1J-L,
Figure 6—figure supplement 2**).** We speculate that the abnormalities in the postsynaptic
structure of the *sema7a$^{-/-}$* mutants arose from reduced contact between the hair cells and the
sensory axons and perturbed topological attributes of the sensory circuit or from the lack of
Sema7A-GPI-mediated juxtracrine signaling onto the apposed neurite terminals (15).

Discussion

The specification of neural circuitry requires precise control of the guidance cues that direct
and restrict neural arbors at their target fields and facilitate synapse formation. In the lateral-
line neuromast of the zebrafish, we have identified Sema7A as an essential cue that regulates
the interaction between hair cells and sensory axons. We have discovered two isoforms of the
sema7a gene *in vivo*: the chemoattractive diffusible form is sufficient to guide the sensory
arbors toward their target, whereas the membrane-attached form likely participates in sculpting
accurate neural circuitry to facilitate contact-mediated formation and maintenance of synapses.
Our results suggest a potential mechanism for hair cell innervation in which a local Sema7A^{sec}
diffusive cue likely consolidates the sensory arbors at the hair cell cluster and the membrane-
anchored Sema7A-GPI molecule guides microcircuit topology and synapse assembly.

Establishing the exact function of each of the *sema7a* isoforms will necessitate studies that

include transcript-specific targeted mutagenesis and rescue of the *sema7a^{-/-}* mutant phenotype

by each of the transcript variants.

The role of Sema7A in regulating diverse topological features of the arborization network

suggests that microcircuits of control and *sema7a^{-/-}* neuromasts have important functional

differences. Loss of Sema7A disrupts the hierarchical organization of the node-degree

distribution along the organ's axis, as well as the preference towards forming loops containing

four nodes with 30-40 μm of arbors. What benefit does the network gain from surrounding the

hair cell cluster with closed loops instead of bare terminals? The network may actively form

loops to contribute to the neuromast's overall structure and function, or else emerge as a

consequence of the arbors navigating the intervening spaces between the hair cells and

support cells. The preferential localization of the postsynaptic aggregates in the loops and an

enrichment near the nodes of the arbor network suggest that the precise patterning of the

network—with characteristic looping behaviors—is not merely a byproduct of neuromast

architecture, but may contribute to the network's ability to gather sensory inputs through

properly placed postsynaptic aggregates and to faithfully transmit sensory information to higher

processing centers in the hindbrain (7). Further studies are necessary to elucidate the potential

role of these topological features in regulating the distribution of synaptic bodies and optimizing

robust neural transmission in the lateral-line microcircuit.

Recent studies have begun to dissect the molecular mechanisms of semaphorin

signaling in axonal navigation. In explants derived from the murine olfactory bulb, Sema7A

interacts directly with integrinβ1 to trigger downstream signaling effectors involving focal

adhesion kinase and mitogen-activated protein kinases, which are critical regulators of the

cytoskeletal network during axonal pathfinding (4,41). Moreover, the interaction between

Sema7A and its receptor plexinC1 activates the rac1-cdc42-PAK pathway, which alters the

actin network to promote axonal guidance and synapse formation (15,42). We speculate that

lateral-line sensory axons utilize similar mechanisms to sense and respond to Sema7A in the

establishment of microcircuits with hair cells.

During early development of the posterior lateral line, the growth cones of sensory axons

intimately associate with the prosensory primordium that deposits neuromasts along the lateral

surface of the larva (43,44). The growth of the sensory fibers with the primordium is regulated

by diverse neurotrophic factors expressed in the primordium (45,46). Impairing brain-derived

or glial cell line-derived neurotrophic factors and their receptors perturbs directed axonal

motility and innervation of hair cells (45,46). Lateral-line hair cells are particularly enriched in

brain-derived neurotrophic factor (47) and other chemotropic cues (Figure 7). Because the

sensory axons of *sema7a$^{-/-}$* mutants reliably branch from the lateral-line nerve to reach hair cell

clusters, these chemoattractive factors might compensate in part for the absence of Sema7A

function. Although sensory axons are strict selectors of hair cell polarity and form synapses

accordingly (5), diverse chemoattractant cues including Sema7A occur uniformly between hair

cell polarities (47). It seems likely that unbiased attractive signals bring sensory arbors toward

hair cells, after which polarity-specific molecules on hair cell surfaces dictate the final

association between axonal terminal and hair cells. Notch-delta signaling and its downstream

effectors regulate hair cell polarity and the innervation pattern (6,8,27). It would be interesting

to determine whether Notch-mediated polarity signatures on the hair cell's surface can refine

the neuromast neural circuitry.

Materials and Methods

Zebrafish strains and husbandry. Experiments were performed on 1.5-4 dpf zebrafish larvae

in accordance with the standards of Rockefeller University's Institutional Animal Care and Use

Committee. Naturally spawned and fertilized eggs were collected, cleaned, staged, and

maintained at 28.5 °C in system-water with 1 µg/mL methylene blue. Wild-type AB and

heterozygous mutant *semaphorin7a^{sa24691}* zebrafish were obtained from the Zebrafish

International Resource Center. In addition, the following transgenic zebrafish lines were used:

Tg(myo6b:actb1-EGFP) (48), *Tg(neurod1:tdTomato)* (27), *Tg(cldnb:LY-mScarlet)* (26), and

TgBAC(neurod1:EGFP) (39).

Embryo dissociation and flow cytometry. Immediately before dissection, 4-day-old

Tg(myo6b:actb1-EGFP) and wild-type larvae were anesthetized in 600 μM 3-aminobenzoic

acid ethyl ester methanesulfonate in system water with 1 μg/mL methylene blue. The anterior

half of each larva was removed with a fine surgical blade. We used approximately 400 GFP-

positive and 50 GFP-negative larvae, the latter for controls of the gating procedure.

Dissociation of the tails were performed using a protocol (22) with minor modification; detailed

protocol is available upon request. The resuspended cells were incubated for 5 min on ice and

sorted with an influx cell sorter (BD Biosciences, San Jose, CA, USA) with an 85 μm nozzle at

20 psi and 1X DPBS (Sigma-Aldrich, St Louis, USA). Laser excitation was conducted at

561 nm and 488 nm. Distinct populations of GFP-positive hair cells were isolated based on

forward scattering, lateral scattering, and the intensity of GFP fluorescence.

RNA extraction and cDNA library preparation. Sorted cells were collected in a lysis-buffer

solution supplemented with β-mercaptoethanol and RNA extraction was performed

immediately using a standard protocol (RNeasy Plus Micro Kit; Qiagen). The yield and quality

of the product were measured with a bioanalyzer (Agilent 2100). Only samples with an RNA

integrity score (RIN) greater than 8.0 were selected for the preparation of cDNA libraries using

iScript cDNA Synthesis Kit (BioRad). One nanogram of total RNA was amplified to several

micrograms of cDNA.

Reverse transcription-polymerase chain reactions (RT-PCR). The *sema7a* gene

expresses two transcript variants: variant 1, *sema7a-GPI* anchored, NM_001328508.1; and

variant 2, *sema7a^{sec}* secreted, NM_001114885.2. The variant 2 lacks several C-terminal

exons, and its 3'-terminal exon extends past a splice site that is used in variant 1. The encoded

variant 2 has a shorter and distinct C-terminus than variant 1. To individually identity these

transcripts, each of them was PCR amplified from the cDNA library using the following primers:

For variant 1, sema7aF= 5′-GGTTTTTCTGAGGCCATTCC-3′, sema7aR1=

5′-GGCACTCGTGACAAATGCTA-3′; and for variant 2, sema7aF=

5′-GGTTTTTCTGAGGCCATTCC-3′, sema7aR2= 5′-TGTGGAGAAAGTCACAAAGCA-3′. To

detect *pvalb8* transcript we used the following primers: pvalb8F=
5´-ATGTCTCTCACATCTATCCT-3´ and pvalb8R= 5´-TCAGGACAGCACCATCGCCT-3´. To
detect *s100t* transcript we used the following primers: s100tF=
5´-ATGCAGCTGATGATCCAGAC-3´ and s100tR= 5´-CTATTTCTTGTCATCCTTCT-3´.

Immunofluorescence microscopy. For the immunofluorescence labeling of wholemounted
wild-type control and *sema7a⁻/⁻* larvae, 1.5-4 dpf larvae were fixed overnight at 4 °C in 4 %
formaldehyde in phosphate-buffered saline solution (PBS) with 1 % Tween-20 (1 % PBST).
Larvae were washed four times with 1 % PBST for 15 min each, then placed for 2 hr in blocking
solution containing PBS, 2 % normal donkey serum (NDS), 0.5 % Tween-20, and 1 % BSA.
Primary antibodies were diluted in fresh blocking solution and incubated with the specimens
overnight at 4 °C. The primary antibodies were goat anti-Sema7A (1:200; AF1835, R&D
Systems) (Figure 1—figure supplement 4) (14), rabbit anti-myosin VI (1:200; Proteus 25-6791),
murine anti-GM130 (1:200; 610822, BD Transduction Laboratories), murine anti-CTBP (1:200;
B-3, SC-55502, Santa Cruz Biotech), and rabbit anti-mCherry (1:200; GTX128508, GeneTex).
Larvae were washed four times with 0.1% PBST for 15 min each. Alexa Fluor-conjugated
secondary antibodies (Invitrogen, Molecular Probes) applied overnight at 1:200 dilutions in
0.2 % PBST included anti-rabbit (488 nm), anti-goat (555 nm), and anti-mouse (647 nm).
Larvae were then washed four times in 0.2 % PBST for 15 min each and stored at 4 °C in
VectaShield (Vector Laboratories).

For the immunofluorescence labeling of postsynaptic density in wholemounted wild-type
control and *sema7a⁻/⁻* larvae, 2-4 dpf larvae were fixed with 4 % formaldehyde in PBS for
4.5-6 hr at 4 °C. Larvae were washed five times with 0.01 % PBST for 5 min each and then
rinsed in distilled water for 5 min. The larvae were permeabilized with ice-cold acetone at -20 °C
for 5 min. They were rinsed in distilled water for 5 min, washed five times with 0.01 % PBST
for 5 min each, and blocked overnight with PBS buffer containing 2 % goat serum and 1 %
BSA. Murine anti-pan-MAGUK antibody (1:500; IgG1 NeuroMab, K28.86, 75-029 and
MABN72, Millipore) was diluted in PBS containing 1 % BSA, added to the larvae, and incubated
for 3 hr at room temperature. Larvae were washed five times for 5 min each with 0.01 % PBST.

Alexa Fluor 555-conjugated anti-mouse secondary antibody (1:1000) diluted in PBS containing

1 % BSA was added to the larvae and incubated for 2 hr at room temperature. Larvae were

washed five times with 0.01 % PBST for 5 min each, rinsed in distilled water for 5 min, and

stored at 4 °C in VectaShield (Vector Laboratories).

Posterior segments of fixed larvae were mounted on a glass slide and imaged on an

Olympus IX81 microscope with a microlens-based, super-resolution confocal microscope

(VT-iSIM, VisiTech international) under 60X and 100X, silicone-oil objective lenses of

numerical apertures 1.30 and 1.35, respectively. Static images at successive focal depths were

captured at 200 nm intervals and deconvolved with the Microvolution software in ImageJ.

Measurement of Sema7A and td-Tomato-positive sensory arbor intensities. The average

Sema7A fluorescence intensities of the hair cells from 1.5 dpf, 2 dpf, 3 dpf, and 4 dpf

neuromasts were measured by determining the mean gray level within each hair cell labeled

by actin-GFP expression. The intensity distribution of Sema7A protein across the basal cell

membrane was measured using the line profile tool in ImageJ (Figure 1—figure

supplement 2G-I). The lines were drawn with 1 μm length, starting from the inside of the hair

cell to the outside. For each hair cell membrane, the intensity values were scaled from 0 to 1

arbitrary units. The intensity distribution of Sema7A protein along the apicobasal axis of the

hair cell was measured using the line profile tool in ImageJ (Figure 1—figure supplement 2J).

Because hair cell lengths differed among samples, lines were drawn with 2 μm widths and

varying lengths of 6-8 μm. For each hair cell, apicobasal length was scaled from 0 to 100

arbitrary units and the intensity values from 0 to 1 arbitrary units. The average Sema7A and td-

Tomato-positive sensory arbor fluorescence intensities at the hair cell bases from 2 dpf, 3 dpf,

and 4 dpf neuromasts were measured by determining the mean gray level within each hair cell

basal area, denoted by the region underneath the hair cell nucleus (Figure 1—figure

supplement 3A-E).

Genotyping of mutant fish. The *semaphorin7a*[sa24691] allele harbors an A-to-T point mutation

in the seventh exon of the *semaphorin7a* gene, which creates a premature stop codon (Figure

2—figure supplement 1A). Genomic DNA was isolated from the tail fins of adult

semaphorin7a[sa24691] (*sema7a[sa91]*) heterozygous mutant zebrafish. The mutant locus in the

genome was PCR amplified using the following primers: sema7a[sa91]F:

5´-AAAGCTGGAAAGCGAATCAA-3´ and sema7a[sa91]R: 5´-ATATCCAAGGATCCGCCTCT-3´.

The 734-base pair amplicons were sequenced, and heterozygous adults were propagated.

Lipophilic styryl fluophore labeling of hair cells. Freely swimming 4 dpf control and

sema7a[-/-] larvae were exposed to the styryl fluorophore FM 4-64

(N-(3-triethylammoniumpropyl)-4-(6-(4-(diethylamino) phenyl) hexatrienyl) pyridinium

dibromide, ThermoFischer Scientific) at 5 μM for 30 s, then rinsed three times in fresh embryo

water and immediately mounted in low-melting-point agarose for live imaging. The average

FM 4-64 intensities within the hair cells of 4 dpf neuromasts of both genotypes were measured

by determining the mean gray level within each hair cell labeled by actin-GFP expression.

Transient transgenesis with the hsp70I:sema7a[sec]-mKate2 construct. The pDNR-LIB

plasmid containing the *sema7a[sec]* coding sequence was purchased from horizon (7140389,

Perkin-Elmer). The coding sequence was PCR amplified using the following primers:

sema7a[sec]F:

5´-GGGGACAAGTTTGTACAAAAAAGCAGGCTTGATGATTCGACATTATTCT-3´   and

sema7a[sec]R:

5´-GGGGACCACTTTGTACAAGAAAGCTGGGTG*CTT*TGTGGAGAAAGTCACAAAGCA-3´.

The italicized nucleotides denote the mutated stop codon. The amplicon was then cloned into

the pDONR221 vector by BP recombination to generate the middle entry pME-sema7a[sec]

vector. To generate the hsp70:sema7a[sec]-mKate2 construct, gateway cloning was performed

by combining the plasmids p5E-hsp70 (49), pME-sema7a[sec], p3E-mKate2-myc no-pA

(Addgene, 80812), and pDESTtol2pACrymCherry (Addgene, 64023) with LR Clonase II Plus

(Invitrogen). Verified constructs (25 ng/µl plasmid DNA) were injected with Tol2 Transposase

mRNA (approximately 25 ng/µl) into one-cell embryos. The transiently transgenic larvae—

identified by the expression of red fluorescent protein (mCherry) in their lenses—were raised

till 3 dpf, heat-shocked in a water bath at 37 °C for 1 hr, incubated at 28 °C for 1 hr to allow

expression, and subsequently mounted for live imaging.

Transient transgenesis with the hsp70I:sema7a-GPI-2A-mCherry construct. The *sema7a-*

GPI coding sequence without the stop codon (*sema7a-GPI^{no-stop}*) was synthesized using

gBlocks HiFi Gene Fragments (Integrated DNA Technologies). The *sema7a-GPI^{no-stop}* coding

sequence was then cloned into the pDONR221 vector by BP recombination to generate the

middle entry pME-sema7a-GPI^{no-stop} vector, henceforth denoted as pME-sema7a-GPI. To

generate the hsp70:sema7a-GPI-2A-mCherry construct, gateway cloning was performed by

combining the plasmids p5E-hsp70 (49), pME-sema7a-GPI, p3E-2A-mcherrypA (Addgene,

26031), and pDESTtol2pACryGFP (Addgene, 64022) with LR Clonase II Plus (Invitrogen).

Verified constructs (25 ng/µl plasmid DNA) were injected with Tol2 Transposase mRNA

(approximately 25 ng/µl) into one-cell embryos. The transiently transgenic larvae—identified

by the expression of green fluorescent protein (GFP) in their lenses—were raised till 3 dpf,

heat-shocked in a water bath at 37 °C for 1 hr, incubated at 28 °C for 1 hr to allow expression,

and subsequently mounted for live imaging.

Microscopy and volumetric rendering of living neuronal arbors and cells. Living larvae

were dechorionated, anaesthetized in 600 µM 3-aminobenzoic acid ethyl ester

methanesulfonate in system-water with 1 µg/mL methylene blue, and mounted in 1 % low-

melting-point agarose on a glass-bottom MetTek dish at specific stages. The larvae were

maintained at 28 °C during imaging in a temperature-controlled stage top chamber (OKO Lab

UNO-T). Static images at successive focal depths were captured at 200 nm intervals and

deconvolved with Microvolution software in ImageJ (NIH). For time-lapse video microscopy,

the ectopic *sema7a^{sec}* integration sites were imaged at intervals of 5 min as Z-stacks acquired

with 1.0 µm steps with laser excitation at 488 nm and 561 nm under a 60X, silicone-oil objective

lens of numerical aperture 1.30. The three-dimensional datasets were processed and volume-

rendered with the surface evolver and filament-tracer tools in Imaris (Bitplane, Belfast, UK).

The Imaris workstation was provided by the Rockefeller University Bio-imaging Resource

Center.

Generation of pseudocolored trajectories of the dwell times of sensory neurites at the

ectopic Sema7A^{sec} sources. Fluorescent intensity values of the sensory neurites and the

ectopic Sema7A^{sec} sources were measured from the maximal-intensity projections at each time

during the time-lapse video microscopy. The intensity values were normalized and binarized

before summing across each timepoint for locations across the XY plane. The resulting

pseudocolored trajectories represent the dwell times of the neurites and the Sema7A^{sec} cells

through the 10 hr time-lapse video sequence.

Generation of skeletonized networks. The labeled sensory axons were traced in three

dimensions with ImageJ's semi-automated framework, SNT (50). Each trace depicts the

skeletonized form of the posterior lateral-line nerve, the posterior lateral-line branch, and the

point of arborization from which the sensory arbors radiate to contact the hair cells

(Figure 2D,E; Figure 2—figure supplement 2B. The three-dimensional pixels or voxels

obtained from the skeletonized sensory arbor traces were processed using custom code written

in Python for visual representation and quantitative analysis.

Generation of combined hair cell clusters and the corresponding combined skeletonized

networks. The hair cell clusters of neuromasts from each developmental stage were aligned

by the centers of their apices, which were identified by the hair bundles, and overlayed to

generate combined hair cell clusters (Figure 2—figure supplement 1E,F) using custom Python

code. The associated skeletonized sensory-arbor traces at each developmental stage were

aligned similarly using custom Python code. The details of the procedures are available upon

request and the corresponding code is available on GitHub.

Quantification of sensory arbor distributions around hair cell clusters. The region from

the center of the combined hair cell cluster to 60 µm in the post-cluster region was divided into

3600 concentric sections of equal area. Each section, as depicted with a distinct color shade,

had an area of π μm^2. Sensory arbor density was defined as Log$_{10}$[(Area occupied by the arbors

in each section)/ (Area of each section)]. The arbor density was then plotted as a histogram

against distance from the center of the combined hair cell cluster. Each bin of the histogram

represents an area of π μm^2 (Figure 2—figure supplement 2A). The histograms were finally

represented as density trace graphs with Kernel Density Estimation (KDE) in Python.

Quantification of contact between sensory arbors and hair cell clusters. For each hair cell

cluster, the voxels of the sensory arbor traces that were inside or within 0.5 μm—the average

neurite radius—of the cluster boundary were denoted as arbors in contact with their

corresponding hair cell cluster (Figure 2—figure supplement 2B). The percentage of arbors in

contact with each hair cell cluster was defined as: (Number of voxels that remained inside the

cluster boundary/Total number of voxels in the arbor) X 100. The data were analyzed through

an automated pipeline with custom Python code.

Quantification of the sensory arborization network topology. We developed a custom

Python script (snt_topology.py; available on GitHub) to convert the skeletonized sensory

arborization networks to network graphs (NetworkX, v3.1). We first used a custom ImageJ

macro to convert SNT skeletons to csv files, then generated the network graph from the

individual arbor paths. As noted previously, we defined five topological attributes: (1) nodes—

junctions where arbors branch, converge, or cross; (2) loops—minimal arbor cycles forming

topological holes; (3) bare terminals—arbors with free ends; (4) node degree—the number of

first-neighbor nodes incident to a node; and (5) arbor curvature—the total curvature of a bare

terminal (Figure 3A). To enumerate loops and quantify their lengths and node numbers, we

computed the minimum cycle basis for each network graph. The relative organ height of each

node in a graph was defined in 3D using Equation (1):

670
$$(\vec{u} \cdot \vec{v}) \ / \ \|\vec{v}\|^2 \, , \tag{1}$$

where $\vec{u}$ is the vector from the arborization initiation point to a node and $\vec{v}$ is the vector from the

arborization initiation point to the hair cell cluster apex. The relative occurrence of nodes (Figure

3—figure supplement 1A-C) was normalized by both the number of hair cells per neuromast

and the total number of neuromasts, for comparison across ages and genotypes. The relative

number of bare terminals and loops (Figure 3B; Figure 3—figure supplement 1D, E) was

normalized by the number of hair cells per neuromast and scaled between 0 and 1. The density

was then normalized by the total number of neuromasts imaged for each developmental stage

and genotype. For Figure 3C and Figure 3—figure supplement 1F and G, the normalized

occurrence was computed as the number of arbors of a given length that formed either a bare

terminal or a loop, divided by the total number of arbors of either feature of that length. Lastly,

we defined bare terminals as having either high or low curvature (Figure 3F; Figure 3—figure

supplement 2E,F) if their absolute total signed curvature [Equation (2)] in three dimensions was

either higher or lower than $\pi/6$, respectively:

684
$$\left| \int \frac{dT}{ds} \cdot N(s)\, ds \right|, \tag{2}$$

where s is the arc-length parameter of the bare terminal, $T(s)$ is the unit tangent vector at each

point along the arbor, $N(s)$ is the unit normal vector at each point, and the integral is evaluated

from the terminal's branching node to the arbor's tip. For a closed curve forming a full loop, this

value will equal 2π. In practice, we first fit B-splines (scipy.interpolate.splprep; SciPy, v1.10.1)

in 3D to the skeletons for each bare terminal before evaluating their derivatives

(scipy.interpolate.splev) and integrating (scipy.integrate.simpson) over their lengths.

Quantification of aberrant sensory arborization in ectopic Sema7A^{sec} expression. The

sensory arbor networks were traced in three dimensions with SNT. The surface of each of the

ectopically expressing Sema7A^{sec}-mKate2 cells was represented by a single point that was

closest to the terminal point of the guided neurite. Distances between two points were

calculated using standard mathematical operations.

Quantification of presynaptic and postsynaptic assembly number and area. Maximal-

intensity projections of consecutive through-focus scans encompassing the entire neuromast

of control and *sema7a$^{-/-}$* mutant larvae at 2 dpf, 3 dfp, and 4 dpf were utilized to capture the

presynaptic and postsynaptic assemblies.   The number of the synaptic densities across

genotypes were measured using ImageJ's cell counter framework. Area of the synaptic

densities from the maximal-intensity projections were measured using ImageJ's polygon

selection tool (Figure 6—figure supplement 2A-D).

Quantification of the distribution of postsynaptic aggregates along the loops and the

bare terminals of the sensory arbor networks. Four-day-old *TgBAC(neurod1:EGFP)* larvae

were fixed and stained for sensory arbors and postsynaptic aggregates marked by GFP and

MAGUK, respectively. The GFP-positive sensory arbors were traced in three dimensions with

ImageJ's semi-automated framework, SNT. The topological attributes of the arbor network,

such as the loop and the bare terminal, were determined as described previously. The

localizations of the MAGUK punctae along the three-dimensional arbor network were

measured using ImageJ's cell counter framework (Figure 6—figure supplement 2E,F). We

measured 567 MAGUK punctae from 16 sensory arbor networks. Whether the MAGUK-

punctae were associated with a loop or a bare terminal, we categorized them in two groups

and quantified their distribution across 16 neuromasts.

Quantification of the distribution of postsynaptic-aggregate density along the distance

from the nearest node. The three-dimensional space surrounding each node of the arbor

network were divided into concentric shells of equal volume (Figure 6—figure supplement 2E,

F). For each node, we computationally generated 100,000 concentric shells that reached a 10

μm radius. We measured the number of MAGUK punctae located within each of those

concentric shells from their nearest nodes.

Statistical analysis. Data visualization and statistical analysis were conducted with GraphPad

Prism (Version 9) and Python (v3.10). The Mann-Whitney test was used for hypothesis testing

unless otherwise noted, and the statistical details are given in the respective figure legends.

Data and code availability. Owing to size restrictions, the experimental image data have not

been uploaded to a repository but are available from the corresponding author on request. The

codes generated during this study are available on GitHub
(https://github.com/agnikdasgupta/Sema7A_regulates_neural_circuitry).

Acknowledgments. The authors thank Nicolas Velez and Emily Atlas for valuable discussions.
Samantha Campbell and Anna Kaczynska provided expert fish husbandry. Katie Kindt kindly
provided various strains of zebrafish. Image processing benefitted from facilities at the
Bio-Imaging Resource Center. A.D. was supported by a Kavli Neural Systems Institute
Postdoctoral Fellowship from The Rockefeller University. C.C.R. is supported by National
Science Foundation Graduate Research Fellowship Grant No. 1946429. A.J., S.P.P., and
L.M.S. were supported by Howard Hughes Medical Institute, of which A.J.H. is an Investigator.

Author Contributions

A.D. initiated the project, conducted experiments, analyzed data, and quantified results. C.C.R
developed the Python codes for the network topology analysis and quantified those results, as
well as analyzed published transcriptomics data. S.P.P. developed the Python codes for three-
dimensional visual representation of sensory arbors and quantified arbor morphologies. L.M.S.
conducted preliminary experiments. A.J. plotted Sema7A intensity measurements. A.D. wrote
the paper with contributions from C.C.R and A.J.H.

Declaration of Interests

The authors declare no competing interests.

References

1. Lykissas M, Batistatou A, Charalabopoulos K, Beris A. The Role of Neurotrophins in
Axonal Growth, Guidance, and Regeneration. Curr Neurovasc Res. 2007 May 1;4(2):143–
51.

2. Vaccaro G, Dumoulin A, Zuñiga NR, Bandtlow CE, Stoeckli ET. The Nogo-66 Receptors
NgR1 and NgR3 Are Required for Commissural Axon Pathfinding. J Neurosci. 2022 May
18;42(20):4087–100.

3. Nguyen-Ba-Charvet KT, Di Meglio T, Fouquet C, Chedotal A. Robos and Slits Control the
Pathfinding and Targeting of Mouse Olfactory Sensory Axons. J Neurosci. 2008 Apr
16;28(16):4244–9.

4. Jeroen Pasterkamp R, Peschon JJ, Spriggs MK, Kolodkin AL. Semaphorin 7A promotes
axon outgrowth through integrins and MAPKs. Nature. 2003 Jul;424(6947):398–405.

5. Nagiel A, Andor-Ardo D, Hudspeth AJ. Specificity of Afferent Synapses onto Plane-
Polarized Hair Cells in the Posterior Lateral Line of the Zebrafish. J Neurosci. 2008 Aug
20;28(34):8442–53.

6. Jacobo A, Dasgupta A, Erzberger A, Siletti K, Hudspeth AJ. Notch-Mediated Determination
of Hair-Bundle Polarity in Mechanosensory Hair Cells of the Zebrafish Lateral Line. Curr
Biol. 2019 Nov;29(21):3579-3587.e7.

7. Valera G, Markov DA, Bijari K, Randlett O, Asgharsharghi A, Baudoin JP, et al. A neuronal
blueprint for directional mechanosensation in larval zebrafish. Curr Biol. 2021
Apr;31(7):1463-1475.e6.

8. Dow E, Jacobo A, Hossain S, Siletti K, Hudspeth AJ. Connectomics of the zebrafish's
lateral-line neuromast reveals wiring and miswiring in a simple microcircuit. eLife. 2018 Jun
12;7:e33988.

9. Lush ME, Diaz DC, Koenecke N, Baek S, Boldt H, St Peter MK, et al. scRNA-Seq reveals
distinct stem cell populations that drive hair cell regeneration after loss of Fgf and Notch
signaling. eLife. 2019 Jan 25;8:e44431.

10.   Carulli D, de Winter F, Verhaagen J. Semaphorins in Adult Nervous System Plasticity
and Disease. Front Synaptic Neurosci. 2021 May 11;13:672891.

11.   Yazdani U, Terman JR. The semaphorins. Genome Biol. 2006;7(3):211.

12.   de Winter F, Kwok JCF, Fawcett JW, Vo TT, Carulli D, Verhaagen J. The
Chemorepulsive Protein Semaphorin 3A and Perineuronal Net-Mediated Plasticity. Neural
Plast. 2016;2016:1–14.

13.   Coate TM, Spita NA, Zhang KD, Isgrig KT, Kelley MW. Neuropilin-2/Semaphorin-3F-
mediated repulsion promotes inner hair cell innervation by spiral ganglion neurons. eLife.
2015 Aug 24;4:e07830.

14.   Carcea I, Patil SB, Robison AJ, Mesias R, Huntsman MM, Froemke RC, et al.
Maturation of cortical circuits requires Semaphorin 7A. Proc Natl Acad Sci. 2014 Sep
23;111(38):13978–83.

15.   Inoue N, Nishizumi H, Naritsuka H, Kiyonari H, Sakano H. Sema7A/PlxnCl signaling
triggers activity-dependent olfactory synapse formation. Nat Commun. 2018
Dec;9(1):1842.

16.   Inoue N, Nishizumi H, Ooyama R, Mogi K, Nishimori K, Kikusui T, et al. The olfactory
786   critical period is determined by activity-dependent Sema7A/PlxnC1 signaling within
787   glomeruli. eLife. 2021 Mar 29;10:e65078.

788   17.   Morote-Garcia JC, Napiwotzky D, Köhler D, Rosenberger P. Endothelial Semaphorin
789   7A promotes neutrophil migration during hypoxia. Proc Natl Acad Sci. 2012 Aug
790   28;109(35):14146–51.

791   18.   Nishide M, Kumanogoh A. The role of semaphorins in immune responses and
792   autoimmune rheumatic diseases. Nat Rev Rheumatol. 2018 Jan;14(1):19–31.

793   19.   Köhler D, Granja T, Volz J, Koeppen M, Langer HF, Hansmann G, et al. Red blood
794   cell-derived semaphorin 7A promotes thrombo-inflammation in myocardial ischemia-
795   reperfusion injury through platelet GPIb. Nat Commun. 2020 Dec;11(1):1315.

796   20.   Pham TL, He J, Kakazu AH, Jun B, Bazan NG, Bazan HEP. Defining a mechanistic
797   link between pigment epithelium–derived factor, docosahexaenoic acid, and corneal nerve
798   regeneration. J Biol Chem. 2017 Nov;292(45):18486–99.

799   21.   Xie J, Wang H. Semaphorin 7A as a potential immune regulator and promising
800   therapeutic target in rheumatoid arthritis. Arthritis Res Ther. 2017 Dec;19(1):10.

801   22.   Baek S, Tran NTT, Diaz DC, Tsai YY, Acedo JN, Lush ME, et al. Single-cell
802   transcriptome analysis reveals three sequential phases of gene expression during
803   zebrafish sensory hair cell regeneration. Dev Cell. 2022 Mar 28;57(6):799-819.e6.

804   23.   Dow E, Siletti K, Hudspeth AJ. Cellular projections from sensory hair cells form
805   polarity-specific scaffolds during synaptogenesis. Genes Dev. 2015 May 15;29(10):1087–
806   94.

807   24.   Pujol-Martí J, Faucherre A, Aziz-Bose R, Asgharsharghi A, Colombelli J, Trapani JG,
808   et al. Converging Axons Collectively Initiate and Maintain Synaptic Selectivity in a
809   Constantly Remodeling Sensory Organ. Curr Biol. 2014 Dec;24(24):2968–74.

810   25.   De Wit J, De Winter F, Klooster J, Verhaagen J. Semaphorin 3A displays a punctate
811   distribution on the surface of neuronal cells and interacts with proteoglycans in the
812   extracellular matrix. Mol Cell Neurosci. 2005 May;29(1):40–55.

813   26.   Dalle Nogare DE, Natesh N, Vishwasrao HD, Shroff H, Chitnis AB. Zebrafish Posterior
814   Lateral Line primordium migration requires interactions between a superficial sheath of
815   motile cells and the skin. eLife. 2020 Nov 25;9:e58251.

816   27.   Ji YR, Warrier S, Jiang T, Wu DK, Kindt KS. Directional selectivity of afferent neurons
817   in zebrafish neuromasts is regulated by Emx2 in presynaptic hair cells. eLife. 2018 Apr
818   19;7:e35796.

819   28.   Diaz-Horta O, Abad C, Sennaroglu L, Foster J, DeSmidt A, Bademci G, et al. ROR1 is
820   essential for proper innervation of auditory hair cells and hearing in humans and mice. Proc
821   Natl Acad Sci U S A. 2016 May 24;113(21):5993–8.

29.   Newman M. The structure of real-world networks [Internet]. Vol. 1. Oxford University
Press; 2018 [cited 2023 May 2]. Available from:
https://academic.oup.com/book/27884/chapter/203817512

30.   Sharma P, Ruel TD, Kocha KM, Liao S, Huang P. Single cell dynamics of embryonic
muscle progenitor cells in zebrafish. Development. 2019 Jan 1;dev.178400.

31.   Keenan SR, Currie PD. The Developmental Phases of Zebrafish Myogenesis. J Dev
Biol. 2019 Jun 2;7(2):12.

32.   Sato A, Koshida S, Takeda H. Single-cell analysis of somatotopic map formation in the
zebrafish lateral line system. Dev Dyn. 2010 Jul;239(7):2058–65.

33.   Shen K, Cowan CW. Guidance Molecules in Synapse Formation and Plasticity. Cold
Spring Harb Perspect Biol. 2010 Apr 1;2(4):a001842–a001842.

34.   Um JW, Ko J. Neural Glycosylphosphatidylinositol-Anchored Proteins in Synaptic
Specification. Trends Cell Biol. 2017 Dec;27(12):931–45.

35.   Uesaka N, Uchigashima M, Mikuni T, Nakazawa T, Nakao H, Hirai H, et al. Retrograde
semaphorin signaling regulates synapse elimination in the developing mouse brain.
Science. 2014 May 30;344(6187):1020–3.

36.   Sheets L, Hagen MW, Nicolson T. Characterization of Ribeye Subunits in Zebrafish
Hair Cells Reveals That Exogenous Ribeye B-Domain and CtBP1 Localize to the Basal
Ends of Synaptic Ribbons. Riley B, editor. PLoS ONE. 2014 Sep 10;9(9):e107256.

37.   Oliva C, Escobedo P, Astorga C, Molina C, Sierralta J. Role of the maguk protein
family in synapse formation and function. Dev Neurobiol. 2012 Jan;72(1):57–72.

38.   Suli A, Pujol R, Cunningham DE, Hailey DW, Prendergast A, Rubel EW, et al.
Innervation regulates synaptic ribbons in lateral line mechanosensory hair cells. J Cell Sci.
2016 Jan 1;jcs.182592.

39.   Obholzer N, Wolfson S, Trapani JG, Mo W, Nechiporuk A, Busch-Nentwich E, et al.
Vesicular Glutamate Transporter 3 Is Required for Synaptic Transmission in Zebrafish Hair
Cells. J Neurosci. 2008 Feb 27;28(9):2110–8.

40.   Taschenberger H, Leão RM, Rowland KC, Spirou GA, von Gersdorff H. Optimizing
Synaptic Architecture and Efficiency for High-Frequency Transmission. Neuron. 2002
Dec;36(6):1127–43.

41.   Myers JP, Gomez TM. Focal Adhesion Kinase Promotes Integrin Adhesion Dynamics
Necessary for Chemotropic Turning of Nerve Growth Cones. J Neurosci. 2011 Sep
21;31(38):13585–95.

42.   O'Donnell M, Chance RK, Bashaw GJ. Axon Growth and Guidance: Receptor
Regulation and Signal Transduction. Annu Rev Neurosci. 2009 Jun 1;32(1):383–412.

43.   Gilmour D, Knaut H, Maischein HM, Nüsslein-Volhard C. Towing of sensory axons by
their migrating target cells in vivo. Nat Neurosci. 2004 May;7(5):491–2.

44.   Metcalfe WK. Sensory neuron growth cones comigrate with posterior lateral line
primordial cells in zebrafish. J Comp Neurol. 1985 Aug 8;238(2):218–24.

45.   Gasanov EV, Rafieva LM, Korzh VP. BDNF-TrkB Axis Regulates Migration of the
Lateral Line Primordium and Modulates the Maintenance of Mechanoreceptor Progenitors.
Proost P, editor. PLOS ONE. 2015 Mar 9;10(3):e0119711.

46.   Schuster K, Dambly-Chaudière C, Ghysen A. Glial cell line-derived neurotrophic factor
defines the path of developing and regenerating axons in the lateral line system of
zebrafish. Proc Natl Acad Sci. 2010 Nov 9;107(45):19531–6.

47.   Germanà A, Laurà R, Montalbano G, Guerrera MC, Amato V, Zichichi R, et al.
Expression of Brain-Derived Neurotrophic Factor and TrkB in the Lateral Line System of
Zebrafish During Development. Cell Mol Neurobiol. 2010 Jul;30(5):787–93.

48.   Kindt KS, Finch G, Nicolson T. Kinocilia Mediate Mechanosensitivity in Developing
Zebrafish Hair Cells. Dev Cell. 2012 Aug;23(2):329–41.

49.   Kwan KM, Fujimoto E, Grabher C, Mangum BD, Hardy ME, Campbell DS, et al. The
Tol2kit: A multisite gateway-based construction kit forTol2 transposon transgenesis
constructs. Dev Dyn. 2007 Nov;236(11):3088–99.

50.   Arshadi C, Günther U, Eddison M, Harrington KIS, Ferreira TA. SNT: a unifying
toolbox for quantification of neuronal anatomy. Nat Methods. 2021 Apr;18(4):374–7.

Figures and Figure Legends

Figure 1. Expression of Semaphorin7A in the zebrafish's lateral line. (A) A volumetric

rendering of a PLL neuromast depicts the sensory axons (magenta) that branch from the lateral

line nerve to arborize around the basolateral surface of the hair cell cluster (green). Additional

cell types in the neuromast are not labeled. Larval age, 3 dpf. A=anterior, P=posterior,

D=dorsal, V=ventral. (B) A schematic drawing of the two variants of the Sema7A protein

molecule depicts the full-length GPI-anchored form and the smaller, potentially secreted form.

Both the molecules include a signal sequence (green) and a conserved sema domain (orange).

(C) *Tg(myo6b:actb1-EGFP)* larvae at 4 dpf were terminally anesthetized and their heads were

removed with fine blades. The tails were then dissociated into the component cells, which were

sorted by flow cytometry. Green cells represent GFP+ hair cells. A cDNA library was prepared

from the isolated RNA. (D) Gel-based RT-PCR analysis indicates the presence of both the

sema7a transcript variants and the expression of hair cell-specific genes, parvalbumin 8

(*pvalb8*) and S100 calcium binding protein T (*s100t*). (E) A surface micrograph of a neuromast

at 3 dpf depicts two pairs of mature hair cells (green dashed lines) and a pair of immature hair

cells (grey dashed lines). Inset: among the three pairs of hair cell apices, the immature pair is

indicated by arrowheads. Immunolabeling reveals that the Sema7A protein (orange) occurs

consistently at the subapical region (arrowheads) and at the basolateral surface (arrows) of a

hair cell. In this and in each of the subsequent neuromast images, anterior is to the left and

dorsal to the top. (F) A plot quantifies developmental changes in the average Sema7A intensity

in both rostrally and caudally polarized hair cells of neuromasts from 1.5 dpf to 4 dpf. The data

stem from 18, 36, 39, and 40 hair cells in neuromasts of respectively 1.5 dpf, 2 dpf, 3 dpf, and

4 dpf larvae. (G) A plot quantifies the distribution of average Sema7A intensity along the hair

cell's apicobasal axis. The results stem from 52, 57, and 54 hair cells of neuromasts from 2 dpf,

3 dpf, and 4 dpf larvae. (H) An immunofluorescence image at the nuclear level of a 4 dpf

neuromast shows the contact of the sensory arbors (magenta) with the basolateral surface of

the hair cells (green). Immunolabeling for Sema7A (orange) reveals enrichment of the protein

at the hair cell bases and sensory-axon interfaces (arrowheads). (I) At the base of the hair cell,

association of the td-Tomato+ sensory arbors positively correlated with the Sema7A intensity.

The results stemmed from 48 hair cells of 13 neuromasts from 3 dpf larvae. HC, hair cell; scale

bars, 5 μm; au, arbitrary unit; means ± SEMs; ρ, Spearman's correlation coefficient; *** implies

p < 0.001 and ** implies *p* < 0.01.

Figure 1—figure supplement 1. Isolation of GFP-expressing hair cells. (A,B)
Representative results from florescence-activated cell sorting demonstrate the efficient
discrimination of GFP+ (green) hair cells from *Tg(myo6b:actb1-EGFP)* larval tails. Around 400
tails were utilized to isolate hair cells. Wild type (WT) controls confirm the selectivity of the
gates. (C,D) Plots represent the RNA integrity numbers (RINs) of the isolated total RNA from
WT and transgenic larvae. High RIN values suggest minimal degradation of isolated RNAs
from both samples.

Figure 1—figure supplement 2. Expression of Semaphorin7A in the lateral line neuromast. (A-C) Surface micrographs show hair cells marked with actin-GFP (green dashed lines) and the localization of Sema7A protein (orange) in the subapical (arrowheads) and basal regions (arrows) of the corresponding hair cells from neuromasts of developing control larvae. The inset in panel A (red arrowheads) depicts the apices of the immature hair cell pair. (D) Immunolabeling with only the secondary antibody fails to detect any Sema7A signal. (E) A surface micrograph depicts hair cells (green, left) whose subapical region (red dashed line) harbors the Golgi network–labeled by the Golgi resident protein GM130 (cyan, middle)–where

the Sema7A protein is enriched (orange, right). (F) A micrograph at a nuclear plane of a 4 dpf

neuromast depicts membrane-tethered Scarlet (green) in all cells of the neuromast and the

Sema7A protein (orange, arrowheads) in centrally clustered hair cells. (G) The Sema7A protein

is located at the basal membrane (red arrowhead). (H) The Sema7A protein is also observed

adjacent to the basal membrane, potentially in a vesicle (cyan arrowhead). (I) A plot quantifies

the distribution of average Sema7A intensity across the hair cell's basal membrane. The results

stem from 100 hair-cell membranes of 10 neuromasts from 4 dpf larvae. (J) A schematic

diagram of a single hair cell depicts the three distinct regions along the apicobasal axis of the

cell. Sema7A intensity was measured along this apicobasal axis (red dashed line). Scale bars,

5 μm; au, arbitrary unit; means ± SEMs.

Figure 1—figure supplement 3. Sema7A progressively enriches at the hair cell base with the sensory axons. (A) A micrograph at a nuclear plane of a 3 dpf neuromast depicts several actin-GFP (green) labeled hair cells. The lateral (white dashed line) and the basal (green dashed line) surfaces of three hair cells are outlined. (B) The td-Tomato-positive sensory arbors contact the basal regions (green dashed line) of three hair cells. (C) The Sema7A is enriched in the basal regions (green dashed line) of the hair cells that are in contact with the sensory arbors. (D,E) At the base of the hair cell, intensities of the td-Tomato-positive sensory arbors positively correlate with the Sema7A intensities. The results stem from 66 hair cells of 15 neuromasts from 2 dpf larvae and 40 hair cells of 11 neuromasts from 4 dpf larvae. Scale bars, 5 μm; au, arbitrary unit; ρ, Spearman's correlation coefficient.

```
Mus        MTPPPPGRAAPSAPRARVLSLPARFGLPLRLRLLLVFWVAAASAQGHSRSGPRISAVWKG
Danio      ------------------MIRHYSSLKFKGNLAYPWLIYGACFQCVFCSCNPRLTLKQD
                         :    .* :: .*    : : .*. *    *     :: :.

Mus        QDHVDFSQPEPHTVLFHEPGSFSVWVGGRGKVYHFNFPEGKNASVRTVNIGSTKGSCQDK
Danio      NAPLNYNTNYSHTVLFHEEGSKSLFVGGSNIVLHFDVDSYEILENFTLNANLPR---CGE
           :  :::.    .******* ** *::*** . * **:. .: .  *:* . .:    .:

Mus        QDCGNYITLLERRGNGLLVCGTNARKPSCWNLVN---DSVMMSLGEMKGYAPFSPDENSL
Danio      SSCENVVTVIETFQNCKFVCGTNGEEPQCWELYPRESNHSVVKSVEGTGISPHSSEQNSL
           ..* * :*::*    *   :*****..:*.**:*      :  ::.  * .* :*.*.::***

Mus        VLFEGDEVYSTIRKQEYNGKIPRFRRIRGESELYTSDTVMQNPQFIKATIVH-QDQAYDD
Danio      SLVADGDLYAAAPLYSDGTLLQFRRKAGKRINVWMHDQWVSEPTFISSFLAKRKNDSLNE
            *. ..::*::    . . :   *:   . ::: *  :.:* **.: :.: :::: ::

Mus        KIYYFFREDNPDKNPEAPLNVSRVAQLCRGDQGGESSLSVSKWNTFLKAMLVCSDAATNR
Danio      KIYVLFREKNSDTSPEADPWISRVARVCKIDEGGSKQFLQNIWTSFLKARLVCGIPKESL
           *** :***.*.*..***   :****::*: *:**...:  . *.:**** ***. .  .

Mus        NFNRLQDVFLLPDPSGQWRDTRVYGVFSNPWNYSAVCVYSLGDIDRVFRTSSLKGYHMGL
Danio      YFNRLQDVFVQH--AEDWRDSLVYALFSSSWNSTAVCIYSLADLDDIFEKSNYKGFSEAI
            ********:    : :***: **.:**..** :***:***.*:* :*...*. **: .:

Mus        PNPRPGMCLPKKQPIPTETFQVADSHPEVAQRVEPMGPLKTPLFHSKYHYQKVVVHRMQA
Danio      PNPRPGKCVENSNSLPIPTLKIIKDYPEMKDWIHPIQ-KDTPFFTSSKNFTKIVVDQIRA
           ****** *: :.:..:*   *:::: ..:**: : :.*:     .**:* *. :: *:**.:::*

Mus        SNGETFHVLYLTTDRGTIHKVVESGDQDHSFVFNIMEIQPFHRAAAIQAISLDADRRKLY
Danio      LDENVYNVLLLATDDGMIFKILQHG----SKPFIISETHLCNGCAPIQSMKLDSKKRKLF
            : :.::** *:** * *.*::: *    *   * * * :   :  .*.**::.**:.:***:

Mus        VTSQWEVSQVPLDMCEVYSGGCHGCLMSRDPYCGWDQDRCVSIYSS--QRSVLQSINPAE
Danio      VGYPGQMSVLELQRCQDYGASCEECVLSRDPYCAWSEKGCTSQTRGGIQNISDGNVTKVC
           *     ::* :  *: *: *...*. *::*******.*.:. *.*     .  *.     .:. .

Mus        PHRECPNPKPDE------APLQKVSLARNSRYYLTCPMESRHATYLWRHEENVEQSCEPG
Danio      DHKSVGRVRRDAQQSSTAPSRIQHTVATDVPFYLSCPVDSHHASYRWEHGG---KSNPCQ
            *:.  . : *       .. : ::* :  :**:**::*:**:* *.*       :*

Mus        HQSPSCILFIENLTARQYGHYRCEAQEGSYLREAQHWELLPE-DRALAEQLMGHARALAA
Danio      QTQSEYLLLIPAMTAENYGNYSCVSQERDYIKVVREYELLAKSVFNDAFKLKIQDRQLMV
           :  ... :*:*  :**.:**:* * :** .*:: .::.:***.:     * :*  : * *  .

Mus        SFWLGVLPTLILGLLVH
Danio      AFVTSAFYLCSL-----
           :*   ..:     *
```

Sequence 1: Mus musculus   664 aa
Sequence 2: Danio rerio   640 aa
Immunogen: Mouse myeloma cell line NS0-derived recombinant mouse Semaphorin 7A Glutamine 45 to Alanine 646. The corresponding region in the zebrafish's Semaphorin 7A is represented by Glutamine 26 to Methionine 627. We predict that the antibody recognizes both variants of the *sema7a* encoded proteins. The Sema domains in both mouse and zebrafish are highlighted in orange.

Figure 1—figure supplement 4. Alignment of the mouse and the zebrafish Sema7A

protein sequences generated by Clustal Omega. (A) Multiple sequence alignment depicts

the conserved Sema domain (orange) and the immunogenic region for the Sema7A antibody

in mouse, and its equivalent region in zebrafish.

Video 1. Sema7A is enriched at the hair cell base and sensory-axon interface. A through-

focus scan of a 4 dpf control neuromast depicts the hair cells (green) that are innervated by the

sensory axons (magenta) at their basolateral surfaces where the Sema7A protein (orange) is

highly enriched. Scale bar, 5 μm.

Figure 2. *sema7a^-/-* mutants display aberrant sensory axon arborizations. (A-D) In micrographs of 3 dpf control and *sema7a^-/-* neuromasts, Myosin VI (green) marks the hair cells (green dashed lines). The level of Sema7A (orange) is highly reduced in the *sema7a^-/-* mutant compared to the control, with sporadic localization in the subapical region (arrowheads) and none in the basolateral (arrow) region. (E) A plot of normalized Sema7A intensity from 99 control and 100 *sema7a^-/-* hair cells quantitates the effect. (F,G) Surface views of a control and a *sema7a^-/-* neuromast at 4 dpf depict the interaction of the sensory arbors (magenta) with hair cell clusters (green). In the control, the arbors intimately contact the hair cells, with a few exceptions (arrowhead). In the *sema7a^-/-* mutant, the arbors direct many aberrant projections

(arrowheads) away from the hair cell cluster. The two immature hair cell pairs in the *sema7a*[-/-]

neuromast are indicated by arrows. (H,I) Skeletonized networks portray the three-dimensional

topology of the sensory arbors from the control and the *sema7a*[-/-] neuromasts depicted in

panels F and G, respectively. The pseudocolored trajectories depict the increase in arbor

contour length from each point of arborization, defined as the point at which the lateral line

branch (magenta) contacts hair cell cluster. (J,K) Micrographs depict the skeletonized networks

of the combined 4 dpf hair cell clusters, whose centers are located at (0,0). The X- and

Y-coordinates represent the anteroposterior (AP) and the dorsoventral (DV) axes of the larva,

respectively. Positive values of the X- and Y-coordinates represent the posterior and ventral

directions, respectively. Combined skeletonized network traces from 27 control and 27

sema7a[-/-] mutant neuromasts are represented. (L) The plot denotes the densities of the

sensory arbors around the center of the combined hair cell clusters for 35 control (magenta)

and 27 *sema7a*[-/-] (red) neuromasts at 4 dpf. The shaded area marks the region proximal to the

boundary of the combined hair cell cluster. (M) The plot quantifies the degree of contact of the

sensory arbors to their hair cell clusters in individual neuromasts from both control (magenta)

and *sema7a*[-/-] mutants (red), each point represents a single neuromast. Thirty-three, 29, and

35 neuromasts were analyzed from 2 dpf, 3 dpf, and 4 dpf control larvae, respectively.

Seventeen, 53, and 27 neuromasts were analyzed from 2 dpf, 3 dpf, and 4 dpf

sema7a[-/-] mutant larvae, respectively. HC, hair cell; Scale bars, 5 μm; au, arbitrary unit;

means ± SEMs; *** signifies $p < 0.001$; ** signifies $p < 0.01$.

1000

1001

1002

1003

1004

1005

1006

Figure 2—figure supplement 1. *sema7a⁻ᐟ⁻* **mutants exhibit aberrant sensory-axon arborization throughout development.** (A) Micrographs depict the overall morphologies and *sema7a* genomic profiles of 2 dpf control (top) and *sema7a⁻ᐟ⁻* larvae (bottom). The schematic diagram on the right depicts the mutation site and the resulting early truncation of the encoded protein. (B) The plot quantitates the number of hair cells from 33, 29, and 35 neuromasts from 2 dpf, 3 dpf, and 4 dpf control larvae, respectively and 17, 53, and 27 neuromasts from 2 dpf,

3 dpf, and 4 dpf *sema7a*$^{-/-}$ mutant larvae, respectively. No significant change in hair cell number was observed between the control and the *sema7a*$^{-/-}$ neuromasts across developmental stages. (C) Micrographs depict maximum intensity projections of 4 dpf neuromasts from control and *sema7a*$^{-/-}$ larvae where the hair cells (green) are stained with FM 4-64 (magenta). The sensory arbors (arrowheads, magenta) tightly enwrap the clustered hair cells in the control neuromast but show aberrant arborization in the *sema7a*$^{-/-}$ larvae. (D) The plot quantitates the average FM 4-64 intensities from the hair cells of the control and the *sema7a*$^{-/-}$ neuromasts. (E,F) Micrographs depict combined 4 dpf hair cell clusters, each of whose centers arise located at (0,0). The X- and Y-coordinates represent the anteroposterior (AP) and dorsoventral (DV) axes of the larva, respectively. Positive values of the X- and Y-coordinates represent posterior and ventral directions, respectively. Combined hair cell clusters from 27 control and 27 *sema7a*$^{-/-}$ mutant neuromasts are represented. (G,H) Micrographs depict the skeletonized networks of the combined 2 dpf hair cell clusters from 17 control and 17 *sema7a*$^{-/-}$ mutant neuromasts. (I,J) Micrographs depict the skeletonized networks of the combined 3 dpf hair cell clusters from 29 control and 29 *sema7a*$^{-/-}$ mutant neuromasts. (K) A plot quantitates the densities of the sensory arbors around the center of the combined hair cell clusters for 33 control (magenta) and 17 *sema7a*$^{-/-}$ (red) neuromasts at 2 dpf. The shaded area marks the region proximal to the boundary of the combined hair cell cluster. (L) A plot quantitates the densities of the sensory arbors around the center of the combined hair cell clusters for 29 control (magenta) and 53 *sema7a*$^{-/-}$ (red) neuromasts at 3 dpf. The shaded area marks the region proximal to the boundary of the combined hair cell cluster. Scale bars, 5 μm; au, arbitrary units; ns, not significant.

Figure 2—figure supplement 2. Quantification of sensory arbor distributions and

contacts with hair cell clusters. (A) A schematic drawing of the combined hair cell cluster

and surrounding region depicts concentric sections of equal area in distinct colors (cyan, green,

yellow, and red). The arbor density is plotted as a histogram against the distance (black arrow)

from the center (black circle) of the combined hair cell cluster. The histogram is represented as

a density trace graph (magenta). (B) A schematic drawing of a posterior lateral-line neuromast

depicts the posterior lateral-line (PLL) nerve (black), posterior lateral-line branch (black), the

point or arborization, the sensory arbors (magenta) that contact the hair cell cluster (green),

and the sensory arbors that project away (blue) from the hair cell cluster.

1045

Videos 2 and 3. FM4-64 labels the hair cells of both control and *sema7a⁻ᐟ⁻* mutant

neuromasts. Through-focus scans of 4 dpf control (Video 2) and *sema7a⁻ᐟ⁻* mutant (Video 3)

neuromasts depict the hair cells (green) that have incorporated FM4-64. The sensory arbors

(magenta) are tightly associated with the clustered hair cells in the control neuromast but show

aberrant arborization in the *sema7a⁻ᐟ⁻* larvae.

Video 4. Three-dimensional arborization patterns in control and *sema7a⁻ᐟ⁻* neuromasts.

Lateral views depict three-dimensionally rendered combined skeletonized network traces from

27 control and 27 *sema7a⁻ᐟ⁻* mutant neuromasts at 4 dpf.

1054

Figure 3. Sema7A patterns the topology of the sensory arbor network. (A) A schematic drawing of the combined hair cell cluster (green) and associated sensory axons (magenta) shows diverse topological features of the arborization network, including nodes, loops, bare terminals, node degree, and bare-terminal curvature. The apex is defined as the center of the hair bundle cluster and the base is denoted by the node from which the arborizations extends. The relative organ height describes the position along the axis formed between the arborization

initiation point and the apex of the hair cell cluster. (B) The correlation between the relative

number of bare terminals and loops from 35 control neuromasts and 27 *sema7a*[-/-] neuromasts

at 4 dpf shows a negative correlation in the control but a scattered distribution in the mutants.

The same neuromasts were analyzed in the subsequent panels. R^2, correlation coefficient.

Dashed red line depicts the regression slope. (C) The relationship of contour lengths to bare

terminals and loops reveals reduced looping in the mutants. (D) Quantification of the number

of nodes per loop demonstrates a decrease in mutants. (E) Mutant larvae manifest a broader

distribution of loop lengths compared to controls. (F) A schematic diagram of the extending

bare terminals depicts regions of high and low curvature along the length of the arbor. The

distributions of bare terminals between low- and high-curvature groups show an enrichment of

long, highly curved arbors in the control and short, linear arbors in the mutants. Statistical

analyses are displayed in Table 1.

1073

Figure 3—figure supplement 1. Quantification of topological features across

developmental stages and genotypes. (A-C) The distribution of node degrees as a function

of the height in hair cell clusters shows the difference in the network's hierarchical organization

between control and *sema7a*⁻/⁻ larvae. Nodes of low degree are commonly observed between

genotypes across developmental stages (A, B), but their distribution along the cluster height

became significantly different at 4 dpf (C), with nodes in control larvae localizing higher along

the axis than in the mutants. In this and the following two panels, 33, 29, and 35 neuromasts

were analyzed from 2 dpf, 3 dpf, and 4 dpf control larvae, respectively and 17, 53, and 27

neuromasts were analyzed from 2 dpf, 3 dpf, and 4 dpf *sema7a*⁻/⁻ mutant larvae, respectively

(D, E) The correlations between the relative number of bare terminals and loops from control

and *sema7a⁻/⁻* neuromasts reveal dispersive distributions at both 2 dpf (D) and 3 dpf (E) stages,

suggesting that the networks are undergoing reorganization and yet to adopt an ordered

topology. (F,G) In the distribution of arbors between bare terminals and loops with respect to

their contour lengths, mutants' arbors display significantly higher propensity to remain as bare

terminals compared to the controls at both 2 dpf (F) and 3 dpf (G). The samples were identical

to those in the previous panel. Statistical analyses are displayed in Table 1.

1090

Figure 3—figure supplement 2. Quantification of topological features between control and *sema7a⁻/⁻* larvae across developmental stages. (A, B) Quantifications of the number of nodes per loop demonstrate a broader, but not significantly different, distribution in the mutants than the controls at both 2 dpf (A) and 3 dpf (B). In this and the following panels, 33 and 29 neuromasts were analyzed from 2 dpf and 3 dpf control larvae, respectively and 17 and 53 neuromasts were analyzed from 2 dpf and 3 dpf *sema7a⁻/⁻* mutant larvae, respectively. (C, D) Quantifications of loop lengths from the same larvae indicate a wider, but not significantly different, distribution in the mutants than the controls at both 2 dpf (C) and 3 dpf (D). (E, F) The distributions of bare terminals between low- and high-curvature groups from the same larvae show higher occurrences of long, curved arbors in the control and short, linear arbors in the mutants at both 2 dpf (E) and 3 dpf (F) stages. Statistical analyses are displayed in Table 1.

Height of node degrees	p-values (control vs $sema7a^{-/-}$)		
	2 dpf	3 dpf	4 dpf
Degree = 3	3.50E-01	1.05E-08***	1.68E-03**
Degree = 4	3.85E-01	7.11E-02	1.71E-03**
Degree = 5	4.60E-01	5.23E-01	4.40E-01
Bare terminals vs Loop	2 dpf	3 dpf	4 dpf
0 to 10 μm	3.84E-01	7.89E-01	2.11E-01
10 to 20 μm	3.50E-02*	3.92E-01	2.94E-02*
20 to 30 μm	2.28E-04***	2.63E-02*	3.82E-09***
30 to 40 μm	5.04E-59***	3.63E-25***	1.28E-33***
40 to 70 μm	———	———	———
Nodes per loop	5.06E-01	1.96E-01	3.50E-02*
Loop length	5.81E-01	2.71E-01	2.21E-02*
Distribution of bare terminal curvature	2 dpf	3 dpf	4 dpf
	4.23E-01	8.91E-01	1.99E-02*

1103

Table 1. Statistical comparison between control and $sema7a^{-/-}$ larvae for topological

features. One-sided Kolmogorov-Smirnov tests were used to assess the difference in node

degree heights, nodes per loop, and loop lengths for control *versus sema7a^{-/-}* neuromasts at

2 dpf, 3 dpf, and 4 dpf. In each of these cases, the difference in the distributions were not

significantly different (p-values in red) between the control and the mutants at 2 dpf and 3 dpf

but became significantly different (p-values in green) at 4 dpf. Chi-square tests were used to

assess the difference in occurrences of bare terminals versus loops, as well as the occurrences

of bare terminals with low versus high curvature. The distribution at 4 dpf shows significant

difference between the control and the mutant larvae. The occurrence of significant differences

at 4 dpf for diverse topological features suggest that Sema7A function is required to achieve

an ordered network topology at a mature stage. Thirty-three, 29, and 35 neuromasts were

1115 analyzed from 2 dpf, 3 dpf, and 4 dpf control larvae, respectively, and 17, 53, and 27

1116 neuromasts were analyzed from 2 dpf, 3 dpf, and 4 dpf *sema7a$^{-/-}$* mutant larvae, respectively.

1117 * Signifies $p < 0.05$, ** signifies $p < 0.01$, and *** signifies $p < 0.001$.

1118

 Figure 4. Ectopic Sema7A^sec diffusive cue provides neural guidance *in vivo*. (A) A

diagrammatic overview depicts the generation of a transgenic animal that expresses the Sema7A^{sec} protein ectopically under the control of a thermally inducible promoter. Larvae with ectopic myotomal-integration near the network of sensory arbors (red rectangle) were imaged to analyze arbor morphology. (B) A schematic drawing of a sensory arbor from a heat-shocked larva depicts an extended axonal projection (arrowhead) that reaches toward the myofibers expressing the ectopic Sema7A^{sec} protein. Parameters that quantitate the accuracy of the extended axonal projections toward the ectopic Sema7A^{sec} sources are denoted. (C) In a micrograph of an ectopically expressing Sema7A^{sec} (orange) larva, the sensory arbor (magenta) extends two aberrant axonal projections. One elongates (cyan arrowhead) along the somite boundary to reach and contact an ectopically integrated muscle progenitor cell (white dashed line) and the other (red arrowhead) reenters the posterior lateral-line nerve while following a second ectopic source. The through-focus scans (i–iii) from the epidermis to the dermomyotome and the three-dimensional (3D) surface reconstruction (iv) reveal the intimate contact between the aberrant sensory arbor (arrowheads) and the muscle progenitor cell. (D) In a micrograph of an ectopically expressing Sema7A^{sec} (orange) larva, the sensory arbor (magenta) extends a single aberrant axonal process (cyan arrowhead) to reach ectopically integrated myofibers (white dashed line). The through-focus scans (i–iii) from the epidermis to the dermomyotome and the 3D surface reconstruction (iv) reveal the proximal association of the aberrant sensory arbor (arrowheads) to the myofibers. Melanocytes (yellow arrowhead) along the horizontal myoseptum intermittently block the visibility of the lateral-line nerve. (E) An injected, but not heat-shocked, control larva does not express ectopic Sema7A^{sec} and does not show aberrant projection from the sensory arbor. (F) A plot demonstrates the accuracy of 18 extended axonal arbors in finding ectopic Sema7A^{sec} sources. Each circle represents a single ectopic integration event. The two pairs of numbers represent the minimal and maximal lengths of the projection path (black) and its corresponding source path (green). (G) A plot quantitates the distribution of projection-proximity length from 18 ectopic integration events. (H) A schematic drawing of a section of the lateral-line nerve between two neuromasts from a heat-shocked larva depicts a few extended neurites (magenta) reaching toward the cells expressing

the ectopic Sema7A^{sec} protein (orange). The Sema7A^{sec} integration site (blue rectangle) with

the sensory axonal projections was imaged for ten hours to visualize axonal dynamics. (I)

Micrographs (i-iii) at three distinct times in the time-lapse video microscopy show directed

branching (cyan arrowheads) of the sensory neurites from the lateral-line nerve toward the

ectopic Sema7A^{sec} sources. Each panel depicts the ectopic integration site and the associated

sensory neurites from three different planes, XY, XZ, and YZ. White arrowheads in XZ and YZ

planes highlight the close association of the sensory axons with the ectopic Sema7A^{sec}

sources. The t=0 hr denotes the onset of imaging, which is two hours post the beginning of

heat shock. (J) The micrographs depict in pseudocolored trajectories the dwell times of the

sensory neurites (left) at the Sema7A^{sec} sources (right) for ten hours. MPC, muscle progenitor

cell; scale bars, 20 μm; means ± SEMs.

1159

Figure 4—figure supplement 1. Ectopic Sema7A^{sec} diffusive cues induce aberrant

sensory neurite extension. (A-A") In a micrograph of an ectopically expressing Sema7A^{sec}

(orange) larva, the sensory arbors (magenta) extend multiple axonal projections (arrowheads)

that reenter the lateral-line nerve while following an ectopic Sema7A source. Maximal-intensity

projections of the sensory arbors and the ectopically expressing Sema7A cells (white dashed

line) are shown in A' and A", respectively. (B-B") Micrographs depict three individual incidents

of ectopic *sema7a^{sec}* (orange) integrations that occurred distant from neuromasts. In each

case, a set of myofibers expresses the ectopic Sema7A^{sec} protein that induces neurite

extensions (arrowheads) from the lateral-line nerve (magenta) toward itself. Scale bars, 20 μm.

1169

Figure 4—figure supplement 2. Ectopic integration of Sema7A^sec distant from the neuromast induces sensory neurite extension. (A) Micrographs of a section of the lateral-line nerve between two neuromasts from a 3-dpf heat-shocked larva depicts a few extended neurites (magenta) reaching toward the cells expressing the ectopic Sema7A^sec (orange). The

Sema7A^{sec} integration site with the sensory axonal projections was imaged for one hour to

visualize axonal dynamics. Micrographs (i-iii) at three distinct time-points from the time-lapse

video microscopy show directed branching (cyan arrowheads) of the sensory neurites from the

lateral-line nerve toward the ectopic Sema7A^{sec} sources. Each panel depicts the ectopic

integration site and the associated sensory neurites from three different planes, XY, XZ, and

YZ. White arrowheads in XZ and YZ planes highlight the close association of the sensory axons

with the ectopic Sema7A^{sec} sources. Melanocytes (yellow arrowheads) along the horizontal

myoseptum intermittently block the visibility of the lateral-line nerve. Scale bar, 20 μm.

1183

Video 5. Sensory axonal projections contact an embryonic muscle progenitor cell

expressing Sema7A^{sec}. A through-focus scan of a 3 dpf larva depicts a muscle progenitor

cell in the dermomyotome that expresses the ectopic Sema7A^{sec} (orange) and the sensory

axonal projection (magenta) that makes intimate contact with it. Scale bar, 20 μm.

Video 6. Volumetric surface reconstruction of a sensory arbor and an embryonic muscle

progenitor expressing Sema7A^{sec}. Reconstruction of the muscle progenitor cell (orange) at

3 dpf reveals its intimate association with an attracted sensory arbor (magenta).

Video 7. Volumetric surface reconstruction of a sensory arbor and the mature myofibers

expressing Sema7A^{sec}. Reconstruction of mature myofibers expressing ectopic Sema7A^{sec}

(orange) at 3 dpf shows their close association with the attracted sensory arbors (magenta).

Video 8. Sensory axonal projections remain in proximity to myofibers expressing

Sema7A^{sec}. A through-focus scan of a 3 dpf larva depicts mature myofibers expressing

exogenous Sema7A^{sec} deep in the myotome (orange) and the sensory axonal projection

(magenta) that remains nearby.

Video 9. Aberrant axonal projections reenter the lateral-line nerve while following an

ectopic Sema7A^{sec} source. A through-focus scan of a 3 dpf larva depicts cells expressing

exogenous Sema7A^{sec} (orange) that guides the aberrant sensory axonal projection (magenta)

back into the lateral-line nerve. Scale bar, 20 μm.

Videos 10, 11, and 12. Ectopic Sema7A^{sec} expression induces aberrant neurite

projections from the lateral-line nerve. Through-focus scans of 3 dpf larvae depict three

individual incidents of ectopic *sema7a^{sec}* (orange) integrations that occurred distant from the

neuromasts. In each case, myofibers express ectopic Sema7A^{sec} protein (orange) that attracts

neurite extensions from the lateral-line nerve (magenta). Scale bars, 20 μm.

Videos 13 and 14. Volumetric reconstructions depict close contacts between the

sensory neurites and the ectopic Sema7a^{sec} expressing cells. Volumetric reconstructions

of 3 dpf larvae depict two individual incidents of ectopic *sema7a^{sec}* (orange) integrations that

occurred distant from the neuromasts. In each case, myofibers or random mosaic of cells in

the dermomyotome express ectopic Sema7A^{sec} protein (orange) that attracts neurite

extensions from the lateral-line nerve (magenta).

Video 15 and 16. Time-lapse video microscopies depict the dynamics between the

sensory neurites and the ectopic Sema7a^{sec} expressing cells. Time-lapse images of 3 dpf

larvae depict two individual incidents where the ectopic Sema7A^{sec} sources (orange) induce

directed branching of the sensory neurites (magenta) from the lateral-line nerve toward itself.

Within each time-lapse videos, individual panel depicts the ectopic integration site and the

associated sensory neurites from three different planes, XY, XZ, and YZ. Scale bars, 20 μm.

1219

Figure 5. Ectopic Sema7A-GPI fails to guide sensory arbors from a distance. (A) A
diagrammatic overview depicts the generation of a transgenic animal that expresses the
Sema7A-GPI protein ectopically under the control of a thermally inducible promoter. Larvae
with ectopic myotomal-integration near the network of sensory arbors (red rectangle) were
imaged to analyze arbor morphology. Uninjected, heat-shocked larvae were used as controls.
(B) Uninjected and heat-shocked larva depicts normal arborization pattern of the sensory axons
(magenta). (C) Injected and heat-shocked larvae robustly express the Sema7A-GPI in the
myofibers (orange). Panels (i-iii) depict the ectopically expressing Sema7A-GPI myofibers that
fail to attract sensory arbors toward itself. Scale bar, 20 μm.

1229

1230 **Videos 17, 18, and 19. Sema7A-GPI is incapable to impart sensory axon guidance from**

1231 **a distance.** Through-focus scans of 3 dpf larvae depict three individual incidents of ectopic

1232 *sema7a-GPI* integrations that occurred near the sensory arbor network. In each case,

1233 myofibers robustly express ectopic Sema7A-GPI protein (orange) but the sensory arborization

1234 pattern (magenta) remain unperturbed. Scale bars, 20 μm.

1235

Figure 6. Sema7A deficiency impairs synaptic assembly. (A,B) Maximal-intensity projections of micrographs from a control and a *sema7a$^{-/-}$* neuromast depict the presynaptic

aggregates marked by CTBP (cyan). The approximate hair cell basal region is outlined by

dashed green circles. (C-F) Plots quantitate the numbers of presynaptic aggregates (C,D) and

their areas (E,F), in each hair cell across developmental stages. One hundred and fifty-two,

317, and 325 hair cells were analyzed from 2 dpf, 3 dpf, and 4 dpf control larvae, respectively.

One hundred and sixty, 216, and 177 hair cells were analyzed from 2 dpf, 3 dpf, and 4 dpf

sema7a$^{-/-}$ mutant larvae, respectively. (G) A micrograph generated from maximal-intensity

projection of ten consecutive through-focus scans across Z-planes of a control 4 dpf neuromast

depict the sensory arbors (magenta) and the postsynaptic aggregates marked by MAGUK

(yellow, white arrowheads). The MAGUK punctae primarily occur on the loop and aggregate

near the nodes. Panels (i-iv) depict maximal-intensity projection of consecutive through-focus

scans of distinct Z-planes to show bare terminals. In each panel, the bare terminal (magenta,

cyan arrowheads) rarely harbors postsynaptic aggregates (yellow, white arrowhead). (H)

Quantification of the distribution of the MAGUK-punctae between the two topological features

of the network depicts enrichment of postsynaptic densities on the loops over the bare

terminals. (I) A histogram shows the distribution of MAGUK-punctae density against the

distance from the node. The histogram is represented as a density trace graph (yellow). For

both H and I, five hundred and sixty-seven MAGUK punctae were analyzed from 16

neuromasts of 4 dpf larvae. (J,K) Maximal-intensity projections from micrographs of a control

and a *sema7a$^{-/-}$* neuromast depict the postsynaptic aggregates marked by MAGUK (yellow).

The approximate hair cell basal region is outlined by dashed green circles. (L-O) Plots

quantitate the numbers of postsynaptic aggregates (L,M) and their areas (N,O), in each hair

cell across developmental stages. One hundred and fifty, 218, and 167 hair cells were analyzed

from 2 dpf, 3 dpf, and 4 dpf control larvae, respectively. Ninety-seven, 141, and 141 hair cells

were analyzed from 2 dpf, 3 dpf, and 4 dpf *sema7a$^{-/-}$* mutant larvae, respectively. Scale bars,

5 μm.

1263

Figure 6—figure supplement 1. ***sema7a*^{-/-} mutants display impaired synaptic assembly.**

(A-F) The plots quantitate the average numbers and areas of presynaptic aggregates from

control and *sema7a*^{-/-} neuromasts. Significant decreases in the numbers and areas of

presynaptic aggregates are observed across development. Three hundred and forty-one, 1229,

and 1286 presynaptic aggregates were analyzed from 2 dpf, 3 dpf, and 4 dpf control larvae,

respectively. Two hundred and thirty-five, 643, and 419 presynaptic aggregates were analyzed

from 2 dpf, 3 dpf, and 4 dpf *sema7a$^{-/-}$* mutant larvae, respectively. (G-L) The plots quantitate

the average numbers and areas of postsynaptic aggregates from control and *sema7a$^{-/-}$*

neuromasts. Significant decreases in the numbers and areas of postsynaptic aggregates

occurred during development. Three hundred and ninety-seven, 1243, and 1300 postsynaptic

aggregates were analyzed from 2 dpf, 3 dpf, and 4 dpf control larvae, respectively. Three

hundred and four, 651, and 451 postsynaptic aggregates were analyzed from 2 dpf, 3 dpf, and

4 dpf *sema7a$^{-/-}$* mutant larvae, respectively. Means ± SEMs; *** implies $p < 0.001$.

1302

 Figure 6—figure supplement 2. Morphologies of presynaptic and postsynaptic

assemblies and the distribution of postsynaptic aggregates across the sensory arbor

network. (A,B) The micrographs depict single presynaptic aggregate marked by CTBP in

control (A) and in sema7a-/- mutant (B) larvae at 4 dpf. (C,D) The micrographs depict single

postsynaptic aggregate marked by MAGUK in control (C) and in sema7a-/- mutant (D) larvae

at 4 dpf. (E) A maximal-intensity projection micrograph depicts the sensory arbor network

(magenta) and the enrichment of the postsynaptic assemblies (yellow, white arrowhead) near

the nodes (cyan arrows). (F) A schematic diagram of the sensory arbor network depicts the

enrichment of the MAGUK punctae near the node (cyan arrow). The three-dimensional space

surrounding each node was divided into concentric shells of equal volume to measure the

distribution of MAGUK-punctae density across the network.

1315 **Video 20. Postsynaptic assemblies are enriched in the loops and near the nodes of a**

1316 **sensory arbor network.** Through-focus scans of a 4 dpf neuromast depict the sensory arbor

1317 network (magenta) where the postsynaptic assemblies (yellow) are primarily located in the

1318 loops of the arbor network and are enriched near the nodes.

1319

Figure 7. Expression of diverse neural guidance cues in developing neuromasts. Analysis of single cell RNA-sequencing data (9) shows the expression of multiple neural guidance genes that are expressed across young hair cells (Y), mature hair cells (M), and support cells (SC). Dot sizes and colors signify the proportion of cells in each cluster that express a gene and the average strength of expression (log[(counts/10,000) +1]), respectively.