CAGEE: computational analysis of gene expression evolution

Jason Bertram[1,2,*], Ben Fulton[1,3], Jay Tourigny[1,4], Yadira Peña-Garcia[1], Leonie C. Moyle[1], and Matthew W. Hahn[1,4,*]

[1]Department of Biology, Indiana University, Bloomington, IN 47405, USA

[2]Department of Mathematics, Western University, London, ON N6A 5B7, Canada

[3]University Information Technology Services, Indiana University, Bloomington, IN 47405, USA

[4]Department of Computer Science, Indiana University, Bloomington, IN 47405, USA

*Corresponding authors: E-mail: jason.bertram@uwo.ca, mwh@indiana.edu

Abstract

Despite the increasing abundance of whole transcriptome data, few methods are available to analyze global gene expression across phylogenies. Here, we present a new software package (CAGEE) for inferring patterns of increases and decreases in gene expression across a phylogenetic tree, as well as the rate at which these changes occur. In contrast to previous methods that treat each gene independently, CAGEE can calculate genome-wide rates of gene expression, along with ancestral states for each gene. The statistical approach developed here makes it possible to infer lineage-specific shifts in rates of evolution across the genome, in addition to possible differences in rates among multiple tissues sampled from the same species. We demonstrate the accuracy and robustness of our method on simulated data, and apply it to a dataset of ovule gene expression collected from multiple self-compatible and self-incompatible species in the genus *Solanum* to test hypotheses about the evolutionary forces acting during mating system shifts. These comparisons allow us to highlight the power of CAGEE, demonstrating its utility for use in any empirical system and for the analysis of most morphological traits. Our software is available at https://github.com/hahnlab/CAGEE/.

Key words: RNA-seq; phylogenetic comparative methods; Brownian motion; Solanum

Introduction

Early studies of gene expression in single genes revealed widespread and frequent changes in the levels, timing, and breadth of expression across species (reviewed in Wray et al. 2003; Fay and Wittkopp 2008; Hill et al. 2021). Such changes in gene expression have been shown to be responsible for many differences between species, and may be a major driver of evolution (King and Wilson 1975). Advances in sequencing technologies (i.e. RNA-seq) have transformed research into gene expression, allowing researchers to cheaply and accurately measure transcript levels for every gene in a genome, in multiple tissues, and across several timepoints or conditions (Wang et al. 2009). There is now a flood of interest in applying RNA-seq to whole clades of organisms in order to identify the genetic changes and evolutionary forces driving species differences (e.g. Brawand et al. 2011; Meisel et al. 2012; Coolon et al. 2014; Harrison et al. 2015; Berthelot et al. 2018; Catalan et al. 2019; Blake et al. 2020; El Taher et al. 2021).

To better understand the importance of changes in gene expression, researchers must be able to characterize the mechanisms and modes by which gene expression evolves. Such work entails understanding the role of natural selection in driving species differences, the stages of development or the tissues that evolve most rapidly, as well as the environments most likely to generate changes in gene expression (Dunn et al. 2013; Hill et al. 2021; Price et al. 2022). Phylogenetic comparative methods enable the rigorous study of traits like gene expression across a species tree (Revell and Harmon 2022). These methods can be used for testing hypotheses about natural selection, the inference of ancestral states (allowing us to polarize the direction of changes), and the estimation of evolutionary rates. Multiple software packages are available that implement a wide variety of comparative methods (e.g. Pennell et al. 2014), including models specifically intended for studying gene expression across a tree (Bedford and Hartl 2009; Rohlfs et al. 2014; Rohlfs and Nielsen 2015; Catalán et al. 2019; Chen et al. 2019).

However, as far as we are aware, all existing comparative methods for analyzing gene expression implement fundamentally single-gene analyses. Each gene is

considered a separate trait, such the evolutionary parameters for each gene are estimated separately. Single-gene analyses can be used to identify tissue-specific or lineage-specific shifts in evolutionary rates, but their power is quite low (Beaulieu et al. 2012). As a result, identifying trends in evolution must be carried out *post hoc* by summing the number of genes found to be individually significant (e.g. Harrison et al. 2015; El Taher et al. 2021). This approach is less than ideal, especially when carrying out comparisons between branches of different lengths or between tissues with different average expression levels (both of which can result in differential statistical power).

Therefore, to better characterize the forces affecting gene expression evolution, we must be able to model effects shared along a lineage, experienced by many genes in the same tissue, or experienced by all genes found in the same environment. In this article, we present a genome-scale platform for the analysis of gene expression data that allows for such shared factors. Our software, CAGEE (Computational Analysis of Gene Expression Evolution), provides a robust set of methods for analyzing expression data across a species tree. CAGEE estimates ancestral states and rates, with rates shared by all or subsets of genes (single-gene analyses can also be carried out). We show that lineage-specific and tissue-specific (or condition-specific) rates can be accurately inferred, and provide principled statistical approaches for model selection. Our current implementation uses a bounded Brownian motion model—in order to account for the fact that expression levels cannot be negative—and assumes expression data are accurate, but the architecture and codebase will easily allow for future extensions that relax these and other assumptions.

New Approaches

We model gene expression evolution as a bounded Brownian motion (BBM) process on a known species tree (cf. Boucher and Démery 2016). Our model has a single bound: trait values must be greater than or equal to zero, since expression levels cannot be negative; there is no upper bound (Figure 1). Previous researchers have often modeled gene expression using an Ornstein-Uhlenbeck (OU) process (e.g. Bedford and Hartl 2009; Rohlfs et al. 2014; Rohlfs and Nielsen 2015; Chen et al. 2019),

a model that includes a force constraining traits about the mean. However, to our knowledge, the OU model has only been compared against an unbounded Brownian motion model (i.e. one that allows negative trait values), making fair comparisons difficult. In addition, OU models may be frequently and incorrectly favored over simpler models due to several biases (e.g. measurement error), especially when the number of tips in a tree is small (Pennell et al. 2015; Silvestro et al. 2015; Boucher and Démery 2016; Cooper et al. 2016; Catalán et al. 2019). Therefore, the initial version of our software models gene expression with the BBM process, which naturally bounds possible values without invoking an additional constraining force.

Let $E_{ij} \geq 0$ be the expression level of gene i in species j. We assume that log-transformed expression $X_{ij} = \ln(E_{ij} + e_{\min})$ evolves as a Brownian motion process with variance σ^2 per unit time, where $e_{\min}$ is a small offset (constant across genes and species) that prevents X_{ij} from taking infinite values if measured values of E_{ij} are zero. We log-transform before assuming Brownian motion because we expect the variance in the evolutionary process to scale with expression level. Assuming that E_{ij} is itself Brownian would unrealistically assume that the rate of evolution is constant across expression levels, even though expression levels vary by many orders of magnitude. We impose a reflecting lower boundary at $x_{\min} = \ln(e_{\min})$, meaning that the Brownian walk immediately bounces back if it reaches $x_{\min}$. Trait values can therefore effectively never reach zero, our theoretical lower bound (Figure 1).

The second major feature of our model (as implemented in CAGEE) is that many genes can share the evolutionary rate parameter, σ^2. This rate may be shared among genes expressed in the same tissue or sample, among genes located on the same chromosome, or among genes evolving along the same lineage of the phylogenetic tree. The simplest model allows σ^2 to be shared among all genes, providing an average rate of evolution across the genome and over time. We explain this model briefly here, with more detail provided in the Materials and Methods.

CAGEE infers the most likely value(s) of σ^2 consistent with an ultrametric tree, T, and a set $E_{\{ij\}}$ of measured expression values at the tips of the tree; i.e. it maximizes

the likelihood $L(\sigma^2|E_{\{ij\}}, T)$. Each gene is assumed to evolve independently, and so the likelihood for each gene $L_i(\sigma^2|E_{i\{j\}}, T)$ is computed independently. The overall likelihood is obtained as the product $L(\sigma^2|E_{\{ij\}}, T) = \Pi_i L_i(\sigma^2|E_{i\{j\}}, T)$ across genes. The likelihood for each gene $L_i(\sigma^2|E_{i\{j\}}, T)$ is computed using the pruning algorithm (Felsenstein 1973). The key ingredient needed to apply the pruning algorithm is the transition probability density $p(x_t|x_{t_0}) = \Pr[X(t) = x_t|X(t_0) = x_{t_0}]$ for log-expression at time t conditional on having log-expression x_{t_0} at time t_0 along a lineage. CAGEE computes the transition density by solving the standard Brownian diffusion equation with reflecting boundary conditions (Materials and Methods). The transition density is used to propagate expression probabilities along the tree: if the probability density of log-expression at time t_0 is $f(x_{t_0})$, then the probability density at time t on the same lineage is $f(x_t) = \int p(x_t|x_{t_0})f(x_{t_0})dx_{t_0}$. At each tip the probability density $f(x_{t_0})$ is a delta function centered at the corresponding measured value of X_{ij}.

Starting with the known tip distributions, the pruning algorithm propagates back to the tips' parent nodes. The distribution at the parent node is then the product of the two backward-propagated child node distributions. Proceeding iteratively across the tree, we ultimately obtain the gene-specific probability density for expression value at the root $f_i(x_R)$. Viewed as a likelihood for σ^2, $f_i(x_R)$ is the gene-specific likelihood conditional on the unknown ancestral root value; i.e. $f_i(x_R) = L_i(\sigma^2|E_{i\{j\}}, T, x_R)$. Therefore, we integrate over all possible x_R to obtain,

$$L_i(\sigma^2|E_{i\{j\}}, T) = \int L_i(\sigma^2|E_{i\{j\}}, T, x_R)\rho(x_R)dx_R, \qquad (1)$$

where $\rho(x_R)$ is the prior distribution for the root value of a randomly selected gene.

The default prior $\rho(x_R)$ is assumed to be a gamma distribution with $k = 0.375$ and $\theta = 1600$, though this distribution can also be set by the user in CAGEE. This choice is based on estimated expression distributions across genes in individual species, which we take as our baseline for the ancestral distribution. CAGEE uses the Nelder-Mead simplex method to find the optimal value(s) of σ^2.

Results

Using CAGEE

The required inputs for CAGEE are a Newick-formatted, rooted, ultrametric tree (with branch lengths) and a tab-delimited data file containing the expression levels of all species or taxa being studied. The data file can consist of data on one gene/transcript or thousands of different genes. The first line of the data file should contain the species' names (matching those used in the Newick tree). In addition, headers for gene names, gene descriptions, and sample IDs (see next section for an explanation of "samples" in CAGEE) can be used. Subsequent lines each correspond to a single gene and contain expression levels for each species. Examples of Newick trees and corresponding data files can be found in the online user manual (https://github.com/hahnlab/CAGEE/docs/manual/cagee_manual.md).

We expect that CAGEE will most often be used to calculate the following outputs: one or more σ^2 values, ancestral states at each internal node (including 95% credible intervals around these states), and the final likelihood associated with a model. However, users do not have to search for σ^2: if a value for this parameter is specified, then the output of CAGEE will just be the ancestral states and a likelihood. In addition to the raw outputs provided in multiple formats (both tab-delimited files and NEXUS-formatted files), CAGEE computes basic statistics about changes in expression levels by comparing values at parent and child nodes. Summaries of these inferred changes for every gene and for every branch of the tree are output, so that the evolutionary history of gene expression changes in every gene are accessible to users. To avoid over-interpretation of small changes in inferred expression levels—especially when there is uncertainty in ancestral states—CAGEE will also compare the credible intervals at parent and child nodes to note if a change is "credible" (i.e. the intervals do not overlap). Such changes are annotated as such in the output.

There are multiple options available for running CAGEE. Users who can take advantage of multiple threads can specify the number to use on the command line. Complex models can also take a long time to converge; by default, CAGEE runs a

maximum of 300 iterations of the Nelder-Mead search, but users can increase this number in subsequent runs if the likelihood is still improving when the limit is hit. As mentioned above, the default prior distribution for the root state is a gamma distribution with $k = 0.375$ and $\theta = 1600$. This distribution can also be specified by the user if desired. Information on how to run more complex evolutionary models, beyond a single σ^2, is given in the next section.

Estimating evolutionary rates in CAGEE

We tested CAGEE's ability to accurately estimate σ^2 by varying both this rate parameter and the number of genes used for inference. We simulated different single values of σ^2 across a tree with constant branch lengths (Supplementary Figure 1) using the simulation tool available within CAGEE. (Note that the total amount of evolution in a tree is determined by the product $\sigma^2 \cdot t$, such that changes in branch lengths will have an effect commensurate with changes in σ^2.) Figure 2 shows the average error associated with estimates of different σ^2 values and using different numbers of genes within each dataset. As can be seen, the error across all parameter values and dataset sizes is quite small (generally less than 2.5%), and is less variable for larger dataset sizes. Fortunately, we expect that most empirical datasets will contain closer to 10,000 genes than 1,000 genes. The results in Figure 2 are for an ancestral state vector of length *N*=200 (the default setting in CAGEE; Materials and Methods); we also estimated σ^2 when allowing the ancestral state vector to have length *N*=500 (Supplementary Figure 2). There appears to be minimal gain from increasing the resolution in this vector, though the computational time is greatly increased (similar to results in Boucher and Démery 2016). Although we have not shown it here, we also expect that the accuracy of σ^2 will be greater for trees with larger numbers of tips (cf. O'Meara et al. 2006). Overall, inferences of σ^2 should be quite accurate when a single rate parameter is shared across the tree and across all genes and lineages.

Variation in the rate of expression can be accommodated by CAGEE in a number of ways, using multi-rate σ^2 models. One type of model allows users to specify that their data come from different "samples": these samples can represent tissues, conditions,

timepoints, and even subsets of the genome (e.g. the X chromosome, or a specific functional class of genes). In the input data file, the "SAMPLETYPE" column is used to indicate which sample each gene is a member of; a separate σ^2 value will be calculated for each sample or set of samples (these values are assumed to be shared among all lineages in the tree). Specifying more than one sample means that an individual gene or transcript name can be used more than once (i.e. once for each sample), but there is no requirement that genes are measured in each sample. For instance, assigning all autosomal genes to sample 1 and all X-linked genes to sample 2 would not permit for any overlap in gene assignment, but is perfectly allowable in CAGEE.

Each additional sample requires another σ^2 parameter to be estimated, and often researchers would like to know if fitting this extra parameter is justified by the data. Under standard information-theoretic criteria (Burnham and Anderson 2002), twice the difference in log-likelihoods between nested models should be χ^2-distributed with degrees of freedom equal to the difference in the number of parameters between models. To test this expectation, we simulated 1000 datasets with a single σ^2 value, but fit models with two σ^2 values (assigning 1000 genes to two equal-sized samples at random; the relative size of the samples should not affect the false positive rate). As anticipated, the results fit a χ^2 distribution with one degree of freedom, with 4.4% of datasets having a difference in 2*log-likelihood greater than 3.84 (5% are expected by chance). This indicates that standard statistical procedures should adequately control the false positive rate when fitting multi-sample σ^2 models.

CAGEE also allows models in which σ^2 varies across branches of the species tree. It does so by fitting separate σ^2 parameters for different parts of the tree. On the command line, CAGEE enables users to specify how multiple σ^2 parameters should be assigned to branches. For n taxa, from 1 to $2n$-2 parameters can be specified, and branches can be grouped together in any way. For instance, a two-parameter model can have all branches that share a rate adjacent to one another in the tree (Supplementary Figure 3A) or spread out across the tree (Supplementary Figure 3B). Similar to the analyses carried out above for the false positive rate associated with

multiple samples, we simulated data with a single σ^2 value and then fit models with multiple σ^2 parameters. Regardless of how we distributed the two rate classes across the tree we observed good control of the false positive rate: 4.5% and 5.4% of 1000 simulated datasets were significant at the P=0.05 level (for the trees shown in Supplementary Figures 3A and 3B, respectively). More limited simulations also showed that we could accurately estimate multiple σ^2 parameters when the data were simulated with multiple rates (Supplementary Table 1). Together, our results suggest that we can estimate multiple types of multi-rate models, and can accurately control the false positive rate when doing so.

Analysis of wild tomato transcriptome data

To demonstrate the utility of CAGEE in an empirical system, we analyzed data from a clade that includes domesticated tomato, *Solanum lycopersicum*. This dataset contains gene expression levels in unfertilized ovules from the flowers of six species, one of which (*S. pennellii*) has two different populations represented (Figure 3). There are 14,556 genes with expression levels measured in all seven accessions. RNA-seq data for five of the seven accessions have been published previously (Moyle et al. 2021; Hibbins and Hahn 2021), while two others are presented here for the first time (Materials and Methods). Note, however, that all data were collected from all samples at the same time (Materials and Methods).

Most species within the tomato clade are self-incompatible (SI), the ancestral state in the family Solanaceae (Igić et al. 2006). Self-incompatibility means that plants must outcross in order to successfully fertilize ovules. However, self-compatibility (SC) has evolved multiple times both within the Solanaceae and within the genus *Solanum* (Goldberg et al. 2010; Bedinger et al. 2011). Self-compatible individuals are able to successfully fertilize ovules using their own pollen, though many also still outcross (Whitehead et al 2018; including in *Solanum*: Vosters et al. 2014 and references therein). Importantly, we have *a priori* expectations about the rate at which reproductive traits—including ovule gene expression—might evolve between groups with different mating systems. Due to conflict within and between the sexes, it is generally expected

that reproductive traits in species that outcross more (i.e. SI taxa) should evolve more rapidly than in species that inbreed more (i.e. SC taxa; Clark et al. 2006). Such patterns are found in some analyses of the rate of protein evolution (e.g. Gossmann et al. 2016; Harrison et al. 2019), but are equivocal in other comparisons (e.g. Gossmann et al. 2014, Moyle et al. 2021). These complex patterns might reflect additional effects that also accompany mating system shifts; for instance, such shifts often lead to reductions in effective population size in more selfing lineages (Charlesworth and Wright 2001). Mating system shifts could also alter global patterns of molecular evolution (including gene expression) by changing the strength and pattern of purifying selection, as morphological changes often accompany mating system changes. The exact effect of shifts in mating system on molecular evolution remains an open question.

The *Solanum* species sampled here represent two independent transitions from SI to SC, with one of the transitions (in accession *S. pennellii* LA0716) occurring recently enough that different populations within this species have different incompatibility systems (Figure 3). We therefore fit a series of nested models within CAGEE to test two related hypotheses about ovule gene expression evolution. First, we would like to know whether the rate of evolution of ovule gene expression is different in SI species than in SC species. Second, given the recent transition to SC within accession *S. pennellii* LA0716, we wanted to know if it shows a pattern of evolution more similar to SI or to SC species. In total, we fit four separate evolutionary models (Table 1; Figure 3). Model A has a single rate parameter for the entire tree. Model B has two rate parameters, one for SI species and one for SC species. This model assigns the branch leading to *S. pennellii* LA0716 as SC. Model C also has two rate parameters, one for SI and one for SC, but assigns *S. pennellii* LA0716 as SI. Model D has three rate parameters: one for SI species, one for longer-term SC species, and one for *S. pennellii* LA0716.

Estimated results from the different models are shown in Table 1. Model A has a worse fit than any other model, with a single σ^2 value of 0.102. For context, this value means that the bounded Brownian motion process the data are fit to has a variance of

0.102 per million years (of log-transformed expression values). This is the average rate across all 14,556 genes and across all branches of the tree. In contrast to a single-rate model, both models B and C are significantly better fits to the data. Contrary to some hypotheses, both models find that SI lineages (σ_1^2) have a lower rate of evolution than SC lineages (σ_2^2; Table 1). There is also a difference between the models, with model C (the one in which *S. pennellii* LA0716 shares a rate with SI species) fitting significantly better. To further examine the evolution of *S. pennellii* LA0716, model D fits a three-parameter model, with this lineage assigned its own rate of evolution. This model is a significantly better fit than model C ($P<0.00001$; χ^2 test with 1 degree of freedom), and demonstrates that *S. pennellii* LA0716 has a rate of evolution (σ_3^2 in Table 1) that is slightly *lower* than SI species. This highly similar rate to SI species implies that it has only recently transitioned to self-compatibility, which is consistent with previous inferences about the timing of transition to SC in this particular accession (e.g. Rick and Tanksley 1981).

CAGEE also allows users to infer the number and direction of changes in gene expression levels along each branch of the tree. Figure 3 reports the number of genes that had "credible" increases and decreases in expression level under model D. Credible changes require that the credible intervals around states at parent and daughter nodes do not overlap, in order to account for uncertainty in our inferences. However, because of this, fewer credible changes will be inferred deeper in the tree, where credible intervals get wider. Therefore, while inferences about the identity of the genes changing along each branch is greatly strengthened by using credible changes (these genes are noted in the raw output from CAGEE), the absolute numbers of credible changes cannot be compared across branches, except for sister branches of equal length. For completeness, we show the total numbers of increases and decreases of gene expression in Supplementary Figure 4; as expected, these total numbers are more uniformly distributed across older and younger branches.

Discussion

Here, we have developed a new software package that enables the estimation of rates of gene expression evolution across a tree, CAGEE. Gene expression levels are much like many other continuous traits, and multiple papers have introduced phylogenetic comparative methods for studying gene expression (Bedford and Hartl 2009; Rohlfs et al. 2014; Rohlfs and Nielsen 2015; Catalán et al. 2019; Chen et al. 2019). However, as far as we are aware none of these methods allows genes to share evolutionary parameters, which precludes the analysis of genome-wide trends, either along the branches of a tree or between tissues/samples/conditions. To overcome this limitation, CAGEE calculates the likelihood of the data using the pruning algorithm (Felsenstein 1973) to facilitate the sharing of evolutionary parameters along branches of the species tree, providing more statistical power to test evolutionary hypotheses. Fortunately, we were able to take advantage of much of the codebase of our existing software, CAFE (Hahn et al. 2005; De Bie et al. 2006; Hahn et al. 2007; Han et al. 2013; Mendes et al. 2020), which implements the pruning algorithm for the analysis of gene family sizes across a tree. While gene expression levels and gene family sizes differ in the type of data they represent (continuous vs. discrete) and their underlying evolutionary models (bounded Brownian motion vs. birth-death), many of the required likelihood calculations and software components are the same.

An important thing to consider for the input to CAGEE is the normalization used to make gene expression levels comparable across species. The data from wild tomatoes used here was normalized using TPM (transcripts per million; Wagner et al. 2012); other published datasets also use this normalization (Berthelot et al. 2018; Chen et al. 2019; El Taher et al. 2021). However, multiple other normalizations have also been used in comparative analyses, including RPKM (Brawand et al. 2011), FPKM (Catalán et al. 2019), and both TMM and CPM (Blake et al. 2020). Each normalization approach has its advantages and disadvantages, and we cannot yet recommend one specific approach as input to CAGEE. The normalization method used will likely depend on the conditions under which samples are collected: if all species can be raised simultaneously in a greenhouse, vivarium, or growth chamber, we expect many fewer

batch effects than samples collected from the field, which will therefore necessitate different normalizations. However, even animals raised in a common environment—but fed different diets—can show many differences in gene expression not due to heritable change (e.g. Somel et al. 2008). Conversely, many between-sample normalization approaches (e.g. TMM, trimmed mean of M values; Robinson and Oshlack 2010) make the assumption that differences in gene expression between samples are rare. While such normalization is sensible in the context of testing for differential expression between samples from the same species, for a set of species that have been evolving independently for millions of years this is likely not an appropriate assumption.

CAGEE currently has multiple limitations, both in the available models that can be applied and in the types of data that can be analyzed. As mentioned earlier, many researchers have modeled gene expression using an OU process (Bedford and Hartl 2009; Rohlfs et al. 2014; Chen et al. 2019). Although OU models may be artifactually preferred over unbounded Brownian motion models due to a number of non-biological factors (see discussion in "New Approaches" above), it would still be helpful to be able to compare such a model to the bounded Brownian motion model used here. However, fitting such a model to genome-wide data is non-trivial: each gene must have its own mean expression value (μ), but possibly shared constraint parameters (α) across genes. We have the goal of implementing such a model in the near future, as well as other models commonly used in comparative methods research (e.g. Landis and Schraiber 2017; Boucher et al. 2018). Beyond the evolutionary model applied to any dataset, there are multiple additional sources of variation that could be modeled. For instance, we have previously accounted for measurement error in a likelihood framework, using an empirically parameterized error model (Han et al. 2013). In addition, not all genes necessarily share the same average rate of evolution; gamma-distributed rate categories can be used to model this variation among genes (cf. Ames et al. 2012; Mendes et al. 2020). Finally, the gene tree discordance found in many phylogenomic datasets implies that complex traits (such as expression levels) will also be controlled by discordant gene trees (Hahn and Nakhleh 2016; Hibbins and Hahn 2021). This underlying discordance can cause evolutionary rates to be overestimated

(Mendes et al. 2018), and should be taken into account when seeking accurate parameter estimates (see discussion of wild tomato data below). Our goal is to include methods for dealing with all these sources of variation in future versions of CAGEE.

In terms of the types of data that can be analyzed, at present CAGEE is limited to positive, continuously varying traits (i.e. the BBM model). However, we also envision different ways to represent and model gene expression data, including as a ratio (e.g. male/female expression). Such a ratio, after log2-transformation, would be most appropriately modeled by an unbounded Brownian motion model since both negative and positive values are possible. This and other data types will be supported in future releases. Moreover, CAGEE does not have to analyze whole-genome or even molecular data: it can be applied to any single trait for which the BBM model is appropriate, even morphological traits. One intriguing application of CAGEE could be to suites of morphological traits that are hypothesized to share a common evolutionary rate parameter. If, for instance, there is a shift in body plan along some lineages, then multiple traits may all increase or decrease their rate of evolution at once, and CAGEE can be used to estimate these shared parameters. Even in the context of single-trait analyses, the pruning algorithm has been hailed as a solution for large-scale comparative analyses (Freckleton 2012). Importantly, the number of branches in a rooted, bifurcating tree with n tips is $2n$-2, so that the number of calculations scales linearly with the number of species. This makes the pruning algorithm ideal for comparative datasets with large numbers of taxa (e.g. Hahn et al. 2005; FitzJohn 2012; Hiscott et al. 2016; Caetano and Harmon 2018; Mitov et al. 2020).

The analysis of data from a clade of wild tomatoes revealed a possibly unexpected result: the rate of ovule gene expression evolution among self-compatible (SC) species is twice as high as the rate among self-incompatible (SI) species (Table 1). This finding is contrary to some prior expectations—informed by research focused on male-female interactions, especially between interacting proteins in the reproductive tract (e.g. Swanson and Vacquier 2002; Clark et al. 2006)— that suggest that lineages might experience slower evolution after transitioning to self-compatibility. However, it is possible that global gene expression levels do not evolve in the same sort of tit-for-tat manner as interacting protein sequences, such that increases/decreases in male-

expressed genes are not matched by increases/decreases in interacting female-expressed genes (or vice versa). Alternatively, only a very small subset of genes may evolve in this manner. Indeed, even prior studies comparing protein evolution have failed to find clear evidence of slower global evolutionary rates in more inbreeding species (e.g. Wong 2011). One caveat to the observed rate differences in our data is that underlying gene tree discordance, whether due to incomplete lineage sorting or introgression, can lead to artifactually higher rate estimates (Mendes et al. 2018; Hibbins and Hahn 2021). However, there is in fact less discordance among the SC lineages sampled here (Pease et al. 2016), which is the reverse of the pattern that would be required to explain our results.

If not due to underlying bias in our estimates, these findings still raise the question: why is ovule gene expression evolving more rapidly in SC than SI species? One possibility is that this increased rate is due to a relaxation of selection in SC species, possibly because genes involved in male-female interactions are no longer needed. If this were the case, we might expect to see a general decrease in expression levels in SC species; however, there appears to be no consistent directionality to the changes along SC branches (Figure 3, Supplementary Figure 4). Instead, an alternative hypothesis is that transitions to SC involve adaptation to new optima of ovule gene expression, compared to SI species that tend to maintain ancestral optima. For example, transitions to SC might favor greater investment in fewer ovules, because self-compatibility decreases the probability that each ovule within a flower will go unfertilized—an otherwise wasted investment under conditions (like SI) where receiving sufficient compatible pollen to fertilize each ovule is less predictable (Burd et al. 2009). The nature of these new optima might be even more complex, as traits like ovule size and number can vary with multiple reproductive and ecological conditions, and often trade-off with each other (Greenway and Harder 2007). Of the species examine here, for example, two SC lineages (*S. pimpinellifolium*, and *S. lycopersicon*—domesticated tomato) have significantly larger seeds than most of the SI lineages and SC *S. pennellii* (unpubl. data). Some of these hypotheses could be evaluated with matching gene expression data from other (non-ovule) reproductive tissues. Analyses including pollen in the same SI and SC lineages, and/or data addressing alternative constraints and

conditions shaping ovule evolution including ovule size and number (e.g. Mione and Anderson 1992), would be useful in teasing apart these hypotheses.

Material and Methods

Bounded Brownian motion model of expression evolution

The probability density of expression, $p(x,t)$, at time t for evolutionary trajectories following a Brownian motion process starting at value x_{t_0} at time t_0 is governed by the diffusion equation

$$\frac{\partial p(x,t)}{\partial t} = \frac{\sigma^2}{2}\frac{\partial^2 p(x,t)}{\partial x^2},\qquad(2)$$

with initial condition $p(x,t_0) = \delta(x - x_{t_0})$ where δ is the Dirac delta function. The reflective boundary condition at $x = x_{\min}$ implies that the probability fluxes into and out of the boundary are balanced, imposing the boundary condition

$$\frac{\partial p(x=x_{\min},t)}{\partial x} = 0.\qquad(3)$$

Note that $p(x,t)$ is identical to the transition density $p(x_t|x_{t_0})$.

Without the reflecting boundary, $p(x,t) \propto e^{-(x-x_{t_0})^2/2\sigma^2(t-t_0)}$ is a normal distribution with variance $\sigma^2(t - t_0)$. The variance therefore scales linearly with elapsed time, $t - t_0$. With the reflecting boundary, $p(x,t)$ is the sum of this spreading normal and its mirror image centered at $2x_{\min} - x_{t_0}$. The analytical solution to this bounded process is helpful for understanding the behavior of $p(x,t)$, but is not used in CAGEE. In anticipation of implementing additional (and possibly more complicated) processes into CAGEE, we instead solve Eq. (2) numerically using the approach described in Boucher and Démery (2016). Briefly, the continuous diffusion equation is converted into a matrix equation by discretizing expression values into N equal bins of width $\delta = \frac{x_{\max}-x_{\min}}{N-1}$. Following Boucher and Démery (2016), we have used a default *N*=200, but this number can be set by the user (see Results). This approach gives

$$\frac{\partial P(t)}{\partial t} = \frac{\sigma^2}{2\delta^2}M \cdot P(t),\qquad(4)$$

where $P(t)$ is the vector obtained by discretizing $p(x,t)$ and $x_{\max}$ is the largest expression value accounted for. The matrix M is tridiagonal with -2 on the diagonal

except at the first and last diagonal entries which are -1. The sub- and supra-diagonal entries are 1. This equation has the matrix exponential solution

$$P(t) = \exp\left(\frac{\sigma^2(t-t_0)}{2\delta^2}M\right), \tag{5}$$

which is evaluated by diagonalizing M.

Implementation of CAGEE

      CAGEE is written in C++ and is compatible with the C++11 standard. A comprehensive manual and extensive unit tests facilitate further code development and maintenance. CAGEE is organized into modular components. A *clade* class, with references to a parent clade and any number of descendant clades, represents a tree structure, and a *gene_transcript* class represents the expression levels observed in the various species. These two classes comprise the fundamental data structures upon which CAGEE performs its analysis (Supplementary Figure 5).

      Calculations are carried out by additional classes. The *optimizer* class has the responsibility of determining the σ^2 value with the highest likelihood, by comparing the likelihood of candidate values and searching the likelihood surface using the Nelder-Mead optimization algorithm. The work of computing the likelihood of a given σ^2 value is performed by a subclass of the *model* class, which for now is limited to a single *Base* model (allowing for further development in the future). After appropriate estimated values are found, the *transcript_reconstructor* class builds a possible set of transcript values for the entire tree (Supplementary Figure 5).

      Performing the likelihood calculations requires extensive matrix operations; it is recommended (though not required) that these be passed off to a specialized library such as Intel's MKL or Nvidia's CUBLAS. If no external library is available, CAGEE will carry out these calculations (slowly) by itself. Creating the diffusion matrix (M) requires calculation of eigenvalues and eigenvectors, and is computationally expensive. This work is performed by the Eigen linear algebra library (https://eigen.tuxfamily.org); various internal data structures also take advantage of Eigen classes. To enable faster searching, the matrix for an ancestral state vector of length 200 (the default in CAGEE)

has been pre-computed and is included with CAGEE. Users who wish to use vectors of different lengths can specify this as an option.

Unit-testing is performed using the Doctest testing framework (https://github.com/doctest/doctest). At the time of writing more than 200 unit tests had been created, comprising more than 1200 individual assertions. For complex logging and debugging cases, CAGEE uses the EasyLogging framework (https://github.com/amrayn/easyloggingpp). C++ development is always made easier by using the Boost C++ libraries (https://www.boost.org/), so we include them as well in CAGEE.

RNA-seq data from wild tomatoes

We briefly describe here the data collected from seven accessions of wild tomatoes (*S. lycopersicum* LA3475, *S. chmielewskii* LA1316, *S. pimpinellifolium* LA1589, *S. habrochaites* LA1777, *S. chilense* LA4117A, *S. pennellii* LA3778, and *S. pennellii* LA0716; all accession ID numbers from tgrc.ucdavis.edu). Further details are given in Moyle et al. (2021). Ovule RNA-seq was performed on between one to four (usually three) biological replicates (individual plants) from each accession. Plants were germinated from seed, and cultivated until flowering. For each replicate individual, ovules were dissected from mature, unpollinated flowers, flash frozen, and maintained at -80C until extraction. For each individual, all ovule collections were pooled into a single sample prior to library construction and sequencing on an Illumina HiSeq 2000. Reads were mapped against the tomato reference genome (ITAG 2.4) and the number of reads mapped onto genic regions were estimated with featureCounts (Liao et al., 2014). We normalized the read counts from each library by calculating TPM (transcripts per million; Wagner et al. 2012) and then calculated the mean normalized read counts across all samples (individuals) within each accession. These means per accession were used as input to CAGEE.

To construct a species tree for use with CAGEE, we started with the topology given in Pease et al. (2016). Specifically, we used the tree found in the supplementary file Pease_etal_TomatoPhylo_RAxMLConcatTree_no1360_Fig2A.nwk, and pruned it to include only the accessions in our study using the software ETE (Huerta-Cepas et al.

2016). Using the "extend" method found in ETE, we converted this tree to ultrametric (same root-to-tip distance for all taxa). Setting the root age to 2.48 million years ago (following Pease et al. 2016), we were able to express all branches in millions of years.

Acknowledgements

We thank Mark Hibbins for assistance with the tomato phylogeny, Matthew Gibson for putting together the tomato gene expression data, and especially Dan Vanderpool for invaluable help in the initial development of CAGEE. This work was supported by National Science Foundation grants DEB-1856469 to L.C.M. and DBI-2146866 to M.W.H.

Data Availability

Raw reads for each sample library are available at NCBI BioProject PRJNA714065. The CAGEE software is available at https://github.com/hahnlab/CAGEE.

References

Ames RM, Money D, Ghatge VP, Whelan S, Lovell SC. 2012. Determining the evolutionary history of gene families. Bioinformatics 28:48-55.

Beaulieu JM, Jhwueng DC, Boettiger C, O'Meara BC. 2012. Modeling stabilizing selection: expanding the Ornstein–Uhlenbeck model of adaptive evolution. Evolution 66:2369-2383.

Bedford T, Hartl DL. 2009. Optimization of gene expression by natural selection. Proceedings of the National Academy of Sciences 106:1133-1138.

Bedinger PA, Chetelat RT, McClure B, Moyle LC, Rose JK, Stack SM, van der Knaap E, Baek YS, Lopez-Casado G, Covey PA. 2011. Interspecific reproductive barriers in the tomato clade: opportunities to decipher mechanisms of reproductive isolation. Sexual Plant Reproduction 24:171-187.

Berthelot C, Villar D, Horvath JE, Odom DT, Flicek P. 2018. Complexity and conservation of regulatory landscapes underlie evolutionary resilience of mammalian gene expression. Nature Ecology & Evolution 2:152-163.

Blake LE, Roux J, Hernando-Herraez I, Banovich NE, Perez RG, Hsiao CJ, Eres I, Cuevas C, Marques-Bonet T, Gilad Y. 2020. A comparison of gene expression and DNA methylation patterns across tissues and species. Genome Research 30:250-262.

Boucher FC, Démery V. 2016. Inferring bounded evolution in phenotypic characters from phylogenetic comparative data. Systematic Biology 65:651-661.

Boucher FC, Démery V, Conti E, Harmon LJ, Uyeda J. 2018. A general model for estimating macroevolutionary landscapes. Systematic Biology 67:304-319.

Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, Weier M, Liechti A, Aximu-Petri A, Kircher M, et al. 2011. The evolution of gene expression levels in mammalian organs. Nature 478:343-348.

Burd M, Ashman T-L, Campbell DR, Dudash MR, Johnston MO, Knight TM, Mazer SJ, Mitchell RJ, Steets JA, Vamosi JC. 2009. Ovule number per flower in a world of unpredictable pollination. American Journal of Botany 96:1159-1167.

Burnham KP, Anderson DR. 2002. Model selection and multimodel inference: A practical information-theoretic approach. New York: Springer.

Caetano DS, Harmon LJ. 2018. Estimating correlated rates of trait evolution with uncertainty. Systematic Biology 68:412-429.

Catalán A, Briscoe AD, Höhna S. 2019. Drift and directional selection are the evolutionary forces driving gene expression divergence in eye and brain tissue of *Heliconius* butterflies. Genetics 213:581-594.

Charlesworth D, Wright SI. 2001. Breeding systems and genome evolution. Current Opinion in Genetics & Development 11:685-690.

Chen J, Swofford R, Johnson J, Cummings BB, Rogel N, Lindblad-Toh K, Haerty W, Di Palma F, Regev A. 2019. A quantitative framework for characterizing the evolutionary history of mammalian gene expression. Genome Research 29:53-63.

Clark NL, Aagaard JE, Swanson WJ. 2006. Evolution of reproductive proteins from animals and plants. Reproduction 131:11-22.

Coolon JD, McManus CJ, Stevenson KR, Graveley BR, Wittkopp PJ. 2014. Tempo and mode of regulatory evolution in *Drosophila*. Genome Research 24:797-808.

Cooper N, Thomas GH, Venditti C, Meade A, Freckleton RP. 2016. A cautionary note on the use of Ornstein Uhlenbeck models in macroevolutionary studies. Biological Journal of the Linnean Society 118:64-77.

De Bie T, Demuth JP, Cristianini N, Hahn MW. 2006. CAFE: a computational tool for the study of gene family evolution. Bioinformatics 22:1269-1271.

Dunn CW, Luo X, Wu Z. 2013. Phylogenetic analysis of gene expression. Integrative and Comparative Biology 53:847-856.

El Taher A, Böhne A, Boileau N, Ronco F, Indermaur A, Widmer L, Salzburger W. 2021. Gene expression dynamics during rapid organismal diversification in African cichlid fishes. Nature Ecology & Evolution 5:243-250.

Fay JC, Wittkopp PJ. 2008. Evaluating the role of natural selection in the evolution of gene regulation. Heredity 100:191-199.

Felsenstein J. 1973. Maximum likelihood and minimum-steps methods for estimating evolutionary trees from data on discrete characters. Systematic Biology 22:240-249.

FitzJohn RG. 2012. Diversitree: comparative phylogenetic analyses of diversification in R. Methods in Ecology and Evolution 3:1084-1092.

Freckleton RP. 2012. Fast likelihood calculations for comparative analyses. Methods in Ecology and Evolution 3:940-947.

Goldberg EE, Kohn JR, Lande R, Robertson KA, Smith SA, Igić B. 2010. Species selection maintains self-incompatibility. Science 330:493-495.

Gossmann TI, Saleh D, Schmid MW, Spence MA, Schmid KJ. 2016. Transcriptomes of plant gametophytes have a higher proportion of rapidly evolving and young genes than sporophytes. Molecular Biology and Evolution 33:1669-1678.

Gossmann TI, Schmid MW, Grossniklaus U, Schmid KJ. 2014. Selection-driven evolution of sex-biased genes is consistent with sexual selection in *Arabidopsis thaliana*. Molecular Biology and Evolution 31:574-583.

Greenway CA, Harder LD. 2007. Variation in ovule and seed size and associated size–number trade-offs in angiosperms. American Journal of Botany 94:840-846.

Hahn MW, De Bie T, Stajich JE, Nguyen C, Cristianini N. 2005. Estimating the tempo and mode of gene family evolution from comparative genomic data. Genome Research 15:1153-1160.

Hahn MW, Demuth JP, Han S-G. 2007. Accelerated rate of gene gain and loss in primates. Genetics 177:1941-1949.

Hahn MW, Nakhleh L. 2016. Irrational exuberance for resolved species trees. Evolution 70:7-17.

Han MV, Thomas GWC, Lugo-Martinez J, Hahn MW. 2013. Estimating gene gain and loss rates in the presence of error in genome assembly and annotation using CAFE 3. Molecular Biology and Evolution 30:1987-1997.

Harrison MC, Mallon EB, Twell D, Hammond RL. 2019. Deleterious mutation accumulation in *Arabidopsis thaliana* pollen genes: a role for a recent relaxation of selection. Genome Biology and Evolution 11:1939-1951.

Harrison PW, Wright AE, Zimmer F, Dean R, Montgomery SH, Pointer MA, Mank JE. 2015. Sexual selection drives evolution and rapid turnover of male gene expression. Proceedings of the National Academy of Sciences 112:4393-4398.

Hibbins MS, Hahn MW. 2021. The effects of introgression across thousands of quantitative traits revealed by gene expression in wild tomatoes. PLoS Genetics 17:e1009892.

Hill MS, Zande PV, Wittkopp PJ. 2021. Molecular and evolutionary processes generating variation in gene expression. Nature Reviews Genetics 22:203-215.

Hiscott G, Fox C, Parry M, Bryant D. 2016. Efficient recycled algorithms for quantitative trait models on phylogenies. Genome Biology and Evolution 8:1338-1350.

Huerta-Cepas J, Serra F, Bork P. 2016. ETE 3: reconstruction, analysis, and visualization of phylogenomic data. Molecular Biology and Evolution 33:1635-1638.

Igić B, Bohs L, Kohn JR. 2006. Ancient polymorphism reveals unidirectional breeding system shifts. Proceedings of the National Academy of Sciences 103:1359-1363.

King M-C, Wilson AC. 1975. Evolution at two levels in humans and chimpanzees. Science 188:107-116.

Landis MJ, Schraiber JG. 2017. Pulsed evolution shaped modern vertebrate body sizes. Proceedings of the National Academy of Sciences 114:13224-13229.

Liao Y, Smyth GK, Shi W. 2013. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics 30:923-930.

Meisel RP, Malone JH, Clark AG. 2012. Disentangling the relationship between sex-biased gene expression and X-linkage. Genome Research 22:1255-1265.

Mendes FK, Fuentes-González JA, Schraiber JG, Hahn MW. 2018. A multispecies coalescent model for quantitative traits. eLife 7:e36482.

Mendes FK, Vanderpool D, Fulton B, Hahn MW. 2020. CAFE 5 models variation in evolutionary rates among gene families. Bioinformatics 36:5516-5518.

Mione T, Anderson GJ. 1992. Pollen-ovule ratios and breeding system evolution in *Solanum* section *Basarthrum* (Solanaceae). American Journal of Botany 79:279-287.

Mitov V, Bartoszek K, Asimomitis G, Stadler T. 2020. Fast likelihood calculation for multivariate Gaussian phylogenetic models with shifts. Theoretical Population Biology 131:66-78.

Moyle LC, Wu M, Gibson MJ. 2021. Reproductive proteins evolve faster than non-reproductive proteins among *Solanum* species. Frontiers in Plant Science 12:635990.

O'Meara BC, Ané C, Sanderson MJ, Wainwright PC. 2006. Testing for different rates of continuous trait evolution using likelihood. Evolution 60:922-933.

Pease JB, Haak DC, Hahn MW, Moyle LC. 2016. Phylogenomics reveals three sources of adaptive variation during a rapid radiation. PLoS Biology 14:e1002379.

Pennell MW, Eastman JM, Slater GJ, Brown JW, Uyeda JC, FitzJohn RG, Alfaro ME, Harmon LJ. 2014. geiger v2. 0: an expanded suite of methods for fitting macroevolutionary models to phylogenetic trees. Bioinformatics 30:2216-2218.

Pennell MW, FitzJohn RG, Cornwell WK, Harmon LJ. 2015. Model adequacy and the macroevolution of angiosperm functional traits. The American Naturalist 186:E33-E50.

Price PD, Palmer Droguett DH, Taylor JA, Kim DW, Place ES, Rogers TF, Mank JE, Cooney CR, Wright AE. 2022. Detecting signatures of selection on gene expression. Nature Ecology & Evolution 6:1035-1045.

Revell LJ, Harmon LJ. 2022. Phylogenetic Comparative Methods in R: Princeton University Press.

Rick CM, Tanksley SD. 1981. Genetic variation in *Solanum pennellii*: Comparisons with two other sympatric tomato species. Plant Systematics and Evolution 139:11-45.

Robinson MD, Oshlack A. 2010. A scaling normalization method for differential expression analysis of RNA-seq data. Genome Biology 11:R25.

Rohlfs RV, Harrigan P, Nielsen R. 2014. Modeling gene expression evolution with an extended Ornstein–Uhlenbeck process accounting for within-species variation. Molecular Biology and Evolution 31:201-211.

Rohlfs RV, Nielsen R. 2015. Phylogenetic ANOVA: the expression variance and evolution model for quantitative trait evolution. Systematic Biology 64:695-708.

Silvestro D, Kostikova A, Litsios G, Pearman PB, Salamin N. 2015. Measurement errors should always be incorporated in phylogenetic comparative analysis. Methods in Ecology and Evolution 6:340-346.

Somel M, Creely H, Franz H, Mueller U, Lachmann M, Khaitovich P, Pääbo S. 2008. Human and chimpanzee gene expression differences replicated in mice fed different diets. PLoS ONE 3:e1504.

Swanson WJ, Vacquier VD. 2002. The rapid evolution of reproductive proteins. Nature Reviews Genetics 3:137-144.

Szövényi P, Ricca M, Hock Z, Shaw JA, Shimizu KK, Wagner A. 2013. Selection is no more efficient in haploid than in diploid life stages of an angiosperm and a moss. Molecular Biology and Evolution 30:1929-1939.

Vosters SL, Jewell CP, Sherman NA, Einterz F, Blackman BK, Moyle LC. 2014. The timing of molecular and morphological changes underlying reproductive transitions in wild tomatoes (*Solanum* sect. *Lycopersicon*). Molecular Ecology 23:1965-1978.

Wagner GP, Kin K, Lynch VJ. 2012. Measurement of mRNA abundance using RNA-seq data: RPKM measure is inconsistent among samples. Theory in Biosciences 131:281-285.

Wang Z, Gerstein M, Snyder M. 2009. RNA-Seq: a revolutionary tool for transcriptomics. Nature Reviews Genetics 10:57-63.

Whitehead MR, Lanfear R, Mitchell RJ, Karron JD. 2018. Plant mating systems often vary widely among populations. Frontiers in Ecology and Evolution 6:38.

Wong A. 2011. The molecular evolution of animal reproductive tract proteins: What have we learned from mating-system comparisons? International Journal of Evolutionary Biology 2011:908735.

Wray GA, Hahn MW, Abouheif E, Balhoff JP, Pizer M, Rockman MV, Romano LA. 2003. The evolution of transcriptional regulation in eukaryotes. Molecular Biology and Evolution 20:1377-1419.

Figures and Tables

Figure 1. Bounded Brownian motion model. An example trait is shown in the bottom graph, evolving along the tree shown above. Although the data input to CAGEE are linear expression levels, internally it logs expression to ensure higher variance among more highly expressed genes. There is also a minimum value, x_{min}, added to all tips.

Figure 2. Accuracy of CAGEE. For five different values of σ^2 we simulated 1000 datasets, with each dataset comprised of either 1000 genes or 10000 genes. All genes in a dataset shared the same σ^2, but their values at the root were drawn independently from the prior. We then provided each simulated dataset to CAGEE in order to infer σ^2. Each box-and-whisker plot shows the mean (horizontal line), 50% interquartile range (box), 1.5X the interquartile range (vertical lines), and outliers (dots).

Figure 3. Changes in gene expression along the tomato phylogeny. Given the set of relationships among the seven *Solanum* accessions used here, we tested multiple models that had branches assigned as different σ^2 parameters (Table 1). In model A, all branches share σ_1^2. In model B, all black branches share σ_1^2, while all red branches share σ_2^2. In model C, all black branches and the dashed red branch share σ_1^2, while all solid red branches share σ_2^2. In model D, all black branches share σ_1^2, all solid red branches share σ_2^2, and the dashed red branch is assigned σ_3^2. Using the results from model D, we inferred the number of genes that had credible increases or decreases in expression level along each branch (results for all changes are shown in Supplementary Figure 4). Numbers are reported as +increases/-decreases for each branch.

Table 1. Model parameters estimated from the tomato data.

Model	Number of rates	-lnL	σ^2_1	σ^2_2	σ^2_3
A	1	67252.4	0.102		
B	2	65883.9	0.074	0.134	
C	2	65124.5	0.075	0.152	
D	3	65108.6	0.077	0.152	0.067

Supplementary Figures

Supplementary Figure 1. The tree used for simulations. The Newick-formatted tree string with branch lengths is:

((((sp1:1,sp2:1):1,sp3:2):1,sp4:3):1,((sp5:2,sp6:2):2)):1,sp7:5):1,sp8:6)

Supplementary Figure 2. Accuracy of CAGEE. This figure is the same as Figure 2 in the main text, but the ancestral state vector has length N=500 (Figure 2 uses N=200).

Supplementary Figure 3. Trees used for simulations with lineage-specific values of σ^2.

A) All black branches share a rate parameter (σ_1^2), and all red branches share a rate parameter (σ_2^2). This "sigma_tree" is specified in CAGEE with the Newick string:

((((sp1:2,sp2:2):2,sp3:2):2,sp4:2):2,((sp5:1,sp6:1):1)):1,sp7:1):1,sp8:1)

B) All black branches share a rate parameter (σ_1^2), and all red branches share a rate parameter (σ_2^2). This "sigma_tree" is specified in CAGEE with the Newick string:

((((sp1:2,sp2:1):1,sp3:2):1,sp4:1):1,((sp5:1,sp6:2):1)):1,sp7:1):1,sp8:1)

Supplementary Figure 4. Changes in gene expression along the tomato phylogeny. This figure is the same as Figure 3 in the main text, but all increases and decreases are reported, regardless of whether they are "credible".

Supplementary Figure 5. Component diagram for the CAGEE software.

Figure 1

Figure 2

Figure 3

SI (σ^2_1)
SC (σ^2_2)
SC* (σ^2_3)

+405/-1047 — *S. habrochaites*

+8/-24

+321/-847 — *S. pennellii* (LA3778)

+68/-16

+406/-309 — *S. pennellii* (LA0716)

+275/-634 — *S. chilense*

+31/-17

+147/-257 — *S. chmielewskii*

+24/-14

+487/-1102

+42/-22 — *S. pimpinellifolium*

+819/-674 — *S. lycopersicum*

| 2.5 mya | 2.0 | 1.5 | 1.0 | 0.5 | 0 |

Supplementary Figure 1

Tree scale: 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Figure 5

Supplementary Table 1

Set of σ² values	Replicate	Simulated σ²	Inferred σ²	% error
0.2, 1.0	1	0.2	0.1926803	-3.66%
		1.0	1.0024905	0.249%
	2	0.2	0.1943218	-2.84%
		1.0	1.0054018	0.54%
	3	0.2	0.2031335	1.57%
		1.0	0.9577028	-4.23%
	4	0.2	0.1967767	-1.61%
		1.0	0.9996745	-0.0325%
	5	0.2	0.1924732	-3.76%
		1.0	1.0075440	0.754%
	6	0.2	0.2004951	0.248%
		1.0	1.0242412	2.42%
	7	0.2	0.1944284	-2.79%
		1.0	0.9733314	-2.67%
	8	0.2	0.2024781	1.24%
		1.0	0.9884935	-1.15%
	9	0.2	0.2018622	0.931%
		1.0	1.0382544	3.83%
	10	0.2	0.1956765	-2.16%
		1.0	1.0065474	0.655%
1.0, 3.0	1	1.0	0.9451556	-5.48%
		3.0	3.0287977	0.96%
	2	1.0	0.9769290	-2.31%
		3.0	3.1000229	3.33%
	3	1.0	0.9861053	-1.39%
		3.0	2.9452423	-1.83%
	4	1.0	0.9724728	-2.75%
		3.0	2.9203197	-2.66%
	5	1.0	0.9226307	-7.74%
		3.0	2.9930805	-0.231%
	6	1.0	0.9939074	-0.609%
		3.0	2.8856497	-3.81%
	7	1.0	0.9871113	-1.29%
		3.0	2.9705234	-0.983%
	8	1.0	0.9684494	-3.16%
		3.0	2.9187219	-2.71%
	9	1.0	0.9667607	-3.32%
		3.0	3.1521085	5.07%
	10	1.0	1.0059515	0.595%
		3.0	3.0232185	0.774%
5.0, 10.0	1	5.0	4.7464719	-5.07%
		10.0	9.9814701	-0.185%
	2	5.0	4.8651485	-2.7%
		10.0	10.0907926	0.908%
	3	5.0	4.7598429	-4.8%
		10.0	9.8058231	-1.94%
	4	5.0	4.9265027	-1.47%
		10.0	10.0593675	0.594%
	5	5.0	4.9239001	-1.52%
		10.0	9.9818898	-0.181%
	6	5.0	4.8684289	-2.63%
		10.0	9.8020147	-1.98%
	7	5.0	4.8752468	-2.5%
		10.0	10.0378953	0.379%
	8	5.0	4.9042695	-1.91%
		10.0	9.8541603	-1.46%
	9	5.0	4.6019327	-7.96%
		10.0	9.9286600	-0.713%
	10	5.0	4.8589572	-2.82%
		10.0	9.8313460	-1.69%