CAGEE: computational analysis of gene expression evolution

Jason Bertram[1,2,*], Ben Fulton[1,3], Jason P. Tourigny[1,4], Yadira Peña-Garcia[1], Leonie C. Moyle[1], and Matthew W. Hahn[1,4,*]

[1]Department of Biology, Indiana University, Bloomington, IN 47405, USA

[2]Department of Mathematics, Western University, London, ON N6A 5B7, Canada

[3]University Information Technology Services, Indiana University, Bloomington, IN 47405, USA

[4]Department of Computer Science, Indiana University, Bloomington, IN 47405, USA

*Corresponding authors: E-mail: jason.bertram@uwo.ca, mwh@indiana.edu

Abstract

Despite the increasing abundance of whole transcriptome data, few methods are available to analyze global gene expression across phylogenies. Here, we present a new software package (CAGEE) for inferring patterns of increases and decreases in gene expression across a phylogenetic tree, as well as the rate at which these changes occur. In contrast to previous methods that treat each gene independently, CAGEE can calculate genome-wide rates of gene expression, along with ancestral states for each gene. The statistical approach developed here makes it possible to infer lineage-specific shifts in rates of evolution across the genome, in addition to possible differences in rates among multiple tissues sampled from the same species. We demonstrate the accuracy and robustness of our method on simulated data, and apply it to a dataset of ovule gene expression collected from multiple self-compatible and self-incompatible species in the genus *Solanum* to test hypotheses about the evolutionary forces acting during mating system shifts. These comparisons allow us to highlight the power of CAGEE, demonstrating its utility for use in any empirical system and for the analysis of most morphological traits. Our software is available at https://github.com/hahnlab/CAGEE/.

Key words: RNA-seq; phylogenetic comparative methods; Brownian motion; Solanum

Introduction

Early studies of gene expression in single genes revealed widespread and frequent changes in the levels, timing, and breadth of expression across species (reviewed in Wray et al. 2003; Fay and Wittkopp 2008; Hill et al. 2021). Such changes in gene expression have been shown to be responsible for many differences between species, and may be a major driver of evolution (King and Wilson 1975). Advances in sequencing technologies (i.e. RNA-seq) have transformed research into gene expression, allowing researchers to cheaply and accurately measure transcript levels for every gene in a genome, in multiple tissues, and across several timepoints or conditions (Wang et al. 2009). There is now a flood of interest in applying RNA-seq to whole clades of organisms in order to identify the genetic changes and evolutionary forces driving species differences (e.g. Brawand et al. 2011; Meisel et al. 2012; Coolon et al. 2014; Harrison et al. 2015; Berthelot et al. 2018; Catalan et al. 2019; Blake et al. 2020; El Taher et al. 2021).

To better understand the importance of changes in gene expression, researchers must be able to characterize the mechanisms and modes by which gene expression evolves. Such work entails understanding the role of natural selection in driving species differences, the stages of development or the tissues that evolve most rapidly, as well as the environments most likely to generate changes in gene expression (Dunn et al. 2013; Hill et al. 2021; Price et al. 2022). Phylogenetic comparative methods enable the rigorous study of traits like gene expression across a species tree (Revell and Harmon 2022). These methods can be used for testing hypotheses about natural selection, the inference of ancestral states (allowing us to polarize the direction of changes), and the estimation of evolutionary rates. Multiple software packages are available that implement a wide variety of comparative methods (e.g. Pennell et al. 2014), including models specifically intended for studying gene expression across a tree (Bedford and Hartl 2009; Rohlfs et al. 2014; Rohlfs and Nielsen 2015; Catalán et al. 2019; Chen et al. 2019; Yang et al. 2019).

However, as far as we are aware, all existing comparative methods for analyzing gene expression implement fundamentally single-gene analyses. Each gene is

considered a separate trait, such the evolutionary parameters for each gene are

estimated separately. Single-gene analyses can be used to identify tissue-specific or

lineage-specific shifts in evolutionary rates, but their power is quite low (Beaulieu et al.

2012). As a result, identifying trends in evolution must be carried out *post hoc* by

summing the number of genes found to be individually significant (e.g. Harrison et al.

2015; El Taher et al. 2021). This approach is less than ideal, especially when carrying

out comparisons between branches of different lengths or between tissues with different

average expression levels (both of which can result in differential statistical power).

Therefore, to better characterize the forces affecting gene expression evolution,

we must be able to model effects shared along a lineage, experienced by many genes

in the same tissue, or experienced by all genes found in the same environment. In this

article, we present a genome-scale platform for the analysis of gene expression data

that allows for such shared factors. Our software, CAGEE (Computational Analysis of

Gene Expression Evolution), provides a robust set of methods for analyzing expression

data across a species tree. CAGEE estimates ancestral states and rates, with rates

shared by all or subsets of genes (single-gene analyses can also be carried out). We

show that lineage-specific and tissue-specific (or condition-specific) rates can be

accurately inferred, and provide principled statistical approaches for model selection.

Our current implementation uses a bounded Brownian motion model and assumes

expression data are accurate, but the architecture and codebase will easily allow for

future extensions that relax these and other assumptions.

New Approaches

We model gene expression evolution as a bounded Brownian motion (BBM)

process on a known species tree (cf. Boucher and Démery 2016). Our model has a

single bound: trait values must be greater than or equal to zero; there is no upper bound

(Figure 1). Previous researchers have often modeled gene expression using an

Ornstein-Uhlenbeck (OU) process (e.g. Bedford and Hartl 2009; Rohlfs et al. 2014;

Rohlfs and Nielsen 2015; Chen et al. 2019), a model that includes a force constraining

traits about the mean. However, to our knowledge, the OU model has only been

compared against an unbounded Brownian motion model (i.e. one that allows negative

expression values), making fair comparisons difficult. In addition, OU models may be

frequently and incorrectly favored over simpler models due to several biases (e.g.

measurement error), especially when the number of tips in a tree is small (Pennell et al.

2015; Silvestro et al. 2015; Boucher and Démery 2016; Cooper et al. 2016; Catalán et

al. 2019). Therefore, the initial version of our software models gene expression with the

BBM process, which naturally bounds possible values without invoking an additional

constraining force.

Let $E_{ij} \geq 0$ be the expression level of gene i in species j. We assume that log-

transformed expression $X_{ij} = \ln(E_{ij} + e_{\min})$ evolves as a Brownian motion process with

variance σ^2 per unit time, where $e_{\min}$ is a small offset (constant across genes and

species) that prevents X_{ij} from taking infinite values if measured values of E_{ij} are zero.

We log-transform before assuming Brownian motion because we expect the variance in

the evolutionary process to scale with expression level. Assuming that E_{ij} is itself

Brownian would unrealistically assume that the rate of evolution is constant across

expression levels, even though expression levels vary by many orders of magnitude.

We impose a reflecting lower boundary at $x_{\min} = \ln(e_{\min})$, meaning that the Brownian

walk immediately bounces back if it reaches $x_{\min}$. Expression can therefore effectively

never reach zero, our theoretical lower bound (Figure 1).

The second major feature of our model (as implemented in CAGEE) is that many

genes can share the evolutionary rate parameter, σ^2. This rate may be shared among

genes expressed in the same tissue or sample, among genes located on the same

chromosome, or among genes evolving along the same lineage of the phylogenetic

tree. The simplest model allows σ^2 to be shared among all genes, providing an average

rate of evolution across the genome and over time; this average may include genes that

vary in their individual rates of evolution. We explain this model briefly here, with more

detail provided in the Materials and Methods.

CAGEE infers the most likely value(s) of σ^2 consistent with an ultrametric tree, T,

and a set $E_{\{ij\}}$ of measured expression values at the tips of the tree; i.e. it maximizes

the likelihood $L(\sigma^2|E_{\{ij\}}, T)$. Each gene is assumed to evolve independently, and so the

likelihood for each gene $L_i(\sigma^2|E_{i\{j\}}, T)$ is computed independently. The overall likelihood

is obtained as the product $L(\sigma^2|E_{\{ij\}}, T) = \Pi_i L_i(\sigma^2|E_{i\{j\}}, T)$ across genes. The likelihood

for each gene $L_i(\sigma^2|E_{i\{j\}}, T)$ is computed using the pruning algorithm (Felsenstein

1973). The key ingredient needed to apply the pruning algorithm is the transition

probability density $p(x_t|x_{t_0}) = \Pr[X(t) = x_t|X(t_0) = x_{t_0}]$ for log-expression at time t

conditional on having log-expression x_{t_0} at time t_0 along a lineage. CAGEE computes

the transition density by solving the standard Brownian diffusion equation with reflecting

boundary conditions (Materials and Methods). The transition density is used to

propagate expression probabilities along the tree: if the probability density of log-

expression at time t_0 is $f(x_{t_0})$, then the probability density at time t on the same lineage

is $f(x_t) = \int p(x_t|x_{t_0})f(x_{t_0})dx_{t_0}$. At each tip the probability density $f(x_{t_0})$ is a delta

function centered at the corresponding measured value of X_{ij}.

Starting with the known tip distributions, the pruning algorithm propagates back

to the tips' parent nodes. The distribution at the parent node is then the product of the

two backward-propagated child node distributions. Proceeding iteratively across the

tree, we ultimately obtain the gene-specific probability density for expression value at

the root $f_i(x_R)$. Viewed as a likelihood for σ^2, $f_i(x_R)$ is the gene-specific likelihood

conditional on the unknown ancestral root value; i.e. $f_i(x_R) = L_i(\sigma^2|E_{i\{j\}}, T, x_R)$.

Therefore, we integrate over all possible x_R to obtain,

$$L_i(\sigma^2|E_{i\{j\}}, T) = \int L_i(\sigma^2|E_{i\{j\}}, T, x_R)\,\rho(x_R)dx_R, \qquad (1)$$

where $\rho(x_R)$ is the prior distribution for the root value of a randomly selected gene.

The default prior $\rho(x_R)$ is assumed to be a gamma distribution with $k = 0.375$ and $\theta =$

1600, though this distribution can also be set by the user in CAGEE. This choice is

based on estimated expression distributions across genes in individual species, which

we take as our baseline for the ancestral distribution. CAGEE uses the Nelder-Mead

simplex method to find the optimal value(s) of σ^2.

Results

Using CAGEE

The required inputs for CAGEE are a Newick-formatted, rooted, ultrametric tree
(with branch lengths) and a tab-delimited data file containing the expression levels of all
species or taxa being studied. The data file can consist of data on one gene/transcript
or thousands of different genes. The first line of the data file should contain the species'
names (matching those used in the Newick tree). In addition, headers for gene names,
gene descriptions, and sample IDs (see next section for an explanation of "samples" in
CAGEE) can be used. Subsequent lines each correspond to a single gene and contain
expression levels for each species. Missing data can be denoted using multiple
characters (-/?/N). Examples of Newick trees and corresponding data files can be found
in the online user manual
(https://github.com/hahnlab/CAGEE/docs/manual/cagee_manual.md).

We expect that CAGEE will most often be used to calculate the following outputs:
one or more σ^2 values, ancestral states at each internal node (including 95% credible
intervals around these states), and the final likelihood associated with a model.
However, users do not have to search for σ^2: if a value for this parameter is specified,
then the output of CAGEE will just be the ancestral states and a likelihood. In addition to
the raw outputs provided in multiple formats (both tab-delimited files and NEXUS-
formatted files), CAGEE computes basic statistics about changes in expression levels
by comparing values at parent and child nodes. Summaries of these inferred changes
for every gene and for every branch of the tree are output, so that the evolutionary
history of gene expression changes in every gene are accessible to users. To avoid
over-interpretation of small changes in inferred expression levels—especially when
there is uncertainty in ancestral states—CAGEE will also compare the credible intervals
at parent and child nodes to note if a change is "credible" (i.e. the intervals do not
overlap). Credible intervals are calculated by summing the probabilities across possible
ancestral states at each node, so that 95% of the probability density is included.
Credible changes on each branch are annotated as such in the output.

We most often expect that an ultrametric species tree will be used as the input

topology, but this is not required by CAGEE. If users wish to specify a gene tree, or

some other bifurcating tree, as input, those can be used in CAGEE as well. However,

the major advantage of CAGEE—incorporating information from multiple genes to

accurately estimate genome-wide rates—will rapidly diminish for trees that represent

the history of only a minority of the genome. Trees that include duplication events

should provide suitable estimates for any genes that follow this topology, but CAGEE

does not have a way to further combine disparate gene trees.

There are multiple options available for running CAGEE. Users who can take

advantage of multiple threads can specify the number to use on the command line.

Complex models can also take a long time to converge; by default, CAGEE runs a

maximum of 300 iterations of the Nelder-Mead search, but users can increase this

number in subsequent runs if the likelihood is still improving when the limit is hit. As

mentioned above, the default prior distribution for the root state is a gamma distribution

with $k = 0.375$ and $\theta = 1600$. This distribution can also be specified by the user if

desired. Information on how to run more complex evolutionary models, beyond a single

σ^2, is given in the next section.

Estimating evolutionary rates in CAGEE

We tested CAGEE's ability to accurately estimate σ^2 by varying this rate

parameter and the number of genes used for inference, as well as the amount of

missing data in each dataset. We simulated different single values of σ^2 across a tree

with constant branch lengths (Supplementary Figure 1) using the simulation tool

available within CAGEE. (Note that the total amount of evolution in a tree is determined

by the product $\sigma^2 \cdot t$, such that changes in branch lengths will have an effect

commensurate with changes in σ^2.) Figure 2 shows the average error associated with

estimates of different σ^2 values and using different numbers of genes within each

dataset. As can be seen, the error across all parameter values and dataset sizes is

quite small (generally less than 2.5%), and is less variable for larger dataset sizes.

Fortunately, we expect that most empirical datasets will contain closer to 10,000 genes

than 1,000 genes. The results in Figure 2 are for an ancestral state vector of length

N=200 (the default setting in CAGEE; Materials and Methods); we also estimated σ^2

when allowing the ancestral state vector to have length *N*=500 (Supplementary Figure

2A). There appears to be minimal gain from increasing the resolution in this vector,

though the computational time is greatly increased (similar to results in Boucher and

Démery 2016). We evaluated the accuracy of CAGEE when different amounts of data

were randomly missing: from 0% to 75% for a dataset of 1,000 genes. As shown in

Supplementary Figure 2B, CAGEE has high accuracy even when large amounts of data

are missing (at random) from a dataset.

219   One major advantage of using CAGEE is that it combines information from

multiple genes to infer a rate of evolution: this is why it can return estimates with high

accuracy even when a large fraction of the data are missing. To further demonstrate this

advantage, we simulated evolution in 1,000 genes using the same parameter value

(σ^2=1) and then estimated σ^2 for each of the 1,000 genes individually. Supplementary

Figure 2C shows that these individual estimates of σ^2 are quite error-prone: although

the mean of all genes is close to the true value, individual estimates can be 7-8X higher

or lower and there is a large amount of variance. Although we have not shown it here,

we do expect that the accuracy of σ^2 will be greater for trees with larger numbers of

tips, even for estimates derived from single genes (cf. O'Meara et al. 2006). On the

other hand, CAGEE is combining information from multiple genes to infer an *average*

rate of evolution, even when the underlying rate may be quite variable. To explore any

effect of underlying rate variation, we carried out further simulations that combined three

simulations of 1,000 genes each with σ^2 equal to 0.5, 3, and 9, respectively (we

repeated these simulations 10 times). When analyzed as single datasets with 3,000

genes total, the average σ^2 inferred was 3.76, approximately 9% lower than the

arithmetic mean rate (Supplementary Figure 2D). It is well-known that single-rate

phylogenetic likelihood models tend to underestimate rates of evolution when there is

underlying variation (Golding 1983; Gillespie 1986; Yang 1996; Mendes et al. 2020),

and we see this effect here. Fortunately, the bias is small, and can be corrected in the

future by including gamma-distributed rate variation into CAGEE. Overall, inferences of

σ^2 should be quite accurate when a single rate parameter is shared across the tree and
across all genes and lineages.

Variation in the rate of expression can currently be accommodated by CAGEE in
a number of ways, using multi-rate σ^2 models. One type of model allows users to
specify that their data come from different "samples": these samples can represent
tissues, conditions, timepoints, and even subsets of the genome (e.g. the X
chromosome, or a specific functional class of genes). In the input data file, the
"SAMPLETYPE" column is used to indicate which sample each gene is a member of; a
separate σ^2 value will be calculated for each sample or set of samples (these values
are assumed to be shared among all lineages in the tree). Specifying more than one
sample means that an individual gene or transcript name can be used more than once
(i.e. once for each sample), but there is no requirement that genes are measured in
each sample. For instance, assigning all autosomal genes to sample 1 and all X-linked
genes to sample 2 would not permit for any overlap in gene assignment, but is perfectly
allowable in CAGEE.

Each additional sample requires another σ^2 parameter to be estimated, and often
researchers would like to know if fitting this extra parameter is justified by the data.
Under standard information-theoretic criteria (Burnham and Anderson 2002), twice the
difference in log-likelihoods between nested models should be χ^2-distributed with
degrees of freedom equal to the difference in the number of parameters between
models. To test this expectation, we simulated 1000 datasets with a single σ^2 value, but
fit models with two σ^2 values (assigning 1000 genes to two equal-sized samples at
random; the relative size of the samples should not affect the false positive rate). As
anticipated, the results fit a χ^2 distribution with one degree of freedom, with 4.4% of
datasets having a difference in 2*log-likelihood greater than 3.84 (5% are expected by
chance). This indicates that standard statistical procedures should adequately control
the false positive rate when fitting multi-sample σ^2 models.

CAGEE also allows models in which σ^2 varies across branches of the species
tree. It does so by fitting separate σ^2 parameters for different parts of the tree. On the

command line, CAGEE enables users to specify how multiple σ^2 parameters should be

assigned to branches. For n taxa, from 1 to $2n$-2 parameters can be specified, and

branches can be grouped together in any way. For instance, a two-parameter model

can have all branches that share a rate adjacent to one another in the tree

(Supplementary Figure 3A) or spread out across the tree (Supplementary Figure 3B).

Similar to the analyses carried out above for the false positive rate associated with

multiple samples, we simulated data with a single σ^2 value and then fit models with

multiple σ^2 parameters. Regardless of how we distributed the two rate classes across

the tree we observed good control of the false positive rate: 4.5% and 5.4% of 1000

simulated datasets were significant at the P=0.05 level (for the trees shown in

Supplementary Figures 3A and 3B, respectively). More limited simulations also showed

that we could accurately estimate multiple σ^2 parameters when the data were simulated

with multiple rates (Supplementary Table 1). Together, our results suggest that we can

estimate multiple types of multi-rate models, and can accurately control the false

positive rate when doing so.

Analysis of wild tomato transcriptome data

To demonstrate the utility of CAGEE in an empirical system, we analyzed data

from a clade that includes domesticated tomato, *Solanum lycopersicum*. This dataset

contains gene expression levels in unfertilized ovules from the flowers of six species,

one of which (*S. pennellii*) has two different populations represented (Figure 3). There

are 14,556 genes with expression levels measured in all seven accessions. RNA-seq

data for five of the seven accessions have been published previously (Moyle et al. 2021;

Hibbins and Hahn 2021), while two others are presented here for the first time

(Materials and Methods). Note, however, that all data were collected from all samples at

the same time (Materials and Methods).

Most species within the tomato clade are self-incompatible (SI), the ancestral

state in the family Solanaceae (Igić et al. 2006). Self-incompatibility means that plants

must outcross in order to successfully fertilize ovules. However, self-compatibility (SC)

has evolved multiple times both within the Solanaceae and within the genus *Solanum*

(Goldberg et al. 2010; Bedinger et al. 2011). Self-compatible individuals are able to

successfully fertilize ovules using their own pollen, though many also still outcross

(Whitehead et al 2018; including in *Solanum*: Vosters et al. 2014 and references

therein). Importantly, we have *a priori* expectations about the rate at which reproductive

traits—including ovule gene expression—might evolve between groups with different

mating systems. Due to conflict within and between the sexes, it is generally expected

that reproductive traits in species that outcross more (i.e. SI taxa) should evolve more

rapidly than in species that inbreed more (i.e. SC taxa; Clark et al. 2006). Such patterns

are found in some analyses of the rate of protein evolution (e.g. Gossmann et al. 2016;

Harrison et al. 2019), but are equivocal in other comparisons (e.g. Gossmann et al.

2014, Moyle et al. 2021). These complex patterns might reflect additional effects that

also accompany mating system shifts; for instance, such shifts often lead to reductions

in effective population size in more selfing lineages (Charlesworth and Wright 2001).

Mating system shifts could also alter global patterns of molecular evolution (including

gene expression) by changing the strength and pattern of purifying selection, as

morphological changes often accompany mating system changes. The exact effect of

shifts in mating system on molecular evolution remains an open question.

The *Solanum* species sampled here represent two independent transitions from

SI to SC, with one of the transitions (in accession *S. pennellii* LA0716) occurring

recently enough that different populations within this species have different

incompatibility systems (Figure 3). We therefore fit a series of nested models within

CAGEE to test two related hypotheses about ovule gene expression evolution. First, we

would like to know whether the rate of evolution of ovule gene expression is different in

SI species than in SC species. Second, given the recent transition to SC within

accession *S. pennellii* LA0716, we wanted to know if it shows a pattern of evolution

more similar to SI or to SC species. In total, we fit four separate evolutionary models

(Table 1; Figure 3). Model A has a single rate parameter for the entire tree. Model B has

two rate parameters, one for SI species and one for SC species. This model assigns the

branch leading to *S. pennellii* LA0716 as SC. Model C also has two rate parameters,

one for SI and one for SC, but assigns *S. pennellii* LA0716 as SI. Model D has three

rate parameters: one for SI species, one for longer-term SC species, and one for *S.*

pennellii LA0716.

Estimated results from the different models are shown in Table 1. Model A has a

worse fit than any other model, with a single σ^2 value of 0.102. For context, this value

means that the bounded Brownian motion process the data are fit to has a variance of

0.102 per million years (of log-transformed expression values). This is the average rate

across all 14,556 genes and across all branches of the tree. In contrast to a single-rate

model, both models B and C are significantly better fits to the data. Contrary to some

hypotheses, both models find that SI lineages (σ_1^2) have a lower rate of evolution than

SC lineages (σ_2^2; Table 1). There is also a difference between the models, with model C

(the one in which *S. pennellii* LA0716 shares a rate with SI species) fitting significantly

better. To further examine the evolution of *S. pennellii* LA0716, model D fits a three-

parameter model, with this lineage assigned its own rate of evolution. This model is a

significantly better fit than model C (P<0.00001; χ^2 test with 1 degree of freedom), and

demonstrates that *S. pennellii* LA0716 has a rate of evolution (σ_3^2 in Table 1) that is

slightly *lower* than SI species. This highly similar rate to SI species implies that it has

only recently transitioned to self-compatibility, which is consistent with previous

inferences about the timing of transition to SC in this particular accession (e.g. Rick and

Tanksley 1981).

CAGEE also allows users to infer the number and direction of changes in gene

expression levels along each branch of the tree. Figure 3 reports the number of genes

that had "credible" increases and decreases in expression level under model D.

Credible changes require that the credible intervals around states at parent and

daughter nodes do not overlap, in order to account for uncertainty in our inferences.

However, because of this, fewer credible changes will be inferred deeper in the tree,

where credible intervals get wider. Therefore, while inferences about the identity of the

genes changing along each branch is greatly strengthened by using credible changes

(these genes are noted in the raw output from CAGEE), the absolute numbers of

credible changes cannot be compared across branches, except for sister branches of

equal length. For completeness, we show the total numbers of increases and decreases

of gene expression in Supplementary Figure 4; as expected, these total numbers are

more uniformly distributed across older and younger branches.

We assessed whether the genes identified as having credible increases or

decreases in expression specifically on any SC branch (solid red branches in Figure 3)

were significantly enriched for any biological process or molecular function gene

ontology (GO) categories compared to genes with credible changes on any SI branch

(black branches in Figure 3). This comparison specifically assesses gene expression

evolution associated with a transition to SC, over and above "background" rates of

expression evolution across the rest of the clade. Although fold enrichment was modest

1.20-1.36X; Supplementary Table 2), there were 11 terms significantly enriched

(FDR<0.05) specifically on SC branches; these terms primarily focused on regulation of

transcription, metabolic processes, and biosynthesis (Supplementary Table 2). Among

the genes in these over-represented categories, a large fraction are transcription factors

associated with development (e.g. WRKY and MADS Box), hormonal responses

(including ethylene- and auxin-responsive transcription factors), and regulation of cell

cycle (e.g. cyclins), in addition to protein kinases (Supplementary Table 2). This

enrichment is consistent with increased expression changes in genes involved in cell

division, differentiation, and development, that could follow transitions to SC.

Discussion

Here, we have developed a new software package that enables the estimation of rates of gene expression evolution across a tree, CAGEE. Gene expression levels are much like many other continuous traits, and multiple papers have introduced phylogenetic comparative methods for studying gene expression (Bedford and Hartl 2009; Rohlfs et al. 2014; Rohlfs and Nielsen 2015; Catalán et al. 2019; Chen et al. 2019). However, as far as we are aware none of these methods allows genes to share evolutionary parameters, which precludes the analysis of genome-wide trends, either along the branches of a tree or between tissues/samples/conditions. To overcome this limitation, CAGEE calculates the likelihood of the data using the pruning algorithm (Felsenstein 1973) to facilitate the sharing of evolutionary parameters along branches of the species tree, providing more statistical power to test evolutionary hypotheses. Fortunately, we were able to take advantage of much of the codebase of our existing software, CAFE (Hahn et al. 2005; De Bie et al. 2006; Hahn et al. 2007; Han et al. 2013; Mendes et al. 2020), which implements the pruning algorithm for the analysis of gene family sizes across a tree. While gene expression levels and gene family sizes differ in the type of data they represent (continuous vs. discrete) and their underlying evolutionary models (bounded Brownian motion vs. birth-death), many of the required likelihood calculations and software components are the same.

An important thing to consider for the input to CAGEE is the normalization used to make gene expression levels comparable across species. The data from wild tomatoes used here was normalized using TPM (transcripts per million; Wagner et al. 2012); other published datasets also use this normalization (Berthelot et al. 2018; Chen et al. 2019; El Taher et al. 2021). However, multiple other normalizations have also been used in comparative analyses, including RPKM (Brawand et al. 2011), FPKM (Catalán et al. 2019), and both TMM and CPM (Blake et al. 2020). Each normalization approach has its advantages and disadvantages, and we cannot yet strongly recommend one specific approach as input to CAGEE. The normalization method used will likely depend on the conditions under which samples are collected: if all species can be raised simultaneously in a greenhouse, vivarium, or growth chamber, we expect

many fewer batch effects than in samples collected from the field, which will therefore

necessitate different normalizations. However, even animals raised in a common

environment—but fed different diets—can show many differences in gene expression

not due to heritable change (e.g. Somel et al. 2008). Conversely, many between-sample

normalization approaches (e.g. TMM, trimmed mean of M values; Robinson and

Oshlack 2010) make the assumption that differences in gene expression between

samples are rare. While such normalization is sensible in the context of testing for

differential expression between samples from the same species, for a set of species

that have been evolving independently for millions of years this is likely not an

appropriate assumption.

CAGEE currently has multiple limitations, both in the available models that can

be applied and in the types of data that can be analyzed. As mentioned earlier, many

researchers have modeled gene expression using an OU process (Bedford and Hartl

2009; Rohlfs et al. 2014; Chen et al. 2019; Yang et al. 2019). Although OU models may

be artifactually preferred over unbounded Brownian motion models due to a number of

non-biological factors (see discussion in "New Approaches" above), it would still be

helpful to be able to compare such a model to the bounded Brownian motion model

used here. However, fitting such a model to genome-wide data is non-trivial: each gene

must have its own mean expression value (μ), but possibly shared constraint

parameters (α) across genes. We have the goal of implementing such a model in the

near future, as well as other models commonly used in comparative methods research

(e.g. Landis and Schraiber 2017; Boucher et al. 2018).

Beyond the evolutionary model applied to any dataset, there are multiple

additional sources of variation that could be modeled. For instance, we have previously

accounted for measurement error in a likelihood framework, using an empirically

parameterized error model (Han et al. 2013). We can imagine both applying a similar

model here to RNA-seq data, as well as extending CAGEE to more error-prone data

such as single-cell sequencing. Such an extension would treat the level of expression in

each cell within a cell type as an error-prone draw from an underlying distribution; one

would then be able to infer the rate of evolution within and across cell-types across

multiple species. The biggest obstacle to this approach may be in identifying

homologous cell types across species (e.g. Tarashansky et al. 2021). In addition, not all

genes necessarily share the same average rate of evolution; gamma-distributed rate

categories can be used to model this variation among genes (cf. Ames et al. 2012;

Mendes et al. 2020). As shown above, not accounting for this rate variation leads to a

slight underestimate of σ^2, but also obscures interesting patterns of evolution among

genes. Finally, the gene tree discordance found in many phylogenomic datasets implies

that complex traits (such as expression levels) will also be controlled by discordant gene

trees (Hahn and Nakhleh 2016; Hibbins and Hahn 2021). This underlying discordance

can cause evolutionary rates to be overestimated (Mendes et al. 2018), and should be

taken into account when seeking accurate parameter estimates (see discussion of wild

tomato data below). Our goal is to include methods for dealing with all these sources of

variation in future versions of CAGEE.

In terms of the types of data that can be analyzed, at present CAGEE is limited to

positive, continuously varying traits (i.e. the BBM model). However, we also envision

different ways to represent and model gene expression data, including as a ratio (e.g.

male/female expression). Such a ratio, after log2-transformation, would be most

appropriately modeled by an unbounded Brownian motion model since both negative

and positive values are possible. This and other data types will be supported in future

releases. Moreover, CAGEE does not have to analyze whole-genome or even

molecular data: it can be applied to any single trait for which the BBM model is

appropriate, even morphological traits. One intriguing application of CAGEE could be to

suites of morphological traits that are hypothesized to share a common evolutionary

rate parameter. If, for instance, there is a shift in body plan along some lineages, then

multiple traits may all increase or decrease their rate of evolution at once, and CAGEE

can be used to estimate these shared parameters. Even in the context of single-trait

analyses, the pruning algorithm has been hailed as a solution for large-scale

comparative analyses (Freckleton 2012). Importantly, the number of branches in a

rooted, bifurcating tree with *n* tips is 2*n*-2, so that the number of calculations scales

linearly with the number of species. This makes the pruning algorithm ideal for

comparative datasets with large numbers of taxa (e.g. Hahn et al. 2005; FitzJohn 2012;

Hiscott et al. 2016; Caetano and Harmon 2018; Mitov et al. 2020).

The analysis of data from a clade of wild tomatoes revealed a possibly

unexpected result: the rate of ovule gene expression evolution among self-compatible

(SC) species is twice as high as the rate among self-incompatible (SI) species (Table

1). This finding is contrary to some prior expectations—informed by research focused

on male-female interactions, especially between interacting proteins in the reproductive

tract (e.g. Swanson and Vacquier 2002; Clark et al. 2006)— that suggest that lineages

might experience slower evolution after transitioning to self-compatibility. However, it is

possible that global gene expression levels do not evolve in the same sort of tit-for-tat

manner as interacting protein sequences, such that increases/decreases in male-

expressed genes are not matched by increases/decreases in interacting female-

expressed genes (or vice versa). Alternatively, only a very small subset of genes may

evolve in this manner. Indeed, even prior studies comparing protein evolution have

failed to find clear evidence of slower global evolutionary rates in more inbreeding

species (e.g. Wong 2011). One caveat to the observed rate differences in our data is

that underlying gene tree discordance, whether due to incomplete lineage sorting or

introgression, can lead to artifactually higher rate estimates (Mendes et al. 2018;

Hibbins and Hahn 2021). However, there is in fact less discordance among the SC

lineages sampled here (Pease et al. 2016), which is the reverse of the pattern that

would be required to explain our results.

If not due to underlying bias in our estimates, these findings still raise the

question: why is ovule gene expression evolving more rapidly in SC than SI species?

One possibility is that this increased rate is due to a relaxation of selection in SC

species, possibly because genes involved in male-female interactions are no longer

needed. If this were the case, we might expect to see a general decrease in expression

levels in SC species; however, there appears to be no consistent directionality to the

changes along SC branches (Figure 3, Supplementary Figure 4). Instead, an alternative

hypothesis is that transitions to SC involve adaptation to new optima of ovule gene

expression, compared to SI species that tend to maintain ancestral optima. For

example, transitions to SC might favor greater investment in fewer ovules, because self-

compatibility decreases the probability that each ovule within a flower will go

unfertilized—an otherwise wasted investment under conditions (like SI) where receiving

sufficient compatible pollen to fertilize each ovule is less predictable (Burd et al. 2009).

The nature of these new optima might be even more complex, as traits like ovule size

and number can vary with multiple reproductive and ecological conditions, and often

trade-off with each other (Greenway and Harder 2007). Of the species examine here,

for example, two SC lineages (*S. pimpinellifolium*, and *S. lycopersicon*—domesticated

tomato) have significantly larger seeds than most of the SI lineages and SC *S. pennellii*

(unpubl. data). Indeed, individual genes identified in our GO analysis are known to

directly influence ovule and/or seed size in *Solanum* (e.g. *NOR-like1*

[SOLYC07G063420.3.1; Han et al, 2014], *GRAS2* [SOLYC07G063940.2.1; Li et al.

2018], and *CRY2* [SOLYC09G090100.3.1; Fantini et al. 2019]). Some of our

hypotheses could be evaluated with matching gene expression data from other (non-

ovule) reproductive tissues. Analyses including pollen in the same SI and SC lineages,

and/or data addressing alternative constraints and conditions shaping ovule evolution

including ovule size and number (e.g. Mione and Anderson 1992), would be useful in

teasing apart these hypotheses.

Material and Methods

Bounded Brownian motion model of expression evolution

The probability density of expression, $p(x,t)$, at time t for evolutionary

trajectories following a Brownian motion process starting at value x_{t_0} at time t_0 is

governed by the diffusion equation

523
$$\frac{\partial p(x,t)}{\partial t} = \frac{\sigma^2}{2}\frac{\partial^2 p(x,t)}{\partial x^2}, \qquad\qquad (2)$$

524   with initial condition $p(x,t_0) = \delta(x - x_{t_0})$ where δ is the Dirac delta function. The

525   reflective boundary condition at $x = x_{\min}$ implies that the probability fluxes into and out

526   of the boundary are balanced, imposing the boundary condition

527
$$\frac{\partial p(x=x_{\min},t)}{\partial x} = 0. \qquad\qquad (3)$$

Note that $p(x,t)$ is identical to the transition density $p(x_t|x_{t_0})$.

Without the reflecting boundary, $p(x,t) \propto e^{-(x-x_{t_0})^2/2\sigma^2(t-t_0)}$ is a normal

distribution with variance $\sigma^2(t - t_0)$. The variance therefore scales linearly with elapsed

time, $t - t_0$. With the reflecting boundary, $p(x,t)$ is the sum of this spreading normal and

its mirror image centered at $2x_{\min} - x_{t_0}$. The analytical solution to this bounded process

is helpful for understanding the behavior of $p(x,t)$, but is not used in CAGEE. In

anticipation of implementing additional (and possibly more complicated) processes into

CAGEE, we instead solve Eq. (2) numerically using the approach described in Boucher

and Démery (2016). Briefly, the continuous diffusion equation is converted into a matrix

equation by discretizing expression values into N equal bins of width $\delta = \frac{x_{\max}-x_{\min}}{N-1}$.

Following Boucher and Démery (2016), we have used a default *N*=200, but this number

can be set by the user (see Results). This approach gives

540
$$\frac{\partial P(t)}{\partial t} = \frac{\sigma^2}{2\delta^2} M \cdot P(t), \qquad\qquad (4)$$

541   where $P(t)$ is the vector obtained by discretizing $p(x,t)$ and $x_{\max}$ is the largest

542   expression value accounted for. The matrix M is tridiagonal with -2 on the diagonal

543   except at the first and last diagonal entries which are -1. The sub- and supra-diagonal

544   entries are 1. This equation has the matrix exponential solution

545
$$P(t) = \exp\left(\frac{\sigma^2(t-t_0)}{2\delta^2} M\right), \qquad (5)$$

which is evaluated by diagonalizing M.

Implementation of CAGEE

CAGEE is written in C++ and is compatible with the C++11 standard. A

comprehensive manual and extensive unit tests facilitate further code development and

maintenance. CAGEE is organized into modular components. A *clade* class, with

references to a parent clade and any number of descendant clades, represents a tree

structure, and a *gene_transcript* class represents the expression levels observed in the

various species. These two classes comprise the fundamental data structures upon

which CAGEE performs its analysis (Supplementary Figure 5).

Calculations are carried out by additional classes. The *optimizer* class has the

responsibility of determining the σ^2 value with the highest likelihood, by comparing the

likelihood of candidate values and searching the likelihood surface using the Nelder-

Mead optimization algorithm. The work of computing the likelihood of a given σ^2 value is

performed by a subclass of the *model* class, which for now is limited to a single *Base*

model (allowing for further development in the future). After appropriate estimated

values are found, the *transcript_reconstructor* class builds a possible set of transcript

values for the entire tree (Supplementary Figure 5).

Performing the likelihood calculations requires extensive matrix operations; it is

recommended (though not required) that these be passed off to a specialized library

such as Intel's MKL or Nvidia's CUBLAS. If no external library is available, CAGEE will

carry out these calculations (slowly) by itself. Creating the diffusion matrix (M) requires

calculation of eigenvalues and eigenvectors, and is computationally expensive. This

work is performed by the Eigen linear algebra library (https://eigen.tuxfamily.org);

various internal data structures also take advantage of Eigen classes. To enable faster

searching, the matrix for an ancestral state vector of length 200 (the default in CAGEE)

has been pre-computed and is included with CAGEE. Users who wish to use vectors of

different lengths can specify this as an option.

Unit-testing is performed using the Doctest testing framework

(https://github.com/doctest/doctest). At the time of writing more than 200 unit tests had

been created, comprising more than 1200 individual assertions. For complex logging

and debugging cases, CAGEE uses the EasyLogging framework

(https://github.com/amrayn/easyloggingpp). C++ development is always made easier by

using the Boost C++ libraries (https://www.boost.org/), so we include them as well in

CAGEE.

RNA-seq data from wild tomatoes

We briefly describe here the data collected from seven accessions of wild

tomatoes (*S. lycopersicum* LA3475, *S. chmielewskii* LA1316, *S. pimpinellifolium*

LA1589, *S. habrochaites* LA1777, *S. chilense* LA4117A, *S. pennellii* LA3778, and *S.*

pennellii LA0716; all accession ID numbers from tgrc.ucdavis.edu). Further details are

given in Moyle et al. (2021). Ovule RNA-seq was performed on between one to four

(usually three) biological replicates (individual plants) from each accession. Plants were

germinated from seed, and cultivated until flowering. For each replicate individual,

ovules were dissected from mature, unpollinated flowers, flash frozen, and maintained

at -80C until extraction. For each individual, all ovule collections were pooled into a

single sample prior to library construction and sequencing on an Illumina HiSeq 2000.

Reads were mapped against the tomato reference genome (ITAG 2.4) and the number

of reads mapped onto genic regions were estimated with featureCounts (Liao et al.,

2014). We normalized the read counts from each library by calculating TPM (transcripts

per million; Wagner et al. 2012) and then calculated the mean normalized read counts

across all samples (individuals) within each accession. These means per accession

were used as input to CAGEE.

To construct a species tree for use with CAGEE, we started with the topology

given in Pease et al. (2016). Specifically, we used the tree found in the supplementary

file Pease_etal_TomatoPhylo_RAxMLConcatTree_no1360_Fig2A.nwk, and pruned it to

include only the accessions in our study using the software ETE (Huerta-Cepas et al.

2016). Using the "extend" method found in ETE, we converted this tree to ultrametric

(same root-to-tip distance for all taxa). Setting the root age to 2.48 million years ago

(following Pease et al. 2016), we were able to express all branches in millions of years.

Analyses of GO enrichment were carried out using ShinyGO (Ge et al. 2020) with a

false discovery rate of 0.05.

Supplementary Material

Supplementary data are available at .

Acknowledgements

We thank Mark Hibbins for assistance with the tomato phylogeny, Matthew Gibson for putting together the tomato gene expression data, and especially Dan Vanderpool for invaluable help in the initial development of CAGEE. Two reviewers provided helpful comments, and Scott Edwards pointed out relevant work that we had previously missed. This work was supported by National Science Foundation grants DEB-1856469 to L.C.M. and DBI-2146866 to M.W.H.

Data Availability

Raw reads for each sample library are available at NCBI BioProject PRJNA714065. The CAGEE software is available at https://github.com/hahnlab/CAGEE.

References

References

Ames RM, Money D, Ghatge VP, Whelan S, Lovell SC. 2012. Determining the
evolutionary history of gene families. Bioinformatics 28:48-55.

Beaulieu JM, Jhwueng DC, Boettiger C, O'Meara BC. 2012. Modeling stabilizing
selection: expanding the Ornstein–Uhlenbeck model of adaptive evolution. Evolution
66:2369-2383.

Bedford T, Hartl DL. 2009. Optimization of gene expression by natural selection.
Proceedings of the National Academy of Sciences 106:1133-1138.

Bedinger PA, Chetelat RT, McClure B, Moyle LC, Rose JK, Stack SM, van der Knaap E,
Baek YS, Lopez-Casado G, Covey PA. 2011. Interspecific reproductive barriers in
the tomato clade: opportunities to decipher mechanisms of reproductive isolation.
Sexual Plant Reproduction 24:171-187.

Berthelot C, Villar D, Horvath JE, Odom DT, Flicek P. 2018. Complexity and
conservation of regulatory landscapes underlie evolutionary resilience of mammalian
gene expression. Nature Ecology & Evolution 2:152-163.

Blake LE, Roux J, Hernando-Herraez I, Banovich NE, Perez RG, Hsiao CJ, Eres I,
Cuevas C, Marques-Bonet T, Gilad Y. 2020. A comparison of gene expression and
DNA methylation patterns across tissues and species. Genome Research 30:250-
262.

Boucher FC, Démery V. 2016. Inferring bounded evolution in phenotypic characters
from phylogenetic comparative data. Systematic Biology 65:651-661.

Boucher FC, Démery V, Conti E, Harmon LJ, Uyeda J. 2018. A general model for
estimating macroevolutionary landscapes. Systematic Biology 67:304-319.

Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, Weier M, Liechti
A, Aximu-Petri A, Kircher M, et al. 2011. The evolution of gene expression levels in
mammalian organs. Nature 478:343-348.

Burd M, Ashman T-L, Campbell DR, Dudash MR, Johnston MO, Knight TM, Mazer SJ,
Mitchell RJ, Steets JA, Vamosi JC. 2009. Ovule number per flower in a world of
unpredictable pollination. American Journal of Botany 96:1159-1167.

Burnham KP, Anderson DR. 2002. Model selection and multimodel inference: A
practical information-theoretic approach. New York: Springer.

Caetano DS, Harmon LJ. 2018. Estimating correlated rates of trait evolution with uncertainty.
Systematic Biology 68:412-429.

Catalán A, Briscoe AD, Höhna S. 2019. Drift and directional selection are the
evolutionary forces driving gene expression divergence in eye and brain tissue of
Heliconius butterflies. Genetics 213:581-594.

Charlesworth D, Wright SI. 2001. Breeding systems and genome evolution. Current
Opinion in Genetics & Development 11:685-690.

Chen J, Swofford R, Johnson J, Cummings BB, Rogel N, Lindblad-Toh K, Haerty W, Di Palma F, Regev A. 2019. A quantitative framework for characterizing the evolutionary history of mammalian gene expression. Genome Research 29:53-63.

Clark NL, Aagaard JE, Swanson WJ. 2006. Evolution of reproductive proteins from animals and plants. Reproduction 131:11-22.

Coolon JD, McManus CJ, Stevenson KR, Graveley BR, Wittkopp PJ. 2014. Tempo and mode of regulatory evolution in *Drosophila*. Genome Research 24:797-808.

Cooper N, Thomas GH, Venditti C, Meade A, Freckleton RP. 2016. A cautionary note on the use of Ornstein Uhlenbeck models in macroevolutionary studies. Biological Journal of the Linnean Society 118:64-77.

De Bie T, Demuth JP, Cristianini N, Hahn MW. 2006. CAFE: a computational tool for the study of gene family evolution. Bioinformatics 22:1269-1271.

Dunn CW, Luo X, Wu Z. 2013. Phylogenetic analysis of gene expression. Integrative and Comparative Biology 53:847-856.

El Taher A, Böhne A, Boileau N, Ronco F, Indermaur A, Widmer L, Salzburger W. 2021. Gene expression dynamics during rapid organismal diversification in African cichlid fishes. Nature Ecology & Evolution 5:243-250.

Fantini E, Sulli M, Zhang L, Aprea G, Jiménez-Gómez JM, Bendahmane A, Perrotta G, Giuliano G, Facella P. 2018. Pivotal roles of cryptochromes 1a and 2 in tomato development and physiology. Plant Physiology 179:732-748.

Fay JC, Wittkopp PJ. 2008. Evaluating the role of natural selection in the evolution of gene regulation. Heredity 100:191-199.

Felsenstein J. 1973. Maximum likelihood and minimum-steps methods for estimating evolutionary trees from data on discrete characters. Systematic Biology 22:240-249.

FitzJohn RG. 2012. Diversitree: comparative phylogenetic analyses of diversification in R. Methods in Ecology and Evolution 3:1084-1092.

Freckleton RP. 2012. Fast likelihood calculations for comparative analyses. Methods in Ecology and Evolution 3:940-947.

Ge SX, Jung D, Yao R. 2020. ShinyGO: a graphical gene-set enrichment tool for animals and plants. Bioinformatics 36:2628-2629.

Gillespie JH. 1986. Variability of evolutionary rates of DNA. Genetics 113:1077-1091.

Goldberg EE, Kohn JR, Lande R, Robertson KA, Smith SA, Igić B. 2010. Species selection maintains self-incompatibility. Science 330:493-495.

Golding G. 1983. Estimates of DNA and protein sequence divergence: an examination of some assumptions. Molecular Biology and Evolution 1:125-142.

Gossmann TI, Saleh D, Schmid MW, Spence MA, Schmid KJ. 2016. Transcriptomes of plant gametophytes have a higher proportion of rapidly evolving and young genes than sporophytes. Molecular Biology and Evolution 33:1669-1678.

Gossmann TI, Schmid MW, Grossniklaus U, Schmid KJ. 2014. Selection-driven
evolution of sex-biased genes is consistent with sexual selection in *Arabidopsis*
thaliana. Molecular Biology and Evolution 31:574-583.

Greenway CA, Harder LD. 2007. Variation in ovule and seed size and associated size–
number trade-offs in angiosperms. American Journal of Botany 94:840-846.

Hahn MW, De Bie T, Stajich JE, Nguyen C, Cristianini N. 2005. Estimating the tempo
and mode of gene family evolution from comparative genomic data. Genome
Research 15:1153-1160.

Hahn MW, Demuth JP, Han S-G. 2007. Accelerated rate of gene gain and loss in
primates. Genetics 177:1941-1949.

Hahn MW, Nakhleh L. 2016. Irrational exuberance for resolved species trees. Evolution
70:7-17.

Han MV, Thomas GWC, Lugo-Martinez J, Hahn MW. 2013. Estimating gene gain and
loss rates in the presence of error in genome assembly and annotation using CAFE
3. Molecular Biology and Evolution 30:1987-1997.

Han QQ, Song YZ, Zhang JY, Liu LF. 2014. Studies on the role of the *SlNAC3* gene in
regulating seed development in tomato (*Solanum lycopersicum*). The Journal of
Horticultural Science and Biotechnology 89:423-429.

Harrison MC, Mallon EB, Twell D, Hammond RL. 2019. Deleterious mutation
accumulation in *Arabidopsis thaliana* pollen genes: a role for a recent relaxation of
selection. Genome Biology and Evolution 11:1939-1951.

Harrison PW, Wright AE, Zimmer F, Dean R, Montgomery SH, Pointer MA, Mank JE.
2015. Sexual selection drives evolution and rapid turnover of male gene expression.
Proceedings of the National Academy of Sciences 112:4393-4398.

Hibbins MS, Hahn MW. 2021. The effects of introgression across thousands of
quantitative traits revealed by gene expression in wild tomatoes. PLoS Genetics
17:e1009892.

Hill MS, Zande PV, Wittkopp PJ. 2021. Molecular and evolutionary processes
generating variation in gene expression. Nature Reviews Genetics 22:203-215.

Hiscott G, Fox C, Parry M, Bryant D. 2016. Efficient recycled algorithms for quantitative
trait models on phylogenies. Genome Biology and Evolution 8:1338-1350.

Huerta-Cepas J, Serra F, Bork P. 2016. ETE 3: reconstruction, analysis, and
visualization of phylogenomic data. Molecular Biology and Evolution 33:1635-1638.

Igić B, Bohs L, Kohn JR. 2006. Ancient polymorphism reveals unidirectional breeding
system shifts. Proceedings of the National Academy of Sciences 103:1359-1363.

King M-C, Wilson AC. 1975. Evolution at two levels in humans and chimpanzees.
Science 188:107-116.

Landis MJ, Schraiber JG. 2017. Pulsed evolution shaped modern vertebrate body sizes.
Proceedings of the National Academy of Sciences 114:13224-13229.

Li M, Wang X, Li C, Li H, Zhang J, Ye Z. 2018. Silencing *GRAS2* reduces fruit weight in
tomato. Journal of Integrative Plant Biology 60:498-513.

Liao Y, Smyth GK, Shi W. 2013. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. Bioinformatics 30:923-930.

Meisel RP, Malone JH, Clark AG. 2012. Disentangling the relationship between sex-
biased gene expression and X-linkage. Genome Research 22:1255-1265.

Mendes FK, Fuentes-González JA, Schraiber JG, Hahn MW. 2018. A multispecies
coalescent model for quantitative traits. eLife 7:e36482.

Mendes FK, Vanderpool D, Fulton B, Hahn MW. 2020. CAFE 5 models variation in
evolutionary rates among gene families. Bioinformatics 36:5516-5518.

Mione T, Anderson GJ. 1992. Pollen-ovule ratios and breeding system evolution in
Solanum section *Basarthrum* (Solanaceae). American Journal of Botany 79:279-
287.

Mitov V, Bartoszek K, Asimomitis G, Stadler T. 2020. Fast likelihood calculation for
multivariate Gaussian phylogenetic models with shifts. Theoretical Population
Biology 131:66-78.

Moyle LC, Wu M, Gibson MJ. 2021. Reproductive proteins evolve faster than non-
reproductive proteins among *Solanum* species. Frontiers in Plant Science
12:635990.

O'Meara BC, Ané C, Sanderson MJ, Wainwright PC. 2006. Testing for different rates of
continuous trait evolution using likelihood. Evolution 60:922-933.

Pease JB, Haak DC, Hahn MW, Moyle LC. 2016. Phylogenomics reveals three sources
of adaptive variation during a rapid radiation. PLoS Biology 14:e1002379.

Pennell MW, Eastman JM, Slater GJ, Brown JW, Uyeda JC, FitzJohn RG, Alfaro ME,
Harmon LJ. 2014. geiger v2. 0: an expanded suite of methods for fitting
macroevolutionary models to phylogenetic trees. Bioinformatics 30:2216-2218.

Pennell MW, FitzJohn RG, Cornwell WK, Harmon LJ. 2015. Model adequacy and the
macroevolution of angiosperm functional traits. The American Naturalist 186:E33-
E50.

Price PD, Palmer Droguett DH, Taylor JA, Kim DW, Place ES, Rogers TF, Mank JE,
Cooney CR, Wright AE. 2022. Detecting signatures of selection on gene expression.
Nature Ecology & Evolution 6:1035-1045.

Revell LJ, Harmon LJ. 2022. Phylogenetic Comparative Methods in R: Princeton
University Press.

Rick CM, Tanksley SD. 1981. Genetic variation in *Solanum pennellii*: Comparisons with
two other sympatric tomato species. Plant Systematics and Evolution 139:11-45.

Robinson MD, Oshlack A. 2010. A scaling normalization method for differential
expression analysis of RNA-seq data. Genome Biology 11:R25.

Rohlfs RV, Harrigan P, Nielsen R. 2014. Modeling gene expression evolution with an
extended Ornstein–Uhlenbeck process accounting for within-species variation.
Molecular Biology and Evolution 31:201-211.

Rohlfs RV, Nielsen R. 2015. Phylogenetic ANOVA: the expression variance and
evolution model for quantitative trait evolution. Systematic Biology 64:695-708.

Silvestro D, Kostikova A, Litsios G, Pearman PB, Salamin N. 2015. Measurement errors
should always be incorporated in phylogenetic comparative analysis. Methods in
Ecology and Evolution 6:340-346.

Somel M, Creely H, Franz H, Mueller U, Lachmann M, Khaitovich P, Pääbo S. 2008.
Human and chimpanzee gene expression differences replicated in mice fed different
diets. PLoS ONE 3:e1504.

Swanson WJ, Vacquier VD. 2002. The rapid evolution of reproductive proteins. Nature
Reviews Genetics 3:137-144.

Szövényi P, Ricca M, Hock Z, Shaw JA, Shimizu KK, Wagner A. 2013. Selection is no
more efficient in haploid than in diploid life stages of an angiosperm and a moss.
Molecular Biology and Evolution 30:1929-1939.

Tarashansky AJ, Musser JM, Khariton M, Li P, Arendt D, Quake SR, Wang B. 2021.
Mapping single-cell atlases throughout Metazoa unravels cell type evolution. eLife
10:e66747.

Vosters SL, Jewell CP, Sherman NA, Einterz F, Blackman BK, Moyle LC. 2014. The
timing of molecular and morphological changes underlying reproductive transitions in
wild tomatoes (*Solanum* sect. *Lycopersicon*). Molecular Ecology 23:1965-1978.

Wagner GP, Kin K, Lynch VJ. 2012. Measurement of mRNA abundance using RNA-seq
data: RPKM measure is inconsistent among samples. Theory in Biosciences
131:281-285.

Wang Z, Gerstein M, Snyder M. 2009. RNA-Seq: a revolutionary tool for
transcriptomics. Nature Reviews Genetics 10:57-63.

Whitehead MR, Lanfear R, Mitchell RJ, Karron JD. 2018. Plant mating systems often
vary widely among populations. Frontiers in Ecology and Evolution 6:38.

Wong A. 2011. The molecular evolution of animal reproductive tract proteins: What
have we learned from mating-system comparisons? International Journal of
Evolutionary Biology 2011:908735.

Wray GA, Hahn MW, Abouheif E, Balhoff JP, Pizer M, Rockman MV, Romano LA.
2003. The evolution of transcriptional regulation in eukaryotes. Molecular Biology
and Evolution 20:1377-1419.

Yang J, Ruan H, Xu W, Gu X. 2019. TreeExp2: an integrated framework for
phylogenetic transcriptome analysis. Genome Biology and Evolution 11:3276-3282.

Yang Z. 1996. Among-site rate variation and its impact on phylogenetic analyses.
Trends in Ecology & Evolution 11:367-372.

Figures and Tables

Figures and Tables

Figure 1. Bounded Brownian motion model. An example trait is shown in the bottom

graph, evolving along the tree shown above. Although the data input to CAGEE are

linear expression levels, internally it logs expression to ensure higher variance among

more highly expressed genes. There is also a minimum value, x_{min}, added to all tips.

Figure 2. Accuracy of CAGEE. For five different values of σ^2 we simulated 1000

datasets, with each dataset comprised of either 1000 genes or 10000 genes. All genes

in a dataset shared the same σ^2, but their values at the root were drawn independently

from the prior. We then provided each simulated dataset to CAGEE in order to infer σ^2.

Each box-and-whisker plot shows the mean (horizontal line), 50% interquartile range

(box), 1.5X the interquartile range (vertical lines), and outliers (dots).

Figure 3. Changes in gene expression along the tomato phylogeny. Given the set of

relationships among the seven *Solanum* accessions used here, we tested multiple

models that had branches assigned as different σ^2 parameters (Table 1). In model A, all

branches share σ_1^2. In model B, all black branches share σ_1^2, while all red branches

share σ_2^2. In model C, all black branches and the dashed red branch share σ_1^2, while all

solid red branches share σ_2^2. In model D, all black branches share σ_1^2, all solid red

branches share σ_2^2, and the dashed red branch is assigned σ_3^2. Using the results from

model D, we inferred the number of genes that had credible increases or decreases in

expression level along each branch (results for all changes are shown in Supplementary

Figure 4). Numbers are reported as +increases/-decreases for each branch.

Table 1. Model parameters estimated from the tomato data.
841

Model	Number of rates	-lnL	σ^2_1	σ^2_2	σ^2_3
A	1	67252.4	0.102		
B	2	65883.9	0.074	0.134	
C	2	65124.5	0.075	0.152	
D	3	65108.6	0.077	0.152	0.067

Supplementary Figures

Supplementary Figure 1. The tree used for simulations. The Newick-formatted tree string with branch lengths is:

((((sp1:1,sp2:1):1,sp3:2):1,sp4:3):1,((sp5:2,sp6:2):2)):1,sp7:5):1,sp8:6)

Supplementary Figure 2. Accuracy of CAGEE. A) This figure is the same as Figure 2 in the main text, but the ancestral state vector has length N=500 (Figure 2 uses N=200). B) For each of three different simulated values of σ^2, we randomly removed different amounts of data from an input dataset with 1,000 genes (the tree is the same as in all other simulations). C) For 1,000 genes simulated with σ^2=1 (dashed vertical line), we ran CAGEE independently on each one to estimate σ^2. D) We combined three datasets of 1,000 genes each simulated with three different values of σ^2 (we repeated these simulations 10 times). The 10 estimates of σ^2 on the combined datasets were slightly downwardly biased compared to the expected value (dashed horizontal line). Each dot represents each of the 10 estimates, with jitter added for clarity,

Supplementary Figure 3. Trees used for simulations with lineage-specific values of σ^2. A) All black branches share a rate parameter (σ_1^2), and all red branches share a rate parameter (σ_2^2). This "sigma_tree" is specified in CAGEE with the Newick string:

((((sp1:2,sp2:2):2,sp3:2):2,sp4:2):2,((sp5:1,sp6:1):1)):1,sp7:1):1,sp8:1)

B) All black branches share a rate parameter (σ_1^2), and all red branches share a rate parameter (σ_2^2). This "sigma_tree" is specified in CAGEE with the Newick string:

((((sp1:2,sp2:1):1,sp3:2):1,sp4:1):1,((sp5:1,sp6:2):1)):1,sp7:1):1,sp8:1)

Supplementary Figure 4. Changes in gene expression along the tomato phylogeny. This figure is the same as Figure 3 in the main text, but all increases and decreases are reported, regardless of whether they are "credible".

872   **Supplementary Figure 5.** Component diagram for the CAGEE software.

Figure 1

Figure 2

Figure 3

SI (σ^2_1)

SC (σ^2_2)

SC* (σ^2_3)

+405/-1047 — *S. habrochaites*

+8/-24

+321/-847 — *S. pennellii* (LA3778)

+68/-16

+406/-309 — *S. pennellii* (LA0716)

+275/-634 — *S. chilense*

+31/-17

+147/-257 — *S. chmielewskii*

+24/-14

+487/-1102

+42/-22 — *S. pimpinellifolium*

+819/-674 — *S. lycopersicum*

2.5 mya   2.0   1.5   1.0   0.5   0

Table 1

Model	Number of rates	-lnL	σ^2_1	σ^2_2	σ^2_3
A	1	67252.4	0.102		
B	2	65883.9	0.074	0.134	
C	2	65124.5	0.075	0.152	
D	3	65108.6	0.077	0.152	0.067

Supplementary Figure 1

Tree scale: 1

sp1 sp2 sp3 sp4 sp5 sp6 sp7 sp8

Supplementary Figure 2A

Supplementary Figure 2B

Supplementary Figure 2C

Supplementary Figure 2D

Supplementary Figure 3

A)

B)

Supplementary Figure 4

Supplementary Figure 5

Supplementary Table 1

Set of σ^2 values	Replicate	Simulated σ^2	Inferred σ^2	% error
0.2, 1.0	1	0.2	0.1926803	-3.66%
		1.0	1.0024905	0.249%
	2	0.2	0.1943218	-2.84%
		1.0	1.0054018	0.54%
	3	0.2	0.2031335	1.57%
		1.0	0.9577028	-4.23%
	4	0.2	0.1967767	-1.61%
		1.0	0.9996745	-0.0325%
	5	0.2	0.1924732	-3.76%
		1.0	1.0075440	0.754%
	6	0.2	0.2004951	0.248%
		1.0	1.0242412	2.42%
	7	0.2	0.1944284	-2.79%
		1.0	0.9733314	-2.67%
	8	0.2	0.2024781	1.24%
		1.0	0.9884935	-1.15%
	9	0.2	0.2018622	0.931%
		1.0	1.0382544	3.83%
	10	0.2	0.1956765	-2.16%
		1.0	1.0065474	0.655%
1.0, 3.0	1	1.0	0.9451556	-5.48%
		3.0	3.0287977	0.96%
	2	1.0	0.9769290	-2.31%
		3.0	3.1000229	3.33%
	3	1.0	0.9861053	-1.39%
		3.0	2.9452423	-1.83%
	4	1.0	0.9724728	-2.75%
		3.0	2.9203197	-2.66%
	5	1.0	0.9226307	-7.74%
		3.0	2.9930805	-0.231%
	6	1.0	0.9939074	-0.609%
		3.0	2.8856497	-3.81%
	7	1.0	0.9871113	-1.29%
		3.0	2.9705234	-0.983%
	8	1.0	0.9684494	-3.16%
		3.0	2.9187219	-2.71%
	9	1.0	0.9667607	-3.32%
		3.0	3.1521085	5.07%
	10	1.0	1.0059515	0.595%
		3.0	3.0232185	0.774%
5.0, 10.0	1	5.0	4.7464719	-5.07%
		10.0	9.9814701	-0.185%
	2	5.0	4.8651485	-2.7%
		10.0	10.0907926	0.908%
	3	5.0	4.7598429	-4.8%
		10.0	9.8058231	-1.94%
	4	5.0	4.9265027	-1.47%
		10.0	10.0593675	0.594%
	5	5.0	4.9239001	-1.52%
		10.0	9.9818898	-0.181%
	6	5.0	4.8684289	-2.63%
		10.0	9.8020147	-1.98%
	7	5.0	4.8752468	-2.5%
		10.0	10.0378953	0.379%
	8	5.0	4.9042695	-1.91%
		10.0	9.8541603	-1.46%
	9	5.0	4.6019327	-7.96%
		10.0	9.9286600	-0.713%
	10	5.0	4.8589572	-2.82%
		10.0	9.8313460	-1.69%