Brief Communication:

A therapeutic hepatitis B mRNA vaccine with strong immunogenicity and persistent

virological suppression

Huajun Zhao[1#✉], Xianyu Shao[2,3#], Yating Yu[1#], Lulu Huang[2,3#], Narh Philip Amor[2,3#], Kun Guo[2,3],

Changzhen Weng[2,3], Weijun Zhao[2,3], Ailu Yang[1], Jiesen Hu[4], Hongbao Yang[3], Zhenguang Liu[5],

Qiuju Han[1], Leilei Shi[6], Shiyu Sun[7], Jian Zhang[1✉], Yong Yang[2,3,8✉], Ang Lin[2,3✉]

Affiliations:

1. Institute of Immunopharmaceutical Sciences, Key Laboratory of Chemical Biology, School
of Pharmaceutical Sciences, Shandong University, Jinan, Shandong, China
2. Vaccine Center, School of Basic Medicine and Clinical Pharmacy, China Pharmaceutical
University, Nanjing, China
3. Center for New Drug Safety Evaluation and Research, China Pharmaceutical University,
Nanjing, China
4. Firestone Biotechnologies, Shanghai, China
5. College of Veterinary Medicine, Nanjing Agricultural University, Nanjing, China
6. Precision Research Center for Refractory Diseases in Shanghai General Hospital, Shanghai
Jiao Tong University, Shanghai, China
7. Key Laboratory of Infection and Immunity, Institute of Biophysics, Chinese Academy of
Sciences, Beijing, China.
8. School of Pharmacy, Xuzhou Medical University, Xuzhou, Jiangsu 221004, P.R. China.
#Authors contributed equally

Correspondence to:

Ang Lin, anglin@cpu.edu.cn; Yong Yang, yy@cpu.edu.cn; Jian Zhang, zhangj65@sdu.edu.cn;

Huajun Zhao, zhaohuajun89@sdu.edu.cn.

Lead contact:

Ang Lin, anglin@cpu.edu.cn

BRIEF ABSTRACT

Here we report on the development and comprehensive evaluations of an mRNA vaccine for chronic hepatitis B (CHB) treatment. In two different HBV carrier mouse models generated by viral vector-mediated HBV transfection (pAAV-HBV1.2 and rAAV8-HBV1.3), this vaccine demonstrates sufficient and persistent virological suppression, and robust immunogenicity in terms of induction of strong innate immune activation, high-level virus-specific antibodies, memory B cells and T cells. mRNA platform therefore holds prospects for therapeutic vaccine development to combat CHB.

INTRODUCTION

CHB is one major cause of liver fibrosis, cirrhosis, and hepatocellular carcinoma and poses a major public health threat[1]. Current CHB treatments such as nucleoside/nucleotide- and interferon-alpha (IFN-α)-based therapy, were unable to achieve sufficient viral clearance[2,3]. Multiple approaches to CHB therapeutic vaccines have been practiced intensively, including recombinant protein-based subunit, adenoviral vectored, and DNA-based vaccines, but yet achieved efficient seroclearance of HBsAg and seroconversion of anti-HBs antibody (Ab)[4,5,6]. mRNA vaccines that contain antigen-encoding mRNAs encapsulated into for example lipid nanoparticle (LNP) have shown superior immunogenicity in eliciting both Ab and cellular immune responses over other types of vaccines, and are also endowed with strong intrinsic adjuvant property to activate innate immune compartment[7]. Several mRNA-based prophylactic or therapeutic vaccines for infections, malignancies or other diseases are currently being evaluated in clinical trials [8, 9, 10, 11]. This platform also holds prospects for the development of therapeutic CHB vaccine that is expected to elicit potent anti-viral immunity mediating efficient virological suppression.

RESUTLS AND DISCUSSION

Recently, we reported a proprietary artificial intelligence-based algorithm that designs mRNA

with optimal folding stability and codon usage that together contribute to a high translation

efficiency [12]. Using this algorithm, a novel mRNA vaccine for CHB treatment was developed,

which is composed of Hepatitis B surface antigen (HBsAg)-encoding mRNAs encapsulated into

an ionizable lipid-based LNP through a well-established microfluidic system[13]. The mRNAs

were modified with N1-Methyl-pseudouridine and showed efficient protein expression reaching

a mean level of 980.6 mIU/ml upon transfection into HEK-293T cells (Supplemental Figure 1A).

Vaccine formulations were well characterized which demonstrated particle diameters of 96.3 ±

2.16 nm with average zeta potential of -1.92 mV and polymer dispersity index (PDI) below 0.2.

In addition, cells incubated with escalating concentrations of mRNA vaccines for 24 hours

maintained high viability, which suggested a limited cytotoxicity of vaccine (Supplemental

Figure 1B).

Therapeutic efficacy of the mRNA vaccine was first evaluated in pAAV-HBV1.2-transduced

HBV-carrier mice. This model shows systemic immune tolerance and long-lasting HBV viremia

that largely resemble asymptomatic chronic HBV-infected individuals and has therefore been

widely used in the study of CHB immunotherapy[14,15]. HBV-carrier mice were administered

intramuscularly (i.m.) with three doses of 5μg or 10μg mRNA vaccines at a 1-week interval

(Figure 1A). Compared to PBS-treated mice showing a high serum HBsAg level, mRNA

vaccine-treated mice demonstrated a rapid decline of serum HBsAg, which was even

undetectable 7 days after the 3[rd] dose (Figure 1B). Clinical management of CHB remains

challenging largely due to HBV recurrence and the failure to achieve seroconversion of anti-HBs

Abs[2, 3]. Magnitude of anti-HBs Ab response was therefore evaluated longitudinally. Three doses

of HBV mRNA vaccines elicited robust levels of anti-HBs Abs reaching a mean titer of 3624.0

mIU/ml and 4804.6 mIU/ml in the 5μg and 10μg vaccine groups at day 59, respectively (Figure

1C).

To evaluate whether the mRNA vaccine could induce sustained protective responses against viral

re-exposure, the mice were re-injected with pAAV-HBV1.2 plasmids on day 60 mimicking viral

challenge. As expected, PBS-treated mice still demonstrated a high level of serum HBsAg. While

all vaccinated mice were fully protected against viral re-exposure showing no detectable level of

serum HBsAg (Figure 1D), which was accompanied by further elevated anti-HBs Ab titers

(Figure 1E). Moreover, copies of intrahepatic HBV cccDNA, total DNA, total RNA, 3.5kb RNA

(Figure 1F) and serum HBV DNA (Figure 1G) were clearly reduced in vaccinated mice and the

expression of HBV core antigen (HBcAg) in liver tissues was largely decreased (Figure 1H).

Since AAV-transduced-HBV-carrier mice cannot produce significant amounts of cccDNA, the

anti-HBV effect on eliminating HBV cccDNA should be further evaluated in specific animal

models, such as rc-cccDNA mouse model[16] or human liver chimeric mouse model[17].

Efficacy of the mRNA vaccine was next compared side-by-side with conventional therapeutic

Entecavir. Compared to the rapid anti-HBs Ab production and serum HBsAg clearance induced

by mRNA vaccine, Entecavir administered orally for consecutive 15 days showed no ability in

inducing anti-HBs Abs or eliminating serum HBsAg (Supplemental Figure 2), which was in line

with previous studies[18]. Considering that viral clearance can be mediated through

vaccine-induced cytotoxic immune responses potentially causing liver damage, serum alanine

transaminase (ALT) and aspartate aminotransferase (AST) levels were monitored longitudinally

(Supplemental Figure 3A-B). A transient elevation of serum ALT was observed in a few animals

7 days post the 1st and 2nd vaccination. While serum AST remained at normal levels in all treated

mice during the period of treatment. In addition, histopathological analysis revealed that there

was no hepatotoxicity or liver injury induced upon the 3-dose vaccine treatment (Supplemental

Figure 3C). These suggested that the mRNA vaccine may potentially function through a

non-cytotoxic mechanism to eliminate virus. Previously, we and others showed that

IFN-γ-producing HBV-specific CD8+ T cells could mediate HBV clearance via a non-cytotoxic

mechanism without causing liver damage[19,20]. Since both cytotoxic and non-cytotoxic immune

responses to HBV are important to eliminate virus, the underlying mechanisms of action of the

mRNA vaccine would merit further investigation.

Therapeutic efficacy of the mRNA vaccine was further evaluated in rAAV8-HBV1.3-transduced

HBV-carrier mouse model (Figure 2A), which shows more efficient and homogeneous HBV

transduction than the aforementioned pAAV-HBV1.2 mice model[21]. As validated in our study,

rAAV8-HBV1.3-transduced mice showed persistent HBV surface antigenemia lasting for more

than 200 days (Supplemental Figure 4) and a typical immunotolerant status represented by high

frequencies of hepatic $CD4^+CD25^+$ $Foxp3^+$ T cells (Supplemental Figure 4B), elevated

expression of inhibitory immune-checkpoint molecules (PD-1, LAG-3, and TIM-3), and

impaired cytokine production by $CD4^+$ T and $CD8^+$ T cells upon stimulation (Supplemental

Figure 4C and D). All these features well mimic the immunotolerant state of human chronic

HBV carriers[21]. Upon three doses of vaccination, HBV1.3-carrier mice demonstrated a rapid and

sufficient serum HBsAg clearance, and the virological suppression was maintained for at least

208 days within the period of observation (Figure 2B). Furthermore, serum HBeAg level was

remarkably reduced in both two groups of vaccinated mice (Figure 2C). Serum HBV DNA

copies were significantly reduced in mice receiving 10μg mRNA vaccine and showed a trend of

decrease in the 5μg dosing group (Figure 2D). Notably, majority of the 10μg mRNA

vaccine-treated mice still presented high levels of anti-HBs Abs when detected 208 days after

treatment initiation (Figure 2E). Since serum HBsAg was sufficiently cleared, but serum HBV

DNA was still present which implied that viral replication may still remain active at a very low

level although was largely restricted. The mRNA vaccine-elicited anti-viral immunity remains to

be further assessed for the efficacy to control viral genome transcription using more suitable

animal models.

To further validate the potency and superiority of the mRNA vaccine, we next performed a

side-by-side comparison with our recently reported recombinant CHB therapeutic vaccines

(Sim+rHBV)[22] in rAAV8-HBV1.3 mice. Three doses of Sim+rHBV vaccine showed a moderate

ability in viral clearance but much less efficient than the mRNA vaccine (Supplemental Figure 5).

Seven days post the 3rd dose, serum HBsAg levels in mice receiving 5μg mRNA vaccine, 10μg

mRNA vaccine and Sim+rHBV vaccine were present at a mean value of 15.8 mIU/ml, 16.0

mIU/ml and 166.0 mIU/ml, respectively. In addition, evidence from other groups testing their

in-house developed therapeutic CHB vaccines using same animal models indirectly supported

the superior therapeutic effect of mRNA vaccine[23, 24].

An optimal therapeutic vaccine for CHB treatment should have the potentials to trigger efficient

innate immune activation leading to robust antigen presentation and the resulting generation of

HBV-specific cellular immunity[12]. To this end, we studied the innate immune responses induced

at early time point (12 hours) after prime immunization in rAAV8-HBV1.3 mice and observed an

increased infiltration of dendritic cell (DC) subsets (CD8α^{+} cDC1s, CD103^{+} cDC1s and CD11b^{+}

cDC2s) and macrophages into spleen, accompanied by potent cell maturation (Supplemental

Figure 6A-D). Other innate immune cell subsets including monocytes and neutrophils

(Supplemental Figure 6E-G) also showed a phenotypic maturation, indicated by elevated

expression of CD80 and CD86. T cell exhaustion is a hallmark of CHB infection and the restored

HBV-specific CD8^{+} T cell function has been widely-accepted to be predictive for the efficacy of

therapeutic vaccines for CHB[25]. We therefore assessed whether the therapeutic mRNA

vaccination could promote HBV-specific T cell responses and break the exhaustion. In

rAAV8-HBV1.3 mice, three doses of mRNA vaccines induced robust levels of Th1-biased CD4^{+}

and CD8^{+} T cells producing IFN-γ or IL-2 favoring viral elimination (Supplemental Figure

7A-C). In addition, frequencies of HBsAg-specific memory B cells (MBCs) in spleens were

obviously elevated upon vaccination (Supplemental Figure 7D) which was associated with the

strong seroconversion and long-term protection as observed earlier in this study (Supplemental

Figure 2).

Collectively, we reported an HBV mRNA vaccine candidate with potent therapeutic efficacy and

strong immunogenicity. To our knowledge, this is the first reported therapeutic mRNA vaccine

candidate for CHB treatment. Three doses of HBV mRNA vaccines could efficiently and

persistently eliminate HBV and achieve a long-term seroconversion of anti-HBs Ab, and most

importantly showed full protection against subsequent viral re-exposure. The strong innate

immune activation and generation of robust functional HBV-specific T cells and MBCs by the

mRNA vaccine may hold prospects for functional cure of CHB and prevention of HBV

recurrence. However, further in-depth assessment of the mRNA vaccine would be needed to

evaluate the effects on restricting viral replication at genomic levels especially the potentiality to

eliminate HBV cccDNA pool. In addition, synergistic efficacy of the mRNA vaccine in

combinatorial use with other types of CHB therapeutics merits further investigation.

METHODS

A detailed description and additional materials and methods are available in supplemental

materials.

Ethics, animals, treatments

C57BL/6J mice (5-6 weeks old, male) were purchased from Beijing HFK Bioscience Co. Ltd.

(Beijing, China). HBV-carrier mouse models were generated either through hydrodynamic

injection of a volume of saline (equivalent to 10% body weight) containing 8μg pAAV-HBV 1.2

plasmid (kindly provided by Pei-Jer Chen; National Taiwan University College of Medicine,

Taipei, Taiwan) or intravenous injection of 1×10^{10} vector genome equivalent of

rAAV8-HBV1.3, as previously described[21,23]. Serum HBsAg levels were measured 6 weeks after

the hydrodynamic or intravenous injection, and mice with serum HBsAg levels **> 500 mIU/mL**

were defined as HBV-carrier mice and used for subsequent in-vivo experiments. HBV-carrier

mice were randomly allocated to different groups and were i.m. injected with three doses of 5μg

or 10μg mRNA vaccines at a one-week interval. HBV-carrier mice administered with PBS were

used as control. In some experiments, HBV-carrier mice were orally treated with Entecavir

(50μg/kg, Selleck Chem) for 15 days consecutively or were immunized with a recombinant

therapeutic vaccine (Sim+rHBV)[21] containing 2 μg rHBVvac adjuvanted with 100 μg simvastatin.

Sera samples were collected at different time points and were stored at -80 ℃ for further use. For

the evaluation of long-term protective response, the immunized HBV-carrier mice were

hydrodynamically re-injected with 8μg pAAV-HBV1.2 plasmids on day 60 after treatment

initiation. All animal experiments were performed in accordance with the Guidelines for the Care

and Use of Laboratory Animals and the Ethical Committee of Shandong University and using

protocols approved by the Institutional Animal Care and Use Committee of Shandong University

(approval number: 20023).

mRNA vaccine preparation

mRNAs encoding for HBsAg (Genbank accession number: MK213856.1) were synthesized by

T7 polymerase-mediated in vitro transcription (IVT) based on a linearized DNA template

containing codon-optimized HBsAg gene flanked with 5' and 3' untranslated regions (UTRs) and

a 100 nt poly-A tail. During IVT procedure, mRNAs were modified with

N1-Methyl-pseudouridine (Synthgene) and capped using CleanCap Reagent (TriLink). After this,

IVT products were purified with Monarch RNA purification columns (NEW ENGLAND

BioLabs Inc. MA, USA) and resuspended in a TE buffer at a desired concentration. For mRNA

encapsulation into LNP, lipid components were dissolved in ethanol at molar ratios of

50:10:38.5:1.5 (ionizable lipid: DSPC: cholesterol: DMG-PEG2000). The novel ionizable lipid

(YX-02) was designed and has been patented by Firestone Biotechnologies. The lipid cocktail

was mixed with mRNAs dissolved in 10mM citrate buffer (pH4.0) at an N/P ratio of 5.3 :1 and a

volume ratio of 3: 1 using a microfluidic-based equipment (INano^{TM}L from Micro&Nano

Biologics) at a total flow rate of 12 mL/min. Formulations were diluted with PBS and

ultrafiltrated using 50-kDa Amicon ultracentrifugal filters. Vaccine formulation was

characterized for particle diameter, polymer dispersity index (PDI) and zeta potentials using

NanoBrook Omni ZetaPlus (Brookhaven Instruments).

DATA AVAILABILITY

All data are available upon reasonable request to the corresponding authors.

ACKNOWLEDGEMENTS

This work was supported by the Natural Science Foundation of Jiangsu Province (BK20221031 to A.L.), the National Natural Science Foundation of China (32200764, 82061138008), the National Postdoctoral Program for Innovative Talents (BX20190192, to J.H.Z.), the Fundamental Research Funds for the Central Universities (2632022YC01, to A.L.), the National Key Research and Development Program (2021YFC2300603, to J.Z.), the National Science Foundation for Young Scientists of China (82001687, to J.H.Z.), the Shandong Provincial Natural Science Foundation for The Excellent Youth Scholars (ZR2022YQ75, to J.Q.H), and the National Natural Science Foundation of China (81972686, to J.Q.H). This study was also supported by the Projects of State Key Laboratory of Natural Medicines (SKLNMZZ2023XX) and State Key Laboratory of Microbial Technology (M2023-13). We thank the Pharmaceutical Biology Sharing Platform and the Translational Medicine Core Facility of Shandong University and China Pharmaceutical University for instrumental support. We also thank Firestone Biotechnology Co., Ltd for technical assistance with the development of novel ionizable lipids and thank Dr. Liang Zhang for the assistance in codon optimization of mRNA using LinearDesign Algorithm.

AUTHOR CONTRIBUTIONS

A.L., H.Z., Y.Y. and J.Z. designed the project; H.Z., X.S., Y.Y., L.H., N.A., C.W., W.Z., A.Y., J.H., Q.H., L.S. and S.S. performed experiments; H.Z., A.L., X.S., T.Y., B.Y. and N.A. performed data analysis; A.L., Y.Y., and H.Z. discussed the data; A.L. H.Z., Y.S. and L.H. wrote the manuscript.

COMPETING INTERESTS

J.H. is a full-time employee at Firestone Biotechnologies, Shanghai. The LinearDesign Algorithm was patented and owned by Baidu USA. China Pharmaceutical University and Shandong University have filed a provisional patent for the HBV mRNA vaccine listing H.Z., A.L., L.H., J.Z., Y.Y. as inventors. Other authors declare no competing interests.

REFERENCES

1. Organization WH. Global progress report on HIV, viral hepatitis and sexually transmitted infections. (2021).
2. Grossi G, Vigano M, Loglio A, Lampertico P. Hepatitis B virus long-term impact of antiviral therapy nucleot(s)ide analogues (NUCs). *Liver Int* **37 Suppl 1**, 45-51 (2017).
3. Konerman MA, Lok AS. Interferon Treatment for Hepatitis B. *Clin Liver Dis* **20**, 645-665 (2016).
4. Wang J, Gujar SA, Cova L, Michalak TI. Bicistronic woodchuck hepatitis virus core and gamma interferon DNA vaccine can protect from hepatitis but does not elicit sterilizing antiviral immunity. *J Virol* **81**, 903-916 (2007).
5. Martin P, *et al.* TG1050, an immunotherapeutic to treat chronic hepatitis B, induces robust T cells and exerts an antiviral effect in HBV-persistent mice. *Gut* **64**, 1961-1971 (2015).
6. Vesikari T, *et al.* Immunogenicity and safety of a tri-antigenic versus a mono-antigenic hepatitis B vaccine in adults (PROTECT): a randomised, double-blind, phase 3 trial. *Lancet Infect Dis* **21**, 1271-1281 (2021).
7. Li C, *et al.* Mechanisms of innate and adaptive immunity to the Pfizer-BioNTech BNT162b2 vaccine. *Nat Immunol* **23**, 543-555 (2022).
8. Lee IT, *et al.* Safety and immunogenicity of a phase 1/2 randomized clinical trial of a quadrivalent, mRNA-based seasonal influenza vaccine (mRNA-1010) in healthy adults: interim analysis. *Nat Commun* **14**, 3631 (2023).
9. Rojas LA, *et al.* Personalized RNA neoantigen vaccines stimulate T cells in pancreatic cancer. *Nature* **618**, 144-150 (2023).

10.   mRNA Vaccine Slows Melanoma Recurrence. *Cancer Discov* **13**, 1278 (2023).

11.   Aliprantis AO*, et al.* A phase 1, randomized, placebo-controlled study to evaluate the
safety and immunogenicity of an mRNA-based RSV prefusion F protein vaccine in
healthy younger and older adults. *Hum Vaccin Immunother* **17**, 1248-1261 (2021).

12.   Zhang H*, et al.* Algorithm for optimized mRNA design improves stability and
immunogenicity. *Nature* **621**, 396-403 (2023).

13.   Yang R*, et al.* A core-shell structured COVID-19 mRNA vaccine with favorable
biodistribution pattern and promising immunity. *Signal Transduct Target Ther* **6**, 213
(2021).

14.   Huang LR, Wu HL, Chen PJ, Chen DS. An immunocompetent mouse model for the
tolerance of human chronic hepatitis B virus infection. *Proc Natl Acad Sci U S A* **103**,
17862-17867 (2006).

15.   Zhang TY*, et al.* Prolonged suppression of HBV in mice by a novel antibody that targets
a unique epitope on hepatitis B surface antigen. *Gut* **65**, 658-671 (2016).

16.   Wang L*, et al.* Discovery of a first-in-class orally available HBV cccDNA inhibitor. *J
Hepatol* **78**, 742-753 (2023).

17.   Yang G*, et al.* HAT1 signaling confers to assembly and epigenetic regulation of HBV
cccDNA minichromosome. *Theranostics* **9**, 7345-7358 (2019).

18.   Michler T*, et al.* Knockdown of Virus Antigen Expression Increases Therapeutic Vaccine
Efficacy in High-Titer Hepatitis B Virus Carrier Mice. *Gastroenterology* **158**, 1762-1775
e1769 (2020).

19.   Xia Y*, et al.* Interferon-gamma and Tumor Necrosis Factor-alpha Produced by T Cells
Reduce the HBV Persistence Form, cccDNA, Without Cytolysis. *Gastroenterology* **150**,
194-205 (2016).

20.   Zhao HJ*, et al.* Poly I:C-based rHBVvac therapeutic vaccine eliminates HBV via
generation of HBV-specific CD8+ effector memory T cells. *Gut.* **68**, 2032-2043 (2019).

21.   Yang D*, et al.* A mouse model for HBV immunotolerance and immunotherapy. *Cell Mol
Immunol* **11**, 71-78 (2014).

22.   Zhao H*, et al.* Cholesterol accumulation on dendritic cells reverses chronic hepatitis B
virus infection-induced dysfunction. *Cell Mol Immunol* **19**, 1347-1360 (2022).

23.   Meng CY*, et al.* Engineered anti-PDL1 with IFNalpha targets both immunoinhibitory and
activating signals in the liver to break HBV immune tolerance. *Gut* **72**, 1544-1554
(2023).

24.   Wang W*, et al.* Dual-targeting nanoparticle vaccine elicits a therapeutic antibody
response against chronic hepatitis B. *Nat Nanotechnol* **15**, 406-416 (2020).

25.   Bengsch B, Martin B, Thimme R. Restoration of HBV-specific CD8+ T cell function by
PD-1 blockade in inactive carrier patients is linked to T cell differentiation. *J Hepatol* **61**,
1212-1219 (2014).

Figure 1

Figure 1. HBV mRNA vaccine induced efficient viral suppression and achieved robust seroconversion in pAAV-HBV1.2 mice.

(A) Study design. pAAV-HBV1.2-transduced HBV-carrier mice (n=5/group) were immunized i.m. with 5μg or 10μg HBV mRNA vaccines three times at a 1-week interval. HBV-carrier mice administered with PBS were used as control. At day 60, mice were hydrodynamically re-injected with 8μg pAAV-HBV1.2 plasmids. Sera samples were collected at the indicated time points. (B, D) Levels of serum HBsAg and (C, E) anti-HBs Abs were measured at the indicated time points. (F) Seven days after viral re-exposure, intrahepatic HBV cccDNA, total DNA, total RNA and

329 3.5-kb RNA were analyzed. (G) Seven days after viral re-exposure, serum HBV DNA was

330 quantified. (H) Seven days after viral re-exposure, HBcAg expression in liver tissues was

331 determined by IHC staining (magnification: ×200; scale bar: 100 µm). GAPDH, glyceraldehyde

332 3-phosphate dehydrogenase; SEM, standard error of the mean. Data are shown as Mean ± SEM.

333 $*p \leq 0.05$, $**p \leq 0.01$, $****p \leq 0.0001$.

Figure 2

Figure 2. HBV mRNA vaccine induced efficient and sustained viral suppression and achieved robust seroconversion in rAAV8-HBV1.3 mice.

(A) Study design. rAAV-HBV1.3-transduced HBV-carrier mice (n=6/group) were immunized i.m. with 5μg or 10μg HBV mRNA vaccines three times at a 1-week interval. HBV-carrier mice administered with PBS were used as control. Sera samples were collected longitudinally. (B) Levels of serum HBsAg were measured at the indicated time points. (C-E) 208 days after treatment start, levels of serum HBeAg (C), serum HBV DNA copies (D), and anti-HBs Abs (E) were measured. Data are shown as Mean ± SEM. *$p \leq 0.05$, **$p \leq 0.01$. PEIU represents Paul-Ehrlich-Institute Unit.

A HBV-carrier mice

Group	Immunization (Day 1,8,15)
●	PBS
▼	5μg mRNA vac.
▲	10μg mRNA vac.

HBV-carrier mice (rAAV8-HBV1.3)

prime boost boost

Serum collection

B Serum HBsAg log$_{10}$ (mIU/mL) vs Day

C Day 208 — Serum HBeAg (PEIU/mL) — PBS, 5μg, 10μg mRNA vac. — ** **

D Day 208 — Serum HBV DNA copies (IU/ml) — PBS, 5μg, 10μg mRNA vac. — ns **

E Day 208 — anti-HBs Ab (mIU/ml) — PBS, 5μg, 10μg mRNA vac. — * * *