1   **Epigenetic and genetic differentiation between *Coregonus* species pairs**

2   Clare J Venney[1], Claire Mérot[1,2], Eric Normandeau[1], Clément Rougeux[1], Martin Laporte[1,3],

3   Louis Bernatchez[1,4]

1. Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, Canada

2. UMR 6553 Ecobio, OSUR, CNRS, Université de Rennes, Rennes, France

3. Ministère des Forêts, de la Faune et des Parcs (MFFP), Québec, Québec

4. This author is deceased.

Author for correspondence: Clare Venney (clarevenney@gmail.com)

Abstract

Phenotypic diversification is classically associated with genetic differentiation and gene

expression variation. However, increasing evidence suggests that DNA methylation is involved

in evolutionary processes due to its phenotypic and transcriptional effects. Methylation can

increase mutagenesis and could lead to increased genetic divergence between populations

experiencing different environmental conditions for many generations, though there has been

minimal empirical research on epigenetically induced mutagenesis in diversification and

speciation. Whitefish, freshwater members of the salmonid family, are excellent systems to study

phenotypic diversification and speciation due to the repeated divergence of benthic-limnetic

species pairs serving as natural replicates. Here we investigate whole genome genetic and

epigenetic differentiation between sympatric benthic-limnetic species pairs in lake and European

whitefish (*Coregonus clupeaformis* and *C. lavaretus*) from four lakes (N=64). We found

considerable, albeit variable, genetic and epigenetic differences between species pairs. All SNP

types were enriched at CpG sites supporting the mutagenic nature of DNA methylation, though

C>T SNPs were most common. We also found an enrichment of overlaps between outlier SNPs

with the 5% highest F_{ST} between species and differentially methylated loci. This could possibly

represent differentially methylated sites that have caused divergent genetic mutations between

species, or divergent selection leading to both genetic and epigenetic variation at these sites. Our

results support the hypothesis that DNA methylation contributes to phenotypic divergence and

mutagenesis during whitefish speciation.

Keywords: DNA methylation, speciation, whitefish, mutagenesis, genetic divergence, genetic

assimilation

Significance statement

DNA methylation is an epigenetic mark known to change in response to the environment

and induce genetic modifications such as point mutations, though its implications for evolution

and speciation have not been thoroughly studied. We find considerable but variable genetic and

epigenetic variation between whitefish benthic-limnetic species pairs, highlighting the potential

for DNA methylation to contribute to mutagenesis and genetic evolution. Our study provides

evidence that DNA methylation could have contributed to whitefish speciation, both through

initially plastic methylation changes and by driving genetic divergence between species pairs.

Introduction

Speciation has long been a focus of evolutionary biology, with recent research expanding

into the role of genomics in reproductive isolation, phenotypic diversification, and species

divergence (Seehausen et al. 2014; Marques et al. 2019). Speciation can occur rapidly despite

slow mutation rates, sometimes due to new combinations of standing genetic variation (Marques

et al. 2019). Phenotypic plasticity can also promote speciation, particularly when allopatric

populations acclimate to their respective environments, phenotypes are partially genetically

controlled, and capacity for plasticity is lost over time (Pfennig et al. 2010). Phenotypic changes

can occur through altered gene expression (Whitehead & Crawford 2006) which can lead to

phenotypic divergence and speciation, as documented between Arctic charr ecotypes (*Salvelinus*

alpinus; Gudbrandsson et al. 2018; Jacobs and Elmer 2021), between Chinook and Coho salmon

and their hybrids (*Oncorhynchus tshawytscha* and *O. kisutch*; Mckenzie et al. 2021), and

between sympatric lake whitefish species (*Coregonus* sp.; Derome et al. 2006; St-Cyr et al.

2008; Rougeux et al. 2019b). Transcriptomic variation associated with lip size has been

identified in the Midas cichlid species complex (*Amphilophus citrinellus;* Manousaki et al. 2013)

and sequence variation in transcribed regions has been reported between *A. astorquii* and *A.*

zaliosus (Elmer et al. 2010). Transcriptomic differences generally increase with taxonomic

distance for closely related species (Whitehead & Crawford 2006) and may contribute to

phenotypic differences among species (Whitehead & Crawford 2006; Pavey et al. 2010).

Therefore, the mechanisms underlying adaptive phenotypic differences during speciation likely

involve both phenotypic plasticity and genetic variants.

There has been increasing interest in the role of DNA methylation as a plastic mechanism

contributing to speciation (Vogt 2017; Ashe et al. 2021). DNA methylation is the addition of a

methyl group to the cytosine of CpG sites in vertebrates (a cytosine followed by a guanine in the

DNA sequence), often resulting in altered transcription without a change in DNA sequence (Bird

2002). DNA methylation is sensitive to the environment and epigenetic marks are known to be

affected by several factors such as temperature (Ryu et al. 2020; Beemelmanns et al. 2021;

McCaw et al. 2020; Venney et al. 2022), salinity (Hu et al. 2021; Artemov et al. 2017; Heckwolf

et al. 2020), and rearing environment (Leitwein et al. 2021; Le Luyer et al. 2017; Wellband et al.

2021; Berbel-Filho et al. 2020; Venney et al. 2020). DNA methylation can alter phenotype

(Anastasiadi et al. 2021; Vogt 2021), with evidence for phenotypically-linked epigenetic

divergence associated with spawning tactics in capelin (*Mallotus villosus*; (Venney et al. 2023)

and between four sympatric Arctic charr morphs (Matlosz et al. 2022). Due to the potential role

of DNA methylation in plasticity and phenotypic diversification, it may contribute to speciation

(Vogt 2017; Ashe et al. 2021; Stajic & Jansen 2021; Laporte et al. 2019). A recent simulation

study showed that epigenetic plasticity can promote speciation when it reduces the fitness of

migrants and hybrids but can prevent genetic adaptation and speciation if epigenetic adaptation

occurs, precluding the need for genetic adaptation (Greenspoon et al. 2022). Empirical evidence

for the role of DNA methylation in reproductive isolation and speciation is also emerging

(Laporte et al. 2019). Methylation differences detected through methylation-sensitive amplified

polymorphism, but not genetic differences, were predictive of behavioural isolation between 16

species of darters (*Ulocentra*, *Nanostoma,* and *Etheostoma*; Smith et al. 2016). Another study

showed considerable epigenetic differences between six phenotypically divergent species of

Lake Malawi cichlids (Vernaz et al. 2022). DNA methylation can affect transcription (Li et al.

2019) and phenotype (Anastasiadi et al. 2021; Vogt 2021) and could result in initial plastic

phenotypic responses that lead to phenotypic diversification and speciation over generations.

DNA methylation is also mutagenic and can generate polymorphism. This is partially due

to the spontaneous hydrolytic deamination of methylated cytosine to uracil which is rapidly

converted to a thymine tautomer if not corrected by DNA repair enzymes (Gorelick 2003).

Spontaneous deamination is ~3.5 times more likely to occur at methylated cytosines than

unmethylated ones (Jones et al. 1992; Gorelick 2003). Different enzymes are involved in base

repair of methylated vs unmethylated cytosines, leading to methylated cytosines having mutation

rates ~20 000 times higher than unmethylated ones after accounting for DNA repair efficiency

(Gorelick 2003). It is estimated that ~8 deamination events occur per day in the ~6 billion bp

diploid human genome (Jones et al. 1992) providing a consequential source of novel mutations,

particularly in larger genomes. C>T transitions are the most common and explainable

epigenetically induced mutation, though there is also evidence for increased C>A and C>G

mutations at CpG sites due to mutagen exposure (Tomkova & Schuster-Böckler 2018). On the

other hand, DNA methylation can also shield sites from mutagenesis depending on the stimulus

or trigger, though the intricacies of when mutagenesis is favoured or prevented remain unclear

(Tomkova & Schuster-Böckler 2018). A study in human cell lines (*Homo sapiens*) showed that

cytosines with intermediate levels of methylation (20-60%) had the highest mutation rate, even

compared to fully methylated sites (Xia et al. 2012). The shielding properties of DNA

methylation could also be due to the inability to discern DNA methylation from other

methylation-related marks. In particular, 5-methylcytosine can be converted to 5-

hydroxymethylcytosine during demethylation (Li et al. 2019). The two cannot be differentiated

through bisulfite sequencing (Li et al. 2019), though 5-hydroxymethylcytosine may protect CpG

sites from mutation (Tomkova & Schuster-Böckler 2018). It is also possible that higher

nucleotide diversity at CpG sites is associated with greater methylation variation at those sites,

suggesting weaker selective constraint at those sites (Ord et al. 2023). Conversely, highly

methylated sites may be under greater selection to maintain consistently high methylation levels

(Ord et al. 2023). Therefore, DNA methylation can not only lead to transcriptional and

phenotypic changes but may also influence mutation rates. DNA methylation has thus been

proposed as a mechanism for genetic assimilation of phenotypes, i.e., when an environmentally

induced phenotype becomes stable and genetically encoded, even in the absence of the original

stimulus (Danchin et al. 2019; Nishikawa & Kinjo 2018). However, empirical evidence for

methylated sites inducing point mutations and contributing to evolution remains sparse.

The two whitefish sister taxa, lake whitefish (*Coregonus clupeaformis*) in North America

and the European whitefish species complex (*C. lavaretus*) in Europe, are well-characterized and

relevant systems to study phenotypic diversification and speciation. Lake and European

whitefish have evolved separately since they became geographically isolated ~500 000 years ago

(Bernatchez & Dodson 1994, 1991; Jacobsen et al. 2012). Both show repeated independent

divergence of sympatric species complexes consisting of a putatively ancestral benthic and

derived limnetic species originating from different glacial refugia during the last glaciation

period (Østbye, Bernatchez, et al. 2005; Bernatchez et al. 2010; Bernatchez & Dodson 1991;

Rougeux et al. 2017; Pigeon et al. 1997) which came into secondary contact ~12 000 years ago

when colonizing postglacial lakes (Rougeux, Gagnaire, Praebel, et al. 2019; Rougeux et al.

2017). In North America, the derived limnetic species evolved to colonize the limnetic zone,

leading to differences in diet (Bernatchez et al. 1999), reduced size (Bernatchez et al. 1999),

slower growth (Trudel et al. 2001), more slender body morphology (Laporte et al. 2015, 2016),

higher metabolic rate (Trudel et al. 2001; Dalziel et al. 2015), and more active swimming

behaviour (Rogers et al. 2002). Benthic and limnetic species often coexist in Europe, though

some Fenno-Scandinavian and alpine lakes contain up to six sympatric whitefish species (De-

Kayne et al. 2022; Østbye, Næsje, et al. 2005). In both North America and Europe, sympatric

whitefish are generally reproductively isolated with variable amounts of gene flow depending on

the lake (Rougeux et al. 2017; Rogers & Bernatchez 2006) which translates into genetic

differentiation (Mérot et al. 2023; Rougeux, Gagnaire, Praebel, et al. 2019; Rogers & Bernatchez

2007; Bernatchez et al. 2010; Dion-Côté et al. 2017; Rougeux et al. 2017; Østbye, Næsje, et al.

2005; De-Kayne et al. 2022; Siwertsson et al. 2013; Østbye et al. 2006), differential transposable

element methylation (Laporte et al. 2019), and transcriptional differences (Rougeux, Gagnaire,

Praebel, et al. 2019; Derome et al. 2006; Jeukens et al. 2009). Benthic-limnetic species pairs in

different lakes thus present a naturally replicated system in which to study the molecular

mechanisms associated with speciation.

We assessed the role of genomic and epigenomic variation in whitefish speciation by

performing whole genome sequencing (WGS) and whole genome bisulfite sequencing (WGBS)

for four benthic-limnetic species pairs (N=64 individuals sequenced in both datasets): lake

whitefish from Cliff and Indian Lake, Maine, USA, and European whitefish from Langfjordvatn

Lake, Norway and Zurich Lake, Switzerland (Figure 1). The objectives of this study were to (i)

determine the extent of whole genome genetic divergence between benthic-limnetic whitefish

species pairs, (ii) measure the level of polymorphism at CpG sites relative to the rest of the

genome, (iii) characterize whole genome differences in DNA methylation among limnetic-

benthic whitefish species pairs, and (iv) assess whether there was an enrichment of outlier SNPs

at differentially methylated loci (DMLs), which would support the hypothesis that epigenetically

influenced mutagenesis may be creating genetic variation and be involved in speciation. As such,

our results provide novel support for the role of DNA methylation in interspecies variation,

driving mutagenesis, and genetic evolution between species.

Results

Genetic divergence between benthic-limnetic species pairs

The extent of genetic differentiation between benthic and limnetic species varied among

lakes. In North America, moderate to high divergence between lakes both within and between

species was observed, with greater genome-wide F_{ST} between species in Cliff Lake than Indian

Lake (Table 1A). In Europe, F_{ST} estimates for Langfjordvatn Lake and Zurich Lake showed

greater differentiation between lakes than between species (Table 1B). In all lakes, genetic

differentiation is widespread along the genome (Figure S1). Genetic differentiation between

benthic-limnetic species pairs have been comprehensively covered in earlier studies (Gagnaire et

al. 2013; Rougeux, Gagnaire & Bernatchez 2019; Rougeux, Gagnaire, Praebel, et al. 2019;

Mérot et al. 2023).

Elevated rate of polymorphism in CpG sites

We analyzed from 11 861 765 to 30 919 358 SNPs per lake at approximately 4X

coverage per sample (Table 2) to assess whether SNPs were enriched in CpG sites relative to the

rest of the genome to investigate the prediction that CpG sites are mutagenic. We also tested

whether the enrichment was specific to (i) C/T and G/A SNPs (to account for the G position of

CpG sites), (ii) C/G and G/C SNPs, and (iii) C/A and G/T SNPs. Permutation tests showed that

SNPs were significantly enriched in CpG sites for all four lakes (p < 0.0001; Table 2), with none

of the 10 000 permutations per test reaching or exceeding the observed rate of polymorphism at

CpG sites. Between 10.5 and 12.3% of SNPs occurred in CpG sites for each lake in contrast with

an average of 3.8% of polymorphic sites across the genome. This means that 2.8 to 3.2 times

more SNPs occur in CpG sites than in the rest of the genome. There was significant enrichment

for all SNP types in CpG sites according to Pearson's chi-squared tests ($p < 0.001$; Table 2). C/T

and G/A SNPs were most common in CpG sites, with 3.0 to 3.4 times more C/T SNPs occurring

in CpG sites than in all C and G sites across the genome. The other SNP types were slightly less

enriched, with an enrichment of 1.1 to 1.2 times more C/G and G/C SNPs and 1.2 to 1.3 times

more C/A and G/T SNPs in CpG sites relative to the rest of the C and G sites in the genome

(Table 2).

Epigenetic differentiation between benthic and limnetic whitefish

We assessed the level of epigenetic differentiation between benthic-limnetic species pairs

by performing differential methylation analysis for each lake. After all quality trimming, we

analyzed between 12 449 354 and 18 999 942 CpG sites per lake with approximately 7.5 to 8.1X

coverage per sample (see Table 3 for detailed information). We identified variable but significant

epigenetic differentiation between benthic-limnetic species pairs in all lakes: 38 060

differentially methylated loci (DMLs, i.e., CpG sites) and 2 891 differentially methylated regions

(DMRs, i.e., prolonged regions of the DNA with differences in methylation between species) in

Cliff Lake, 24 949 DMLs and 2 300 DMRs in Indian Lake, 3 537 DMLs and 367 DMRs in

Langfjordvatn Lake, and 7 140 DMLs and 703 DMRs in Zurich Lake (Figure 2, Table 3, Tables

S1-S8). The DMRs covered between 2 296 and 24 524 CpG sites in each lake (Table 3). Overall

epigenetic differentiation between species was greater in North America than in Europe,

consistent with higher interspecies genetic differentiation as measured by F_{ST} in North America

relative to Europe.

Enrichment for overlaps between outlier SNPs and DMLs

For each lake, we assessed whether the most differentiated SNPs (5% highest F_{ST}) in

each lake overlap more often than expected by chance with CpG sites identified as DMLs

between species. Such overlaps in genetic and epigenetic differentiation could represent potential

sites of ongoing genetic divergence where divergent methylation patterns could be undergoing

epigenetically influenced mutagenesis into stable genetic variants between species. We

considered only polymorphic CpG sites in this analysis to account for the elevated mutation rate

at CpG sites. Pearson's chi-squared tests showed that the number of outlier SNP-DML overlaps

was much greater than expected based on the combined probability of finding both an outlier

SNP and a DML in a polymorphic CpG site (Table 3; see methods). We found between an 8.4 to

a 65.9-fold enrichment of overlaps depending on lake. The enrichment was greater in North

America (65.9-fold in Cliff Lake and 29.4-fold in Indian Lake) than in Europe (8.4-fold in

Langfjordvatn Lake and 15.9-fold in Zurich Lake).

Gene ontology enrichment analysis for the outlier SNP-DML overlaps showed

enrichment for 12 terms mostly involved in immune function in Cliff Lake, no terms in Indian

Lake, five molecular function terms in Langfjordvatn Lake, and 10 terms in Zurich Lake

associated with DNA binding and transcription (Table 4).

Gene-level parallelism for epigenetic variation among lakes

We assessed parallelism in the previously identified DMLs, DMRs, and outlier SNP-

DML overlaps between benthic and limnetic species among lakes. We were unable to compare

exact genomic locations for these markers since we used different reference genomes for

European and lake whitefish. Instead, we determined which markers directly overlapped with

gene transcripts in the annotated whitefish genomes and considered parallelism in genes across

lakes (Figure 3). There was some parallelism in the genes associated with DMLs and DMRs (111

and three common genes among all four lakes representing 2.0% and 0.23% of genes,

respectively), especially shared between two or three lakes. There was little parallelism with

respect to outlier SNP-DML overlaps with common genes among all lakes.

Discussion

Recent speciation research has expanded to include epigenetic mechanisms such as DNA

methylation (Pál & Miklós 1999; Richards et al. 2010; Greenspoon et al. 2022; Ashe et al. 2021)

which could contribute to phenotypic diversification and reproductive isolation (Laporte et al.

2019). Here we provide evidence for genetic and epigenetic differentiation between sympatric

benthic-limnetic whitefish species pairs from two continents with limited epigenetic parallelism

(i.e., shared genes) between lakes. We show that polymorphism is enriched at CpG sites,

including high overlap of outlier SNPs and CpG sites showing differential methylation between

species which may represent sites of ongoing mutagenesis. Together, our results provide support

for the proposed contributions of DNA methylation to phenotypic diversification (Anastasiadi et

al. 2021; Vogt 2021), mutagenesis (Tomkova & Schuster-Böckler 2018), and speciation (Pál &

Miklós 1999; Richards et al. 2010; Greenspoon et al. 2022; Ashe et al. 2021).

Considerable but variable genetic divergence between species and lakes

We observed genetic differentiation between species in all four lakes, consistent with

previous studies (Rougeux et al. 2017; Bernatchez et al. 2010; Dion-Côté et al. 2017; Rogers &

Bernatchez 2007; Rougeux, Gagnaire, Praebel, et al. 2019; Østbye, Næsje, et al. 2005; Østbye et

al. 2006; Siwertsson et al. 2013; De-Kayne et al. 2022; Feulner & Seehausen 2019; Mérot et al.

2023). There was greater genetic differentiation between species in Cliff Lake than Indian Lake

in North America, as previously reported in our sister study which reported on the same lake

whitefish data and conclusions (Mérot et al. 2023) and in another study with lower genomic

resolution (Gagnaire et al. 2013). Genetic differentiation was greater between lakes than between

species in Europe (Table 1B), likely due to the Zurich Lake and Langfjordvatn Lake populations

having occupied two different glacial refugia during the last glaciation period (Østbye,

Bernatchez, et al. 2005). However, our within-lake genome-wide F_{ST} estimates are slightly low

compared to other estimates between European whitefish species pairs (0.037 to 0.12 based on

RAD sequencing (Feulner & Seehausen 2019), 0.042 to 0.096 based on 16 microsatellite loci

(Siwertsson et al. 2013) and 0.01 to 0.075 based on six microsatellite loci (Østbye et al. 2006)),

likely due to the use of whole genome vs. targeted or reduced representation methods and the

low MAF filter (MAF > 0.05) including many low frequency SNPs in this analysis. Variable

interspecies F_{ST} between lakes is consistent with varying degrees of reproductive isolation/gene

flow between species in different systems (Lu & Bernatchez 1999; Gagnaire et al. 2013; De-

Kayne et al. 2022; Rougeux, Gagnaire & Bernatchez 2019; Østbye et al. 2006). They are also

consistent with estimated lake whitefish divergence history in Cliff and Indian Lakes (32 000

years in allopatric speciation and 9 200 years since secondary contact for Cliff, 29 000 years in

allopatric speciation and 8 500 years since secondary contact for Indian) (Rougeux et al. 2017).

European whitefish divergence is estimated to have occurred over a longer timeline (121 000

years in allopatry and 12 000 years since secondary contact for Langfjordvatn, 107 000 years in

allopatry and 28 600 years since secondary contact for Zurich) (Rougeux, Gagnaire &

Bernatchez 2019). Despite European whitefish having a longer period of allopatric speciation,

lake whitefish show greater genetic divergence. The variation in interspecies genetic divergence

between lakes could be due to the extent of differences between the species' trophic niches, with

species occupying vastly different trophic niches also showing greater morphological differences

(Lu & Bernatchez 1999). Overall, genetic differentiation between species was widespread along

the genome consistent with previous work (De-Kayne et al. 2022; Feulner & Seehausen 2019;

Mérot et al. 2023), though large-effect loci (De-Kayne et al. 2022) and genomic islands of

divergence (Gagnaire et al. 2013) have also been reported. This genome-wide genetic

differentiation between species may be associated with phenotypic divergence and reproductive

isolation between species (Coyne & Orr 2004).

Epigenetic divergence between whitefish species pairs

We found considerable epigenetic divergence in liver tissue between species, though the

extent of divergence varied between lakes and was greater in North America than in Europe

(Figure 2, Table 3). Our results support the idea that DNA methylation could provide an

additional mechanism for phenotypic diversification and speciation, similar to results in lake

whitefish showing differential methylation of transposable elements between species in liver

tissue, which can affect transposable element activity (Laporte et al. 2019). These methylation

changes may have arisen due to environmental differences between benthic and limnetic habitats

which could contribute to character displacement between species since the environment can

have profound effects on the methylome. Previous studies have shown that methylation is

affected by temperature (Ryu et al. 2018; Venney et al. 2022; Metzger & Schulte 2017; McCaw

et al. 2020; Anastasiadi et al. 2017), salinity (Heckwolf et al. 2020; Artemov et al. 2017), rearing

environment (Leitwein et al. 2021; Le Luyer et al. 2017; Wellband et al. 2021; Gavery et al.

2018), and other factors in various systems. Differences in habitat use can influence DNA

methylation, as observed between capelin (*Mallotus villosus*) utilizing beach- and demersal-

spawning life history tactics (Venney et al. 2023), among sympatric Arctic charr morphs

occupying different habitats and dietary niches (Matlosz et al. 2022), between phenotypically

divergent cichlid species (Vernaz et al. 2022), and in freshwater snails (*Potamopyrgus*

antipodarum) which exhibit differences in shell shape associated with water current speed

(Thorson et al. 2017).

DNA methylation can also be influenced by genetic variation (Richards 2006; Lallias et

al. 2021), therefore epigenetic differences may in part be driven by genetic divergence between

species. Given the greater epigenetic divergence and F_{ST} between species pairs in North America

than in Europe, it is possible that the methylation differences are a function of genetic divergence

between species. Parallel transcriptional differences were previously reported between benthic-

limnetic whitefish species pairs in both captive and natural conditions, indicating that some

transcriptional differences between species are under genetic control and may have been subject

to selection (St-Cyr et al. 2008). Transcriptional divergence between benthic-limnetic species

pairs was also shown in 48 of 64 samples used in this study (6 samples per species per lake) and

differences were often parallel across lakes in both lake whitefish and European whitefish (see

Rougeux, Gagnaire, Praebel, et al. 2019). We generally observed similar ratios of interspecies

DMRs among lakes from our study and differentially expressed gene (DEG) counts in the earlier

transcriptome study by Rougeux et al. (2 891 DMRs and 3 175 DEGs in Cliff Lake, 2 300 DMRs

and 238 DEGs in Indian Lake, 367 DMRs and 276 DEGs in Langfjordvatn Lake, and 703 DMRs

and 1 392 DEGs in Zurich Lake). This indicates that Cliff Lake and Zurich Lake have greater

epigenomic and transcriptomic differences between species relative to the other lakes, though

there are considerably more DMRs than DEGs in Indian Lake, possibly due to greater genetic

and epigenetic differentiation between species. Thus, DNA methylation divergence between

species may be due to differences in habitat and genetic background and could provide a

mechanistic basis for phenotypic diversification between these nascent species pairs due to its

effects on transcription (Bird 2002) and phenotype (Anastasiadi et al. 2021; Vogt 2021).

Mutational enrichment at CpG sites supports epigenetically influenced mutagenesis

Our results suggest that polymorphism is enriched at CpG sites, possibly due to the

mutagenic nature of DNA methylation (Flores et al. 2013; Tomkova & Schuster-Böckler 2018;

Ashe et al. 2021) generating genetic variation between species pairs. We show that CpG sites,

the main sites where DNA methylation occurs in vertebrate genomes, have higher levels of

polymorphism compared to the rest of the genome (Table 2). This suggests that DNA

methylation may be inducing point mutations that could accumulate between species over

generations. We observed a 2.8- to 3.2-fold polymorphism enrichment at CpG sites which is

comparable to the previously reported 3.5-fold increased mutation in methylated relative to

unmethylated cytosines (Gorelick 2003; Jones et al. 1992). It is likely that more mutations have

occurred yet were not detected due to selection against them since epigenetically induced

mutations are generally inferred to be deleterious based on mismatches between experimentally

observed epigenetically induced mutation rate and the observed rates of these mutations in

populations (Danchin et al. 2019). However, some mutations could be retained through random

genetic drift if neutral or mildly deleterious, or selected for if they prove beneficial, leading to

increased frequency of the novel mutation over time. While epigenetically induced C>T

transitions are most common due to spontaneous deamination of cytosines to uracil (Tomkova &

Schuster-Böckler 2018), we show that there is significant enrichment of all types of point

mutations at CpG sites (Table 2). The exact mechanisms behind spontaneous C>A and C>G

mutations without the involvement of mutagens are less clear in the context of epigenetically

influenced mutagenesis (Tomkova & Schuster-Böckler 2018), though we provide evidence that

all types of polymorphism are enriched at CpG sites.

Co-occurrence of genetic and epigenetic divergence at CpG sites

Our results showed that SNPs with high F_{ST} between species are enriched at DMLs.

These sites could potentially reflect ongoing mutagenesis or genetic assimilation contributing to

genetic evolution between benthic-limnetic species pairs, wherein one species is in the process of

losing the CpG site. While the ancestral species is unknown, it is also possible that some of these

sites have mutated from a non-CpG site to a CpG site (e.g., from TpG to CpG), though a

causative mechanism for this is not immediately clear. The outlier SNP-DML overlaps showed

both differential liver methylation between species (i.e., when CpG sites are still present in both

species) and high genetic differentiation between species (i.e., when one species has partially

assumed the "assimilated" state and a cytosine has mutated to another nucleotide). There was a

greater enrichment for outlier SNP-DML overlaps in North America than Europe (see Table 3)

consistent with greater genetic differentiation and more pronounced reproductive isolation

between species pairs in North America, though this could also reflect differences in mutational

signatures between species (e.g., Goldberg and Harris 2022). It is also likely that the degree of

enrichment would differ among tissues and through ontogeny since the methylome is tissue-

specific (Venney et al. 2016; Christensen et al. 2009; Gavery et al. 2018) and affected by age

(Venney et al. 2016; Christensen et al. 2009). Gene ontology analysis of the overlaps showed

that they occurred in genes associated with behaviour, immune function, metabolism, DNA

binding, cellular processes, oxio-reductase activity, and transcription factor activity depending

on the lake (Table 4), though there were no shared GO terms among lakes in our study.

Nevertheless, our findings were consistent with previous transcriptomic studies in whitefish

showing enrichment of similar functions in DEGs (Rougeux, Gagnaire, Praebel, et al. 2019; St-

Cyr et al. 2008), with potential implications for immune function and growth (Rougeux,

Gagnaire, Praebel, et al. 2019). To our knowledge, this one of the first studies to relate epigenetic

divergence and putatively epigenetically induced polymorphism to speciation, providing support

for previous theories on the role of DNA methylation in genetic assimilation. Stajic et al. (2019)

previously showed that mutational assimilation was dependent on the capacity of *S. cerevisiae* to

modify histone acetylation, showing the involvement of epigenetic mechanisms in genetic

assimilation. A recent study in threespine stickleback (*Gasterosteus aculeatus*) showed that

DMLs between freshwater and marine stickleback were also associated with high nucleotide

diversity (Ord et al. 2023). Interestingly, a study in great apes (*Homo*, *Pan*, *Gorilla*, and *Pongo*

sp.) found that mutational signatures across the genome differed between species and that

epigenetically induced mutations were influenced by chromatin state and cytosine

hydroxymethylation (Goldberg & Harris 2022). The role of epigenetic processes in inducing

mutagenesis is becoming clearer, though we provide some initial support for DNA methylation

influencing mutagenesis in the context of ecological speciation.

While the idea of genetic assimilation is exciting, there are other explanations for the

association between outlier SNPs and DMLs. An alternative hypothesis is that both DNA

methylation and genetic polymorphism at the same CpG site could have similar effects on

transcription and thus provide two different, simultaneous molecular mechanisms for controlling

transcription at the same site in different individuals. Genetic variation at proximal linked sites

could also determine methylation state at these genetically and epigenetically differentiated CpG

sites, though it is unclear why this would cause an enrichment of outlier SNP-DML overlaps

given widespread genomic differentiation between species in all lakes. It is also possible that

selection maintains both genetic and epigenetic state at these sites and the two are not related.

Future long-term experimental evolution studies have the potential to distinguish between these

possibilities and facilitate real-time observation of epigenetically induced mutagenesis and

genetic assimilation.

A hypothetical role for DNA methylation in whitefish speciation

When environments change and then remain stable (e.g., in range expansions, habitat

colonization, or shifts to novel habitat use), DNA methylation may serve as an initial plastic

response to the new environment, with the new methylation state being maintained by natural

selection (Ashe et al. 2021). Given the rapid postglacial speciation rate (~3-4K generations)

between benthic and limnetic species in all lakes, methylation differences could have arisen

quickly, contributing to the multi-trait rapid evolution that occurred between species. Over time,

epigenetically influenced mutagenesis could have led to methylation changes inducing genetic

divergence (Danchin et al. 2019; Ashe et al. 2021). Therefore, epigenetically induced

mutagenesis could have contributed to genetic differentiation between benthic and limnetic

whitefish. Divergent selection could then have acted on both epigenetic and genetic marks if they

affected phenotype. This is consistent with heritable differences in behaviour (e.g., Rogers et al.

2002) and minimal plasticity in morphological traits differentiating benthic and limnetic lake

whitefish (Laporte et al. 2016). Transcriptomic differences between benthic-limnetic species

pairs were also stable across environments in lake whitefish (St-Cyr et al. 2008) and related to

parallel genetic differences between species in both lake and European whitefish (Rougeux,

Gagnaire, Praebel, et al. 2019), suggesting that these putatively adaptive traits might be

genetically controlled. However, further study would be needed for any direct test of

environmental differences between species pairs or a link between (epi)genetic variation and

phenotype.

Conclusions

We found substantial genetic and epigenetic divergence between independently derived

benthic-limnetic species pairs in lake and European whitefish. We provide evidence that DNA

methylation may lead to increased polymorphism due to its mutagenic nature, potentially

contributing to early phenotypic diversification, genetic divergence, and speciation. We

characterized potential sites of ongoing genetic assimilation wherein differential methylation

levels between species may be influencing polymorphism and resulting in divergent genetic

variation between benthic-limnetic species pairs. As such, our results shed light on the diverse

ways DNA methylation can contribute to plasticity and evolution, from plastic responses to

environmental changes to the induction of mutagenesis leading to genetic divergence. Future

studies using experimental evolution or evolve and resequence approaches are needed to confirm

the mutagenic nature of DNA methylation and its role in genetic assimilation.

Materials and Methods

Sample preparation and sequencing

We sampled limnetic-benthic whitefish species pairs from two lakes in North America

and two lakes in Europe: lake whitefish (*C. clupeaformis*) from Cliff and Indian Lake, Maine,

USA, and European whitefish (*C. lavaretus*) from Langfjordvatn Lake, Norway and Zurich

Lake, Switzerland (Fig. 1). Animal care was performed humanely under Université Laval animal

care permit 126316. Fish were caught with gillnets, humanely euthanized, and immediately

dissected to obtain fresh tissue samples. Liver tissue was sampled from 64 fish, including six per

lake previously used in Rougeux et al. (2019): eight individuals per species per lake except for

Indian Lake where we sampled seven benthic and nine limnetic fish. Samples were stored either

at -80°C or in RNAlater; all samples from Europe were stored in RNAlater. Liver tissue was

chosen due to its homogeneous tissue characteristics and involvement in growth and metabolism

(Trefts et al. 2017).

Genomic DNA was isolated using a modified salt extraction protocol (Aljanabi &

Martinez 1997). DNA was checked on a 1% agarose gel and quantified on a NanoDrop

spectrophotometer. WGS and WGBS libraries were built at the McGill University and Genome

Quebec Innovation Centre (Montreal, Canada) using in-house protocols. WGS was performed

using paired end 150 bp sequencing on an Illumina HiSeq4000 with estimated 5X coverage. The

North American samples were previously sequenced in Mérot et al. (2022), and WGBS was

performed in this study only using paired end 150 bp sequencing on the Illumina HiSeqX with

samples randomly distributed across 16 lanes (four per lane) with ~10X coverage.

Whole genome sequencing analysis

Reads were trimmed and quality filtered with fastp (Chen et al. 2018). The North

American samples were aligned to the *Coregonus clupeaformis* genome (ASM1839867v1;

(Mérot et al. 2023) and the European samples were aligned to the European whitefish genome

(*Coregonus* sp. Balchen; LR778253.1; De-Kayne et al. 2020) using BWA-MEM (Li 2021).

Aligned reads were filtered to require mapping quality over 10 with Samtools v1.8 (Li et al.

2009). Duplicate reads were removed with MarkDuplicates (PicardTools v1.119.,

http://broadinstitute.github.io/picard). We realigned around indels with GATK IndelRealigner

(McKenna et al. 2010) and soft clipped overlapping read ends using clipOverlap in bamUtil

v1.0.14 (Breese & Liu 2013). The pipeline is available at

https://github.com/enormandeau/wgs_sample_preparation.

Bam alignments were analysed with the program ANGSD v0.931 (Korneliussen et al.

2014) which accounts for genotype uncertainty and is appropriate for low and medium coverage

WGS (Lou et al. 2021). Reads were filtered to remove low-quality reads and to keep mapping

quality above 30 and base quality above 20. We ran ANGSD on each lake separately to detect

polymorphic positions (SNPs), estimate the spectrum of allele frequency, minor allele frequency

(MAF), and genotype likelihoods. Genotype likelihoods were estimated with the GATK method

(-GL 2). The major allele was the most frequent allele (-doMajorMinor 1). We kept positions

covered by at least one read in at least 75% of individuals, with a total coverage below 400 (25

times the number of individuals) to avoid including repeated regions in the analysis. After such

filtering, we exported a list of covered positions for each lake for further analysis, including

1 998 994 058 positions in Cliff Lake, 1 981 247 030 in Indian Lake, 1 658 793 591 in

Langfjordvatn, and 1 702 852 520 in Zurich Lake.

From this list of variant and invariant positions, we extracted a list of SNPs as the

variable positions with an MAF above 5% and subsequently used this list with their respective

major and minor alleles for most analyses (i.e., F_{ST} between benthic and limnetic, overlap with

CpG sites). Differentiation between benthic and limnetic species in each lake was measured with

F_{ST} statistics, using ANGSD to estimate joint allele frequency spectrum, realSFS functions to

compute F_{ST} in sliding windows of 100 KB with a step of 25 KB. Positions were restricted to the

polymorphic SNPs (>5% MAF) previously polarized as major or minor allele (options –sites and

–doMajorMinor 3), and which were covered in at least 75% of the samples in each species. F_{ST}

estimates for Cliff Lake and Indian Lake in North America were previously published in Mérot

et al. (2022). F_{ST} estimates between lakes were performed with the same analyses at the

continent level to ensure a consistent polarisation of the major allele. The ANGSD pipeline is

available at https://github.com/clairemerot/angsd_pipeline.

Frequency of SNPs in CpG sites

We determined whether CpG sites were more polymorphic than the rest of the genome,

which would be indicative of CpG methylation influencing mutagenesis (Jones et al. 1992;

Gorelick 2003). We used fastaRegexFinder.py (https://github.com/dariober/bioinformatics-

cafe/blob/master/fastaRegexFinder/) to find all CpG sites in each reference genome. Next, we

restricted the CpG site lists to include only sites with sufficient coverage in the WGS data using

bedtools *intersect* (Quinlan & Hall 2010) with the list of covered positions for each lake exported

earlier (>75% of samples covered). We used regioneR (Gel et al. 2016) to determine how many

SNPs fell within CpG sites using the *numOverlaps* command. We assumed a uniform

distribution of SNPs across the genome as a null hypothesis for both tests, thus a higher

proportion of SNPs in CpGs relative to average polymorphism in the genome would indicate an

enrichment of point mutations in CpGs.

$$\frac{n(SNPs\ in\ CpGs)}{n(CpG\ nucleotides)\ covered\ by\ WGS\ data}\ v.s.\ \frac{n(SNPs)}{length\ of\ genome\ covered\ by\ WGS\ data}$$

We also used a permutation test to determine if the enrichment of SNPs in CpG sites

would occur by chance over 10 000 iterations. We determined the proportion of the genome

made up of CpG sites by dividing the number of CpG nucleotides by the total length of the

genome with sufficient WGS coverage. We assumed a uniform distribution of SNPs across the

genome as a null hypothesis for both tests, thus a higher proportion of SNPs in CpG sites relative

to the proportion of the genome made up of CpG sites would indicate an enrichment of point

mutations in CpGs.

$$\frac{n(SNPs\ in\ CpGs)}{n(SNPs)} \quad v.s. \quad \frac{n(CpG\ nucleotides)\ covered\ by\ WGS\ data}{length\ of\ genome\ covered\ by\ WGS\ data}$$

We tested whether there was an enrichment of specific SNP substitution types as C/T

polymorphisms are expected to be more common due to deamination of methylated cytosines

(Tomkova & Schuster-Böckler 2018; Gorelick 2003; Jones et al. 1992). For this analysis, we

only considered C and G sites in the genome due to the increased mutagenicity of these

nucleotides (Kiktev et al. 2018). We split the list of SNPs into three separate files for each

possible mutation: (i) all C/T and G/A SNPs (where G/A SNPs would indicate a C/T mutation at

the G position in the reverse complement of the DNA), (ii) all C/A and T/G SNPs, and (iii) all

C/G SNPs. We also generated a list of all C and G sites in the genome with sufficient WGS

coverage. We compared the rate of finding each SNP type in a CpG site to the rate of finding that

SNP type in C and G sites, then used Pearson's chi-squared test to determine if the proportions

were significantly different. All scripts are available at

https://github.com/cvenney/ga_permutation.

$$\frac{n(specific\ SNPs\ in\ CpGs)}{n(CpG\ nucleotides)} \quad v.s. \quad \frac{n(specific\ SNPs\ in\ genome)}{n(C\ and\ G\ sites\ in\ the\ genome\ covered\ by\ WGS\ data)}$$

Whole genome DNA methylation tabulation

Raw methylation data were trimmed using fastp (Chen et al. 2018) to remove sequences

with phred quality less than 25, length less than 100 bp, and to remove the first and last

nucleotides which have high sequencing error rate. After trimming, lake and European whitefish

methylation data were analysed separately. Trimmed sequences were aligned to the same

reference genomes as described in the WGS methods using bwa-meth

(https://github.com/brentp/bwa-meth). Alignments with mapping quality greater than 10 were

outputted to a BAM file using samtools (Li et al. 2009) and duplicate sequences were removed

using Picard tools. Methyldackel's *mbias* function was used to inform trimming of biased

methylation calls at the beginning and end of reads (https://github.com/dpryan79/MethylDackel).

CpG-specific methylation calling was performed on bias-trimmed data using methyldackel's

extract function while removing any detected variant sites where SNPs could affect methylation

calling in that individual (*--maxVariantFrac 0.1 --minOppositeDepth 1*). The pipeline is

available at https://github.com/enormandeau/bwa-meth_pipeline.

517   *Individual-level whitelisting of methylation data using SNP data*

C/T SNPs cannot be discerned from true methylation reads because bisulfite conversion

leaves methylated cytosine unchanged but converts unmethylated cytosine to uracil which is

sequenced as a thymine. Therefore, CpG sites which overlap with C/T or G/A SNPs must be

removed from the analysis. We used an individual-level approach where we filtered CpG sites

separately for each sample based on that sample's genotype (Figure 1), in addition to

methylDackel's built-in *--maxVariantFrac* SNP removal function described above. Therefore,

we filter problematic SNPs out of this dataset using both the SNP and the methylation data.

Relying on our WGS data, we thus created a whitelist of CpG sites for each sample, keeping the

sites homozygous for C and G and masking individuals heterozygous or homozygous for the T

or A allele at that site. To do so, we kept CpG sites that were covered in at least 12 individuals in

a given lake (75% of samples for each lake). Then we filtered the whitelists, requiring sites to be

either (i) non-variant positions, (ii) not a C/T SNP for the C position of the CpG in the lake, (iii)

not a G/A SNP for the G position of the CpG in the lake, or (iv) a C/T SNP or a G/A SNP in

other individuals within the lake, but where this individual has a likelihood greater than 0.7 to be

a C/C and G/G homozygote. This resulted in individual-level SNP masking for each sample

where CpG sites with C/T and G/A SNPs were removed in heterozygous individuals, but not

blindly across all samples (see Figure 1 for a visual representation). C/C homozygotes and

individuals with C/A and C/G SNPs at a given CpG site were retained in the analysis as these

genotypes do not affect methylation calling. BedGraph files were filtered to include only CpG

sites covered in the whitelist (i.e., where both the C and G positions were whitelisted). The

pipeline for SNP masking and subsequent methylation analysis is available at

https://github.com/cvenney/ga_methyl.

Coverage filtration and differential methylation analysis

Whitelisted bedGraph files were filtered to exclude CpG sites with less than five and

more than 100 reads. The files were imported into R (R Core Team 2022) where we kept only

CpG sites with sufficient coverage in at least four limnetic and four benthic samples in the lake.

BedGraph files were reformatted for further analysis with DSS (Park & Wu 2016). We then

assessed the level of epigenetic differentiation between benthic-limnetic species pairs by

performing differential methylation analysis for each lake. Methylation data were smoothed over

500 bp regions using the built-in moving average algorithm in DSS to control for spatial

correlation of methylation levels among proximal CpGs. We ran generalized linear models in

DSS to identify differentially methylated loci (DMLs; i.e., CpG sites with significantly different

methylation levels between limnetic and benthic species) and differentially methylated regions

(DMRs; regions of the genome showing differences in methylation levels between experimental

groups) between species. DMLs were considered significant if the false discovery rate (FDR)

corrected p-value was less than 0.05. DMRs were identified in DSS as regions with many

statistically significant DMLs with p-values less than 0.05.

Overlap between DMLs and outlier SNPs

We assessed whether there was an enrichment of outlier SNP-DML overlaps compared to

the number of overlaps expected by chance given the frequency of polymorphic DMLs and the

frequency of outlier SNPS in CpG sites. Enrichment could indicate epigenetically influenced

mutagenesis and genetic assimilation of methylation changes into stable genetic variants. We

generated a list of highly differentiated SNPs for each lake, retaining only the top 5% of SNPs

with the greatest F_{ST} between species, hereafter called "outlier SNPs". DML test files were

converted to bed format for input into bedtools, then *intersect* was used to determine the number

of overlaps between outlier SNPs and DMLs. We used only CpG sites that were (i) covered by

WGS data, (ii) covered by WGBS data, and (iii) polymorphic to account for the observed

elevated mutation rate in CpG sites compared to the rest of the genome. We then used Pearson's

chi-squared test to determine if there was an enrichment of observed DML-outlier SNP overlaps

in polymorphic CpG sites compared to the expected rate of finding overlaps.

$$\frac{n(DMLs\ in\ SNPs)}{n(polymorphic\ CpG\ sites)}\ x\ \frac{n(outlier\ SNPs\ in\ covered\ CpG\ sites)}{n(polymorphic\ CpG\ sites)}\ vs.\ \frac{n(overlaps\ between\ DMLs\ and\ outlier\ SNPs)}{n(polymorphic\ CpG\ sites)}$$

We also performed gene ontology (GO) enrichment analysis for the outlier SNP-DML

overlaps for each lake. The whitefish genomes were first annotated with the GAWN pipeline

(https://github.com/enormandeau/gawn) using the *Salvelinus namaycush* (GCF_018398675.1)

transcriptome from GenBank, which is more complete than the available whitefish

transcriptomes. We subsetted the annotation to include only the genes covered by both SNP and

methylation data which represents the list of genes that could possibly contribute to enriched

terms based on our data. We then retrieved the genes that overlapped the SNP-DML overlap

positions for each lake and proceeded to GO enrichment tests using the pipeline at

https://github.com/enormandeau/go_enrichment). Results were filtered to remove broad terms

(depths 0 and 1) and to require a Benjamini-Hochberg FDR corrected p-value of 0.1 or less.

Parallelism across lakes

We tested for parallelism in (epi)genetic variation among lakes for DMLs, DMRs, and

outlier SNP-DML overlaps between benthic and limnetic species. Genomic positions of genes

were identified using the annotated whitefish genomes from GO enrichment. Bedtools *intersect*

was used to find direct overlaps between the gene positions and the markers of interest for each

lake. Overlaps for each type of marker were visualized using UpSetR (Conway et al. 2017).

Acknowledgements

We dedicate this article to the memory of Louis Bernatchez, who passed away in

September 2023 during the review process. Louis had a tireless passion for science and

whitefish, made outstanding contributions to the field, and was a wonderful mentor and friend.

We thank Kim Præbel, Shripathi Bhat and the Freshwater ecology group at UiT (Tromso) for

providing samples from Norway, and Ole Seehausen for providing samples from Switzerland.

This work was supported by a Discovery research grant from the Natural Sciences and

Engineering Research Council of Canada (NSERC) to L.B. CJV is supported by an NSERC

postdoctoral fellowship.

Data availability statement

European whitefish and lake whitefish whole genome sequencing data are available

through SRA accessions PRJNA906116 and PRJNA820751, respectively. Whole genome

bisulfite sequencing data are available through SRA accession PRJNA559821. All scripts are

available on GitHub as indicated in the methods.

References

Aljanabi SM, Martinez I. 1997. Universal and rapid salt-extraction of high quality genomic DNA

for PCR-based techniques. Nucleic Acids Res. 25:4692–4693.

Anastasiadi D, Díaz N, Piferrer F. 2017. Small ocean temperature increases elicit stage-

dependent changes in DNA methylation and gene expression in a fish, the European sea bass.

Sci. Rep. 7:12401. doi: 10.1038/s41598-017-10861-6.

Anastasiadi D, Venney CJ, Bernatchez L, Wellenreuther M. 2021. Epigenetic inheritance and

reproductive mode in plants and animals. Trends Ecol. Evol. 36:1124–1140. doi:

10.1016/j.tree.2021.08.006.

Artemov A V. et al. 2017. Genome-wide DNA methylation profiling reveals epigenetic

adaptation of stickleback to marine and freshwater conditions. Mol. Biol. Evol. 34:2203–2213.

doi: 10.1093/molbev/msx156.

Ashe A, Colot V, Oldroyd BP. 2021. How does epigenetics influence the course of evolution?

Philos. Trans. R. Soc. B Biol. Sci. 376:20200111. doi: 10.1098/rstb.2020.0111.

Beemelmanns A et al. 2021. DNA methylation dynamics in Atlantic salmon (*Salmo salar*)

challenged with high temperature and moderate hypoxia. Front. Mar. Sci. 7:3. doi:

10.3389/fmars.2020.604878.

Berbel-Filho WM et al. 2020. Environmental enrichment induces intergenerational behavioural

and epigenetic effects on fish. Mol. Ecol. 29:2288–2299. doi: 10.1111/mec.15481.

Bernatchez L et al. 2010. On the origin of species: insights from the ecological genomics of lake

whitefish. Philos. Trans. R. Soc. B Biol. Sci. 365:1783–1800. doi: 10.1098/rstb.2009.0274.

Bernatchez L, Chouinard A, Lu G. 1999. Integrating molecular genetics and ecology in studies

of adaptive radiation: whitefish, *Coregonus* sp., as a case study. Biol. J. Linn. Soc. 68:173–194.

Bernatchez L, Dodson JJ. 1994. Phylogenetic relationships among Palearctic and Nearctic

whitefish (*Coregonus* sp.) populations as revealed by mitochondrial DNA variation. Can. J. Fish.

Aquat. Sci. 51:240–251. doi: 10.1139/f94-310.

Bernatchez L, Dodson JJ. 1991. Phylogeographic structure in mitochondrial DNA of the lake

whitefish (*Coregonus clupeaformis*) and its relation to Pleistocene glaciations. Evolution (N. Y).

45:1016–1035. doi: 10.1111/j.1558-5646.1991.tb04367.x.

Bird A. 2002. DNA methylation patterns and epigenetic memory. Genes Dev. 16:6–21. doi:

10.1101/gad.947102.6.

Breese MR, Liu Y. 2013. NGSUtils: a software suite for analyzing and manipulating next-

generation sequencing datasets. Bioinformatics. 29:494–496. doi:

10.1093/bioinformatics/bts731.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 34:i884–i890. doi: 10.1093/bioinformatics/bty560.

Christensen BC et al. 2009. Aging and environmental exposures alter tissue-specific DNA

methylation dependent upon CpG island context. PLoS Genet. 5:e1000602. doi:

10.1371/journal.pgen.1000602.

Conway JR, Lex A, Gehlenborg N. 2017. UpSetR: An R package for the visualization of

intersecting sets and their properties. Bioinformatics. 33:2938–2940. doi:

10.1093/bioinformatics/btx364.

Coyne JA, Orr HA. 2004. *Speciation*. Sinauer: Sunderland.

Dalziel AC, Martin N, Laporte M, Guderley H, Bernatchez L. 2015. Adaptation and acclimation

of aerobic exercise physiology in lake whitefish ecotypes (*Coregonus clupeaformis*). Evolution

(N. Y). 69:2167–2186. doi: 10.1111/evo.12727.

Danchin E, Pocheville A, Rey O, Pujol B, Blanchet S. 2019. Epigenetically facilitated mutational

assimilation: epigenetics as a hub within the inclusive evolutionary synthesis. Biol. Rev. 94:259–

282. doi: 10.1111/brv.12453.

De-Kayne R et al. 2022. Genomic architecture of adaptive radiation and hybridization in Alpine

whitefish. Nat. Commun. 13:1–13. doi: 10.1038/s41467-022-32181-8.

De-Kayne R, Zoller S, Feulner PGD. 2020. A de novo chromosome-level genome assembly of

Coregonus sp. "Balchen": one representative of the Swiss Alpine whitefish radiation. Mol. Ecol.

Resour. 20:1093–1109. doi: 10.1111/1755-0998.13187.

Derome N, Duchesne P, Bernatchez L. 2006. Parallelism in gene transcription among sympatric

lake whitefish (*Coregonus clupeaformis* Mitchill) ecotypes. Mol. Ecol. 15:1239–1249. doi:

10.1111/j.1365-294X.2005.02968.x.

Dion-Côté AM et al. 2017. Standing chromosomal variation in lake whitefish species pairs: the

role of historical contingency and relevance for speciation. Mol. Ecol. 26:178–192. doi:

10.1111/mec.13816.

Elmer KR et al. 2010. Rapid evolution and selection inferred from the transcriptomes of

sympatric crater lake cichlid fishes. Mol. Ecol. 19:197–211. doi: 10.1111/j.1365-

294X.2009.04488.x.

Feulner PGD, Seehausen O. 2019. *Genomic insights into the vulnerability of sympatric whitefish*

species flocks. doi: 10.1111/mec.14977.

Flores KB, Wolschin F, Amdam G V. 2013. The role of methylation of DNA in environmental

adaptation. Integr. Comp. Biol. 53:359–372. doi: 10.1093/icb/ict019.

Gagnaire PA, Pavey SA, Normandeau E, Bernatchez L. 2013. The genetic architecture of

reproductive isolation during speciation-with-gene-flow in lake whitefish species pairs assessed

by RAD sequencing. Evolution (N. Y). 67:2483–2497. doi: 10.1111/evo.12075.

Gavery MR, Nichols KM, Goetz GW, Middleton MA, Swanson P. 2018. Characterization of

genetic and epigenetic variation in sperm and red blood cells from adult hatchery and natural-

origin steelhead, Oncorhynchus mykiss. G3 Genes, Genomes, Genet. 8:3723–3736. doi:

10.1534/g3.118.200458.

Gel B et al. 2016. RegioneR: an R/Bioconductor package for the association analysis of genomic

regions based on permutation tests. Bioinformatics. 32:289–291. doi:

10.1093/bioinformatics/btv562.

Goldberg ME, Harris K. 2022. Mutational signatures of replication timing and epigenetic

modification persist through the global divergence of mutation spectra across the great ape

phylogeny. Genome Biol. Evol. 14:1–18. doi: 10.1093/gbe/evab104.

Gorelick R. 2003. Evolution of dioecy and sex chromosomes via methylation driving Muller's

ratchet. Biol. J. Linn. Soc. 80:353–368. doi: 10.1046/j.1095-8312.2003.00244.x.

Greenspoon PB, Spencer HG, M'Gonigle LK. 2022. Epigenetic induction may speed up or slow
down speciation with gene flow. Evolution (N. Y). 76:1170–1182. doi: 10.1111/evo.14494.

Gudbrandsson J et al. 2018. Differential gene expression during early development in recently
evolved and sympatric Arctic charr morphs. PeerJ. 2018:1–35. doi: 10.7717/peerj.4345.

Heckwolf MJ et al. 2020. Two different epigenetic information channels in wild three-spined
sticklebacks are involved in salinity adaptation. Sci. Adv. 6. doi: 10.1126/sciadv.aaz1138.

Hu J et al. 2021. Heritability of DNA methylation in threespine stickleback (*Gasterosteus*
aculeatus). Genetics. 217. doi: 10.1093/GENETICS/IYAB001.

Jacobs A, Elmer KR. 2021. Alternative splicing and gene expression play contrasting roles in the
parallel phenotypic evolution of a salmonid fish. Mol. Ecol. 30:4955–4969. doi:
10.1111/mec.15817.

Jacobsen MW et al. 2012. Mitogenome sequencing reveals shallow evolutionary histories and
recent divergence time between morphologically and ecologically distinct European whitefish
(*Coregonus* spp.). Mol. Ecol. 21:2727–2742. doi: 10.1111/j.1365-294X.2012.05561.x.

Jeukens J, Bittner D, Knudsen R, Bernatchez L. 2009. Candidate genes and adaptive radiation:
insights from transcriptional adaptation to the limnetic niche among coregonine fishes
(*Coregonus* spp., *Salmonidae*). Mol. Biol. Evol. 26:155–166. doi: 10.1093/molbev/msn235.

Jones PA, Rideout WM, Shen J-C, Spruck CH, Tsai YC. 1992. Methylation, mutation and
cancer. BioEssays. 14:33–36.

Kiktev DA, Sheng Z, Lobachev KS, Petes TD. 2018. GC content elevates mutation and
recombination rates in the yeast *Saccharomyces cerevisiae*. Proc. Natl. Acad. Sci. U. S. A.
115:E7109–E7118. doi: 10.1073/pnas.1807334115.

Korneliussen TS, Albrechtsen A, Nielsen R. 2014. ANGSD: analysis of next generation
sequencing data. BMC Bioinformatics. 15:1–13. doi: 10.1186/s12859-014-0356-4.

Lallias D et al. 2021. Sources of variation of DNA methylation in rainbow trout: combined
effects of temperature and genetic background. Epigenetics. 16:1031–1052. doi:
10.1080/15592294.2020.1834924.

Laporte M et al. 2019. DNA methylation reprogramming, TE derepression, and postzygotic
isolation of nascent animal species. Sci. Adv. 5:1–9. doi: 10.1126/sciadv.aaw1644.

Laporte M et al. 2015. RAD-QTL mapping reveals both genome-level parallelism and different
genetic architecture underlying the evolution of body shape in lake whitefish (*Coregonus*
clupeaformis) species pairs. G3 Genes, Genomes, Genet. 5:1481–1491. doi:
10.1534/g3.115.019067.

Laporte M, Dalziel AC, Martin N, Bernatchez L. 2016. Adaptation and acclimation of traits
associated with swimming capacity in lake whitefish (*Coregonus clupeaformis*) ecotypes. BMC
Evol. Biol. 16:1–13. doi: 10.1186/s12862-016-0732-y.

Leitwein M et al. 2021. Epigenomic modifications induced by hatchery rearing persist in germ
line cells of adult salmon after their oceanic migration. Evol. Appl. 14:2402–2413. doi:
10.1111/eva.13235.

Li H. 2021. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
Mol. Ecol. 30:5966–5993.

Li H et al. 2009. The sequence alignment/map format and SAMtools. Bioinformatics. 25:2078–
2079. doi: 10.1093/bioinformatics/btp352.

Li L, Gao Y, Wu Q, Cheng ASL, Yip KY. 2019. New guidelines for DNA methylome studies
regarding 5-hydroxymethylcytosine for understanding transcriptional regulation. Genome Res.
29:543–553. doi: 10.1101/gr.240036.118.

Lou RN, Jacobs A, Wilder A, Therkildsen NO. 2021. A beginner's guide to low-coverage whole
genome sequencing for population genomics. Mol. Ecol. 30:5966–5993.
https://www.authorea.com/users/380682/articles/496574-a-beginner-s-guide-to-low-coverage-
whole-genome-sequencing-for-population-genomics.

Lu G, Bernatchez L. 1999. Correlated trophic specialization and genetic divergence in sympatric
lake whitefish ecotypes (*Coregonus clupeaformis*): support for the ecological speciation
hypothesis. Evolution (N. Y). 53:1491–1505. doi: 10.1111/j.1558-5646.1999.tb05413.x.

Le Luyer J et al. 2017. Parallel epigenetic modifications induced by hatchery rearing in a Pacific
salmon. Proc. Natl. Acad. Sci. 114:12964–12969. doi: 10.1101/148577.

Manousaki T et al. 2013. Parsing parallel evolution: ecological divergence and differential gene

expression in the adaptive radiations of thick-lipped Midas cichlid fishes from Nicaragua. Mol.

Ecol. 22:650–669. doi: 10.1111/mec.12034.

Marques DA, Meier JI, Seehausen O. 2019. A combinatorial view on speciation and adaptive

radiation. Trends Ecol. Evol. 34:531–544. doi: 10.1016/j.tree.2019.02.008.

Matlosz S et al. 2022. DNA methylation differences during development distinguish sympatric

morphs of Arctic charr (*Salvelinus alpinus*). Mol. Ecol. 31:4739–4761. doi: 10.1111/mec.16620.

McCaw BA, Stevenson TJ, Lancaster LT. 2020. Epigenetic responses to temperature and

climate. Integr. Comp. Biol. 60:1469–1480. doi: 10.1093/icb/icaa049.

McKenna A et al. 2010. The genome analysis toolkit: a MapReduce framework for analyzing

next-generation DNA sequencing data. Genome Res. 20:1297–1303. doi:

10.1101/gr.107524.110.20.

Mckenzie JL et al. 2021. Incomplete reproductive isolation and strong transcriptomic response to

hybridization between sympatric sister species of salmon. Proc. R. Soc. B Biol. Sci.

288:20203020.

Mérot C et al. 2023. Genome assembly, structural variants, and genetic differentiation between

lake whitefish young species pairs (Coregonus sp.) with long and short reads. Mol. Ecol.

32:1458–1477. doi: 10.1111/mec.16468.

Metzger DCH, Schulte PM. 2017. Persistent and plastic effects of temperature on DNA

methylation across the genome of threespine stickleback (*Gasterosteus aculeatus*). Proc. R. Soc.

B Biol. Sci. 284:20171667. doi: 10.1098/rspb.2017.1667.

Nishikawa K, Kinjo AR. 2018. Mechanism of evolution by genetic assimilation. Biophys. Rev.

10:667–676. doi: 10.1007/s12551-018-0403-x.

Ord J, Gossmann TI, Adrian-Kalchhauser I. 2023. High nucleotide diversity accompanies

differential DNA methylation in naturally diverging populations. Mol. Biol. Evol. 40:1–16. doi:

10.1093/molbev/msad068.

Østbye K et al. 2006. Parallel evolution of ecomorphological traits in the European whitefish

Coregonus lavaretus (L.) species complex during postglacial times. Mol. Ecol. 15:3983–4001.

doi: 10.1111/j.1365-294X.2006.03062.x.

Østbye K, Bernatchez L, Næsje TF, Himberg KJM, Hindar K. 2005. Evolutionary history of the

European whitefish *Coregonus lavaretus* (L.) species complex as inferred from mtDNA

phylogeography and gill-raker numbers. Mol. Ecol. 14:4371–4387. doi: 10.1111/j.1365-

294X.2005.02737.x.

Østbye K, Næsje TF, Bernatchez L, Sandlund OT, Hindar K. 2005. Morphological divergence

and origin of sympatric populations of European whitefish (*Coregonus lavaretus* L.) in Lake

Femund, Norway. J. Evol. Biol. 18:683–702. doi: 10.1111/j.1420-9101.2004.00844.x.

Pál C, Miklós I. 1999. Epigenetic inheritance, genetic assimilation and speciation. J. Theor. Biol.

200:19–37. doi: 10.1006/jtbi.1999.0974.

Park Y, Wu H. 2016. Differential methylation analysis for BS-seq data under general

experimental design. Bioinformatics. 32:1446–1453. doi: 10.1093/bioinformatics/btw026.

Pavey SA, Collin H, Nosil P, Rogers SM. 2010. The role of gene expression in ecological

speciation. Ann. N. Y. Acad. Sci. 1206:110–129. doi: 10.1111/j.1749-6632.2010.05765.x.

Pfennig DW et al. 2010. Phenotypic plasticity's impacts on diversification and speciation.

Trends Ecol. Evol. 25:459–467. doi: 10.1016/j.tree.2010.05.006.

Pigeon D, Chouinard A, Bernatchez L. 1997. Multiple modes of speciation involved in the

parallel evolution of sympatric morphotypes of lake whitefish (*Coregonus clupeaformis*,

Salmonidae). Evolution (N. Y). 51:196–205. doi: 10.1111/j.1558-5646.1997.tb02401.x.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics. 26:841–842. doi: 10.1093/bioinformatics/btq033.

R Core Team. 2022. R: A language and environment for statistical computing. https://www.r-

project.org.

Richards CL, Bossdorf O, Pigliucci M. 2010. What role does heritable epigenetic variation play

in phenotypic evolution? Bioscience. 60:232–237. doi: 10.1525/bio.2010.60.3.9.

Richards EJ. 2006. Inherited epigenetic variation - revisiting soft inheritance. Nat. Rev. Genet.

7:395–402.

Rogers SM, Bernatchez L. 2007. The genetic architecture of ecological speciation and the

association with signatures of selection in natural lake whitefish (*Coregonus* sp. Salmonidae)

species pairs. Mol. Biol. Evol. 24:1423–1438. doi: 10.1093/molbev/msm066.

Rogers SM, Bernatchez L. 2006. The genetic basis of intrinsic and extrinsic post-zygotic

reproductive isolation jointly promoting speciation in the lake whitefish species complex

(*Coregonus clupeaformis*). J. Evol. Biol. 19:1979–1994. doi: 10.1111/j.1420-9101.2006.01150.x.

Rogers SM, Gagnon V, Bernatchez L. 2002. Genetically based phenotype-environment

association for swimming behavior in lake whitefish ecotypes (*Coregonus clupeaformis*

Mitchill). Evolution (N. Y). 56:2322–2329.

Rougeux C, Bernatchez L, Gagnaire PA. 2017. Modeling the multiple facets of speciation-with-

gene-flow toward inferring the divergence history of lake whitefish species pairs (*Coregonus*

clupeaformis). Genome Biol. Evol. 9:2057–2074. doi: 10.1093/gbe/evx150.

Rougeux C, Gagnaire PA, Bernatchez L. 2019. Model-based demographic inference of

introgression history in European whitefish species pairs'. J. Evol. Biol. 32:806–817. doi:

10.1111/jeb.13482.

Rougeux C, Gagnaire PA, Praebel K, Seehausen O, Bernatchez L. 2019. Polygenic selection

drives the evolution of convergent transcriptomic landscapes across continents within a Nearctic

sister species complex. Mol. Ecol. 28:4388–4403. doi: 10.1111/mec.15226.

Ryu T et al. 2020. An epigenetic signature for within-generational plasticity of a reef fish to

ocean warming. Front. Mar. Sci. 7:1–15. doi: 10.3389/fmars.2020.00284.

Ryu T, Veilleux HD, Donelson JM, Munday PL, Ravasi T. 2018. The epigenetic landscape of

transgenerational acclimation to ocean warming. Nat. Clim. Chang. 8:504–509. doi:

10.1038/s41558-018-0159-0.

Seehausen O et al. 2014. Genomics and the origin of species. Nat. Rev. Genet. 15:176–192. doi:

10.1038/nrg3644.

Siwertsson A et al. 2013. Discrete foraging niches promote ecological, phenotypic, and genetic

divergence in sympatric whitefish (*Coregonus lavaretus*). Evol. Ecol. 27:547–564. doi:

10.1007/s10682-012-9607-x.

Smith TA, Martin MD, Nguyen M, Mendelson TC. 2016. Epigenetic divergence as a potential

first step in darter speciation. Mol. Ecol. 25:1883–1894. doi: 10.1111/mec.13561.

St-Cyr J, Derome N, Bernatchez L. 2008. The transcriptomics of life-history trade-offs in

whitefish species pairs (*Coregonus* sp.). Mol. Ecol. 17:1850–1870. doi: 10.1111/j.1365-

294X.2008.03696.x.

Stajic D, Jansen LET. 2021. Empirical evidence for epigenetic inheritance driving evolutionary

adaptation. Philos. Trans. R. Soc. B Biol. Sci. 376:20200121. doi: 10.1098/rstb.2020.0121.

Stajic D, Perfeito L, Jansen LET. 2019. Epigenetic gene silencing alters the mechanisms and rate

of evolutionary adaptation. Nat. Ecol. Evol. 3:491–498. doi: 10.1038/s41559-018-0781-2.

Thorson JLM et al. 2017. Epigenetics and adaptive phenotypic variation between habitats in an

asexual snail. Sci. Rep. 7:1–11. doi: 10.1038/s41598-017-14673-6.

Tomkova M, Schuster-Böckler B. 2018. DNA modifications: naturally more error prone? Trends

Genet. 34:627–638. doi: 10.1016/j.tig.2018.04.005.

Trefts E, Gannon M, Wasserman DH. 2017. The liver. Curr. Biol. 27:R1147–R1151. doi:

10.1016/j.cub.2017.09.019.

Trudel M, Tremblay A, Schetagne R, Rasmussen JB. 2001. Why are dwarf fish so small? An

energetic analysis of polymorphism in lake whitefish (*Coregonus clupeaformis*). Can. J. Fish.

Aquat. Sci. 58:394–405. doi: 10.1139/cjfas-58-2-394.

Venney CJ et al. 2023. Genome-wide DNA methylation predicts environmentally driven life

history variation in a marine fish. Evolution (N. Y). 77:186–198. doi:

https://doi.org/10.1093/evolut/qpac028.

Venney CJ et al. 2022. Thermal regime during parental sexual maturation, but not during

offspring rearing, modulates DNA methylation in brook charr (*Salvelinus fontinalis*). Proc. R.

Soc. B Biol. Sci. 289:20220670. doi: 10.1098/rspb.2022.0670.

Venney CJ, Johansson ML, Heath DD. 2016. Inbreeding effects on gene-specific DNA

methylation among tissues of Chinook salmon. Mol. Ecol. 25:4521–4533. doi:

10.1111/mec.13777.

Venney CJ, Wellband KW, Heath DD. 2020. Rearing environment affects the genetic

architecture and plasticity of DNA methylation in Chinook salmon. Heredity (Edinb). 38–49.

doi: 10.1038/s41437-020-0346-4.

Vernaz G et al. 2022. Epigenetic divergence during early stages of speciation in an African crater

lake cichlid fish. Nat. Ecol. Evol. 1–12. doi: 10.1038/s41559-022-01894-w.

Vogt G. 2021. Epigenetic variation in animal populations: sources, extent, phenotypic

implications, and ecological and evolutionary relevance. J. Biosci. 46:1–24. doi:

10.1007/s12038-021-00138-6.

Vogt G. 2017. Evolution of epigenetic mechanisms in animals and their role in speciation. In:

Handbook of Epigenetics: The New Molecular and Medical Genetics. Elsevier Inc. pp. 409–426.

doi: 10.1016/B978-0-12-805388-1.00027-4.

Wellband K, Roth D, Linnansaari T, Curry RA, Bernatchez L. 2021. Environment-driven

reprogramming of gamete DNA methylation occurs during maturation and is transmitted

intergenerationally in Atlantic salmon. G3 Genes, Genomes, Genet. 11:jkab353. doi:

10.1093/g3journal/jkab353.

Whitehead A, Crawford DL. 2006. Variation within and among species in gene expression: raw

material for evolution. Mol. Ecol. 15:1197–1211. doi: 10.1111/j.1365-294X.2006.02868.x.

Xia J, Han L, Zhao Z. 2012. Investigating the relationship of DNA methylation with mutation

rate and allele frequency in the human genome. BMC Genomics. 13 Suppl 8:S7. doi:

10.1186/1471-2164-13-s8-s7.

Tables

Table 1: Pairwise F_{ST} matrices for (A) *C. clupeaformis* and (B) *C. lavaretus* reveal variable but
considerable genetic divergence between lakes and species.

A)

Cliff benthic	0.175		
Indian limnetic	0.084	0.162	
Indian benthic	0.146	0.182	0.098
	Cliff limnetic	**Cliff benthic**	**Indian limnetic**

B)

Langfjordvatn benthic	0.034		
Zurich limnetic	0.240	0.242	
Zurich benthic	0.237	0.239	0.043
	Langfjordvatn limnetic	**Langfjordvatn benthic**	**Zurich limnetic**

871 **Table 2:** Pearson's chi-squared test results for the rate of polymorphism in CpG sites (i) for all

872 SNP types relative to the rate of polymorphism across the entire genome, and (ii) for specific

873 SNP types relative to all C and G positions in the genome. Fold change indicates the ratio

874 between observed and expected polymorphism rates.

SNP Type	Value	Cliff	Indian	Langfjordvatn	Zurich
All	SNPs	11 861 765	12 727 697	30 919 358	24 221 888
	SNPs in CpGs	1 250 313	1 349 717	3 795 855	2 879 669
	Polymorphic CpGs (%)	10.5	10.6	12.3	11.9
	Polymorphic sites (%)	3.8	3.8	3.8	3.8
	Fold change	2.8	2.8	3.2	3.1
	p-value (Chi-squared test)	*< 0.001*	*< 0.001*	*< 0.001*	*< 0.001*
	p-value (permutation test)	*< 0.001*	*< 0.001*	*< 0.001*	*< 0.001*
C/T and G/A	SNPs	2 804 547	3 080 961	8 321 890	6 251 941
	SNPs in CpGs	750 311	819 314	2 487 283	1 834 278
	SNPs in CpGs (%)	0.98	1.09	3.94	2.81
	SNPs in all Cs and Gs (%)	0.32	0.36	1.17	0.85
	Fold change	3.0	3.0	3.4	3.3
	p-value (Chi-squared test)	*< 0.001*	*< 0.001*	*< 0.001*	*< 0.001*
C/G and G/C	SNPs	956 737	1 012 702	2 253 168	1 810 523
	SNPs in CpGs	95 721	100 779	239 633	198 157
	SNPs in CpGs (%)	0.13	0.13	0.38	0.30
	SNPs in all Cs and Gs (%)	0.11	0.12	0.32	0.25
	Fold change	1.1	1.1	1.2	1.2
	p-value (Chi-squared test)	*< 0.001*	*< 0.001*	*< 0.001*	*< 0.001*
C/A and G/T	SNPs	2 124 638	2 314 058	5 422 615	4 126 260
	SNPs in CpGs	234 886	248 499	590 917	476 256
	SNPs in CpGs (%)	0.31	0.33	0.94	0.73
	SNPs in all Cs and Gs (%)	0.25	0.27	0.76	0.56
	Fold change	1.3	1.2	1.2	1.3
	p-value (Chi-squared test)	*< 0.001*	*< 0.001*	*< 0.001*	*< 0.001*

876 **Table 3:** Statistics from differential methylation analysis, including coverage depth and number

877 of CpGs analysed after all coverage trimming, number of DMLs and DMRs identified using a

878 significance level of $p < 0.05$, and information on outlier SNP-DML overlaps and Pearson chi-

879 squared test results.

Statistic	Cliff	Indian	Langfjordvatn	Zurich
Average coverage depth (WGBS)	7.5	7.6	8.1	8.1
Total CpGs analyzed	16 668 063	18 999 942	12 449 354	17 896 958
DMLs	38 060	24 949	3 537	7 140
% CpGs that are DMLs	0.2	0.1	0.03	0.04
DMRs	2891	2300	367	703
CpGs in DMRs	24524	19566	2296	4547
%CpGs in DMRs	0.15	0.10	0.02	0.03
Outlier SNPs (5% highest F_{ST})	593088	636384	1545963	1211091
Outlier SNP-DML overlaps	180	178	45	132
Expected frequency of overlaps	1.38e-05	2.62e-05	9.12e-06	1.30e-05
Observed frequency of overlaps	9.08e-04	7.71e-04	7.65e-05	2.07e-04
Fold enrichment	65.9	29.4	8.4	15.9
p-value	*<0.001*	*<0.001*	*<0.001*	*<0.001*

881 **Table 4:** Gene ontology results for outlier SNP-DML overlaps in lake and European whitefish.
882 There were 12 enriched terms from Cliff Lake, none from Indian Lake, five from Langfjordvatn
883 Lake, and 10 from Zurich Lake. Significant results had a Benjamini-Hochberg false discovery
884 rate corrected p-value less than 0.1 and were filtered for GO term depth greater than 1 to remove
885 broad terms.

GO term	Subontology	GO term name	GO term depth	p-value (BH-FDR)
		Cliff Lake		
GO:0050912	BP	detection of chemical stimulus involved in sensory perception of taste	5	0.047
GO:0002414	BP	immunoglobulin transcytosis in epithelial cells	6	0.044
GO:0001580	BP	detection of chemical stimulus involved in sensory perception of bitter taste	6	0.047
GO:0002415	BP	immunoglobulin transcytosis in epithelial cells mediated by polymeric immunoglobulin receptor	7	0.044
GO:0071745	CC	IgA immunoglobulin complex	3	0.003
GO:0042571	CC	immunoglobulin complex, circulating	3	0.009
GO:0071746	CC	IgA immunoglobulin complex, circulating	4	0.003
GO:0071749	CC	polymeric IgA immunoglobulin complex	5	0.003
GO:0071751	CC	secretory IgA immunoglobulin complex	6	0.003
GO:0019763	MF	immunoglobulin receptor activity	4	0.042
GO:0005154	MF	epidermal growth factor receptor binding	5	0.013
GO:0001792	MF	polymeric immunoglobulin receptor activity	5	0.013
		Langfjordvatn Lake		
GO:0016840	MF	carbon-nitrogen lyase activity	3	0.052
GO:0016842	MF	amidine-lyase activity	4	0.014
GO:0016715	MF	oxidoreductase activity, acting on paired donors, with incorporation or reduction of molecular oxygen, reduced ascorbate as one donor, and incorporation of one atom of oxygen	4	0.015
GO:0004504	MF	peptidylglycine monooxygenase activity	5	0.001
GO:0004598	MF	peptidylamidoglycolate lyase activity	5	0.001
		Zurich Lake		
GO:0003700	MF	DNA-binding transcription factor activity	2	0.027
GO:0000981	MF	DNA-binding transcription factor activity, RNA polymerase II-specific	3	0.030
GO:0001067	MF	transcription regulatory region nucleic acid binding	4	0.051
GO:0043565	MF	sequence-specific DNA binding	5	0.051
GO:0003690	MF	double-stranded DNA binding	5	0.090
GO:1990837	MF	sequence-specific double-stranded DNA binding	6	0.061

GO:0000976	MF	transcription cis-regulatory region binding	7	0.051
GO:0000987	MF	cis-regulatory region sequence-specific DNA binding	8	0.010
GO:0000977	MF	RNA polymerase II transcription regulatory region sequence-specific DNA binding	8	0.030
GO:0000978	MF	RNA polymerase II cis-regulatory region sequence-specific DNA binding	9	0.010

Figures

Fig. 1: Overview of the sampling design and analysis for sympatric benthic-limnetic species

pairs sampled from four lakes. Lake whitefish (*C. clupeaformis*) were sampled from two lakes in

North America (Cliff Lake and Indian Lake in Maine, USA) and European whitefish (*C.*

lavaretus) were sampled from two lakes in Europe (Langfjordvatn Lake, Norway and Zurich

Lake, Switzerland). The analysis pipelines for whole genome sequencing and whole genome

bisulfite sequencing are outlined in the flowchart. C/T and G/A SNPs that affect methylation

calling are filtered (1) during methylation calling based on settings in methylDackel, and (2)

through our whitelisting approach which uses the SNP data for an additional layer of stringent

filtration. The *C. clupeaformis* samples were previously analysed in Mérot et al. (2022).

Fig. 2: Differential methylation analysis comparing limnetic-benthic species pairs identified (A) 2 891 DMRs in Cliff Lake, (B) 2 300 DMRs in Indian Lake, (C) 367 DMRs in Langfjordvatn Lake, and (D) 703 DMRs in Zurich Lake. Limnetic (navy) and benthic (gold) species are generally neatly differentiated in the dendrograms based on Euclidean distance shown on the x-axes. Percent methylation is displayed for each DMR from 0% (yellow) to 100% (indigo) with clear DNA methylation differences between species. Each column represents a sample and each row represents a DMR.

Fig. 3: Parallelism among lakes was assessed at the gene level based on the number of unique
genes overlapping (A) DMLs, (B) DMRs, and (C) outlier SNP-DML overlaps. Missing
comparisons indicate that no genes are shared between those lakes. The connected dots on the x-
axis represent the populations considered in each comparison and the intersection size shows the
number of parallel genes for each comparison. Set size gives the number of unique genes
overlapping the marker in each lake.

912