In silico docking screen identifies airway host protease targets for human

SERPINs

Joaquin Rodriguez Galvan[1], Maren de Vries[1], Ashley Fisher[1], Rachel A Prescott[1],

Keaton M Crosse[1], Ralf Duerr[1] and Meike Dittmann[1]*

[1] New York University Grossman School of Medicine, New York, NY

* Corresponding author

Abstract

Proteases play key roles in viral replication cycles. They can provide cleavage

maturation of viral glycoproteins, processing of viral polyproteins, or disassembly of viral

capsids. Thus, proteases constitute ideal targets for antiviral intervention –

pharmaceutically, by small molecule inhibitors, or naturally, by host immune responses.

Indeed, we and others have shown that individual members of the Serine protease

inhibitor (SERPIN) family have specific antiviral function by blocking proteolytic steps

inherent to viral replication cycles. Whether additional members of the large SERPIN

family possess antiviral activity and whether SERPINs function as part of the antiviral

cell-intrinsic immune response, is currently unknown. Here, we found that specific

SERPINs are produced upon infection with clinically relevant respiratory viruses *in vitro*

and *in vivo*, and in concert with classical interferon-stimulated genes. We next

developed a structure-based *in silico* screen to uncover non-canonical SERPIN-

protease pairs. We identified several SERPINs with potential antiviral function,

including: SERPINE1 targeting cathepsin L, required for SARS-CoV-2 entry; SERPINB8

targeting furin, required for glycoprotein maturation cleavage of numerous viruses; and

SERPINB2 targeting adenovirus protease, which suggests the first direct-acting antiviral

SERPIN. Our study demonstrates how proteolysis is modulated for antiviral defense

and how this process could inform antiviral targets against clinically relevant respiratory

pathogens.

Introduction

Respiratory viruses constitute a serious risk to human health. For example, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and influenza A virus (IAV) have originated major pandemics leading to severe respiratory impairment and death for millions of people and have caused considerable collateral damage to the economy and society [1]. Respiratory viruses encompass viruses from diverse families that show inherent tropism for the respiratory tract. Despite this commonality, and due to their considerable genetic and structural diversity, they can cause different clinical outcomes [2]. In particular, infections with IAV and SARS-CoV-2 are associated with a more severe prognosis in adults, whereas infections with other viruses, such as adenovirus or parainfluenza virus, tend to be milder in this age group [2,3]. These differences in pathogenesis may be attributed to a virus' ability to evade or downmodulate host immune responses or to viral tropism to the upper or lower respiratory tract.

Although respiratory viruses are highly heterogeneous in their replication strategies and lifecycles within a host cell, they all rely on a critical group of enzymes: proteases [4]. The origin of these proteases may be host or viral, and the roles they play in viral replication differ for each virus [4,5]. For example, most enveloped viruses rely on cleavage of their fusogenic glycoprotein for entry via fusion with host cell membranes, as is the case for parainfluenza viruses, influenza viruses, and coronaviruses, among others. The proteases executing viral glycoprotein cleavage are typically host-encoded, and are present at the plasma membrane (either intra- or extracellularly) or within endocytic or secretory pathways, aligning with the compartment in which the cleavage

event occurs [3,6-9]. Another protease function is for viral polyprotein processing, which

is the cleavage of viral polyprotein precursors into individual functional subunits upon

translation [9,10].This is the case for 3C-like and papain-like proteases (3CL^{pro} and

PL^{pro}), which are viral proteases encoded within the SARS-CoV-2 genome [9]. Other

proteases play roles in assembly/disassembly of viral particles inside the cell, as is the

case for the adenovirus-encoded protease (Ad^{pro}), which allows for assembly of the

protein capsid by cleaving capsid subunits, among other functions [11,12].

The multifaceted and integral role proteases play within the life cycles of many viruses

has led to their targeting for antiviral therapeutics. In fact, multiple antiviral FDA-

approved antiviral drugs are in fact small molecule protease inhibitors. These drugs

target the viral proteases of HIV-1, HCV and most recently SARS-CoV-2. The targeted

viral proteases all cleave large poly- chains into functional subunits, which are an

essential steps of these viral life cycles [13-15]. The clinical success of these antivirals

further illustrates the importance of proteases within viral lifecycles and highlights the

potential of protease inhibition for successful antiviral intervention.

Similar to pharmacological inhibition, humans encode genes to regulate and inhibit

proteolytic activity [16]. In this study, we focus on the largest family of human-encoded

protease inhibitors: serine protease inhibitors (SERPINs) [16]. SERPINs are structurally

and mechanistically conserved, consisting of a globular domain with critical central beta

sheets and a reactive center loop (RCL) that protrudes from the main SERPIN body [16].

The SERPIN mechanism of action is well-established at the molecular level. As a

SERPIN's RCL inserts into the protease active site, mimicking a substrate, the protease

catalytic triad residues initiate the proteolysis mechanism, resulting in a covalent bond

between the protease main active residue (nucleophile) and the P1 residue of the

SERPIN RCL [16]. Subsequently, the SERPIN loop triggers its insertion into the central

beta sheets, thereby trapping the protease in a covalent, suicide-like inhibitory complex

[16]. Specificity for a given SERPIN-mediated protease inhibition stems mainly from

appropriate fitting of a SERPIN's RCL into the protease active site. Other factors

determining SERPIN-protease specificity are actions that stabilize the SERPIN-protease

complex, which is intrinsically unstable. Stabilization is achieved through the action of

co-factors, or through characteristics inherent to the SERPIN loop structure itself, both

of which can prevent SERPIN collapse, aggregation, or cleavage [16-18].

Given the protease requirement for most respiratory viruses to successfully complete

their replication cycles, it seems intuitive that SERPINs may elicit antiviral activity.

Indeed, our group was the first to describe that SERPINE1 (plasminogen activator

inhibitor 1, PAI-1) is antiviral against influenza A H1 and H3 viruses, by preventing the

proteolytic maturation of those low pathogenic hemagglutinins [19]. Moreover, we found

that *SERPINE1* is an interferon-stimulated gene, indicating its role in the innate immune

response against viruses [19]. Since then, other groups have described the antiviral

effect of SERPINA1 against SARS-CoV-2 by inhibiting TMPRSS2, a protease required

for the plasma membrane-mediated route of entry of SARS-CoV-2 [20,21]. However,

beyond these two studies, little is known about other SERPIN family members in the

context of viral infection. Therefore, it would not be surprising to identify additional

SERPINs with different protease targets that may elicit antiviral activity against diverse

viruses in the human airways. Here, we investigate the following questions: which

SERPINs are expressed in the human airways, either at steady-state or upon viral

infection? What are their target proteases beyond those previously identified? Lastly,

based on this information, could they have antiviral potential against clinically relevant

respiratory viruses?

We found that multiple SERPINs that are differentially expressed to higher levels in

COVID-19 patient-derived bronchoalveolar lavage fluid (BALF) via readily available

scRNA-seq data. Complimentarily, we found 10 SERPINs upregulated in a model of the

human airway epithelium upon infection with a diverse panel of respiratory viruses. With

this information we then established a protein-protein docking screen that enabled us to

identify novel SERPIN-protease pairs based on structure. More specifically, based on

the fit of the SERPIN RCL into the protease active site, we predicted interaction and

potential inhibition of the protease by a given SERPIN. Using this approach, we found

several predicted SERPIN-protease pairs that may be key to understanding unexplored

aspects of the innate antiviral response against respiratory viruses.

Results

Results

SERPINs are differentially expressed in airway epithelial cells of COVID-19
patients.

SERPINs have been shown to play defined roles during inflammation, cancer
metastasis, angiogenesis, and neurodegenerative disease [22]. We were the first to
report that *SERPINE1* expression is enhanced on the mRNA level by interferon
signaling in human cell lines and primary human airway epithelial cultures [19]. While
most SERPIN functions to date have been characterized at steady-state, select
SERPINs are upregulated upon specific stimuli, including interferons, tumor necrosis
factors, or tumor growth factors [23-25]. All these stimuli can be produced during viral
infections [2,26,27].

We first asked whether, in addition to *SERPINE1*, other members of the SERPIN family
are also upregulated in response to viral infections. To begin answering this question,
we mined a published single cell RNA-seq (scRNA-seq) dataset of bronchoalveolar
lavage fluids (BALF) from patients with mild or severe COVID-19, or uninfected control
patients [28] (Figure 1A). This dataset allowed us to investigate mRNA expression levels
of individual *SERPIN*s in different cell types at the site of and during an acute viral
infection in humans. We first focused on epithelial cells, as they are the primary
replication site for acute SARS-CoV-2 infection and the main site of action for direct-
acting antiviral genes (Figure 1B). We found that *IRF1*, a bona fide interferon-stimulated
gene with antiviral function [29], was highly expressed in epithelial cells from patients

with mild and severe COVID-19, serving as positive control. A number of *SERPIN*s

were expressed at low levels in epithelial cells of uninfected patients, and at higher

levels in patients with mild or severe COVID-19, most notably *SERPINB2*, *SERPINB5*,

SERPINB11 and *SERPINF1*. Others were already robustly present in epithelial cells of

uninfected patients and further upregulated in COVID-19 patients, namely *SERPINB1*,

SERPINB3, *SERPINB6*, and *SERPING1*. *SERPINA1*, somewhat counterintuitively,

followed an inverse pattern and was an exception, as it was highly expressed in non-

infected patient epithelial cells but downregulated during SARS-CoV-2 infections.

Expression of *SERPINA3*, *SERPINA5*, *SERPINA9*, *SERPINA11*, *SERPINB7*,

SERPINC1, *SERPIND1*, *SERPINE2*, *SERPINE3* and *SERPINI1* changed less than 2-

fold in COVID-19 patients as compared to control patients (Figure 1B). Correlation

analysis revealed that *SERPIN* mRNA expression levels correlate with those of

canonical TNF-alpha and interferon-stimulated genes [29] (Supplementary Figure 1),

suggesting that *SERPINs* are expressed during viral-induced inflammation. Taken

together, our analyses show that a subset of *SERPIN*s are expressed high mRNA levels

during SARS-CoV-2 infection in airway epithelial cells.

Next, we asked whether this upregulation of *SERPIN*s in COVID-19 patients could also

be observed in other BALF cell types, and we determined log2-fold mRNA expression

changes compared to uninfected patients. Monocytes, innate lymphoid cells,

neutrophils, and dendritic cells, while readily detected in COVID-19 patients, were not

detected in uninfected patients, and thus -fold changes could not be determined for

these cell types. We found a greater number of highly-expressed *SERPINs* in epithelial

cells in comparison to immune cells such as macrophages, T cells, B cells, and NK cells

(Figure 1 C). In addition, the magnitude of *SERPIN* mRNA expression was higher

overall in epithelial cells from both mild and severe COVID-19 patients. Given that major

role of SERPINs is to inhibit proteases, we next analyzed expression of proteases in

epithelial cells. While we found only minimal changes in mRNA expression of airway

proteases upon SARS-CoV-2 infection, the protease family with the most robust

expression at baseline was cathepsins (Supplementary Figure 2). A notable exception

was *TMPRSS4*, which was >10-fold upregulated in COVID-19 patients as compared to

uninfected patients. Interestingly, TMPRSS4 has been shown to process SARS-CoV-2

Spike protein [7,30].

Figure 1. Integrated data from GEO accession number GSE145926 analyzed for *SERPIN* mRNA

expression. A. Single-cell RNAseq (scRNA-seq) was performed on bronchioalveolar lavage fluids

(BALF) of uninfected patients (n=2), and COVID-19 patients with mild (n=3) or severe (n=6) symptoms. **B.**

Expression of *SERPIN*s and of interferon-stimulated gene *IRF1* in epithelial cells. TPM, reads per million.

C. Fold-change of *SERPIN* expression across cell types relative to control patients. Number of

upregulated (>2 fold) SERPINs is annotated above, and names of most up- or downregulated *SERPIN*s

are annotated in the graph.

Together, our analyses suggest that *SERPIN*s are expressed to high levels in patients with COVID-19, predominantly in epithelial cells, and that *SERPIN* expression correlates with markers of the antiviral and inflammatory immune response.

SERPINs are expressed in response to respiratory virus infection in human airway epithelial cultures

While the COVID-19 pandemic allowed us to obtain unique samples such as BALF from acutely SARS-CoV-2-infected patients to study *SERPIN* and protease mRNA expression, such specimens are not readily available for other respiratory virus infections. Therefore, we next analyzed SERPIN expression *in vitro*, using a human airway epithelial culture (HAEC) model, which closely mimics the human bronchial epithelium in terms of cell composition (basal, ciliated, goblet and club cells), architecture, and secreted extracellular environment, including mucus and proteases [15]. We infected HAEC with a panel of clinically relevant respiratory viruses, including adenovirus 5 (AdV5), human reovirus (ReoV), human rhinovirus A (RhV), human parainfluenzavirus 3 (HPIV3), influenza A/California/2009 H1N1 virus (IAV) and SARS-CoV-2 ancestral isolate WA-1. At 24 or 72 h post infection, we harvested the cultures, isolated RNA, and quantified specific transcripts via RT-qPCR (Fig 2A).

Figure 2. *SERPIN* mRNA expression upon viral infection of human airway epithelial cultures

(HAEC). A. Schematic representation of a panel of respiratory viruses and of a trans-well containing a

polarized human airway epithelial culture (HAEC) growing in air-liquid interface. HAEC were infected

apically with human rhinovirus A (RhV), influenza A/Californa/09 H1N1 virus (IAV), human

parainfluenzavirus 3 (HPIV3), or SARS-CoV-2 WA-1 (SARS2), or basolaterally with human adenovirus 5

(AdV5) or human reovirus (ReoV). Cultures were lysed at indicated time points post-infection, total RNA

isolated and transcripts quantified by RT-qPCR. Prototype interferon-stimulated gene *IFITM3* mRNA

levels shown as -fold change over uninfected cultures over time. **B.** *SERPIN* and *IFITM3* mRNA, shown

as -fold change over uninfected cultures at indicated time points. n.d., not detected. **C.** Frequency of

upregulated (2-fold) SERPINs across viruses.

We first determined expression of the interferon-stimulated gene *IFITM3* (10) upon

infection with the different viruses. As expected, we found *IFITM3* upregulated upon

infection with all viruses tested (Figure 2B), validating the induction of an antiviral host

response in HAEC upon infection. Of note, robust (>8-fold) *IFITM3* upregulation for both

AdV5 and SARS-CoV-2 occurred later than for the other viruses, at the 72 h timepoint,

which may reflect different viral replication rates, differential viral immune escape

mechanisms, or both. Next, we examined *SERPIN* expression. Across all viral

infections, we found 10 *SERPIN*s differentially expressed to higher levels (Figure 2B):

SERPINB1, SERPINB2, SERPINB4, SERPINB8, SERPINB9, SERPINB10,

SERPINB13, SERPINC1, SERPINE1, and *SERPING1.* Others, for example *SERPINF1*,

are not differentially expressed upon infection with any virus, suggesting that they may

not be part of the induced antiviral response in airway epithelial cells. Interestingly, and

in agreement with the scRNA-seq data from COVID-19 patients (Figure 1), some

*SERPIN*s are downregulated, as is the case for *SERPINA1*. Furthermore, by analyzing

the frequencies of *SERPIN* expression across our panel of viruses, we observe that a

subset of *SERPIN*s that are upregulated upon infection with each of the 6 viruses (e.g.

SERPING1), which may indicate a broad antiviral function of these *SERPINs* (Figure

2C). In contrast, other *SERPIN*s are differentially expressed upon infection by select

viruses (e.g. *SERPINA1*), suggesting a different regulatory mechanism, and potentially

select specificity against viruses.

Table 1. SERPINs and published targets

SERPIN	protease(s)	target(s) involved in the lifecycle of*
A1	elastase, thrombin, trypsin, chymotrypsin, uPA	picornaviruses, RhV, HSV-1, HSV-2
A4	tissue kallikrein	IAV
A5	protein C, uPA, kallikrein	IAV
A10	factor Xa	HIV, SARS1, IAV, IBV, HSV, PIV-1, AdV, CMV, RSV, DENV
B1	protease 3, cathepsin G, elastase, Cathepsin L	RPV, HIV1, IAV, SARS1, SARS2, RhV
B2	uPA, tPA	HPV, IAV
B3	cathepsin G, cathepsin L, cathepsin K	SARS1, SARS2
B4	cathepsin G, chymase	IAV
B6	cathepsin G, thrombin, chymotrypsin, plasmin	HIV-1, HSV-2, IAV, HEV, HCV
B8	furin	EBOV, SINV, CoV, RSV, IAV, SFV, BDV, PIV, HSV-1, HSV-2, SARS2
B9	granzyme B, subtilisin A	HSV-1, HSV-2, IAV
B10	trypsin	IAV
B13	cathepsin L, cathepsin K	SARS1, SARS2
C1	thrombin, factor Xa	HIV-1, HSV-2, PIV, IAV, IBV, RSV, AdV, CMV
D1	thrombin	HSV-1, HSV-2
E1	uPA/tPA, TMPRSS2, HAT, trypsin	DENV, IAV, IBV, SeV, HCV, HSV-1, HSV-2
F2	plasmin	IAV
G1	complement C1 esterase	IAV
I1	uPA, urokinase, plasmin	IAV

Abbreviations: AdV, adenovirus; BDV, borna disease virus; CMV, cytomegalovirus; CoV, coronavirus; DENV, dengue virus; HCV, hepatitis C virus; HEV, hepatitis E virus; HSV, herpes sim- plex virus; IAV, influenza A virus; IBV, influenza B virus; PIV, parainfluenzavirus; RPV, rabbitpox virus; RSV, respiratory syncytial virus; RV, rotavirus; SARS1, SARS-CoV; SARS2, SARS-CoV-2; SFV, semliki forest virus; tPA, tissue plasminogen activator; uPA, urokinase plasminogen activator. * assembled from [19,37-62]

235
236
237
With this,
our data
indicate
that a
subgroup of
*SERPIN*s
are
upregulated
in response
to viral
infection,
which
correlates
with the
induction of
a reference

ISG, thus indicating that some *SERPIN* family members may be part of the cell-intrinsic

antiviral response in the human airway epithelial epithelium.

Development of an *in silico* protein-protein docking screen to identify SERPIN-

protease pairs

So far, we have shown that *SERPIN*s such as *SERPINB1, SERPINB2, SERPINB4,*

SERPINB8, SERPINB9, SERPINB10, SERPINB13, SERPINC1, SERPINE1, and

SERPING1 are upregulated upon diverse viral infections, suggesting that they may be

part of the cell-intrinsic antiviral response. Due to their function in inhibiting proteases,

we hypothesize that these SERPINs inhibit viruses by hindering proteases that are

required in their respective viral life cycles. Although SERPINs are not recognized

broadly as antiviral proteins, remarkably, many canonical SERPINs target proteases are

identical to those required during viral life cycles (Table 1). However, the list of

experimentally identified SERPIN targets may not be complete. Indeed, we previously

found that SERPINE1, well-characterized to inhibit uPA and tPA [16], also inhibits

human airway trypsin and TMPRSS2 [19]. Subsequently, it is feasible that we may yet

discover SERPINs with previously unappreciated target proteases that are critical in

viral infections.

To identify novel SERPIN-protease pairs, we developed an *in silico* screen based on 3D

protein structures and leveraging our molecular knowledge of the SERPIN mechanism

of action. A SERPIN's specificity for a given target protease is predominantly

determined by the fit of the protruding SERPIN RCL into the protease active site [16,62].

A good fit increases the likelihood for the formation of a SERPIN-protease Michaelis

complex [18], followed by a nucleophilic attack by the protease active residue on the

SERPIN RCL, and, ultimately, for the formation of a covalent link resulting in irreversible

protease inhibition. Therefore, structural modeling of the fit between a SERPIN's RCL

and a protease active site can be used to predict a favorable interaction, and thus

identify novel SERPIN targets.

Given that our previous data has focused on SERPIN expression in airway epithelial

cells, and that respiratory viruses are a clinically important group, we focused our

search on host proteases expressed in the human airways, and proteases encoded by

respiratory viruses. We first curated a database of 52 proteases expressed in the

respiratory tract, based on the LungMAP human expression database (Figure 3A) [63].

The final database contained members of the cysteine, serine, aspartic and threonine

protease families. Next, we obtained, if publicly available, the 3D structures of each

individual protease and each of the 10 SERPIN candidates. If not publicly available, we

homology-modeled the 3D structure using Swiss Modeler 2.0 [64-66] (Figure 3A). Next,

we mapped all protease active sites and all SERPIN RCL by assessing individual

entries on UniProt [67], with the goal of performing protein-protein docking of each

individual protease with each individual SERPIN via the HADDOCK 2.4 server [68,69].

HADDOCK is an informed protein-protein docking server, which docks regions of

interest and predicts many structural conformations of the complex, then ranks them by

score (Figure 3A). The calculated HADDOCK score is proportional to the Van der

Waals, electrostatic, desolvation, and restraint energy of the complex, therefore allowing

for the identification of complexes with favorable interaction energy [70]. Since these

energy values are of negative nature, a lower HADDOCK score corresponds to a higher

likelihood of a favorable interaction. We used this information to guide our docking

screen. While specifying the eight central active residues of each SERPIN (P4-P4') that

dictate specificity [16,71], we docked the entire SERPIN RCL to the complete protease

active site, and identified putative binding pairs based on low HADDOCK scores.

To test the feasibility of our strategy, we docked known target and known non-target

proteases with SERPINB1 and determined their HADDOCK scores. For example,

SERPINB1 is a known inhibitor of Cathepsin G, but does not inhibit Granzyme A [72,73].

As expected, HADDOCK identified that the SERPINB1-Cathepsin A complex represents

a good fit of the RCL into the protease active site. In contrast, the SERPINB1 RCL fails

to fit into the active site of Granzyme A (Figure 3B). Indeed, the distance between active

residues of Cathepsin G and SERPINB1 is <5A, whereas the distance between active

residues of Granzyme A and SERPINB1 is multiple times broader (Figure 3B),

suggesting that the positive control pair exhibits better positioning of active residues for

initiation of the nucleophilic attack by the protease and furthering the SERPIN inhibitory

mechanism [16].

Figure 3. Establishing the *in silico* screen for identification of SERPIN-protease pairs. A. Workflow of the *in silico* docking screen. **B.** Validation of docking using as an example the structures of SERPINB1 with positive (Cathepsin G) or negative (Granzyme A) controls. **C.** Validation of raw scoring of known positive and negative hits. Example of raw HADDOCK scores for SERPINB1 positive controls (Cathepsin G, Elastase) or negative controls (RCL mutant + elastase, Granzyme A). **D.** Normalized scoring to

facilitate the detection of new SERPIN-protease interaction pairs. Violin plots of HADDOCK scores for all

SERPIN-protease pairs. Red line, mean negative control; shaded red, non-hit range; green line, mean

positive control, shaded green, hit range; red data points, negative controls; green data points, positive

controls. n=10 SERPINsm n=52 proteases.

We further validated our strategy by docking two known positive and two known

negative controls for each SERPIN and quantifying their interaction via HADDOCK. If no

negative controls were known, we generated SERPIN RCL *in silico* mutants by

converting P4-P4' residues to alanines, determining the structure of the mutated

SERPIN using Swiss Modeler 2.0 [64-66], and docking the mutant structure to a positive

control protease. For example, SERPINB1 docked to the positive controls Cathepsin G

and Elastase yielded more negative HADDOCK scores, while SERPINB1 docked to the

negative control Granzyme A or the SERPINB1 RCL mutant docked to Elastase yielded

more positive scores (Figure 3C). These results validated our strategy to assess

interactions between individual SERPINs and proteases.

Upon validation of our strategy, we performed docking for each of the 52 individual

proteases, with each of the 10 individual SERPINs or RCL mutant SERPINs, and

calculated a raw HADDOCK score for each of the pairs. We then adjusted our dataset

for SERPIN-SERPIN variation and mathematically discriminated pairs based on positive

and negative controls. To be able to define a positive (likely inhibitory) threshold

common to all SERPINs in this study, we created a normalized Z-score centered to the

mean of both positive and negative controls for each SERPIN (Figure 3D). Using this

modified HADDOCK score (Figure 3D) combined with visual assessment of the

SERPIN RCL fit into the protease active site (Figure 3B), we proceeded to screen our

panel of respiratory protease-SERPIN pairs in greater detail one by one (Figure 4A).

In silico **screen identifies potential SERPIN-protease pairs relevant to viral life**

cycles.

To visualize novel SERPIN targets that may have antiviral activity in the airways, we

plotted modified HADDOCK scores for each SERPIN-protease pair in a heatmap

(Figure 4A). Proteases are listed on the x-axis and include 52 host airway proteases

and 3 virally encoded proteases. The 10 SERPINs that we previously identified to be

expressed in HAEC upon viral infections (Figure 2) are listed on the y-axis.

First, we observe that known SERPIN-protease pairs display a low HADDOCK score

(shown in dark red; i.e. SERPINB1 and Cathepsin A) and known non-binders display a

high HADDOCK score (shown in dark grey; i.e. SERPINC1 and TMPRSS11C), as

expected. Second, we observe that individual SERPINs, such as SERPINC1,

SERPINB8 and SERPINB9, display low HADDOCK scores with only few host

proteases, suggesting that these SERPINs have high specificity. In contrast, other

SERPINs, such as SERPINB1 or SERPINB13, display low HADDOCK scores with

several host proteases along with proteases from different families, suggesting that

these SERPINs may be more promiscuous. Third, we observe that some host

proteases, notably most TMPRSS family members and Cathepsin E, do not show

negative HADDOCK scores with most SERPINs, suggesting that these proteases may

be generally less sensitive to the action of SERPINs. This may be particularly relevant

for viruses that rely on these proteases (particularly TMPRSS2 and TMPRSS4) to

complete their life cycles, such as SARS-CoV-2 and IAV [7,30]. In contrast, other host

proteases, most notably trypsin, plasma kallikrein and cathepsin A show low HADDOCK

scores with almost all SERPINs, suggesting that they may be targeted by multiple

SERPINs redundantly. Finally, we examined relevant viral proteases such as SARS-

CoV-2 3CL^{pro} and PL^{pro}, as well as Ad^{pro}. We observe a strikingly low HADDOCK score

for AdV5 protease with SERPINB2. Indeed, the docked structure of the Ad^{pro}-

SERPINB2 complex shows a good fit of the SERPINB2 RCL into the Ad^{pro} active site

(Figure 4B). Moreover, the structure reveals the presence of exosites, which are defined

as residues outside the active regions in both SERPINs and proteases that further

stabilize the interaction [62,71]. When looking at the protease active site in detail, we

observe that the active cysteine is positioned in proximity for a nucleophilic attack to the

P1-P1' central active residues of SERPINB2 (Figure 4C). This gives us confidence of a

good fit and potential subsequent completion of the SERPIN inhibitory mechanism. To

our knowledge, this is the first indication of a human SERPIN potentially targeting a

protease of viral origin.

Figure 4. *In silico* screen of 10 SERPINs against a panel of host respiratory or viral proteases.

A. Heatmap-formatted results of docking screen, where Z-scores centered to Haddock mean of control

SERPIN-protease pairs are shown in a color scale. Negative Z-scores (red) are identified hits, positive Z-

scores (grey) are non-hits. B. HADDOCK v2.4 complex of Adenovirus 5 Protease docked to SERPINB2,

showing appropriate Michaelis SERPIN-protease fitting and exosites. Interacting residues are highlighted

in sticks; visualized via Pymol v2.0. C. Zoomed-in view of the residues within the catalytic triad of

Adenovirus 5 Protease. Blue, nitrogen; red, oxygen; yellow, sulfur. Visualized on Pymol v2.0.

Discussion

Our group was the first to show that a SERPIN could target viral proteolysis and cause an antiviral effect, particularly by inhibiting IAV H1 cleavage [19]. Moreover, we found that SERPINE1 was able to inhibit proteases beyond its 'canonical' target uPA [19]. However, several questions remained: Is the antiviral phenotype SERPIN- and virus-specific? Are other SERPINs involved in responding to respiratory viral infection? Which other respiratory proteases are targeted by SERPINs? We show here that select SERPINs are upregulated upon SARS-CoV-2 infection in humans, particularly in SARS-CoV-2-permissive airway epithelial cells (Figure 1). Using a human airway epithelial culture *in vitro* model, we further identified SERPINs that are differentially expressed upon infection with other respiratory viruses (Fig 2). We developed and executed an *in silico* docking screen which identified novel SERPIN target proteases with relevance in respiratory virus lifecycles (Fig 4). In all, our study shows that SERPINs are upregulated upon viral infection and may target several proteases involved in the life cycles of respiratory viruses.

SERPINs have been studied in many contexts of human disease and sequencing efforts have shown tissue specificities in their expression [16,74]. Increasing data shows that SERPINs can be expressed in response to inflammatory cytokines [19,23-25,75]. In agreement with these data, we show that *SERPINs* are expressed in response to inflammation both in COVID-19 patients and in virus-infected human airway epithelial cultures. Interestingly, seven clade B *SERPINs* are differentially expressed to higher levels in our *in vitro* HAEC model; namely, *SERPINB1*, *SERPINB2*, *SERPINB4*,

SERPINB8, *SERPINB9*, *SERPINB10*, and *SERPINB13*. This suggests that they are produced in response to respiratory viral infection; however, the mechanism of SERPIN gene regulation remains unclear. Moreover, clade B SERPINs have been shown to emerge through gene duplications, indicative of host-pathogen mediated evolution. Interestingly, we observed a difference between our *in vitro* HAEC model and data obtained from BALF of SARS-CoV-2 infected patients, as we see additional SERPINs upregulated *in vivo*. This may be due to the presence of activated immune cells in the respiratory tract, which are not present in the HAEC model, because, as previously shown, specific SERPIN levels can be stimulated by macrophage-derived proinflammatory cytokines [76]. This represents a limitation of the HAEC approach, as it lacks non-epithelial sources of signaling molecules. However, our *in vitro* model still captures intra-epithelial signals, making it highly informative for respiratory virus biology at the host-pathogen interface. Our scRNA-seq data identified cathepsins as the most robustly expressed family of proteases, even in uninfected control patients, indicating their relevance in respiratory virus infections. Altogether, we show that SERPINs are upregulated upon infection with diverse respiratory viruses, suggesting that they may be part of the respiratory antiviral arsenal.

Molecular docking has been widely used in drug discovery, for instance to visualize and predict the fit of a small molecule inhibitor inside the catalytic site of an enzyme [77]. Additionally, docking screens are used to screen small molecule compounds to inhibit or enhance a relevant enzymatic reaction [77]. In contrast to this vast body of studies using docking to predict protein-drug interactions, there is limited data docking protein-protein

interactions. The HADDOCK platform allows for informed protein-protein docking,

optimizing the docked complex by "informing" the process; namely, using only the

regions of interests on both molecules to guide the interaction [68]. This greatly benefits

our study as it is known that the SERPIN mechanism of protease inhibition is specific to

the RCL reaching into the active site. However, to validate each score that HADDOCK

generates, we must visually assess whether the fit is appropriate, which is feasible for

the scale of this investigation, but may be a limitation for high-throughput use of this

method. Future optimization of this screen may focus on quantifying the fit-specific

interaction energies to help select appropriate complexes.

The HADDOCK platform provides a score of the putative complexes, which allows us to

quantitatively assess a SERPIN's ability to fit into the protease active site and predict

candidates for inhibition based on an informed protein-protein docking approach. This is

particularly important as a single amino acid change in a SERPIN RCL can drastically

shift protease targets [71]. Thus, structural modeling and docking is a powerful tool to

narrow down candidates when characterizing SERPIN activity, and our approach

provides a wide lens for screening candidate binding pairs. However, some limitations

are inherent to our strategy. For example, is does not consider the action of cofactors

on SERPIN specificity, as cofactors may not be co-crystalized with SERPINs or

proteases. One such example is SERPINC1, which requires the cofactor heparin to

mediate the interaction in a bridge-like manner between SERPINC1 and its canonical

target thrombin [16]. While SERPINC1-heparin-thrombin complex has been crystallized

and can be used as a template for modeling, other cofactors may not be known or have

been co-crystallized with any of the relevant molecules [78]. It is also important to note

that our screen does not take into account intracellular localization of SERPIN and

target protease. Both molecules need to meet and locate within the same subcellular

compartment or extracellular space for inhibition to occur. For instance, SERPINB8

targets furin, which localizes in the golgi apparatus [52,79,80]. When expressed,

SERPINB8 is located along the secretory pathway and primarily in the golgi apparatus,

therefore facilitating productive inhibition [79]. Despite this limitation of our approach, the

localization of our ten SERPINs of interest is described in the literature, as well as most

proteases. Moreover, some of the large protease families are localized in the same

compartment, such as TMPRSSs and cathepsins, which localize to the plasma

membrane and endosomes, respectively [30,81,82]. We used this information, when

available, to prioritize hits that we expect to colocalize or temporarily meet throughout

the same pathway, based on literature.

Viruses can utilize redundant host proteolytic activities; thus, a combination of SERPINs

may be required for an antiviral effect. Redundant proteolytic pathways have been

shown in SARS-CoV-2 and IAV, as their fusion glycoproteins can be processed by a

wide range of proteases [7,30,39,81]. This is facilitated by the presence of a polybasic

cleavage site in their fusogenic glycoproteins, which renders them accessibility to a

broad range of host respiratory proteases for activation [83]. From this, we can infer that

a combination of SERPINs may be required for a measurable antiviral effect. For these

reasons, hits from our screen will require future experimental validation.

Our *in silico* screen revealed that out of our panel of ten SERPINs upregulated during

viral infections, all were predicted to inhibit more than one protease, but some were

predicted to be more promiscuous than others. Particularly promiscuous is SERPINB1,

which is the most ubiquitously expressed SERPIN of the B clade [74] and has been

shown to block cathepsin proteases such as cathepsin G [16]. In contrast, other

SERPINs are more specific, as is the case for SERPINC1, which does not bind

favorably with many respiratory proteases *in silico*, likely because its targets are more

focused to blood clotting and coagulation pathways [16].

We identified several previously unknown SERPIN-protease pairs with potential function

in viral restriction. First, we identified a human SERPIN potentially targeting a viral

protease: SERPINB2 and Ad^{pro}. If confirmed experimentally, this would be the first

report of a direct-acting antiviral SERPIN, which not only represents cross-species

modulation of enzymes at the host-pathogen interface but could inform a novel antiviral

target. Second, we identified SERPINs potentially targeting host proteases that process

surface glycoproteins of multiple clinically impactful respiratory viruses. This includes

SERPINB8 inhibiting furin, which serves to mature glycoproteins of high-pathogenic

IAV, HPIV3 and SARS-CoV-2 [5,7,81,84-86]. Third, we identified additional putative

inhibitors of the SARS-CoV-2 lifecycle. This is of particular interest, as the current

COVID-19 pandemic has highlighted the need for tools that have the potential to inform

novel therapeutics. SARS-CoV-2 encodes two proteases: 3CL^{pro} PL^{pro}, which both

cleave polyprotein intermediates [9]. Inhibition of such proteases indeed stops viral

replication at a single cycle time frame[15]. We show that clade B SERPINs such as

SERPINB1, SERPINB2 and SERPINB13 may inhibit these proteases. Further, SARS-

CoV-2 relies on host proteases to mediate fusion with the plasma or endosomal

membranes and enter cells by cleaving the Spike (S) protein and triggering its fusion

peptide [9]. A key protease mediating this endosomal entry is cathepsin L, however

other cathepsin family members such as cathepsins V, B, K and S have also been

shown to cleave S [39]. The omicron lineage has been described to use the

cathepsin/endosomal pathway of entry as opposed to the plasma membrane, and

omicron mutations in S have been associated with this phenotype [87]. Interestingly, we

found cathepsins to be more prone to interaction with our panel of SERPINs. This adds

relevance towards describing all putative inhibitors of cathepsin L as a way to target the

currently predominant omicron clade.

We believe these studies to be an important initial step in characterizing the broad and

important role of SERPINs as an antiviral mechanism. Future directions include testing

for physical inhibition of the select proteases by their SERPIN hits. Furthermore, we will

characterize the mechanisms of the regulation of these SERPINs, and whether they are

part of particular antiviral signaling pathways. Lastly, we will perform antiviral assays in

overexpression models to assess the magnitude of antiviral activity elicited by

SERPINs.

Material and Methods

COVID19 Patient scRNA-seq data acquisition and analysis

Publicly available COVID-19 patient scRNA-seq data were obtained from the Gene

Expression Omnibus database under the accession code GSE145926 [28]. Aligned and

normalized transcript per million (TPM) values were obtained from the integrated

ToppCell web server [63]. Frequencies and log2 fold changes were calculated from the

normalized TPM values by the healthy controls. Upregulated SERPINs were defined as

those expressed to >2-fold higher levels in COVID-19 patients compared to uninfected

controls. Plots were generated in GraphPad Prism v9.

Differentiation of HAEC

Differentiation of hTert-immortalized bronchial basal progenitor cells into stratified

human airway epithelial cultures was performed as stated in the following publication

[15].

Infection of HAEC with respiratory virus panel

Bci-NS1.1 cells (kind gift of Dr. Ronald Crystal) were seeded into Pneumacult™-Ex Plus

Medium (StemCell) and passaged at least two times before plating ($7x10^4$ cells/well) on

rat-tail collagen type-1 coated permeable transwell membrane supports (6.5mm;

Corning Inc) to generate human airway epithelial cultures (HAE). HAE were maintained

in Pneumacult™-Ex Plus Medium until confluent, then grown at an air-liquid interface

with Pneumacult™-ALI Medium in the basal chamber for approximately 3 weeks to form

differentiated, polarized cultures that resemble *in vivo* pseudostratified mucociliary epithelium. For infection with influenza A/California/07/2009 virus (BEI resources, NR-13663), HPIV3-GFP (kind gift of Dr. Peter Collins), and Rhinovirus A16 (kind gift of Dr. Ann Palmenberg), cultures were washed twice with room-temperature PBS, then virus was added apically ($5x10^6$ PFU for IAV, $1.12x10^7$ FFU for PIV, and 0.39 $TCID_{50}$ for Rhinovirus) in 50 µl of PBS and incubated for one hour at 37°C for IAV and PIV, and at 4°C for 90 minutes for Rhinovirus. Viruses were aspirated and HAE cultures were washed apically once with room-temperature PBS, then incubated at 37°C until endpoint. For infection with human Reovirus T3 SA (kind gift of Dr. Terrence Dermody) and Adenovirus 5-GFP (kind gift of Dr. Patrick Hearing), transwells were placed inverted in a sterile 5cm petri dish bottom and all remaining basal media was aspirated from the basal side of the transwells. 25 µl of each virus was added to the basal side of the transwells ($1.3x10^8$ PFU for Reovirus and $8.85x10^5$ PFU for Adenovirus) and incubated for one hour at 37°C for Adenovirus and room temperature for Reovirus. Viruses were aspirated from the basal side of the transwells, and transwells were placed back into ALI media and incubated at 37°C until endpoint. At each endpoint, transwell membranes were submerged in 375 µl Buffer RLT (Qiagen) and cell lysate was frozen at -80°C for future RNA extraction.

RT-qPCR of virus-infected HAEC

Infected HAEC were lysed as described above, and RNA was isolated using the Qiagen RNeasy kit following manufacturer's instructions. Extracted RNA was then quantified via NanoDrop and diluted to correct difference in yield. Relative mRNA levels, normalized

to housekeeping gene *RPS11*, were determined by RT-qPCR (SuperScript III First

Strand Synthesis System, Life Technologies and PowerUP SYBR Green Master Mix,

Thermo Fisher Scientific). Primers used are listed in Supplementary Table 1.

Protease and SERPIN database & structure modeling

Protein structures were obtained directly from the Protein Data Bank (PDB) or modeled

by homology using Swiss PDB Modeler [64-66]. If obtained from the PDB, structures

were refined by removing solvents, co-factors, small molecules, and co-crystallized

atoms. Selection of proteases for later docking was based on the LungMAP database

[63] of human airway expressed proteases and protease families, mainly prioritizing

those expressed to >10 Log2TPM on epithelial cells of donors.

Active sites and SERPIN RCLs were mapped on to refined structures using the UniProt

server [67].

Protein-protein docking

Docking of complexes was performed using the HADDOCK v2.4 server [68,69].

Selection of primary residues to perform docking was based on the residues specified

on UniProt as the RCL's 8 critical residues found to dictate SERPIN specificity - P4-P4'

for SERPINs and the catalytic triad within the protease active site for proteases.

Secondary residues were defined on a range of 13A radius from the primary residues,

this ensures to consider the protease pockets that may fit RCL sidechains and SERPIN-

protease exosites that may stabilize interactions and complex formation. Furthermore,

the full RCL was given full flexibility, starting at the poly-alanine segment and ending at

the PF loop end motif. Upon submission and finish of an entry on the HADDOCK v2.4

server, we evaluated the resulting complex position and specifically selected complexes

that had RCLs fit inside the protease active site. If no such complex was found, the

protease was removed from the screen. The score of the visually fitting complexes was

then used to guide the initial optimization and calculate thresholds for the Z-scores.

Structures of the complexes were then saved and visualized on PyMoL v2.0 for

distance calculation and deeper examination of the positioned residues (Figure 4) [88].

Statistical analysis and docking screen scoring

To define the thresholds of SERPIN-protease docking, Haddock scores were

determined from at least two positive and two negative control proteases for each

SERPIN, known to bind or not bind SERPINs *in vitro,* respectively. For each SERPIN,

average HADDOCK scores were calculated separately for the positive and negative

controls, and the mean of both averaged scores served as the in-silico threshold of

SERPIN-protease interaction. The SERPIN-specific HADDOCK thresholds were used to

normalize the full data set of HADDOCK scores and to generate Z-scores, i.e., the

thresholds were set to zero. Violin plots were generated using GraphPad Prism v.9, and

clustered heatmaps of the normalized, Z-scored data were generated using the

ComplexHeatmap package in program R v.4.1.2 [89].

Acknowledgements

We thank all members of the Dittmann lab for their critical input on data and the manuscript. We also thank Dr. Ronald Crystal for the kind gift of Bci-NS1.1 cells, Dr. Terrence Dermody for Reovirus T3 SA, Dr. Ann Palmenberg for Rhinovirus A16, Dr. Peter Collins for PIV3-GFP, and Dr. Patrick Hearing for Adenovirus 5-GFP. Research was supported by the following grants from NIH/NIAID: R01AI143639 and R21AI139374 to M.D., as well as T32AI100853 to J.R.G. Work was further supported by The Vilcek Institute of Graduate Biomedical Sciences, and by NYU Grossman School of Medicine Startup funds.

References

1. Organization, W.H. WHO COVID-19 Dashboard. **2020**.
2. Weston, S.; Frieman, M.B. Respiratory Viruses. *Encyclopedia of Microbiology* **2019**, 85-101, doi:doi:10.1016/B978-0-12-801238-3.66161-5.
3. Henrickson, K.J. Parainfluenza viruses. *Clin Microbiol Rev* **2003**, *16*, 242-264, doi:10.1128/cmr.16.2.242-264.2003.
4. Kemp, G.; Webster, A.; Russell, W.C. Proteolysis is a key process in virus replication. *Essays Biochem* **1992**, *27*, 1-16.
5. Aguilar, H.C.; Buchholz, D.W. Finding proteases that make cells go viral. *J Biol Chem* **2020**, *295*, 11408-11409, doi:10.1074/jbc.H120.015153.
6. Forsythe, H.M.; Galvan, J.R.; Yu, Z.; Pinckney, S.; Reardon, P.; Cooley, R.B.; Zhu, P.; Rolland, A.D.; Prell, J.S.; Barbar, E. Multivalent binding of the partially disordered SARS-CoV-2 nucleocapsid phosphoprotein dimer to RNA. *Biophys J* **2021**, doi:10.1016/j.bpj.2021.03.023.
7. Bertram, S.; Glowacka, I.; Steffen, I.; Kühl, A.; Pöhlmann, S. Novel insights into proteolytic cleavage of influenza virus hemagglutinin. *Rev Med Virol* **2010**, *20*, 298-310, doi:10.1002/rmv.657.
8. Chang, A.; Dutch, R.E. Paramyxovirus fusion and entry: multiple paths to a common end. *Viruses* **2012**, *4*, 613-636, doi:10.3390/v4040613.
9. Masters, P.S. The molecular biology of coronaviruses. *Adv Virus Res* **2006**, *66*, 193-292, doi:10.1016/s0065-3527(06)66005-3.
10. Jensen, L.M.; Walker, E.J.; Jans, D.A.; Ghildyal, R. Proteases of human rhinovirus: role in infection. *Methods Mol Biol* **2015**, *1221*, 129-141, doi:10.1007/978-1-4939-1571-2_10.
11. Kulanayake, S.; Tikoo, S.K. Adenovirus Core Proteins: Structure and Function. *Viruses* **2021**, *13*, doi:10.3390/v13030388.
12. Greber, U.F.; Webster, P.; Weber, J.; Helenius, A. The role of the adenovirus protease on virus entry into cells. *EMBO J* **1996**, *15*, 1766-1777.
13. de Leuw, P.; Stephan, C. Protease inhibitors for the treatment of hepatitis C virus infection. *GMS Infect Dis* **2017**, *5*, Doc08, doi:10.3205/id000034.
14. Lv, Z.; Chu, Y.; Wang, Y. HIV protease inhibitors: a review of molecular selectivity and toxicity. *HIV AIDS (Auckl)* **2015**, *7*, 95-104, doi:10.2147/HIV.S79956.
15. de Vries, M.; Mohamed, A.S.; Prescott, R.A.; Valero-Jimenez, A.M.; Desvignes, L.; O'Connor, R.; Steppan, C.; Devlin, J.C.; Ivanova, E.; Herrera, A.; et al. A comparative analysis of SARS-CoV-2 antivirals characterizes 3CL. *J Virol* **2021**, doi:10.1128/JVI.01819-20.
16. Law, R.H.; Zhang, Q.; McGowan, S.; Buckle, A.M.; Silverman, G.A.; Wong, W.; Rosado, C.J.; Langendorf, C.G.; Pike, R.N.; Bird, P.I.; et al. An overview of the serpin superfamily. *Genome Biol* **2006**, *7*, 216, doi:10.1186/gb-2006-7-5-216.
17. Sanrattana, W.; Maas, C.; de Maat, S. SERPINs-From Trap to Treatment. *Front Med (Lausanne)* **2019**, *6*, 25, doi:10.3389/fmed.2019.00025.
18. Ye, S.; Cech, A.L.; Belmares, R.; Bergstrom, R.C.; Tong, Y.; Corey, D.R.; Kanost, M.R.; Goldsmith, E.J. The structure of a Michaelis serpin-protease complex. *Nat Struct Biol* **2001**, *8*, 979-983, doi:10.1038/nsb1101-979.

19. Dittmann, M.; Hoffmann, H.H.; Scull, M.A.; Gilmore, R.H.; Bell, K.L.; Ciancanelli, M.;
Wilson, S.J.; Crotta, S.; Yu, Y.; Flatley, B.; et al. A serpin shapes the extracellular
environment to prevent influenza A virus maturation. *Cell* **2015**, *160*, 631-643,
doi:10.1016/j.cell.2015.01.040.
20. Rosendal, E.; Mihai, I.S.; Becker, M.; Das, D.; Frängsmyr, L.; Persson, B.D.; Rankin, G.D.;
Gröning, R.; Trygg, J.; Forsell, M.; et al. Serine Protease Inhibitors Restrict Host
Susceptibility to SARS-CoV-2 Infections. *mBio* **2022**, *13*, e0089222,
doi:10.1128/mbio.00892-22.
21. Azouz, N.P.; Klingler, A.M.; Callahan, V.; Akhrymuk, I.V.; Elez, K.; Raich, L.; Henry, B.M.;
Benoit, J.L.; Benoit, S.W.; Noé, F.; et al. Alpha 1 Antitrypsin is an Inhibitor of the SARS-
CoV-2-Priming Protease TMPRSS2. *Pathog Immun* **2021**, *6*, 55-74,
doi:10.20411/pai.v6i1.408.
22. Heit, C.; Jackson, B.C.; McAndrews, M.; Wright, M.W.; Thompson, D.C.; Silverman, G.A.;
Nebert, D.W.; Vasiliou, V. Update of the human and mouse SERPIN gene superfamily.
Hum Genomics **2013**, *7*, 22, doi:10.1186/1479-7364-7-22.
23. Prada, A.E.; Zahedi, K.; Davis, A.E. Regulation of C1 inhibitor synthesis. *Immunobiology*
1998, *199*, 377-388, doi:10.1016/S0171-2985(98)80042-9.
24. He, W.; Tan, R.; Dai, C.; Li, Y.; Wang, D.; Hao, S.; Kahn, M.; Liu, Y. Plasminogen activator
inhibitor-1 is a transcriptional target of the canonical pathway of Wnt/beta-catenin
signaling. *J Biol Chem* **2010**, *285*, 24665-24675, doi:10.1074/jbc.M109.091256.
25. Dong, C.; Zhu, S.; Wang, T.; Yoon, W.; Goldschmidt-Clermont, P.J. Upregulation of PAI-1
is mediated through TGF-beta/Smad pathway in transplant arteriopathy. *J Heart Lung*
Transplant **2002**, *21*, 999-1008, doi:10.1016/s1053-2498(02)00403-5.
26. Lee, A.J.; Ashkar, A.A. The Dual Nature of Type I and Type II Interferons. *Front Immunol*
2018, *9*, 2061, doi:10.3389/fimmu.2018.02061.
27. Liu, T.; Zhang, L.; Joo, D.; Sun, S.C. NF-κB signaling in inflammation. *Signal Transduct*
Target Ther **2017**, *2*, doi:10.1038/sigtrans.2017.23.
28. Liao, M.; Liu, Y.; Yuan, J.; Wen, Y.; Xu, G.; Zhao, J.; Cheng, L.; Li, J.; Wang, X.; Wang, F.; et
al. Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19. *Nat*
Med **2020**, *26*, 842-844, doi:10.1038/s41591-020-0901-9.
29. Wang, W.; Xu, L.; Brandsma, J.H.; Wang, Y.; Hakim, M.S.; Zhou, X.; Yin, Y.; Fuhler, G.M.;
van der Laan, L.J.; van der Woude, C.J.; et al. Convergent Transcription of Interferon-
stimulated Genes by TNF-α and IFN-α Augments Antiviral Activity against HCV and HEV.
Sci Rep **2016**, *6*, 25482, doi:10.1038/srep25482.
30. Zang, R.; Gomez Castro, M.F.; McCune, B.T.; Zeng, Q.; Rothlauf, P.W.; Sonnek, N.M.; Liu,
Z.; Brulois, K.F.; Wang, X.; Greenberg, H.B.; et al. TMPRSS2 and TMPRSS4 promote SARS-
CoV-2 infection of human small intestinal enterocytes. *Sci Immunol* **2020**, *5*,
doi:10.1126/sciimmunol.abc3582.
31. Walters, M.S.; Gomi, K.; Ashbridge, B.; Moore, M.A.; Arbelaez, V.; Heldrich, J.; Ding, B.S.;
Rafii, S.; Staudt, M.R.; Crystal, R.G. Generation of a human airway epithelium derived
basal cell line with multipotent differentiation capacity. *Respir Res* **2013**, *14*, 135,
doi:10.1186/1465-9921-14-135.

32.   Lazarowitz, S.G.; Choppin, P.W. Enhancement of the infectivity of influenza A and B
723   viruses by proteolytic cleavage of the hemagglutinin polypeptide. *Virology* **1975**, *68*,
724   440-454, doi:10.1016/0042-6822(75)90285-8.

725   33.   Klenk, H.D.; Rott, R.; Orlich, M.; Blödorn, J. Activation of influenza A viruses by trypsin
726   treatment. *Virology* **1975**, *68*, 426-439, doi:10.1016/0042-6822(75)90284-6.

727   34.   Hamilton, B.S.; Whittaker, G.R. Cleavage activation of human-adapted influenza virus
728   subtypes by kallikrein-related peptidases 5 and 12. *J Biol Chem* **2013**, *288*, 17399-17407,
729   doi:10.1074/jbc.M112.440362.

730   35.   Golden, J.W.; Schiff, L.A. Neutrophil elastase, an acid-independent serine protease,
731   facilitates reovirus uncoating and infection in U937 promonocyte cells. *Virol J* **2005**, *2*,
732   48, doi:10.1186/1743-422X-2-48.

733   36.   Molla, A.; Hellen, C.U.; Wimmer, E. Inhibition of proteolytic activity of poliovirus and
734   rhinovirus 2A proteinases by elastase-specific inhibitors. *J Virol* **1993**, *67*, 4688-4695,
735   doi:10.1128/JVI.67.8.4688-4695.1993.

736   37.   Gong, D.; Farley, K.; White, M.; Hartshorn, K.L.; Benarafa, C.; Remold-O'Donnell, E.
737   Critical role of serpinB1 in regulating inflammatory responses in pulmonary influenza
738   infection. *J Infect Dis* **2011**, *204*, 592-600, doi:10.1093/infdis/jir352.

739   38.   Huang, I.C.; Bosch, B.J.; Li, F.; Li, W.; Lee, K.H.; Ghiran, S.; Vasilieva, N.; Dermody, T.S.;
740   Harrison, S.C.; Dormitzer, P.R.; et al. SARS coronavirus, but not human coronavirus NL63,
741   utilizes cathepsin L to infect ACE2-expressing cells. *J Biol Chem* **2006**, *281*, 3198-3203,
742   doi:10.1074/jbc.M508381200.

743   39.   Bollavaram, K.; Leeman, T.H.; Lee, M.W.; Kulkarni, A.; Upshaw, S.G.; Yang, J.; Song, H.;
744   Platt, M.O. Multiple sites on SARS-CoV-2 spike protein are susceptible to proteolysis by
745   cathepsins B, K, L, S, and V. *Protein Sci* **2021**, *30*, 1131-1143, doi:10.1002/pro.4073.

746   40.   Avril, L.E.; Di Martino-Ferrer, M.; Pignede, G.; Séman, M.; Gauthier, F. Identification of
747   the U-937 membrane-associated proteinase interacting with the V3 loop of HIV-1 gp120
748   as cathepsin G. *FEBS Lett* **1994**, *345*, 81-86, doi:10.1016/0014-5793(94)00410-2.

749   41.   Zhirnov, O.P.; Golyando, P.B.; Ovcharenko, A.V. Replication of influenza B virus in
750   chicken embryos is suppressed by exogenous aprotinin. *Arch Virol* **1994**, *135*, 209-216,
751   doi:10.1007/BF01309780.

752   42.   Visseren, F.L.; Bouwman, J.J.; Bouter, K.P.; Diepersloot, R.J.; de Groot, P.H.; Erkelens,
753   D.W. Procoagulant activity of endothelial cells after infection with respiratory viruses.
754   *Thromb Haemost* **2000**, *84*, 319-324.

755   43.   Du, L.; Kao, R.Y.; Zhou, Y.; He, Y.; Zhao, G.; Wong, C.; Jiang, S.; Yuen, K.Y.; Jin, D.Y.;
756   Zheng, B.J. Cleavage of spike protein of SARS coronavirus by protease factor Xa is
757   associated with viral infectivity. *Biochem Biophys Res Commun* **2007**, *359*, 174-179,
758   doi:10.1016/j.bbrc.2007.05.092.

759   44.   Sutherland, M.R.; Friedman, H.M.; Pryzdial, E.L. Thrombin enhances herpes simplex
760   virus infection of cells involving protease-activated receptor 1. *J Thromb Haemost* **2007**,
761   *5*, 1055-1061, doi:10.1111/j.1538-7836.2007.02441.x.

762   45.   Knabb, M.T.; Danielsen, C.A.; McShane-Kay, K.; Mbuy, G.K.; Woodruff, R.I. Herpes
763   simplex virus-type 2 infectivity and agents that block gap junctional intercellular
764   communication. *Virus Res* **2007**, *124*, 212-219, doi:10.1016/j.virusres.2006.11.006.

46.   Stokol, T.; Yeo, W.M.; Burnett, D.; DeAngelis, N.; Huang, T.; Osterrieder, N.; Catalfamo, J.
Equid herpesvirus type 1 activates platelets. *PLoS One* **2015**, *10*, e0122640,
doi:10.1371/journal.pone.0122640.

47.   Stieneke-Gröber, A.; Vey, M.; Angliker, H.; Shaw, E.; Thomas, G.; Roberts, C.; Klenk, H.D.;
Garten, W. Influenza virus hemagglutinin with multibasic cleavage site is activated by
furin, a subtilisin-like endoprotease. *EMBO J* **1992**, *11*, 2407-2414, doi:10.1002/j.1460-
2075.1992.tb05305.x.

48.   Esumi, M.; Ishibashi, M.; Yamaguchi, H.; Nakajima, S.; Tai, Y.; Kikuta, S.; Sugitani, M.;
Takayama, T.; Tahara, M.; Takeda, M.; et al. Transmembrane serine protease TMPRSS2
activates hepatitis C virus infection. *Hepatology* **2015**, *61*, 437-446,
doi:10.1002/hep.27426.

49.   Tang, T.; Jaimes, J.A.; Bidon, M.K.; Straus, M.R.; Daniel, S.; Whittaker, G.R. Proteolytic
Activation of SARS-CoV-2 Spike at the S1/S2 Boundary: Potential Role of Proteases
beyond Furin. *ACS Infect Dis* **2021**, *7*, 264-272, doi:10.1021/acsinfecdis.0c00701.

50.   Baron, J.; Tarnow, C.; Mayoli-Nüssle, D.; Schilling, E.; Meyer, D.; Hammami, M.;
Schwalm, F.; Steinmetzer, T.; Guan, Y.; Garten, W.; et al. Matriptase, HAT, and TMPRSS2
activate the hemagglutinin of H9N2 influenza A viruses. *J Virol* **2013**, *87*, 1811-1820,
doi:10.1128/JVI.02320-12.

51.   Zhang, X.; Fugère, M.; Day, R.; Kielian, M. Furin processing and proteolytic activation of
Semliki Forest virus. *J Virol* **2003**, *77*, 2981-2989, doi:10.1128/jvi.77.5.2981-2989.2003.

52.   Brennan, S.O.; Nakayama, K. Furin has the proalbumin substrate specificity and serpin
inhibitory properties of an in situ hepatic convertase. *FEBS Lett* **1994**, *338*, 147-151,
doi:10.1016/0014-5793(94)80353-6.

53.   Follis, K.E.; York, J.; Nunberg, J.H. Furin cleavage of the SARS coronavirus spike
glycoprotein enhances cell-cell fusion but does not affect virion entry. *Virology* **2006**,
350, 358-369, doi:10.1016/j.virol.2006.02.003.

54.   Goto, H.; Kawaoka, Y. A novel mechanism for the acquisition of virulence by a human
influenza A virus. *Proc Natl Acad Sci U S A* **1998**, *95*, 10224-10228,
doi:10.1073/pnas.95.17.10224.

55.   Moon, K.B.; Turner, P.C.; Moyer, R.W. SPI-1-dependent host range of rabbitpox virus
and complex formation with cathepsin G is associated with serpin motifs. *J Virol* **1999**,
73, 8999-9010, doi:10.1128/JVI.73.11.8999-9010.1999.

56.   Choi, H.K.; Tong, L.; Minor, W.; Dumas, P.; Boege, U.; Rossmann, M.G.; Wengler, G.
Structure of Sindbis virus core protein reveals a chymotrypsin-like serine proteinase and
the organization of the virion. *Nature* **1991**, *354*, 37-43, doi:10.1038/354037a0.

57.   Nava, V.E.; Rosen, A.; Veliuona, M.A.; Clem, R.J.; Levine, B.; Hardwick, J.M. Sindbis virus
induces apoptosis through a caspase-dependent, CrmA-sensitive pathway. *J Virol* **1998**,
72, 452-459, doi:10.1128/JVI.72.1.452-459.1998.

58.   Sugrue, R.J.; Brown, C.; Brown, G.; Aitken, J.; McL Rixon, H.W. Furin cleavage of the
respiratory syncytial virus fusion protein is not a requirement for its transport to the
surface of virus-infected cells. *J Gen Virol* **2001**, *82*, 1375-1386, doi:10.1099/0022-1317-
82-6-1375.

59.   Ortmann, D.; Ohuchi, M.; Angliker, H.; Shaw, E.; Garten, W.; Klenk, H.D. Proteolytic
cleavage of wild type and mutants of the F protein of human parainfluenza virus type 3

809 by two subtilisin-like endoproteases, furin and Kex2. *J Virol* **1994**, *68*, 2772-2776,
810 doi:10.1128/JVI.68.4.2772-2776.1994.

60. Richt, J.A.; Fürbringer, T.; Koch, A.; Pfeuffer, I.; Herden, C.; Bause-Niedrig, I.; Garten, W. Processing of the Borna disease virus glycoprotein gp94 by the subtilisin-like endoprotease furin. *J Virol* **1998**, *72*, 4528-4533, doi:10.1128/JVI.72.5.4528-4533.1998.

61. Hahm, B.; Han, D.S.; Back, S.H.; Song, O.K.; Cho, M.J.; Kim, C.J.; Shimotohno, K.; Jang, S.K. NS3-4A of hepatitis C virus is a chymotrypsin-like protease. *J Virol* **1995**, *69*, 2534-2539, doi:10.1128/JVI.69.4.2534-2539.1995.

62. Gettins, P.G.; Olson, S.T. Inhibitory serpins. New insights into their folding, polymerization, regulation and clearance. *Biochem J* **2016**, *473*, 2273-2293, doi:10.1042/BCJ20160014.

63. Ardini-Poleske, M.E.; Clark, R.F.; Ansong, C.; Carson, J.P.; Corley, R.A.; Deutsch, G.H.; Hagood, J.S.; Kaminski, N.; Mariani, T.J.; Potter, S.S.; et al. LungMAP: The Molecular Atlas of Lung Development Program. *Am J Physiol Lung Cell Mol Physiol* **2017**, *313*, L733-L740, doi:10.1152/ajplung.00139.2017.

64. Guex, N.; Peitsch, M.C.; Schwede, T. Automated comparative protein structure modeling with SWISS-MODEL and Swiss-PdbViewer: a historical perspective. *Electrophoresis* **2009**, *30 Suppl 1*, S162-173, doi:10.1002/elps.200900140.

65. Bienert, S.; Waterhouse, A.; de Beer, T.A.; Tauriello, G.; Studer, G.; Bordoli, L.; Schwede, T. The SWISS-MODEL Repository-new features and functionality. *Nucleic Acids Res* **2017**, *45*, D313-D319, doi:10.1093/nar/gkw1132.

66. Waterhouse, A.; Bertoni, M.; Bienert, S.; Studer, G.; Tauriello, G.; Gumienny, R.; Heer, F.T.; de Beer, T.A.P.; Rempfer, C.; Bordoli, L.; et al. SWISS-MODEL: homology modelling of protein structures and complexes. *Nucleic Acids Res* **2018**, *46*, W296-W303, doi:10.1093/nar/gky427.

67. Consortium, U. UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Res* **2021**, *49*, D480-D489, doi:10.1093/nar/gkaa1100.

68. de Vries, S.J.; van Dijk, M.; Bonvin, A.M. The HADDOCK web server for data-driven biomolecular docking. *Nat Protoc* **2010**, *5*, 883-897, doi:10.1038/nprot.2010.32.

69. Honorato, R.V.; Koukos, P.I.; Jiménez-García, B.; Tsaregorodtsev, A.; Verlato, M.; Giachetti, A.; Rosato, A.; Bonvin, A.M.J.J. Structural Biology in the Clouds: The WeNMR-EOSC Ecosystem. *Front Mol Biosci* **2021**, *8*, 729513, doi:10.3389/fmolb.2021.729513.

70. Siebenmorgen, T.; Zacharias, M. Evaluation of Predicted Protein-Protein Complexes by Binding Free Energy Simulations. *J Chem Theory Comput* **2019**, *15*, 2071-2086, doi:10.1021/acs.jctc.8b01022.

71. Marijanovic, E.M.; Fodor, J.; Riley, B.T.; Porebski, B.T.; Costa, M.G.S.; Kass, I.; Hoke, D.E.; McGowan, S.; Buckle, A.M. Reactive centre loop dynamics and serpin specificity. *Sci Rep* **2019**, *9*, 3870, doi:10.1038/s41598-019-40432-w.

72. van der Burgh, R.; Meeldijk, J.; Jongeneel, L.; Frenkel, J.; Bovenschen, N.; van Gijn, M.; Boes, M. Reduced serpinB9-mediated caspase-1 inhibition can contribute to autoinflammatory disease. *Oncotarget* **2016**, *7*, 19265-19271, doi:10.18632/oncotarget.8086.

73. Cooley, J.; Takayama, T.K.; Shapiro, S.D.; Schechter, N.M.; Remold-O'Donnell, E. The serpin MNEI inhibits elastase-like and chymotrypsin-like serine proteases through

efficient reactions at two active sites. *Biochemistry* **2001**, *40*, 15762-15770,
doi:10.1021/bi0113925.

74. Silverman, G.A.; Whisstock, J.C.; Askew, D.J.; Pak, S.C.; Luke, C.J.; Cataltepe, S.; Irving,
J.A.; Bird, P.I. Human clade B serpins (ov-serpins) belong to a cohort of evolutionarily
dispersed intracellular proteinase inhibitor clades that protect cells from promiscuous
proteolysis. *Cell Mol Life Sci* **2004**, *61*, 301-325, doi:10.1007/s00018-003-3240-3.

75. Lappin, D.; Whaley, K. Regulation of C1-inhibitor synthesis by interferons and other
agents. *Behring Inst Mitt* **1989**, 180-192.

76. Schroder, W.A.; Hirata, T.D.; Le, T.T.; Gardner, J.; Boyle, G.M.; Ellis, J.; Nakayama, E.;
Pathirana, D.; Nakaya, H.I.; Suhrbier, A. SerpinB2 inhibits migration and promotes a
resolution phase signature in large peritoneal macrophages. *Sci Rep* **2019**, *9*, 12421,
doi:10.1038/s41598-019-48741-w.

77. Meng, X.Y.; Zhang, H.X.; Mezei, M.; Cui, M. Molecular docking: a powerful approach for
structure-based drug discovery. *Curr Comput Aided Drug Des* **2011**, *7*, 146-157,
doi:10.2174/157340911795677602.

78. Jin, L.; Abrahams, J.P.; Skinner, R.; Petitou, M.; Pike, R.N.; Carrell, R.W. The anticoagulant
activation of antithrombin by heparin. *Proc Natl Acad Sci U S A* **1997**, *94*, 14683-14688,
doi:10.1073/pnas.94.26.14683.

79. Leblond, J.; Laprise, M.H.; Gaudreau, S.; Grondin, F.; Kisiel, W.; Dubois, C.M. The serpin
proteinase inhibitor 8: an endogenous furin inhibitor released from human platelets.
Thromb Haemost **2006**, *95*, 243-252, doi:10.1160/TH05-08-0561.

80. Izaguirre, G.; Arciniega, M.; Quezada, A.G. Specific and Selective Inhibitors of Proprotein
Convertases Engineered by Transferring Serpin B8 Reactive-Site and Exosite
Determinants of Reactivity to the Serpin α1PDX. *Biochemistry* **2019**, *58*, 1679-1688,
doi:10.1021/acs.biochem.8b01295.

81. Zhao, M.M.; Yang, W.L.; Yang, F.Y.; Zhang, L.; Huang, W.J.; Hou, W.; Fan, C.F.; Jin, R.H.;
Feng, Y.M.; Wang, Y.C.; et al. Cathepsin L plays a key role in SARS-CoV-2 infection in
humans and humanized mice and is a promising target for new drug development.
Signal Transduct Target Ther **2021**, *6*, 134, doi:10.1038/s41392-021-00558-8.

82. Pišlar, A.; Mitrović, A.; Sabotič, J.; Pečar Fonović, U.; Perišić Nanut, M.; Jakoš, T.; Senjor,
E.; Kos, J. The role of cysteine peptidases in coronavirus cell entry and replication: The
therapeutic potential of cathepsin inhibitors. *PLoS Pathog* **2020**, *16*, e1009013,
doi:10.1371/journal.ppat.1009013.

83. Peacock, T.P.; Goldhill, D.H.; Zhou, J.; Baillon, L.; Frise, R.; Swann, O.C.; Kugathasan, R.;
Penn, R.; Brown, J.C.; Sanchez-David, R.Y.; et al. The furin cleavage site in the SARS-CoV-
2 spike protein is required for transmission in ferrets. *Nat Microbiol* **2021**, *6*, 899-909,
doi:10.1038/s41564-021-00908-w.

84. Gamblin, S.J.; Haire, L.F.; Russell, R.J.; Stevens, D.J.; Xiao, B.; Ha, Y.; Vasisht, N.;
Steinhauer, D.A.; Daniels, R.S.; Elliot, A.; et al. The structure and receptor binding
properties of the 1918 influenza hemagglutinin. *Science* **2004**, *303*, 1838-1842,
doi:10.1126/science.1093155.

85. Yin, H.S.; Paterson, R.G.; Wen, X.; Lamb, R.A.; Jardetzky, T.S. Structure of the uncleaved
ectodomain of the paramyxovirus (hPIV3) fusion protein. *Proc Natl Acad Sci U S A* **2005**,
102, 9288-9293, doi:10.1073/pnas.0503989102.

86.   Porotto, M.; Murrell, M.; Greengard, O.; Moscona, A. Triggering of human parainfluenza
898   virus 3 fusion protein (F) by the hemagglutinin-neuraminidase (HN) protein: an HN
899   mutation diminishes the rate of F activation and fusion. *J Virol* **2003**, *77*, 3647-3654,
900   doi:10.1128/jvi.77.6.3647-3654.2003.
901   87.   Du, X.; Tang, H.; Gao, L.; Wu, Z.; Meng, F.; Yan, R.; Qiao, S.; An, J.; Wang, C.; Qin, F.X.
902   Omicron adopts a different strategy from Delta and other variants to adapt to host.
903   *Signal Transduct Target Ther* **2022**, *7*, 45, doi:10.1038/s41392-022-00903-5.
904   88.   pymol.bib<source-app name="BibDesk" version="1.5.2">BibDesk</source-app><rec-
905   number>1</rec-number><ref-type>34</ref-type>Schrodinger, LLC<secondary-
906   authors><subsidiary-authors></subsidiary-authors></secondary-authors>2015<pub-
907   dates>November</pub-dates>AxPyMOL<pdf-urls><related-urls></related-urls></pdf-
908   urls>unpublished19/08/2010<modified-date>22/12/2015</modified-date>
909   pymol.bib<source-app name="BibDesk" version="1.5.2">BibDesk</source-app><rec-
910   number>2</rec-number><ref-type>34</ref-type>Schrodinger,, LLC<secondary-
911   authors><subsidiary-authors></subsidiary-authors></secondary-authors>2015<pub-
912   dates>November</pub-dates>JyMOL<pdf-urls><related-urls></related-urls></pdf-
913   urls>unpublished19/08/2010<modified-date>22/12/2015</modified-date>
914   pymol.bib<source-app name="BibDesk" version="1.5.2">BibDesk</source-app><rec-
915   number>3</rec-number><ref-type>34</ref-type>Schrodinger,, LLC<secondary-
916   authors><subsidiary-authors></subsidiary-authors></secondary-authors>2015<pub-
917   dates>November</pub-dates>PyMOL<pdf-urls><related-urls></related-urls></pdf-
918   urls><research-notes>PyMOL The PyMOL Molecular Graphics System, Version 1.8,
919   Schrödinger, LLC.</research-notes>unpublished19/08/2010<modified-
920   date>22/12/2015</modified-date>.
921   89.   Team, R.C. A Language and Environment for Statistical Computing. **2014**.
922