***Streptomyces* alleviate abiotic stress in plant by producing pteridic acids**

Zhijie Yang[1], Emil Strøbech[1], Yijun Qiao[1], Naga Charan Konakall[2], Pernille Harris[3], Gundela Peschel[4],

Miriam Agler-Rosenbaum[4], Tilmann Weber[5], Erik Andreasson[2], Ling Ding[1,*]

[1] Department of Biotechnology and Biomedicine, Technical University of Denmark, Søltofts Plads,

Building 221, 2800 Kgs. Lyngby, Denmark.

[2] Department of Plant Protection Biology, Swedish University of Agricultural Sciences, Sundsvägen

14, SE-230 53, Alnarp, Sweden.

[3] Department of Chemistry, Technical University of Denmark, Søltofts Plads, Building 206, 2800 Kgs.

Lyngby, Denmark.

[4] Leibniz Institute for Natural Product Research and Infection Biology - Hans Knöll Institute (HKI),

Beutenbergstr. 11a, 07745 Jena, Germany.

[5] Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark,

Kemitorvet, Building 220, 2800 Kgs. Lyngby, Denmark.

*Corresponding author: Ling Ding lidi@dtu.dk

Abstract

Soil microbiota can confer fitness advantages to plants and increase crop resilience to drought and

other abiotic stressors. However, there is little evidence on the mechanisms correlating a microbial

trait with plant abiotic stress tolerance. Here, we report that *Streptomyces* effectively alleviates the

drought and salinity stress by producing spiroketal polyketide pteridic acid H (**1**) and its isomer F (**2**),

both of which promote root growth in *Arabidopsis* at a concentration of 1.3 nM under abiotic stress.

Pteridic acids induce stress response genes expression in salinity-stressed *Arabidopsis* seedlings. The

bifunctional biosynthetic gene cluster of pteridic acids and antimicrobial elaiophylin is confirmed in

vivo and mainly disseminated by vertical transmission which is geographically distributed in various

environments. This discovery reveals a perspective for understanding plant-*Streptomyces* interactions

and provides a promising approach for utilising beneficial *Streptomyces* and their secondary

metabolites in agriculture to mitigate the detrimental effects of climate change.

Introduction

According to the Food and Agriculture Organization of the United Nations, climate change generates considerable uncertainty about future water availability in many regions. Increased water scarcity under climate change will present a major challenge for climate adaptation, while sea-level rise will affect the salinity of surface and groundwater in coastal areas. Stress caused by climate change has led to increased agricultural losses and threatened global food security.[1,2] Drought is considered the most damaging environmental stress, which directly affects the entire growth period of plant seeds from germination to final fruiting.[3] Drought stress can lead to increased plant osmotic regulators, the inhibition of photosynthesis, and the change of plant endogenous hormone content.[4-6] Drought stress also induces reactive oxygen species (ROS), such as superoxide radicals, hydrogen peroxide, and hydroxyl radicals, leading to oxidative stress.[7] At high concentrations, ROS can cause damage at various tissue levels, for example, triggering lipid peroxidation and membrane degradation in plants.[8,9] Crop loss due to soil salinisation is another increasing threat to agriculture worldwide, which is more severe in agricultural land in coastal and arid regions.[10] Irrigation with saline water, low precipitation, and high evapotranspiration are key factors of the rapid salinisation of agricultural land.[11] These abiotic stresses of drought and salinity have brought unprecedented challenges to the development of crop farming. Compared to the heavy use of chemical fertilisers, the use of plant growth-promoting bacteria to improve plant growth under drought and salinity environments is more sustainable and gaining more attention.[12,13]

Soil microbial communities are critical to plant health and their resistance to both biotic and abiotic stressors, such as pathogens, drought, salinity, and heavy metal pollution.[14] A few studies have demonstrated that many beneficial soil bacteria harbour plant growth-promoting activities, *e.g.* by helping plants with disease suppression,[15] nutrient acquisition,[16] phosphorus uptake[17] and nitrogen fixations.[18] Beneficial root microbiota also regulates biosynthetic pathways in the plant itself, leading to differential alterations in the plant metabolome in response to stresses.[19] *Streptomyces* are Gram-positive filamentous bacteria, widely distributed in soil and marine environments. While they have long been considered the richest source of bioactive secondary metabolites,[20] *Streptomyces* have recently drawn attention as a class of plant growth-promoting bacteria that help plants respond to adversity stress.[21] The growing evidence showed that *Streptomyces* can promote plant growth or tolerance to stressors in direct or indirect ways, by secreting plant growth regulator auxin (indole-3-acetic acid, IAA) and siderophores, inducing systemic resistance in plants, and regulating the rhizosphere microbiome via producing antibacterial compounds or signalling molecules.[22-24] Notably, the commercial product Actinovate® and Mycostop® are two *Streptomyces*-based formulations that

have been widely used to suppress a wide range of diseases in a variety of crop groups as a biological fungicide/bactericide for the long term. Recently, the enrichment of *Streptomyces* has also been shown to play a subsequent role in the drought/salt tolerance of plants.[25] Despite the widespread claims of efficacy of inoculation of plant growth-promoting *Streptomyces*, the molecular basis of the growth-promoting effects and the key role of secondary/specialised metabolites in this process are largely unknown.

Here, we report that *Streptomyces iranensis* HM 35 has profound beneficial effects on helping barley alleviate osmotic, drought and salinity stress. The active components were identified as a bioactive spiroketal polyketide pteridic acids H (**1**) and its isomer F (**2**) through large-scale fermentation and bioactivity-directed purification followed by NMR, MS, and X-ray crystallography. The abiotic stress mitigating effect of pteridic acids H and F have been confirmed on the model plant *Arabidopsis thaliana,* where it effectively reversed both drought and salinity stress as phytohormone-like small biomolecules at concentrations as low as 0.5 ng mL^{-1} (1.3 nM). Moreover, the Biosynthetic Gene Cluster (BGC) of pteridic acids (*pta*) was identified and analysed in silico and functionally confirmed by in vivo CRISPR-based genome editing. We have furthermore conducted a survey of 81 potential producers of pteridic acids, which are widely distributed around the world. Phylogenetic and comparative genomic analysis of *pta*-containing *Streptomyces* suggested that these strains have evolutionary convergence in disseminating *pta* BGC through main vertical transmission and occasional horizontal gene transfer. In summary, we reveal a strategy of *Streptomyces* to secret plant growth regulators that help plants cope with abiotic stresses, which is a promising alternative solution for plant development and crop yields under the current climate change-induced environmental stresses.

Results

Abiotic Stress-Mitigating Activities Exhibited by *S. iranensis*. The promotion of barley growth induced by *S. iranensis* was tested under multiple abiotic stresses including osmotic, salinity and drought. The osmotic stress experiment was simulated using soils supplemented with 20% (w/v) PEG-6000 by transiently reducing the water potential (Ψ_W) of the plant. We found that the *S. iranensis* played a significant role in alleviating osmotic stress in barley seedlings. The treated seedlings showed a significant increase in height, fresh weight, and dry weight compared to the control group without any extra treatment (Fig. 1a). The culture broth of *S. iranensis* also showed considerable activity for the growth of barley in alleviating salinity stress mediated by 100 mM NaCl (Fig. 1b), while *S. iranensis* was not significantly enriched in the soil around the roots of barley seedlings (Supplementary Fig. 1). Additionally, based on the analysis of barley seedling phenotypes, the treatment with *S.*

iranensis resulted in a significant improvement in plant growth recovery from drought stress (Fig. 1c).

Surprisingly, *S. iranensis* promoted the growth of barley seedlings even in non-stress growth condition

and thus indicate that *S. iranensis* may have potential for use in biocontrol (Supplementary Fig. 2).

Genomic and metabolomic profiles of *S. iranensis*. To reveal the potential bioactive components,

we first annotated BGCs responsible for the biosynthesis of secondary metabolites in *S. iranensis* using

antiSMASH 6.0.[26] Genome sequence analysis of *S. iranensis* revealed the presence of 50 putative

secondary metabolites BGCs with a variety of biosynthetic categories (Supplementary Tab. 1). The

clusters 4, 7, 8, 9, 24, 32, 36 and 41 were annotated to have greater than 85% similarity with BGCs

responsible for the biosynthesis of coelichelin, azalomycin, nigericin, elaiophylin, desferrioxamin,

ectoine, rapamycin and hygrocin, respectively. Their corresponding products were also detected and

identified through High-resolution Liquid Chromatography-tandem Mass Spectrometry (HR-LC-

MS/MS) as well as Global Natural Products Social (GNPS) molecular networking (Fig. 1d,

Supplementary Fig. 3).[27] However, there are still a number of metabolites of *S. iranensis* are still

unknown. Since none of the previously identified compounds have been associated with mitigating

abiotic stress in plants, we were prompted to expand the fermentation process and identify the potential

bioactive compounds.

Characterisation of the bioactive compound pteridic acid. To uncover the bioactive components,

fermentation of *S. iranensis* was scaled up to 175 litres and the culture broth was subjected to

separation through open-column chromatography on Amberchrom CG161Me resin, silica gel, and

Sephadex LH-20. Bioactivity-guided fractionation led to the isolation of bioactive compound **1** (15.0

mg) together with another structurally related compound **2** (4.0 mg) (Fig. 1e).

The bioactive component compound **1** was isolated as a white solid. Its formula of $C_{21}H_{34}O_6$ was

deduced by *m/z* 383.2439 $[M+H]^+$ (calculated for 383.2428, Δ 2.84 ppm). The ^1H NMR spectrum

exhibited signals for four olefinic protons (δ 7.33, 6.26, 6.13, 5.90) corresponding to two conjugated

double bonds, four oxygen-bearing methines (δ 3.85, 3.69, 3.59, 3.43), five methyls, and other aliphatic

protons. The ^{13}C NMR spectrum indicated the presence of one carbonyl group (δ 168.7) and one

oxygen-bearing quaternary carbon (δ 103.2, C-11). The COSY spectrum established two partial

substructures, which could be connected via a spiral function by analysing HSQC and HMBC

correlations (Supplementary Fig. 4-11). Compound **1** was crystallised in methanol solution, and the

structure was determined via X-ray crystallography (Supplementary Fig. 12). Therefore, compound **1**

was identified as a new *Streptomyce*s-derived natural product named pteridic acid H.

Compound **2**, a white solid, is an isomer of **1** since MS deduced the same molecular formula of

$C_{21}H_{34}O_6$. The ^1H NMR spectrum exhibited signals for four olefinic protons (δ 7.16, 6.25, 6.07, 5.97)

corresponding to two conjugated double bonds, four oxygen-bearing methines (δ 3.88, 3.66, 3.56,
3.32), five methyls, and other aliphatic protons. The ^{13}C NMR spectrum indicated the presence of one
carbonyl group (δ 170.2) and one oxygen-bearing quaternary carbon (δ 103.2, C-11). HSQC and
HMBC correlations confirmed a spiroketal skeleton. NOESY spectrum confirmed its relative
configurations, where correlations between H-21 and H-13 and H-15, H-7 and H-12a, and H6, H-10,
and Me-18 were observed. The key NOESY correlations between H-7 and H-12a revealed a different
spiroketal structure than **1** (Supplementary Fig. 11, 13-20). This can be reflected by the relative up-
field NMR data for C-12 (δ 33.8 vs δ 37.5 in **1**). Compound **2** was identified as pteridic acid F,
previously isolated from *Streptomyces pseudoverticillus* YN17707 and a marine-derived *Streptomyces*
sp. SCSGAA 0027.[28,29]

Abiotic stress mitigation of pteridic acids *in planta*. Initially, we tested the effects of different
concentrations of pteridic acids H and F on *Arabidopsis* growth in the absence of abiotic stress. A
concentration of 0.5 ng ml^{-1} of both pteridic acids H and F was found to significantly promote the
growth of *Arabidopsis* seedlings (Supplementary Fig. 21). Under drought stress, pteridic acid H at a
concentration of 0.5 ng mL^{-1} increased the root length and fresh weight of *Arabidopsis* seedlings by
54.5% and 89%, respectively, and its activity was significantly better than the IAA and ABA at the
same molar concentration (Fig. 2a, 2c). The treatment of pteridic acid F also showed great activity in
alleviating drought stress, and the root length and fresh weight were increased by 30.5% and 56.7%,
respectively (Fig. 2c). Pteridic acids H and F also showed significant activity in alleviating NaCl-
mediated salinity stress (Fig. 2b. 2d). Compared to the non-treated groups, the treatment of 0.5 ng mL^{-1}
pteridic acids H and F increased root length of *Arabidopsis* seedlings by 74.0% and 61.8%, as well
as fresh weight by 126.2% and 110.9%, respectively (Fig. 2d).
To gain insights into how plants respond to pteridic acids, we performed quantitative real-time PCR
(qRT-PCR) analysis to measure the expression levels of stress response genes TONOPLAST
INTRINSIC PROTEIN 2;3 (TIP2;3) and SALT OVERLY SENSITIVE 1 (SOS1) in *Arabidopsis*
seedlings, induced by pteridic acids H and F under salinity stress. Compared to the control, the relative
expression level of TIP2;3 was significantly increased following both 1-hour and 24-hour treatments,
while the transcript levels of SOS1 was down-regulated at 1-hour while up- regulated at 24-hour (Fig.
2e). This result indicated that pteridic acids H and F regulated the TIP2;3 and SOS1 to activate stress-
resistance responses in plants.
A previous study suggested that pteridic acids A and B might have a plant growth-promoting effect
like IAA and could stimulate the formation of adventitious roots in kidney beans.[28] However, we
observed that pteridic acids H and F displayed the different IAA-induced phenotypes. Pteridic acids

did not exhibit the function to significantly promote lateral root growth of *Arabidopsis* seedlings like

IAA (Fig. 2f, 2g).[30] We also found that pteridic acids were also not capable of promoting the formation

of adventitious roots in kidney beans, as shown in Supplementary Fig. S22. Except for drought and

salinity stress, we also tested the $CuSO_4$-mediated heavy metal stress alleviation activity of pteridic

acids in mung beans. The results showed that the 1 ng mL^{-1} pteridic acid H was effective as ABA in

helping mung beans to relieve heavy metal stress (Supplementary Fig. S23). In conclusion, pteridic

acids H and F are widely applicable potent plant growth regulators produced by *Streptomyces* to assist

plants in coping with different abiotic stress.

Biosynthesis of pteridic acids. The retro-biosynthesis analysis indicated that pteridic acids could

derive from a modular type I polyketide synthase. A putative *pta* BGC was identified in the whole

genome sequence of S. *iranensis*, which shows 87% antiSMASH similarity to the BGC of elaiophylin

(BGC0000053 in MiBiG database).[31] The *pta* BGC spans approximately 56 kb and encodes 20

individual biosynthetic genes responsible for the biosynthesis of core polyketide backbones, precursor,

glycosylated substituents, transporters and regulators (Supplementary Tab. 2). The five consecutive

Type I polyketide synthase (PKS) encoding genes within the *pta* BGC consist of one loading module

and seven extender modules, which are sequentially extended to form a linear polyketide chain by

ketosynthase (KS) domain, acyltransferase (AT) domain, acyl carrier protein (ACP), with additional

ketoreductase (KR), dehydratase (DH), and enoyl reductase (ER) domains. The substrate specificity

predictions for individual AT domains fit well with the structure of pteridic acids (Supplementary Tab.

3). The last "Asn" residue is absent in the conserved Lys-Ser-Tyr-Asn tetrad of the KR domain in

module 3 (PtaB), which is predicted to be inactive (Supplementary Fig. 24). This is consistent with the

nonreduced carbonyl group on the α-carbon in module 3. The first DH domain in module 1 is inactive

since it does not have a conserved active motif LxxHxxGxxxxP (Supplementary Fig. 25). Following

the thioesterase-mediated release of the polyketide chain, the 6,6-spiroketal core structure is likely

formed by spontaneous spiroketalisation of the carbonyl group on C11 and the two hydroxyl groups

on C17 and C25. Following a loss of H_2O, two differentially oriented spirocyclic rings were formed

to yield pteridic acids F and H (Fig. 3a). Remarkably, pteridic acid H showed molecular instability

under extreme conditions. In the water solution with high temperature (65°C) or acidity (pH = 3),

pteridic acid H is transformed into pteridic acid F (Supplementary Fig. 26). A similar spontaneous

transformation from (*S*) to (*R*) chirality at the centre of the spiroketal ring was also observed in 6,6-

spiroketal avermectin.[32]

CRISPR base editing in *S. iranensis*. To validate the *in silico* prediction of BGC, we utilized the efficient base editing tool CRISPR-cBEST to further confirm *pta* BGC.[33] As a non-model *Streptomyces* strain, *S. iranensis* is hard to genetically manipulate through intergeneric conjugation.[34] Therefore, the conjugation process was systematically optimised in this study (Supplementary Fig. 27). The core polyketide synthase *ptaA* was targeted and inactivated by converting a TGG (Trp) codon at position 916 into the stop codon TAA using CRISPR base editing. The editing event was confirmed by PCR amplification and Sanger sequencing of the editing site (Fig. 3c). As expected, the production of both pteridic acids and elaiophylin was abolished in *S. iranensis*/Δ*ptaA* (Fig. 3c). Plant experiments showed that the treatment of *S. iranensis*/Δ*ptaA* fermentation suspension led to the abolishment of the abiotic stress mitigating effects (Fig. 1a, 1b, 1c). To further confirm *pta* gene cluster, a bacterial artificial chromosome (BAC) library of *S. iranensis* was constructed. BAC-based cross-complementation of the *ptaA* in *S. iranensis*/Δ*ptaA* restored the production of pteridic acids and elaiophylin (Supplementary Fig. 28).

Interestingly, based on isotope-labelled precursor feeding and partial cosmid sequencing-based bioinformatics prediction, this BGC has long been inferred to be responsible for the biosynthesis of the antibacterial elaiophylin.[35,36] In 2015, Zhou et al. reported that thioesterase in the last module catalyse the formation of symmetrical macrodiolide using two units of linear elaiophylin monomeric seco acid (Fig. 3b).[37] To confirm whether the biosynthesis of pteridic acids is also thioesterase-dependent, site-specific mutations of residues Met-Glu-Asp to Ile-Lys-Met were introduced into the active sites of the TE domain *in vivo* (Supplementary Fig. S29).[38] The HR-LC-MS/MS analysis showed that the mutant strain (M2089I+E2090K+D2091M) no longer produced pteridic acids and elaiophylin (Fig. 3d). Hence, we provide additional evidence via in vivo inactivation and site-directed mutagenesis. Co-production of the plant growth-regulating pteridic acids and the antimicrobial elaiophylin through a shared BGC is intriguing and points to possible joint efforts in helping plants cope with biotic and abiotic stress.

Geographical distribution of pteridic acid producers. We surveyed available gene cluster family (GCF) data for all bacteria in the BiG-FAM database.[39] We found that *pta* BGC (GCF_02696) is strictly restricted to the *Streptomyces* genus. In addition, a total of 55 BGCs with high similarity to the *pta* BGC were detected by BiG-SCAPE,[40] among a total of 9,386 type I polyketides BGCs in 1,965 *Streptomyces* from the NCBI assembly database. Through literature supplementation and data dereplication of other reported producers without sequence information, at least 81 *Streptomyces* are known to produce pteridic acids/elaiophylin or have specific *pta* BGC up to date (Supplementary Tab. 4). Based on the known sampling information, the *pta*-containing *Streptomyces* display a variety of

geographic distribution and biological origins (Fig. 4). We selected two available *Streptomyces* strains (*Streptomyces violaceusniger* Tu 4113 and *Streptomyces rapamycinicus* NRRL 5491) to test the potential plant growth-promoting activity of these potential pteridic acid producers. The HR-LC-MS/MS analysis of both culture broths revealed that they shared similar metabolite profiles, and both produced pteridic acids H and F (Supplementary Fig. 30). Treatment with both culture broths on barley seedlings also exhibited significant plant growth-promoting activities under osmotic, salinity and drought stress (Fig. 4b, 4c and 4d). This evidence suggests that this class of *Streptomyces* and its specific secondary metabolite pteridic acids have unique ecological significance involved in plant abiotic stress resistance.

Phylogeny and evolution of *pta* BGC. To explore the evolutionary clues of pteridic acid producers, 16S rRNA genes were initially used to assess the relatedness of the collected 34 potential producers of pteridic acids with other *Streptomyces* strains that do not contain *pta* BGC. (Fig. 5a; Supplementary Fig. 31). The results revealed that, except for *Streptomyces albus* DSM 41398 and *Streptomyces* sp. GMR22, and other *pta*-containing *Streptomyces* strains cluster together and are distinct with divergent lineages. To further confirm this hypothesis, two high-resolution *Streptomyces* housekeeping genes, tryptophan synthase subunit beta (*trpB*) and RNA polymerase subunit beta (*rpoB*) were employed to analyse the phylogeny relationship among these strains.[41] Consequently, only *S. albus* DSM 41398 was classified in a distinct phylogenetic lineage among the *Streptomyces* strains containing *pta* BGC. The strict congruence among the clades of the housekeeping genes indicated dominant vertical transmission and potential horizontal gene transfer of the *pta* BGC in *Streptomyces*.

A total of 15 *pta*-containing *Streptomyces* with complete genome sequence information were selected to conduct the comparative genomics investigation. The genetic diversity in these *Streptomyces* strains was initially revealed using genome sequence similarity analysis. Except for *S. albus* DSM 41398 and *Streptomyces* sp. NA02950, we observed a high degree of similarity in the aligned region, as indicated by both the average nucleotide identity (ANI) and the alignment percentage (AP) among these strains (Supplementary Fig. 32). Genome synteny analysis revealed that partial genome rearrangements happened among strains even with high sequence similarities (Supplementary Fig. 33). Notably, the *pta* BGC in *S. albus* DSM 41398 (*pta-alb*) is located at the end of the chromosome, a high variable region in *Streptomyces*, suggesting its existence by accepting heterologous biosynthetic gene fragments. The nucleotide sequence alignment of *pta* BGC results showed that *pta-alb* is relatively complete, and the similarity of core genes is proportional to evolutionary relatedness (Fig. 5b). The metabolite profile of *S. albus* DSM 41398 also confirmed the integrity of the *pta-alb* by detecting the production of elaiophylin and pteridic acid H (Supplementary

Fig. 34). To further assess the biosynthesis diversity in remaining genetically related strains, we

performed a BGCs similarity analysis (Supplementary Fig. 35, 36). The connections between their

secondary metabolites, BGCs, revealed that these vertically inherited *Streptomyces* strains also harbour

striking similarities. Combining phylogenetic and comparative genomics analysis, we expect that *S.*

albus DSM 41398 is evolutionarily the most distinct member from other *pta*-containing *Streptomyces*

strains and obtained the *pta* BGC via horizontal gene transfer. However, most *pta*-containing

Streptomyces strains have vertically inherited *pta* and other gene clusters from their ancient ancestors

that may be ecologically important and rarely studied.

Discussion

Drought and salinisation of soil are increasing globally, driving a reduction in crop yields that threatens

food security. Plant growth-promoting bacteria is a class of beneficial microorganisms that positively

interact with the plant to confer environmental stresses.[42,43] Although some *Streptomyces* species have

been reported to have plant growth-promoting activity, the molecules mediating such positive effects

are poorly understood. Deciphering the molecular mechanism is key to understanding the complex

plant microbiota interaction. In this study, we present an example of *S. iranensis* secreting a family of

secondary metabolites, pteridic acids, to assist plants to cope with abiotic stresses like osmotic, salinity

and drought. Pteridic acids H and F were chemically isolated, structurally characterised and functional

validated as plant-beneficial molecules.

Plants respond to the harsh environment by changing their physiological processes for better

survival.[44] Salt and drought stress signal transduction consists of ionic and osmotic homeostasis

signaling pathways, detoxification (i.e., damage control and repair) response pathways, and pathways

for growth regulation.[45] Aquaporins are membrane channels that facilitate the uptake of soil water and

mediate the regulation of root hydraulic conductivity in response to a large variety of environmental

stresses. [46] TIPs represent one family of aquaporins, and they are involved in regulating water flow in

response to osmotic challenges like drought or salinity for a plant cell.[47] In our study, treatment of

pteridic acids H and F resulted in a 2.1-fold and 2-fold increase in TIP2;3 transcripts at 1-hour and a

2.1-fold and 2.9-fold increase at 24-hour, respectively (Fig. 2e). SOS1 is a plasma membrane Na^+/H^+

antiporter, which is important for plant to alleviate salinity stress.[48] In response to the ionic toxicity

triggered by salt stress, plants rely on the Salt Overly Sensitive (SOS) pathway to transport excessive

Na^+ from the cytoplasm to the apoplast, thus ensuring endogenous ionic homeostasis.[49] The

overexpression of SOS1 could significantly increase the salt tolerance of *Arabidopsis* plants.[50] We

noticed that the relative expression of SOS1 was significantly enhanced after the 24-hour treatment of

pteridic acids, which preliminary revealed the molecular basis that endows it with stress resistance.

However, the exact mode of action of pteridic acids remains to be further investigated by for example genome-wide transcriptomics screening for pteridic acids insensitive mutant plants.

Horizontal gene transfer is an integral driver of BGC evolution, revealing the independent processes of species phylogeny and BGCs distribution.[51] However, vertical inheritance also influences BGCs evolutionary dynamics, evident from BGCs conservation among closely related strains.[52] We found that the *pta* BGC in *Streptomyces* are widely dispersed geographically and mainly inherited through vertical gene transmission. Some of these strains have also been described to have remarkable biocontrol capabilities. For example, *Streptomyces* sp. AgN23 activates *Arabidopsis* defence responses to fungal pathogen infection by secreting plant elicitors,[53] *Streptomyces rhizosphaericus* 0250 and *Streptomyces* sp. 5-10 displayed significant biocontrol potential to fusarium wilt of bitter gourd.[54,55] The family of *pta*-containing *Streptomyces* was also previously described as a specific phylogenetic lineage with the highest BGC abundance and largest genome size across diverse *Streptomyces-type* strains.[56] Although the biosynthetic diversity of these *Streptomyces* is likely due to horizontal transfer events that occurred relatively recently in their evolutionary history instead of genetic diversification through a vertical transfer of BGCs. The multiple Type I PKSs presenting among these strains are highly conserved based on genetic similarity network analysis. There is currently some evidence supporting potential complex cross-BGC regulation in this class of *Streptomyces* strains. Jiang et al. demonstrated that a TetR family transcriptional regulator, GdmRIII, controls the biosynthesis of geldanamycin and elaiophylin meanwhile, in *Streptomyces autolyticus* CGMCC 0516.[57] Recently, He et al. found that the rapamycin BGC-situated LAL family regulator RapH co-ordinately regulated the biosynthesis of both rapamycin and elaiophylin in *S. rapamycinicus* NRRL 5491.[58] Although these reports correspond to cross-regulation between evolutionarily conserved BGCs, more details of these communications need to be investigated.

In conclusion, pteridic acids are secondary metabolites produced by *Streptomyces* enhancing plant resistance to abiotic stress. This is a useful illustration of the bacterial metabolite-mediated alteration of plants in response to environmental stress. It will open avenues for utilising *Streptomyces* to rewild plant microbiomes and improve plant abiotic stress resistance to tackle climate change.[59]

Methods

Strains, plasmids, and cultivation. All strains and plasmids used in this study were listed in Supplementary Tab. 5. All *Streptomyces* strains were obtained from the German Collection of Microorganisms and Cell Cultures GmbH (DSMZ, Germany). All *Escherichia coli* strains were grown in liquid/solid LB medium (5.0 g L^{-1} yeast extract, 10.0 g L^{-1} peptone, 10.0 g L^{-1} NaCl) at 37°C. All *Streptomyces* strains were grown on SFM medium (20.0 g L^{-1} mannitol, 20.0 g L^{-1} soya flour, 20.0 g

L^{-1} agar), and the SFM medium with the addition of 120 mM calcium chloride solution was used for

the step of conjugation at 28°C. The ISP2 medium (4.0 g L^{-1} yeast extract, 10.0 g L^{-1} malt extract, 4.0

g L^{-1} dextrose, and 1.0 L distilled water) was used for liquid fermentation of all *Streptomyces* strains

used in plant assay and metabolomics analysis. Appropriate antibiotics were supplemented with the

following working concentrations: apramycin (50 µg mL^{-1}), chloramphenicol (25 µg mL^{-1}), and

kanamycin (50 µg mL^{-1}). All chemicals utilized in this study were from Sigma-Aldrich, USA.

Metabolomics analyses. High performance liquid chromatography was carried out on the Agilent

Infinity 1290 UHPLC system (Agilent Technologies, USA). The 250 × 2.1 mm i.d., 2.7 µm, Poroshell

120 Phenyl Hexyl column (Agilent Technologies, USA) was used for separation. The 2-µL samples

were eluted at a flow rate of 0.35 mL min^{-1} using a linear gradient from 10% acetonitrile in Milli-Q

water buffered with 20 mM formic acid increasing to 100% in 15 min. Each starting condition was

held for 3 min before the next run. Mass spectrometry detection was performed on an Agilent 6545

QTOF MS equipped with Agilent Dual Jet Stream electrospray ion source (ESI) with a drying gas

temperature of 160°C, a gas flow of 13 L min^{-1}, sheath gas temperature of 300°C, and flow of 16 L

min^{-1}. The capillary voltage was set to 4000 V and the nozzle voltage to 500 V in positive mode. MS

spectra were recorded as centroid data at an m/z of 100-1700, and auto MS/HRMS fragmentation was

performed at three collision energies (10, 20, and 40 eV) on the three most intense precursor peaks per

program. Data were analysed with MassHunter software (Agilent Technologies, USA) and compared

with known compounds and crude extract spectral libraries stored in the GNPS platform.[45] The

precursor and fragment ion mass tolerance were set as 0.1 Da and 0.02 Da, respectively. In addition,

the minpairs cos was set as 0.65, and the minimum matched fragment ions were set as 6.0. The

metabolites profile of wild-type *S. iranensis* was visualised by MS-Dial 4.9.2.[60]

Large-scale fermentation and isolation. *S. iranensis* was cultivated in medium 2 (3.0 g L^{-1}

$CaCl_2 \cdot 2H_2O$, 1.0 g L^{-1} citric acid/Fe III, 0.2 g L^{-1} $MnSO_4 \cdot H_2O$, 0.1 g L^{-1} $ZnCl_2$, 0.025 g L^{-1}

$CuSO_4 \cdot 5H_2O$, 0.02 g L^{-1} $Na_2B_4O_2 \cdot 10H_2O$, 0.01 g L^{-1} $NaMoO_4 \cdot 2H_2O$, and 20.0 g L^{-1} oatmeal in 1.0 L

distilled water), at 175 L filling volume in a 300 L fermentation vessel (Sartorius, Germany). The

fermentation was carried out for 6 days with aeration of 25-50 L min^{-1}, stirring at 200 rpm with a

temperature of 28°C and at a pH range of 5.4-6.4. The fermentation broth was separated, filtered, and

loaded onto an Amberchrom CG161Me resin LC column (200 × 20 cm, 6 L). Elution with a linear

gradient of H_2O-MeOH (from 30% to 100% v/v, flow rate 0.5 L min^{-1}, in 58 min) afforded seven

fractions (A-G). Fraction G was firstly fractionated by silica gel chromatography with a

CH_2Cl_2/CH_3OH gradient to yield 16 fractions, F01-F16. F07 was separated by a Sephadex LH-20

(MeOH) column and twelve sub-fractions F07a-l were obtained. From F07e, **1** (15.0 mg) and **2** (4.0

mg) were obtained by repeated HPLC RP-C_{18} (CH_3CN/H_2O as gradient).

Pteridic acid H (**1**): white solid; $[\alpha]_D^{20}$ 81 (0.32 mg/mL, CH_3OH), ^1H NMR (800 MHz, MeOD):

7.33 (dd, 15.4 Hz, 11.1 Hz, 1H), 6.26 (dd, 15.2 Hz, 10.8 Hz, 1H), 6.13 (dd, 15.3 Hz, 8.8 Hz, 1H), 5.90

(d, 15.4 Hz, 1H), 3.85 (dd, 10.2 Hz, 2.2 Hz, 1H), 3.69 (dd, 11.4 Hz, 4.9 Hz, 1H), 3.59 (m, 1H), 3.43

(m, 1H), 2.48 (m, 1H), 2.29 (dd, 14.9 Hz, 6.1 Hz, 1H), 2.01 (m, 1H), 1.64 (dd, 14.9 Hz, 1.9 Hz, 1H),

1.52 (m, 1H), 1.49 (m), 1.21 (d, 6.1 Hz, 3H), 1.01 (d, 6.8 Hz, 3H), 0.96 (d, 6.8 Hz, 3H), 0.93 (t, 7.3

Hz, 3H), 0.91 (d, 7.0 Hz, 3H); ^{13}CNMR (200 MHz, MeOD): 168.7, 151.0, 146.7, 129.6, 120.6, 103.2,

75.5, 72.8, 72.7, 70.3, 51.3, 42.2, 40.4, 37.5, 37.4, 24.8, 20.9, 16.0, 12.1, 11.1, 5.0; UV/vis

(CH_3CN/H_2O) λ_{max} 262 nm; IR (ATR) ν_{max} 2967, 2934, 2879, 1712, 1642, 1600, 1458, 1410, 1383,

1300, 1266, 1223, 1187, 1142, 1109, 1058, 1002, 973 cm^{-1}; (+)-HR-ESI-MS (*m/z*) [M+H]$^+$ calcd for

$C_{21}H_{35}O_6$, 383.2428; found, 383.2439. ^1H NMR and ^{13}C NMR see Supplementary Tab. 6.

Pteridic acid F (**2**): white solid; $[\alpha]_D^{20}$ -18 (10 mg/mL, CH_3OH), ^1H NMR (800 MHz, MeOD): 7.16

(dd, 15.1 Hz, 10.9 Hz, 1H), 6.25 (dd, 15.1 Hz, 10.9 Hz, 1H), 6.07 (dd, 15.1 Hz, 8.6 Hz, 1H), 5.97 (d,

15.1 Hz, 1H), 3.88 (m, 1H), 3.66 (td, 10.8 Hz, 4.3 Hz, 1H), 3.56 (dd, 11.5 Hz, 4.7 Hz, 1H), 3.32 (m,

1H), 2.49 (m, 1H), 2.19 (dd, 13.1 Hz, 4.3 Hz, 1H), 2.02 (m, 1H), 1.69 (m, 1H), 1.60 (m, 1H), 1.44 (m,

1H), 1.32 (dd, 13.2 Hz, 11.2 Hz, 1H), 1.14 (d, 6.2 Hz, 3H), 1.02 (m, 1H), 1.02 (d, 6.8 Hz, 3H), 0.95

(d, 6.9 Hz, 3H), 0.98 (d, 6.8 Hz, 3H), 1.60 (m, 1H), 1.44 (m, 1H), 0.82 (t, 7.6 Hz, 3H); ^{13}CNMR (200

MHz, MeOD): 170.2, 148.1, 143.4, 129,6, 122.9, 103.2, 77.7, 74.7, 69.8, 66.3, 52.2, 42.1, 40.5, 37.7,

33.8, 20.4, 19.7, 15.9, 12.7, 10.5, 5.3; UV/vis (CH_3CN/H_2O) λ_{max} 264 nm; IR (ATR) ν_{max} 2968, 2931,

2877, 1692, 1643, 1618, 1458, 1410, 1380, 1299, 1270, 1188, 1138, 1106, 1059, 1002, 973, 850 cm^{-1};

(+)-HR-ESI-MS (*m/z*): [M+H]$^+$ (calcd for $C_{21}H_{35}O_6$, 383.2428; found, 383.2433. ^1H NMR and ^{13}C

NMR see Supplementary Tab. 7.

Structure identification. NMR spectra were recorded on an 800 MHz Bruker Avance III spectrometer

equipped with a TCI CryoProbe using standard pulse sequences. NMR data were processed using

MestReNova 11.0. UHPLC-HRMS was performed on an Agilent Infinity 1290 UHPLC system

equipped with a diode array detector. UV-Vis spectra were recorded from 190 to 640 nm. Specific

rotations were acquired using Perkin-Elmer 241 polarimeter. IR data were acquired on Bruker Alpha

FTIR spectrometer using OPUS version 7.2. TLC analysis was performed on silica gel plates (Sil

G/UV$_{254}$, 0.20 mm, Macherey-Nagel). The Biotage Isolera One Flash Chromatography system was

used for flash chromatography and performed on silica gel 60 (Merck, 0.04–0.063 mm, 230–400 mesh

ASTM). Sephadex LH-20 was from Pharmacia.

Crystal Structure Determination. X-ray data collection of **1** was performed on an Agilent Supernova Diffractometer using CuKα radiation. Data were processed and scaled using the CrysAlisPro software (Agilent Technologies, USA). The structure was solved using SHELXS and refined using SHELXL. Hydrogen atoms were included in ideal positions using riding coordinates. The absolute configuration was determined based on the Flack parameter. Crystal Data for **1**: $C_{21}H_{34}O_6$, Mr = 382.50, monoclinic, a = 8.4619(1) Å, b = 15.6161(2) Å, c = 8.4994(1) Å, α = 90.00°, β = 107.768(1)°, γ = 90.00°, V = 1069.55(2) Å3, T = 120(2) K, space group P21, Z = 2, μ(Cu Kα) = 0.698 mm–1, 17514 reflections collected, 4275 independent reflections (Rint = 0.0226, Rsigma = 0.0155). The final R1 values were 0.0249 (I > 2σ(I)). The final wR2 values were 0.0648 (I > 2σ(I)). The final R1 values were 0.0252 (all data). The final wR2 values were 0.0651 (all data). The goodness of fit on F2 was 1.057. The Flack parameter is 0.13(10).

Genetic manipulation. All primers used were synthesised by IDT (Integrated DNA Technologies, USA) and listed in Supplementary Tab. 8. Plasmids and genomic DNA purification, PCR and cloning were conducted according to standard procedures using manufacturer protocols. PCR was performed using OneTaq Quick-Load 2X Master Mix with Standard Buffer (New England Biolabs, USA). DNA assembly was done by using NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs, USA). DNA digestion was performed with FastDigest restriction enzymes (Thermo Fisher Scientific, USA). NucleoSpin Gel and PCR Clean-up Kits (Macherey-Nagel, Germany) were used for DNA clean-up from PCR products and agarose gel extracts. One Shot Mach1 T1 Phage-Resistant Chemically Competent *E. coli* (Thermo Fisher Scientific, USA) was used for cloning. NucleoSpin Plasmid EasyPure Kit (Macherey-Nagel, Germany) was used for plasmid preparation. Sanger sequencing was carried out using a Mix2Seq Kit (Eurofins Scientific, Luxembourg). All DNA manipulation experiments were conducted according to standard procedures using manufacturer protocols.

Gene inactivation and site-directed mutagenesis. To use the pCRISPR-cBEST for base editing applications, an oligo was designed as Del-ptaA by the online tool CRISPy-web, and the pCRISPR-cBEST plasmid was linearised by *Nco*I. Mixing the linearised pCRISPR-cBEST plasmid and Del-ptaA with the NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs, USA). The linearised pCRISPR-cBEST plasmid was then bridged by Del-ptaA, ending with the desired pCRISPR-cBEST/Δ*ptaA*. Chemically competent *E. coli* were transformed with the recombinant plasmid and confirmed via PCR amplification (program: 94°C for 30 s, followed by 30 cycles consisting of 94°C for 15 s; 54°C for 15 s; 68°C for 40 s, and 68°C for 2 min) and Sanger sequencing. The experimental

procedure for site-directed mutagenesis for the TE domain is the same as described above. The *E. coli-*

Streptomyces conjugation experiment was conducted according to the modified protocol in this study,

and the mutant *Streptomyces* strains were also confirmed by PCR and Sanger sequencing (Eurofins,

France).

Construction of BAC and genetical complementation. The BAC library of *S. iranensis* was

constructed using pESAC13-A from Bio S&T (Montreal, Canada). Based on a high-throughput

screening method (unpublished), we selected two BACs 1J23 and 6M10 that cross-cover the *ptaA* gene

(Supplementary Fig. 28). The selected BAC clones were further confirmed using four sets of primers,

including ID-1J23-right-F/R, ID-1J23-left-F/R, ID-6M10-right-F/R, and ID-6M10-left-F/R

(Supplementary Tab. 8). Subsequently, we introduced these two BAC clones 1J23 and 6M10 into *S.*

iranensis/Δ*ptaA* (remove the pCRISPR-cBEST/Δ*ptaA* to obtain antibiotics resistance free strain)

separately by conjugation. Exconjugants of mutants were further validated by apramycin-resistance

screening and PCR.

Enrichment evaluation of *S. iranensis* in rhizosphere soil. The *S. iranensis* (with the apramycin

resistance gene) spore suspension was well mixed with fully sterilised soil and was transferred to a

250 mL flask. The sterilised germinated barley seed was placed in the centre position of soil in flask

and grown at $24 \pm 2°C$, 8 h dark/16 h light in the growth chamber for 7 days. Samples were collected

from soil within 0.1 cm and 3 cm distance from the barley root, with a specification of 0.1 g soil per

sample. Then, these samples were transferred to sterilised 1.5 mL Eppendorf tubes and mixed with

500 μL sterilised H_2O. 200 μL of each sample was Spread evenly over the solid MS medium with the

addition of 50 μg mL^{-1} apramycin and grown at 28°C for 7 days. The number of *Streptomyces* colonies

grown on each plate were counted and statistically analysed.

Arabidopsis **growth assays.** *A. thaliana* ecotype Columbia (Col-0) was used to test the effects of

pteridic acid treatment under drought stress mediated by PEG-6000 (Duchefa Biochemie BV) and

salinity stress mediated by NaCl (Duchefa Biochemie BV). The modified Murashige & Skoog medium

(2.2 g L^{-1} Murashige & Skoog medium including B5 vitamins, 5.0 g L^{-1} of sucrose, 250 mg L^{-1} MES

monohydrate, 7.0 g L^{-1} agar, and adjusted pH to 5.7 with KOH) was used in this study. PEG-6000 (15%

w/v) was dissolved in water and filtered through 0.2-micron Sartorius Minisart™ Plus Syringe Filters

(Fisher Scientific). 50 mL of the filtered solution was overlaid onto the surface of solidified Murashige

& Skoog medium. The plates were left for 24 h to diffuse the PEG into the Murashige & Skoog medium.

NaCl was added to the medium to a final concentration of 80 mM for the salinity stress alleviation test.

The pure compound pteridic acids H and F (0.50 ng/mL), IAA (0.23 ng/mL), and ABA (0.34 ng/mL)

were mixed with different media to a final concentration of approximately 1.3 nM and poured into the

plates. Seeds were surface sterilised by washing with 70% ethanol for 2 min, then in sterilisation

solution (10% bleach) for 1 min by inverting the tubes, and finally washed five times with sterilised

water. The seeds were stratified for 2 days at 4°C in the dark. Sterilised seeds were placed in Petri

dishes (approx. 100 seeds per Petri dish) on Murashige & Skoog medium and grown for 3-4 days in

the vertical position in a culture chamber at 22°C under standard long-day conditions (16/8 h light/dark

photoperiod). After three days of growth, seedlings with similar root lengths (7-10 mm) were

transferred to square plates containing Murashige & Skoog medium (control) or Murashige & Skoog

medium supplemented with 15% (w/v) PEG-6000 or 80 mM NaCl. 16 seedlings were used per

replicate for each treatment. The initial position of the plant root tip was marked with a marker. The

plants were grown in the vertical position under standard long day conditions (22°C, 16/8h light/dark)

for 8 days, and then each plate was scanned using Image Scanner. Primary and lateral root lengths and

the total plant weight were then scored. The primary and lateral root length measurements were

performed by analysing pictures with the Image J software. Fresh weight measurements were

estimated using a precision balance. Whole 8-day-old plants grown in the medium were removed using

forceps, dried in tissue paper, and then weighed using a precision balance.

Barley and mung bean growth assays. The pteridic acid *Streptomyces* producers were tested for their

effects on barley cultivars Guld grown in soil. *S. iranensis* HM 35, *S. rapamycinicus* NRRL 5491, and

S. violaceusniger Tu 4113 were cultivated in ISP2 medium for 7 days at 28°C. 500 µL culture broth

(ca. 3 x 10^9 CFU/mL) was added to 100 g sand soil and well mixed. Treated soils were infiltrated with

Milli-Q water, 20% PEG-6000 solutions to simulate osmotic stress, and 100 mM NaCl solutions to

simulate salinity stress in soil environments. To simulate drought stress during barley growth, the plant

was initially watered for 7 days, then subjected to 7 days of drought stress, and finally allowed to

recover with various treatments for another 7 days. Barley seeds were rinsed in distilled water and

sterilised with 1% sodium hypochlorite for 15 min. They were then washed with distilled water and

germinated in distilled water at 24°C for 2 days. Barley seeds were planted in each plastic pot (5cm x

5cm x 6cm, six seedlings per pot) supplemented with different treated soil and grown at 24 ± 2°C, 8 h

dark/16 h light in the growth chamber for 7 days. Plant height (cm) was measured as the aerial part of

the plant, and the fresh shoot weight (g) and fresh root weight (g) of each seedling were measured

separately. Then, the seedlings were dried in the hot-air oven at 70°C for 6 h to obtain the dry shoot

and dry root weights (g). For heavy metal stress experiment, mung beans were pre-germinated and

placed on top of the modified Murashige & Skoog medium agar, supplemented with 10 mM CuSO4,

with 1.0 ng/mL pure substances. All mung beans were grown in the dark at 24 ± 2°C for 4 days.

Kidney beans growth assays. The seeds of kidney beans (appr. 2 cm in length, from organic farming) were firstly sterilised successively with ethanol (70% v/v) and sodium hypochlorite (5% v/v), each for 2 min and then rinsed with sterile Milli-Q water (three times). The sterilised seeds were cultivated on modified Murashige & Skoog agar plates for 3-4 days. After germination, the seedlings (with 1.5-2 cm-long roots) were soaked in 10 mL aliquots of testing compounds (pteridic acids H and F, 1.0 ng/mL, dissolved in sterile Milli-Q water) in ultra-clear polypropylene containers (ø 34 mm, vol. 20 mL) with polyethylene caps. The control group was treated with 10 mL sterile Milli-Q water. For each treatment, three repetitions (containers) were used, and each repetition included four seedlings. After 24 h, the seeds were transferred into a cut square petri dish, put on the top layer of sandy soil, and then incubated vertically in a growth chamber (24/22°C, day/night cycle of 16/8 h, 50%, 60%, 70%, 100% of circulated wind velocity for 12 h, 2 h, 2h, 8h) for 7 days. Solutions of pteridic acids H and F (2 mL, 1.0 ng/mL for both) were added separately into corresponding containers, with sterile Milli-Q water as control, and an extra 8 mL of Milli-Q water was added to each petri dish every other day.

RNA extraction and quantitative real-time PCR. *Arabidopsis* seeds were sterilized, and seven to ten seeds were cultured in each well of a 24 well plate in 2 mL medium containing modified Murashige & Skoog and 30 mM sucrose with 16 hours light, at 24 °C in a controlled environment room. After 7 days, seedlings were washed in 20 mL modified Murashige & Skoog medium and then moved to 20 mL of fresh medium and cultured in 125ml E-flasks with 1 ng mL^{-1} of each pteridic acids H and F. Samples were collected at three Total RNA was extracted using RNeasy Plant Mini Kit (Qiagen, Germany) according to the manufacturer's instructions. Reverse transcription was performed using SuperScript™ III First-Strand Synthesis SuperMix for qRT-PCR (Invitrogen, USA). qRT-PCR was conducted on CFX96 Touch Real-Time PCR Detection System (Bio-Rad, USA) using Platinum SYBR Green qPCR SuperMix-UDG with ROX (Invitrogen, USA). The thermal cycling program consisted of an initial denaturation at 95 °C for 3 min, followed by 39 cycles of 95 °C for 15 s, 60 °C for 30 s. Three independent biological replicates were performed for each condition. TIP2;3 and SOS1 were used as target genes, and primers sequences were given in Supplementary Tab. 8. The expression level of target gene after treatments was calculated using the $2^{\wedge}\Delta\Delta Ct$ method, with ACT1 as the internal control.

Bioinformatics analyses. The identification and annotation of all BGCs of *Streptomyces* secondary metabolites were carried out with antiSMASH 6.0.[26] The threshold for similar BGCs was selected as greater than 45% sequence similarity. The BiG-FAM database and BiG-SCAPE software were used to identify the distribution of *pta* BGC in different bacteria and generate the *pta* family similarity

network.[39,40] A cut-off of 0.3 was used as a raw index similarity metric for the BiG-SCAPE analysis.

The alignment of 15 *pta*-containing *Streptomyces* genome sequences was performed using the whole-

genome alignment plugin of the CLC Genomics Workbench version 22.0.2 (Qiagen). The minimum

initial seed length was 15 bp, and the minimum alignment block length was 100 bp. The networks of

Big-SCAPE and GNPS analysis were visualised using Cytoscape 3.9. The phylogenetic analysis was

conducted by the online multiple sequence alignment tool MAFFT and visualised by iTOL v5. [61,62]

Statistical analysis. Statistical significance was assessed by one-way ANOVA with post hoc

Dunnett's multiple comparisons test, one-way ANOVA with Tukey test or t test (see each figure

legends). All analyses were performed using GraphPad Prism version 9. P-values < 0.05 were

considered significant. Asterisks indicate the level of statistical significance: $*P < 0.05$, $**P < 0.01$,

$***P < 0.001$, and $****P < 0.0001$. For all relevant figures, source data and exact P values are provided

in the Source Data file.

Data availability. The gene sequences used in this study were collected by searching National Center

for Biotechnology Information (NCBI) or extracting from assembled genome sequences (attached in

Supplementary information files). All genome sequences used in this study are publicly available on

NCBI. The crystal structure was deposited at Cambridge Crystallographic Data Centre

(https://www.ccdc.cam.ac.uk/) with CCDC deposition number 1984025. Metabolomics data in this

study were deposited at https://massive.ucsd.edu with the identifier MSV000090745. Source data of

each figure are provided with this paper.

References

1. Naylor, R., et al. A 20-year retrospective review of global aquaculture. *Nature* **591**, 551-563 (2021).

2. Wheeler, T. & von Braun, J. Climate change impacts on global food security. *Science* **341**, 508-513

(2013).

3. Zhang, H., Zhu, J., Gong, Z. & Zhu, J. Abiotic stress responses in plants. *Nat. Rev. Genet.* **23**, 104-

119 (2022).

4. Kuromori, T., Fujita, M., Takahashi, F., Yamaguchi-Shinozaki, K. & Shinozaki, K. Inter-tissue and

inter-organ signaling in drought stress response and phenotyping of drought tolerance. *Plant J.* **109**,

342-358 (2022).

5. Dos Santos, V., et al. Causes of reduced leaf-level photosynthesis during strong El Niño drought in

a central Amazon forest. *Global Change Biol.* **24**, 4266-4279 (2018).

6. Waadt, R., et al. Plant hormone regulation of abiotic stress responses. *Nat. Rev. Mol. Cell Biol.* **23**,

680-694 (2022).

7. Kerchev, P. & Van Breusegem, F. Improving oxidative stress resilience in plants. *Plant J.* **109**, 359-

372 (2022).

8. Abid, M., et al. Physiological and biochemical changes during drought and recovery periods at

tillering and jointing stages in wheat (*Triticum aestivum* L.). *Sci. Rep.* **8**, 1-15 (2018).

9. Hasanuzzaman, M., et al. Regulation of ROS metabolism in plants under environmental stress: a

review of recent experimental evidence. *Int. J. Mol. Sci.* **21**, (2020).

10. Van Zelm, E., Zhang, Y. & Testerink, C. Salt tolerance mechanisms of plants. *Annu. Rev. Plant*

Biol. **71**, 403-433 (2020).

11. Corwin, D. Climate change impacts on soil salinity in agricultural areas. *Eur. J. Soil Sci.* **72**, 842-

862 (2021).

12. Hong, Y., Zhou, Q., Hao, Y. & Huang, A. C. Crafting the plant root metabolome for improved

microbe-assisted stress resilience. *New Phytol.* **234**, 1945-1950 (2022).

13. De Vries, F., Griffiths, R., Knight, C., Nicolitch, O. & Williams, A. Harnessing rhizosphere

microbiomes for drought-resilient crop production. *Science* **368**, 270-274 (2020).

14. Arif, I., Batool, M. & Schenk, P. Plant microbiome engineering: expected benefits for improved

crop growth and resilience. *Trends Biotechnol.* **38**, 1385-1396 (2020).

15. Berlanga-Clavero, M., et al. *Bacillus subtilis* biofilm matrix components target seed oil bodies to

promote growth and anti-fungal resistance in melon. *Nat. Microbiol.* **7**, 1-15 (2022).

16. Das, P., et al. Plant-soil-microbes: A tripartite interaction for nutrient acquisition and better plant

growth for sustainable agricultural practices. *Environ. Res.* **214**, 113821 (2022).

17. Hiruma, K., et al. Root endophyte *Colletotrichum tofieldiae* confers plant fitness benefits that are

phosphate status dependent. *Cell* **165**, 464-474 (2016).

18. Henneron, L., Kardol, P., Wardle, D., Cros, C. & Fontaine, S. Rhizosphere control of soil nitrogen

cycling: a key component of plant economic strategies. *New Phytol.* **228**, 1269-1282 (2020).

19. Etalo, D., Jeon, J. & Raaijmakers, J. Modulation of plant chemistry by beneficial root microbiota.

Nat. Prod. Rep. **35**, 398-409 (2018).

20. Quinn, G., Banat, A., Abdelhameed, A. & Banat, I. *Streptomyces* from traditional medicine: sources

of new innovations in antibiotic discovery. *J. Med. Microbiol.* **69**, 1040-1048 (2020).

21. Viaene, T., Langendries, S., Beirinckx, S., Maes, M. & Goormachtig, S. *Streptomyces* as a plant's

best friend? *FEMS Microbiol. Ecol.* **92**, (2016).

22. Myo, E., et al. Indole-3-acetic acid production by *Streptomyces fradiae* NKZ-259 and its

formulation to enhance plant growth. *BMC Microbiol.* **19**, 1-14 (2019).

23. Sadeghi, A., et al. Plant growth promoting activity of an auxin and siderophore producing isolate

of *Streptomyces* under saline soil conditions. *World J. Microbiol. Biotechnol.* **28**, 1503-1509 (2012).

24. Singh, S. & Gaur, R. Endophytic *Streptomyces* spp. underscore induction of defense regulatory

genes and confers resistance against *Sclerotium rolfsii* in chickpea. *Biol. Control* **104**, 44-56 (2017).

25. Fitzpatrick, C., et al. Assembly and ecological function of the root microbiome across angiosperm

plant species. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E1157-E1165 (2018).

26. Blin, K., et al. antiSMASH 6.0: improving cluster detection and comparison capabilities. *Nucleic*

Acids Res. **49**, W29-W35 (2021).

27. Wang, M., et al. Sharing and community curation of mass spectrometry data with Global Natural

Products Social Molecular Networking. *Nat. Biotechnol.* **34**, 828-837 (2016).

28. Igarashi, Y., Iida, T., Yoshida, R. & Furumai, T. Pteridic acids A and B, novel plant growth

promoters with auxin-like activity from *Streptomyces hygroscopicus* TP-A0451. *J. Antibiot.* **55**, 764-

767 (2002).

29. Nong, X., Wei, X. & Qi, S. Pteridic acids C-G spirocyclic polyketides from the marine-derived

Streptomyces sp. SCSGAA 0027. *J. Antibiot.* **70**, 1047-1052 (2017).

30. Lavenus, J., et al. Lateral root development in *Arabidopsis*: fifty shades of auxin. *Trends Plant Sci.*

18, 455-463 (2013).

31. Kautsar, S., et al. MIBiG 2.0: a repository for biosynthetic gene clusters of known function. *Nucleic*

Acids Res. **48**, D454-D458 (2020).

32. Sun, P., et al. Spiroketal formation and modification in avermectin biosynthesis involves a dual

activity of AveC. *J. Am. Chem. Soc.* **135**, 1540-1548 (2013).

33. Tong, Y., et al. Highly efficient DSB-free base editing for streptomycetes with CRISPR-BEST.

Proc. Natl. Acad. Sci. U. S. A. **116**, 20366-20375 (2019).

34. Netzker, T., et al. An efficient method to generate gene deletion mutants of the rapamycin-

producing bacterium *Streptomyces iranensis* HM 35. *Appl. Environ. Microbiol.* **82**, 3481-3492 (2016).

35. Gerlitz, M., Hammann, P., Thiericke, R. & Rohr, J. The biogenetic origin of the carbon skeleton

and the oxygen atoms of elaiophylin, a symmetric macrodiolide antibiotic. *J. Org. Chem.* **57**, 4030-

4033 (1992).

36. Haydock, S., Mironenko, T., Ghoorahoo, H. & Leadlay, P. The putative elaiophylin biosynthetic

gene cluster in *Streptomyces* sp. DSM4137 is adjacent to genes encoding adenosylcobalamin-

dependent methylmalonyl CoA mutase and to genes for synthesis of cobalamin. *J. Biotechnol.* **113**,

55-68 (2004).

37. Zhou, Y., Prediger, P., Dias, L., Murphy, A. & Leadlay, P. Macrodiolide formation by the

thioesterase of a modular polyketide synthase. *Angew. Chem. Int. Ed.* **54**, 5232-5235 (2015).

38. Du, L. & Lou, L. PKS and NRPS release mechanisms. *Nat. Prod. Rep.* **27**, 255-278 (2009).

39. Kautsar, S., Blin, K., Shaw, S., Weber, T. & Medema, M. BiG-FAM: the biosynthetic gene cluster

families database. *Nucleic Acids Res.* **49**, D490-D497 (2021).

40. Navarro-Munoz, J., et al. A computational framework to explore large-scale biosynthetic diversity. *Nat. Chem. Biol.* **16**, 60-68 (2020).

41. Komaki, H. Resolution of housekeeping gene sequences used in MLSA for the genus *Streptomyces* and reclassification of *Streptomyces anthocyanicus* and *Streptomyces tricolor* as heterotypic synonyms of *Streptomyces violaceoruber*. *Int. J. Syst. Evol. Microbiol.* **72**, 005370 (2022).

42. Coban, O., De Deyn, G. & van der Ploeg, M. Soil microbiota as game-changers in restoration of degraded lands. *Science* **375**, eabe0725 (2022).

43. De Souza, R., Ambrosini, A. & Passaglia, L. M. P. Plant growth-promoting bacteria as inoculants in agricultural soils. *Genet. Mol. Biol.* **38**, 401-419 (2015).

44. Lopes, M., Dias, M. & Gurgel, E. Successful plant growth-promoting microbes: inoculation methods and abiotic factors. *Front. sustain. food syst.* **5**, 606454 (2021).

45. Zhu, J. Salt and drought stress signal transduction in plants. *Annu. Rev. Plant Biol.* **53**, 247-273 (2002).

46. Maurel, C., et al. Aquaporins in plants. *Physiol. Rev.* **95**, 1321-1358 (2015)

47. Kaldenhoff, R. & Fischer, M. Functional aquaporin diversity in plants. *Biochim. Biophys. Acta Biomembr.* **1758**, 1134-1141 (2006).

48. Shi, H., Ishitani, M., Kim, C. & Zhu, J. The *Arabidopsis thaliana* salt tolerance gene SOS1 encodes a putative Na^+/H^+ antiporter. *Proc. Natl. Acad. Sci. U. S. A.* **97**, 6896-6901 (2000).

49. Ji, H., et al. The Salt Overly Sensitive (SOS) pathway: established and emerging roles. *Mol. Plant* **6**, 275-286 (2013).

50. Shi, H., Lee, B., Wu, S. & Zhu, J. Overexpression of a plasma membrane Na^+/H^+ antiporter gene improves salt tolerance in *Arabidopsis thaliana*. *Nat. Biotechnol.* **21**, 81-85 (2003).

51. McDonald, B. & Currie, C. Lateral gene transfer dynamics in the ancient bacterial genus *Streptomyces*. *Mbio* **8**, e00644-17 (2017).

52. Chase, A., Sweeney, D., Muskat, M., Guillen-Matus, D. & Jensen, P. Vertical inheritance facilitates interspecies diversification in biosynthetic gene clusters and specialized metabolites. *Mbio* **12**, e02700-21 (2021).

53. Vergnes, S., et al. Phyllosphere colonization by a soil *Streptomyces* sp. promotes plant defense responses against fungal infection. *Mol. Plant-Microbe Interact.* **33**, 223-234 (2020).

54. Li, X., Tian, Y., Peng, H., He, B. & Gao, K. Isolation, screening and identification of an antagonistic actinomycetes to control *Fusarium* wilt of *Momordica charantia*. *Ying Yong Sheng Tai Xue Bao* **31**, 3869-3879 (2020).

55. Yun, T., et al. Anti-Foc RT4 activity of a newly isolated *Streptomyces* sp. 5-10 from a medicinal

plant (*Curculigo capitulate*). *Front. Microbiol.* **11**, 610698 (2021).

56. Chung, Y., et al. Comparative genomics reveals a remarkable biosynthetic potential of the

Streptomyces phylogenetic lineage associated with rugose-ornamented spores. *Msystems* **6**, e00489-21

(2021).

57. Jiang, M., et al. GdmRIII, a TetR family transcriptional regulator, controls geldanamycin and

elaiophylin biosynthesis in *Streptomyces autolyticus* CGMCC0516. *Sci. Rep.* **7**, 4803 (2017).

58. He, W., et al. Crossregulation of rapamycin and elaiophylin biosynthesis by RapH in *Streptomyces*

rapamycinicus. *Appl. Microbiol. Biotechnol.* **106**, 2147-2159 (2022).

59. Raaijmakers, J. & Kiers, E. Rewilding plant microbiomes. *Science* **378**, 599-600 (2022).

60. Tsugawa, H., et al. MS-DIAL: data-independent MS/MS deconvolution for comprehensive

metabolome analysis. *Nat. Methods* **12**, 523-526 (2015).

61. Madeira, F., et al. Search and sequence analysis tools services from EMBL-EBI in 2022. *Nucleic*

Acids Res. **50**, W276-W279 (2022).

62. Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. *Nucleic Acids Res.* **49**, W293-W296 (2021).

Acknowledgements

We acknowledge Dr. Yaojun Tong for the discussions on CRISPR-BEST. We thank the support from

Dr. Charlotte Held Gotfredsen (DTU NMR Centre), and Dr. Aaron John Christian Andersen (DTU

Metabolomics Core). We acknowledge financial support from Carlsberg Infrastructure (CF20-0177),

Novo Nordisk Foundation Proof of Concept (NNF20OC0062267), DTU Enable Program,

InnoExplorer Grant, Innovation Fund Denmark, and Danish National Research Foundation (DNRF137)

for support towards the Centre for Microbial Secondary Metabolites (CeMiSt). Z.Y. acknowledges

funding from the China Scholarship Council (202004910340). T.W. acknowledges funding from the

Novo Nordisk Foundation (NNF20CC0035580).

Author contributions

Z.Y. and T.W. designed and carried out genetic experiments and bioinformatic analyses. E.S. isolated

the metabolites and characterised the structures; L.D. did structure elucidation, compounds crystalision,

and preliminary plant assays; Z.Y., Y.Q., N.C.K. and E.A. performed the plant and qRT-PCR assays;

G. P. and M. A. carried out large scale fermentation and downstream processing; P.H. carried out X-

ray crystallography and data analysis. All the authors discussed the results and commented on the

manuscript.

Competing interests

The authors declare no competing interests.

Correspondence and requests for materials should be addressed to Ling Ding.

Fig. 1. The effect of *S. iranensis* on barley seedlings under the abiotic stress and the bioactive components produced by *S. iranensis*. **a**, the box plots (middle bar=median, box limit=upper and lower quartile, extremes=min and max values) depict the plant height, fresh weight and dry weight of barley seedlings growing under osmotic stress mediated by 20% (w/v) PEG-6000 (*n*=18). Abbreviation: ***Mock***, control; ***Si***, treatment of *S. iranensis* culture broth; ***D***, treatment of *S. iranensis*/Δ*ptaA* culture broth; ***M***, treatment of blank medium (ISP2); Statistical significance was assessed by one-way ANOVA with post hoc Dunnett's multiple comparisons test. **b**, the box plots depict the plant height, fresh weight and dry weight of barley seedlings growing under salinity stress mediated by 100 mM NaCl (*n*=18). Abbreviation: ***Mock***, control; ***Si***, treatment of *S. iranensis* culture broth; ***D***, treatment of *S. iranensis*/Δ*ptaA* culture broth; ***M***, treatment of blank medium (ISP2). Statistical significance was assessed by one-way ANOVA with post hoc Dunnett's multiple comparisons test. **c**, the box plots depict the plant height, fresh weight and dry weight of barley seedlings growing under drought stress (*n*=18). Different colours of box plots indicate different growing conditions: green, 21 days water; brown, 7 days treatment after 7 days water followed by 7 days drought; red, 7 days water followed by 14 days drought. Statistical significance was assessed by one-way ANOVA with Tukey test. Abbreviation: ***Water***, well water for 21 days; ***CK***, 7 days treatment of water after 7 days water + 7 days drought; ***Si***, 7 days treatment of *S. iranensis* culture broth after 7 days water + 7 days drought; ***D***, 7 days treatment of *S. iranensis*/Δ*ptaA* culture broth after 7 days water + 7 days drought; ***M***, treatment of blank medium (ISP2); ***Drought***, 14 days drought after 7 days water. Statistical significance was assessed by one-way ANOVA with Tukey test. **d**, the metabolite profile of the native *S. iranensis* growing in liquid ISP2 medium, the known secondary metabolites were identified by HR-LC-MS/MS and highlighted; **e**, the bioactive components pteridic acid H (**1**) and pteridic acid F (**2**) isolated from *S. iranensis*.

Fig. 2. The effect of pteridic acids H and F on *Arabidopsis* seedlings under abiotic stress. *a*, phenotype of *Arabidopsis* seedlings growing under drought stress mediated by 15% (w/v) PEG-6000 using other treatments (bars=1 cm); *b*, phenotype of *Arabidopsis* seedlings growing under salinity stress mediated by 80 mM NaCl using other treatments (bars=1 cm); *c*, the box plots (middle bar=median, box limit=upper and lower quartile, extremes=min and max values) depict the primary root length and fresh weight of *Arabidopsis* seedlings growing on non-stress condition (n=16). Statistical significance was assessed by one-way ANOVA with Tukey test; *d*, the box plots depict the primary root length and fresh weight of *Arabidopsis* seedlings growing on drought stress condition (n=16). Statistical significance was assessed by one-way ANOVA with Tukey test; *e*, the relative expression level of TIP2;3 and SOS1 in *Arabidopsis* seedlings under salinity stress were measured after 1 hour and 24 hours of different treatments. Each treatment contains three biological replicates. Statistical significance was assessed by one-way ANOVA with post hoc Dunnett's multiple comparisons test; *f*, the lateral root number of *Arabidopsis* seedlings growing in different conditions (n=16). Statistical significance was assessed by one-way ANOVA with post hoc Dunnett's multiple comparisons test. *g*, the phenotype differences of lateral root growth of *Arabidopsis* seedlings growing in other conditions. Abbreviations: ***Mock***, control; ***PH***, treatment of 0.5 ng mL^{-1} pteridic acid H; ***PF***, treatment of 0.5 ng mL^{-1} pteridic acid F; **IAA**, treatment of 1.3 nM indole-3-acetic acid; ***ABA***, treatment of 1.3 nM abscisic acid.

Fig. 3. Biosynthesis mechanism of pteridic acids and CRISPR base editing application in *S. iranensis*. ***a***, the proposed
biosynthetic pathway of pteridic acids H (**1**) and F (**2**); ***b***, the proposed biosynthesis mechanism of elaiophylin (**3**),
and its macrodiolide formation is catalysed by thioesterase (TE) domain. ***c***, Sanger sequencing and LC-MS/MS
output of CRISPR base editing application of STOP codon introduction targeting the *ptaA* of *S. iranensis*. The 20-nt
protospacer sequence is highlighted in light green, whereas the 3-nt PAM sequence is shown in yellow. The codons
and corresponding amino acids are indicated, and the black double-headed arrow represents the position of the editing
window; ***c***, Extracted Ion Chromatography (EIC) for **1** and **2** (*m/z* 383.2428 [M+H]$^+$) and **3** (*m/z* 1047.5863 [M+Na]$^+$)
in the wild type *S. iranensis* (Control) and the mutant *S. iranensis*/Δ*ptaA*; ***d***, Sanger sequencing and LC-MS/MS
output of CRISPR base editing application of site-directed mutagenesis targeting the TE domain of *pta* BGC. EIC
for **1** an **2** (*m/z* 383.2428 [M+H]$^+$) and **3** (*m/z* 1047.5863 [M+Na]$^+$) in the wild type *S. iranensis* (Control) and the
mutant *S. iranensis*/M2089I + E2090K + D2091M.

Fig. 4. Geographical distribution of *pta*-containing *Streptomyces* strains possessing *pta* BGC. A total of 62 *Streptomyces* strains were displayed on the map and distinguished by different colours. For detailed strain information, see Supplementary Tab. 4.

Fig. 5. Abiotic stresses alleviation led by *S. violaceusniger* Tu 4113 and *S. rapamycinicu*s NRRL 5491. **a**, the box plots (middle bar=median, box limit=upper and lower quartile, extremes=min and max values) shows different growth of barley seedlings in osmotic stress mediated by 20% (w/v) PEG-6000 (n=16). Abbreviation: **Mock**, control; **Sv**, treatment of *S. violaceusniger* Tu 4113 culture broth; **Sr**, treatment of *S. rapamycinicus* NRRL 5491 culture broth; **b**, different growth of barley seedlings in salinity stress mediated by 100 mM NaCl (n=16). Abbreviation: **Mock**, control; **Sv**, treatment of *S. violaceusniger* Tu 4113 culture broth; **Sr**, treatment of *S. rapamycinicus* NRRL 5491 culture broth; **c**, different growth of barley seedlings in drought stress (n=16). Abbreviation: **CK**, 7 days treatment of water after 7 days water + 7 days drought; **Sv**, 7 days treatment of *S. violaceusniger* Tu 4113 culture broth after 7 days water + 7 days drought; **Sr**, 7 days treatment of *S. rapamycinicus* NRRL 5491 culture broth after 7 days water + 7 days drought. Statistical significance was assessed by one-way ANOVA with post hoc Dunnett's multiple comparisons test.

Fig. 6. The phylogenetic and secondary metabolites BGCs analysis of *pta*-containing *Streptomyces* strains. *a*, the
phylogenetic tree of 16S rRNA nucleotides sequences of *pta*-containing *Streptomyces* and other *Streptomyces* strains;
b, the heatmap depicts similarity differences of core biosynthetic genes of *pta* BGC between *S. iranensis* and other
14 *pta*-containing *Streptomyces* strains.

755
756
757
758
759