Biofilm characterization in the maize pathogen, *Fusarium verticillioides*

Chizné Peremore[1,2,] Brenda Wingfield[1,2], Quentin Santana[3], Emma T Steenkamp[1,2,]

Thabiso E Motaung[1,2, *]

[1]Division of Microbiology, Department of Biochemistry, Genetics and Microbiology,

University of Pretoria, Private Bag X20, Hatfield Pretoria 0028, South Africa

[2]Forestry and Agricultural Biotechnology Institute, University of Pretoria, Hatfield

0083, Pretoria, South Africa.

[3] Biotechnology Platform, Agricultural Research Council, Onderstepoort, South

Africa

[*]Correspondence: Email: thabiso.motaung@up.ac.za

Thabiso E Motaung orcid.org/0000-0002-8813-7671

University of Pretoria

Private Bag X20

Hatfield 0028

Pretoria

South Africa

Abstract

Nearly all microbes, including fungal pathogens, form biofilms, which are structured communities of microbial aggregates enclosed in self-produced extracellular polymeric substances (EPS) and attached to a surface. Studying plant-associated fungal biofilms can enhance understanding of fungal biology and knowledge of the links between fungal diseases and plants. However, only a few plant-associated fungi are reported to form biofilms. This study aimed to examine the ability of a mycotoxigenic fungus of maize, *Fusarium verticillioides*, to form biofilms under laboratory conditions. During our investigation, *F. verticillioides* stationary phase cultures incubated in liquid media developed a biofilm-like pellicle with a hyphal assemblage that appears in the form of a cloudy and thin slime material. Under the microscope, the biofilms exhibited a highly heterogeneous architecture made of dense, entangled, and compact hyphae, which were accompanied by a quantifiable EPS and extracellular DNA (eDNA). The biofilm was also found to respond to different abiotic conditions including pH and temperature, suggesting their relevance in a field setting. We further demonstrate the biofilm structural maintenance role of eDNA through treatment with DNase, which was only marginally effective during late biofilm stages, suggesting that it forms complex interactions with the EPS during biofilm maturation. Based on these results, we propose that *F. verticillioides* forms a 'true' biofilm that may act as a potential virulence factor.

Keywords: Biofilm, *Fusarium verticillioides*, extracellular polymeric substances, extracellular DNA, virulence factors

1. Introduction

After rice and wheat, maize (*Zea mays*) is the third most abundant grain crop, feeding millions of people, especially in Sub-Saharan Africa (FAO. 2021). However, its development and yield are threatened by *Fusarium verticillioides*, which systemically colonizes leaves, stems, roots, and kernels. The fungus can therefore induce serious damage that often manifests in *Fusarium* ear and stalk rot (Gai et al., 2018; Oren et al., 2003). These diseases have food safety and security implications due to mycotoxin contamination that is associated with them. Primarily, *F. verticillioides* secretes the mycotoxin fumonisin B_1, which contaminates symptomatic and asymptomatic maize kernels and stored grains (Munkvold and Desjardins, 1997). The toxicity of this compound is due to the inhibition of ceramide synthase and subsequent toxic intracellular accumulation of sphingosine and other sphingoid bases (Marin et al., 2013), ultimately imposing detrimental health effects on consumer populations (Wild and Gong, 2010). Indeed, studies from around the globe, including some from Africa, Asia, and Latin America, paint a disconcerting picture of how the prevalence of mycotoxins leads to a variety of human pathologies, including oesophageal and liver cancer in adults who consumed contaminated maize (Wild and Gong, 2010).

The mechanisms by which *F. verticillioides* invades maize have been outlined (Gai et al., 2018; Oren et al., 2003), providing important clues on the circumstances leading to infection symptoms and the precise anatomical locations of the maize plant that would probably harbour mycotoxins. Undoubtedly, the production of mycotoxins in *F. verticillioides* has been heavily emphasized over virulence factor production in numerous papers on mycotoxigenic fungi. The current study posits the development of biofilms as a virulence factor is somehow closely related to the accumulation of mycotoxins in fungi. Tell-tale is the biofilm 3D structure and biomass, both of which covered in extracellular polymeric substances (EPS). This structural design gives rise to emergent characteristics that are only seen in the biofilm mode of microbial life, such as surface adhesion-cohesion, spatial organization, physical and social interactions, chemical heterogeneity, and greater tolerance to antimicrobials (Karygianni et al., 2021). In the case of mycotoxigenic fungi, the biofilm EPS may exert a substantial effect on mycotoxin production; it may permit mycotoxins to stably accumulate and persist for longer periods of time due to the fact that that it glues the

cells together, in the process creating pockets and channels through which mycotoxins
can be concentrated and distributed within a biofilm, respectively. *Aspergillus*
fumigatus biofilms, for instance, augment the production of gliotoxin, a sulphur-
containing mycotoxin with immunosuppressive properties (Bruns et al., 2010). An
interesting area of research will be determining to what extent the components of a
biofilm, including the EPS and its associated cell-free components such as the
extracellular DNA (eDNA), influence mycotoxin production. Therefore, research on
biofilms will provide a fresh perspective on mycotoxin synthesis in economically
important fungi.

As biofilms present a cross-sectoral challenge, affecting a wide range of sectors
including healthcare, built environment, food and agriculture (Cámara et al., 2022), our
lack of understanding of how filamentous fungal biofilms originate, and how they adapt
to their microenvironments once developed, will restrict our capacity to identify and
counteract their detrimental impacts. This is apparent in the clinical setting, where
most clinical infections associated with medical instruments, including indwelling
devices (e.g., catheters, pacemakers, dentures, orthopedic prostheses, and heart
valves) colonized by biofilms are difficult to treat due to antifungal resistance of these
cell community structures (Bryers, 2008; Donlan, 2001; Lindsay and von Holy, 2006).
Marine biofilms, on the other hand, which are created by both microorganisms and
macroorganisms (such as algae), play a significant role in the environmental effects
of biological fouling, also known as marine biofouling, which is the accumulation of
undesirable biological matter on artificial submerged surfaces. Ships and underwater
surfaces (such as undersea cables and acoustic instruments) are examples of
colonized surfaces, and their treatment is difficult due to concurrent and intolerable
environmental impacts on non-target species (Callow and Callow, 2011; Bixler and
Bushan, 2012). The same type of challenges might apply to agriculture, where some
of the essential and most used tools and machinery are contaminated with harmful
fungal biofilms that are challenging to remove. When employed across many fields,
farming equipment colonized by biofilms has the potential to contaminate unaffected
fields with biofilm-derived propagules that may have acquired novel traits within a
biofilm, including resistance to fungicides.

Somewhat formal descriptions of biofilms in plant fungal pathogens have started to emerge (Motaung et al., 2020), with the latest being provided for *F. graminearum* (Shay et al., 2022). However, to date, we have no knowledge pertinent to the role of biofilm formation in many disease-causing plant fungi, including key biofilm components such as eDNA and EPS. The current study sought to see if the morphologies of surface-associated development by *F. verticillioides* might be included in current biofilm descriptions. Our research will contribute to a better understanding of how filamentous plant fungal pathogens coordinate survival by forming a community structure.

2. Materials and Methods

2.1. Fungal strains and maintenance

Strains of *F. verticillioides* were obtained from the Culture collection in the Forestry and Agricultural Biotechnology Institute (FABI), University of Pretoria, following isolation from maize samples in taken from fields in the Eastern Cape province of South Africa. Two strains (CMWF 1196, and CMWF 1197) were screened for their ability to form biofilms in this study. The strains were allowed to grow for two weeks at room temperature by plating on ¼ strength PDA (Potato Dextrose Agar; 10 g of PDA powder and 12 g of Difco agar in 1 L distilled H_2O). They were then used to develop a biofilm in liquid media as described below.

2.2. Screening for biofilm formation

Rapid screening of biofilm formation was performed for all fungal isolates by cutting a block of agar (5 mm x 5 mm) from the sporulating culture and inoculating it into 15 ml of three different growth media. These were: ¼ Potato Dextrose Broth (PDB, Potato Dextrose Broth; 6 g of PDB powder (Difco) in 1 L distilled H_2O), Roswell Park Memorial Institute-1640 broth (RPMI) and Sabouraud Dextrose Broth (SDB or Sabouraud liquid medium) in 50 ml falcon tubes. After mixing, the inoculate solution was poured into petri dishes (100 mm × 15 mm) and incubated at 25 and 30 °C for 24-72 hrs. They were visually analysed every 24 hrs for biofilm formation which is a hyphal assemblage that appears in the form of a cloudy and thin slime material. Photos were taken with an Epson Perfection V700 Photo scanner.

Biofilm formation for the remaining experiments was conducted using a cell counting

method. To do this, the inoculum for counting cells was prepared by adding 2 ml of 1x

Phosphate-buffered saline (PBS, 137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, and

1.8 mM KH_2PO_4) onto sporulating culture of *F. verticillioides* growing on ¼ PDA for

seven days. The plate was swirled to allow spores to be released into the PBS buffer,

and then counted using a haemocytometer placed under a Zeiss Axioskop 2 plus Light

Microscope. A volume of 40 µL of fungal cell culture (i.e., PBS spore suspension) was

diluted by adding 40 µL of 0.4% Trypan Blue solution (Sigma-Aldrich) into an

Eppendorf tube, in order to distinguish between dead and living cells. The

concentration of the harvested conidia was adjusted to 1 x 10^5 conidia/ml in ¼ PDB.

Of this 200 µL and left to form a biofilm in 48 wells plates (Corning® Costar® TC-

Treated Multiple Well Plates from Sigma-Aldrich) or chamber slides (Lab- Tek®

Chamber SlideTM System, 8 Well Permanox® Slide) and incubated for seven days at

25 °C. Once biofilms had matured, the culture was then analysed using different

microscopy techniques as explained later.

148 **2.3.   Analyses of biofilm-like structures**

To better describe the biofilms formed by *F. verticillioides*, colony and cell morphology

were first analysed on plates containing liquid media (PDB), where the former was

examined visually every 24 hrs for 7 days. Following the visual screening, the spores

were cultured in ¼ PDB under shaking (180 rpm) (Shake-O-Mat, LABOTEC) and

stationary conditions for 72 hrs in chamber slides, harvested and then analysed under

a light microscope (Zeiss Axioskop 2 plus Light Microscope (LM)). A scanning electron

microscope (SEM) was used to study the ultrastructure of biofilms formed as

previously described, in chamber slides or 48 well plates with glass coverslips (LASEC

SA). Sample preparation for SEM was performed according to Harding et al. (2010).

Samples were examined with Zeiss Ultra PLUS FEG SEM Confocal Laser Scanning

Microscopy (CLSM). This was used to analyse of *F. verticillioides* biofilms by forming

biofilms in chamber slides at different time points (24 and 7 days) and staining them

for 30 mins in a dark room with 100 µl of FUN-1, which fluorescent viability probe for

fungi (Chandra et al., 2001). Biofilms were then visualized using Zeiss LM 880 CLSM

with excitation wavelength at 488nm and emission at 650nm.

2.4. Analysis of biofilm-derived cells

This study considered whether biofilm-derived cells are different from planktonic cells in *F. verticillioides.* To do this, biofilms of *F. verticillioides* were formed as previously mentioned. After 7 days of incubation, cells were extracted from the biofilm by scraping and briefly agitated (30 s) to loosen the cells from the EPS matrix. This suspension was filtered through Mira cloth (Sigma-Aldrich) to separate the cells from the EPS material. In parallel, *F. verticillioides* planktonic cells were scraped from the sporulating cultures on ¼ PDA and added to PBS. The cells were then counted and adjusted to the desired concentration (1×10^5 spres/ml). A volume of 10 µl was aliquoted onto ¼ PDA and incubated at 25 °C for 7 days. The agar plates were then examined to study differences in growth and colony morphology, with three sections of each biofilm-derived and planktonic cells examined with Zeiss Ultra PLUS FEG SEM.

2.5. Quantification of biomass, EPS and metabolic activity

Biofilm biomass was determined in relation to the production of EPS and metabolic activity. To measure the biofilm biomass, crystal violet was used as it is known to be a good indicator of the amount of cellular biomass (Cruz et al., 2018). Biomass was then quantified according to the method described by Mello et al. (2016). Biofilms in 96 well plates (Corning® Costar® TC- Treated Multiple Well Plates (Sigma-Aldrich) were fixed for 15 min in 200 µl of 99% methanol. The supernatant was then removed, and the biofilm in microtiter plates was air-dried for 5 min prior to adding 200 µl of 0.3% crystal violet solution (stock solution diluted in PBS; Sigma-Aldrich) to each well. The stained biofilms in micotiter plates were then incubated for 20 min at room temperature prior to being rinsed twice with PBS to remove excess dye. The biomass in each well was then decolourized with 200 µl of 99% ethanol for 5 min. A volume of 100 µl of this solution was then transferred to a new 96-well plate and the absorbance was measured at 590 nm using a microplate reader (SpectraMax ® Paradigm® Multi-Mode Detection Platform).

Non-fixed biofilms were stained for 5 min at room temperature with 20 µl of 0.1% safranin (stock solution diluted in PBS:Sigma-Aldrich). After that, the stained biofilms were rinsed with PBS till the supernatants became transparent. With 200 µl of 99% ethanol, the extracellular matrix was decolorized. A volume of 10 µl of the supernatant

was transferred to a fresh 96-well plate, and the absorbance at 530 nm was measured using a microplate reader as previously reported. To measure the metabolic activity of the biofilm, biofilms were formed in 96 well plates as before and incubated at 25 °C. Once biofilms had been formed, their metabolic activity was then quantified using a colorimetric assay, XTT (sodium 3´- [1- (phenylaminocarbonyl)- 3,4- tetrazolium]-bis (4- methoxy6-nitro) benzene sulfonic acid hydrate) (Sigma-Aldrich), according to the manufacturer's recommended specifications. The activity of fungal mitochondrial dehydrogenase converts XTT tetrazolium salt to XTT formazan resulting in a colour change that can be measured using the microplate reader as previously described, with the absorbance from each well measured at 492 nm.

2.6.   The impact of abiotic conditions on biofilm growth

The impact of abiotic conditions on biofilm production by *F. verticillioides* was assessed at different pH values (2, 3, 4, 5, 6, 7, 8) and temperatures (10, 15, 20, 25 and 35 °C) in ¼ PDB. The pH was adjusted using HCl and NaOH. To conduct these experiments, standardized spore suspensions were inoculated as previously described and biofilms were allowed to develop under the aforementioned abiotic conditions for 7 days. Quantification of biomass, EPS, and metabolic activity was then performed using the microplate reader as previously described. Following this, EPS and biomass were determined, with EPS expressed per biomass (i.e., EPS/Biomass) and metabolic activity percentages calculated to compare the response of biofilms under the different physical factors.

2.7.   Effect of DNase treatment on biofilms formation

The eDNA release was measured using a microplate fluorescence assay using a DNA binding dye (SYBR® Green I), as previously described by Rajendran et al. (2014). First, EPS was extracted following a method described by Rajendran et al. (2013). For this study, biofilms were cultivated in ¼ PDB for 7 days at 25 °C before being scraped from the plates using sterile cell scrapers and rinsed with PBS. The EPS was extracted from the disaggregated biofilm using 0.2 M EDTA. Following this the samples were centrifuged at 10,000 x g for 30 min, the EDTA supernatant was collected and filtered through a 0.45 m syringe filter (Millipore). SYBR® Green I (Invitrogen), whose binding results in fluorescence that is directly proportional to DNA content, was applied at a

1:4 ratio to biofilm supernatants in a black well microtiter plate (Costar3603; Corning).
The levels of eDNA were then quantified using a fluorescence plate reader
(SpectraMax® Paradigm® Multi-Mode Detection Platform) at 485 and 525 nm,
respectively. The concentration of eDNA in the sample was quantified using the DNA
standard curve as previously described by Leggate et al. (2006).

The role of eDNA in *F. verticillioides* biofilm formation was investigated by depletion of
eDNA within the biofilm using the hydrolytic enzyme (Rajendran et al., 2013), DNase
I from bovine pancreas (Sigma-Aldrich). The DNase I was prepared in 0.15 M NaCl
supplemented with 5 mM of $MgCl_2$. To assess the effect of DNase I on biofilm
formation, biofilms were formed in ¼ PDB as described above and were incubated
with DNase I at the concentration of 0.25, 1, and 2 mg/ml, and incubated at 25°C for
72 hrs and 7 days. Untreated controls were included for direct comparison. After each
treatment, the biofilms were washed in PBS and their metabolic activity, biomass, and
EPS were quantified as mentioned prior.

2.8.   Statistical analysis

All experiments were performed in duplicate, in two independent experimental sets.
The data were expressed as mean ± standard deviation (SD). The results were
evaluated using the GraphPad Prism 9 computer program. A *p*-value of 0.05 or below
was deemed statistically significant in all analyses (ns= $p > 0.05$; *= $p \leq 0.05$; **= $p \leq$
0.01; *** =$p \leq 0.001$; ****= $p \leq 0.0001$).

3.   Results

3.1.   Colony morphology of *in vitro* biofilm-like structures

The ability of *F. verticillioides* to form biofilm-like structures was observed in both
strains examined in this study (Figure 1). Then, using visual inspection of liquid
cultures, colonies emerging from a biofilm culture were observed in PDB, SDB and
RMPI, incubated at 25 °C from 24 hrs to 7 days (data not shown). The biofilm is
normally distinguished from planktonic cells by their dense, highly hydrated clusters of
cells enmeshed in a gelatinous matrix (Coraça-Huber et al., 2020; Hurlow and Bowler,
2009, Metcalf and Bowler et al., 2013). Indeed, *F. verticillioides* biofilm-like colonies
displayed a dense, thin, and cloudy material (Figure 1). Based on these morphological

traits, the biofilm-like formations will be referred to as simply biofilms from this point
on.

3.2. *Fusarium verticillioides* biofilm development

Since, morphologically, both the strains appeared to be forming similar biofilms, only
one CMW 1196 was then selected and utilized for subsequent studies. It was
anticipated that a biofilm would form most effectively under stationary conditions and
assumed that the shear stress from shaking would prevent the formation of the EPS
matrix, a distinguishing feature of microbial biofilms. Therefore, the cells from *F.*
verticillioides CMW 1196 were cultured under both shaking and stationary conditions.
Following this, it was found that planktonic cells incubated under shaking conditions
did not typically clump together when observed under a light microscope (Figure 2A),
but those incubated under stationary conditions developed a community of cells
resembling a biofilm (Figure 2B). When these cells were analysed under SEM, little to
no EPS formation in cells incubated under shaking conditions was observed (Figure
2C). However, cells that were incubated without shaking formed a visible EPS (Figure
2D), which was not surprising to us given that EPS have been observed often after
growth under non-shaking conditions in a variety of biofilm investigations (Cavalheiro
and Teixeira, 2018).

3.3. *Fusarium verticillioides* biofilms and impact on cells therein

The development of a biofilm in *F. verticillioides* may influence the metabolic status of
cells and by extension, phenotype (Ramage et al., 2012). Confirming this were that
the spores at the dispersion stage (Figure 3D) appeared to be morphologically distinct
from the normal microconidia spores initially used as inoculum to initiate a biofilm
(indicated in Figure 3A). Usually, microconidia of *F. verticillioides* are club- or elliptical-
shaped or pointed at both ends (Figure 3A). However, the biofilm-derived spores
appeared to be more globose/lemon-shaped and slightly larger than typical conidial
cells (Figure 3D). Therefore, biofilm-derived cells, as indicated in Figure 3D, may
influence phenotypic diversity in *F. verticillioides*. For instance, when these cells were
harvested from a biofilm and plated on ¼ strength PDA, they displayed a colony
morphology that is different from cells not derived from a biofilm i.e., they formed a
colony smaller than that of their planktonic counterpart (Figure 3). This implies that the

physiological response of cells contained within a biofilm is greatly influenced by the biofilm ecosystem. However, the morphology of cells derived from a biofilm had no apparent differences when compared to the morphology of normal cells (Supplementary Figure 2), suggesting that the differences between these cells might largely be in their response to environmental signals, as in the case with observations in Figure 3, as opposed to their morphology.

3.4 EPS/Biomass and metabolic activity as indicators of biofilm response

The complexity of the biofilm is for the most part brought on by the release of EPS (Ramage et al., 2011). This means that the biofilm may possess the ability to affect the physiology of the cells within it by virtue of holding them in place, thus maintaining the biofilm's 3D structure while also optimising the exchange of nutrients and genetic material. As previously indicated, the results showed that unlike cells cultured under shaking conditions (Figure 2A, C), the generation of a visible EPS occurs concurrently with the establishment of a mature biofilm (Figure 2B, D). Since establishing that *F. verticillioides* biofilms produce EPS, in this study, we were also interested in how much of this material was being produced during biofilm formation and to what extent metabolically active cells contributed to the total biofilm biomass. For this reason, colorimetric assays were applied, namely crystal violet and XTT, to analyse the biomass and EPS by the absorption of safranin, and to assess the metabolic activity (cell viability) of the biofilm, respectively. In the case of the XTT reduction assay, the production of soluble coloured formazan salts by sessile cells is a direct reflection of cellular metabolic activity. According to the results, an increase in cell mass and EPS production (Figure 4A) seems to be accompanied by an increase in metabolic activity of the cells inside a biofilm (Figure 4B); EPS/Biomass increased significantly from 13% (3 days) to 45% (7 days) (p = 0,002), suggesting the more the biofilm matures the more EPS is produced.

3.5 Biofilm formation in response to abiotic factors

Having shown that *F. verticillioides* asexual cells could develop into a biofilm, this study then investigated how the biofilm will react to different environmental conditions, namely different temperature and pH conditions. As shown in Figure 5A, EPS/Biomass

was highest at pH 5, suggesting that *F. verticillioides* may prefer pH 5 for biofilm
formation. The media that was used in the initial experiments (¼ PDB) has a pH of
around 5, and it was in this medium that all the stages of a biofilm were observed
(Figure 2). The EPS and metabolic activity were essentially the same at acidic pH
levels (i.e., 2, 3, and 4, below the optimal pH of 5), whereas at pH levels higher than
the optimal (i.e., pH 6, 7, and 8, higher than pH 5), the EPS was produced at lower
levels than at pH 5 and essentially stayed at these levels. From the optimal pH (pH 5)
point of view, *F. verticillioides* biofilms seem to produce significantly more
EPS/Biomass than at pH 6 and pH 8 ($p \le 0.05$). However, this biofilm had a
significantly lower metabolic activity at a range of pHs from 2-8 (2, 3, 4, 6, 7, and 8) (p
≤ 0.001). This suggests that, although pH 5 permits better biofilm formation, the biofilm
response to pH is versatile, spanning a range of pH conditions, which could influence
the adaptability of this pathogen to a range of field conditions.

Figure 5C and D depict the influence of temperature on the production of *F.*
verticillioides biofilms. The best temperature tested for metabolic activity is 25 °C, and
the fungus displayed similar metabolic activities and formed robust biofilms at 20 and
25 °C with no significant differences. The fungus also formed biofilms at 10, 15 and
35 °C, but these were not as robust as 20 and 25 °C and had lower metabolic activity.
Different temperatures do not seem to significantly affect the EPS/Biomass
percentage except for 10 °C where the biomass was beyond detection.

3.6 The structural maintenance role of extracellular DNA

The biofilm was treated with DNase I to show how eDNA maintains the structure of
the biofilm. The DNAse caused the collapse of biofilm formation during early stages of
development i.e., at 72 hrs (Figure 6A). Also, the structural integrity of the biofilm was
revealed to be significantly impacted by the addition of DNase I in a concentration-
dependent manner. Unfortunately, the EPS/Biomass percentages could not be
determined during the early biofilm maturation phase (72 hrs) as the biofilm was too
weak to perform the relevant assays. In comparison to the biofilm growth control at 7
days, the application of 0.25 and 1 mg/ml, DNase I slightly reduced EPS/Biomass by
17% ($p > 0.05$) while 2 mg/ml significantly reduced it by 40% ($p = 0.0095$) (Figure 6B).
Interestingly, although DNase I had an impact on the EPS/Biomass, the metabolic
activity of the biofilm was not drastically affected (Data not shown), which could show

that eDNA has more impact on the structure of *F. verticillioides* biofilm than on cellular

activity, as has been demonstrated in other fungi

4. Discussion

Members of the genus *Fusarium* cause economically important and hard to control

diseases including cankers, crown rot, head blight, scabs, and wilts. Although many

of these diseases likely have strong links with biofilm formation (Harding et al., 2010;

Harding and Daniels, 2017; Motaung et al., 2020) given that microbes predominantly

exist in a state of a biofilm in their natural environments, biofilm formation has been

formally described in just a few *Fusarium* species including *F. oxysporum* f.

sp. *cucumerinum* and *F. graminearum* (Li et al., 2014; Shay et al., 2022). Therefore,

for many fungal pathogens of plants, including those belonging to *Fusarium*, it is

unclear how biofilms form, let along how they impact infections and disease outcomes.

Seven years ago, Miguel and his colleagues saw what looked like *F. verticillioides*

biofilms, in which the mycelium was structured in an extracellular material around the

hyphae (Miguel et al., 2015). These researchers also discovered a flocculent

substance over the cells or small fibrils of hyphae connecting to one another, like a

biofilm. To the best of my knowledge, this is the only time that evidence of a biofilm-

like structure for *F. verticillioides* has been reported *in vitro*. There is no formal

description of biofilms by this fungal pathogen. In the current study, we addressed this

knowledge deficit by describing how *F. verticillioides* forms biofilms under *in vitro*

conditions.

Information that describes biofilm colony morphologies in stationary liquid cultures is

generally lacking in fungal plant pathogen biofilm studies. The filamentous fungus *F.*

graminearum has recently been described as being able to form biofilm colonies at the

air/liquid interface that are growing as pellicle–floating masses of cells that cling to

each other and move as a unit (Shay et al., 2022). In Figure 1 of this study, it is shown

that floating masses for *F. verticillioides* can be distinguished from free-living

(planktonic) cells by forming colonies displaying a dense, thin, and cloudy material.

The results of this study are consistent with several other studies also reporting similar

features in plant and animal tissues, as well as surgically implanted equipment such

as catheters and pacemakers colonized by fungal and bacterial biofilms (Coraça-

Huber et al., 2020; Hurlow and Bowler, 2009; Santos et al., 2011; Trautner and
Darouiche, 2004).

In the current study, it was also observed that biofilms in *F. verticillioides* also
developed most efficiently under stationary conditions, while shear stress from shaking
conditions prevented proper biofilm formation. Cells incubated in the stationary
conditions without shaking had hyphal cells tangled with EPS that appeared to behave
like a matrix binding the hyphae together, and cells cultured under agitated conditions
seldomly clumped together or not at all. These findings are similar to those of Hawser
et al. (1998), who demonstrated that only a few cells are visible on the surface of
cultures when they are shaking, in contrast to stationary conditions, whereby cultures
are characterized by dense networks of hyphae. This might be due to the severe
shaking influencing cell architecture, matrix deposition, and biofilm formation (Soll and
Daniel, 2016). In *C. albicans*, shaking at a speed of 60 rpm prevents biofilm growth,
with biofilms exposed to shear stress being thinner than those exposed to non-shaking
conditions (Cavalheiro and Teixeira, 2018). Hawker et al. (1998) also found that lower
speeds result in the production of biofilms with no extracellular matrix but only hyphae,
while shaking at higher-speed results in a biofilm which consists of a few cells on the
surface. In this study, it was observed that some EPS material is present in planktonic
cultures (Figure 2C); however, this was not as abundant as in biofilm cells (Figure 2D),
suggesting that under conditions causing agitation of the fungal cells, the cells struggle
to produce the EPS matrix.

In light of the above, the findings presented here align with the biofilm concept
previously proposed (Córdova-Alcántara et al 2019; Harding et al. 2009, 2010;
Motaung et al., 2020). The biofilms of *F. verticillioides* seem to develop through spore
adhesion, microcolony formation, maturation, and dispersion. Although many fungal
and bacterial species have recorded comparable developmental stages for biofilm
formation, filamentous fungal biofilm formation seems to differ from strain to strain (Li
et al., 2014; Mowat et al., 2007; Ramage et al., 2011,2012). In contrast to unicellular
life forms such as yeast and bacteria, most fungi contain many planktonic forms that
can disperse and continue the cycle (e.g., sporangia asexual spores, sexual spores,
and hyphal fragments), and these dispersive forms most usually float in air rather than
water (Harding et al., 2009). In this study, the dispersal phase of biofilms leads to a

substantial number of free-living cells in the form of conidia, but these cells appeared to be morphologically distinct from the normal microconidial spores initially used as inoculum to initiate a biofilm (Supplementary Figure 1). This suggests that the dispersed biofilm cells differ from normal microconidia. Similar findings were reported in a study on *Bacillus cereus* where the cells in the biofilm have different cell-surface characteristics than their planktonic counterparts. For example, the structure of a polysaccharide linked to peptidoglycan in *B. cereus* has been discovered to change during biofilm development (Majed et al., 2016). In addition, according to Boles et al. (2004), the short-term development of *P. aeruginosa* in biofilms causes considerable genetic diversity in the resident bacteria. They also discovered that genetic diversity forms bacterial subpopulations in biofilms with specialized functions and that functional diversity improves the biofilm community's capacity to endure physiological stress. Equivalent diversity has not been shown in fungal biofilms and should be investigated further.

The biofilm detachment phase is important in disease development, and there has been growing evidence that specific techniques of dispersion can result in releasing biofilm cells that are more pathogenic than their planktonic counterpart (Beitelshees et al. 2018). The dispersed biofilm-derived conidia may be critical to the pathogen's invasive nature during fungal infections. In other studies, biofilm-derived cells have frequently been shown to be resistant to antifungals and stress (Guilhen et al., 2017). Therefore, *F. verticillioides* biofilm-derived conidia need to be further investigated for agriculturally relevant traits including heightened antifungal resistance against fungicides currently employed in agriculture, which could make the disease caused by this fungus challenging to treat.

Traditional methods for probing biofilm biomass, extracellular matrix, and metabolic were used in this study. In *A. fumigatus* biofilm have been previously linked with metabolism, biomass, and hyphal growth (Mowat et al. 2007). It was also shown that metabolic activity and biomass increased during the first 24 hours of biofilm formation and subsequently reaching a plateau of development (Mowat et al. 2007). Mello et al. (2016) focused on the assessment of biofilm formation by *Scedosporium* species and reported similar outcomes in some species in terms of biomass, metabolic activity, and EPS production. The production of the matrix has a high energetic cost, which may be

evolutionary justified given the matrix's structural and physicochemical significance in
the growth and operation of the biofilm, without which the beneficial emergent
properties of biofilms would not be possible (Flemming et al., 2016; Saville et al.,
2011). This goes to show that the formation of biofilms is closely linked to the formation
of the matrix, the bulk of which is extracellular material (Li et al., 2014).

As biofilms grow, metabolic activity and EPS production also rise, suggesting that they
may become more resistant to abiotic stress (Motaung et al., 2020). In the current
study, it was found that *F. verticillioides* developed biofilms under a range of pH and
temperature conditions, and a similar trend was previously reported (Li et al., 2014).
These results show that *F. verticillioides* biofilms are produced under different
conditions and have an optimum pH of 5 and an optimum temperature of 25 °C. Under
field conditions it has been reported that the optimum temperature for spore
development is 27 °C a temperature at which biofilm development could occur.
Evaluation of biofilm formation under different pH conditions showed that biofilms were
much more robust in somewhat acidic (pH 4-5) conditions than in acidic (pH 2, 3),
neutral (pH 7) or alkaline (pH 8) environments. These findings are consistent with other
reports (Cornet and Gaillardin, 2014) since the pH range for fungal development is
fairly broad, ranging from pH 3.0 to more than pH 8.0, with the optimum at pH 5.0
assuming nutritional needs are met. The capacity to develop a biofilm under varying
physical conditions may provide the fungus with survival benefits in inhospitable
environments.

Unlike other components of the biofilm EPS matrix, the eDNA has by far attracted the
most attention and is considered a useful tool to study the biology of biofilms. Many
studies have uncovered that eDNA plays many important roles in bacteria such as in
biofilm structural maintenance, assisted by the action of DNA binding proteins (Buzzo
et al., 2021; Kavanaugh et al., 2019; Whitchurch et al., 2002), antimicrobial resistance
(Okshevsky and Meyer, 2015; Rajendran et al., 2013), and acting as a reservoir for
the interexchange of genes through natural transformation (Merod and Wuertz, 2014).
Furthermore, eDNA assumes more unusual roles, including acting as a source of
energy and nutrients (e.g., carbon, nitrogen and phosphorus) (Ibáñez de Aldecoa et
al., 2017; Mulcahy et al., 2010; Pinchuk et al., 2008), and forming higher-order
conformations (e.g., G-quadruplex DNA) that further strengthen the biofilm through

extracellular EPS-eDNA networks (Seviour et al., 2021). Given these roles, eDNA is
an attractive target of antimicrobial drugs to manage biofilm-related infections.
However, there are only a few studies exploring the existence and function of eDNA
in filamentous fungi, with no studies conducted in plant pathogenic fungi. In the current
study, it was demonstrated that DNase I has an impact on the biofilm stability. In a
similar study using *A. fumigatus*, DNase I was effective at all concentrations (0.25, 1,
4 mg/ml) (Rajendran et al., 2013, 2014), but the maximal effect was observed with 4
mg/ml of DNase I (*p* value < 0.001) which is similar to the observations in this study.
The discovery that eDNA contributes significantly to the biofilm EPS in both bacteria
and fungi suggests that this may be a conserved and possibly active microbial biofilm
process. Nucleic acids are incorporated early in the development of the EPS matrix in
F. graminearum, and they similarly appear to function as a scaffold, likely regulating
the entire matrix structure of the biofilm (Shay et al., 2022). The finding presented in
the current study imply that eDNA plays a significant structural maintenance role
during biofilm development in a filamentous plant fungal pathogen and may contribute
to the severity of disease.

5. Conclusions

The study showed that the maize fungal pathogen, *F. verticillioides*, can develop a
biofilm under laboratory conditions. Even though the description described here of
fungal biofilms may have limitations when the natural environment is considered, at
the very least *F. verticillioides* biofilm development seems to follow a typical model
that entails attachment, colonization, development (including EPS synthesis),
maturity, and dissociation from EPS, with the last phase producing conidia to restart
the biofilm cycle. However, in the natural environment, plant stem and leaf surfaces
can be sparsely or densely colonized by diverse fungal biofilms and are likely more
complex than conditions used in this study. Laboratory-grown biofilms are a simple
surface-covering, frequently exhibiting confluent and compact uniformity that is
consistent with the original definition of biofilms (Motaung et al., 2020). Documenting
biofilm formation in the natural environment by analysing heavily infected maize plants
and a population of field strains, as opposed to a few ones, is needed to better
understand what is happening in field conditions. By better understanding the
complexity of plant-associated fungal biofilms and their phenotypic traits, it would be

possible to develop novel antifungal drug targets and treatment alternatives to decrease the prevalence of fungal infections. This study thus establishes a baseline with regards to *F. verticillioides* biofilms, showing its intricate structure and response to the environment. The findings presented further showed that eDNA degradation causes biofilms to collapse, suggesting it is a candidate for antifungal development. These findings represent the first investigation on eDNA analysis in a plant fungal pathogen. Taken together, *F. verticillioides* ability to form biofilms may give it an ecological edge in its battle to keep its place as a commensal and pathogen of maize. The biofilm might enable this fungus to evade host immunity, withstand antifungal treatment and competition from other microbes. The current study, together with earlier studies, therefore, will deepen the understanding of the relationship between disease outcomes and biotic interactions in *F. verticillioides*.

6. Declaration of competing interest

The authors affirm that they have no known financial or personal interests.

7. Acknowledgements

Funding from the National Research Foundation (South Africa) through the Thuthuka funding instrument (grant no. 129580) is acknowledged. Any conclusions and/ or recommendations made here are solely the responsibility of the author(s), and the NRF accepts no accountability in this regard.

8. Reference

Allesen-Holm, M., Barken, K.B., Yang, L., Klausen, M., Webb, J.S., Kjelleberg, S., Molin, S., Givskov, M., Tolker-Nielsen, T., 2006. A characterization of DNA release in *Pseudomonas aeruginosa* cultures and biofilms. *Molecular Microbiology* **59**, 1114-1128.

Beitelshees, M., Hill, A., Jones, C.H., Pfeifer, B.A., 2018. Phenotypic variation during biofilm formation: Implications for anti-biofilm therapeutic design. *Materials* **11**, 1086.

Berne, C., Kysela, D.T., Brun, Y.V., 2010. A bacterial extracellular DNA inhibits settling of motile progeny cells within a biofilm. *Molecular Microbiology* **77**, 815-829.

Bixler, G.D., Bhushan, B., 2012. Biofouling: lessons from nature. *Philosophical*
Transactions of the Royal Society A: Mathematical, Physical and Engineering
Sciences **370**, 2381-2417.

Bruns, S., Seidler, M., Albrecht, D., Salvenmoser, S., Remme, N., Hertweck, C.,
Brakhage, A.A., Kniemeyer, O., Müller, F.M., 2010. Functional genomic
profiling of *Aspergillus fumigatus* biofilm reveals enhanced production of the
mycotoxin gliotoxin. *Proteomics* **10**, 3097-3107.

Bryers, J.D., 2008. Medical biofilms. *Biotechnology and bioengineering* **100 1**, 1-18.

Buzzo, J.R., Devaraj, A., Gloag, E.S., Jurcisek, J.A., Robledo-Avila, F., Kesler, T.,
Wilbanks, K., Mashburn-Warren, L., Balu, S., Wickham, J., Novotny, L.A.,
Stoodley, P., Bakaletz, L.O., Goodman, S.D., 2021. Z-form extracellular DNA
is a structural component of the bacterial biofilm matrix. *Cell* **184**, 5740-
5758.e17.

Callow, J.A., Callow, M.E., 2011. Trends in the development of environmentally
friendly fouling-resistant marine coatings. *Nature Communications* **2**, 244.

Cámara, M., Green, W., MacPhee, C.E., Rakowska, P.D., Raval, R., Richardson,
M.C., Slater-Jefferies, J., Steventon, K., Webb, J.S., 2022. Economic
significance of biofilms: a multidisciplinary and cross-sectoral challenge. *npj*
Biofilms and Microbiomes **8**, 42.

Cavalheiro, M., Teixeira, M.C., 2018. *Candida* Biofilms: Threats, challenges, and
promising strategies. *Frontiers in Medicine* **5**, 28.

Chandra, J., Kuhn, D.M., Mukherjee, P.K., Hoyer, L.L., McCormick, T., Ghannoum,
M.A., 2001. Biofilm formation by the fungal pathogen *Candida albicans*:
development, architecture, and drug resistance. *Journal of Bacteriology* **183**,
5385-5394.

Coraça-Huber, D.C., Kreidl, L., Steixner, S., Hinz, M., Dammerer, D., Fille, M., 2020.
Identification and Morphological Characterization of Biofilms Formed by
Strains Causing Infection in Orthopedic Implants. *Pathogens (Basel,*
Switzerland) **9**, 649.

Córdova-Alcántara, I.M., Venegas-Cortés, D.L., Martínez-Rivera, M.Á., Pérez, N.O.,
Rodriguez-Tovar, A.V., 2019. Biofilm characterization of *Fusarium solani*

keratitis isolate: increased resistance to antifungals and UV light. *Journal of Microbiology* **57**, 485-497.

Cornet, M., Gaillardin, C., 2014. pH signaling in human fungal pathogens: a new target for antifungal strategies. *Eukaryotic Cell* **13**, 342-52.

Costa, D., Johani, K., Melo, D.S., Lopes, L., Lopes Lima, L., Tipple, A.F.V., Hu, H., Vickery, K., 2019. Biofilm contamination of high-touched surfaces in intensive care units: epidemiology and potential impacts. *Letters in applied microbiology* **68**, 269-276.

Costa-Orlandi, C.B., Sardi, J.C., Santos, C.T., Fusco-Almeida, A.M., Mendes-Giannini, M.J., 2014. In vitro characterization of *Trichophyton rubrum* and *T. mentagrophytes* biofilms. *Biofouling* **30**, 719-727.

Cruz, C.D., Shah, S., Tammela, P., 2018. Defining conditions for biofilm inhibition and eradication assays for Gram-positive clinical reference strains. *BMC Microbiology* **18**, 173.

Di Domenico, E.G., Farulla, I., Prignano, G., Gallo, M.T., Vespaziani, M., Cavallo, I., Sperduti, I., Pontone, M., Bordignon, V., Cilli, L., De Santis, A., Di Salvo, F., Pimpinelli, F., Lesnoni La Parola, I., Toma, L., Ensoli, F., 2017. Biofilm is a major virulence determinant in bacterial colonization of chronic skin ulcers independently from the multidrug-resistant phenotype. *International Journal of Molecular Sciences* **18**, 1077.

Donlan, R.M., 2001. Biofilms and device-associated infections. *Emerging Infectious Diseases* **7**, 277-281.

Essary, B.D., Marshall, P.A., 2009. Assessment of FUN-1 vital dye staining: Yeast with a block in the vacuolar sorting pathway have impaired ability to form CIVS when stained with FUN-1 fluorescent dye. *Journal of Microbiological Methods* **78**, 208-212.

FAO, I., UNICEF, WFP and WHO, 2021. The State of Food Security and Nutrition in the World 2021, *Transforming food systems for food security, improved nutrition and affordable healthy diets for all*. FAO, Rome.

Flemming, H.-C., Wingender, J., 2010. The biofilm matrix. *Nature Reviews Microbiology* **8**, 623-633.

Flemming, H.-C., Wingender, J., Szewzyk, U., Steinberg, P., Rice, S.A., Kjelleberg,
S., 2016. Biofilms: an emergent form of bacterial life. *Nature Reviews*
Microbiology **14**, 563-575.

Gai, X., Dong, H., Wang, S., Liu, B., Zhang, Z., Li, X., Gao, Z., 2018. Infection cycle of
maize stalk rot and ear rot caused by *Fusarium verticillioides. PLoS One* **13**,
e0201588.

Guilhen, C., Forestier, C., Balestrino, D., 2017. Biofilm dispersal: multiple elaborate
strategies for dissemination of bacteria with unique properties. *Molecular*
Microbiology **105**, 188-210.

Gupta, A., Gupta, R., Singh, R.L., 2016. Microbes and Environment. In: Singh, R. (eds)
Principles and Applications of Environmental Biotechnology for a Sustainable
Future. *Applied Environmental Science and Engineering for a Sustainable*
Future. Springer, Singapore, pp 43-84.

Harding, M.W., Marques, L.L., Howard, R.J., Olson, M.E., 2009. Can filamentous fungi
form biofilms? *Trends in Microbiology* **17**, 475-480.

Harding, M.W., Marques, L.L., Howard, R.J., Olson, M.E., 2010. Biofilm morphologies
of plant pathogenic fungi. *The Americas Journal of Plant Science and*
Biotechnology **4**, 43-47.

Harding, M.W., Daniels, G.C., 2017. In Vitro assessment of biofilm formation by soil-
and plant-associated microorganisms, Biofilms in Plant and Soil Health, pp.
253-273.

Hawser, S.P., Baillie, G.S., Douglas, L.J., 1998. Production of extracellular matrix by
Candida albicans biofilms. *Journal of Medical Microbiology* **47**, 253-256.

Henriques, A., Fraqueza, M., 2017. Biofilm-forming ability and biocide susceptibility of
Listeria monocytogenes strains isolated from the ready-to-eat meat-based
food products food chain. *LWT-Food Science and Technology* **81**, 180-187.

Hurlow, J., Bowler, P.G., 2009. Clinical experience with wound biofilm and
management: a case series. *Ostomy Wound Management* **55**, 38-49.

Ibáñez de Aldecoa, A.L., Zafra, O., González-Pastor, J.E., 2017. Mechanisms and
regulation of extracellular DNA release and Its biological roles in microbial
communities. *Frontiers in Microbiology* **8**, 1390.

Kaplan, J.B., 2010. Biofilm dispersal: mechanisms, clinical implications, and potential
therapeutic uses. *Journal of Dental Research* **89**, 205-218.

Karygianni, L., Ren, Z., Koo, H., Thurnheer, T., 2020. Biofilm Matrixome: Extracellular
Components in Structured Microbial Communities. *Trends in Microbiology* **28**,
668-681.

Kavanaugh, J.S., Flack, C.E., Lister, J., Ricker, E.B., Ibberson, C.B., Jenul, C.,
Moormeier, D.E., Delmain, E.A., Bayles, K.W., Horswill, A.R., 2019.
Identification of extracellular DNA-Binding proteins in the biofilm matrix. *mBio*
10.

Kuhn, D.M., George, T., Chandra, J., Mukherjee, P.K., Ghannoum, M.A., 2002.
Antifungal susceptibility of *Candida* biofilms: unique efficacy of amphotericin
B lipid formulations and echinocandins. *Antimicrobial agents and*
chemotherapy **46**, 1773-1780.

LaFleur, M.D., Kumamoto, C.A., Lewis, K., 2006. *Candida albicans* biofilms produce
antifungal-tolerant persister cells. *Antimicrobial agents and chemotherapy* **50**,
3839-3846.

Leggate, J., Allain, R., Isaac, L., Blais, B.W., 2006. Microplate fluorescence assay for
the quantification of double stranded DNA using SYBR Green I dye.
Biotechnology Letters **28**, 1587-1594.

Lewis, K., 2007. Persister cells, dormancy and infectious disease. *Nature Reviews*
Microbiology **5**, 48-56.

Li, P., Xiaoming, P., Huifang, S., Jingxin, Z., Ning, H., Birun, L., 2014. Biofilm formation
by Fusarium oxysporum f. sp. cucumerinum and susceptibility to
environmental stress. *FEMS Microbiology Letters* **350**, 138-145.

Lindsay, D., von Holy, A., 2006. Bacterial biofilms within the clinical setting: what
healthcare professionals should know. *Journal of Hospital Infection* **64**, 313-
325.

López, D., Vlamakis, H., Kolter, R., 2010. Biofilms. *Cold Spring Harbor Perspectives*
in Biology **2**, a000398.

Majed, R., Faille, C., Kallassy, M., Gohar, M., 2016. *Bacillus cereus* Biofilms-same,
only different. *Frontiers in Microbiology* **7**, 1054.

Marin, S., Ramos, A., Cano-Sancho, G., Sanchis, V., 2013. Mycotoxins: Occurrence,
toxicology, and exposure assessment. *Food and chemical toxicology* **60**, 218-
237.

Martinez, L.R., Casadevall, A., 2007. *Cryptococcus neoformans* biofilm formation
depends on surface support and carbon source and reduces fungal cell
susceptibility to heat, cold, and UV light. *Applied and Environmental*
Microbiology **73**, 4592-4601.

Martins, M., Uppuluri, P., Thomas, D.P., Cleary, I.A., Henriques, M., Lopez-Ribot, J.L.,
Oliveira, R., 2010. Presence of extracellular DNA in the C*andida albicans*
biofilm matrix and its contribution to biofilms. *Mycopathologia* **169**, 323-331.

Mello, T.P., Aor, A.C., Gonçalves, D.S., Seabra, S.H., Branquinha, M.H., Santos,
A.L.S., 2016. Assessment of biofilm formation by *Scedosporium*
apiospermum, S. aurantiacum, S. minutisporum and *Lomentospora*
prolificans. Biofouling **32**, 737-749.

Merod, R.T., Wuertz, S., 2014. Extracellular polymeric substance architecture
influences natural genetic transformation of Acinetobacter baylyi in biofilms.
Applied and Environmental Microbiology **80**, 7752-7757.

Metcalf, D.G., Bowler, P.G., 2013. Biofilm delays wound healing: A review of the
evidence. *Burns & trauma* **1**, 5-12.

Miguel, T.d.Á., Bordini, J.G., Saito, G.H., Andrade, C.G.d.J., Ono, M.A., Hirooka, E.Y.,
Vizoni, É., Ono, E., 2015. Effect of fungicide on F*usarium verticillioides*
mycelial morphology and fumonisin B 1 production. *Brazilian journal of*
microbiology **46**, 293-299.

Moënne-Loccoz, Y., Mavingui, P., Combes, C., Normand, P., Steinberg, C., 2015.
Microorganisms and Biotic Interactions, in: Bertrand, J.-C., P. Caumette, P.
Lebaron, R. Matheron, P. Normand, T. Sime-Ngando (Eds.), Environmental
Microbiology: Fundamentals and Applications: Microbial Ecology. Springer
Netherlands, Dordrecht, pp. 395-444.

Motaung, T., Peremore, C., Wingfield, B., Steenkamp, E., 2020. Plant-associated
fungal biofilms – knowns and unknowns. *FEMS Microbiology Ecology* **96**, 224.

Mowat, E., Butcher, J., Lang, S., Williams, C., Ramage, G., 2007. Development of a
694   simple model for studying the effects of antifungal agents on multicellular
695   communities of *Aspergillus fumigatus*. *Journal of Medical Microbiology* **56**,
696   1205-1212.

697   Mulcahy, H., Charron-Mazenod, L., Lewenza, S., 2008. Extracellular DNA chelates
698   cations and induces antibiotic resistance in *Pseudomonas aeruginosa*
699   biofilms. *PLoS pathogens* **4**, e1000213.

700   Mulcahy, L.R., Burns, J.L., Lory, S., Lewis, K., 2010. Emergence of *Pseudomonas*
701   *aeruginosa* strains producing high levels of persister cells in patients with
702   cystic fibrosis. *Journal of Bacteriology* **192**, 6191-6199.

703   Müller, F.-M.C., Seidler, M., Beauvais, A., 2011. *Aspergillus fumigatus* biofilms in the
704   clinical setting. *Medical Mycology* **49**, S96-S100.

705   Munkvold, G.P., McGee, D.C., Carlton, W.M., 1997. Importance of different pathways
706   for maize kernel Infection by *Fusarium moniliforme*. *Phytopathology* **87**, 209-
707   217.

708   Murillo-Williams, A., Munkvold, G., 2008. Systemic infection by *Fusarium verticillioides*
709   in maize plants grown under three temperature regimes. *Plant Disease* **92**,
710   1695-1700.

711   Okshevsky, M., Regina, V.R., Meyer, R.L., 2015. Extracellular DNA as a target for
712   biofilm control. *Current Opinion in Biotechnology* **33**, 73-80.

713   Oren, L., Ezrati, S., Cohen, D., Sharon, A., 2003. Early events in the *Fusarium*
714   *verticillioides*-maize interaction characterized by using a green fluorescent
715   protein-expressing transgenic isolate. *Applied and Environmental*
716   *Microbiology* **69**, 1695-1701.

717   Pinchuk, G.E., Ammons, C., Culley, D.E., Li, S.M., McLean, J.S., Romine, M.F.,
718   Nealson, K.H., Fredrickson, J.K., Beliaev, A.S., 2008. Utilization of DNA as a
719   sole source of phosphorus, carbon, and energy by *Shewanella spp*.:
720   ecological and physiological implications for dissimilatory metal reduction.
721   *Applied and Environmental Microbiology* **74**, 1198-1208.

Rajendran, R., Williams, C., Lappin, D., Millington, O., Martins, M., Ramage, G., 2013.
Extracellular DNA release acts as an antifungal resistance mechanism in
mature A*spergillus fumigatus* biofilms. *Eukaryotic Cell* **12**, 420–429.

Rajendran, R., Sherry, L., Lappin, D.F., Nile, C.J., Smith, K., Williams, C., Munro, C.A.,
Ramage, G., 2014. Extracellular DNA release confers heterogeneity in
Candida albicans biofilm formation. *BMC Microbiology* **14**, 303.

Ramage, G., Mowat, E., Jones, B., Williams, C., Lopez-Ribot, J., 2009. Our current
understanding of fungal biofilms. *Critical Reviews in Microbiology* **35**, 340-355.

Ramage, G., Rajendran, R., Gutierrez-Correa, M., Jones, B., Williams, C., 2011.
Aspergillus biofilms: clinical and industrial significance. *FEMS Microbiology*
Letters **324**, 89-97.

Ramage, G., Rajendran, R., Sherry, L., Williams, C., 2012. Fungal biofilm resistance.
International journal of microbiology **2012**, 528521.

Ramage, G., Robertson, S.N., Williams, C., 2014. Strength in numbers: antifungal
strategies against fungal biofilms. *International Journal of Antimicrobial*
Agents **43**, 114-120.

Santos, A.P., Watanabe, E., Andrade, D., 2011. Biofilm on artificial pacemaker: fiction
or reality? *Arquivos Brasileiros de Cardiologia* **97**, e113-20.

Sardi, J.D.C.O., Pitangui, N.D.S., Rodríguez-Arellanes, G., Taylor, M.L., Fusco-
Almeida, A.M., Mendes-Giannini, M.J.S., 2014. Highlights in pathogenic
fungal biofilms. *Ibero-American Journal of Mycology* **31**, 22-29.

Saville, R.M., Rakshe, S., Haagensen, J.A.J., Shukla, S., Spormann, A.M., 2011.
Energy-dependent stability of *Shewanella oneidensis* MR-1 biofilms. *Journal*
of Bacteriology **193**, 3257-3264.

Seviour, T., Winnerdy, F.R., Wong, L.L., Shi, X., Mugunthan, S., Foo, Y.H., Castaing,
R., Adav, S.S., Subramoni, S., Kohli, G.S., Shewan, H.M., Stokes, J.R., Rice,
S.A., Phan, A.T., Kjelleberg, S., 2021. The biofilm matrix scaffold of
P*seudomonas aeruginosa* contains G-quadruplex extracellular DNA
structures. *npj Biofilms and Microbiomes* **7**, 27.

Shay, R., Wiegand, A.A., Trail, F., 2022. Biofilm formation and structure in the filamentous fungus *Fusarium graminearum*, a plant pathogen. *Microbiology Spectrum* **10**, e0017122.

Sheppard, D.C., Howell, P.L., 2016. Biofilm exopolysaccharides of pathogenic fungi: lessons from bacteria. *Journal of Biological Chemistry* **291**, 12529-12537.

Soll, D.R., Daniels, K.J., 2016. Plasticity of *Candida albicans* biofilms. *Microbiology and Molecular Biology Reviews* **80**, 565-595.

Steinberger, R.E., Holden, P.A., 2005. Extracellular DNA in single- and multiple-species unsaturated biofilms. *Applied and Environmental Microbiology* **71**, 5404-5410.

Subroto, E., van Neer, J., Valdes, I., de Cock, H., 2022. Growth of *Aspergillus fumigatus* in Biofilms in comparison to C*andida albicans*. *Journal of Fungi* **8**, 48.

Trautner, B.W., Darouiche, R.O., 2004. Role of biofilm in catheter-associated urinary tract infection. *American journal of infection control* **32**, 177-183.

Whitchurch, C.B., Tolker-Nielsen, T., Ragas, P.C., Mattick, J.S., 2002. Extracellular DNA required for bacterial biofilm formation. *Science* **295**, 1487-1487.

Wild, C.P., Gong, Y.Y., 2010. Mycotoxins and human disease: a largely ignored global health issue. *Carcinogenesis* **31**, 71-82.

Wuyts, J., Van Dijck, P., Holtappels, M., 2018. Fungal persister cells: The basis for recalcitrant infections? *PLoS pathogens* **14**, e1007301-e1007301.

Zarnowski, R., Westler, W.M., Lacmbouh, G.A., Marita, J.M., Bothe, J.R., Bernhardt, J., Lounes-Hadj Sahraoui, A., Fontaine, J., Sanchez, H., Hatfield, R.D., Ntambi, J.M., Nett, J.E., Mitchell, A.P., Andes, D.R., 2014. Novel entries in a fungal biofilm matrix encyclopedia. *mBio* **5**, e01333-14.

Zarnowski, R., Sanchez, H., Covelli, A.S., Dominguez, E., Jaromin, A., Bernhardt, J., Mitchell, K.F., Heiss, C., Azadi, P., Mitchell, A., Andes, D.R., 2018. *Candida albicans* biofilm–induced vesicles confer drug resistance through matrix biogenesis. *PLOS Biology* **16**, e2006872.

Figure 1: *Fusarium verticillioides* strains that formed biofilm-like cultures in Petri-dishes containing ¼ strength Potato Dextrose Broth. The strains were left to grow, without shaking for 7 days at 25 °C.

Figure 2: *Fusarium verticillioides* cultured for 7 days under shaking conditions (**A, C**), remained in the planktonic (free-living) state, and without shaking (**B, D**), formed biofilms with observable extracellular polymeric substances (EPS, indicated by red arrows in **D**).

Figure 3: Biofilm-derived and planktonic cultured *Fusarium verticillioides* in chamber slides for (**A**) 24 hrs and (**C**) 7 days. The cells were stained with FUN-1 to determine biofilm developments. *F. verticillioides* cells from shaking (**B**) and non-shaking (**D**) cultures established colonies on ¼PDA. Colonies derived from biofilm cells of asexual cells that were formed were significantly smaller (**=$p \leq 0.01$) than those derived from planktonic cells (**E**). This is despite the fact that the respective cells were plated at the same concentrations and incubated under the same conditions. Each dot on the bar graphs represents an independent biological replicate. ns= $p > 0.05$; *= $p \leq 0.05$; **= $p \leq 0.01$; *** = $p \leq 0.001$; ****= $p \leq 0.0001$.

Figure 4: *Fusarium verticillioides* biofilm formation was assessed based on **(A)** metabolic activity, evaluated by XTT reduction assay and **(B)** biomass and extracellular polymeric substances (expressed as EPS/Biomass) evaluated using crystal violet (OD_{590nm}) and safranin (OD_{530nm}), respectively (*p*-value=0.0013). An increase in cell mass and EPS production is observed. Each dot on the bar graphs represents an independent biological replicate. ns= $p > 0.05$; *= $p \leq 0.05$; **= $p \leq 0.01$; *** =$p \leq 0.001$; ****= $p \leq 0.0001$.

Figure 5: *Fusarium verticillioides* biofilm formation assessed at various pH and temperature conditions by measuring **(A, C)** metabolic activity (XTT reduction assay (OD_{475nm}; p value<0.05) (expressed as metabolic activity percentage)), and **(B, D)** biomass and extracellular polymeric substances (expressed as EPS/Biomass percentage), evaluated using crystal violet (OD_{590nm}) and safranin (OD_{530nm}), respectively (p value<0.05). This data shows that pH 5 permits better biofilm formation, and the biofilm response to pH is versatile, spanning a range of pH conditions. The optimum temperatures evaluated for metabolic activity were 20 and 25°C. It also generated biofilms at 10, 15, and 35°C, but these were less robust and had lower metabolic activity. Different temperatures do not appear to have a significant impact on the EPS/Biomass %, except for 10°C, when the biomass could not be quantified. Each dot on the bar graphs represents an independent biological replicate. ns= $p >$ 0.05; *= $p \leq 0.05$; **= $p \leq 0.01$; *** =$p \leq 0.001$; ****= $p \leq 0.0001$.

Figure 6: *Fusarium verticillioides* biofilm response to DNase treatment. **(A)** The response of a 72hr-old biofilm to DNase at different concentrations (0.25, 1 and 2 mg/ml). **(B)** The response of biofilms measured in biomass and extracellular polymeric substances (expressed as EPS/Biomass percentage). This data shows that DNase I caused the collapse of biofilm formation at the early stages of growth, i.e., at 72 hrs, and the structural integrity of the biofilm was found to be strongly influenced by DNase I in a concentration-dependent manner. Each dot on the bar graphs represents an independent biological replicate. ns= $p > 0.05$; *= $p \leq 0.05$; **= $p \leq 0.01$; *** =$p \leq 0.001$; ****= $p \leq 0.0001$.