Characterization of HcaA, a novel autotransporter protein in *Helicobacter cinaedi*, and

its role in host cell adhesion

Sae Aoki,[a] Shigetarou Mori,[a] Hidenori Matsui,[a] Keigo Shibayama,[b] Tsuyoshi Kenri,[a] and

Emiko Rimbara[a#]

[a]Department of Bacteriology II, National Institute of Infectious Diseases, Tokyo, Japan

[b]Department of Bacteriology, Nagoya University Graduate School of Medicine, Nagoya,

Aichi, Japan

Running head: *Helicobacter cinaedi* autotransporter HcaA

[#]Address correspondence to Emiko Rimbara, rimbara@niid.go.jp
14

ABSTRACT

Helicobacter cinaedi infects the human gut and causes invasive infections such as

bacteremia and cellulitis through bacterial translocation. However, the mechanism by which

H. cinaedi attaches to host cells and establishes infection remains unclear. This study aimed to

investigate the relationship between a novel putative autotransporter protein, *H. cinaedi*

autotransporter protein A (HcaA), and its role in pathogenicity. The cytotoxicity of *H. cinaedi*

infection in colon epithelial cell lines (Caco-2 and HT-29) was assessed using a lactate

dehydrogenase assay, and it was found that cytotoxicity significantly decreased upon HcaA

knockout. Adhesion assays further revealed that the HcaA-knockout strain showed

significantly reduced attachment to the human epithelial colorectal adenocarcinoma cell line

(Caco-2) compared to that of the wild-type strain. Moreover, the recombinant HcaA protein

demonstrated strong adhesion properties to the human monocytic cell line (U937). The

adhesive activity was diminished when the RGD motif in HcaA was replaced with RAD,

indicating that the RGD motif in HcaA is crucial for host cell adhesion. To determine the role

of HcaA in *H. cinaedi* infection *in vivo*, C57BL/6 mice were orally infected with wild-type

and HcaA-knockout *H. cinaedi* strains. Bacterial colonization was assessed 7, 14, and 28 days

post-infection. At 7 days post-infection, colonization was significantly lower in mice infected

with the HcaA-knockout strain compared to those infected with the wild-type strain. In

conclusion, our findings suggest that HcaA, a novel putative autotransporter protein in *H.*

cinaedi, plays a significant role as an adhesin in establishing colonization.

IMPORTANCE

Helicobacter species are classified as gastric or enterohepatic according to their habitat. Among enterohepatic Helicobacter species, which inhabit the intestine, colon and liver, H. cinaedi has been most frequently isolated from humans. H. cinaedi often causes bacteremia and cellulitis in immunocompromised hosts. Here, we focused on the H. cinaedi autotransporter protein A (HcaA), a novel virulence factor in H. cinaedi. We discovered that HcaA contributes to cell adhesion via its RGD motif. Furthermore, in animal experiments, bacterial colonization was reduced in mice infected with HcaA-knockout strains, supporting the hypothesis that HcaA contributes to H. cinaedi adhesion to host cells. Our study provides a novel mechanism for the establishment of H. cinaedi infections and provides new insights into the role of autotransporter proteins in the establishment of Helicobacter infection.

INTRODUCTION

Helicobacter species are gram-negative, helical bacilli isolated from humans and mammals. They are classified into gastric or enterohepatic Helicobacter species based on their habitat. Helicobacter pylori, a typical gastric Helicobacter species, infects the human stomach and is a major cause of gastric cancer (1). H. cinaedi is the most frequently isolated enterohepatic Helicobacter species strain from humans, mainly found in immunocompromised patients with bacteremia. H. cinaedi is responsible for recurrent cellulitis, infected aortic aneurysm, subdural hematoma, and hepatic cyst infection (2–6). H. cinaedi was first isolated in 1984 from a homosexual man with intestinal symptoms in Seattle and it was first isolated in Japan in 2003 (7, 8). Though H. cinaedi infections are predominantly reported in Japan, there have been some case reports from Denmark (9, 10). A retrospective study at a single hospital in Japan indicated that, 2.2 % of all blood culture-positive samples turned out to be H. cinaedi-

positive when the blood culture period was extended, suggesting that *H. cinaedi* infections

might be overlooked in other countries (11).

Cytolethal distending toxin (CDT) is a virulence factor in a variety of gram-negative

pathogenic bacteria, including *H. cinaedi*. It consists of three polypeptide subunits, CdtA,

CdtB, and CdtC (12–14). CdtB possesses nuclease activity and causes DNA damage, leading

to senescence and apoptosis in host cells. CdtA and CdtC play a role in delivering CdtB into

host cells. Additionally, another virulence factor, *H. cinaedi* antigen CAIP (15), causes the

differentiation and maintenance of the pro-inflammatory profile of human macrophages and

triggers foam cell formation. Consequently, inflammation in atherosclerosis is induced.

Studies have been conducted on *H. cinaedi* biology and host-host interactions protection

against oxidative stress and changes in virulence via the production of alkyl hydroperoxide

reductase (16) as well as changes in oxygen demand during infection (17). However, the

mechanisms by which *H. cinaedi* interacts with host cells and establishes infection have not

been fully elucidated.

Autotransporter proteins (ATs) are type V secretion systems (T5SSs) comprising an

essential passenger domain that functions as the secreted portion, and an autotransporter

domain with a β-barrel that exports the passenger domain from the periplasm to the

extracellular environment. ATs are widely found in gram-negative bacteria and have diverse

functions, acting as toxins and adhesion factors (18). The *H. cinaedi* genome encodes a

previously uncharacterized putative autotransporter protein. In this study, we named this

uncharacterized protein *H. cinaedi* autotransporter protein A (HcaA) and investigated its

contribution to *H. cinaedi* pathogenicity.

RESULTS

Structure and localization of HcaA

HcaA of *H. cinaedi* MRY 08-1234 [1,835 amino acids (aa), 197 kDa] is predicted to be

composed of a signal peptide (1-27 aa), passenger domain (27-1425 aa), and β-barrel domain

(1426-1825 aa, Fig. 1A). The β-barrel structure, characteristic of ATs, was highly supported

based on the pLDDT score, indicating the high confidence of the model (Fig. 1A). Therefore,

HcaA is predicted to be ATs. We constructed knockout mutants (Fig. 1B, top and middle) to

identify the function of HcaA. *H. cinaedi* MRY08-1234 and MRY12-0027 strains were used

as host strains, and *hcaA* was knocked out by replacing *hcaA* with *aphA* by homologous

recombination (Fig. 1B, bottom). No *hcaA* complemented strains were constructed in this

study, because no suitable plasmids for constructing complementation strains have been

developed for *H. cinaedi*, and several trials have not succeeded in creating a *hcaA*

complemented strain by inserting the *hcaA* gene on the chromosome. Therefore, we generated

two mutants from strains with different backgrounds to confirm the effect of HcaA deletion.

We had previously demonstrated the role of efflux pumps in *H. cinaedi* by creating several

knockout strains in the same way (19). We believe that our approach (i.e., the use of multiple

knockout strains) is suitable for investigating gene function. The knockout of *hcaA* was

verified by polymerase chain reaction (PCR) amplification and immunoblotting using an anti-

HcaA antibody (Fig. 1C and D). Immunoblot analysis revealed that HcaA was localized in the

membrane fraction and was not secreted (Fig. 1D). Fluorescence immunostaining further

confirmed the presence of HcaA on the bacterial surface (Fig. S1). No growth defects by the

knockout of *hcaA* were observed (Fig. S2).

HcaA knockout reduced cytotoxicity and adherence of human colon epithelial cells *in*

vitro

The effects of the HcaA knockout on cellular cytotoxicity were evaluated *in vitro* using a

lactate dehydrogenase (LDH) assay. The human epithelial colorectal adenocarcinoma (Caco-

2) cell line and human colon cancer (HT29) cell line were infected with *H. cinaedi* MRY08-

1234 and MRY12-0027 strains. LDH production was significantly lower in cells infected with

the HcaA-knockout strain than in those with the wild-type strain (Fig. 2A). In Caco-2 cells,

the differences in LDH production were small, although yet significant differences were

observed ($P < 0.05$). Therefore, HcaA may contribute to the cytotoxic effects of *H. cinaedi*

infection on epithelial cells. HcaA knockout did not affect LDH production by mucus-

secreting HT29-MTX cells (Fig. 2A), suggesting that the protective role of mucin in reducing

the cytotoxicity of *H. cinaedi* was not affected by the knockout of HcaA. To investigate

whether CDT is involved in the reduction of cytotoxicity, we analyzed its expression of *cdt* at

the transcriptional level and observed no difference between the wild-type and HcaA-

knockout strains (Figure S3).

To determine whether HcaA contributes to cell adhesion and invasion, we performed a

gentamicin protection assay, and compared the rate of adherent and invaded *H. cinaedi* cells

between the wild-type and knockout strains. The adhesion rate of *H. cinaedi* MRY08-1234 to

Caco-2 cells decreased significantly by HcaA knockout (Fig. 2B, upper panel). A significant

reduction in the adhesion rate was also observed when *H. cinaedi* MRY12-0027 was used for

gene knockout (Fig. 2B, bottom panel). However, it is important to note that this assay

finding includes both bacteria adhered to the surface and those internalized within host cells.

The invasion rates of MRY08-1234 and MRY12-0027 were also reduced by HcaA knockout;

however, the differences were not significant (Fig. 2C). The invasion rate of adherent bacteria

was higher in the HcaA-knockout strain and was significantly higher in MRY12-0027 (*P*

<0.05, Fig. 2D).

Adhesion was reduced in the recombinant mutant HcaA protein

HcaA contains an Arg-Gly-Asp (RGD) motif at amino acid residues 348-350. To determine

the contribution of the RGD motif to adhesion, recombinant passenger regions of HcaA

spanning 27-1,425 aa (out of total of 1,835 aa) of MRY08-1234 (WT HcaA) and mutant

HcaA (with RGD motif replaced by RAD) were generated (Fig. 3A). Mutant HcaA showed a

significant reduction in binding to U937 cells compared to that of WT HcaA ($P < 0.05$),

indicating that the RGD motif contributes to binding to host cells (Fig. 3B). The predicted

protein structure of HcaA of MRY08-1234 determined using AlphaFold2 indicated that the

RGD motif is located on the surface of the protein, supporting the contribution of this motif to

the adherence function of HcaA in *H. cinaedi* (Fig. 3C, left). The RGD motif was conserved

in all *H. cinaedi* strains registered at the National Center for Biotechnology Information

(NCBI), including clinical isolates (Fig. S4). HcaA homologs have been reported in *H. bilis*,

H. fennelliae, *H. canicola*, and *H. canis*, all of which are closely related to *H. cinaedi* (20–22).

The structural predictions for HcaA homologs determined using AlphaFold2 suggest that

HcaA and its homologs have structural similarities in the passenger region; however, the

RGD motif was present only in HcaA (Fig. 3C and D). In addition, the HcaA homologs are

predicted to have a structure protruding from the surface, similar to the RGD motif of HcaA.

However, the protruding structure did not contain the RGD motif (Fig. 3C).

Effects of HcaA on the establishment of *H. cinaedi* colonization in mice

To clarify the role of HcaA in *H. cinaedi* infection *in vivo*, C57BL/6 female mice were

orally infected with wild-type and HcaA-knockout *H. cinaedi* strains. The recovery of *H.

cinaedi* from the colon, cecum, cecal contents, biceps femoris muscle, and trapezius muscle

was analyzed at 7, 14, and 28 days post-infection (dpi). Because *H. cinaedi* is known to cause

cellulitis and infected aortic aneurysm (23), muscle samples were also collected in addition to

the intestinal tract samples. Animal experiments were performed twice, yielding consistent

results. Results of the first experiment, including only the MRY08-1234 strain, are shown in

Fig. S5. The results of the second experiment, including both MRY08-1234 and MRY12-

0027 strains, are shown in Figs. 4 and S6. To confirm colonization, reactivity to *H. cinaedi*

was measured in mouse serum. An increase in reactivity against whole cells of *H. cinaedi* was

observed in both groups of infected mice. Reactivity to *H. cinaedi* tended to be higher in the

group of mice infected with the wild-type strain than in the mice infected with the knockout

strain (Fig. 4A). In the MRY08-1234 strain, the reactivity at 14 dpi was significantly higher

than that in the knockout strain ($P < 0.01$). Using the culture method, *H. cinaedi* was

recovered from tissues other than cecal contents in both groups of mice infected with the

wild-type and HcaA-knockout strains (Table 1). The amount of *H. cinaedi* was highest in the

cecum from mice infected with the wild-type strains (MRY08-1234 WT and MRY12-0027

WT) at 7 dpi. The recovery rate of *H. cinaedi* from the cecum was lower in mice infected with

the HcaA-knockout strain (0%) than in mice infected with the wild-type strain (MRY08-1234

WT, 2/6, 33.3%; MRY12-0027 WT, 2/5, 40.0%, $P = 0.12$). At 14 dpi, there were lower rates

of colonization in the cecum of mice infected with the HcaA-knockout strain (MRY08-1234

KO, 0/5, 0%; MRY12-0027 KO, 0/5, 0%) strain than in mice infected with the wild-type

strain (MRY08-1234, 4/5, 80.0%; MRY12-0027, 2/5, 40.0%, $P = 0.06$). In MRY08-1234,

there was a trend towards a lower rate of colonization in the colon of mice infected with the

HcaA-knockout strain than with the wild-type strain at 14 dpi. Similarly, in MRY12-0027,

there were also lower rates of colonization in mice infected with the HcaA-knockout strain

(MRY12-0027 ΔHcaA, 0/5, 0%) than in mice infected with the wild-type strain (MRY12-

0027 WT, 1/5, 20%). *H. cinaedi* was not isolated from the cecal contents. For the cecum and

colon, in which bacteria were frequently detected by the culture method, mRNA was

extracted and bacteria were detected by quantitative real-time PCR (qPCR). The frequency of

positive samples was higher in the group of mice infected with the wild-type strain than in

mice infected with the knockout strain (Fig. S6). Therefore, a combined analysis of the results

of the culture and qPCR assays showed that the frequency of positive detection was higher for

mice infected with the wild-type strain than in mice infected with the HcaA-knockout (Fig.

4B). A reduction in the number of HcaA-knockout bacterial cells was also observed by an

immunofluorescence analysis of the colon infected with *H. cinaedi* using an anti-*H. cinaedi*

antibody (Figs. 4C and S5). Regardless of the presence or absence of HcaA and the duration

of infection, *H. cinaedi* was observed in the mucosal epithelium. However, in samples from

mice infected with the HcaA-knockout strain, the number of bacteria tended to be lower

compared to samples from mice infected with the wild-type strain.

DISCUSSION

Enterohepatic *Helicobacter* species are increasingly recognized as critical pathogens in

mammals. Among them, *H. cinaedi* causes bacteremia, cellulitis, and infected aortic

aneurysms in humans (2, 4, 24). Although *H. cinaedi* infections are rarely reported, cases

have been increasing in Japan since 2003 (8, 11, 24). However, little is known about the

virulence factors of *H. cinaedi*. ATs have various functions that contribute to the

establishment of infections by pathogenic gram-negative bacteria. In this study, we found that

HcaA, a putative AT, contributes to adhesion to host cells and to the colonization of *H.*

cinaedi. This is the first report to elucidate the role of ATs in enterohepatic *Helicobacter*

species.

T5SS proteins are classified into subgroups labelled Va–Ve; however, this classification is

based on protein architecture and does not reflect the functions of the secreted passengers.

VacA produced by *H. pylori* is classified as a type Va AT; it binds to host cells and is

internalized, causing vacuolation characterized by the accumulation of large vesicles (25).

Serine protease autotransporters of *Enterobacteriaceae* (SPATEs) produced by

Enterobacteriaceae are also classified as type Va ATs (26). SPATEs exhibit protease activity

to cleave host proteins. HcaA is structurally similar to EspP, SPATEs in *Escherichia coli*. We

did not detect protease activity against casein and pepsin, the major substrates for SPATEs

(Fig. S7a and b). Furthermore, no significant differences in CDT expression were observed in

the presence or absence of HcaA (Fig. S3). In contrast, the number of bacterial cells adhering

to human epithelial cells and the level of cytotoxicity were significantly reduced by HcaA

knockout. These findings suggest that the reduced cytotoxicity observed in the HcaA-

knockout strain can be explained by the decrease in the number of adherent *H. cinaedi* cells,

rather than a direct cytotoxic effect of HcaA itself.

HcaA contains the RGD motif in its passenger region. The RGD motif acts as a fibronectin

cell adhesion site. It plays a role in adhesion to host epithelial cells that express receptors of

the RGD motif, such as integrins (27). Cell adhesion experiments using purified WT HcaA

with the RGD motif and mutant HcaA with the RAD motif showed that the adhesion ability

decreased significantly when RGD was replaced by RAD. Structural prediction indicated that

the RGD motif is located on the surface of HcaA at an externally protruding site. The RGD

motif is known to bind to integrins, which are thought to mediate the adhesion of HcaA to

cells. Our results strongly suggest that the RGD motif of HcaA plays a role in cell adhesion.

However, the adhesion ability of HcaA was not completely diminished by replacing RGD

with RAD. Therefore, HcaA may function as an adhesin independent of the RGD motif, and

the presence of the RGD motif enhances its ability to adhere to cells. HcaA homologs in other

enterohepatic *Helicobacter* species, such as *H. bilis*, *H. canis*, *H. fennelliae,* and *H. canicola,*

lacked the RGD motif (Fig. 3). In addition, the RGD motif of HcaA was highly conserved

among *H. cinaedi* strains, regardless of strain origin (Fig. S4). *H. canicola* is genetically most

closely related to *H. cinaedi* (22). However, a phylogenetic analysis indicated that HcaA is

most similar to the HcaA homolog of *H. fennelliae* and not to that of *H. canicola*. *H.*

fennelliae is the second most common enterohepatic *Helicobacter* species causing invasive

infections in humans. The structural similarity between HcaA and HcaA homologs of *H.*

fennelliae suggests that HcaA plays a critical role in the pathogenesis of enterohepatic

Helicobacter species in humans. The results of comparative genomic analyses indicate that *H.*

cinaedi is a human-adapted lineage, unlike *H. canicola* (22). The acquisition of the RGD

motif of HcaA during the evolution of *H. cinaedi* may have contributed to the adaptation of *H.*

cinaedi to humans, resulting in a higher isolation rate of *H. cinaedi* from humans compared to

other enterohepatic *Helicobacter* species. Further investigation is needed to identify the

HcaA-binding target in the host cell.

The mouse infection experiment also supported the results of *in vitro* analyses, showing

that HcaA knockout reduced the number of infected bacterial cells in the cecum and colon of

H. cinaedi-infected mice. In particular, the HcaA-knockout strain had significantly fewer

bacterial cells in culture than that in the wild-type strain at 7 dpi. Therefore, HcaA plays an

important role in the established phase of *H. cinaedi* infection. A similar effect was observed

for FaaA (flagella-associated autotransporter A), an AT possessed by *H. pylori* contributing to

gastric colonization, and an FaaA deficiency has been reported to significantly reduce the

number of colonized bacteria in the early stages of infection (28).

ATs have various functions. For example, ImaA, another ATs in *H. pylori,* is essential for

host colonization and modulates the host inflammatory response via the inflammatory

cytokines IL-8 and TNF-α (29). In addition, EspC, SPATEs, degrades various host proteins,

such as hemoglobin and mucin, and also to prevent cytotoxicity during the initial stage of

bacterial infection by regulating pore formation (26, 30). Thus, similar to other ATs, HcaA

can also have multiple functions related to the establishment of chronic *H. cinaedi* infections

in humans. Further analyses are needed to elucidate additional functions of HcaA beyond its

role in cell adhesion.

This study identified a novel virulence factor of *H. cinaedi*, HcaA, as an adhesin protein.

HcaA was shown to play an important role in establishing *H. cinaedi* infection both *in vitro*

and *in vivo*. On the other hand, adhesion was also observed in the HcaA-knockout strain,

although it was reduced compared to that of the wild-type strain, suggesting that factors other

than HcaA contribute to host adhesion. Further studies of HcaA and other factors are needed

to understand the interaction between *H. cinaedi* and host cells.

MATERIALS AND METHODS

Condition of bacterial strains and host cells

H. cinaedi strains MRY08-1234 (MLST: ST10) and MRY12-0027 (MLST: ST15) isolated

from the blood of patients with bacteremia in Japan were used throughout the study. The *H.*

cinaedi strains were cultured as described previously (19). Briefly, strains were subcultured

on Brucella agar (Becton, Dickinson, Franklin Lakes, NJ, USA) with 5% horse blood at 37 °C

under microaerobic conditions with hydrogen generated by the gas exchange method using an

anaerobic gas mixture (H_2, 10%; CO_2, 10%; and N_2, 80%). For liquid culture, bacteria were

suspended in Brucella broth (Becton, Dickinson) with 10% fetal bovine serum (FBS;

NICHIREI), and the culture was incubated at 37 °C under microaerobic conditions with

hydrogen.

Human epithelial colorectal adenocarcinoma cells (Caco-2), a human myelomonocyte cell

line (U937), a human colon cancer cell line (HT29), and mucus-secreting HT29 cells (HT29-

MX) were used in this study. Caco-2 cells were maintained in Eagle's minimum essential

medium (EMEM, Sigma Aldrich, St. Louis, MO, USA) containing 2 mM L-glutamine, 1 mM

sodium pyruvate, 1.5 mg/L sodium bicarbonate, 10 mg/L phenol red, 10% heat-inactivated

(56 °C, 30 min) FBS, 1% non-essential amino acids. U937, HT29, and HT29-MX cells were

maintained in RPMI 1640 (Sigma Aldrich) with 10% heat-inactivated FBS. All media

contained an antimicrobial cocktail (penicillin-streptomycin), except for the *H. cinaedi*

infection experiments. All cells were incubated in a 5% CO_2, humidified atmosphere at 37 °C,

in a stationary manner.

Generation of *hcaA* deletion mutants and antibodies to HcaA

Knockout mutants were generated according to previously described methods (19). Briefly,

1,000 base pairs up- and downstream of *hcaA* were PCR-amplified. The primers contained a

linker sequence overlapping with *aphA*. To obtain the construct of *aphA* sandwiched by the

target gene sequence, three purified PCR products, up- and downstream of *hcaA* and *aphA*,

were mixed and amplified. The donor fragment was electroporated into *H. cinaedi*. The

transformed *H. cinaedi* was plated on agar plates containing 10 µg/mL kanamycin and 3%

agar. After incubation, the growing colonies were selected as knockout mutants. The

knockout of *hcaA* was confirmed by PCR and an immunoblot analysis. Primers used for

cloning and PCR are listed in Table S1. For the immunoblot analysis, the whole bacterial cell

lysates, culture supernatant, soluble fraction, and membrane fraction were prepared as follows.

The bacterial culture was incubated until an optical density at 600 nm (OD600) of 0.4, and the

supernatant was collected by centrifugation (8,600 ×*g*, 10 min, 4 °C). Then, the insoluble and

the soluble fractions were prepared by extracting proteins from the pellet using the

EzBactYeast Crusher (ATTO Corp., Tokyo, Japan) and by centrifugation. The insoluble

fraction was used as the membrane fraction. These samples were separated and transferred to

Immobilon^TM -FL PVDF (pore size, 0.45 um; Millipore Inc., Burlington, MA, USA) using a

semidry electroblotting system. An anti-HcaA antibody was obtained by immunizing rabbits

with a KLH-conjugated peptide (RGDMPQESKTLEQN corresponding to positions 348–361

of HcaA in MRY08-1234), followed by affinity purification against the peptide. The resulting

rabbit polyclonal HcaA antibody was used as the primary antibody for the immunoblot

analysis (1:1,000), and peroxidase-conjugative AffiniPure donkey anti-mouse IgG (H+L)

(Jackson ImmunoResearch Inc., 1:50,000) was used as the secondary antibody. The bands

were visualized using ECL Select (Amersham Biosciences, Amersham, UK) and

Amersham^{TM} ImageQuant 800 (Amersham Biosciences).

Cell infection experiments

Strains were grown in Brucella broth with 10% FBS at 37 °C under microaerobic

conditions. After incubation, the bacterial culture was inoculated into 10 mL of fresh broth,

and the culture was incubated until an OD600 of 0.4. Bacterial cells were harvested by

centrifugation ($3,350 \times g$, 10 min, 25 °C) and resuspended in the appropriate medium for each

cell. Infections were performed at a multiplicity of infection (MOI) of 100. Immediately after

the addition of bacteria, the plates were centrifuged at $300 \times g$ for 5 min at 25 °C, followed by

incubation in 5% CO_2 humidified atmosphere, at 37 °C for 4 h.

Cytotoxicity was evaluated by measuring lactate dehydrogenase using the Cytotoxicity

LDH Assay Kit-WST (Dojindo Laboratories, Tokyo, Japan). The assay was performed

according to the manufacturer's instructions.

Cell adhesion and invasion ability were evaluated using a gentamicin assay. To evaluate

adhesion ability, cells were liberated by trypsinization, seeded into a 24-well tissue culture-

treated polystyrene flat-bottom microtiter plate (FALCON), and incubated to confluence.

Caco-2 cells were infected by adding 1 mL of bacterial suspension (MOI 100) directly to each

well. After 4 h of incubation, the un-bound bacteria were removed by washing with pre-

warmed PBS. To evaluate invasive ability, *H. cinaedi* adherent to Caco-2 cells was incubated

in EMEM with gentamicin at 250 μg/mL. After 1.5 h, the infected cells were washed with

pre-warmed PBS. For both assays, host cells were washed with PBS containing 0.1% saponin

(FUJIFILM Wako Pure Chemical Co., Osaka, Japan). Serial dilutions of the suspension were

plated on Brucella agar containing 5% horse blood and 3% agar. After incubation, colonies

were counted, and the adhesion rate (%), invasion rate (%), and the invasion rate of adherent

bacteria (%) were calculated.

337 Purification of HcaA

To produce recombinant HcaA, we constructed expression vectors; *hcaA* was amplified by

PCR and introduced into pCold™ I DNA (TaKaRa, Shiga, Japan) using the In-Fusion® HD

Cloning Kit. gDNA from *H. cinaedi* strain MRY08-1234 was used to amplify the passenger

region of HcaA (27-1,425/1,835 aa). The expression vector containing the passenger region of

HcaA was transferred into *Escherichia coli* BL21 (DE3) pLysS cells and plated on LB agar

plates containing 100 µg/mL ampicillin. The growing colonies were inoculated into 100 mL

of LB broth containing 100 µg/mL ampicillin and incubated at 37 °C with shaking at 200 rpm

for 3 h. The cells were then transferred to 900 mL of fresh medium and grown in a 2 L flask

at 37 °C until OD600 of 0.6. Protein expression was induced by cooling in ice water for 1 h

and adding 400 µL of 1 mM isopropyl β-D-thiogalactopyranoside. Recombinant HcaA was

expressed as a fusion protein containing a histidine tag. The cell cultures were further

incubated at 15 °C overnight. Bacteria were harvested by centrifugation at 5,000 $\times g$ for 10

min at 4 °C and resuspended in 20 mM Tris-HCl (Nacalai, Kyoto, Japan), 150 mM NaCl

(FUJIFILM Wako), and 40 mM imidazole (FUJIFILM Wako) at pH 7.6 with cOmplete™

Protease Inhibitor Cocktail (Roche Molecular Systems). The suspended solution was

sonicated on ice, and the supernatant was collected by centrifugation at 5,000 $\times g$ for 45 min at

4 °C. The recombinant protein was purified in two steps using an affinity column (HisTrap FF

crude, Cytiva), followed by size exclusion chromatography (Superdex™ 200 Increase 10/300

GL, Cytiva). To replace the RGD motif of the HcaA protein with RAD, site-directed

mutagenesis of the HcaA expression vector was performed using PrimeSTAR® GXL DNA

Polymerase. The primers used to generate expression vectors are listed in Table S1.

360   Adhesion assay of HcaA to U937 cells

Purified HcaA and mutant HcaA were prepared at a concentration of 50 µg/mL in PBS,

dispensed in 100 µL aliquots to 96-well plates, and fixed to the bottom of the plates by

incubation overnight at 4 °C. The wells were then washed with PBS and blocked by adding

2% BSA in PBS and incubated again overnight at 4 °C. U937 cells were incubated until

confluence and then diluted with MEM to a concentration of 1×10^5 cells/mL. The use of

U937 cells in this assay was based on previous analyses of their activity in relation to *H.*

pylori (31). Fluorescein (FUJIFILM Wako) was added to the cell suspension at a final

concentration of 1 mg/mL and incubated at 37 °C for 30 min. After washing, MEM

containing 2 mM Ca and 2 mM Mg was added and allowed to react for 30 min on ice in the

dark. The cell suspension, diluted to a concentration of 5×10^5 /mL, was then added to HcaA-

coated plates and incubated at 37 °C for 1 h. After washing and replacement with phenol red-

free MEM medium (Sigma), fluorescence was measured, and the number of cells bound to

purified proteins was observed.

375   Sequence alignment, phylogenetic analysis, and structure prediction of HcaA and HcaA

376   homologs

Amino acid sequences of HcaA homologs in other enterohepatic *Helicobacter* species were

obtained from the whole genome sequences of *H. bilis* ATCC43879 (Accession No.

KI392040), and *H. fennelliae* MRY12-0050 (Accession No. BSAD00000000), and *H.*

canicola PAGU1409 (Accession no. DRR092854) and *H. canis* CCUG32756 (Accession No.

GCF008693005) were obtained from the National Center for Biotechnology Information

(NCBI). The passenger regions of the amino acid sequences of the HcaA and HcaA homologs

were aligned using ClustalW ver. 2.1 (32), and visualized using CLC Genomics Workbench

Version 22.0.1 (QIAGEN). A phylogenetic tree based on the aligned amino acid sequences of

the passenger domain of HcaA and HcaA homologs was constructed using RAxML-NG

v1.1.0 (https://github.com/amkozlov/raxml-ng) and visualized using MEGA X v 10.2.6 (33).

The prediction of the structures of HcaA and HcaA homologs were performed using

AlphaFold2 Ver.2.2.2 on the supercomputer system of the Institute for Chemical Research,

Kyoto University, with a full dBs preset.

Animal experiments and infection

Animal experiments were performed twice; the results of the first animal experiment are

shown in the Supplementary Appendix, and the results of the second experiment are reported

in the manuscript. Six-week-old female C57BL/6 mice (Japan SLC, Inc., Shizuoka, Japan)

maintained under specific pathogen-free conditions were used in this study. Mice were fed a

standard rodent chow, given ad libitum access to water, housed in microisolator cages, and

isolated from specific pathogens.

Strains were suspended in PBS at 1×10^9 cells/mL. The mice were then orally administered

0.2 mL of each bacterial suspension through a 1 mL syringe attached to an oral probe every

other day for a total of three times. The same volume of PBS was administered to mice as a

negative control. There were 14 mice in the control group, 15 mice in the wild-type strain

infection group, and 15 mice in the HcaA-knockout strain infection group.

On days 7, 14, and 28 after inoculation, five or six mice from each group were sacrificed

for analyses. The blood, cecal contents, colon, biceps femoris muscle, and trapezius muscle

were collected aseptically for further analysis. All animal experiments and the study protocol

were approved by the Committee for Animal Experimentation of the National Institute of

Infectious Diseases (approval number 121126).

ELISA

To measure reactivity to *H. cinaedi* in mouse serum, an enzyme-linked immunosorbent

assay (ELISA) was performed as described previously (34). To prepare serum, whole blood

collected at autopsy was incubated at 4 °C and then centrifuged at 2,000 ×*g* for 5 min at 4 °C.

The supernatant was collected, mixed with glycerol (50% final concentration), and designated

as serum. To prepare whole-cell lysates, *H. cinaedi* MRY08-1234 cells were disrupted by

sonication. Each well of Nunc-Immuno 96-well microtiter plates (No. 439454; Thermo Fisher

Scientific, Waltham, MA, USA) was coated with of whole-cell lysates (4 ug/mL in 0.1 M

carbonate/bicarbonate buffer, pH 9.4). After overnight incubation at 4 °C, the plate was

washed with PBS containing 0.05% (vol/vol) Tween 20 (PBS-T), and the wells were

saturated with blocking buffer (PBS containing 1% BSA) and incubated for 1 h at 37 °C while

shaking at 500 rpm. After washing with PBS-T, the wells were filled with serum samples,

which were diluted 100-fold with a blocking buffer. The dilution of serum was determined by

preliminary experiments (Fig. S8). Plates were incubated for 1 h at 37 °C with shaking at 500

rpm. Horseradish peroxidase-conjugated goat anti-mouse IgG secondary antibody (Jackson,

Bar Harbor, ME, USA) diluted 5,000-fold in blocking buffer was used. Absorbance was

measured at 450 nm using a Multiskan FC microplate reader (Thermo Scientific).

Isolation and detection of *H. cinaedi* from infected mice

Culture and PCR were used to assess *H. cinaedi* infection in each organ. For the cultivation

of *H. cinaedi*, agar plates containing Skirrow (Oxoid), 5% horse blood, 3% agar, and 4 µg/mL

ciprofloxacin (FUJIFILM Wako) were inoculated with the cecal contents, colon, biceps

femoris muscle and trapezius muscle that had been homogenized in Brucella broth. MRY08-

1234 is resistant to ciprofloxacin; accordingly, ciprofloxacin was added to eliminate

commensal intestinal bacteria in mice and isolate only *H. cinaedi*. Plates were incubated at

37 °C under microaerobic conditions. The growth of the colonies was quantified and

confirmed via detection of *cdt* using PCR to determine whether they were colonies of *H.*

cinaedi. Data for MRY08-1234 WT and MRY08-1234 KO at 7 dpi were collected from six

mice in each group, while data for other strains were collected from five mice in each group.

We also evaluated total RNA of *H. cinaedi*. Samples were homogenized with 1 mL of Isogen

(Nippon Gene, Toyama, Japan). The homogenates were supplemented with 0.2 mL of

chloroform (Sigma Aldrich) and mixed. The samples were then centrifuged (12,000 $\times g$, 15

min, 4 °C), and the aqueous layer was transferred to new tubes. RNA was precipitated by

combining 500 µL of isopropanol with the aqueous layer, followed by incubation at room

temperature for 10 min and centrifugation as described above. Total RNA was extracted from

the supernatant using the PureLink® RNA Mini Kit (Thermo Fisher Scientific, Waltham,

USA) according to the manufacturer's instructions. The extracted RNA was used for

quantitative real-time PCR (qPCR). Reverse transcription and reverse transcription

quantitative PCR (RT-qPCR) were performed using ReverTra Ace qPCR RT Master Mix

with gDNA Remover (Toyobo, Osaka, Japan), THUNDERBIRD™ SYBR qPCR MIX

(Toyobo), and Applied Biosystems® 7500Fast (Thermo Fisher Scientific). We assessed the

colonization of *H. cinaedi* by measuring the expression of *cdt* at the transcriptional level.

Primers for PCR and qPCR are listed in Table S2.

Histology and fluorescence images

Colons from mice were harvested and fixed with a 4% paraformaldehyde phosphate buffer

solution (FUJIFILM Wako). Sections obtained from fixed specimens were permeabilized and

incubated with an anti-*H. cinaedi* antibody (1:1,000) as the first antibody at room temperature

for 1 h. The anti-*H. cinaedi* antibody was obtained by immunizing rabbits with heat-

inactivated *H. cinaedi* MRY08-1234 as an antigen, followed by protein A-purification. The

sections were then rinsed and stained with Alexa Fluor 488 (*H. cinaedi*) and 4',6-diamidino-2-

phenylindole (nuclei). Fluorescence signals were visualized using an automated slide

scanning system (Olympus VS200, Olympus, Tokyo, Japan) and images were analyzed using

ImageJ ver.1.53t.

Statistical analysis

Differences in the recovery of *H. cinaedi* by culture from infected mice were analyzed

using Fisher's exact tests. For other experiments, the Mann–Whitney test was used for

comparison. All statistical analyses were performed using GraphPad Prism version 9.4.1 and

differences were considered statistically significant when the *P*-value was <0.05.

ACKNOWLEDGMENTS

This work was supported by JSPS KAKENHI Grant Numbers 16K18928, 19K07074 (E.

R.) and 22K15465 (S. A.), the GSK Japan Research Grant 2017 (E. R.) and by AMED

JP20fk0108148 (E. R.).

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

References

1. Polk DB, Peek RM. 2010. *Helicobacter pylori*: Gastric cancer and beyond. 6. Nat Rev

Cancer 10:403–414. https://doi.org/10.1038/nrc2857.

2. Matsuo T, Mori N, Mizuno A, Sakurai A, Kawai F, Starkey J, Ohkushi D, Abe K,

Yamasaki M, Ito J, Yoshino K, Mikami Y, Uehara Y, Furukawa K. 2020. Infected aortic

aneurysm caused by *Helicobacter cinaedi*: Case series and systematic review of the literature.

BMC Infect Dis 20:854. https://doi.org/10.1186/s12879-020-05582-7.

3. Akiyama T, Imamura H, Fukui N, Sakai N. 2021. *Helicobacter cinaedi*-infected chronic

subdural hematoma mimicking an expanding hematoma: A case report. Surg Neurol Int

12:288. https://doi.org/10.25259/SNI_387_2021.

4. Kawamura Y, Tomida J, Morita Y, Fujii S, Okamoto T, Akaike T. 2014. Clinical and

bacteriological characteristics of *Helicobacter cinaedi* infection. J Infect 0Chemother 20:517–

526. https://doi.org/10.1016/j.jiac.2014.06.007.

5. Suzuki T, Kutsuna S, Tsuboi M, Ota M, Hayakawa K, Ohmagari N. 2019. *Helicobacter*

cinaedi hepatic cyst infection with bacteremia. Emerg Infect Dis 25:603–604.

https://doi.org/10.3201/eid2503.180936.

6. Saito J, Rimbara E, Inaguma S, Hasegawa C, Kamiya S, Mizuno A, Sone Y, Ogawa T,

Numata Y, Takahashi S, Asano M. 2022. Determining infected aortic aneurysm treatment

using focused detection of *Helicobacter cinaedi*. Emerg Infect Dis 28:1494–1498.

https://doi.org/10.3201/eid2807.212505.

7. Totten PA, Fennell CL, Tenover FC, Wezenberg JM, Perine PL, Stamm WE, Holmes KK.

1985. *Campylobacter cinaedi* (sp. nov.) and *Campylobacter fennelliae* (sp. nov.): Two New

Campylobacter species Associated with Enteric Disease in Homosexual Men. J Infect Dis

151:131–139. https://doi.org/10.1093/infdis/151.1.131.

8. Murakami H, Iwata M, Goto M, Takahashi T, Ono E, Sawabe E, Okuzumi K, Yamaguchi

K, Takahashi T. 2003. Isolation of *Helicobacter cinaedi* from blood of an

immunocompromised patient in Japan. J Infect Chemother 9:344–347.

https://doi.org/10.1007/s10156-003-0265-3.

9. Holst H, Andresen K, Blom J, Højlyng N, Kemp M, Krogfelt KA, Christensen JJ. 2008. A

case of *Helicobacter cinaedi* bacteraemia in a previously healthy person with cellulitis. Open

Microbiol J 2:29–31. https://doi.org/10.2174/1874285800802010029.

10. Nielsen HL, Prag J, Krogfelt KA. 2015. *Helicobacter cinaedi* knee infection after

arthroscopy in an immunocompetent patient. BMJ Case Rep 2015:bcr2014208637.

https://doi.org/10.1136/bcr-2014-208637.

11. Araoka H, Baba M, Kimura M, Abe M, Inagawa H, Yoneyama A. 2014. Clinical

characteristics of bacteremia caused by *Helicobacter cinaedi* and time required for blood

cultures to become positive. J Clin Microbiol 52:1519–1522.

12. Guerra L, Cortes-Bratti X, Guidi R, Frisan T. 2011. The biology of the cytolethal

distending toxins. Toxins 3:172–190. https://doi.org/10.3390/toxins3030172.

13. Taylor NS, Ge Z, Shen Z, Dewhirst FE, Fox JG. 2003. Cytolethal distending toxin： A

potential virulence factor for *Helicobacter cinaedi*. J Infect Dis 188:1892–1897.

14. Scuron MD, Boesze-Battaglia K, Dlakić M, Shenker BJ. 2016. The Cytolethal

distending toxin contributes to microbial virulence and disease pathogenesis by acting as a tri-

perditious toxin. Front Cell Infect Microbiol 6:168.

15. D'Elios MM, Vallese F, Capitani N, Benagiano M, Bernardini ML, Rossi M, Rossi GP,

Ferrari M, Baldari CT, Zanotti G, de Bernard M, Codolo G. 2017. The *Helicobacter cinaedi*

antigen CAIP participates in atherosclerotic inflammation by promoting the differentiation of

macrophages in foam cells. Sci Rep 7:40515. https://doi.org/10.1038/srep40515.

16. Charoenlap N, Shen Z, McBee ME, Muthupalani S, Wogan GN, Fox JG, Schauer DB.

2012. Alkyl hydroperoxide reductase is required for *Helicobacter cinaedi* intestinal

colonization and survival under oxidative stress in BALB/c and BALB/c interleukin-10–/–

mice. Infect Immun 80:921–928. https://doi.org/10.1128/IAI.05477-11.

17. Schmitz JE, Taniguchi T, Misawa N, Cover TL. 2016. Epithelial coculture and l-lactate

promote growth of *Helicobacter cinaedi* under H2-free aerobic conditions. Appl Environ

Microbiol 82:6701–6714. https://doi.org/10.1128/AEM.01943-16.

18. Meuskens I, Saragliadis A, Leo JC, Linke D. 2019. Type V Secretion Systems: An

Overview of Passenger Domain Functions. Frontiers in Microbiology 10.

30. Rimbara E, Mori S, Kim H, Suzuki M, Shibayama K. 2018. Mutations in genes

encoding penicillin-binding proteins and efflux pumps play a role in β-lactam resistance in

Helicobacter cinaedi. Antimicrob Agents Chemother 62:e02036-17.

https://doi.org/10.1128/AAC.02036-17.

19. Flahou B, Haesebrouck F, Smet A, Yonezawa H, Osaki T, Kamiya S. 2013. Gastric and

enterohepatic Non-*Helicobacter pylori* Helicobacters. Helicobacter 18 Supplement 1:66–72.

https://doi.org/10.1111/hel.12072.

20. Saito S, Tsukahara M, Ohkusu K, Kurai H. 2016. *Helicobacter fennelliae* bacteremia

Three Case Reports and Literature review. Med (Baltim) 95:e3556.

https://doi.org/10.1097/MD.0000000000003556.

21. Gotoh Y, Atsuta Y, Taniguchi T, Nishida R, Nakamura K, Ogura Y, Misawa N, Hayashi

T 2022. *Helicobacter cinaedi* is a human-adapted lineage in the *Helicobacter*

cinaedi/canicola/'magdeburgensis' complex. Microbial Genomics 8.

https://doi.org/10.1099/mgen.0.000830.

22. Taniguchi T, Yamazaki W, Saeki Y, Takajo I, Okayama A, Hayashi T, Misawa N. 2016.

The pathogenic potential of *Helicobacter cinaedi* isolated from non-human sources:

Adherence, invasion and translocation ability in polarized intestinal epithelial Caco-2 cells in

vitro. 4. J Vet Med Sci 78:627–632. https://doi.org/10.1292/jvms.15-0595.

23. Matsumoto T, Goto M, Murakami H, Tanaka T, Nishiyama H, Ono E, Okada C, Sawabe

E, Yagoshi M, Yoneyama A, Okuzumi K, Tateda K, Misawa N, Yamaguchi K. 2007.

Multicenter study to evaluate bloodstream infection by *Helicobacter cinaedi* in Japan. J Clin

Microbiol 45:2853–2857. https://doi.org/10.1128/JCM.00465-07.

25. Palframan SL, Kwok T, Gabriel K. 2012. Vacuolating cytotoxin A (VacA), a key toxin

for *Helicobacter pylori* pathogenesis. Front Cell Infect Microbiol 2:92.

https://doi.org/10.3389/fcimb.2012.00092.

25. Dautin N. 2010. Serine protease autotransporters of *Enterobacteriaceae* (SPATEs):

Biogenesis and Function. Toxins 2:1179–1206. https://doi.org/10.3390/toxins2061179.

26. D'Souza SE, Ginsberg MH, Plow EF. 1991. Arginyl-glycyl-aspartic acid (RGD): A cell

adhesion motif. Trends Biochem Sci 16:246–250. https://doi.org/10.1016/0968-

0004(91)90096-E.

27. Radin JN, Gaddy JA, González-Rivera C, Loh JT, Algood HMS, Cover TL. 2013.

Flagellar localization of a *Helicobacter pylori* autotransporter protein. mBio 4:e00613-e00612.

https://doi.org/10.1128/mBio.00613-12.

28. Sause WE, Castillo AR, Ottemann KM. 2012. The *Helicobacter pylori* autotransporter

ImaA (HP0289) modulates the immune response and contributes to host colonization. Infect

Immun 80:2286–2296. https://doi.org/10.1128/IAI.00312-12.

29. Guignot J, Segura A, Nhieu GTV. 2015. The serine protease EspC from

enteropathogenic *Escherichia coli* regulates pore formation and cytotoxicity mediated by the

Type   III   secretion   system.   PLOS   Pathog   11:e1005013.

https://doi.org/10.1371/journal.ppat.1005013.

31. Shibayama K, Takeuchi H, Wachino J, Mori S, Arakawa Y. 2011. Biochemical and

pathophysiological characterization of *Helicobacter pylori* asparaginase. Microbiol Immunol

55:408–417. https://doi.org/10.1111/j.1348-0421.2011.00333.x.

32. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H,

Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG. 2007.

Clustal   W   and   Clustal   X   version   2.0.   Bioinformatics   23:2947–2948.

https://doi.org/10.1093/bioinformatics/btm404.

33. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular

evolutionary genetics analysis across computing platforms. Mol Biol Evol 35:1547–1549.

https://doi.org/10.1093/molbev/msy096.

34. Matsui H, Suzuki M, Isshiki Y, Kodama C, Eguchi M, Kikuchi Y, Motokawa K, Takaya

A, Tomoyasu T, Yamamoto T. 2003. Oral immunization with ATP-dependent protease-

deficient mutants protects mice against subsequent oral challenge with virulent *Salmonella*

enterica serovar Typhimurium. Infect Immun 71:30–39. https://doi.org/10.1128/IAI.71.1.30-

39.2003.

 Table and figure legends

Table 1. Number of *H. cinaedi* isolated from tissues of *H. cinaedi*-infected mice

		Bacterial count (CFU g^{-1}) by infection duration											
		MRY08-1234 WT			MRY08-1234 KO			MRY12-0027 WT			MRY12-0027 KO		
Tissue[#]		7 d	14 d	28 d	7 d	14 d	28 d	7 d	14 d	28 d	7 d	14 d	28 d
Cecum													
	1	0	2.0×10^3	0	0	0	0	0	0	0	0	0	0
	2	0	0	0	0	0	0	9.5×10^5	0	0	0	0	1.5×10^3
	3	1.5×10^5	1.0×10^3	1.0×10^3	0	0	0	0	1.0×10^3	0	0	0	0
	4	0	1.0×10^3	0	0	0	0	1.5×10^5	0	0	0	0	0
	5	0	1.5×10^5	2.0×10^3	0	0	1.5×10^3	0	2.0×10^3	0	0	0	0
	6	9.5×10^5	-	-	0	-	-	-	-	-	-	-	-
Colon													
	1	0	0	2.5×10^3	0	0	1.0×10^3	0	0	9.0×10^3	0	0	1.0×10^3
	2	0	2.5×10^3	2.0×10^3	0	0	2.0×10^2	0	0	0	0	0	2.0×10^2
	3	3.0×10^3	0	2.0×10^3	0	0	0	5.0×10^2	0	0	0	0	0
	4	0	0	0	0	0	0	3.0×10^3	2.5×10^3	0	0	0	0
	5	5.0×10^2	0	9.5×10^3	0	0	0	0	0	0	0	0	0
	6	0	-	-	0	-	-	-	-	-	-	-	-

Table 1. (Continued.)

Tissue	MRY08-1234 WT			MRY08-1234 KO			MRY12-0027 WT			MRY12-0027 KO		
	\multicolumn Bacterial count (CFU g^{-1}) by infection duration											
	7 d	14 d	28 d	7 d	14 d	28 d	7 d	14 d	28 d	7 d	14 d	28 d
Biceps femoris muscle												
1	0	0	0	0	0	0	0	0	0	0	0	0
2	0	0	2.0×10^2	0	5.0×10^2	1.0×10^1	0	0	0	0	0	0
3	0	0	0	0	0	0	0	0	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0	0	0
6	0	-	-	0	-	-	-	-	-	-	-	-
Trapezius muscle												
1	0	0	0	0	0	0	0	0	0	0	0	0
2	0	0	0	0	0	0	0	0	0	0	0	0
3	0	5.0×10^2	0	0	0	0	0	5.0×10^2	0	0	0	0
4	0	0	0	0	0	0	0	0	0	0	0	0
5	0	0	0	0	0	0	0	0	0	0	0	0
6	0	-	-	0	-	-	-	-	-	-	-	-

592 [#]Cecal contents of all mice were cultured, and *H. cinaedi* was not isolated. WT, wild type; KO, HcaA-knockout

Figure legends

Fig. 1. Construction of *hcaA*-knockout strains.

(A) HcaA contains features of putative autotransporter proteins. The three-dimensional structure of HcaA predicted by AlphaFold2 is color-coded by pLDDT values, which indicate the confidence level of the prediction. (B) Schematic diagram of the knockout strategy for the *hcaA* gene. The *hcaA* locus on *H. cinaedi* MRY12-0027 and MRY08-1234 chromosomes, the donor fragment used to disrupt the *hcaA* gene locus, and the predicted *hcaA* gene locus after homologous recombination are shown. (C, D) Confirmation of *hcaA* gene knockout. (C) PCR using genomic DNA from wild-type (MRY08-1234 WT and MRY12-0027 WT) and HcaA-knockout strains (MRY08-1234 Δ*hcaA* and MRY12-0027 Δ*hcaA*) as templates. Lane 1, MRY12-0027 WT; Lane 2, MRY12-0027 Δ*hcaA*; Lane 3, MRY08-1234 WT; Lane 4, MRY08-1234 Δ*hcaA*; Lane 5, OneSTEP Marker 6 (λ/Sty I digest) (Nippon Gene Co., Ltd.). White arrows indicate amplicons of the *hcaA* gene (5,478 bp). Absence of the amplicons in HcaA-knockout strains was confirmed. (D) Immunoblot of HcaA. The concentration of the extracted protein was measured using the BCA protein assay. The samples were adjusted to a concentration of 0.5 mg/mL with PBS and separated by 5–20% sodium dodecyl sulfate-polyacrylamide gel electrophoresis. The top and bottom panels show results for MRY08-1234 and MRY12-0027, respectively. Lane M, Precision Plus Protein Dual Color Standards (Bio-Rad); Lane 1 and 5, whole cell lysate; Lane 2 and 6, culture supernatant fraction; Lane 3 and 7, membrane fraction; Lane 4 and 8, soluble fraction. The expected size of HcaA was 150 kDa, and a band corresponding to 150 kDa was observed in the WT (lanes 1 and 3) but not in the ΔHcaA strain (lanes 5 and 7), confirming the knockout and localization of HcaA.

Fig. 2. Knockout of HcaA reduced *H. cinaedi* cytotoxicity and adherence to cells.

(A) LDH assay to evaluate cytotoxicity. Each cell was infected with wild-type (MRY08-

1234 WT or MRY12-0027 WT) and HcaA-knockout (MRY08-1234 ΔHcaA or MRY12-0027

ΔHcaA) strains for 24 h (MOI 100). Cytotoxicity (%) was calculated using the method

described in the instructions. (B), (C), and (D): Adhesion and invasion assays against Caco-2

cells. Caco-2 cells were infected with wild-type (MRY08-1234 WT or MRY12-0027 WT)

and HcaA knockout (MRY08-1234 ΔHcaA or MRY12-0027 ΔHcaA) strains for 4 h (MOI

100). The adhesion rate (B) and invasion rate (C) were calculated by dividing the number of

adherent bacterial cells and invading bacterial cells, respectively, by the total number of

infected bacterial cells. The invasion rate of adherent bacteria (D) was calculated by dividing

the number of invading bacteria cells by the number of adherent bacteria cells. Results are

presented as the mean ± standard deviation (SD) of six independent experiments. *,

$P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$ ****, $P < 0.0001$; ns, not specified.

Fig. 3. RGD motif on HcaA contributes to the adhesion of *H. cinaedi*.

(A) SDS-PAGE of the purified WT HcaA and mutant HcaA, stained with Coomassie Brilliant

Blue (CBB). HcaA (120 kDa) with the RGD motif (WT HcaA) and HcaA with glycine

replaced with alanine in the RGD motif (Mutant HcaA) were expressed in *E. coli* and purified.

Proteins were adjusted to 1 mg/mL and applied. (B) Adhesion of purified WT HcaA (with the

RGD motif) and Mutant HcaA (with the RAD motif) to U937 cells. Fluorescence-labeled

U937 cells attached to HcaA were compared by measuring the fluorescence at 490 nm. Means

and SD reflect values of six per group. **, $P < 0.01$. (C) Three-dimensional structure of

HcaA and HcaA homologues predicted by AlphaFold2. Of the five models generated, the top-

ranked model was used in this study. The predicted structure was visualized using PyMOL

Molecular Graphics System Ver. 2.2.5. HcaA homologs from *H. bilis*, *H. fennelliae*, *H.*

canicola, and *H. canis* were analyzed and compared. The RGD motif is located at the

protrusion in HcaA (within the square). HcaA homologs had a structure protruding from the

surface, like HcaA; however, the protruding structure did not contain the RGD motif. (D)

Phylogenetic tree based on the alignment of the passenger domain of HcaA and HcaA

homologs. The alignment was generated using ClustalW ver. 2.1. The maximum-likelihood

tree was constructed using RAxML (https://github.com/amkozlov/raxml-ng) and visualized

using MEGA-X v 10.2.6. A comparison of the amino acid sequence of HcaA showed that the

RGD motif was present only in HcaA from *H. cinaedi* (red text).

Fig. 4. Knockout of HcaA reduced *H. cinaedi* colonization in mice.

(A) Reactivity to *H. cinaedi* in infected mice, as determined by ELISA. Sera from mice at 7,

14, and 28 days after inoculation were used. Squares and triangles indicate serum from the

group of infected wild-type (black square, MRY08-1234 WT; gray square, MRY12-0027

WT) or HcaA knockout (black triangle, MRY08-1234 ΔHcaA; gray triangle, MRY12-0027

ΔHcaA) strains, respectively. Means and SD reflect values from five of six mice per group.

**, $P < 0.01$. (B) Heatmap combining the results of the culture method and qPCR assays.

Columns indicate each mouse and rows indicate the duration of infection. Positive results for

both culture method and qPCR assays are shown in red, positive results for either are shown

in pink, and negative results for both are shown in light blue. (C) Quantitative bacteria count

per length of colon and representative images of the colon of *H. cinaedi*-infected mice.

Adhesion of *H. cinaedi* to the colon was quantified using ImageJ ver. 1.53t, by counting the

number of *H. cinaedi* in images taken with a 20× objective lens. Results are expressed as the

number of *H. cinaedi* per length of colon. Means and SD reflect the values of five or six mice

per group. **, $P < 0.01$. Green, *H. cinaedi*; blue, nucleus. Scale bar 100 um.

(A) Cytotoxicity

(B) The adhesion rate

(C) The invasion rate

(D) The invasion rate of adherent bacteria

(A)

kDa	M	WT HcaA	Mutant HcaA

150
100

(B)

(C)

HcaA   HcaA paralogs

RGD   DNNTL   NGSWT   NGSNG   NALLT

H. cinaedi   *H. fennelliae*   *H. canis*   *H. canicola*   *H. bilis*
MRY08-1234   MRY12-0050   CCUG32756ᵀ   PAGU1409   ATCC43879

(D)

Strain	Sequence (position)
H. cinaedi MRY08-1234	³³⁶ W T L N G N N L K F E S **R G D** M P Q E S K T L E Q N K D I I ³⁶⁵
H. cinaedi MRY12-0027	³³⁵ W T L N G N N L K F E S **R G D** M P Q E S K T L E Q N K D I I ³⁶⁴
H. cinaedi CCUG11818ᵀ	³⁴⁰ W T L N N T T L K F E K **R G D** T P D K T Y N L E A N K D L I ³⁶⁹
H. fennelliae MRY12-0050	²⁹² W T L N G N A L D - - - - N N T L S V D Q T I Q G N K D I I ³¹⁷
H. canis CCUG32756ᵀ	³³⁴ W T F D G S G F Q - - - N G S - - - - W T G K Q Q N K D I V ³⁶¹
H. canicola PAGU1409	³⁴⁰ W T F D N S N F - - - - N G S N G N G W V G K Q N N K D L V ³⁶⁵
H. bilis ATCC43879	³³⁶ W T L S N - - - - - - - N A L L T A G W E T L Q G D K D I V ³⁵⁸

68
92
100
63

(A)

PBS / MRY08-1234 WT / MRY08-1234 ΔHcaA

Optical density at 450 nm vs Post infection (days)

PBS / MRY12-0027 WT / MRY12-0027 ΔHcaA

(B)

Cecum   **Colon**

MRY08-1234   MRY12-0027   MRY08-1234   MRY12-0027
WT ΔHcaA   WT ΔHcaA   WT ΔHcaA   WT ΔHcaA

Days: 7, 14, 28

■ Culture and qPCR were both positive   ■ Either culture or qPCR were positive   ■ Culture and qPCR were both negative

(C)

7 days   **14 days**   **28 days**

Count/length of colon

Mock / MRY08-1234 WT / MRY08-1234 ΔHcaA / MRY12-0027 WT / MRY12-0027 ΔHcaA

Mock
MRY08-1234 WT
MRY08-1234 ΔHcaA
MRY12-0027 WT
MRY12-0027 ΔHcaA