Characterization of HcaA, a novel autotransporter protein in *Helicobacter cinaedi*, and

its role in host cell adhesion

Short title: *Helicobacter cinaedi* autotransporter HcaA

Sae Aoki[1], Shigetarou Mori[1], Hidenori Matsui[1], Keigo Shibayama[2], Tsuyoshi Kenri[1], Emiko

Rimbara[1*]

[1]Department of Bacteriology II, National Institute of Infectious Diseases, Tokyo, Japan

[2]Department of Bacteriology, Nagoya University Graduate School of Medicine, Nagoya,

Aichi, Japan

*Corresponding author:

Emiko Rimbara

Department of Bacteriology II, National Institute of Infectious Diseases

4-7-1 Gakuen, Musashimurayama-shi, Tokyo 208-0011, JAPAN

Tel: +81-42-561-0771

Email: rimbara@niid.go.jp

ABSTRACT

Helicobacter cinaedi infects the human gut and causes invasive infections such as

bacteremia and cellulitis by bacterial translocation. However, how *H. cinaedi* attaches to host

cells and establishes infection has not been elucidated. In this study, we focused on a novel

autotransporter protein, i.e., *H. cinaedi* autotransporter protein A (HcaA) to examine its

relationship with *H. cinaedi* pathogenicity. The cytotoxicity of *H. cinaedi* infection in the

colon epithelial cell lines (Caco-2 and HT-29) by lactate dehydrogenase assay showed a

significant reduction in cytotoxicity by HcaA knockout. Adhesion assays further revealed that

the HcaA-knockout strain showed significantly reduced attachment of *H. cinaedi* to the

human epithelial colorectal adenocarcinoma cell line (Caco-2) compared to the wild-type

strain. To determine the role of HcaA in *H. cinaedi* infection *in vivo*, *H. cinaedi* wild-type and

HcaA-knockout strains were orally infected C57BL/6 mice. The colonized bacteria were then

measured 7, 14, and 28-days post-infection. The number of colonized *H. cinaedi* cells was

significantly lower in HcaA-knockout strain infections than in wild-type strain infections at 7

days post-infection. Recombinant HcaA protein showed strong adhesion characteristics to the

human monocytic cell line (U937) by adhesion assay used recombinant purified HcaA protein.

The adherent activity was diminished by the replacement of the RGD motif in HcaA with

RAD, indicating the contribution of the RGD motif in HcaA to host cell adherence. These

results suggest that HcaA, a novel autotransporter protein in *H. cinaedi*, plays a significant

role in establishing infection as an adhesin.

Keywords: *Helicobacter cinaedi*, autotransporter protein, type V secretion system, non-

Helicobacter pylori Helicobacter species, cell adhesion

IMPORTANCE

Helicobacter species are classified as gastric or enterohepatic according to their habitat.

Enterohepatic *Helicobacter* species inhabit the intestine, colon and liver; among them, *H.*

cinaedi has been most frequently isolated from humans. *H. cinaedi* commonly causes

bacteremia and cellulitis in immunocompromised hosts. Here, we focused on the *H. cinaedi*

autotransporter protein A (HcaA), a novel virulence factor in *H. cinaedi*. We discovered that

HcaA contributed to cell adhesion via its RGD motif. Furthermore, in animal experiments, the

number of colonized bacteria was decreased in HcaA knockout strain infections, supporting

the hypothesis that HcaA contributes to the adhesion of *H. cinaedi* to host cells. Our study

provides a novel mechanism for the establishment of *H. cinaedi* infections and is expected to

provide new insights into the role of autotransporter proteins in the establishment of

Helicobacter species infection.

INTRODUCTION

Helicobacter species, gram-negative and helical bacilli, are isolated from humans and

mammals such as dogs, cats, pigs, and birds. They are grouped into gastric or enterohepatic

Helicobacter species according to their habitat. *Helicobacter pylori*, a typical gastric

Helicobacter species, is a pathogenic bacterium that infects the human stomach and is a major

cause of gastric cancer (1).

Among enterohepatic *Helicobacter* species*, H. cinaedi* is the most frequently isolated

strain from humans. This strain is mainly isolated from patients with bacteremia in

immunocompromised hosts. *H. cinaedi* is known to cause recurrent cellulitis, and in recent

years, it has been reported to cause an infected aortic aneurysm, subdural hematoma, and

hepatic cyst infections (2–6). The first isolation of *H. cinaedi* was reported in a homosexual

man with intestinal symptoms in Seattle in 1984, and the first isolation in Japan was reported

in 2003 (7,8). For unknown reasons, *H. cinaedi* infection is mainly reported in Japan,

followed by several case reports from Denmark (9,10). A retrospective study in a single

hospital in Japan indicated that, of all blood culture-positive samples, 2.2% (126/5,769 sets of

blood cultures) became *H. cinaedi*-positive when the period of blood culture was extended,

indicating that *H. cinaedi* infections may be overlooked in other countries (11).

Cytolethal distending toxin (CDT) is a virulence factor in *H. cinaedi*. CDT is composed of

three polypeptide subunits encoded by CdtA, CdtB and CdtC (12,13). Among these proteins,

CdtB has function of nuclease activity and cause DNA damage. As a result, the host cells

were led to cell senescence and apoptosis. CdtA and CdtC play a role in delivering CdtB into

host cells. Another virulence factor *H. cinaedi* antigen CAIP contributes atherosclerosis

inflammation by the differentiation and maintenance of the pro-inflammatory profile of

human macrophages and triggers the formation of foam cells (14). However, the mechanism

by which *H. cinaedi* establishes infection has not been fully elucidated.

Autotransporter proteins (ATs) are type V secretion systems (T5SS) that consist of a

passenger domain, which plays a significant role, a functional secreted portion, and an

autotransporter domain involving a β-barrel, which exports the passenger domain from the

periplasm to the extracellular environment. ATs are widely found in gram-negative bacteria

and have diverse functions in their pathogenesis, acting as toxins and adhesion factors (15).

For example, the vacuolating cytotoxin A (VacA) of *H. pylori* is an ATs that acts as a toxin

contributing *to H. pylori* pathogenesis by inducing vacuolation in the host cells, altering

mitochondrial membrane permeability and apoptosis, and stimulation of pro-inflammatory

signaling (16). *H. pylori* possesses three other autotransporter proteins, ImaA, VlpC, and

FaaA, which contribute to establishing infection and motility (17). In *H. cinaedi*, one

autotransporter protein is encoded in its chromosome, which has not previously been

characterized. In this study, we named this uncharacterized protein *H. cinaedi* autotransporter

protein A (HcaA) and investigated its contribution to *H. cinaedi* pathogenicity.

RESULTS

HcaA knockout strain showed reduced cytotoxicity and adherence and did not invade

human colon epithelial cells *in vitro*

We constructed deletion knockout mutants of *hcaA* gene. HcaA is a 197 kDa protein

encoded by the 5,478 bp *hcaA* gene. A donor fragment containing an antibiotic selection

marker (*aphA* gene) flanked by a paralog sequence approximately 1,000 bp upstream and

downstream of *hcaA* was created (Fig. 1A, upper and middle). *H. cinaedi* MRY12-0027 and

MRY08-1234 strains were used as host strains, and *hcaA* was knocked out by replacing *hcaA*

with *aphA* by paralog recombination (Fig. 1A, bottom). Knockout of *hcaA* was verified by

polymerase chain reaction (PCR) amplification and immunoblot analysis (Fig. 1B and C). The

results of immunoblot analysis suggested that HcaA was localized in the membrane fraction

(Fig. 1C).

The effects of the HcaA-knockout strain on cellular cytotoxicity were evaluated *in vitro*

using a lactate dehydrogenase (LDH) assay. When the human epithelial colorectal

adenocarcinoma (Caco-2) cell line and human colon cancer (HT29) cell line were infected

with *H. cinaedi*, LDH production in cells infected with the HcaA-knockout strain was

significantly lower than that of the wild-type strain (Fig. 2A). Therefore, HcaA may play a

significant role in the cytotoxic effects of *H. cinaedi* infection on the epithelial cells. HcaA

knockout did not affect LDH production when mucus-secreting HT29-MTX cells were used

(Fig. 2A). This suggests mucin's protective role in reducing the cytotoxicity caused by *H.*

cinaedi was not affected by the knockout of HcaA.

To determine whether HcaA contributes to cell adhesion and invasion, we performed a

gentamicin assay, and compared the rate of adherents and invaded *H. cinaedi* cells between

the wild-type and knockout strains. The adhesion rate of *H cinaedi* MRY12-0027 strain to

Caco-2 cells was significantly decreased by HcaA knockout (Fig. 2B, upper panel). A

significant reduction in the adhesion rate was also observed when *H. cinaedi* MRY08-1234

was used (Fig. 2B, bottom panel). The invasion rate of MRY12-0027 and MRY08-1234 was

also reduced by HcaA knockout. However, no significant difference was observed (Fig. 2C).

In contrast, the invasion rate of adherent bacteria was higher in the HcaA-knockout strain (Fig.

2D).

HcaA effects on the establishment of *H. cinaedi* infection in mice

To clarify the role of HcaA in *H. cinaedi* infection *in vivo*, C57BL/6 female mice were

orally infected with *H. cinaedi* wild-type and HcaA-knockout strains. The recovery of *H.*

cinaedi from cecal contents, colon, biceps femoris muscle, trapezius muscle, and heart was

analyzed at 7, 14, and 28 days post-infection (Fig. 3A). As *H. cinaedi* is known as a causative

agent of cellulitis and infected aortic aneurysm (18), muscle and heart samples were also

taken in addition to the intestinal tract samples. To confirm the infection, antibody titers

against *H. cinaedi* in mouse serum were measured. An increase in antibody titers against

whole cells of *H. cinaedi* was observed in both groups of mice infected with the wild-type

strain and HcaA knockout strain. The antibody titers against *H. cinaedi* at 28 days post-

infection were significantly higher in the group of mice infected with the wild-type strain than

in the mice infected with the knockout strain ($P < 0.01$, Fig. 3B). Using the culture method, *H.*

cinaedi was recovered from the colon and cecal contents in both groups of mice infected with

the wild-type strain and HcaA-knockout strain. The recovery rate of *H. cinaedi* in cecal

contents was significantly higher in mice infected with the wild-type strain (4/5, 80%) than in

mice infected with the HcaA-knockout strain at 7 days post-infection (0/6, 0%, $P < 0.01$,

Table 1). Interestingly, *H. cinaedi* was detected in muscle only in the group of mice infected

with the wild-type strain at 7 and 28 days post-infection (Table 1). *H. cinaedi* was detected in

the cecal contents of all *H. cinaedi*-infected mice by quantitative real-time PCR (qPCR).

Bacterial cells in the cecal contents from mice infected with the HcaA-knockout strain were

significantly fewer than those from mice infected with the wild-type strain at 7 days post-

infection ($P < 0.05$, Fig. 3C). At 14 and 28 days post-infection, there was a trend toward fewer

bacterial cells in mice infected with the HcaA-knockout strain. However, no significant

differences were observed between groups ($P = 0.15$ and 0.42, respectively). The reduction in

the number of HcaA-knockout bacterial cells was also observed through immunofluorescence

analysis of the colon infected with *H. cinaedi* using an anti-*H. cinaedi* antibody (Fig. 3D). *H.*

cinaedi was observed in the mucosal epithelium regardless of the presence or absence of

HcaA and the duration of infection; however, the number of bacteria tended to be lower in

samples from mice infected with the HcaA-knockout strain than in samples from mice

infected with the wild-type strain.

Adhesive ability was reduced in the mutant HcaA recombinant protein

The HcaA protein of *H. cinaedi* MRY 08-1234 [1,835 amino acids (aa), 197 kDa] is

predicted to be structured with a signal peptide (1–27 aa), passenger domain (27–1425 aa),

and β-barrel domain (1426–1825 aa). HcaA of *H. cinaedi* contains an Arg–Gly–Asp (RGD)

motif at amino acid positions 348–350. To understand the contribution of the RGD motif to

adhesion, a recombinant protein of the passenger region of HcaA of MRY08-1234 (WT

HcaA) and mutant HcaA (with the RGD motif replaced by RAD) was created (Fig. 4A). The

ability of the cells to adhere to U937 cells was compared between WT HcaA and mutant

HcaA. Mutant HcaA revealed a significant reduction in binding to U937 cells compared to

WT HcaA (*P* < 0.05), indicating the role of the RGD motif in binding to host cells (Fig. 4B).

Structural prediction of HcaA of MRY08-1234 by AlphaFold2 indicated that the RGD motif

of HcaA is located on the surface of the protein, supporting the contribution of the RGD motif

to the adherence function of HcaA in *H. cinaedi* (Fig. 4C, left). The RGD motif was

conserved in all *H. cinaedi* strains registered at the National Center for Biotechnology

Information (NCBI), including clinical isolates (Fig. S1). HcaA paralogs have been in *H. bilis*,

H. fennelliae, *H. canicola*, and *H. canis*, which are all related to *H. cinaedi* (19–21). The

structural predictions of HcaA paralogs by AlphaFold2 suggested that HcaA and its paralogs

had structural similarities in the passenger region; however, the RGD motif was present only

in HcaA of *H. cinaedi* (Fig. 4C and D). In addition, the HcaA paralogs had a structure

protruding from the surface, similar to the RGD motif of HcaA. However, the protruding

structure did not contain the RGD motif (Fig. 4C).

DISCUSSION

Enterohepatic *Helicobacter* species are increasingly being recognized as critical

mammalian pathogens. Among these, *H. cinaedi* has been reported to cause bacteremia,

cellulitis, and infected aortic aneurysms in humans (2,4,22). Although *H. cinaedi* infections

have rarely been reported, *H. cinaedi* infection cases in Japan have been increasing since 2003

(8,11,22). *H. cinaedi* infections have been reported to show diverse pathologies; however,

knowledge concerning the virulence factors of *H. cinaedi* is limited. ATs have various

functions that contribute to the establishing infections by pathogenic gram-negative bacteria.

In this study, we found that HcaA, the only ATs in *H. cinaedi*, is responsible for adhesion to

host cells and contributes to the pathogenesis of *H. cinaedi*. This is the first report to elucidate

the role of ATs in enterohepatic *Helicobacter* species.

Based on their protein architecture, T5SS proteins are classified into subgroups labelled

Va–Ve; this does not reflect the functions of the secreted passengers. Pertactin produced by

Bordetella pertussis is classified as a type Va ATs; it contributes to adhesion to cell surfaces

and receptors (24). VacA produced by *H. pylori* is likewise classified as a type Va AT; it

binds to host cells and is internalized, causing vacuolation characterized by the accumulation

of large vesicles (16). Serine protease autotransporters of *Enterobacteriaceae* (SPATEs)

produced by *Enterobacteriaceae* are also classified as type Va ATs (23). The effect of

SPATEs on cells varies, with the primary reported function being protease activity that

cleaves host proteins. HcaA is structurally similar to EspP, one of the SPATEs in *Escherichia*

coli. We tested protease activity against casein and pepsin, the major substrates for SPATEs;

however, no protease activity against these proteins was observed (Fig. S2A and B). In

contrast, the number of bacterial cells adhered to human epithelial cells and cytotoxicity

levels were significantly decreased by HcaA knockout. These results indicate that the

reduction in cytotoxicity in the HcaA knockout strain is due to the decrease in the number of

attached *H. cinaedi* cells and that HcaA itself may not be cytotoxic.

The mouse infection experiment also supported the *in vitro* data, showing that HcaA

knockout reduces the number of infected bacterial cells in the cecal contents of *H. cinaedi*-

infected mice at 7 days post-infection. Because the number of bacterial cells was comparable

between the wild-type and HcaA-knockout strains after 28 days of infection, HcaA could be

involved in cell adhesion and may play a significant role in the established phase of *H.*

cinaedi infection. Recovery of *H. cinaedi* from cecal contents and biceps femoris muscle at 7

days post-infection was only observed in mice infected with the wild-type strain, suggesting

that HcaA contributes to the establishment of *H. cinaedi* infection. Further studies are needed

in terms of bacterial translocation, as isolates from muscle suggested that it may be associated

with cellulitis caused by *H. cinaedi*. Interestingly, HcaA knockout did not affect the number

of invaded bacterial cells/total number of infected bacterial cells. In contrast, the number of

invaded bacterial cells/adhered bacterial cells increased with HcaA knockout (Fig. 2D). ATs

have been found to have a variety of functions. For example, ImaA, an AT in *H. pylori,* has

been reported to be an essential factor for host colonization and to reduce host inflammatory

responses by modulating the host inflammatory response by controlling the expression of

inflammatory cytokines IL-8 and TNF-α (25). In addition, EspC, an SPATEs, has been

reported to degrade various host proteins, such as hemoglobin and mucin, and also to prevent

cytotoxicity during the initial stage of bacterial infection by regulating pore formation (23,26).

Thus, HcaA may have a similar function in escaping from the host immune mechanism to

contribute to the establishment of chronic *H. cinaedi* infections in humans. Further analysis is

needed to elucidate additional functions of HcaA other than cell adhesion.

HcaA contains an RGD motif in its passenger region. The RGD motif acts as the

fibronectin cell adhesion site and plays a role in adhesion to host epithelial cells expressing

receptors of the RGD motif, such as integrins. Cell adhesion experiments using purified HcaA

with the RGD motif and mutant HcaA with the RAD motif showed that adhesion ability

decreased significantly upon replacing RGD with RAD. Structural prediction of the HcaA

structure indicated that the RGD motif is located on the surface of the protein and that the

motif is located at an externally protruding site. The RGD motif is known to bind to integrins,

and it is assumed that integrins mediate the adhesion of HcaA to cells. Our results strongly

suggest that the RGD motif of HcaA plays a role in cell adhesion. However, the adhesion

ability of HcaA was not completely diminished by replacing RGD with RAD. Therefore,

HcaA possibly functions as an adhesin regardless of the RGD motif, and the presence of the

RGD motif enhances its ability to attach to cells. HcaA paralogs are also found in

enterohepatic *Helicobacter* species, such as *H. bilis*, *H. canis*, *H. fennelliae,* and *H. canicola.*

However, none of these have an RGD motif (Fig. 4). In addition, the RGD motif of HcaA was

highly conserved among *H. cinaedi* strains, regardless of the origin of the strains (Fig. S1). *H.*

canicola was formerly identified as *H. cinaedi* and is genetically closest to *H. cinaedi* among

Helicobacter species (21). However, phylogenetic analysis of HcaA indicated that HcaA is

closest to the HcaA paralog of *H. fennelliae*, but not that of *H. canicola*. *H. fennelliae* is the

second most prevalent enterohepatic *Helicobacter* species causing invasive infections in

humans. The structural similarity between the HcaA and HcaA paralogs of *H. fennelliae*

suggests that HcaA plays a critical role in the pathogenesis of enterohepatic *Helicobacter*

species in humans. The results of comparative genomic analyses indicate that *H. cinaedi* is a

human-adapted lineage, whereas *H. canicola* is not (21). The acquisition of the RGD motif of

HcaA during the evolution of *H. cinaedi* may have contributed to the adaptation of *H. cinaedi*

to humans, resulting in a higher isolation rate of *H. cinaedi* from humans compared to other

enterohepatic *Helicobacter* species. Further investigation is needed to identify the HcaA-

binding target in the host cell.

This study identified a novel virulence factor of *H. cinaedi*, HcaA, as an adhesin protein.

HcaA plays an essential role in the establishing *H. cinaedi* infection both *in vitro* and *in vivo*

and is expected to provide essential data for understanding the pathogenicity of *H. cinaedi*

and the pathogenesis of *H. cinaedi* infection in the future.

MATERIALS AND METHODS

Condition of bacterial strains and host cells

H. cinaedi (MRY08-1234 and MRY12-0027) strains isolated from the blood of patients

with bacteremia in Japan were used throughout the study. The *H. cinaedi* strain was cultured

as previously described (27). Briefly, strains were subcultured on Brucella agar (Becton,

Dickinson, Franklin Lakes, NJ, USA) with 5% horse blood at 37 °C under microaerobic

conditions with hydrogen generated by the gas replacement method using an anaerobic gas

mixture (H_2, 10%; CO_2, 10%; and N_2, 80%). For liquid culture, bacteria were suspended in

Brucella broth (Becton, Dickinson, Franklin Lakes, NJ) with 10% fetal bovine serum (FBS;

NICHIREI), and the culture was incubated for 2 days at 37 °C under microaerobic conditions

with hydrogen.

Human epithelial colorectal adenocarcinoma cells (Caco-2), human myelomonocyte cell

line (U937), human colon cancer cell line (HT29), and mucus-secreting HT29 cells (HT29-

MX) were used in this study. Caco-2 was maintained in Eagle's minimum essential medium

(EMEM, Sigma Aldrich, USA) containing 2 mM l-glutamine, 1 mM sodium pyruvate, 1.5

mg/L sodium bicarbonate, 10 mg/L phenol red, 10% heat-inactivated (56 °C, 30 min) FBS,

1% non-essential amino acids. U937, HT29, and HT29-MX cells were maintained in RPMI

1640 (Sigma Aldrich) containing 10% heat-inactivated FBS. All media contained an

antimicrobial cocktail (penicillin-streptomycin), except for the *H. cinaedi* infection

experiments. All cells were incubated in a 5% CO_2, humidified atmosphere at 37 °C.

Generation of deletion mutants of *hcaA* gene in *H. cinaedi*

Knockout mutants were generated by paralog recombination according to previously

described methods (27). Briefly, 1,000 base pairs of DNA sequences upstream and

downstream of *hcaA* gene were PCR-amplified. The primers Upstream-R and Downstream-F

contained a linker sequence that overlapped with *aphA* gene. To obtain the construct of *aphA*

sandwiched by the target gene sequence (donor fragment), three purified PCR products, up-

and downstream of *hcaA* and *aphA*, were mixed and amplified using the Upstream-F and

Downstream-R primers. The donor fragment was electroporated into *H. cinaedi* MRY08-1234

or MRY12-0027. The transformed *H. cinaedi* was spread onto agar plates containing 10

µg/mL kanamycin and 3% agar. After incubation, the growing colonies were selected as

knockout mutants. Knockout of *hcaA* was confirmed by PCR and immunoblot analysis.

Primers used for cloning and PCR are listed in Table S1. For the immunoblot analysis, whole

bacterial cell lysates, culture supernatant, soluble fraction, and membrane fraction were

prepared as follows: bacterial culture was incubated until an optical density at 600 nm

(OD600) of 0.4 and the supernatant was collected by centrifugation (8,600 × g, 10 min, 4 °C).

The pellet was resuspended and washed by phosphate-buffered saline (PBS). Then, samples

for the analysis were prepared by extracting proteins from the bacterial cells using,

EzBactYeast Crusher (ATTO Corp., Tokyo, Japan). The concentration of the extracted

protein was measured using the BCA protein assay kit (Thermo Fisher Scientific K.K., Tokyo,

Japan). These samples were adjusted to a concentration of 0.5 mg/mL by PBS and separated

by 5–20% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), and the

separated proteins were transferred to Immobilon[TM] -FL PVDF (pore size, 0.45 um; Millipore

Inc., Burlington, MA, USA) using a semidry electroblotting system (ATTO, Tokyo, Japan).

Anti-HcaA antibody was obtained by immunization of rabbits with KLH-conjugated peptide

(RGDMPQESKTLEQN corresponding to positions 348–361 of HcaA in MRY08-1234),

followed by affinity purification against the peptide. The obtained HcaA rabbit polyclonal

antibody was used as the primary antibody for immunoblot analysis (1:1,000), and peroxidase

AffiniPure donkey anti-mouse IgG (H+L) (Jackson ImmunoResearch Inc., 1:50,000) were

used as secondary antibodies. The bands were visualized using ECL Select (Amersham

Biosciences, Amersham, UK) and Amersham[TM] ImageQuant 800 (Amersham Biosciences).

Cell infection experiments

H. cinaedi strains were grown for three days in Brucella broth with 10% FBS at 37 °C

under microaerobic conditions. After incubation, the bacterial culture was inoculated in 10

mL of fresh broth, and the culture was incubated until an OD600 of 0.4. Bacterial cells were

harvested by centrifugation (3,350 × g, 10 min, 25 °C) and resuspended in a suitable medium

for each cell. Infections were performed at a multiplicity of infection (MOI) of 100.

Immediately after addition of bacteria, the plates were centrifuged for 5 min at $300 \times g$ at

25 °C, followed by incubation in 5% CO_2 humidified atmosphere, at 37 °C for 24 h.

Cytotoxicity was assessed by measuring lactate dehydrogenase (LDH release assay) using

the Cytotoxicity LDH Assay Kit-WST (Dojindo Laboratories, Tokyo, Japan). The assay was

performed according to the manufacturer's instructions. Caco-2, HT29, and HT29-MX cells

were used in this assay.

Cell adhesion and invasive ability were assessed using a gentamicin assay. To evaluate

adhesion ability, cells were liberated by trypsinization, seeded into 24-well microtiter plates

with tissue-culture-treated polystyrene, flat-bottom (FALCON), and incubated for confluency.

Caco-2 cells were infected by adding 1 mL of bacterial suspension (MOI 100) directly to each

well. After 4 h of incubation, the non-bound bacteria were removed by washing three times

with pre-warmed PBS. To evaluate invasive ability, *H. cinaedi* adherent to Caco-2 cells was

incubated in EMEM with gentamicin at 250 µg/mL. After 1.5 h, the infected cells were

washed thrice with pre-warmed PBS. In both assays, host cells were washed with PBS

containing 0.1% saponin (FUJIFILM). Serial dilutions of the suspension were plated onto

Brucella agar containing 5% horse blood and 3% agar (Becton Dickinson, Franklin Lakes, NJ,

USA). After incubation, the number of colonies was counted, and the adhesion rate (%) and

invasion rate (%) were calculated by dividing the number of adherents and invading bacterial

cells by the total number of infected bacterial cells, respectively. The invasion rate of adherent

bacteria (%) was calculated by dividing the number of invading bacterial cells by the number

of adherent bacterial cells.

Animal experiment designs and infection

Six-week-old C57BL/6 female mice maintained under specific pathogen-free conditions

(Japan SLC, Inc., Shizuoka, Japan) were used in this study. The mice were fed a standard

rodent diet, provided with water ad libitum, housed in microisolator cages, and isolated from

specific pathogens.

Strains were suspended into PBS became 1×10^9 cells/mL. Then, mice were orally

administered 0.2 mL of each bacterial suspension by a 1 mL syringe attached to an oral sonde,

every other day for total of three times. As a negative control, the mice were administered the

same volume of PBS. The experiment used 14 mice in the control group, 15 mice in the wild-

type *H. cinaedi* MRY08-1234 infection group, and 16 mice in the *hcaA*-knockout strain

infection group. They were kept in cages with five or six mice.

On days 7, 14, and 28 after inoculation, five or six mice from each group were sacrificed

for analysis. Serum, cecal content, colon, biceps femoris muscle, trapezius muscle, and heart

were collected aseptically for further analysis. All animal experiments and the study protocol

were conducted with the approval of the Committee for Animal Experimentation of the

National Institute of Infectious Diseases (Acceptance number 121126).

ELISA assay

To measure antibody titers against *H. cinaedi* in mouse serum, an enzyme-linked

immunosorbent assay (ELISA) was performed as previously described (28). Serum was

prepared as follows: whole blood collected during autopsy was incubated at 4 °C and then

centrifuged at 2,000 × *g* for 5 min at 4 °C. The supernatant was collected and designated as

serum. Serum was mixed with glycerol (50% final concentration) and stored at −20 °C until

use. To prepare whole-cell lysates, *H. cinaedi* MRY08-1234 cells were collected, washed

twice, and suspended in PBS. The cells were then disrupted by sonication with a Bioruptor II

(BMBio, Tokyo, Japan). Each well of Nunc-Immuno 96-well microtiter plates (No. 439454;

Thermo Fisher Scientific, Waltham, MA, USA) was coated with 100 µl of whole-cell lysates

of *H. cinaedi* MRY08-1234 (4 ug/mL in 0.1 M carbonate/bicarbonate buffer, pH 9.4). After

overnight incubation at 4°C, plate washed three times with PBS containing 0.05% (vol/vol)

Tween 20 (PBS-T), the wells were saturated with 200 µL of blocking buffer (PBS containing

1% BSA) and incubated for 1 h at 37°C while shaking at 500 rpm. After the three times of

washing with PBS-T, the wells were filled with 50 µL of serum samples were diluted 100-

fold with a blocking buffer (PBS containing 1% BSA), and the plates were incubated for 1 h

at 37 °C while shaking at 500 rpm. Horseradish peroxidase (HRP)-conjugated goat anti-

mouse IgG secondary antibody (Jackson, Bar Harbor, ME, USA) diluted 5,000-fold in

blocking buffer was used. Absorbance at 450 nm (reference wavelength, 620 nm) was

measured using a Multiskan FC microplate reader (Thermo Scientific, Waltham, MA, USA).

Isolation and detection of *H. cinaedi* from infected mice

Culture and PCR were used to evaluate *H. cinaedi* infection in each organ. For cultivation

of *H. cinaedi*, agar plates containing Skirrow (Oxoid), 5% horse blood, 3% agar, and 4 µg/mL

ciprofloxacin (FUJIFILM Wako Pure Chemical Co.) were inoculated with cecal contents,

colon, biceps femoris muscle, trapezius muscle, and heart that had been homogenized in 300

µL of Brucella broth. Since MRY08-1234 is resistant to ciprofloxacin, ciprofloxacin was

added to eliminate indigenous intestinal bacteria in mice and isolate only *H. cinaedi*. The

plates were incubated at 37 °C under microaerobic conditions for 7 days. The growth of the

colonies was confirmed by the detection of *cdt* genes using PCR. The primers used are listed

in Table S2. Two, we assessed using genome DNA (gDNA) of *H. cinaedi*. gDNA was

extracted using the QIAamp PowerFecal Pro DNA Kit (QIAGEN) for cecal contents, and the

remaining tissue was extracted using the DNeasy Blood and Tissue Kit (QIAGEN). The

extracted genomic DNA was used for quantitative real-time PCR (qPCR). qPCR was

performed using THUNDERBIRD^{TM} SYBR qPCR MIX (Toyobo) and Applied Biosystems$^{®}$

7500Fast (Thermo Fisher Scientific). The primers used are listed in Table S2. To quantify

bacterial colonization, correlation curves between Ct and $\log_{10}$ CFU/mL were generated. This

correlation was used to transform Ct values into $\log_{10}$ CFU/g of cecal content.

Histology, microscopy, and Fluorescence intensity quantification

Colons from two mice in each group were collected and fixed with a 4% paraformaldehyde

phosphate buffer solution (FUJIFILM Wako Pure Chemical Co.). The sections obtained from

fixed samples were permeabilized and incubated with anti-*H. cinaedi* antibody (1:1,000) as

the first antibody at room temperature for 1 h. Anti-*H. cinaedi* antibody was obtained by

immunizing rabbits with heat-inactivated *H. cinaedi* MRY08-1234 as an antigen, followed by

protein A-purification. The sections were then rinsed and stained with Alexa Fluor 488 (*H.*

cinaedi) and 4',6-diamidino-2-phenylindole (DAPI, nuclei). The fluorescence signals were

visualized using a laser scanning confocal microscope (Olympus FV3000, Olympus, Tokyo,

Japan).

HcaA purification

To create recombinant HcaA, we constructed expression vectors: *hcaA* was amplified by

PCR and introduced into pCold™ I DNA (TaKaRa, Shiga, Japan) using the In-Fusion® HD

Cloning Kit. The primers used to generate the fragments are listed in Table S1. Genomic

DNA of the *H. cinaedi* strain MRY08-1234 was used to amplify the partial *hcaA* gene. The

expression vector containing the partial *hcaA* gene was transferred into *Escherichia coli* BL21

(DE3) pLysS cells and spread on LB agar plates containing 100 µg/mL ampicillin. The

growing colonies were inoculated into 100 mL of LB broth containing 100 µg/mL ampicillin

and incubated at 37 °C with constant shaking at 200 rpm for 3 h. Cells were further

transferred into 900 mL of fresh medium grown in a 2 L flask at 37 °C until OD600 of 0.6.

Protein expression was induced by cooling in ice water for 1 h and adding 400 µL of 1 mM

isopropyl β-d-thiogalactopyranoside (IPTG). Recombinant HcaA was expressed as a fusion

protein containing a histidine tag (His). The cell cultures were further incubated for overnight

at 15 °C. Bacteria were harvested by centrifugation at 5,000 × g for 10 min at 4 °C and

resuspended in 10 mL of 20 mM Tris-HCl (Nakarai), 150 mM NaCl (FUJIFILM Wako Pure

Chemical Co.), and 40 mM imidazole (FUJIFILM Wako Pure Chemical Co.) at pH 7.6 with

protease inhibitor. Suspended solution was sonicated on ice and supernatant was harvested by

centrifugation at 5,000 × g for 45 min at 4 °C. The recombinant protein was purified in two

steps using an affinity column (HisTrap FF crude, Cytiva), followed by size exclusion

chromatography (Superdex™ 200 Increase 10/300 GL, Cytiva). To replace the RGD motif of

the HcaA protein with RAD, site-directed mutagenesis of the HcaA expression vector was

performed using PrimeSTAR® GXL DNA Polymerase and specific primers (Table S1).

Purified wild-type and mutant HcaA proteins were verified using SDS-PAGE.

Adhesion assay of HcaA protein to U937 cells

Purified HcaA and mutant HcaA were prepared at a concentration of 50 µg/mL in PBS,

dispensed 100 uL/well into 96-well plates, and fixed by incubation overnight at 4 °C. This

was then washed with pre-warmed PBS; 100 µL of 2% BSA in PBS was added and incubated

again overnight at 4 °C (blocking). U937 cells were incubated until confluence and then

diluted with MEM to a concentration of 1×10^5 cells/mL. Fluorescein (FUJIFILM Wako) was

added to the cell suspension at a final concentration of 1 mg/mL and incubated for 30 min at

37 °C. After washing with MEM, MEM containing 2 mM Ca and 2 mM Mg was added and

allowed to react for 30 min on ice in the dark. The cell suspension, diluted to a concentration

of 5×10^5 /mL, was then dispensed onto plates coated with HcaA and incubated at 37 °C for 1

h. After washing and replacement with phenol red-free MEM medium (Sigma), fluorescence

was measured, and the number of cells bound to purified proteins was observed.

Alignment and phylogenetic analysis of HcaA and HcaA paralogs

Amino acid sequences of HcaA paralogs in other enterohepatic *Helicobacter* species were

obtained from the whole genome sequences of *H. bilis* ATCC43879 (Accession No.

KI392040), and H. fennelliae MRY12-0050 (Accession No. BSAD00000000), *and H.*

canicola PAGU1409 (accession no. DRR092854) and *H. canis* CCUG32756 (Accession No.

GCF008693005) was obtained from the National Center for Biotechnology Information

(NCBI). The passenger regions of the amino acid sequences of the HcaA and HcaA paralogs

were aligned using ClustalW ver. 2.1 (29), and visualized using CLC Genomics Workbench

Version 22.0.1 (QIAGEN). A phylogenetic tree using the aligned amino acid sequences of the

passenger domain of HcaA and HcaA paralogs was constructed using RAxML-NG v1.1.0

(https://github.com/amkozlov/raxml-ng) and visualized using MEGA X v 10.2.6 (30).

Prediction of HcaA structure

Three-dimensional structures of the HcaA sequences from *H. cinaedi* MRY08-1234 were

predicted using AlphaFold2 Ver.2.2.2 on the supercomputer system of the Institute for

Chemical Research, Kyoto University, with a full dBs preset. Of the five models generated,

the ranked zero model was used in this study. Visualization of the predicted structure was

performed using PyMOL Molecular Graphics System Ver. 2.2.5.

Statistical analysis

Differences in the recovery of *H. cinaedi* by culture from infected mice were analyzed

using Fisher's exact test. For other experiments, the Mann–Whitney test was used for

comparison. All statistical analyses were performed using GraphPad Prism version 9.4.1 and

were considered statistically significant when the P value was < 0.05.

ACKNOWLEDGMENTS

This work was supported by JSPS KAKENHI Grant Numbers 16K18928, 19K07074 (E.

R.) and 22K15465 (S. A.), the GSK Japan Research Grant 2017 (E. R.) and by AMED

JP20fk0108148 (E. R.).

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

References

1. Polk DB, Peek RM. 2010. *Helicobacter pylori*: gastric cancer and beyond. 6. Nat Rev Cancer 10:403–414. https://doi.org/10.1038/nrc2857.

2. Matsuo T, Mori N, Mizuno A, Sakurai A, Kawai F, Starkey J, Ohkushi D, Abe K, Yamasaki M, Ito J, Yoshino K, Mikami Y, Uehara Y, Furukawa K. 2020. Infected aortic aneurysm caused by *Helicobacter cinaedi*: case series and systematic review of the literature. BMC Infect Dis 20:854. https://doi.org/10.1186/s12879-020-05582-7.

3. Akiyama T, Imamura H, Fukui N, Sakai N. 2021. *Helicobacter cinaedi*-infected chronic subdural hematoma mimicking an expanding hematoma: A case report. Surg Neurol Int 12:288. https://doi.org/10.25259/SNI_387_2021.

4. Kawamura Y, Tomida J, Morita Y, Fujii S, Okamoto T, Akaike T. 2014. Clinical and bacteriological characteristics of *Helicobacter cinaedi* infection. Journal of Infection and Chemotherapy 20:517–526. https://doi.org/10.1016/j.jiac.2014.06.007.

5. Suzuki T, Kutsuna S, Tsuboi M, Ota M, Hayakawa K, Ohmagari N. 2019. *Helicobacter cinaedi* Hepatic Cyst Infection with Bacteremia. Emerg Infect Dis 25:603–604. https://doi.org/10.3201/eid2503.180936.

6. Saito J, Rimbara E, Inaguma S, Hasegawa C, Kamiya S, Mizuno A, Sone Y, Ogawa T, Numata Y, Takahashi S, Asano M. 2022. Determining Infected Aortic Aneurysm Treatment Using Focused Detection of *Helicobacter cinaedi*. Emerg Infect Dis 28:1494–1498. https://doi.org/10.3201/eid2807.212505.

7. Totten PA, Fennell CL, Tenover FC, Wezenberg JM, Perine PL, Stamm WE, Holmes KK. 1985. *Campylobacter cinaedi* (sp. nov.) and *Campylobacter fennelliae* (sp, nov.): Two New *Campylobacter* Species Associated with Enteric Disease in Homosexual Men.

The   Journal   of   Infectious   Diseases   151:131–139.

https://doi.org/10.1093/infdis/151.1.131.

8.   Murakami H, Iwata M, Goto M, Takahashi T, Ono E, Sawabe E, Okuzumi K,

Yamaguchi K, Takahashi T. 2003. Isolation of *Helicobacter cinaedi* from blood of an

immunocompromised patient in Japan. Journal of Infection and Chemotherapy 9:344–

347. https://doi.org/10.1007/s10156-003-0265-3.

9.   Holst H, Andresen K, Blom J, Højlyng N, Kemp M, Krogfelt KA, Christensen JJ. 2008.

A Case of *Helicobacter cinaedi* Bacteraemia in a Previously Healthy Person with

Cellulitis. Open Microbiol J 2:29–31. https://doi.org/10.2174/1874285800802010029.

10.   Nielsen HL, Prag J, Krogfelt KA. 2015. *Helicobacter cinaedi* knee infection after

arthroscopy in an immunocompetent patient. BMJ Case Rep 2015:bcr2014208637.

https://doi.org/10.1136/bcr-2014-208637.

11.   Araoka H, Baba M, Kimura M, Abe M, Inagawa H, Yoneyama A. 2014. Clinical

Characteristics of Bacteremia Caused by *Helicobacter cinaedi* and Time Required for

Blood Cultures To Become Positive. Journal of Clinical Microbiology 52:1519–1522.

12.   Taylor NS, Ge Z, Shen Z, Dewhirst FE, Fox JG. 2003. Cytolethal Distending Toxin: A

Potential Virulence Factor for *Helicobacter cinaedi*. The Journal of Infectious Diseases

188:1892–1897.

13.   Scuron MD, Boesze-Battaglia K, Dlakić M, Shenker BJ. 2016. The Cytolethal

Distending Toxin Contributes to Microbial Virulence and Disease Pathogenesis by

Acting As a Tri-Perditious Toxin. Frontiers in Cellular and Infection Microbiology 6.

14.   D'Elios MM, Vallese F, Capitani N, Benagiano M, Bernardini ML, Rossi M, Rossi GP,

Ferrari M, Baldari CT, Zanotti G, de Bernard M, Codolo G. 2017. The *Helicobacter*

cinaedi antigen CAIP participates in atherosclerotic inflammation by promoting the

differentiation of macrophages in foam cells. Sci Rep 7:40515.

https://doi.org/10.1038/srep40515.

15. Meuskens I, Saragliadis A, Leo JC, Linke D. 2019. Type V Secretion Systems: An

Overview of Passenger Domain Functions. Frontiers in Microbiology 10.

16. Palframan S, Kwok T, Gabriel K. 2012. Vacuolating cytotoxin A (VacA), a key toxin for

Helicobacter pylori pathogenesis. Frontiers in Cellular and Infection Microbiology 2.

17. Radin JN, Gaddy JA, González-Rivera C, Loh JT, Algood HMS, Cover TL. 2013.

Flagellar Localization of a *Helicobacter pylori* Autotransporter Protein. mBio 4:e00613-

12. https://doi.org/10.1128/mBio.00613-12.

18. Taniguchi T, Yamazaki W, Saeki Y, Takajo I, Okayama A, Hayashi T, Misawa N. 2016.

The pathogenic potential of *Helicobacter cinaedi* isolated from non-human sources:

adherence, invasion and translocation ability in polarized intestinal epithelial Caco-2

cells in vitro. J Vet Med Sci 78:627–632. https://doi.org/10.1292/jvms.15-0595.

19. Flahou B, Haesebrouck F, Smet A, Yonezawa H, Osaki T, Kamiya S. 2013. Gastric and

Enterohepatic Non-*Helicobacter pylori* Helicobacters. Helicobacter 18:66–72.

https://doi.org/10.1111/hel.12072.

20. Saito S, Tsukahara M, Ohkusu K, Kurai H. 2016. *Helicobacter fennelliae* Bacteremia.

Medicine (Baltimore) 95:e3556. https://doi.org/10.1097/MD.0000000000003556.

21. Gotoh Y, Atsuta Y, Taniguchi T, Nishida R, Nakamura K, Ogura Y, Misawa N, Hayashi

T 2022. *Helicobacter cinaedi* is a human-adapted lineage in the *Helicobacter*

cinaedi/canicola/'magdeburgensis' complex. Microbial Genomics 8:000830.

https://doi.org/10.1099/mgen.0.000830.

22. Matsumoto T, Goto M, Murakami H, Tanaka T, Nishiyama H, Ono E, Okada C, Sawabe

E, Yagoshi M, Yoneyama A, Okuzumi K, Tateda K, Misawa N, Yamaguchi K. 2007.

Multicenter Study To Evaluate Bloodstream Infection by *Helicobacter cinaedi* in Japan.

J Clin Microbiol 45:2853–2857. https://doi.org/10.1128/JCM.00465-07.

23. Aricò B, Nuti S, Scarlato V, Rappuoli R. 1993. Adhesion of *Bordetella pertussis* to

eukaryotic cells requires a time-dependent export and maturation of filamentous

hemagglutinin. Proceedings of the National Academy of Sciences 90:9204–9208.

https://doi.org/10.1073/pnas.90.19.9204.

24. Dautin N. 2010. Serine Protease Autotransporters of *Enterobacteriaceae* (SPATEs):

Biogenesis and Function. 6. Toxins 2:1179–1206. https://doi.org/10.3390/toxins2061179.

25. Sause WE, Castillo AR, Ottemann KM. 2012. The *Helicobacter pylori* Autotransporter

ImaA (HP0289) Modulates the Immune Response and Contributes to Host Colonization.

Infect Immun 80:2286–2296. https://doi.org/10.1128/IAI.00312-12.

26. Guignot J, Segura A, Nhieu GTV. 2015. The Serine Protease EspC from

Enteropathogenic *Escherichia coli* Regulates Pore Formation and Cytotoxicity Mediated

by the Type III Secretion System. PLOS Pathogens 11:e1005013.

https://doi.org/10.1371/journal.ppat.1005013.

27. Rimbara E, Mori S, Kim H, Suzuki M, Shibayama K. 2018. Mutations in Genes

Encoding Penicillin-Binding Proteins and Efflux Pumps Play a Role in β-Lactam

Resistance in *Helicobacter cinaedi*. Antimicrobial Agents and Chemotherapy

62:e02036-17. https://doi.org/10.1128/AAC.02036-17.

28. Matsui H, Suzuki M, Isshiki Y, Kodama C, Eguchi M, Kikuchi Y, Motokawa K, Takaya

A, Tomoyasu T, Yamamoto T. 2003. Oral Immunization with ATP-Dependent Protease-

Deficient Mutants Protects Mice against Subsequent Oral Challenge with Virulent

Salmonella enterica Serovar Typhimurium. Infection and Immunity 71:30–39.

https://doi.org/10.1128/IAI.71.1.30-39.2003.

29. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H,

Valentin F, Wallace IM, Wilm A, Lopez R, Thompson JD, Gibson TJ, Higgins DG.

2007. Clustal W and Clustal X version 2.0. Bioinformatics 23:2947–2948.

https://doi.org/10.1093/bioinformatics/btm404.

30. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular

Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol 35:1547–

1549. https://doi.org/10.1093/molbev/msy096.

Table and figure legends

Table 1. Recovery of *H. cinaedi* from tissues of *H. cinaedi*-infected mice

Tissue	No. of positive samples/no. of mice tested*					
	Post-infection (days)					
	7		14		28	
	WT	KO	WT	KO	WT	KO
Heart	0/5	0/6	0/5	0/5	0/5	0/5
Biceps femoris muscle	1/5	0/6	0/5	0/5	1/5	0/5
Trapezius muscle	0/5	0/6	0/5	0/5	0/5	0/5
Colon	1/5	0/6	1/5	1/5	1/5	0/5
Cecal contents	4/5[†]	0/6	2/5	1/5	0/5	2/5

* Numerator is the number of mice positive for *H. cinaedi* by culture; denominator is the total number of mice sampled. [†] Significantly different vs. KO by Fisher's extract test (*P* <0.01). WT, wild type; KO, HcaA-knockout

Figure legends

Fig. 1. Construction of *hcaA*-knockout strain.

(A) Schematic diagram of the knockout strategy for the *hcaA* gene. The *hcaA* locus on the

H. cinaedi MRY12-0027 and MRY08-1234 chromosomes, the donor fragment used to disrupt

the *hcaA* gene locus, and the predicted *hcaA* gene locus after homologous recombination are

shown. (B, C) Confirmation of *hcaA* gene knockout. (B) PCR using genomic DNA from wild

type (MRY12-0027 WT and MRY08-1234 WT) and HcaA-knockout (MRY12-0027 Δ*hcaA*

and MRY08-1234 Δ*hcaA*) strains as template. Lane 1, MRY12-0027 WT; Lane 2, MRY12-

0027 *ΔhcaA*; Lane 3, MRY08-1234 WT; Lane 4, MRY08-1234 *ΔhcaA*; Lane 5, OneSTEP

Marker 6 (λ/Sty I digest) (Nippon Gene Co., Ltd.). White arrows indicate amplicons of *hcaA*

gene (5,478 bp). The absence of the amplicons in HcaA-knockout strains was confirmed. (C)

Immunoblot of HcaA. The upper and bottom panels show MRY12-0027 and MRY08-1234,

respectively. Lane M, Precision Plus Protein Dual Color Standards (Bio-Rad); Lane 1 and 5,

whole cell lysate; Lane 2 and 6, culture supernatant fraction; Lane 3 and 7, membrane

fraction; Lane 4 and 8, soluble fraction. The size of HcaA was expected to be 150 kDa, and a

band corresponding to 150 kDa was observed in the WT (lanes 1 and 3) but not in the ΔHcaA

strain (lanes 5 and 7), confirming the knockout and localization of HcaA.

Fig. 2. Knockout of HcaA reduced the cytotoxicity caused by *H. cinaedi* infection and the

adherence of *H. cinaedi* to cells.

(A) LDH assay for evaluating cytotoxicity. Each cell was infected with wild type (MRY12-

0027 WT) and HcaA-knockout (MRY12-0027 ΔHcaA) strains for 24 h (MOI 100).

Cytotoxicity (%) was calculated using the method described in the instructions. The results

are presented as the mean ± standard deviation (n = 6/group). (B), (C) and (D): Adhesion and

invasion assay experiments against Caco-2 cells. Caco-2 infected with wild type (MRY12-

0027 WT or MRY08-1234 WT) and HcaA knockout (MRY12-0027 ΔHcaA or MRY08-1234

ΔHcaA) strains for 4 h (MOI 100). The adhesion rate (B) and invasion rate (C) were

calculated by dividing the number of adherent bacteria cells and invading bacteria cells by the

number of total infected bacteria cells, respectively. Invasion rate of adhering bacteria (D)

was calculated by dividing the number of invading bacteria cells by the number of adherent

bacteria cells. *, $P < 0.05$; ***, $P < 0.001$; ns, not specified.

Fig. 3. Knockout of HcaA reduced *H. cinaedi* colonization in mice.

(A) Schematic of animal experiments. C57BL/6 mice were infected with *H. cinaedi*

MRY08-1234 wild type or MRY08-1234 ΔHcaA strains for 28 days after oral administration.

The bacteria were inoculated with 0.2 mL of each bacterial suspension (10^9 CFU) every three

days for a total of three inoculations. For negative control, the same volume of PBS was

inoculated. Cecal contents, colon, biceps femoris muscle, trapezius muscle, heart, and blood

were collected and analyzed at days 7, 14, and 28 days, based on the day of the last infection.

(B) Measurement of anti-*H. cinaedi* antibody titers in infected mice by ELISA. Sera from

mice at 7, 14 and 28 days after inoculation were used. Black squares and gray triangles

indicate serum from the group of infected wild type (black square, WT) or HcaA knockout

(gray triangle, ΔHcaA) strains, respectively. Means and SD reflect values of five mice per

group. **, $P < 0.01$. (C) CFU was estimated by qPCR using DNA obtained from cecal

contents of mice infected with wild-type or HcaA-knockout *H. cinaedi* for 7, 14, and 28 days

after infection. CFU was calculated the as threshold cycle (Ct) values using a correlation

curve between Ct values and CFU/mL, and $\log_{10}$ CFU/gram of cecal contents was used for the

comparison. Means and SD reflect values of five mice per group. *, $P < 0.05$; ns, not specified.

(D) Representative confocal laser-scanning microscopy images of the colon from *H. cinaedi*-

infected mice. Green, *H. cinaedi*; blue, nucleus. Tissues were imaged on an inverted laser

scanning fluorescent microscope (Olympus) using a 40× objective lens.

Fig. 4. The RGD motif on HcaA contributes to the adhesion of *H. cinaedi*.

(A) SDS-PAGE of the purified WT HcaA and mutant HcaA, stained with Coomassie brilliant

blue (CBB). HcaA (120 kDa) with RGD motif (WT HcaA) and HcaA with glycine replaced

with alanine in the RGD motif (Mutant HcaA) were expressed in *E. coli* and purified. Proteins

were adjusted to 1 mg/mL concentrations and applied. (B) The adhesion ability of purified

HcaA proteins with RGD motif (WT HcaA) and HcaA with RAD motif (Mutant HcaA) to

U937 cells. Fluorescent-labeled U937 cells attached to HcaA proteins were compared by

measuring the fluorescence at 490 nm. **, $P < 0.01$. (C) Phylogenetic tree constructed from

the alignment of the passenger domain of the HcaA and HcaA paralogs. Alignment was

performed using ClustalW ver. 2.1. The maximum-likelihood tree was constructed using

RAxML (https://github.com/amkozlov/raxml-ng) and visualized using MEGA-X v 10.2.6. A

comparison of the amino acid sequence of HcaA around the position on the RGD motif

showed that the RGD motif was present only in HcaA of *H. cinaedi* (red text). (D) Three-

dimensional structure of HcaA and HcaA paralogs predicted by AlphaFold2. HcaA paralogs

of *H. bilis*, *H. fennelliae*, *H. canicola*, and *H. canis* were analyzed and compared. The RGD

motif is located at the protrusion in HcaA (inside the square). HcaA paralogs had a structure

protruding from the surface like HcaA; however, the protruding structure did not contain the

RGD motif.

(A)

(B)

(C)

(A)

Days -9   0   7   14   28

↑ Orally administration (10^9 CFU/mL, 0.2 mL)   ⇧ Autopsy

(B)

○ PBS
■ MRY08-1234 WT
△ MRY08-1234 ΔHcaA

Optical density at 450 nm

**

Post infection (days)

(C)

■ MRY08-1234 WT
△ MRY08-1234 ΔHcaA

log (CFU/gram of cecal contents)

*   ns   ns

Post infection (days)

(D)

7days   14 days   28 days

MRY08-1234 WT

MRY08-1234 ΔHcaA