Cell-free biosynthesis combined with deep learning accelerates de novo-development of antimicrobial peptides

Amir Pandi*[1], David Adam[1,2], Amir Zare[1], Van Tuan Trinh[3], Stefan L. Schaefer[4], Marie Wiegand[5], Björn Klabunde[5], Elizaveta Bobkova[1], Manish Kushwaha[6], Yeganeh Foroughijabbari[1], Peter Braun[2,7], Christoph Spahn[8], Christian Preußer[9,10], Elke Pogge von Strandmann[9,10], Helge B. Bode[8,11,12,13,14], Heiner von Buttlar[2,7], Wilhelm Bertrams[5], Anna Lena Jung[5,15], Frank Abendroth[3], Bernd Schmeck[5,14,15,16,17,18], Gerhard Hummer[4,19], Olalla Vázquez[3,14], and Tobias J. Erb*[1,14]

[1] Department of Biochemistry and Synthetic Metabolism, Max Planck Institute for Terrestrial Microbiology, Marburg, Germany
[2] Bundeswehr Institute of Microbiology, Munich, Germany
[3] Department of Chemistry, Philipps-University Marburg, Germany
[4] Department of Theoretical Biophysics, Max Planck Institute of Biophysics, Frankfurt am Main, Germany
[5] Institute for Lung Research, Universities of Giessen and Marburg Lung Center, Philipps-University Marburg, German Center for Lung Research (DZL), Marburg, Germany
[6] Université Paris-Saclay, INRAe, AgroParisTech, Micalis Institute, Jouy-en-Josas, France
[7] German Center for Infection Research (DZIF), Munich, Germany
[8] Department of Natural Products in Organismic Interactions, Max Planck Institute for Terrestrial Microbiology, Marburg, Germany
[9] Institute for Tumor Immunology, Center for Tumor Biology and Immunology, Philipps-University Marburg, Marburg, Germany
[10] Core Facility Extracellular Vesicles, Center for Tumor Biology and Immunology, Philipps University of Marburg, Marburg, German
[11] Molecular Biotechnology, Department of Biosciences, Goethe University Frankfurt, Frankfurt am Main, Germany
[12] Department of Chemistry, Chemical Biology, Phillips-University Marburg, Germany
[13] Senckenberg Gesellschaft für Naturforschung, Frankfurt, Germany
[14] SYNMIKRO Center of Synthetic Microbiology, Marburg, Germany
[15] Core Facility Flow Cytometry – Bacterial Vesicles, Philipps-University Marburg, Marburg, Germany
[16] Department of Medicine, Pulmonary and Critical Care Medicine, University Medical Center Marburg, Philipps-University Marburg, Marburg, Germany
[17] Institute for Lung Health (ILH), Giessen, Germany
[18] Member of the German Center for Infectious Disease Research (DZIF), Marburg, Germany
[19] Institute for Biophysics, Goethe University Frankfurt, Frankfurt am Main, Germany

* To whom correspondence should be addressed: toerb@mpi-marburg.mpg.de and amir.pandi@mpi-marburg.mpg.de

Abstract

Bioactive peptides are key molecules in health and medicine. Deep learning holds a big promise for the discovery and design of bioactive peptides. Yet, suitable experimental approaches are required to validate candidates in high throughput and at low cost. Here, we established a cell-free protein synthesis (CFPS) pipeline for the rapid and inexpensive production of antimicrobial peptides (AMPs) directly from DNA templates. To validate our platform, we used deep learning to design thousands of AMPs de novo. Using computational methods, we prioritized 500 candidates that we produced and screened with our CFPS pipeline. We identified 30 functional AMPs, which we characterized further through molecular dynamics simulations, antimicrobial activity and toxicity. Notably, six de novo-AMPs feature broad-spectrum activity against multidrug-resistant pathogens and do not develop bacterial resistance. Our work demonstrates the potential of CFPS for production and testing of bioactive peptides within less than 24 hours and <10$ per screen.

Main

According to the world health organization, antimicrobial resistance (AMR) is among the top 10 global health threats[1]. In 2019 alone, multidrug-resistant bacteria including pathogenic *Escherichia coli*, ESKAPE pathogens (*Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, *Enterobacter* spp.), *Streptococcus pneumoniae*, and *Mycobacterium tuberculosis* caused 1.27 million deaths[2]. This number is predicted to reach 10 million annually by 2050[2]. Despite this looming threat, the development of new antimicrobials is lagging behind. While more than 4,000 immuno-oncology compounds were in clinical trials in 2021, only 40 antimicrobials (of which none is active against multi-drug resistant Gram-negative bacteria) were subjected to clinical studies[3], highlighting the urgent need to increase the development of novel antimicrobial compounds.

One promising class of antimicrobial compounds are antimicrobial peptides (AMPs)[4–8]. AMPs are peptides of 12-50 amino acids (AA), which have evolved as part of nature's antimicrobial arsenal of bacteria as well as the innate immune system of multicellular organisms, millions of years before humans started to use antibiotics[4,6,8]. Compared to classical antibiotics, AMPs show decreased resistance development mainly because (i) most AMPs act directly at the cell membrane, (ii) show a relatively high killing rate, and (iii), resistance against AMPs is conferred by rather non-specific mechanisms, which reduces the chances of mutational and/or horizontal gene transfer events[4]. Overall, this makes AMPs interesting candidates for next-generation antimicrobials.

About 5,000 AMPs have been characterized to date, most of which are of natural origin. However, these 5,000 AMPs span only a tiny fraction of the possible solution space that nature could have explored (~20^{30} for a 30 AA AMP). Yet, our ability to discover or develop new AMPs from this *terra incognita* are still limited. One possibility to overcome this challenge is the use of deep learning models, which are increasingly employed in protein and peptide design[9–13]. In these approaches, known as generative deep learning, models "learn" the natural protein sequence landscape in training sets with unlabeled data to propose new-to-nature protein sequences[14]. This approach is distinct from predictive modeling that uses labeled data to predict specific properties (labels) of proteins from their sequence[15]. Generative and predictive deep learning have been recently used for the discovery of novel AMP sequences, which have been subsequently created and validated through chemical synthesis of the individual candidates[6,16–19]. While this proof-of-principle showcased the potential of deep learning in AMP discovery, a broader application of this approach has been limited due to the lack of convenient methods for the production and screening of more AMP candidates in medium to high-throughput.

One possibility to increase the throughput in AMP production is to switch from chemical synthesis to DNA-based bioproduction methods. However, heterologous expression of AMPs in microorganisms, such as *E. coli*, features several disadvantages: (i) it is time- and labor-intensive, (ii) it requires the cloning, production and purification of AMPs from cell cultures, and most importantly, (iii) many (potent) AMP candidates might not be available, as they potentially kill the producer strain upon induction. Cell-free protein synthesis (CFPS) offers a promising solution to these challenges. CFPS systems are in vitro transcription translation (TX-TL) systems that directly use DNA templates for protein biosynthesis[20–22], which allows the production of peptides outside of living cells. Thus, these systems can help overcome potential cellular toxicity effects, and open up the way for the rapid, small-scale production of several hundreds of peptides from linear DNA in parallel.

Here, we combined deep learning and CFPS for de novo-design, rapid production and screening of AMPs at small scale within 24 hours, and less than 10$ per individual AMP production assay

(excluding cost for the DNA fragment). Having explored ~500,000 theoretical sequences, we
screened 500 AMP candidates to identify 30 functional AMPs, which are completely unrelated to
any natural sequences. Notably, six of these AMPs exhibited high antimicrobial activity against
multidrug-resistant pathogens, showed no emergence of resistance and only minimal toxicity on
human cells.

Results

<u>De novo AMP design using deep learning</u>

For de novo-design of AMPs, we adapted two versions of deep generative variational
autoencoders (VAE) from a previous study[23]. Generative VAE are unsupervised learning models,
which take as input only AMP sequences, and comprise an encoder, a latent space, and a
decoder. During model training, the encoder compresses the input sequences into a low-
dimensional space ("latent space"), while the decoder aims at reconstructing sequences from this
latent space (**Fig. 1a**). We first pretrained the VAEs using ~1.5 million peptide sequences from
UniProt as a generic dataset. Second, performed transfer learning on the pretrained VAEs using
a dataset of ~5,000 known AMPs to set up the latent space to be used for de novo AMP generation
(**Methods** and **Supplementary Table 1**).

To reduce the number of AMPs for experimental testing, we set up a method to select potential
candidates according to their predicted bioactivity, i.e., their Minimum Inhibitory Concentration
(MIC). To that end, we established predictive deep learning models that we trained with the
sequence and respective experimental MIC values of ~5,000 known AMPs (sequence-MIC
relationship, **Fig. 1b**). As regressors, we used convolutional neural networks (CNN) and recurrent
neural networks (RNN) (**Methods** and **Supplementary Table 1**).

To identify interesting AMP candidates, we first generated new AMPs by sampling points from
the latent space and subsequently feeding them into the decoder, which yielded peptide
sequences that share the same properties but are novel compared to the training dataset. These
de novo AMPs were then prioritized by the regressors according to their predicted MICs. In five
rounds, using different versions of models, we generated ~500,000 new peptides by sampling
from the latent space. We filtered these peptides by length and viability to ~50,000 candidates
and prioritized 500 AMP candidates for wet lab bioactivity test (**Fig. 1c** and **Supplementary Table
2**).

119
Fig. 1: The workflow for de novo-development of AMPs via deep learning and cell-free biosynthesis. a,
Generative variational autoencoders (VAE) for de novo-design of AMPs after being trained on known AMP sequences.
b, Predictive convolutional or recurrent neural networks as regressors for the MIC prediction after being trained on
known AMPs and their MIC. **c,** Trained generative and predictive models are used for sampling from the latent space
(de novo-design of AMPs) and prioritization of AMPs (predicting their MIC), respectively. **d,** Experimental pipeline for
rapid cell-free biosynthesis of the designed AMPs from synthetic DNA fragments and direct testing of produced AMPs
in the cell-fee mix to bacterial cultures followed by overnight continuous growth assay.

<u>Cell-free biosynthesis enables rapid screening for functional AMPs</u>

To establish CFPS-based screening of AMPs, we designed an experimental pipeline for the high-
throughput synthesis and testing of AMPs in 384-well format (**Fig. 1d**). The system is based on
linear DNA templates, which comprise a T7 promoter and a ribosome binding site (RBS), to initiate
transcription (TX) and translation (TL), followed by the AMP coding region, and a T7 terminator.
After adding the DNA template (10 nM) directly into 10 µL of a cell-free TX-TL system, AMPs were
produced within 4 hours (**Methods**). To test the antimicrobial activity of the in vitro-produced
peptides, 4 µL of the cell-free mix was added into a final volume of 20 µL cultures of *E. coli* (Gram-
negative) and *Bacillus subtilis* (Gram-positive). Following the OD_{600} measurement for 20 hours
allowed identification of those peptides that show antimicrobial activity by suppressing growth.
Overall, the entire process of CFPS with subsequent bioactivity tests takes ~24 hours, as the
system works with linear DNA and does not require any extensive cloning or peptide purification
steps.

First, we validated the screening pipeline with two known AMPs, BP100[24] and Cecropin B[25], and
then screened 500 AMP candidates (see above) in five subsequent design-predict-build-test
cycles to identify 30 functional de novo AMPs (**Fig. 2**). During these five rounds, the success of
functional AMP discovery increased from 0% to 12.7% from the first to the fifth round, respectively
(**Supplementary Table 2**). Because translation initiation rates (TIRs) strongly affect protein
yield[26] we tested whether our screen had biased against candidates with low TIR. We calculated
the TIR for all sequences tested[27], but could not find a significant difference between the 500
candidates tested and the 30 functional AMPs identified (**Supplementary Note 1**,
Supplementary Fig. 3). Functional AMPs were re-validated in biological triplicates with 10 µL of
cell-free mix added to the final volume of 20 µL cultures (**Fig. 2b** and **Supplementary Fig. 1**) and
production of AMPs was analyzed through SDS-PAGE (**Supplementary Fig. 2**)

151
Fig. 2: Cell-free production of de novo-generated and prioritized AMPs and activity screening against *B. subtilis*
and *E. coli*. a, We used different generative and regressor models (**Supplementary Table 2**) to design and prioritize
AMPs in five rounds, produced and screened a total number of 500 AMPs from synthetic DNA fragments and found 30
functional candidates. **b,** The charge and AlphaFold-predicted structure of the functional AMPs with associated
slowed/stopped growth curves for *B. subtilis* and *E. coli*. All (including control) AMPs were produced using CFPS and
no peptide purification was carried out prior to the activity test. Growth curves (OD_{600} 0-0.45 over time 4-20 h for all)
are the average of n = 3 independent experiments. Growth curves with error bars as standard deviation are provided in
Supplementary Fig. 1. **c,** 2D-projections of the 50-dimensional latent space were obtained by principal component
analysis (PCA) for the two generative variational autoencoders (VAE) that were used for de novo-design of AMPs. Blue

color intensity represents the frequency of training AMPs in the latent space. Functional AMPs (red), BP100 and
Cecropin B (black) annotated back into the latent space. Source data for b are provided as a **Source Data** file.

<u>Functional AMPs are unique but share properties with natural counterparts</u>

We next analyzed our de novo-designed AMPs in more detail. AlphaFold[28] predicted that 27 out
of the 30 sequences form a helical structure (**Fig. 2b**), which is a common feature of many
AMPs[4,5]. Interestingly, AMPs #1-19, generated by VAE 1, showed more structural diversity
compared to AMPs #20-30, generated by VAE 2, which is in line with the fact that the two VAEs
create two different latent spaces (**Fig. 2c**), thus generating AMPs with distinct structural,
physicochemical, and sequence features (**Supplementary Fig. 4**). One of the main
characteristics of AMPs is an amphiphilic character that results from alternating cationic and
hydrophobic amino acids in the AMP core[4,5]. In contrast to natural AMPs, which mainly feature
aliphatic amino acids, the hydrophobic core of our de novo AMPs was mostly aromatic
(**Supplementary Fig. 5a**). Phenylalanine was particularly overrepresented at the cost of leucine,
which was underrepresented (**Supplementary Fig. 5b**, **Supplementary Table 3**). BLAST
searching showed that our de novo AMPs were unique in their sequence. No significant similarity
was observed against the UniProt database, encompassing ~240 million entries, nor an AMP
sequence from the training data set. (**Supplementary Tables 4-6**, **Supplementary Note 2** for
detailed BLAST sequence similarity analyses). Altogether, these results demonstrated that our
de novo AMPs shared the physico-chemical building principles with their natural counterparts, but
were distinct from them in their amino acid sequences.

<u>De novo AMPs prefer bacterial over human membranes</u>

Structural and sequence analysis suggested that our de novo AMPs act as amphipathic helices
that insert into membranes. We used molecular dynamics (MD) simulations to study the
interaction of our AMPs with models of a negatively charged inner membrane of bacteria (IM) and
the human plasma membrane (PM) (**Fig. 3a**, **Supplementary Note 3**). According to our
simulations, all AMPs bind much stronger to the IM interface than to the PM (**Fig. 3b**). Binding of
the AMPs at the IM progressed rapidly, taking at most 200 ns to fully insert into the membrane
interface (**Supplementary Fig. 6a**). Once bound, AMPs stayed tightly bound to the IM for the
remainder of the simulations. In some cases, we observed a reorientation of the AMP after a few
hundred nanoseconds from a shallowly bound state to a binding mode that resided deeper in the
membrane (e.g., AMP #29). Several AMPs also partially bind the PM. However, in most cases,
this binding is transient with frequent un- and rebinding and without penetrating deeper into the
PM, as seen for the IM (**Supplementary Fig. 6b**). Furthermore, all AMPs show a higher number
of (mainly electrostatic) interactions with the IM and remain in an ordered structure, compared to
the PM system, where fewer interactions were observed and AMPs tended to unfold over time
(**Supplementary Fig. 6c-d**). The MD simulations suggested that all our AMPs generally target
bacterial membranes over the human plasma membrane, naturally however, the degree of
preference is dependent on the individual AMP.

199
Fig. 3: Molecular dynamics (MD) simulations of AMP interactions with membranes. a, Overview of simulation
setup and exemplary snapshot of AMP #5 after 1 μs of simulated time on the IM. The peptide is shown in red cartoon
representation, solvent (water + ions) is shown as transparent blue surface and the membrane is shown as gray licorice
representation with lipid headgroup phosphates shown as spheres. **b,** Distributions of distances between the centers
of mass of the AMPs and the membrane midplane along the direction of the membrane normal (y axis). Distributions
are calculated from the last 940 ns of 1 μs long replicates, run with different lateral membrane tensions (x axis) and
with different membranes (blue: PM; red: IM). The thin dotted line indicates the headgroup phosphate positions. AMP
#2 was simulated with and without disulfide bond (**Methods**). **c,** Rendered simulation snapshots of potent AMPs (**Fig.
4**) on IM (top) and PM (bottom). The interacting leaflet of the membrane is depicted with gray lipid tails and orange
(phosphates) and green (cholesterol oxygen) spheres.

De novo AMPs show favorable MIC to toxicity ratios

To obtain pure compounds for cellular assays, we chemically synthesized the functional AMPs and characterized their bioactivity, in particular the minimum inhibitory concentration (MIC)[29], as well as hemolysis (HC50) and cytotoxicity (CC50), both expressed as 50% toxic concentration. Of the 30 candidates, 22 peptides were successfully produced by chemical synthesis. Twenty AMPs showed a MIC of ≤6 µM on *B. subtilis*, and fifteen AMPs showed a MIC of ≤25 µM on *E. coli* (**Fig. 4a** and **Supplementary Table 7**). HC50 and CC50 were significantly higher in most cases, with fifteen AMPs showing HC50 >100 µM (thirteen >250 µM) against fresh human red blood cells, and thirteen AMPs showing CC50 >100 µM (six >250 µM) against HCT116 human colon cells (**Fig. 4a** and **Supplementary Table 7**). This indicates that the bioactivity versus toxicity relationship was very favorable for several of our de novo AMPs. We decided to continue with sixteen AMPs that showed a favorable bioactivity to toxicity ratio and excluded six AMPs because of high MIC and/or low HC50/CC50 values (AMP #1, #7, #10, #18, #23, and #26).

Fig. 4: Bioactivity characterization of chemically synthesized functional AMPs. a, Minimum inhibitory concentration (MIC) of the AMPs against *E. coli* and *B. subtilis* (average of n = 3 independent experiments) and HC50 (hemolysis) and CC50 (cytotoxicity) values of the AMPs on human red blood cells and HCT116 human colon cells, respectively (average of n = 2 independent experiments). MIC, HC50 and CC50 values are provided in **Supplementary Table 7**. **b,** MIC values of the AMPs tested against ESKAPE pathogens including *E. faecium*, Methicillin-resistant *S. aureus* (MRSA), *K. pneumoniae*, *A. baumannii*, *P. aeruginosa*, *Enterobacter* spp measured in duplicates (n = 2 independent experiments) and MIC of the six potent, broad-band AMPs on *Y. pestis* and *B. anthracis* as the average of triplicates (n = 3 independent experiments).

De novo AMPs show broad-band activities

In the following, we tested the sixteen remaining de novo AMPs against clinically relevant strains, and in particular multidrug-resistant ESKAPE pathogens (i.e., *Enterococcus faecium*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa*, and *Enterobacter* spp.; **Fig. 4b**). Our AMPs were most potent against *E. faecium* and *A. baumannii*, resulting in fifteen AMPs with MIC ≤25 μM (thirteen ≤6.3 μM), and thirteen AMPs with MIC ≤25 μM (ten ≤6.3 μM), respectively (**Fig. 4b**). For the rest of the ESKAPE pathogens, *K. pneumoniae*, *P. aeruginosa*, *S. aureus*, and *Enterobacter* spp., ten, nine, eight, and three AMPs showed MIC ≤25 μM, respectively (**Fig. 4b**). While some AMPs showed distinct activity profiles against individual strains, six AMPs classified as "broad-spectrum" antimicrobials that showed favorable therapeutic window i.e., antimicrobial activity at relevantly low hemolysis and cytotoxicity (AMP #3, #5, #13, #15, #16, and #27). Notably, these AMPs were also active against the notorious biothreat agents *Yersinia pestis* and *Bacillus anthracis* (**Fig. 4b**).

Of the six broad-band candidates, AMP #5 was the most potent one against multiple pathogens (**Fig. 4b**) and showed a highly favorable bioactivity to cytotoxicity ratio. Thus, we further decided to test AMP #5 in a clinically relevant infection model. We studied the impact of AMP #5 treatment on the macrophage response to infection with *Streptococcus pneumoniae* (**Fig. 5a**)[30,31]. AMP #5 resulted in only low cytotoxic damage (**Fig. 5b**), but strongly reduced interleukin-6 (IL-6), interleukin-8 (IL-8) and tumor necrosis factor alpha (TNF-α) cytokine levels in *S. pneumoniae*-infected macrophages compared to untreated macrophages (**Fig. 5c,d**). These results indicated that AMP #5 had successfully reduced *S. pneumoniae* pro-inflammatory activation of the macrophage, confirming AMP #5 as a promising antimicrobial compound.

Fig. 5: Analysis of the macrophage response upon infection by *Streptococcus pneumoniae* and AMP treatment. **a,** We preincubated blood-derived macrophages for one hour with AMP #5, infected with *S. pneumoniae* D39 at MOI 1, and collected RNA and supernatant samples after 24 hours. **b,** Cytotoxicity testing of the supernatant by LDH (lactate dehydrogenase) assay for 100% kill control (1% saponin), (AMP) solvent control, AMP #5, *S. pneumoniae* infection, and *S. pneumoniae* infection and AMP #5 treatment. **c, d,** Macrophage response to infection by measuring protein (**c**) and mRNA (**d**) of interleukin-6 (IL-6), interleukin-8 (IL-8) and tumor necrosis factor alpha (TNF-α) cytokines (**Methods**).

No resistance was developed against de novo AMPs

Next, we tested the emergence of resistance against the six broad-band AMPs identified in this study (AMP #3, #5, #13, #15, #16, and #27). To that end, we performed *E. coli* serial passaging experiments with the peptides. As control, we added imipenem, a broad-spectrum antibiotic that is generally considered a "last resort" against multidrug-resistant pathogens[32]. During 21 days of serial passage, we did not observe a significant increase in MIC of our AMPs, while the imipenem MIC gradually increased up to 8-fold and exceeded the susceptibility breakpoint of clinically

relevant resistance defined by EUCAST (**Fig. 6a** and **Supplementary Fig. 7**). The fact that we
did not observe resistance is in line with the fact that our AMPs do not act on a specific cellular
target, but rather globally (i.e., at the membrane, **Fig. 3c**), which makes them less likely to cause
resistance development. This mode of action was further confirmed by propidium iodide staining
and microscopy (**Fig. 6c-d**), which clearly indicated membrane disruption upon AMP treatment.
Finally, we also investigated the effect of AMP treatment on the release of outer membrane
vesicles (OMVs) in *E. coli*. OMVs are naturally released by Gram-negative bacteria and can be a
reaction against surface attacking agents neutralizing the effect of membrane targeting
antibiotics[33–35]. Notably, all six broad-band AMPs showed no significant alteration of OMV release
compared to untreated *E. coli*. In contrast, an AMP candidate with limited antimicrobial efficacy
(AMP #10) that we had excluded early on in this work, significantly induced OMV release (**Fig.**
6b). Altogether, these experiments suggested that our six broad-band AMPs are able to escape
resistance development and self-defense reactions of bacteria.

284
Fig. 6: Resistance and mode of action test of AMPs on *E. coli* and the mode of action assays. a, 21 days MIC
measurement of daily culture passage each day from the cells grown in half-MIC concentrations of AMPs. The MIC

values were normalized by the minimum of 21 daily averages for each AMP. Bars are the average of n = 3 independent
experiments. See **Supplementary Fig. 7** for raw MIC data. **b,** Impact of different AMPs on the relative number of *E.coli*
outer membrane vesicles (OMVs) for untreated *E. coli* (control), (AMP) solvent, six potent AMPs (black bars) as well
as one less potent AMP (red bar). Bars are the average of n = 3 independent experiments. Statistics: one-way ANOVA;
****p <0.0001, compared to control and each of the other AMPs (black bars). **c,** Plate reader measured fluorescence
of propidium iodide (DNA staining after membrane disruption) of *E. coli* cells untreated (control) or treated with AMPs.
Bars are the average of n = 3 independent experiments. *P* values are for paired Student's t-test of each AMP with
untreated *E. coli* cells. **d,** Phase-contrast microscopy of *E. coli* cells untreated or treated with six potent AMPs and
Cecropin B (a membrane disruptive AMP) and stained with propidium iodide (red). Source data for **a-c** and six images
(n = 3 independent experiments) of which one was used in **d** are provided as a **Source Data** file.

Discussion

In this work, we describe the design and validation of 30 de novo AMPs, of which six show broad-
band activity. Recent work has impressively demonstrated the power of deep learning methods
in discovering and/or developing novel AMPs[6,19]. However, these efforts still suffer from the limited
number of peptides that can be synthesized and tested, and the relatively long time it takes from
design to validation. Here we used CFPS to dramatically advance the design-build-test cycle in
AMP development. Compared to recent approaches[19], our success rate is in the same range (6%
versus 10%). However, the total number of functional AMPs discovered is an order of magnitude
higher (30 AMPs versus 2) and at massively increased rate (24 hours versus 28 days) and
reasonably low cost (<10$ for production of one AMP for screening on two strains in parallel).

While being unique and diverse, our de novo AMPs share common properties with known AMPs.
They are predicted to be mostly α-helical peptides rich in cationic and hydrophobic amino acids
and preferably act on negatively-charged IMs, showing that our pipeline was able to design new-
to-nature sequences that follow the general building principles of AMPs. While we put our focus
onto AMPs with broad-band activity, we note that our pipeline is in principle also suited to develop
and/or iteratively optimize peptides with more specific activities or desired functionalities (e.g.,
narrow spectrum, increased stability, etc.). A unique advantage of CFPS is the possibility to
incorporate modified and unnatural amino acids more rapidly and conveniently than by cell-based
methods.

Overall, our work provides a proof-of-principle, how CFPS can be used to leverage the full
potential of machine learning approaches in the future. Especially in the light of ever-decreasing
DNA synthesis costs, our combined approach of deep learning and CFPS provides a time-, cost-
, and labor-effective approach for peptide production and screening. Thus, our work holds the
potential to explore the design-function space of AMPs at increased rate and depth. This will
hopefully lead to the increased discovery and development of peptide-based drug candidates in
the future.

Methods

Pretraining and training datasets

Pretraining data. To gather a large corpus of protein sequences representing a general protein
grammar, we downloaded all protein sequences shorter than 49 amino acids from UniProt[36] (as
of July 2021). After removing duplicate sequences and entries with unknown amino acid
characters, 3,104,952 unique sequences remained of which a random subset of half of them was
used for pretraining.

AMP data. For experimentally validated AMP sequences, we used the Giant Repository of AMP
Activities (GRAMPA)[37] which has combined sequence and activity data from several public AMP
databases; APD[38], DADP[39], DBAASP[40], DRAMP[41], and YADAMP[42]. This database consists of

6,760 unique sequences and 51,345 total MIC measurements, spanning several bacterial and
nonbacterial target species. We filtered the MIC measurements to the ten most abundant bacterial
species (*E. coli, P. aeruginosa, Salmonella typhimurium, K. pneumoniae, A. baumannii, S.*
aureus, B. subtilis, S. epidermidis, Micrococcus luteus and E. faecalis) and omitted all peptides
containing any chemical modification other than C-terminal amidation, for feasible in-laboratory
expression. After removing duplicate sequences and entries longer than 48 amino acids, 5,319
unique AMP sequences were left.

Non-AMP data. We searched the UniProtKB[36,43] for proteins labeled as "NOT antimicrobial,
antibiotic, antiviral or antifungal" (downloaded as of July 2021), removed entries containing
ambiguous amino acids, and kept only unique sequences shorter than 49 amino acids. This
resulted in a dataset containing 10,612 unique non-AMP sequences.

Generator variational autoencoder (VAE)

We used VAEs because they have previously been used for de novo AMP design[16,17,19,44]. The
generative VAE consists of an encoder, a latent vector, and a decoder. The encoder feeds the
input data (one-hot encoded amino acid letter of peptides) into a latent vector that is an information
bottleneck, and the decoder aims to reconstruct the input data from the latent vector. Training the
VAE and minimizing the difference between the input data and the reconstructed data acts
twofold; the encoder learns to map the training dataset into a lower-dimensional space and the
decoder learns to generate samples similar to the training data from any vector in the latent space.
Thus, each peptide in the training dataset lands on a point in the multi-dimensional latent space.
Picking vectors from the empty regions in this space and feeding them into the decoder yield
peptide sequences that share the same grammar but are novel and not seen in the training
dataset. After pretraining and training (transfer learning), we generated new AMPs by sampling
from the latent space using different strategies in particular by exploring the neighborhood of a
control functional AMP, gradient descent, or random sampling (**Supplementary Tables 1, 2**).

Generator VAE models. We adapted the neural network architecture of the CNN-RNN hybrid VAE
model from Hawkins-Hooker et al.[23]. The encoder consists of 5 consecutive one-dimensional
convolution layers fed into a dense layer of size 50, which is the latent vector. The decoder is
made from 4 deconvolution layers that samples the latent vector and a GRU layer of 512 cells
outputs a sequence in the same dimension as the input. The model loss is the weighted sum of
a reconstruction loss and the Kullback-Leibner (KL) loss. The total loss function can be
dynamically changed in the training process for KL-term annealing[45]; as standard practice when
working with discrete data such as language. Our two final models were trained without and with
the KL-term annealing (VAE_v1 and VAE_v2, respectively). The models were compiled using the
Adam optimizer.

Pretraining and training. Our training dataset of ~5,000 AMP sequences is not sufficient for
learning what makes a protein sequence distinct from a random string of amino acid characters
and what makes a protein sequence an AMP. In such cases, pretraining with a much bigger
generic dataset is needed to enhance the model performance. We pretrained the generator
models with ~1.5 million protein sequences from UniProtKB for 600 epochs. We then trained the
models on the AMP data for 400 epochs. Model training metrics are provided in **Supplementary**
Table 1.

Regressor convolutional and recurrent neural networks

Regressor neural networks. We adapted a regressor model previously reported[37]. First, a CNN
regressor was built with two consecutive one-dimensional convolutional layers, a max pooling

layer, a flattening layer, a dropout layer (0.5), and three dense layers. Second, we built a simple
RNN regressor with an LSTM layer and two dense layers. The models were compiled with mean
squared error loss and the Adam optimizer.

Gram-specific regressors. Due to structural differences in the cell membrane, we assumed there
are differences between Gram-positive and Gram-negative bacteria in their response to AMPs.
To capture these differences, we trained Gram-specific regressor models. We trained the Gram-
negative model on 4619 unique AMP sequences and corresponding MIC measurements on *E.*
coli; and the Gram-positive model on 4089 AMP sequences and corresponding MIC
measurements on *B. subtilis*. This approach improved the accuracy and efficiency of the
regressor models (**Supplementary Tables 1, 2**).

Training. We trained these models on pairs of data containing AMP sequences and their
corresponding MIC measurements (in log 10). Based on a previous study[37], the nonAMP
sequences were labeled to have a log MIC of 4. We interpreted the predicted MIC value as
follows; below 3.5 as AMP, between 3.5-3.9 as potential AMP and above 3.9 as non-AMP. The
regressor models were trained by the pooled AMP and non-AMP data for 200 epochs. Model
training metrics are provided in **Supplementary Table 1**.

<u>Sampling and prioritizing of AMPs</u>

In each round of peptide synthesis, we selected thousands of random points from the VAEs latent
space and from each reconstructed a peptide sequence. We omitted non-viable sequences and
kept AMPs with 36-48 amino acids. This left us with viable peptide sequences (**Supplementary**
Table 2). We then fed these sequences into the regressors and prioritized them based on the
MIC prediction. 50-150 AMPs were chosen for wet-lab experiments in five rounds making a list of
over 500 AMPs tested in this work (**Supplementary Table 2**).

<u>Molecular dynamics (MD) simulations</u>

All MD simulations were performed using Gromacs 2020.3[46] and the CHARMM36m forcefield[47]
using an integration time step of 2 fs (partly 1 fs in the membrane equilibration; see
Supplementary Table 9). Bonds including hydrogens were constrained using the LINCS
algorithm[48]. Electrostatic interactions were computed using the Particle-Mesh Ewald (PME)
algorithm[49] with a real-space cut-off for pairs further apart than 1.2 nm. Lennard-Jones
interactions were smoothly switched to zero between 1 and 1.2 nm using the force-switch
algorithm.

The protein, the membrane and the solvent (water and ions) were individually coupled to thermal
baths set to 37 °C (310.15 K) using the v-rescale algorithm[50] with a time-constant of 1 ps. During
the equilibration runs pressure coupling was handled by the Berendsen barostat[51], which was
switched to the Parrinello-Rahman barostat[52] for all production runs. The barostat time-constant
and compressibility factor were consistently set to 5 ps and 4.5 x 10^{-5} bar^{-1}, respectively. Pressure
coupling was applied semiisotropically (with the x and y dimensions coupled together) for systems
with membrane and isotropically otherwise. The reference pressure was set to 1 bar. To
counteract an energetic penalty for peptide insertion into the membrane due to finite size effects,
we simulated each membrane system with three different lateral membrane tensions (0, 9, 17.1
bar). Therefore, the diagonal elements of the pressure tensor (P_{XX}, P_{YY}, P_{ZZ}) were set to

419
$$P_{XX} = P_{YY} = P - \frac{\Delta P}{3} \qquad\qquad P_{ZZ} = P + 2\frac{\Delta P}{3}$$

with P as the reference pressure (1 bar) and ΔP as the desired lateral membrane tension. To
ensure uncorrelated runs for the different tensions, the starting velocities of the atoms were
randomly initialized according to the Maxwell-Boltzmann distribution. The MD simulations of the
AMPs on membranes were performed for 1 µs each.

Visual analysis and renders used the VMD[53], PyMOL[54] and ChimeraX[55] software.

Membrane setup. Following earlier work and results from lipidomics experiments, we modeled
membranes resembling the outer leaflet of the human plasma membrane (PM)[56,57] and the *E. coli*
inner membrane (IM)[58]. The detailed compositions are summarized in **Supplementary Tables**
10, 11.

Using the CHARMM-GUI membrane builder[59,60], we generated 12.5 x 12.5 nm^2 patches of these
model membranes, energy minimized them with a steepest descent algorithm until the largest
force acting on any atom was below 1000 kJ mol^{-1} and subsequently equilibrated them following
the CHARMM-GUI equilibration scheme (summarized in **Supplementary Table 9**).

AMP system setup. Structures of the selected 30 AMPs and of Cecropin B were predicted using
AlphaFold[28] while the short BP100 was modeled as a coil with initial angles φ=-60° and ψ=30°,
using the Molefacture Protein Builder plugin for VMD[53]. Since several of our AMPs have more
than one cysteine (AMPs #2, #4, #6, #8, #10, #11, #12, #14, #18, #19), we next performed 1-2
µs long MD simulations of them surrounded by only water and ions (150 mM NaCl plus counter
ions for overall neutralization) and with no disulfide bonds imposed. Based on the frequency and
distance with which two cysteines in the respective structure interacted with each other in these
simulations, we assigned a disulfide bond for AMP #2 (Cys11, Cys47) and disulfide bonds
connecting the predicted β-sheets of AMP #8 (Cys29, Cys35) and AMP #14 (Cys10, Cys32). Due
to the particularly high abundance of cysteine in AMP #2 and the potential structural bias from
imposing one specific disulfide bond, we additionally simulated it without any imposed disulfide
bonds.

These structures were then orientated so that their first principal axis was orthogonal to the z-axis
and placed in proximity, but not yet bound to the equilibrated PM and IM. The box was
subsequently solvated with TIP3P water[61] and NaCl ions were added to a concentration of 150
mM, ensuring overall neutrality by adding additional counter ions to the systems. The systems
were energy minimized in the same way as the pure membrane systems and were then
equilibrated for 5 ns. During minimization and equilibration, we applied position restraints on the
peptide heavy atom with a force constant of 1000 kJ mol^{-1} nm^{-2}.

Cell-free production and activity test of AMPs

DNA fragments encoding AMPs were designed with T7 promoter
(GAATTTAATACGACTCACTATAGGGAGA), RBS (TCTAGAGATTAAAGAGGAGAATACTAG)
sequences upstream of the AMP coding region, and a T7 terminator
(TACTCGAACCCCTAGCCCGCTCTTATCGGGCGGCTAGGGGTTTTTTGT) downstream. 500
DNA fragments were purchased from Twist Bioscience. A final concentration of 10 nM of each
fragment was used for cell-free transcription and translation of AMPs using PUREfrex®2.0 kits
(GeneFrontier #PF201-0.25-5-EX). In 384-well plates (BRAND, #781687), 30 µL volume of the
cell-free reaction was made for each AMP and incubated for 4 hours at 37 °C. The cell-free mix
was directly used for the activity test on *E. coli* and *B. subtilis* or for the SDS-PAGEl of the
functional AMPs.

E. coli MG1655 and *B. subtilis* PY79 were used as representatives of Gram-negative and Gram-
positive bacteria. From LB agar plates into LB medium, three overnight cultures for each strain
were made from three different colonies and grown while shaking at 37 °C. The next day, each
was subcultured in LB (1:1000) and grown while shaking at 37 °C to OD ≈ 1. Cells were diluted
in LB to 10^4 cfu mL^{-1}, and 16 µL of diluted cells were added to wells of a 384-well plate (Greiner
Bio-One, #781185) in which 4 µL of the cell-free reaction mix (with AMPs produced) had been
added beforehand. Cultures were mixed and the plate was sealed by a gas-permeable film (Carl
Roth, #T093.1). OD_{600} was measured every 10 min in a plate reader (Tecan Infinite® 200 PRO)
shaking at 37 °C for 20 hours. Growth curves were analyzed for AMPs impairing bacterial growth.

SDS-PAGE of AMPs produced in the cell-free system

SDS-PAGE was used to detect produced functional AMPs in the cell-free reaction as previously
described[22]. Cell-free reactions were boiled for 3 min in 2x Tricine buffer (Bio-Rad, #1610739),
loaded in 16.5% Mini-PROTEAN polyacrylamide Tris/Tricine gels (Bio-Rad, #4563065), and run
for 5 hours at 200 mA in the running buffer (10 mM Tris, 10 mM Tricine, and 0.01% SDS). Gels
were then fixed for one hour in 12% trichloroacetic acid and one hour in 40% EtOH, 10% acetic
acid, followed by overnight staining in QC Colloidal Coomassie (Bio-Rad, #161-0803), 24 hours
of de-staining in water, and imaged using Intas GelStick Touch Imager.

Measurement of minimum inhibitory concentration (MIC) and resistance test

Strains used for MIC measurements are *E. coli* MG1655, *B. subtilis* PY79, *E. faecium* (isolate
from the gut of the cow), *S. aureus* DSM 11729, *K. pneumoniae* DSM 30104, *A. baumannii* (isolate
from the human abdominal wall), *P. aeruginosa* DSM 1117, *Enterobacter* spp., *Y. pestis* EV76
and *B. anthracis* Sterne. A commonly used standard protocol for determination of MIC for
antimicrobials was used to measure the MIC of the AMPs taking into account all suggestions for
cationic AMPs in the protocol[29]. Chemically synthesized peptides were dissolved in BSA (0.2%
w/v) acetic acid (0.01% v/v) solution to have 10x of the highest concentration to be tested. In 96-
well PCR plates (Axygen, #PCR-96-SG-C) two-fold serial dilutions were made from columns 1-
10 in each row specified to each peptide and BSA (0.2% w/v) acetic acid (0.01% v/v) solution
was pipetted into columns 11 and 12. Triplicates of 7.5 µL of each dilution were pipetted into
polypropylene 96-well plates (Corning, #3359). Triplicates of bacterial overnight cultures in
Mueller-Hinton broth 2 (MHB 2, Sigma-Aldrich, #90922) were prepared from three different
colonies the day before and grown shaking at 37 °C, subcultured in the morning by diluting 1000x
in MHB 2 grown shaking at 37 °C to OD ≈ 1. Bacterial cultures were then diluted with MHB 2 to
10^5 cfu mL^{-1} and 67.5 µL of each triplicate was added on top of peptides in columns 1-11 of 96-
well plates. MHB 2 was added to column 12. The plates were sealed by adhesive films (VWR,
#391-1262) and incubated at 37 °C for 20 hours. MIC values were reported as the highest
concentration of each AMP in which no visible growth was observed.

The same procedure was used for the resistance test except for cultures that from the second
day on, each of the triplicate cells grown in the highest AMP concentration (half MIC) was diluted
10,000x in MHB 2 and added to newly prepared peptides dilutions.

Measurement of cytotoxicity (CC50)

Cytotoxicity assay was performed on HCT116 human colon cells (ATCC, #CCL-247™) using
CellTiter 96® AQueous One Solution Cell Proliferation Assay (Promega, #G3580) which is a
colorimetric method based on MTS (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-
(4-sulfophenyl)-2H-tetrazolium) for determining cell viability. The MTS tetrazolium compound

(Owen's reagent) is reduced into a colored formazan product by NADPH or NADH produced by
dehydrogenase enzymes in metabolically active cells.

On the first day, when cells reached the density of 50-80% of the covered surface, gently washed
twice using 10 mL of DPBS (Gibco, #14190367). 1 mL trypsin (Capricorn, #TRY-1B) was added,
5 min incubated at 37 °C and 9 mL medium was added including DMEM high glucose (Capricorn,
#DMEM-HPA), 10 % v/v fetal bovine serum (FBS, Capricorn, #FBS-11A), and antibiotic mix for
cell culture (Capricorn, #PS-B). The culture was transferred into a 15 mL falcon, and spun down
at 1000 rpm for 3 min. The supernatant was sucked out and 10 mL of fresh media was added and
transferred. Cells were diluted by the medium to have 5000 cells in 36 µL. 36 µL of the cell culture
was pipetted into wells of a 384-well plate (Greiner, #781185). The last well received only media.
Cells were incubated at 37 °C, 5% CO_2, for 24 hours. On the second day, peptides were prepared
in two-fold serial dilutions starting from a final concentration of 250 µM. Columns 11 and 12
received only water. 4 µL of each peptide dilution was added to wells of the cell culture plate
prepared on the first day and the plate was put at 37 °C, 5% CO_2, for 24 hours. On the third day,
8 µL of CellTiter 96® AQueous One Solution and 10 µL SDS 10% were added to each well and
after 90 min incubation at 37 °C, absorbance at 490 nm was measured and corrected by the value
of wells with only medium. CC50 values, the concentration of each AMP killing 50% of cells, were
calculated using Graphpad Prism 9.

<u>Measurement of hemolytic activity (HC50)</u>

Human blood from a healthy donor was washed three times with PBS and afterwards
resuspended in 2V PBS. AMPs with an initial concentration of 250 µM were titrated in 96-well
polypropylene plates (V-bottom, Greiner Bio-One GmbH). 5 µL AMP dilutions were overlaid with
45 µL of washed and concentrated human erythrocytes, and the plates were sealed and incubated
at 37°C for 1 h. 40 µL of supernatant were transferred after final centrifugation at 1000 g for 5 min
at room temperature to ELISA plates and absorbance was measured at 405 nm. Triton X 100
treated erythrocytes served as positive control. HC50 values, the concentration of each AMP
lysing 50% of RBCs, were calculated using Graphpad Prism.

<u>Mode of action assay and microscopy using propidium iodide (PI)</u>

Plate reader assay. Three colonies were picked to culture *E. coli* MG1655 cells in LB at 37 °C to
the exponential phase. Cells were harvested by centrifugation at 4000 g and washed three times
in 10 mM PBS (pH=7.0), and adjusted to OD_{600}=1 with 10 mM PBS. 10 µL of the cells in PBS
were mixed with 10 µL of AMPs to a final concentration of 4×MIC and incubated at 37 °C for 1
hour. 20 µM final concentration of PI[62] was added to each of the AMP-treated and untreated
samples and incubated at 37 °C for 30 min in the dark. Fluorescence was recorded at an excitation
of 535 nm and emission of 615 nm was measured using a Tecan Infinite® 200 PRO plate reader.

Sample preparation for microscopy. *E. coli* was grown in LB at 37 °C to the exponential phase
and diluted to 10^8 cfu mL^{-1} in fresh LB. 50 µL of diluted cells were pipetted in a 1.5 mL tube and
50 µL of AMPs was added at 4×MIC final concentration together with 20 µM PI. The mixture was
incubated at 37 °C for 1 h while shaking. 1 µL of samples was loaded onto agarose pads (2%
agarose in PBS) and imaged using a Zeiss AxioPlan 2 upright widefield microscope equipped
with a 100x NeoFluor phase contrast objective and a FluoArc HBO lamp. Fluorescence of PI was
recorded using TxRed HC Filter set (AHF, F36-504).

Image processing. The phase contrast and fluorescence images were overlaid. The dynamic
range of the fluorescence channel was set to a minimum of 125AU to remove background
fluorescence, while the contrast of the phase contrast channel was manually adjusted for better
visualization. For comparison of the different phenotypes, 232 pixels x 232 pixels regions of
interest were cropped. Raw image files are provided in this study.

<u>Outer membrane vesicle (OMV) release of *E.coli* after AMP treatment</u>

Escherichia coli MG1655 was grown on MacConkey agar (Carl Roth, Karlsruhe, Germany) plates
overnight. For overnight culture, a single colony was used for inoculation of 2 mL LB media at 37
°C, 160 rpm (MaxQ 6000, Thermo Fisher Scientific, Karlsruhe, Germany). Bacterial culture was
transferred to 10 mL fresh LB media and incubated (1 h, 37 °C, 160 rpm). The required amount
of bacteria was treated with ¼ MIC of the AMPs #3, #5, #10, #13, #15, #16, and #27, left untreated
for control or was treated with acetic acid and BSA as solvent control (90 min, 37°C, 160 rpm).
To determine bacterial colony forming units and check for possible contamination, serial dilutions
of the bacteria were plated. Samples were centrifuged (4,500 g, 15 min, 4 °C; Multifuge X3R,
Thermo Fisher Scientific), the supernatant was sterile filtered (0.22 µm) and afterwards
concentrated by the factor of 20 using 100 kDa molecular weight cut-off filters (Merck KGaA,
Darmstadt, Germany). To determine the number of released vesicles, samples were measured
by nano-flow cytometry (nFCM) using a NanoAnalyzer (NanoFCM Co., Ltd, Nottingham, UK) as
previously described[63]. The relative vesicle release per cell was calculated.

<u>Macrophage infection experiment</u>

We employed a macrophage infection model to test whether the bactericidal effect of the AMP is
relevant for immune cells, leading to an attenuated pro-inflammatory activation due to lower
bacterial burden. As also dead bacteria might trigger cells with their components (e.g., via the
Jarisch-Herxheimer reaction[64]), it is crucial to evaluate to what extent reduction of live bacteria by
the AMP impacts macrophage activation. The model of macrophage infection with *Streptococcus*
pneumoniae in the context of AMP treatment has been established before[30].

Monocytes were isolated from donor Buffy coats by CD14 positive selection according to the
manufacturer's protocol (Miltenyi Biotec). Monocytes were left to adhere to plastic and cultured
for seven days in serum-free X-VIVO medium and 15 ng mL^{-1} GM-CSF. On day seven, maturation
to macrophages was verified by microscopy, and cells were used for infection. For the infection
experiment, *S. pneumoniae* D39 was streaked on blood agar and cultivated overnight at 37 °C/5%
CO_2. The next day, single colonies were picked to inoculate a liquid culture in THY medium to an
OD_{600}=0.06 and cultivated at 37 °C/5% CO_2. Upon reaching OD_{600}=0.35, bacteria were used to
infect macrophages at a multiplicity of infection (MOI) of one, either untreated or treated with 25
µM (MIC in THY medium) AMP #5. RNA and Supernatant samples were taken after 24 hours.
For measurement of cytotoxicity, supernatant was used to measure the release of lactate
dehydrogenase (LDH) as described before[65] with a TECAN Infinite M200 Pro plate reader.

For flow cytometry measurement of secreted cytokines, cell culture supernatant from infected and
uninfected samples was prepared according to the manufacturer's protocol (Miltenyi Biotec, 12

Plex human cytokine panel). Samples were measured on a BD FACS Canto II, and APC signal
was used to quantify the cytokine signal by referencing the respective standard curve.
Commercial ELISA kits (OptEIA, BD Biosciences, Heidelberg, Germany) were used to detect IL-
8 in cell supernatant according to the manufacturer's instructions. The measurement was carried
out on a Tecan Infinite M200 Pro (Thermo Fisher Scientific). For qPCR, cDNA from infected and
uninfected samples was used for mRNA analysis of IL-6, IL-8, and TNFα with the following forward
and reverse primers: AATTCGGTACATCCTCGACGG and TTGGAAGGTTCAGGTTGTTTTCT
for IL-6, ACTGAGAGTGATTGAGAGTGGAC and AACCCTCTGCACCCAGTTTTC for IL-8, and
GCTGCACTTTGGAGTGATCG and TCACTCGGGGGTTCGAGAAGA for TNF-α.

<u>Chemical peptide synthesis of AMPs</u>

Materials. All commercially available reagents were purchased from the following companies, and
used without further purification: thioanisol, 1,2-ethandithiol (EDT) from Sigma Aldrich (USA);
piperidine, 2 mL polypropylene reactors with plunger and frit pore size 25 µm (#7926.1) from Carl
Roth (Germany); 2,6-lutidine, palladium acetate, phenylsilane, trifluoracetic acid (TFA) from Acros
(USA); microscale columns from Intavis (#35.091) (Germany); Fmoc-Gly-OH, Fmoc-L-Asn(Trt)-
OH, Fmoc-L-Asp(tBu)-OH, Fmoc-L-His(Trt)-OH, Fmoc-L-Ile-OH, Fmoc-L-Met-OH, Fmoc-L-Phe-
OH, Fmoc-L-Pro-OH*H20, Fmoc-L-Thr(tBu)-OH, Fmoc-L-Tyr(tBu)-OH, Fmoc-L-Val-OH, peptide
grade dimethylformamide (DMF), TentaGel S RAM resin from Iris Biotech (Germany); Fmoc-L-
Ala-OH-OH, Fmoc-L-Arg(Pbf)-OH, Fmoc-L-Cys(Trt)-OH, Fmoc-L-Gln(Trt)-OH, Fmoc-L-
Glu(OtBu)-OH, Fmoc-L-Leu-OH, Fmoc-L-Lys(Boc)-OH, Fmoc-L-Ser(tBu)-OH, Fmoc-L-Trp(Boc)-
OH, N,N′-diisopropylcarbodiimid (DIC), and Oxyma from Carbolution (Germany); acetic anhydride
(Ac$_2$O) from Grüssing GmbH (Germany); HPLC grade acetonitrile from Merck (Germany);
Ultrapure water of type 1 was obtained with a MicroPure Water Purification System from TKA
(Germany).

Solid-phase peptide synthesis. All peptides were synthesized via the Fmoc-solid phase strategy.
The synthesis was carried out by an automated peptide synthesizer using INTAVIS ResPep SLi
instrument for a 5 µmol scale. Higher amounts of peptides were achieved by running multiple 5
µmol syntheses in parallel. For all peptides, the TentaGel S RAM (0.22 mmol/g) resin was used.

Automated solid-phase peptide synthesis (INTAVIS ResPep SLi):

- The conditions, reagents, and corresponding volumes of this synthesis protocol
correspond to a 5 µmol scale synthesis. No mixing was performed during incubation or
reaction time. In the coupling step, the temperature was set to 40°C.

- Swelling: The appropriate amount of resin (23 mg) was swelling in 200 µL DMF for 30 min.

- Deprotection of temporal Fmoc protecting groups: Piperidine (150 µL, 20% in DMF) was
added to the resin and incubated for 5 min. This step was repeated, and the resin was
filtered off, and washed with DMF (1 x 300 µL, 3 x 225 µL).

- Coupling of amino acids: In a mixing vial the machine automatically added 53 µL of the
corresponding Fmoc-amino acid (4 eq, 0.5 M), 15 µL Oxyma (4 eq, 2 M), 13 µL DIC (4 eq,
2 M) each in DMF and 29 µL NMP. The resulting solution was activated by waiting for 1
min before addition to the resin. This suspension was incubated for 15 min. Next, the resin
was filtered off, and the coupling was repeated. No washing was performed after the
coupling.

- Capping: 150 µL of a lutidine/Ac$_2$O/DMF 6:5:89 solution was added to the resin and
incubated for 8 min. The resin was filtered off and washed with DMF (3 x 225 µL).

- After the last coupling, the resin was washed with DMF (1 x 300 µL, 3x 225 µL), ethanol
(4 x 150 µL) and CH$_2$Cl$_2$ (5 x 150 µL). The resin was finally dried under continuous air flow
for 5 min.

Final Cleavage, Purification, and Characterization:

- Cleavage and deprotection of the amino acid side chains: All peptides have been cleaved
from a dry resin previously washed with 5x DMF and 10x CH$_2$Cl$_2$ following the last step of
the synthesis protocol. Depending on the total number of Cys, Met, or Trp, one of the
cleavage cocktails in **Supplementary Table 12** was utilized. Cleavage cocktail A was
used as the initial test cleavage after complete synthesis[66]. In the cases where oxidation
was observed after cleavage, cocktail B was applied[67,68]. For 5 µmol of resin, 2 mL of
cleavage cocktail was prepared. The total volume was increased by 1.5-fold for peptides
containing more than eight arginine residues. The dry resin was loaded into 2 mL reactors
with a plunger and the frit was treated with the corresponding mixture and shaken for 2.5
h and filtered off. The resin was washed with 1 mL of TFA and the filtrates were combined.
The TFA content of the filtrate was reduced via a gentle nitrogen flow. Next, ice-cold diethyl
ether (DEE) (1.00 mL of DEE for 100 µL cocktail) was added to precipitate the final
peptide. The precipitated peptide was centrifuged (8000 rpm, 4°C, 5 min), the supernatant
was discarded and the pellet washed once more; i.e. redissolved, and precipitated with
cold DEE. Afterward, the peptide pellet was dissolved in ultrapure water/MeCN (70:30)
with 0.1% of TFA (more MeCN was added when insoluble, not exceeding 1:1) to be
purified.

- Purification: The peptides were purified by reverse-phase (RP)-HPLC using a preparative
Agilent 1260 Infinity II Series HPLC-system (Agilent Technologies) with column 1
(**Supplementary Table 13**). An isocratic regime during the first five minutes for column
equilibration, followed by the respectively stated linear gradient in 25 min (gradient is
specified at the respective peptide). The detection was carried out by measuring
absorption at the wavelengths: 220 nm and 260 nm. Ultrapure water (A) and MeCN (B)
were employed as eluents with an addition of 0.1% of TFA in both solvents.

- Characterization: The freeze-dried products were identified via analytical HPLC-MS on an
Agilent 1260 Infinity II Series HPLC-system (Agilent Technologies) using column 2
(**Supplementary Table 13**). The detection was carried out by measuring absorption at the
wavelengths: 220 nm and 260 nm. Ultrapure water (A) with an addition of 0.05% of TFA
and MeCN (B) addition of 0.03% of TFA were employed as eluents. HR-ESI-MS was
performed for identification on an LTQ-FT Ultra device (Thermo Fischer Scientific). HPLC
chromatogram of purified peptides are provided in **Supplementary Figs. 8-34**.

Acknowledgments

This work was supported by a European Molecular Biology Organization (EMBO) long-term postdoctoral fellowship to A.P. (ALTG 165-2020), the Max Planck Society, German Research Council DFG; AB 792/1-1 (423428279) to F.A., German Research Council TRR81: "Chromatin Changes in Differentiation and Malignancies" TRR81/3, Z04 (109546710) to V.T.T. and O.V., the Bundesministerium für Bildung und Forschung (Federal Ministry of Education and Research, PermedCOPD – FKZ 01EK2203A; ERACoSysMed2 – SysMed-COPD – FKZ 031L0140), the Deutsche Forschungsgemeinschaft (SFB/TR-84 TP C01) to B.S., the Hessisches Ministerium für Wissenschaft und Kunst (LOEWE Diffusible Signals LOEWE-Schwerpunkt Diffusible Signals) to A.L.J and B.S. S.LS. and G.H. acknowledge financial support by the Clusterprojekt ENABLE funded by the Hessian Ministry for Science and the Arts, and the Collaborative Research Center 1507 funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), and thank the Max Planck Computing and Data Facility (MPCDF) for computational resources. We thank R. Weiss, P. Wichmann, A.M. Küffner, S. Scholz, R. Inckemann, C. Diehl, A. Yazdizadeh Kharrazi, J. Zarzycki, and H. He for technical and experimental support, as well as fruitful discussion. Figures were created with Biorender.com.

Contributions

T.J.E and A.P. conceived the study and wrote the manuscript with contributions from other authors, as indicated in the following. A.Z. developed deep learning codes and simulations and wrote the corresponding method section and performed BLAST analysis. A.P. and Y.F. performed cell-free protein synthesis and the initial bioactivity tests. M.K. performed the RBS calculator and kinfolk calculations. S.L.S. performed and, together with G.H analyzed molecular dynamics simulation and wrote the corresponding methods and results section. A.P., D.A., P.B. and H.v.B. performed and analyzed MIC, hemolysis, and cytotoxicity tests. V.T.T. performed chemical peptide synthesis and analysis via automated solid-phase and wrote the corresponding methods section. A.P., E.B, and C.S. performed and analyzed the mode of action assays. M.W. performed and analyzed the outer membrane vesicle experiments and wrote the corresponding results and methods sections. C.P. established the OMV quantification and performed the OMV quantification in the Core Facility for Extracellular Vesicles. B.K. performed and analyzed the macrophage infection experiment and wrote the corresponding results and methods sections. E.P.v.S., H.B.B., H.v.B., W.B., A.L.J., F.A., B.S, G.H., O.V., and T.J.E supervised the work. All authors provided input on the manuscript and confirmed the final draft.

References

1.   10 global health issues to track in 2021. https://www.who.int/news-room/spotlight/10-global-

health-issues-to-track-in-2021.

2.   Antimicrobial Resistance Collaborators. Global burden of bacterial antimicrobial resistance

in 2019: a systematic analysis. *Lancet* **399**, 629–655 (2022).

3.   Miethke, M. *et al.* Towards the sustainable discovery and development of new antibiotics.

Nat Rev Chem **5**, 726–749 (2021).

4.   Lazzaro, B. P., Zasloff, M. & Rolff, J. Antimicrobial peptides: Application informed by

evolution. *Science* **368**, (2020).

5.   Jaumaux, F., P Gómez de Cadiñanos, L. & Gabant, P. In the Age of Synthetic Biology, Will

Antimicrobial Peptides be the Next Generation of Antibiotics? *Antibiotics (Basel)* **9**, (2020).

6.   Ma, Y. *et al.* Identification of antimicrobial peptides from the human gut microbiome using

deep learning. *Nat. Biotechnol.* **40**, 921–931 (2022).

7.   Fjell, C. D., Hiss, J. A., Hancock, R. E. W. & Schneider, G. Designing antimicrobial

peptides: form follows function. *Nat. Rev. Drug Discov.* **11**, 37–51 (2011).

8.   Dijksteel, G. S., Ulrich, M. M. W., Middelkoop, E. & Boekema, B. K. H. L. Review: Lessons

Learned From Clinical Trials Using Antimicrobial Peptides (AMPs). *Front. Microbiol.* **12**,

616979 (2021).

9.   Huang, P.-S., Boyken, S. E. & Baker, D. The coming of age of de novo protein design.

Nature **537**, 320–327 (2016).

10. Pan, X. & Kortemme, T. Recent advances in de novo protein design: Principles, methods,

and applications. *J. Biol. Chem.* **296**, 100558 (2021).

11. Dauparas, J. *et al.* Robust deep learning-based protein sequence design using

ProteinMPNN. *Science* eadd2187 (2022).

12. Ferruz, N., Schmidt, S. & Höcker, B. ProtGPT2 is a deep unsupervised language model for

protein design. *Nat. Commun.* **13**, 4348 (2022).

13. Nijkamp, E., Ruffolo, J., Weinstein, E. N., Naik, N. & Madani, A. ProGen2: Exploring the

boundaries of protein language models. (2022) doi:10.48550/ARXIV.2206.13517.

14. Strokach, A. & Kim, P. M. Deep generative modeling for protein design. *Curr. Opin. Struct.*

Biol. **72**, 226–236 (2022).

15. Li, F. *et al.* Deep learning-based kcat prediction enables improved enzyme-constrained

model reconstruction. *Nat. Catal.* **5**, 662–672 (2022).

16. Dean, S. N. & Walper, S. A. Variational Autoencoder for Generation of Antimicrobial

Peptides. *ACS Omega* **5**, 20746–20754 (2020).

17. Dean, S. N., Alvarez, J. A. E., Zabetakis, D., Walper, S. A. & Malanoski, A. P. PepVAE:

Variational Autoencoder Framework for Antimicrobial Peptide Generation and Activity

Prediction. *Front. Microbiol.* **12**, 725727 (2021).

18. Wang, C., Garlick, S. & Zloh, M. Deep Learning for Novel Antimicrobial Peptide Design.

Biomolecules **11**, (2021).

19. Das, P. *et al.* Accelerated antimicrobial discovery via deep generative models and

molecular dynamics simulations. *Nature Biomedical Engineering* **5**, 613–623 (2021).

20. Garenne, D. *et al.* Cell-free gene expression. *Nat Rev Methods Primers* **1**, (2021).

21. Silverman, A. D., Karim, A. S. & Jewett, M. C. Cell-free gene expression: an expanded

repertoire of applications. *Nat. Rev. Genet.* **21**, 151–170 (2020).

22. Pardee, K. *et al.* Portable, On-Demand Biomolecular Manufacturing. *Cell* **167**, 248–259.e12

(2016).

23. Hawkins-Hooker, A. *et al.* Generating functional protein variants with variational

autoencoders. *PLoS Comput. Biol.* **17**, e1008736 (2021).

24. Badosa, E. *et al.* A library of linear undecapeptides with bactericidal activity against

phytopathogenic bacteria. *Peptides* **28**, 2276–2285 (2007).

25. Jarvis, D. L., Summers, M. D., Garcia, A., Jr & Bohlmeyer, D. A. Influence of different signal

peptides and prosequences on expression and secretion of human tissue plasminogen

activator in the baculovirus system. *J. Biol. Chem.* **268**, 16754–16762 (1993).

26. Espah Borujeni, A. & Salis, H. M. Translation initiation is controlled by RNA folding kinetics

via a ribosome drafting mechanism. *J. Am. Chem. Soc.* **138**, 7016–7023 (2016).

27. Salis, H. M., Mirsky, E. A. & Voigt, C. A. Automated design of synthetic ribosome binding

sites to control protein expression. *Nat. Biotechnol.* **27**, 946–950 (2009).

28. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**,

583–589 (2021).

29. Konur, O. *Handbook of Algal Science, Technology and Medicine.* (Academic Press, 2020).

30. Lindhauer, N. S. *et al.* Antibacterial activity of a defensin in an infection model of. *Virulence*

10, 902–909 (2019).

31. Bertrams, W. *et al.* Tribolium castaneum defensin 1 kills Moraxella catarrhalis in an in vitro

infection model but does not harm commensal bacteria. *Virulence* vol. 12 1003–1010

Preprint at https://doi.org/10.1080/21505594.2021.1908741 (2021).

32. Rodloff, A. C., Goldstein, E. J. C. & Torres, A. Two decades of imipenem therapy. *J.*

Antimicrob. Chemother. **58**, 916–929 (2006).

33. Kulp, A. & Kuehn, M. J. Biological functions and biogenesis of secreted bacterial outer

membrane vesicles. *Annu. Rev. Microbiol.* **64**, 163–184 (2010).

34. Bauwens, A., Kunsmann, L., Karch, H., Mellmann, A. & Bielaszewska, M. Antibiotic-

Mediated Modulations of Outer Membrane Vesicles in Enterohemorrhagic Escherichia coli

O104:H4 and O157:H7. *Antimicrob. Agents Chemother.* **61**, (2017).

35. Manning, A. J. & Kuehn, M. J. Contribution of bacterial outer membrane vesicles to innate

bacterial defense. *BMC Microbiol.* **11**, 258 (2011).

36. UniProt Consortium. UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids*

Res. **49**, D480–D489 (2021).

37. Witten, J. & Witten, Z. Deep learning regression model for antimicrobial peptide design.

bioRxiv 692681 (2019) doi:10.1101/692681.

38. Wang, G., Li, X. & Wang, Z. APD3: the antimicrobial peptide database as a tool for

research and education. *Nucleic Acids Res.* **44**, D1087–93 (2016).

39. Novković, M., Simunić, J., Bojović, V., Tossi, A. & Juretić, D. DADP: the database of anuran

defense peptides. *Bioinformatics* **28**, 1406–1407 (2012).

40. Pirtskhalava, M. *et al.* DBAASP v.2: an enhanced database of structure and

antimicrobial/cytotoxic activity of natural and synthetic peptides. *Nucleic Acids Res.* **44**,

6503 (2016).

41. Fan, L. *et al.* DRAMP: a comprehensive data repository of antimicrobial peptides. *Sci. Rep.*

6, 1–7 (2016).

42. Piotto, S. P., Sessa, L., Concilio, S. & Iannelli, P. YADAMP: yet another database of

antimicrobial peptides. *Int. J. Antimicrob. Agents* **39**, 346–351 (2012).

43. UniProt. https://www.uniprot.org/help/uniprotkb.

44. Ghorbani, M., Prasad, S., Brooks, B. R. & Klauda, J. B. Deep attention based variational

autoencoder for antimicrobial peptide discovery. Preprint at

https://doi.org/10.1101/2022.07.08.499340.

45. Bowman, S. R. *et al.* Generating sentences from a continuous space. (2015)

doi:10.48550/ARXIV.1511.06349.

46. Abraham, M. J. *et al.* GROMACS: High performance molecular simulations through multi-

level parallelism from laptops to supercomputers. *SoftwareX* **1-2**, 19–25 (2015).

47. Huang, J. *et al.* CHARMM36m: an improved force field for folded and intrinsically

disordered proteins. *Nat. Methods* **14**, 71–73 (2017).

48. Hess, B., Bekker, H., Berendsen, H. J. C. & Fraaije, J. G. E. M. LINCS: A linear constraint

solver for molecular simulations. *J. Comput. Chem.* **18**, 1463–1472 (1997).

49. Essmann, U. *et al.* A smooth particle mesh Ewald method. *J. Chem. Phys.* **103**, 8577–8593

(1995).

50. Bussi, G., Donadio, D. & Parrinello, M. Canonical sampling through velocity rescaling. *J.*

Chem. Phys. **126**, 014101 (2007).

51. Berendsen, H. J. C., Postma, J. P. M., van Gunsteren, W. F., DiNola, A. & Haak, J. R.

Molecular dynamics with coupling to an external bath. *J. Chem. Phys.* **81**, 3684 (1984).

52. Parrinello, M. & Rahman, A. Polymorphic transitions in single crystals: A new molecular

dynamics method. *J. Appl. Phys.* **52**, 7182 (1981).

53. Humphrey, W., Dalke, A. & Schulten, K. VMD: visual molecular dynamics. *J. Mol. Graph.*

14, 33–8, 27–8 (1996).

54. Schrödinger, L., & DeLano, W. PyMOL, Available at: http://www.pymol.org/pymol, (2020).

55. Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators,

and developers. *Protein Sci.* **30**, 70–82 (2021).

56. Schaefer, S. L., Jung, H. & Hummer, G. Binding of SARS-CoV-2 Fusion Peptide to Host

Endosome and Plasma Membrane. *J. Phys. Chem. B* **125**, 7732–7741 (2021).

57. Lorent, J. H. *et al.* Plasma membranes are asymmetric in lipid unsaturation, packing and

protein shape. *Nat. Chem. Biol.* **16**, 644–652 (2020).

58. Berglund, N. A. *et al.* Interaction of the antimicrobial peptide polymyxin B1 with both

membranes of E. coli: a molecular dynamics study. *PLoS Comput. Biol.* **11**, e1004180

(2015).

59. Jo, S., Kim, T., Iyer, V. G. & Im, W. CHARMM-GUI: a web-based graphical user interface

for CHARMM. *J. Comput. Chem.* **29**, 1859–1865 (2008).

60. Jo, S., Kim, T. & Im, W. Automated builder and database of protein/membrane complexes

for molecular dynamics simulations. *PLoS One* **2**, e880 (2007).

61. Jorgensen, W. L., Chandrasekhar, J., Madura, J. D., Impey, R. W. & Klein, M. L.

Comparison of simple potential functions for simulating liquid water. *J. Chem. Phys.* **79**, 926

(1983).

62. Blaskovich, M. A. T. *et al.* Protein-inspired antibiotics active against vancomycin- and

daptomycin-resistant bacteria. *Nat. Commun.* **9**, 22 (2018).

829 63. Ponath, V. *et al.* Secreted ligands of the NK cell receptor NKp30: B7-H6 is in contrast to

830   BAG6 only marginally released via extracellular vesicles. *Int. J. Mol. Sci.* **22**, 2189 (2021).

831 64. Butler, T. The Jarisch-Herxheimer Reaction After Antibiotic Treatment of Spirochetal

832   Infections: A Review of Recent Cases and Our Understanding of Pathogenesis. *Am. J.*

833   *Trop. Med. Hyg.* **96**, 46–52 (2017).

834 65. Griss, K. *et al.* MicroRNAs Constitute a Negative Feedback Loop in Streptococcus

835   pneumoniae-Induced Macrophage Activation. *J. Infect. Dis.* **214**, 288–299 (2016).

836 66. King, D. S., Fields, C. G. & Fields, G. B. A cleavage method which minimizes side reactions

837   following Fmoc solid phase peptide synthesis. *Int. J. Pept. Protein Res.* **36**, 255–266

838   (1990).

839 67. Fujii, N., Otaka, A., Sugiyama, N., Hatano, M. & Yajima, H. Studies on Peptides. CLV.

840   Evaluation of Trimethylsilyl Bromide as a Hard-Acid Deprotecting Reagent in Peptide

841   Synthesis. *Chem. Pharm. Bull.* **35**, 3880–3883 (1987).

842 68. Pearson, D. A., Blanchette, M., Baker, M. L. & Guindon, C. A. Trialkylsilanes as scavengers

843   for the trifluoroacetic acid deblocking of protecting groups in peptide synthesis. *Tetrahedron*

844   *Lett.* **30**, 2739–2742 (1989).