Spatiotemporal brain hierarchies of auditory memory recognition and predictive coding

Bonetti, L.[1,2,3,4*], Fernández-Rubio, G.[1], Carlomagno, F.[1,5], Dietz, M. J.[6], Pantazis, D.[7], Vuust, P.[1], Kringelbach, M. L.[1,2,3]

[1] Center for Music in the Brain, Department of Clinical Medicine, Aarhus University & The Royal Academy of Music, Aarhus/Aalborg, Denmark

[2] Centre for Eudaimonia and Human Flourishing, Linacre College, University of Oxford, Oxford, United Kingdom

[3] Department of Psychiatry, University of Oxford, Oxford, United Kingdom

[4] Department of Psychology, University of Bologna

[5] Department of Education, Psychology, Communication, University of Bari Aldo Moro, Italy

[6] Center of Functionally Integrative Neuroscience, Institute of Clinical Medicine, Aarhus University, Denmark

[7] McGovern Institute for Brain Research, Massachusetts Institute of Technology (MIT), MA 02139, Cambridge, USA

*Corresponding author: leonardo.bonetti@psych.ox.ac.uk

Abstract

Our brain is constantly extracting, predicting, and recognising key spatiotemporal features of

the physical world in order to survive. While neural processing of visuospatial patterns has

been extensively studied, the hierarchical brain mechanisms underlying conscious recognition

of auditory sequences and the associated prediction errors remain elusive. Using

magnetoencephalography (MEG), we studied the brain functioning of 83 participants during

recognition of previously memorised musical sequences and systematic variations. The

results showed feedforward connections originating from auditory cortices, and extending to

the hippocampus, anterior cingulate gyrus, and medial cingulate gyrus. Simultaneously, we

observed backward connections operating in the opposite direction. Throughout the

sequences, the hippocampus and cingulate gyrus maintained the same hierarchical level,

except for the final tone, where the cingulate gyrus assumed the top position within the

hierarchy. The evoked responses of memorised sequences and variations engaged the same

hierarchical brain network but systematically differed in terms of temporal dynamics,

strength, and polarity. Furthermore, induced-response analysis showed that alpha and beta

power was stronger for the variations, while gamma power was enhanced for the memorised

sequences. This study expands on the predictive coding theory by providing quantitative

evidence of hierarchical brain mechanisms during conscious memory and predictive

processing of auditory sequences.

Keywords

Recognition memory, Auditory sequences, Predictive coding (PC), Magnetoencephalography

(MEG)

Introduction

The spatiotemporal features of hierarchical processing in the brain are essential to fully grasp the neural substrates of human perception and cognition, as suggested by the predictive coding theory (PCT) [1-5, 6, 7]. To elucidate such brain mechanisms, much research has focused on the visual system [8, 9], which primarily relies on the recognition of patterns arranged in space. Conversely, the auditory system extracts information from patterns and sequences that develop over time [10], providing unique opportunities to understand the temporal hierarchies of the brain.

Extensive research spanning decades has established the hierarchical organisation of auditory perception, with a particular emphasis on the processing of elementary auditory stimuli. This hierarchical progression starts at the peripheral level within the cochlea and moves forward to the auditory pathway, which encompasses the brainstem, pons, trapezoid body, superior olivary complex, lateral lemniscus, inferior and medial geniculate nucleus of the thalamus, and ends in the primary auditory cortex [11, 12]. However, how auditory information is integrated from the auditory cortex to the whole brain has not been fully established yet.

Aiming to bridge this gap, prior investigations focused on automatic predictive processes within the framework of PCT. These studies assessed auditory automatic prediction error, often relying on well-established event-related potential/field (ERP/F) components such as N100, mismatch negativity (MMN), and error-related negativity (ERAN) [13-21]. They demonstrated that such components were automatically evoked in response to auditory stimuli, deviations from expected sound features, likelihood of occurrence of musical tones (N100, MMN), and varied harmonic properties (ERAN). Additional studies employed dynamic causal modelling (DCM) to investigate the brain hierarchical architecture during automatic predictive processes, as indexed by MMN. This research has yielded quantitative evidence demonstrating the flow of information from the primary auditory cortex to the superior temporal gyrus and the inferior frontal gyrus [22].

Expanding upon these investigations, Rocchi and colleagues provided evidence of hierarchical organisation in the auditory system using direct electrical brain stimulation. They showed effective connectivity from the auditory cortex to regions in the medial temporal lobe and prefrontal cortex, including the hippocampus and ventro-lateral prefrontal cortex [23]. Yet, it remains uncertain how the integration of auditory information across the whole brain

relates to complex cognitive functions, such as the conscious recognition and prediction of
auditory sequences evolving over time.

To address this question, a unique perspective has arisen through the integration of
musical memory paradigms with magnetoencephalography (MEG). Music is a complex
artform acquiring meaning through the combination of its constituent elements extended over
time [24], while MEG, particularly when used in combination with MRI, allows to track the
brain activity with excellent temporal resolution in the order of milliseconds and acceptable
spatial accuracy [25]. It is precisely due to these attributes that this combination offers a unique
framework for investigating the rapid memory and predictive processes of temporal
information in the human brain [7, 26].

Indeed, recent studies employing MEG and musical paradigms investigated the brain
mechanisms associated with perception, manipulation and recognition of sound sequences.
As an example, Albouy and colleagues [27] explored the brain activity underlying memory
retention, showing that theta oscillations in the dorsal stream of the participants' brain
predicted their auditory WM performance. Along this line, we recently uncovered a vast
network of brain areas responsible for encoding and recognising previously memorised and
novel musical sounds. This network extended from the auditory cortex to the medial
cingulate, inferior temporal cortex, insula, frontal operculum, and hippocampus [28-30]. We also
observed that the complexity of music [31] and individual cognitive differences [32] influenced
the activity recorded in this network.

In our previous studies, we utilised relatively rapid musical tones with a duration of 250
ms. However, this approach prevented us from closely tracking the neural dynamics
associated with each sound within the sequences, leading to frequent overlap with the brain
responses to subsequent tones. For this reason, we were also unable to study the rapid
hierarchies between the brain regions that we revealed, which is necessary to understand the
neurophysiology of conscious recognition of auditory sequences.

Moreover, while our previous works showed differences in brain activity when comparing
previously memorised versus completely novel auditory sequences, we did not investigate the
scenario in which an original sequence was systematically varied. This prevented us from
exploring how the brain generates prediction errors within the framework of a conscious
long-term memory recognition task, as opposed to the several instances of automatic
prediction errors described in the literature [13-21].

To address these questions, in the present study we made two essential modifications to
our previously employed experimental paradigm: (i) we compared the original five-tone
melodies with systematic variations occurring at the 2^{nd}, 3^{rd}, 4^{th}, or 5^{th} tone; (ii) we increased
the duration of the tones from 250 ms to 350 ms. These seemingly minor modifications were
of great importance, as they enabled us to directly address our three main hypotheses, stated
as follows. First, the conscious recognition of previously memorised auditory sequences
relied on the hierarchical recruitment of a widespread brain network including auditory
cortex, hippocampus and cingulate gyrus. We expected to observe that these regions
responded to each tone forming the sequence, with the auditory cortex preceding the
hippocampus and cingulate gyrus. Second, the detection of the varied auditory sequences
elicited a prediction error, originating in the auditory cortex and then propagating to
hippocampus, anterior cingulate and ventromedial prefrontal cortex. While we expected a
consistent response in the auditory cortex to the introduction of varied sounds at any point in
the sequence, we hypothesised that the anterior cingulate gyrus and ventromedial prefrontal
cortex exhibited their strongest response exclusively when the variation in the sequence was
introduced. Third, sequence recognition relied on a brain hierarchy characterised by
feedforward connections from auditory cortices to hippocampus and cingulate gyrus,
simultaneous with feedback connections in the opposite direction. Moreover, recognition of
both previously memorised and varied sequences relied on the same hierarchy, while the
temporal dynamics, strength and polarity of brain responses differed.

These hypotheses were confirmed by the results of the experiment, which showed that the
brain distinctively responds to each tone forming the previously memorised sequence, with
the auditory cortex preceding hippocampus and anterior and medial cingulate gyrus. In
addition, the detection of the sequence variations elicits a conscious prediction error in the
brain, originating in the auditory cortex and propagating to hippocampus, anterior and medial
cingulate. While the auditory cortex has a constant response to the varied sounds introduced
at any point in the sequence, the hippocampus and anterior and medial cingulate gyrus exhibit
their strongest response exclusively to the sound which introduced the variation in the
sequence. Finally, we provide quantitative evidence of the brain functional hierarchies during
long-term auditory recognition. Here, as hypothesised, feedforward connections originate
from the auditory cortices and extend to the hippocampus, anterior cingulate gyrus, and
medial cingulate gyrus, along with feedback connections in the opposite direction. Notably,
throughout the sequence, the hippocampus and cingulate gyrus maintain the same

hierarchical level, except for the final tone, where the cingulate gyrus assumes the top

position within the hierarchy.

Results

Experimental design and behavioural results

Eighty-three participants performed an old/new auditory recognition task during MEG recordings. After learning a short musical piece (**Figure S1**), participants were presented with 135 five-tone musical sequences lasting 1750 ms each and were requested to state whether each sequence belonged to the original music ('memorised' sequence (M), old) or it was a varied sequence ('novel' sequence (N), new) (**Figure 1a**).

Twenty-seven sequences were extracted from the original musical piece and 108 were variations of the original melodies. We organised these variations in four categories depending on whether changes involved every musical tone of the sequence after the first (NT1), second (NT2), third (NT3) or fourth (NT4) tone (**Figure 1b** and **Figure S2**).

Figure 1. Experimental design, stimuli, and analysis pipeline

a – The brain activity of 83 participants was collected using magnetoencephalography (MEG) while they
performed an auditory old/new recognition task. One at a time, five-tone auditory sequences were presented in
randomised order and participants were instructed to respond with button presses whether they were 'old'
*(memorised musical sequences, 'M') or 'new' (novel musical sequences, 'N'). **b** – Five types of auditory*
*sequences (M, NT1, NT2, NT3, NT4) were used in the study (see **Figure S2** for the full set of sequences). The N*
sequences were created through systematic variations of the M sequences. This procedure consisted of changing
every musical tone of the sequence after the first (NT1), second (NT2), third (NT3) or fourth (NT4) tone, as
*illustrated by the red musical tones. **c** – After MEG data pre-processing, multivariate pattern analysis was used*
to assess whether it was possible to discriminate the experimental conditions based on the neural activity
*recorded with MEG. **d** – Univariate analyses on the MEG channels that primarily contributed to the decoding*
algorithm were performed using t-tests to compare the M condition with each of the N conditions (i.e. M versus
*NT1, M versus NT2, etc.). **e** – The MEG data was co-registered with the individual anatomical magnetic*

resonance imaging (MRI) data, and source reconstructed using a beamforming algorithm. This resulted in one
time series for either the 3559 reconstructed brain sources (8-mm parcellation) or for the 90 non-cerebellar
automated anatomical labelling (AAL) regions of interest (ROIs). The reconstructed brain activity was
*contrasted across experimental conditions (M versus each category of N). **f** – Evoked and induced responses*
*were computed for the AAL ROIs and contrasted across experimental conditions. **g** – Dynamic causal modelling*
was applied to a restricted set of AAL ROIs: left Heschl's gyrus (LHG), right Heschl's gyrus (RHG), left
hippocampus (LHP), right hippocampus (RHP), anterior cingulate gyrus (ACC), and medial cingulate gyrus
(MC), to assess the functional hierarchical organisation of the brain during the recognition of auditory
sequences.

We performed statistical analyses on the MEG task behavioural data (see **Table 1** for
descriptive statistics). We computed two independent Kruskal-Wallis H tests to assess
whether the five categories of temporal sequences (M, NT1, NT2, NT3, and NT4) differed in
terms of response accuracy and reaction times (**Figure S3**).

The Kruskal-Wallis H test for response accuracy was significant ($H(4) = 36.38$, $p < .001$),
indicating a difference between categories in the number of correct responses. The Tukey-
Kramer correction for multiple comparisons highlighted that NT4 trials were correctly
identified with a lower frequency than M ($p = .001$), NT1 ($p = .001$), NT2 ($p = .0003$), and
NT3 trials ($p < .0001$).

The Kruskal-Wallis H test for the reaction times was also significant ($H(4) = 22.53$, $p =
.0002$). The Tukey-Kramer correction for multiple comparisons highlighted that NT4 trials
were correctly identified with greater reaction times than M ($p = .0016$), NT1 ($p = .0013$),
NT2 ($p = .0054$), NT3 trials ($p = .0008$).

Behavioral variables	M	NT1	NT2	NT3	NT4
Correct recognition	22.33±5.30	22.36±4.27	21.58±5.31	21.66±5.34	17.04±7.12
Reaction times (ms)	2426±226	2407±284	2431±282	2415±272	2578±259

196 *Table 1. MEG task behavioural results*
197 *Mean and standard deviations across participants of number of correctly recognised trials and reaction times*
198 *(ms) for the five experimental conditions (previously memorised [M], novel T1 [NT1], NT2, NT3, NT4).*
199 *Participants were presented with 27 M, 27 NT1, 27 NT2, 27 NT3, and 27 NT4 trials.*

Multivariate pattern analysis on the MEG channels

Using a support vector machine (SVM) classifier (see details in Methods, and **Figure 1c**), we performed multivariate pattern analyses to decode different neural activity associated with the recognition of M versus N. Specifically, we computed four independent analyses, decoding M versus each of the four categories of novel sequences (i.e., M versus NT1, M versus NT2, M versus NT3, and M versus NT4).

As shown in **Figure 2a** and **Figure S4**, each of these analyses returned a decoding time series showing how the neural activity differentiated the pair of experimental conditions. Overall, the results showed that the SVM was able to detect significant differences between M and N sequences. As illustrated in **Figure 2a**, decoding M versus NT1 returned the following main significant time windows: 0.53 - 0.73 sec; 0.91 – 0.95 sec; 1.27 – 1.30 sec; 1.62 – 1.88 sec ($p < .012$, false-discovery rate [FDR]-corrected). Decoding M versus NT2 gave rise to the following main significant time windows: 0.89 – 1.18 sec; 1.26 – 1.42 sec; 1.54 – 1.89 sec ($p < .012$, FDR-corrected). Decoding M versus NT3 returned one main significant time window: 1.25 - 2.07 sec ($p < .012$, FDR-corrected). Finally, decoding M versus NT4 showed the following main significant time window: 1.64 - 2.07 ($p < .012$, FDR-corrected). The contribution of each MEG channel to the decoding algorithm in the significant time-windows is depicted in **Figure 2b**. Detailed statistical results are reported in **Table S1** and illustrated in **Figure S4**.

To evaluate the persistence of discriminable neural information over time, we used a temporal generalisation approach by training the SVM classifier at a given time-point *t* and testing it across all time-points. This was calculated for the four pairs of experimental conditions described above. The signed-rank test against chance level and cluster-based Monte-Carlo simulations [30, 33-36] (MCS; $\alpha = .05$, MCS *p-value* $= .001$) showed that the performance of the classifier was significantly above chance even a few hundreds of milliseconds beyond the diagonal, for all pairs of conditions. Notably, the neural difference between M and N was comparable across diverse musical tones, as shown by the recurrent patterns depicted in **Figure 2c** and highlighted by the red graphs. Detailed statistical results are reported in **Table S2**.

233
234

Figure 2. Multivariate pattern analysis and cluster-based univariate analysis on MEG channels

a - Multivariate pattern analysis decoding the different neural activity associated with memorised versus novel
musical sequences. The plot shows the decoding time series (50% chance level) for four rounds of pairwise
decoding (memorised [M] versus novel T1 [NT1], M versus NT2, M versus NT3, M versus NT4). The sketch of
the musical tones represents the onset of the sounds forming the temporal sequences. The blue-black lines
indicate the significant time-points when the algorithm successfully decoded the experimental conditions (p <
*.012, false-discovery rate [FDR]-corrected). **b** – Activation patterns estimated by the decoding weights (without*
unit of measure) representing the relative contribution of MEG channels during significant time-points (as
*indicated in **a**. **c** – Temporal generalisation of pairwise decoding results, tested against a 50% chance level.*
Plots display the significant time-points (either yellow or dark blue) where conditions were successfully
*distinguished after cluster-based Monte-Carlo simulations [MCS] correction for multiple comparisons. **d** –*
Univariate analysis computed on the magnetometers which primarily contributed (one standard deviation plus
the mean over magnetometers) to the decoding algorithm. After averaging the magnetometers channels in two
independent groups (see Methods for details), t-tests were conducted for each time-point and M versus NTs pair
(i.e., M versus NT1, M versus NT2, M versus NT3, M versus NT4), and corrected for multiple comparisons
using cluster-based MCS (MCS, α = 0.05, MCS p-value = 0.001). Blue-black lines highlight the temporal extent
of the significant differences between M and N conditions. Different hues of blue-black show specific M versus
*N condition comparisons. **e** – Same procedure observed for **d** was performed on gradiometers. All the time*
series illustrated in this figure correspond to averages over participants, while shaded areas indicate standard
errors. For all plots, musical tone sketches indicate sounds sequence onset.

255
256

Univariate analysis on the MEG channels
Pairwise decoding is a powerful technique, but it can only tell if two conditions are
characterised by significantly different brain activity and does not provide information about
which condition presented the strongest brain activity. To answer this question, we computed
an additional analysis focusing on the MEG channels showing the strongest weights (one
standard deviation plus the mean) in the decoding algorithm (**Figure 1d**). This was done
independently for magnetometers and gradiometers and returned 16 magnetometers and 32
gradiometers (see Methods for the channels number).
These MEG channels were then averaged based on the channel type (magnetometers or
gradiometers) and the polarity of the signal (positive or negative orientation of the N100 to
the first sound of the sequence). Using this procedure, four distinct groups were created: (i)
magnetometers negative N100 (**Figure 2d**, left), (ii) magnetometers positive N100 (**Figure
2d**, right), (iii) gradiometers negative N100 (**Figure 2e**, left), (iv) gradiometers positive N100
(**Figure 2e**, right) (further details are provided in the Methods section).

Then, independently for each group, we computed one t-test for each time-point and each

combination of M versus Ns (i.e., M versus NT1, M versus NT2, M versus NT3, M versus

NT4). We corrected for multiple comparisons using a one-dimensional (1D) cluster-based

MCS (MCS α = .05, MCS p-value = .001). As reported in detail in **Table S3**, the results

showed significant differences between experimental conditions which aligned closely in

timing with the decoding analyses reported in the previous section, indicating consistency

between the two methods.

To provide full disclosure of our data, we complemented these computations with the

same univariate analysis calculated independently for each MEG channel. As shown in

Figure S5$_{a-z2}$ and **Table S4**, the significant differences between experimental conditions were

coherent and spread across a large array of MEG channels.

Neural sources of the MEG channels peak activity

First, we computed source reconstruction by employing a local-spheres forward model and

a beamforming approach as inverse solution in an 8-mm grid (corresponding to 3559 brain

voxels), using magnetometers only (**Figure 1e**). This procedure returned a time series for

each of the reconstructed bran voxel, which is commonly referred to as neural activity index

(see Methods for extensive details on this standard procedure).

Second, we extracted the time indices of the minimum and maximum value recorded by

the magnetometers shown in **Figures 2d** and **3** in a 400 ms time-window following the onset

of each tone. The duration of this time window reflects the period during which independent

responses to each sound occurred. This procedure returned the time index of two neural peaks

for each experimental condition and each tone of the sequence (with exclusion of the 1st one

which did not differ across conditions). This revealed that one peak of the neural activity

occurred around 350 ms after the onset of each tone (highlighted in yellow in **Figure 3a** and

3b). With regards to this peak, condition M showed the strongest response among all

conditions. The other peak occurred in response to the tone which introduced the variation in

the musical sequence (highlighted in purple in **Figure 3a** and **3b**). In this case, the N

condition which introduced the variation reported the strongest response among all other

conditions. This occurred approximately at 250 ms after the varied tone (i.e. NT1 showed the

strongest response to tone two among all conditions, NT2 to tone three, NT3 to tone four and

NT4 to tone five). Extensive details on these neural activity peaks are reported in **Table 2**.

Third, for each of the defined peaks, we computed contrasts in MEG source space between

M and N. More specifically, with regards to the peaks occurring approximately at 350 ms

after each tone, we contrasted M versus NT1 (illustrated in the yellow boxes in **Figure 3**).

This was done because the peaks occurring 350 ms after each tone were always greater for

M, and NT1 was the only condition comprising sounds that were always different from M.

Similarly, with regards to the peaks occurring approximately at 250 ms after the onset of the

varied tones, we contrasted the N condition which showed the strongest peak against M

(illustrated in the purple boxes in **Figure 3**). Here, for each condition in the pair, we averaged

the absolute value of the activity of each of the 3559 reconstructed brain voxels in the ± 20

ms time-window around the correspondent time index. Then, we computed t-tests for each of

the 3559 brain voxels by contrasting the pairs of conditions and FDR-corrected for multiple

comparisons. The results of these analyses are shown in **Figure 3** and reported in detail in

Table 2 and **Table S5**. In addition, we reported the neural activity recorded by the MEG

channels in topographic maps in **Figure S6**.

Figure 3. Neural activity peaks recorded by magnetometers and contrasts in magnetoencephalography
(MEG) 8-mm source space

a – Exactly as illustrated in **Figure 2**, *the waveform plot displays the time series averaged over participants for*
the following groups of MEG channels: (i) magnetometers negative N100, and (ii) magnetometers positive
N100. Shaded areas indicate the standard errors. In addition, here we have marked the maximum and minimum
peaks of neural activity following the onset of each tone (excluding the first tone). These peaks occurred
approximately at 250 ms (purple circle) and 350 ms (yellow circle) after each tone's onset. **b -** *Within a time*
window of ±20 ms around each peak, we averaged the brain activity across all 3559 reconstructed brain voxels
for each condition. In the yellow boxes, we represent the significant brain voxels obtained by contrasting the
activity associated with memorised (M) versus novel T1 (NT1). These contrasts were done by using t-tests and
false discovery rate (FDR) to correct for multiple comparisons. Purple boxes depict the same analysis, but for
comparisons involving M versus NT1 (tone 2), M versus NT2 (tone 3), M versus NT3 (tone 4), and M versus
NT4 (tone 5). In both cases, the red colour shows the brain voxels (t-values) characterised by a stronger activity
for M versus NTs, while the blue colour indicates the opposite pattern. For a detailed statistical report, consult
Table S5.

336
337

338 Furthermore, the brain activity recorded by the magnetometers for each peak and
339 hemisphere was correlated with the measure of participants' musical training provided by the
340 Goldsmith Musical Sophistication Index (GOLD-MSI) [37] and corrected for multiple
341 comparisons using FDR (**Table 2**).
342

Tone #	Tone 2		Tone 3		Tone 4		Tone 5	
Peak	Max	Min	Max	Min	Max	Min	Max	Min
Magnetometers in the left hemisphere								
Time (s)	0.708 (M)	0.612 (NT1)	1.092 (M)	0.992 (NT2)	1.452 (M)	1.340 (NT3)	1.804 (M)	1.736 (NT4)
Musical expertise (FDR-corrected)	r = .093 p = .473	r = -.218 p = .084	r = .239 p = .066	r = -.186 p = .132	r = .286 p = .040	r = -.261 p = .050	r = .344 p = .014	r = -.034 p = .768
Contrasts in MEG source space								
Conditions	M versus NT1	M versus NT1	M versus NT1	M versus NT2	M versus NT1	M versus NT3	M versus NT1	M versus NT4
Max t-val	5.79	n.s	5.45	-6.09	5.81	-6.56	5.1	-6.03
FDR-adjusted p-value	.003	n.s	.009	.008	.01	.01	.01	.02
MNI coordinates	-6 -22 64	n.s	-6 -30 40	2 34 0	-6 -38 32	2 18 -16	26 -30 -8	-6 26 -16
AAL region	Supp Motor Ar	n.s	Cingulum Mid	Cingulum Ant	Cingulum Mid	Subgenual	Hippocampus	Front Med Orb
Magnetometers in the right hemisphere								

Time (s)	0.620 (NT1)	0.832 (M)	0.988 (NT2)	1.088 (M)	1.352 (NT3)	1.456 (M)	1.732 (NT4)	1.920 (M)
Musical expertise (FDR-corrected)	r = .242 p = .061	r = .051 p = .819	r = .428 p < .001	r = -.034 p = .819	r = .459 p < .001	r = -.050 p = .819	r = .257 p = .050	r = -.026 p = .819
Contrasts in MEG source space								
Conditions	M versus NT1	M versus NT1	M versus NT2	M versus NT1	M versus NT3	M versus NT1	M versus NT4	M versus NT1
Max t-val	n.s	4.41	-6.14	5.50	-7.13	5.61	-6.17	5.24
FDR-adjusted p-value	n.s	.001	.008	.01	.02	.01	.02	.01
MNI coordinates	n.s	58 -14 -16	2 34 -8	-6 -30 40	2 18 -16	-6 -38 32	-6 26 -16	10 -46 48
AAL region	n.s	Temporal Mid	Cingulum Ant	Cingulum Mid	Subgenual	Cingulum Mid	Front Med Orb	Precuneus

Table 2. Neural activity peaks recorded by magnetometers, musical expertise and contrasts in MEG 8-mm source space

The table shows the time index of the maximum and minimum peaks recorded by the magnetometers (refer to Figure 3) for each tone (excluding the 1st one). Additionally, the table indicates the condition for which the strongest peaks were observed (in parenthesis). The table also illustrates the correlations between the absolute value of the neural peaks and the participants' musical training (r = Pearson's correlation coefficient, p-values are adjusted by the false discovery rate [FDR] correction for multiple comparisons). Finally, the table provides details of the MEG source contrasts in 8-mm space. This includes the conditions which were contrasted, t-value, FDR-adjusted p-value, Montreal Neurological Institute (MNI) coordinates, and automated anatomical labelling (AAL) region of the brain voxel which showed the strongest difference between the experimental conditions.

Automated anatomical labelling (AAL) time series

After evaluating the neural sources of the different brain activity associated with the experimental conditions using the fine-grained 8-mm parcellation of the brain (**Figure 1e**, *"8-mm parcellation"*), we focused on a set of anatomically defined regions of interest (ROIs) (**Figure 1e**, *"AAL parcellation"*). Here, we used a standard anatomical parcellation method known as automated anatomical labelling (AAL) [38] and calculated a time series for each of the 90 non-cerebellar ROIs of AAL (**Figure 1f**, left, *"Evoked responses"*).

As specified above, the source reconstruction was conducted in an 8-mm space, resulting in 3559 brain voxels, each with its own time series. Afterwards, for each of the 90 ROIs, we identified the corresponding brain voxels and then averaged the time series of these associated voxels. This allowed us to obtain a final time series for each AAL ROI.

Then, we computed one t-test for each AAL ROI, each time-point and each combination
of M versus Ns, and we corrected for multiple comparisons using 1D cluster-based MCS
(MCS, α = .05, MCS *p*-value = .001). Confirming the results from the 8-mm brain
parcellation shown in **Figure 3**, this analysis returned several significant clusters of
differential brain activity over time between M and Ns. These were primarily localised in
auditory cortices (e.g. Heschl's gyrus, superior temporal gyrus), hippocampus and para-
hippocampal gyrus, inferior temporal cortex, anterior, medial and posterior cingulate gyrus,
and ventromedial prefrontal cortex. **Figure S7$_{a-g}$** and **Tables S6** and **S7** show all AAL ROIs
and the significant differences between conditions. **Figures 4** and **S8** shows instead a selected
array of ROIs which were particularly relevant for this study. Here, we identified the two
ROIs that showed the maximum activity in absolute value among auditory regions (i), medial
temporal lobe (ii) and cingulate and prefrontal cortices (iii). We selected these broad brain
regions based on the cognitive processes involved in the experimental task used in this study:
audition (i), memory (ii), evaluation and decision-making (iii). As shown by **Figures 4** and
S8, the selected AAL ROIs were left and right Heschl's gyrus (LHG, RHG), left and right
hippocampus (LHP, RHP), anterior cingulate gyrus (ACC, left and right averaged together)
and medial cingulate gyrus (MC, left and right averaged together), respectively.

In essence, the contrasts showed that M versus N was characterized by stronger activity in
RHG, LHP, RHP and ACC at 350 – 450 ms after the onset of each tone. Similarly, M
presented stronger negative activity than N in the MC at 400 – 500 ms after the onset of each
tone (**Figure 4**). Conversely, late N100 responses localised in LHG were stronger for N
versus M. Moreover, LHP, RHP and ACC showed a stronger response for N versus M
occurring at 250 – 300 ms after altering the original sequences (**Figure 4**). **Table 3** reports
detailed statistics of the largest significant clusters, while complete statistical results of these
six ROIs are described in **Table S7**.

Contrast	ROI	Temporal extent of the largest clusters from the 1st tone of the sequence (ms)	Peak t-value	P-value
Positive activity				
M versus NT1 (onset deviation NT1: 350 ms)	ACC	580 – 830	7.45	< .001
	LHG	520 – 630	7.17	< .001
	RHP	580 – 780	7.66	< .001
M versus NT2 (onset deviation NT2: 700 ms)	ACC	940 – 1160	7.48	< .001
	LHG	890 – 970	6.26	< .001
	RHP	950 – 1160	6.24	< .001

M versus NT3 (onset deviation NT3: 1005 ms)	ACC	1310 – 1540	6.13	< .001
	RHG	1660 – 1760	5.37	< .001
	RHP	1320 – 1520	6.72	< .001
M versus NT4 (onset deviation NT4: 1400 ms)	ACC	1700 – 1890	6.54	< .001
	LHG	1640 – 1720	6.58	< .001
	RHP	1680 – 1890	6.56	< .001
Negative activity				
M versus NT1 (onset deviation NT1: 350 ms)	MC	680 – 860	-6.06	< .001
M versus NT2 (onset deviation NT1: 700 ms)	MC	980 – 1180	-6.17	< .001
M versus NT3 (onset deviation NT1: 1005 ms)	MC	1360 – 1550	-5.96	< .001
M versus NT4 (onset deviation NT1: 1400 ms)	MC	1720 – 1920	-6.17	< .001

Table 3. Largest clusters of stronger activity of memorised (M) versus novel sequences (Ns).

Largest clusters of significantly stronger activity of M versus Ns computed for the six selected automated

anatomical (AAL) regions of interest (ROIs): left Heschl's gyrus (LHG), right Heschl's gyrus (RHG), left

hippocampus (LHP), right hippocampus (RHP), anterior cingulate gyrus (ACC), and medial cingulate gyrus

(MC). The table shows the contrast (including the onset of the first varied note in the N sequence), the

correspondent ROI, the temporal extent (in ms) of the largest cluster, the peak t-value of the cluster and the

associated Monte-Carlo simulations (MCS) p-value. The MC shows stronger negativity since the polarity of the

MC signal was primarily negative, thus stronger activity in the MC for M versus N was indicated by a more

pronounced negativity for M compared to N. All the other significant clusters, for both the selected six and the

*remaining AAL ROIs, are reported in detail in **Tables S6** and **S7**.*

In addition, we applied the multivariate source leakage correction method proposed by

Colclough and colleagues [39] (see Methods for details) to the ROIs time series and computed

the same statistical analysis described above. As shown in **Figure S9** and **Table S8**, the

results aligned with our initial findings, thereby corroborating their robustness.

Finally, as a further reliability measure, we reported in supplementary materials the

description of an alternative, functional parcellation derived from the data. As shown by

supplementary **Figures S7a-g, S10** and **S11** and **Tables S9**, the contrasts across experimental

conditions when using the functional parcellation returned results which were highly

comparable with AAL, demonstrating that the significance of our findings did not depend on

the chosen parcellation. All the subsequent analyses reported in the manuscript were

computed on the AAL parcellation. Moreover, we complemented them by performing the

same analyses using the functional parcellation, as reported in the supplementary materials

(**Figures S15, S16, S17, S18** and **Tables S14, S15**).

420
421

422   *Figure 4. Source-localised differences in evoked responses across experimental conditions*

 This figure illustrates the source-localised brain activity averaged over participants for each experimental
condition (memorised [M], novel T1 [NT1], NT2, NT3, NT4) within six selected automated anatomical labelling
(AAL) regions of interest (ROIs): left Heschl's gyrus (LHG), right Heschl's gyrus (RHG), left hippocampus
(LHP), right hippocampus (RHP), anterior cingulate gyrus (ACC), and medial cingulate gyrus (MC). The
shaded areas indicate standard errors. Musical tone sketches indicate the onset of the sounds forming the
sequences. Brain templates depict the spatial extent of the selected ROIs. Blue-black lines highlight the temporal
extent of the significant differences between M versus each category of Ns (i.e. M versus NT1, M versus NT2, M
versus NT3, M versus NT4). Different hues of blue-black correspond to specific M versus N condition
comparisons. For a comprehensive depiction of all AAL ROIs, please refer to **Figure S7a-g**. *For a detailed*
statistical report on significant differences between experimental conditions in all ROIs, consult **Table S6**.

Prediction error across brain regions of interest
After differentiating the brain activity of M versus N, we conducted an additional analysis to
investigate the peak responses elicited by the varied tones within the N conditions. Regarding
Heschl's gyrus (left and right), the peaks corresponded to the late N100, occurring
approximately at 150 ms after the onset of each varied tone. Regarding the hippocampus (left
and right), anterior and medial cingulate gyrus, the peaks occurred approximately at 250 ms
after the onset of each varied tone (these responses were negative for the hippocampus and
anterior cingulate gyrus and positive for the medial cingulate gyrus). We computed the
average neural activity within a ±20 ms time window centred around the designated time
index (depicted in **Figure 5** by red circles). Subsequently, we performed separate one-way
analyses of variance (ANOVAs) for each ROI and selected condition, with the sequential
neural peaks as independent variable and the neural activity as the dependent variable. Since
this approach led to a total of 18 ANOVAs (six ROIs for NT1, NT2, NT3), we applied FDR
correction to account for multiple comparisons. Moreover, the Tukey-Kramer correction was
applied to the post-hoc analysis computed for each ANOVA (**Table S10**). Results revealed
that the responses in Heschl's gyri to subsequent varied tones were not significantly different:
LHG_{NT1}: $F(3, 331) = 0.87$, $p = .51$; LHG_{NT2}: $F(2, 248) = 1.33$, $p = .37$; LHG_{NT3}: $F(1, 165) =$
0.68, $p = .51$; RHG_{NT1}: $F(3, 331) = 0.90$, $p = .51$; RHG_{NT2}: $F(2, 248) = 0.18$, $p = .83$;
RHG_{NT3}: $F(1, 165) = 0.33$, $p = .60$. In contrast, for the hippocampus and cingulate gyrus, the
responses to different varied tones were significantly different, highlighting the stronger
activity following the initial tone which introduced the sequence variation: LHP_{NT1}: $F(3, 331)$
$= 2.82$, $p = .05$; LHP_{NT2}: $F(2, 248) = 7.29$, $p = .002$; LHP_{NT3}: $F(1, 165) = 7.86$, $p = .01$;
RHP_{NT1}: $F(3, 331) = 7.40$, $p = 2.9e-04$; RHP_{NT2}: $F(2, 248) = 14.33$, $p = 1.3e-06$; RHP_{NT3}:

$F(1, 165) = 27.16$, $p = 5.56\text{e-}07$; ACC_{NT1}: $F(3, 331) = 4.98$, $p = .005$; ACC_{NT2}: $F(2, 248) =$

11.54, $p = 1.6\text{e-}05$; ACC_{NT3}: $F(1, 165) = 17.55$, $p = 2.05\text{e-}04$; MC_{NT1}: $F(3, 331) = 3.21$, $p =$

$.02$; MC_{NT2}: $F(2, 248) = 8.40$, $p = 2.9\text{e-}04$; MC_{NT3}: $F(1, 165) = 6.24$, $p = .02$.

Figure 5. Focus on prediction error: differentiating Heschl's gyrus from hippocampus and cingulate gyrus

This figure focuses on prediction error responses in three novel (N) experimental conditions (NT1, NT2, NT3)

within six selected automated anatomical labelling (AAL) regions of interest (ROIs): left Heschl's gyrus (LHG),

right Heschl's gyrus (RHG), left hippocampus (LHP), right hippocampus (RHP), anterior cingulate gyrus

(ACC), and medial cingulate gyrus (MC). The figure shows the time series averaged over participants. Shaded

*areas indicate standard errors. Musical tone sketches indicate the onset of the sounds forming the sequences. Brain templates illustrate the spatial extent of the selected ROIs. For each ROI, the circles highlight the peak responses indexing the prediction error to each of the varied tones in the sequences (occurring approximately at 150 ms after tone onset for LHG and RHG, and at 250 ms for LHP, RHP, ACC, MC. To be noted, LHP, RHP and ACC exhibit negative peaks, while MC displays positive peaks). Beneath each waveform, violin plots display the distribution of the amplitude of the peaks highlighted by the circles (each dot represents a participant). The different amplitude of these peaks was statistically tested using analyses of variance (ANOVAs), independently for each ROI. The stars indicate the significant ANOVAs after false discovery rate (FDR) correction for multiple comparisons (FDR-adjusted p-values: * < .05; ** < .01; *** < .001; n.s. = not significant).*

Dynamic causal modelling

In this study, DCM was used to test our hypothesised model of brain hierarchies during recognition of memorised and varied musical sequences against five competing models, which are described in detail in the Methods section and illustrated in **Figure 1g**. These models comprised the six AAL ROIs described in the previous section and depicted in **Figure S8**. Our hypothesised model consisted of feedforward connections from LHG and RHG to LHP, RHP, ACC, and MC, and feedback connections from LHP, RHP, ACC and MC to LHG and RHG (**Figure 6a**, red box).

Since DCM for MEG is usually computed on relatively short time-windows (e.g. 350 – 400 ms) and for one experimental perturbation (i.e. sound) at a time [40, 41], we computed a series of DCM analyses independently for each tone of the sequences, in a 350 ms time-window from each tone onset. Specifically, for each participant, we computed an independent DCM for tones two, three, four and five for condition M. In relation to the N conditions, we were interested in the tone that introduced the variation in the sequences. Thus, we computed four DCMs, organised as follows: tone two for NT1; tone three for NT2; tone four for NT3; tone five for NT4.

To compare alternative models, we estimated the free energy *F* associated with the computation of each model (model evidence). To study the consistency of *F* across the population, we used a random-effects Bayesian model selection (RFX BMS) procedure, which provides the posterior probability and protected exceedance probability of each model, indicating which of the alternative models has the strongest model evidence within the whole population. In addition, we computed the Bayesian omnibus risk (BOR), which represents the probability that the protected exceedance probability associated with each model comparison

is attributable to chance within the sample of participants (**Figure 6b** and **6c**) [40]. This
procedure was observed independently for the eight DCM analyses described above (four for
M and four for Ns).

Results indicated that our hypothesised model was the one with the highest evidence for
all the DCM analyses of tones two, three and four (**Figure 6**). This result showed a high
probability of non-being driven by random chance, as demonstrated by the BOR parameter:
$BOR_{M\text{-}tone2} = 2.91e\text{-}07$; $BOR_{M\text{-}tone3} = 4.33e\text{-}04$; $BOR_{M\text{-}tone4} = .008$; $BOR_{NT1\text{-}tone2} = 4.49e\text{-}12$;
$BOR_{NT2\text{-}tone3} = .010$; $BOR_{NT3\text{-}tone4} = .003$.

Regarding tone five, the model with the highest evidence for M showed a hierarchical
architecture which consisted of the auditory cortex leading to the hippocampus and anterior
cingulate gyrus and subsequently to the medial cingulate gyrus ($BOR_{M\text{-}tone5} = .006$). For NT4,
the model with the highest evidence presented a hierarchy which started in the auditory
cortex and led to the hippocampus and medial cingulate gyrus and then to the anterior
cingulate gyrus ($BOR_{NT4\text{-}tone5} = .036$).

519
520

Figure 6. Brain hierarchies during recognition of auditory sequences revealed by Dynamic Causal Modelling
(DCM)

a – Graphical depiction of the six alternative models employed in DCM analysis. The connections between the
six selected automated anatomical labelling (AAL) regions of interest (ROIs) (left Heschl's gyrus [LHG], right
Heschl's gyrus [RHG], left hippocampus [LHP], right hippocampus [RHP], anterior cingulate gyrus [ACC],
medial cingulate gyrus [MC]) are illustrated within a brain template, and through a graphical representation.
Our hypothesised model of brain hierarchies during recognition of memorised and varied musical sequences is
enclosed in a red box. b – Posterior probability, protected exceedance probability and Bayesian omnibus risk
(BOR) indicating the model with the highest evidence across the population (inference computed using random-
effects Bayesian model selection) for the memorised (M) sequence. We conducted four independent DCM
analyses, one for each tone (excluding the first tone, which was common across all experimental conditions).
Our hypothesised model (model one) had the highest evidence for tones two, three and four, while model three
reported the highest evidence for tone five. c – Posterior probability, protected exceedance probability and BOR
indicating the model with the highest evidence across the population (inference computed using random-effects
Bayesian model selection) for the novel (N) conditions. Here, we focused on the first tone which introduced the
variation in the sequence (i.e. tone 2 for NT1, tone 3 for NT2, tone 4 for NT3, tone 5 for NT4). Our hypothesised
model (model one) had the highest evidence for tones two, three and four, while model six reported the highest
evidence for tone five.

539
540

Time-frequency analysis for induced responses
We computed time-frequency analysis using complex Morlet wavelet transform (from 1 to 60
Hz with 1-Hz intervals). This analysis was conducted for induced responses. First, we
estimated independent time-frequency decompositions for each trial and each voxel of the six
ROIs described above (LHG, RHG, LHP, RHP, ACC, MC) plus left and right occipital
superior lobe for comparison purposes. Then, the computed power spectra were averaged
over voxels and over trials within each ROI. Finally, in line with the previous analyses, we
calculated four contrasts (M versus NT1, M versus NT2, M versus NT3, and M versus NT4).
Specifically, we computed a t-test for each frequency and time-point and corrected for
multiple comparisons using 2D cluster-based MCS (MCS, $\alpha = .05$, MCS p-value = .001). As
shown in **Figure 7**, results were similar across ROIs and displayed a generalised stronger
power for N versus M between 2 and 20 Hz (corresponding to, approximately, theta, alpha,
and beta bands), in the time window 1.0 – 3.0 seconds ($p < .001$). In addition, a few
significant clusters showed a moderate yet significant increase of power for M versus N ($p <$
.001) for frequencies higher than 30 Hz, which are normally referred to as gamma [42]. Detailed

statistical results about this procedure are extensively reported in **Table S11** and depicted in

Figure S12.

Finally, to strengthen the reliability of our results in MEG source space, we computed the

same analysis for eight fronto-temporal and occipital MEG channels. **Figure S13** and **Table**

S12 shows that the outcome of this procedure aligned with the results in MEG source space.

561
562
Figure 7. Induced responses during the recognition of memorised (M) and novel T1 (NT1) sequences

For each of the selected automated anatomical labelling (AAL) regions of interest (ROIs; left Heschl's gyrus
[LHG], right Heschl's gyrus [RHG], left hippocampus [LHP], right hippocampus [RHP], anterior cingulate
gyrus [ACC], medial cingulate gyrus [MC], plus left and right occipital superior lobe for comparison
purposes), three plots are provided. First, we illustrate the power spectrum computed using complex Morlet
wavelet transform for the M and NT1 conditions. Here, the colorbar indicates squared femtotesla (fT^2). Second,
we illustrate the significant results emerging from the contrasts between the power spectra of M versus NT1 (t-
tests and cluster-based Monte-Carlo simulations [MCS] correction for multiple comparisons). In this case, the
colorbar indicates the t-values obtained by contrasting M versus NT1.

Discussion

This study is rooted in the foundation of previous investigations and provides a deeper understanding of the neural mechanisms and brain's functional hierarchies underlying the long-term recognition of auditory sequences. Here, in contrast to prior studies, we deliberately reduced the speed of the musical stimuli [28-32]. This allowed us to reveal that the recognition of auditory sequences relied on distinct brain responses to each tone forming the sequence, as opposed to the slow response lasting for the entire duration of the sequence detected by previous studies [28-32]. This evidence suggests that the previously described slow signal might have been the summation of the neural responses to each sound of the sequence occurring at a faster tempo (i.e. each sound lasting 250 ms [28-32] as opposed to the 350 ms employed in the current study).

Furthermore, here we introduced systematic variations to the original memorised sequences to examine the brain mechanisms associated with conscious prediction error. The declarative nature of the prediction error, where participants were asked to identify differences in the sequences and provide a conscious response, extends our understanding beyond the existing body of research focused on automatic prediction error indexed by well-known ERP/Fs components such as N100, MMN, and ERAN [16, 17, 19-21]. Those studies revealed that such components were primarily generated by the auditory cortex [43, 44], reporting a complementary yet much reduced recruitment of the medial cingulate gyrus, inferior frontal gyrus, and frontal and hippocampal regions [14, 44]. Conversely, in our study we revealed an initial recruitment of the auditory cortex (100 – 150 ms) and then of the anterior and medial cingulate gyrus, ventromedial prefrontal cortex, and bilateral hippocampus (250 – 500 ms). Notably, our results showed that Heschl's gyrus discriminated memorised melodies versus systematic variations but did not distinguish the strength of the errors (i.e., the response was not significantly different across varied tones). Conversely, the prediction error observed in the bilateral hippocampus, anterior and medial cingulate gyrus was significantly stronger in response to the tone that introduced the variations compared to all other tones. Given the different brain responses observed in our study in comparison to previous literature on automatic prediction error, we argue that the brain signature underlying the awareness of the variation might be represented by the responses recorded in the anterior cingulate gyrus, ventromedial prefrontal cortex and hippocampus and their specific temporal dynamics.

Our results fit the well-known PCT [6, 7, 26], which states that the brain is constantly updating internal models to predict external information and stimuli. Recently, this theory has been

proposed to explain the brain mechanisms underlying complex cognitive processes [45], finding

a remarkable example in the neuroscience of music [7, 26]. However, while PCT holds promise

for understanding complex cognitive processes, the quantitative evidence in its favour has

been limited. Thus, our study filled this gap by providing quantitative evidence of the brain

functional hierarchies during long-term recognition of previously memorised and novel

sequences. The results showed feedforward connections from the auditory cortices,

representing sensory regions [46, 47], to the hippocampus, anterior cingulate gyrus, and medial

cingulate gyrus, brain areas which were previously associated with memory and predictive

processes [48, 49], and evaluation [50]. Simultaneously, we observed backward connections

operating in the opposite direction. This hierarchical alignment persisted consistently

throughout the entire sequence, except for the final tone. Here, the cingulate gyrus, which

holds a pivotal role in decision-making and evaluation [50, 51], assumed the top position within

the hierarchy. For the memorised sequences, the hierarchy consisted of the auditory cortex

sending feedforward signals to the hippocampus and anterior cingulate gyrus, and

subsequently to the medial cingulate gyrus (and receiving feedback signals in the opposite

direction). Similarly, for the varied sequences, the hierarchy showed feedforward connections

from the auditory cortex to the hippocampus and medial cingulate and then to the anterior

cingulate (and feedback in the opposite direction). This result suggests that, upon hearing the

last tone, the brain might prepare the evaluation of the sequence (i.e. categorising it as

'memorised' or 'novel') and exactly for this reason either the anterior or medial cingulate

gyri, both relevant regions for decision-making and evaluative processes [50, 51], occupy the

highest position in the hierarchy. In addition, our results present a significant dissociation

between anterior and medial cingulate gyri, suggesting a prominent role of the medial part of

the cingulate for recognition of memorised sequences and of the anterior part for novelty

detection.

These findings align with previous research demonstrating the flow of information from

sensory cortices to the medial temporal lobe, associative areas, and prefrontal regions of the

brain [23, 52]. Moreover, our results are coherent with the several studies which used PCT to

investigate perceptual [53] and automatic memory mechanisms in the brain [22], as reviewed by

Spratling [45]. These studies described how information processing, orientation tuning, contour

integration, and binocular rivalry in the visual system relied on the hierarchical transition of

the psychophysical and neurophysiological signals from the retina to the lateral geniculate

nucleus and V1 [54-57]. In the context of automatic sensory memory, PCT has also lent support

to the interpretation of the MMN as an error signal which aids the brain in adjusting its

internal predictive model, returning quantitative evidence of information flow from the

primary auditory cortex to the superior temporal gyrus and inferior frontal gyrus [22]. Our

findings expand on these studies by providing quantitative evidence of the hierarchical

organisation of the brain during a complex cognitive task, involving episodic memory

recognition for musical sequences. In addition, while the hierarchical architecture involved in

recognising both memorised and varied sequences remained largely unchanged, the temporal

dynamics, strength, and polarity of the brain signal sharply differed. Here, our results suggest

that when the upcoming sound of the sequence matched the predicted sound based on the

previously stored memory trace, first the auditory cortex and then hippocampus, anterior and

medial cingulate gyri responded with positive components. Conversely, when the upcoming

sound was incoherent with the prediction made by the brain, a pathway of primarily negative

and faster components emerged within the same brain network. It is worth noting that also the

inferior frontal gyrus, typically linked to short-term and working memory for music [58, 59],

displayed significantly different activity across experimental conditions. However, this region

exhibited overall reduced responses compared to the other brain regions within the network.

This observation further underscores the differentiation between long-term memory

recognition for auditory sequences and other memory subsystems.

An additional finding pertains to our utilisation of multivariate pattern analysis and

temporal generalisation techniques, as described by King and Dehaene [60]. Here, we revealed

that when decoding memorised from novel sequences, the algorithm could differentiate the

conditions starting from the first tone that was varied in the novel sequences. This learning

could then be generalised by the algorithm with significant (yet reduced) accuracy for all the

subsequent tones. This pattern was consistently observed across all categories of variations,

and it implies that the differential processes between the two conditions remained consistent

over time, indicating that the brain not only identifies the initial varied tone but also monitors

the entire auditory sequence. This monitoring might have served the purpose of assessing

whether the varied tone represented a transient mistake or the beginning of a proper varied

sequence.

Our findings revealed a positive overall correlation between musical training and neural

activity associated with the recognition of memorised and varied musical sequences. This

correlation was particularly evident with regards to the neural responses to the sounds that

introduced variations. This aligns with previous research indicating a connection between

musical expertise and altered brain characteristics, including enhanced neural responses (e.g.
enhanced auditory automatic prediction error) [14, 26, 61] and structural changes [62]. In this study,
what distinguishes our findings was the extension of this relationship to the domain of long-
term recognition of musical sequences and the underlying brain mechanisms.

On a behavioural level, both reaction times and accuracy remained consistent across
memorised and three categories of novel sequences: NT1, NT2, and NT3. However, accuracy
was significantly reduced, and reaction times increased for NT4. This shift might be
attributed to the altered final tone in the sequence, eliciting a slower behavioural prediction
error. Alternatively, the reduced accuracy and prolonged reaction times in NT4 could be
linked to chunking; the mental grouping process that occurs when sequences are presented
with a beat every four tones. In this view, after listening to four tones of the memorised
sequence (corresponding to a complete beat), the perception that the sequence belonged to the
group of previously learned sequences was very strong, and especially much stronger than
after only three tones. For this reason, we did not observe a linear increase in reaction times
and accuracy but only a difference between T4 and all the other conditions. Currently, we do
not have enough data to make definitive claims and future studies are needed to
systematically vary the length of the sequences.

Finally, the induced-response analysis showed that, after the end of the sequence, alpha
and beta bands were stronger for the varied compared to the memorised sequences.
Considering the modulation of alpha and beta activity in encoding and working memory tasks
described in the scientific literature [63], this finding may suggest that post-stimulus increase of
alpha-beta power might be relevant to process novel information. In addition, this analysis
revealed a weak yet significant increase in gamma power for the memorised compared to the
varied sequences. This is coherent with previous studies which reported increased gamma
power during recognition of target stimuli [64, 65] and, more generally, a modulation of the brain
oscillations associated with memory load and complex cognitive functions [66]. Although this
study provided initial insights on the time-frequency modulation occurring in the brain during
long-term recognition of auditory sequences, further research employing MEG and additional
tools such as stereo-electroencephalography (SEEG) is needed to expand our experimental
design and conduct cross-frequency coupling analysis, testing whether gamma-theta coupling
is connected to long-term recognition of auditory sequences. This would also provide
supporting evidence of the quality of the source reconstruction performed in this study since

SEEG data is less sensitive to source leakage artifacts than MEG source reconstruction

algorithms [67].

In conclusion, this study reveals the brain mechanisms and hierarchical dynamics of long-

term memory recognition of previously memorised and varied auditory sequences, effectively

integrating the PCT in the domain of complex cognitive tasks.

Materials and methods

Participants

The participant sample consisted of 83 volunteers [33 males and 50 females (sex, biological

attribute)] aged 19 to 63 years old (mean age: 28.76 ± 8.06 years). The sample was recruited

in Denmark and participants came from Western countries. All participants were healthy and

reported normal hearing. Their educational background was overall homogeneous.

Specifically, 73 participants had either a university degree (bachelor's or master's degree, n =

54) or were university students (n = 19). The remaining 10 were divided as follows: five had

a professional degree obtained after finishing high-school, while the remaining five had only

the high-school diploma. The project was approved by the Institutional Review Board (IRB)

of Aarhus University (case number: DNC-IRB-2020-006). The experimental procedures were

carried out in compliance with the Declaration of Helsinki – Ethical Principles for Medical

Research. All participants gave the informed consent before starting the experimental

procedure.

Experimental stimuli and design

In this study, we used an old/new paradigm auditory recognition task [28, 29, 31, 32] during

magnetoencephalography (MEG) recordings. First, participants listened to a short musical

piece twice and were asked to memorise it as much as possible. The musical piece consisted

of the first four bars of the right-hand part of Johann Sebastian Bach's Prelude No. 2 in C

Minor, BWV 847. In this piece, each bar included 16 tones. Thus, the total number of tones

was 16*4 = 64. Each tone lasted approximately 350 ms for a total of 22400 ms. In addition,

to provide a sense of musical closure, we included a final tone after the four bars which lasted

1000 ms. Thus, the total duration was 23400 ms (23.4 seconds). The stimulus is illustrated in

musical notation in **Figure S1**. A MIDI version of the piece was created using Finale

(MakeMusic, Boulder, CO). Second, participants were presented with 135 five-tone musical

excerpts that lasted 1750 ms each. Participants were requested to state whether each excerpt

belonged to the original music ('memorised' sequence [M], old) or were a varied musical

sequence ('novel' sequence [N], new) (**Figure 1a**). Twenty-seven excerpts were drawn from

the original musical piece and 108 were variations of the original melodies (**Figure S2** shows

all the sequences used in the study). The two categories of stimuli (M and N) were created as

follows. The M sequences were comprised by the first five tones of the first three measures of

the musical piece. These sequences were presented nine times each, for a total of 27 trials.
The N sequences were created through systematic variations of the three M sequences
(**Figure 1b**). This procedure consisted of changing every musical tone of the sequence after
the first (NT1), second (NT2), third (NT3) or fourth (NT4) tone. We created nine variations
for each of the original M sequences and each of the four categories of N. This resulted in 27
N sequences for each category, and 108 N in total. To be noted, the variations were created
according to the following rules:

- Inverted melodic contours (used twice): the melodic contour of the variation was
inverted with respect to the original M sequence (i.e., if the M sequence had the
following melodic contour: down-down-up-down, the N sequence would be: up-
up-down-up)

- Same tone scrambled (used three times): the remaining tones of the M sequence
were scrambled (e.g., M sequence: C-E-D-E-C, was converted into NT1 sequence:
C-C-E-E-D). When this was not possible (e.g., in the case of NT4, where only the
last tone is different from the M sequence), we substituted the last tone of the M
sequence with a random tone.

- Same tone (used three times): the same tone was repeatedly used, in some cases
varying only the octave (e.g., M sequence: C-E-D-E-C, was transformed into NT1
sequence: $C-E^{8-} E^{8-} E_8^- E_8$).

- Scrambling intervals (used once): the intervals between the tones were scrambled
(e.g., M sequence: $6^{th}m - 2^{nd}m - 2^{nd}m - 3^{rd}m$, was adapted to NT1 sequence:
$2^{nd}m, 6^{th}m, 3^{rd}m, 2^{nd}m$).

To be noted, the harmonic structure of the N sequences with regards to the original M
sequences was preserved in most of the cases. The only exceptions occurred for a few
instances of the 'inverted melodic contours' and 'scrambling intervals' strategies, which were
however important to control for other musical features such as melodic contour and
intervals. This strategy was implemented to avoid potential confounding variables stemming
from changes in harmony. However, investigating harmonic alterations concerning conscious
brain prediction errors could prove relevant for future studies centred on diverse types of
prediction errors.

This procedure allowed us to investigate (i) the brain dynamics underlying the recognition of
previously memorised auditory sequences and (ii) the conscious detection of the sequence
variation.

Data acquisition

The MEG recordings were acquired in a magnetically shielded room at Aarhus University
Hospital (AUH), Aarhus, Denmark, using an Elekta Neuromag TRIUX MEG scanner with
306 channels (Elekta Neuromag, Helsinki, Finland). The data was recorded at a sampling rate
of 1000 Hz with an analogue filtering of 0.1 – 330 Hz. Before the recordings, the head shape
of the participants and the position of four Head Position Indicator (HPI) coils were
registered with respect to three anatomical landmarks using a 3D digitizer (Polhemus Fastrak,
Colchester, VT, USA). This recording was later used to co-register the MEG data with the
MRI anatomical scans. For the entire duration of the MEG recordings, the HPI coils
registered the continuous localization of the head, which was subsequently employed for
movement correction. In addition, two sets of bipolar electrodes were used to record cardiac
rhythm and eye movements. This allowed us to remove the electrocardiography (ECG) and
electrooculography (EOG) artifacts in a later stage of the analysis pipeline.

The MRI scans were recorded on a CE-approved 3T Siemens MRI-scanner at AUH. The
recorded data consisted of structural T1 (mprage with fat saturation) with a spatial resolution
of 1.0 x 1.0 x 1.0 mm and the following sequence parameters: echo time (TE) = 2.61 ms,
repetition time (TR) = 2300 ms, reconstructed matrix size = 256 x 256, echo spacing = 7.6
ms, bandwidth = 290 Hz/Px.

The MEG and MRI recordings were acquired in two separate days.

Behavioural data

We obtained behavioural data (number of correctly recognised trials and correspondent
reaction times) from the experimental task carried out during the MEG recording.

Since the data was not normally distributed, we computed two independent Kruskal-Wallis H
tests (non-parametric one-way analysis of variance) to assess whether the five categories of
temporal sequences (M, NT1, NT2, NT3, NT4, NT5) differed in terms of correct responses
and reaction times. Multiple comparisons were corrected using the Tukey-Kramer correction
[68].

MEG data pre-processing

The raw MEG sensor data (204 planar gradiometers and 102 magnetometers) was first pre-processed by MaxFilter [69] to attenuate external interferences. We applied signal space separation (MaxFilter parameters: spatiotemporal signal space separation [SSS], down-sample from 1000Hz to 250Hz, movement compensation using cHPI coils [default step size: 10 ms], correlation limit between inner and outer subspaces used to reject overlapping intersecting inner/outer signals during spatiotemporal SSS: 0.98). We opted to downsample the data by a factor of four because MEG is not particularly sensitive to high-gamma frequency (e.g. 100 Hz) [25, 70], while it provides much better outcomes for lower frequency bands (e.g. 0.1 – 60 Hz, as done in the current study) [25, 70]. Moreover, our research focused on the event-related broadband brain data during memory recognition, which are often expressed in slower bands [27, 28]. For these reasons, as commonly done in the literature [25, 70], we downsampled the data, a procedure which significantly accelerated the computations by reducing data volume without substantial data loss [25, 70].

The data was then converted into Statistical Parametric Mapping (SPM) format and further pre-processed and analysed in MATLAB (MathWorks, Natick, MA, USA) using a combination of in-house-built codes (LBPD, https://github.com/leonardob92/LBPD-1.0.git) and the Oxford Centre for Human Brain Activity (OHBA) Software Library (OSL) [71] (https://ohba-analysis.github.io/osl-docs/), a freely available software that builds upon Fieldtrip [72], FSL [73], and SPM [74] toolboxes.

The continuous MEG data was visually inspected to identify and remove large artifacts using the OSLview tool. The data that was removed was less than 0.1% of the amount of collected data. Independent component analyses (ICA) were used (OSL implementation) to discard the interference of eyeblinks and heart-beat artefacts from the brain data [75]. First, we decomposed the original signal into independent components. Second, we correlated all the components with the activity recorded by the EOG and ECG channels. This procedure identified the ICA component X that strongly correlated with either the EOG or the ECG (strongly correlated means that the correlation coefficient for component X was at least three times higher than the correlation coefficients for the remaining components). To further validate the accuracy of this procedure, we also visually inspected the highly correlated components. We assessed whether their topographic distribution across MEG channels matched the typical distribution associated with eyeblink and heartbeat activity. When both the correlations and visual inspections converged, indicating that the ICA component strongly

reflected eyeblink or heartbeat activity, those components were discarded. Third, the signal

was rebuilt using the remaining components. Finally, the signal was epoched in 135 trials (27

M, 27 NT1, 27 NT2, 27 NT3, 27 NT4) and baseline-corrected by removing the mean signal

recorded in the baseline from the post-stimulus brain signal. Each trial lasted 4500 ms (4400

ms plus 100 ms of baseline time).

Multivariate pattern analysis (decoding) – MEG sensors

We performed multivariate pattern analyses to decode different neural activity associated

with the recognition of M versus N (**Figure 1c**). Here, we computed four independent

analyses, decoding M from each of the four categories of the novel sequences (i.e., M versus

NT1, M versus NT2, M versus NT3, M versus NT4). This was our first analysis since

decoding relies on all the available data and makes no assumptions about which MEG

channels or time-points should be selected. We used support vector machines (SVMs) [76] and

calculated independent analyses for each participant. The MEG data was rearranged in a 3D

matrix (channels x time-points x trials) and submitted to the SVM algorithm. To avoid

overfitting, a leave-one-out cross-validation approach was adopted to train the SVM classifier

to decode the two experimental conditions. This procedure divided the trials into N different

groups (here N = 8). Then, for each time-point, it assigned N – 1 groups to the training set

and the remaining N_{th} group to the testing set. After that, the classifier ability to separate the

two conditions was evaluated. This process was performed 100 times with random

reassignment of the data to training and testing sets. To summarise, the decoding accuracy

time series were averaged to obtain a final time series showing the performance of the

classifier for each participant. In addition to the decoding accuracy time series (**Figure 2a**),

this analysis returned weights which can be used to learn which MEG channels provided the

strongest contribution to the decoding analysis [77] (**Figure 2b**).

To test the significance of the decoding results (chance level set at 50%), we employed a

sign permutation test against the chance level for each time-point and then corrected for

multiple comparisons using false-discovery rate (FDR) correction (α = .05; FDR-adjusted q <

.012).

To assess whether each pair of conditions were differentiated by neural patterns which

were stable over time, we computed four temporal generalisation multivariate analyses

(**Figure 2c**). The algorithm was the same as the one previously described. However, in this

case we utilised each time-point of the training set to predict not only the same time-point in

the testing set, but all time-points [78]. Here, the significance was tested using a signed

permutation test against the chance level (50%) for each time-point, as the previous analyses.
Then, we corrected for multiple comparisons using two-dimensional (2D) cluster-based
Monte-Carlo simulations (MCS, α = .05, MCS p-value = .001). First, we computed the
clusters size of the continuous, binarized, significant values over time. The MCS α = .05
means that to binarize the values outputted by the signed permutation test against the chance
level that we described above, we used a *p-value* threshold of .05 (i.e. values below .05 were
considered significant and inputted to the MCS as '1s' while values above .05 as '0s'). Thus,
the first step of the MCS algorithm consisted of detecting the clusters of '1s'. Second, we
made 1000 permutations of these binarized values. For each permutation, we computed the
size of the maximum emerging cluster and built a reference distribution using those values.
Finally, we considered significant the original clusters that were bigger than the 99.9% of the
permuted data maximum cluster sizes (in other words, the MCS *p-value* = .001 refers to the
percentage of the cluster sizes in the reference distribution that the original clusters should
exceed to be considered significant). This procedure was observed for all the analogous MCS
described in the study. Moreover, additional details on this MCS algorithm can be found in [28, 31, 32, 79].
891

Univariate analysis – MEG sensors
Pairwise decoding is a powerful technique, but it can only tell if two conditions are
characterised by significantly different brain activity. It does not provide information about
which condition presented the strongest brain activity. To answer this question, we computed
an additional analysis using the brain activity recorded by the MEG channels which mainly
contributed to the decoding algorithm. This was done following two steps: (i) averaging the
weights of the MEG channels in all the significant time-windows emerged from the decoding
algorithm (i.e. FDR-corrected decoding accuracy time series for M versus NT1, M versus
NT2, M versus NT3, M versus NT4); (ii) selecting the MEG channels whose weight was
higher than the average (over the MEG channels) plus one standard deviation. This was done
independently for magnetometers and gradiometers and returned the following MEG
channels; magnetometers: 1321, 0211, 1441, 0131, 1411, 1221, 1331, 0121, 1311, 0341,
2611, 0241, 1511, 1341, 0111, 0221; gradiometers: 1442, 0242, 0222, 1332, 2612, 0232,
1513, 1312, 0312, 1222, 1342, 0342, 0123, 1423, 2422, 2613, 0213, 1212, 1323, 0413, 0132,
0142, 0113, 1233, 2033, 1413, 1123, 0143, 0323, 1343, 1612, 1412.

These MEG channels were then averaged into four distinct groups, based on the channel
type and the polarity of the signal: (i) magnetometers negative N100 (**Figure 2d**, left), (ii)
magnetometers positive N100 (**Figure 2d**, right), (iii) gradiometers negative N100 (**Figure
2e**, left), and (iv) gradiometers positive N100 (**Figure 2e**, right). The reasons why these four
groups were considered are as follows: (i) magnetometers and gradiometers are different
types of sensors and have a difference scale so they are usually analysed independently [25, 70];
(ii) because of the relationship between electrical activity and the correspondent magnetic
field, when the brain generates a truly negative response (e.g. N100 to the first sound of the
sequence), the MEG channels record a magnetic field characterised by both negative and
positive values (i.e. the same electric activity generates both a positive and negative polarity
of the magnetic field that are picked up independently by different MEG channels). The
result of this well-known effect is that some channels show the 'true' polarity of the brain
signal (e.g. the N100 to the first sound is negative) and other show the inverted polarity (e.g.
the N100 to the first sound is positive). When averaging MEG channels together, this must be
considered to prevent averaging out the signal. Thus, as depicted in **Figure 2**, we averaged
the MEG channels into the above-described four groups. As follows, we report the selected
MEG channels divided into the four groups. (i) Magnetometers negative N100: 0131, 0211,
0121, 0341, 0241, 1511, 0111; (ii) magnetometers positive N100: 1321, 1441, 1411, 1221,
1331, 1311, 2611, 1341; (iii) gradiometers negative N100: 0213, 0232, 0312, 0342, 0132,
0222, 1423, 1212, 1123, 0413, 1413, 1233, 0323, 2613; (iv) gradiometers positive N100:
1442, 0242, 1332, 2612, 1513, 1312, 1222, 1342, 0123, 2422, 1323, 0142, 0113, 2033, 0143,
1343, 1612, 1412.
Then, independently for each group, we computed one t-test for each time-point and each
combination of M versus Ns (i.e., M versus NT1, M versus NT2, M versus NT3, M versus
NT4). We corrected for multiple comparisons using a one-dimensional (1D) cluster-based
MCS (MCS, α = .05, MCS *p-value* = .001) (**Table S3**). First, we identified the clusters of
significant continuous values in time. Second, we computed 1000 permutations, randomising
the significant values obtained from the t-tests. For each permutation, we extracted the
maximum cluster size and built their reference distribution. Finally, we considered significant
the original clusters that were larger than 99.9% of the permuted ones.
To provide full disclosure of our data, we have complemented these computations with a
further analysis. In this case, we computed the same procedure described above, contrasting
M versus each category of N (i.e. M versus NT1, M versus NT2, M versus NT3, M versus

NT4) and correcting for multiple comparisons with cluster-based MCS (MCS, α = .05, MCS

p-value = .001). However, here we computed the contrasts independently for each MEG

channel. The results are depicted in **Figure S5$_{a-z2}$** and reported in detail in **Table S4**.

Source reconstruction

MEG is a powerful tool to detect the whole-brain activity with excellent temporal resolution.

However, to obtain a complete picture of the whole-brain activity underlying complex

cognitive tasks the spatial component of the brain activity must be also identified. Here, we

employed the established beamforming method [80-82] (**Figure 1e**), built upon a combination of

in-house-built codes and codes available in OSL, SPM, and FieldTrip.

To reconstruct the brain sources that generated the MEG signal, an inverse problem must

be solved. The MEG recording shows the activity of the neural signals outside the head but

cannot provide information on the specific brain sources which generated it. Thus, we used

beamforming algorithms to solve this problem, implementing the two following steps: (i)

designing a forward model and (ii) computing the inverse solution.

The forward model is a theoretical model which considers each brain source as an active

dipole (brain voxel). It describes how the unitary strength of each dipole would be reflected

over all MEG sensors. Here, we employed only the magnetometer channels and an 8-mm

grid, which returned 3559 dipole locations (voxels) within the whole brain. This was done

because the magnetometers are the MEG sensors that better capture the activity coming from

deep sources of the brain, while the gradiometers are more sensitive to the activity originated

in the portion of the cortex nearest to each gradiometer [25, 70]. Since our hypothesis was that

regions in the medial temporal lobe (e.g. hippocampus) played a role in long-term auditory

recognition, we opted for the magnetometers to compute the source reconstruction. After co-

registering the individual structural T1 data with the fiducial points (i.e., information about

head landmarks), we computed the forward model by adopting the widely used method called

"Single Shell", which is presented in detail in Nolte [83]. The output of this computation,

referred to as "leadfield model", was stored in the matrix L (sources x MEG channels). In the

three cases where the structural T1 was not available we performed the leadfield computation

using a template (MNI152-T1 with 8-mm spatial resolution).

Then, we computed the inverse solution. As mentioned above, we chose the beamforming,

which is one of the most popular and effective algorithms available in the field. This

procedure employs a different set of weights which are sequentially applied to the source

locations for isolating the contribution of each source to the activity recorded by the MEG

channels. This is done for each time-point of the recorded brain data. The beamforming

inverse solution can be summarized by the following main steps.

The data recorded by MEG sensors (B) at time t, can be described by the following

equation (1):

$$B_{(t)} = L * Q_{(n_i,t)} + \qquad \qquad (1)$$

981   where L is the leadfield model, Q is the dipole matrix carrying the activity of each active

982   dipole (q) over time and $\square$ is noise (see Huang and colleagues [82] for details). To solve the

983   inverse problem, Q must be computed. In the beamforming algorithm, weights are computed

984   and then applied to the MEG sensors at each time-point, as shown for the single dipole q in

985   equation (2):

$$q_{(t)} = W^T * B_{(t)} \qquad \qquad (2)$$

988   To obtain q, the weights W should be computed (the subscript T refers to transpose

989   matrix). To this goal, the beamforming relies on the matrix multiplication between L and the

990   covariance matrix between MEG sensors I, which is calculated on the concatenated

991   experimental trials. Specifically, for each brain source n, the weights W_n are computed as

992   shown in equation (3):

$$W_{(n)} = (L_{(n)}{}^T * C^{-1} * L_{(n)})^{-1} * L_{(n)}{}^T * C^{-1} \qquad \qquad (3)$$

995   To be noted, the computation of the leadfield model was performed for the three main

996   orientations of each brain source (dipole), according to Nolte [83]. Before computing the

997   weights, the orientations were reduced to one using the singular value decomposition

998   algorithm on the matrix multiplication reported in equation (4). This procedure is widely

999   adopted to simplify the beamforming output [67,68].

1000

$$L = svd(l^T * C^{-1} * l)^{-1} \qquad \qquad (4)$$

1001

Here, l represents the leadfield model with the three orientations, while L is the resolved one-orientation model that was utilized in (3). Finally, the weights were applied to each brain source and time-point. To be noted, the covariance matrix C was computed on the continuous signal, which was estimated by concatenating the trials of all experimental conditions. The weights were applied to the brain data associated with each condition and normalised according to Luckhoo and colleagues [80, 84-86]. For counterbalancing the reconstruction bias towards the centre of the head. The weights were applied to the neural activity averaged over trials for the evoked responses and to the neural activity of each independent trial for the induced responses. This procedure returned a time series for each of the 3559 brain sources (and each trial in the case of induced responses), commonly referred to as neural activity index [80, 84]. The sign ambiguity of the evoked responses time series was adjusted for each brain source using its sign in correspondence with the N100 response to the first tone of the auditory sequences [28, 31, 32].

Neural sources of the peak activity (MEG magnetometers) and musical training

After showing evidence of the differential brain activity in relation to the experimental conditions, we wanted to characterise the neural sources of the main peaks of brain activity recorded by the magnetometers in response to each tone of the musical sequences, as described in the section on the univariate analysis of MEG channels.

To detect the peaks of the brain activity recorded by the magnetometers for each tone (except the 1st one) of the musical sequence, we extracted the time indices of the minimum and maximum value recorded by the magnetometers shown in **Figures 2d** and **3** in a 400 ms time-window following the onset of each tone. The duration of this time window reflects the period during which independent responses to each sound occurred. This procedure returned the time index of two neural peaks for each experimental condition and each tone of the sequence (with exclusion of the 1st one which did not differ across conditions). This revealed that one peak of the neural activity occurred approximately at 350 ms after the onset of each tone (highlighted in yellow in **Figure 3a**, top row and **Figure 3b**, bottom row). With regards to this peak, condition M showed the strongest response among all conditions. The other peak occurred in response to the tone which introduced the variation in the musical sequence (highlighted in purple in **Figure 3a**, bottom row and **Figure 3b**, top row). In this case, the N condition which introduced the variation reported the strongest response among all other conditions. This occurred approximately at 250 ms after the varied tone (i.e. NT1 showed the

strongest response to tone two among all conditions, NT2 to tone three, NT3 to tone four and NT4 to tone five). Extensive details on these neural activity peaks are reported in **Table 2**.

Then, for each of the defined peaks, we computed contrasts in MEG source space between M and N. More specifically, with regards to the peaks occurring approximately at 350 ms after each tone, we contrasted M versus NT1 (illustrated in the yellow boxes in **Figure 3**). This was done because the peaks occurring at 350 ms after each tone were always greater for M, and NT1 was the only condition comprising sounds that were always different from M. Similarly, with regards to the peaks occurring approximately at 250 ms after the onset of the varied tones, we contrasted the N condition which showed the strongest peak against M (illustrated in the purple boxes in **Figure 3**). Here, for each condition in the pair, we averaged the absolute value of the activity of each of the 3559 reconstructed brain voxels in the ±20 ms time-window around the correspondent time index. Then, we computed t-tests for each of the 3559 brain voxels by contrasting the pairs of conditions and FDR-corrected for multiple comparisons. The results of these analyses are shown in **Figure 3** and reported in detail in **Table 2** and **Table S5**. In addition, we reported the neural activity recorded by the MEG channels in topographic maps in **Figure S6**.

Furthermore, the brain activity recorded by the MEG magnetometers for each neural peak and hemisphere was correlated with the measure of participants' musical training provided by the Goldsmith Musical Sophistication Index (GOLD-MSI) [37] and corrected for multiple comparisons using FDR. The GOLD-MSI is a widely used standardised test which measures the ability of engaging with music providing five subscales: (i) Active engagement with music, (ii) Perceptual abilities, (iii) Musical training, (iv) Emotions), and (v) Singing abilities. Here, we focused on the *Musical training* subscale which encompasses data related to music education, formal training, average time spent playing musical instruments at different life stages, etc [37]. The Results are reported in **Table 2**.

Statistical analysis – automated anatomical labelling (AAL) time series

After evaluating the neural sources of the different brain activity associated with the experimental conditions using the fine-grained 8-mm parcellation of the brain (**Figure 1e**, *"8-mm parcellation"*), we focused on a set of anatomically defined regions of interest (ROIs) (**Figure 1e**, *"AAL parcellation"*). Here, we used a standard anatomical parcellation method known as automated anatomical labelling (AAL) [38] and calculated a time series for each of the 90 non-cerebellar ROIs of AAL (**Figure 1f**, *"Evoked responses"*).

As specified above, our source reconstruction was conducted in an 8-mm space, resulting in

3559 brain voxels, each with its own time series. Thus, for each of the 90 ROIs, we identified

the corresponding brain voxels and then averaged the time series of these associated voxels.

This allowed us to obtain a final time series for each AAL ROI.

Then, we computed one t-test for each AAL ROI, each time-point and each combination

of M versus Ns. Finally, we corrected for multiple comparisons using one-dimensional (1D)

cluster-based MCS (MCS, α = .05, MCS *p-value* = .001). **Figure S7$_{a-g}$** and **Table S6** show

all AAL ROIs and the significant differences between conditions. **Figures 4** and **S8** shows

instead a selected array of ROIs which were particularly relevant for this study. Here, we

identified the two ROIs that showed the strongest activity in absolute terms among auditory

regions (i), medial temporal lobe (ii) and cingulate and prefrontal cortices (iii). We selected

these regions based on the cognitive processes involved in the experimental task used in this

study: audition (i), memory (ii), evaluation and decision-making (iii). As shown by **Figures 4**

and **S8**, these ROIs were left and right Heschl's gyrus (LHG, RHG), left and right

hippocampus (LHP, RHP), anterior cingulate gyrus (ACC) and medial cingulate gyrus (MC),

respectively. These six regions were also used for the Dynamic Causal Modelling (DCM)

analysis described later.

In addition, since it has been shown that the source reconstruction can produce leakage of

the brain signals between the ROIs [39], we took additional measures to ensure the robustness

of our results. Specifically, we applied the multivariate source leakage correction method

proposed by Colclough and colleagues [39] to the ROIs time series. This method consists of

removing the zero-lag correlations between the ROIs time series through a symmetric,

multivariate orthogonalization process. After computing this correction, we performed the

exact same statistical tests described above. The outcomes of this analysis are illustrated in

Figure S9 and described in **Table S8**.

Finally, as a further measure to enhance the reliability of our findings, in supplementary

materials we reported the description of an alternative, functional parcellation derived from

the data (**Figures S7$_{a-g}$, S10, S11, S15, S16, S17, S18** and **Tables S9, S14, S15**).

1096

Prediction error – differentiating Heschl's gyrus from hippocampus and cingulate

gyrus

After differentiating the brain activity of M versus N, we conducted an additional analysis to

investigate the responses elicited by the varied tones within the N conditions. Regarding

Heschl's gyrus (left and right), we focused on the late N100 peaking approximately at 150 ms
after the onset of each varied tone. Regarding the hippocampus (left and right), anterior and
medial cingulate gyrus, we focused on the responses peaking approximately at 250 ms after
the onset of each varied tone (these responses were negative for hippocampus and anterior
cingulate gyrus and positive for the medial cingulate gyrus). Our goal was to compare the
responses to each individual tone independently within each ROI, for N conditions containing
at least two varied tones. To achieve this, we computed the average neural activity within a
±20 ms time window centred around the designated time index (150 ms for Heschl's gyri and
250 ms for the hippocampus and cingulate gyrus), as depicted in **Figure 5** using red circles.
Subsequently, we performed separate one-way analyses of variance (ANOVAs) for each ROI
and N condition containing at least two varied tones to assess whether the responses to
subsequent varied tones differed, independently for each ROI. Here, we used the sequential
neural peaks as independent variable and the neural activity as the dependent variable. Since
this approach led to a total of 18 ANOVAs, we applied an FDR correction to account for
multiple comparisons. Moreover, the post-hoc analyses computed for each of the nine
ANOVAs were corrected for multiple comparisons using Tukey-Kramer correction [68].
1117

Dynamic causal modelling (DCM)
DCM aims to infer the causal structure of interconnected or dispersed dynamical systems
such as a network of brain regions (**Figure 1g**). It employs a Bayesian model comparison
approach that involves evaluating competing models which explain the generation of time
series data associated with alternative network architectures. DCM has been widely used in
the neuroscientific literature [40, 41, 87-90], especially but not limited to the framework of
predictive coding, and it is particularly useful for estimating hierarchical relationships
between brain regions in relation to experimental stimuli. In brief, a dynamic causal model
for MEG responses starts with a reasonably plausible neuronal model which describes how
neuronal populations generate the activity of the large brain regions that are source
reconstructed using MEG. Here, we employed the widely used neural-mass model developed
by David and colleagues [88]. This model characterises large brain regions by averaging the
post-synaptic membrane depolarization of different neuronal populations, which are
interconnected in a vertical pattern, mimicking the canonical cortical microcircuit (CMC)
structure outlined by Bastos and colleagues [91]. Then, the neural-mass model is integrated with
a network architecture indicating the hypothesised feedforward and feedback connections

between the brain regions that were identified from the MEG data. Finally, the DCM is fitted
to the time series of those brain regions and the marginal likelihood of the model (model
evidence), as well as the posterior and protected exceedance probabilities, are estimated.
Typically, a first model representing the hypothesised network architecture/brain hierarchy is
formulated, and its evidence is compared to the evidence of a few alternative, competing
models. This is done to establish which is the network architecture which shows the highest
model evidence given the experimental data.

DCM is a popular and established method and extensive details on its functioning can be
found in multiple experimental, theoretical and review papers [40, 41, 87-90]. Additional details on
the Bayesian model comparison are available in the next section.

In this study, DCM was used to test our hypothesised model of brain hierarchies during
recognition of memorised and varied musical sequences against five competing models
(**Figure 6a**).

These six models comprised the six AAL ROIs showed in **Figure S8** and illustrated in the
previous sections of the paper (LHG, RHG, LHP, RHP, ACC, MC). The models are
described as follows:

1. **Model 1**: this is the model that we hypothesised would be true and consisted of feedforward connections from LHG, RHG to LHP, RHP, ACC, MC, and feedback connections from LHP, RHP, ACC, MC to LHG, RHG.

2. **Model 2**: this alternative model consisted of feedforward connections from LHG, RHG to LHP, RHP, ACC, MC. No feedback connections were indicated. This model served to evaluate if there was evidence for a loop of feedforward and feedback connectivity between the auditory cortices and the other brain regions or if only feedforward connectivity was the most likely explanation of the data.

3. **Model 3**: this alternative model consisted of feedforward connections from LHG, RHG to LHP, RHP and from LHP, RHP to ACC, MC. The model also hypothesised feedback connections from ACC, MC to LHP, RHP and from LHP, RHP to LHG, RHG. This alternative model distinguished HP from ACC and MC. This model was formulated based on the anatomical contiguity of ACC and MC [92], as opposed to HP and wished to establish if there was an additional hierarchical layer between LHP, RHP and ACC, MC.

4. **Model 4**: this alternative model consisted of feedforward connections from LHG, RHG to ACC, MC and from ACC, MC to LHP, RHP. The model also hypothesised

feedback connections from LHP, RHP to ACC, MC and from ACC, MC to LHG, RHG. This model was formulated with the same rationale behind model 3, but it tested the opposite direction of the connections between LHP, RHP and ACC, MC.

5. **Model 5**: this alternative model consisted of feedforward connections from LHG, RHG to MC and from MC to ACC, LHP, RHP. The model also hypothesised feedback connections from ACC, LHP, RHP to MC and from MC to LHG, RHG. This alternative model distinguised HP and ACC from MC. This model was formulated based on the higher correlations between the time series of ACC, LHP, RHP ($r_{ACC-LHP}$ = .542; $r_{ACC-RHP}$ = .731; $r_{LHP-RHP}$ = .486) compared to the correlations between MC and ACC (r = -.565), LHP (r = .256), and RHP (-.163). It aimed to establish if there was an additional hierarchical layer between MC and LHP, RHP, ACC.

6. **Model 6**: this alternative model consisted of feedforward connections from LHG, RHG to ACC, LHP, RHP and from ACC, LHP, RHP to MC. The model also hypothesised feedback connections from MC to ACC, LHP, RHP and from ACC, LHP, RHP to LHG, RHG. This model was formulated with the same rationale behind model 5, but it tested the opposite direction of the connections between LHP, RHP, ACC and MC.

Based on the extensive evidence that the auditory cortices are the first brain regions to respond to sounds in the context of ERP/Fs [46, 47], all the models hypothesised that the external auditory inputs reached LHG and RHG.

We aimed to test the network architecture of this array of brain regions for both recognition of previously memorised and novel musical sequences [i.e. for conditions M and N(s)]. Since DCM for MEEG is usually computed on relatively short time-windows (e.g. 350 – 400 ms) and for one experimental perturbation at a time (e.g. one sound) [40, 41, 87-90], we computed a series of DCM analyses independently for each tone of the sequences, in a 350 ms time-window from each tone onset. Specifically, we computed an independent DCM for tones two, three, four and five for condition M. Tone one was excluded because it was the same across all conditions.

In relation to the N conditions, we were interested only in the tone that introduced the variation in the sequences. Thus, we computed four DCMs, organised as follows: NT1 for tone two; NT2 for tone three; NT3 for tone four; NT4 for tone five. In conclusion, we

computed a total of eight DCM analyses, always testing the same alternative models since we
hypothesised the same hierarchy in the network architecture of M and Ns.

As a final note, following standard DCM procedures, the DCMs were estimated
independently for each participant. The consistency of the results across participants was
tested using random-effects Bayesian model selection (RFX BMS), which is described in
detail in the following section.

1206

Bayesian model comparison across participants

Dynamic causal models are estimated with variational Laplace [93]. The variational Laplace is
used for deriving both the posterior distribution and the marginal likelihood of the model
itself, which is also known as the model evidence. Employing a Laplace approximation to the
multivariate posterior distribution of model parameters, denoted as $q(\theta|y, m)$, we iteratively
compute conditional means and covariances by maximizing a lower bound on the logarithm
of the model evidence, referred to as the log-evidence.

This optimisation utilises the Newton's method and a Fisher scoring scheme to maximize
the (negative) variational free-energy F of the model (5):

1216

$$F = E_q[ln\ p(y|\theta, m)] - D_{KL}[q(\theta|y, m)||p(\theta|m)] \qquad (5)$$

1217

Where $E_q[\cdot]$ indicates the expectation under the variational posterior density $q(\theta|y, m)$,
while $D_{KL}[\cdot]$ refers to the relative entropy [or Kullback-Leibner (KL) divergence]. This
makes the free energy a lower-bound approximation to the log-evidence.

To gain a deeper understanding of the utility of free energy in model comparison, we can
break it down into two components. The initial component comprises the anticipated log-
likelihood of the data, denoting the model's precision. The subsequent component is the
relative entropy, or KL divergence, between the multivariate posterior and prior probability
distributions, signifying the model's complexity. This formulation of complexity as a KL
divergence is based on the premise that the posterior need not deviate significantly from the
prior to accommodate new data. In essence, a proficient model should yield posterior beliefs
consistent with its prior beliefs when exposed to new data. Consequently, free energy can be
dissected into accuracy minus complexity, jointly encapsulating a model's ability to explain
the data. Our set of alternative models were then compared using their free energy, which
indexes each model evidence [89]. To study the consistency of the free energy across the

participants, we used a RFX BMS model selection procedure, which provides the posterior

probability and protected exceedance probability of each model for model comparison within

the entire sample of participants [90].

The posterior probability is determined using normalised Bayesian factors (BF) and

reflects the likelihood of each model being independently a good fit based on the observed

data. The protected exceedance probability describes the likelihood that a particular model is

the best of explaining the data when compared to all the alternative models. In addition, by

accounting for variations in model frequencies that could arise purely by chance [94], the

protected exceedance probability keeps into account potential issues associated with multiple

comparisons, enhancing the robustness of the results. Finally, in addition to posterior

probability and protected exceedance probability, we also calculated a statistical measure

called the Bayesian omnibus risk (BOR). The BOR is a metric that quantifies the statistical

risk associated with performing Bayesian model comparisons at a group level, taking into

account potential variations in the likelihood of the models among different participants. In

other words, the BOR indicates the likelihood that the protected exceedance probability

observed in each model comparison is due to random chance within the participant sample.

This parameter is a recent addition to the DCM framework and serves as a further tool to

assess the reliability of the DCM results [40]. Taken together, the posterior probability,

protected exceedance probability and BOR provide a state-of-the-art, conservative approach

for determining which of the alternative DCM models has the strongest evidence within the

entire population. Additionally, they offer a robust measure of the probability that this

evidence is not simply a result of random chance [40, 87, 90, 94].

This procedure was observed independently for the eight DCM analyses described in the

previous section.

1256

Time-frequency analysis for induced responses

We computed a time-frequency analysis using complex Morlet wavelet transform (from 1 to

60 Hz with 1-Hz intervals) [95]. This analysis was conducted for induced responses,

independently for the six ROIs previously described and for the four contrasts considered in

this study (i.e., M versus NT1, M versus NT2, M versus NT3, M versus NT4). Specifically,

the time-frequency decomposition was done independently for each trial, brain voxel, and

participant. Then, as shown in **Figures 7** and **S12**, the power spectrum of each trial and each

brain voxel was averaged within each of the six ROIs. In addition, to provide results which

can more easily compared with previous literature, we have also computed the power

spectrum for left and right occipital superior lobe. Finally, we computed a t-test for each

frequency and time-point, making four contrasts: M versus NT1, M versus NT2, M versus

NT3, M versus NT4. The emerging *p*-values were binarized ($\alpha = .05$) and then submitted to a

2D MCS (MCS *p*-value = .001). Here, we calculated the clusters size of continuous

significant values in time and frequency. Then, we made 1000 permutations of the binarized

p-values. For each permutation, we measured the size of the maximum emerging cluster and

built a reference distribution with one value for each permutation. Finally, the original

clusters were considered significant when they were bigger than the 99.9% of the permuted

data maximum cluster sizes.

To strengthen the reliability of our results in MEG source space, we computed the same

analysis for eight fronto-temporal and occipital MEG channels. In this case, the power

spectra were computed independently for each trial and then averaged (**Figure S13** and

Table S12)**.**

1279

1280

1281

1282 *Data availability*

1283 The neuroimaging data related to the experiment is available in Zenodo at the following link:

1284 https://doi.org/10.5281/zenodo.8046524.

1285

1286   *Code availability*

1287   The codes are available at the following links: https://github.com/leonardob92/LBPD-1.0.git

1288   and https://github.com/leonardob92/HierarchicalPredictiveCoding_Music_MEG.git.

1289

1290

Acknowledgements

The Center for Music in the Brain (MIB) is funded by the Danish National Research Foundation (project number DNRF117). Additionally, we thank the Fundación Mutua Madrileña for the economic support provided to author GFR.

LB is supported by Carlsberg Foundation (CF20-0239), Lundbeck Foundation (Talent Prize 2022), Center for Music in the Brain, Linacre College of the University of Oxford (Lucy Halsall fund), Society for Education and Music Psychology (SEMPRE's 50th Anniversary Awards Scheme), and Nordic Mensa Fund.

MLK is supported by Center for Music in the Brain and Centre for Eudaimonia and Human Flourishing, which is funded by the Pettit and Carlsberg Foundations.

Author contributions

LB, GFR and MLK conceived the hypotheses. LB designed the study. LB, MLK and PV

recruited the resources for the experiment. LB, GFR and FC collected the data. LB and GFR

performed pre-processing, statistical analysis and modelling. DP provided MATLAB codes

for decoding analysis. MJD provided initial MATLAB codes and assistance for the DCM

analysis. DP, MJD, MLK, GFR and PV provided essential help to interpret and frame the

results within the neuroscientific literature. GFR and LB wrote the first draft of the

manuscript. LB, GFR, FC and MLK prepared the figures. All the authors contributed to and

approved the final version of the manuscript.

1311

1312 *Competing interests' statement*

1313 The authors declare no competing interests.

1314

References

1. Deco, G., Vidaurre, D. & Kringelbach, M.L. Revisiting the global workspace orchestrating

the hierarchical organization of the human brain. *Nat Hum Behav* **5**, 497-511 (2021).

2. Mesulam, M.M. From sensation to cognition. *Brain* **121 (Pt 6)**, 1013-1052 (1998).

3. Preti, M.G., Bolton, T.A. & Van De Ville, D. The dynamic functional connectome: State-

of-the-art and perspectives. *Neuroimage* **160**, 41-54 (2017).

4. Scalabrini, A., Mucci, C. & Northoff, G. The nested hierarchy of self and its trauma: In

search for a synchronic dynamic and topographical re-organization. *Front Hum Neurosci* **16**,

980353 (2022).

5. Kringelbach, M.L*., et al.* Dynamic coupling of whole-brain neuronal and neurotransmitter

systems. *Proc Natl Acad Sci U S A* **117**, 9566-9576 (2020).

6. Friston, K. Predictive coding, precision and synchrony. *Cognitive neuroscience* **3**, 238-239

(2012).

7. Koelsch, S., Vuust, P. & Friston, K. Predictive processes and the peculiar case of music.

Trends in cognitive sciences **23**, 63-77 (2019).

8. Ip, I.B. & Bridge, H. Investigating the neurochemistry of the human visual system using

magnetic resonance spectroscopy. *Brain Struct Funct* **227**, 1491-1505 (2022).

9. King, J.-R. & Wyart, V. The human brain encodes a chronicle of visual events at each

instant of time through the multiplexing of traveling waves. *Journal of Neuroscience* **41**,

7224-7233 (2021).

10.   Zatorre, R.J. Sound analysis in auditory cortex. *Trends Neurosci* **26**, 229-230 (2003).

11.   Moore, J.K. Organization of the human superior olivary complex. *Microsc Res Tech*

51, 403-412 (2000).

12.   Pandya, D.N. Anatomy of the auditory cortex. *Rev Neurol (Paris)* **151**, 486-494

(1995).

13.   Bonetti, L*., et al.* Brain predictive coding processes are associated to COMT gene

Val158Met polymorphism. *NeuroImage* **233**, 117954 (2021).

14.   Bonetti, L*., et al.* Whole-brain computation of cognitive versus acoustic errors in

music: A mismatch negativity study. *Neuroimage: Reports* **2**, 100145 (2022).

15.   Bonetti, L*., et al.* Auditory sensory memory and working memory skills: association

between frontal MMN and performance scores. *Brain research* **1700**, 86-98 (2018).

16.   Bonetti, L., Haumann, N., Vuust, P., Kliuchko, M. & Brattico, E. Risk of depression

enhances auditory Pitch discrimination in the brain as indexed by the mismatch negativity.

Clinical Neurophysiology **128**, 1923-1936 (2017).

17.   Brattico, E., Winkler, I., Naatanen, R., Paavilainen, P. & Tervaniemi, M.

Simultaneous storage of two complex temporal sound patterns in auditory sensory memory.

Neuroreport **13**, 1747-1751 (2002).

18.   Conley, E.M., Michalewski, H.J. & Starr, A. The N100 auditory cortical evoked

potential indexes scanning of auditory short-term memory. *Clin Neurophysiol* **110**, 2086-

2093 (1999).

19.   Naatanen, R. & Picton, T. The N1 wave of the human electric and magnetic response

to sound: a review and an analysis of the component structure. *Psychophysiology* **24**, 375-425

(1987).

20.   Koelsch, S. Towards a neural basis of processing musical semantics. *Phys Life Rev* **8**,

89-105 (2011).

21.   Koelsch, S. Music□syntactic processing and auditory memory: Similarities and

differences between ERAN and MMN. *Psychophysiology* **46**, 179-190 (2009).

22.   Garrido, M.I., Kilner, J.M., Stephan, K.E. & Friston, K.J. The mismatch negativity: a

review of underlying mechanisms. *Clin Neurophysiol* **120**, 453-463 (2009).

23.   Rocchi, F.*, et al.* Common fronto-temporal effective connectivity in humans and

monkeys. *Neuron* **109**, 852-868 e858 (2021).

24.   Peretz, I. & Zatorre, R.J. *The cognitive neuroscience of music* (OUP Oxford, 2003).

25.   Hansen, P., Kringelbach, M. & Salmelin, R. *MEG: An introduction to methods*

(Oxford university press, 2010).

26.   Vuust, P., Heggli, O.A., Friston, K.J. & Kringelbach, M.L. Music in the brain. *Nature*

Reviews Neuroscience **23**, 287-305 (2022).

27.   Albouy, P., Weiss, A., Baillet, S. & Zatorre, R.J. Selective entrainment of theta

oscillations in the dorsal stream causally enhances auditory working memory performance.

Neuron **94**, 193-206. e195 (2017).

28.   Bonetti, L.*, et al.* Brain recognition of previously learned versus novel temporal

sequences: a differential simultaneous processing. *Cerebral Cortex*, bhac439 (2022).

29.   Bonetti, L.*, et al.* Spatiotemporal brain dynamics during recognition of the music of

Johann Sebastian Bach. *Biorxiv* (2020).

30. Bonetti, L*., et al.* Rapid encoding of musical tones discovered in whole-brain
connectivity. *NeuroImage* **245**, 118735 (2021).

31. Fernandez-Rubio, G*., et al.* Magnetoencephalography recordings reveal the
spatiotemporal dynamics of recognition memory for complex versus simple auditory
sequences. *Commun Biol* **5**, 1272 (2022).

32. Fernández-Rubio, G., Carlomagno, F., Vuust, P., Kringelbach, M.L. & Bonetti, L.
Associations between abstract working memory abilities and brain activity underlying long-
term recognition of auditory sequences. *PNAS Nexus* **1**, pgac216 (2022).

33. Bonetti, L*., et al.* Spatiotemporal brain dynamics during recognition of the music of
Johann Sebastian Bach. *bioRxiv* (2020).

34. Bruzzone, S.E.P*., et al.* Dissociated brain functional connectivity of fast versus slow
frequencies underlying individual differences in fluid intelligence: a DTI and MEG study. *Sci*
Rep **12**, 4746 (2022).

35. Fernández Rubio, G., Carlomagno, F., Vuust, P., Kringelbach, M.L. & Bonetti, L.
Associations between abstract working memory abilities and brain activity underlying long-
term recognition of auditory sequences. *bioRxiv* (2022).

36. Bonetti, L*., et al.* Temporal pattern recognition in the human brain: a dual
simultaneous processing. *bioRxiv* (2021).

37. Müllensiefen, D., Gingras, B., Stewart, L. & Musil, J.J. Goldsmiths Musical
Sophistication Index (Gold-MSI) v1. 0: Technical Report and Documentation Revision 0.3.
(2013).

38. Tzourio-Mazoyer, N*., et al.* Automated anatomical labeling of activations in SPM
using a macroscopic anatomical parcellation of the MNI MRI single-subject brain.
Neuroimage **15**, 273-289 (2002).

39. Colclough, G.L., Brookes, M.J., Smith, S.M. & Woolrich, M.W. A symmetric
multivariate leakage correction for MEG connectomes. *Neuroimage* **117**, 439-448 (2015).

40. Dietz, M.J., Nielsen, J.F., Roepstorff, A. & Garrido, M.I. Reduced effective
connectivity between right parietal and inferior frontal cortex during audiospatial perception
in neglect patients with a right-hemisphere lesion. *Hearing Research* **399**, 108052 (2021).

41. Garrido, M.I*., et al.* The functional anatomy of the MMN: a DCM study of the roving
paradigm. *Neuroimage* **42**, 936-944 (2008).

42. Fitzgibbon, S.P., Pope, K.J., Mackenzie, L., Clark, C.R. & Willoughby, J.O.
Cognitive tasks augment gamma EEG power. *Clin Neurophysiol* **115**, 1802-1809 (2004).

43.   Brattico, E*., et al.* Neural discrimination of nonprototypical chords in music experts

and laymen: an MEG study. *J Cogn Neurosci* **21**, 2230-2244 (2009).

44.   Näätänen, R., Paavilainen, P., Rinne, T. & Alho, K. The mismatch negativity (MMN)

in basic research of central auditory processing: a review. *Clinical neurophysiology* **118**,

2544-2590 (2007).

45.   Spratling, M.W. Predictive coding as a model of cognition. *Cognitive processing* **17**,

279-305 (2016).

46.   Brattico, E. & Pearce, M. The neuroaesthetics of music. *Psychology of Aesthetics,*

Creativity, and the Arts **7**, 48 (2013).

47.   Zatorre, R.J., Belin, P. & Penhune, V.B. Structure and function of auditory cortex:

music and speech. *Trends Cogn Sci* **6**, 37-46 (2002).

48.   Hasselmo, M.E. A Handbook for Modeling Hippocampal Circuits. (Frontiers

Research Foundation, 2011).

49.   Stern, C.E., Sherman, S.J., Kirchhoff, B.A. & Hasselmo, M.E. Medial temporal and

prefrontal contributions to working memory tasks with novel and familiar stimuli.

Hippocampus **11**, 337-346 (2001).

50.   Merkley, T.L., Larson, M.J., Bigler, E.D., Good, D.A. & Perlstein, W.M. Structural

and functional changes of the cingulate gyrus following traumatic brain injury: relation to

attention and executive skills. *J Int Neuropsychol Soc* **19**, 899-910 (2013).

51.   Kennerley, S.W., Walton, M.E., Behrens, T.E., Buckley, M.J. & Rushworth, M.F.

Optimal decision making and the anterior cingulate cortex. *Nat Neurosci* **9**, 940-947 (2006).

52.   Plakke, B. & Romanski, L.M. Auditory connections and functions of prefrontal

cortex. *Frontiers in neuroscience* **8**, 199 (2014).

53.   Denham, S.L. & Winkler, I. Predictive coding in auditory perception: challenges and

unresolved questions. *Eur J Neurosci* **51**, 1151-1160 (2020).

54.   Denison, R.N., Piazza, E.A. & Silver, M.A. Predictive Context Influences Perceptual

Selection during Binocular Rivalry. *Front Hum Neurosci* **5**, 166 (2011).

55.   Jehee, J.F. & Ballard, D.H. Predictive feedback can account for biphasic responses in

the lateral geniculate nucleus. *PLoS computational biology* **5**, e1000373 (2009).

56.   Spratling, M.W. Predictive coding as a model of response properties in cortical area

V1. *Journal of neuroscience* **30**, 3531-3543 (2010).

57.   Spratling, M.W. A single functional model of drivers and modulators in cortex. *J*

Comput Neurosci **36**, 97-118 (2014).

58.   Zatorre, R.J., Evans, A.C. & Meyer, E. Neural mechanisms underlying melodic
perception and memory for pitch. *J Neurosci* **14**, 1908-1919 (1994).

59.   Zatorre, R.J., Perry, D.W., Beckett, C.A., Westbury, C.F. & Evans, A.C. Functional
anatomy of musical processing in listeners with absolute pitch and relative pitch. *Proc Natl*
Acad Sci U S A **95**, 3172-3177 (1998).

60.   King, J.R. & Dehaene, S. Characterizing the dynamics of mental representations: the
temporal generalization method. *Trends Cogn Sci* **18**, 203-210 (2014).

61.   Vuust, P., Brattico, E., Seppanen, M., Naatanen, R. & Tervaniemi, M. The sound of
music: differentiating musicians using a fast, musical multi-feature mismatch negativity
paradigm. *Neuropsychologia* **50**, 1432-1443 (2012).

62.   Criscuolo, A., Pando-Naude, V., Bonetti, L., Vuust, P. & Brattico, E. An ALE meta-
analytic review of musical expertise. *Sci Rep* **12**, 11726 (2022).

63.   Proskovec, A.L., Wiesman, A.I., Heinrichs-Graham, E. & Wilson, T.W. Load effects
on spatial working memory performance are linked to distributed alpha and beta oscillations.
Hum Brain Mapp **40**, 3682-3689 (2019).

64.   Lenz, D., Schadow, J., Thaerig, S., Busch, N.A. & Herrmann, C.S. What's that sound?
Matches with auditory long-term memory induce gamma activity in human EEG.
International Journal of Psychophysiology **64**, 31-38 (2007).

65.   Slobounov, S., Tutwiler, R., Slobounova, E., Rearick, M. & Ray, W. Human
oscillatory brain activity within gamma band (30–50 Hz) induced by visual recognition of
non-stable postures. *Cognitive Brain Research* **9**, 177-192 (2000).

66.   Sauseng, P., Griesmayr, B., Freunberger, R. & Klimesch, W. Control mechanisms in
working memory: a possible function of EEG theta oscillations. *Neurosci Biobehav Rev* **34**,
1015-1022 (2010).

67.   Afnan, J.*, et al.* Validating MEG source imaging of resting state oscillatory patterns
with an intracranial EEG atlas. *Neuroimage* **274**, 120158 (2023).

68.   Tukey, J.W. Comparing individual means in the analysis of variance. *Biometrics* **5**,
99-114 (1949).

69.   Taulu, S. & Simola, J. Spatiotemporal signal space separation method for rejecting
nearby interference in MEG measurements. *Phys Med Biol* **51**, 1759-1768 (2006).

70.   Gross, J.*, et al.* Good practice for conducting and reporting MEG research.
Neuroimage **65**, 349-363 (2013).

71.   Woolrich, M., Hunt, L., Groves, A. & Barnes, G. MEG beamforming using Bayesian

PCA for adaptive data covariance matrix regularization. *Neuroimage* **57**, 1466-1479 (2011).

72.   Oostenveld, R., Fries, P., Maris, E. & Schoffelen, J.-M. FieldTrip: open source

software for advanced analysis of MEG, EEG, and invasive electrophysiological data.

Computational intelligence and neuroscience **2011**, 1-9 (2011).

73.   Woolrich, M.W.*, et al.* Bayesian analysis of neuroimaging data in FSL. *Neuroimage*

45, S173-S186 (2009).

74.   Penny, W.D., Friston, K.J., Ashburner, J.T., Kiebel, S.J. & Nichols, T.E. *Statistical*

parametric mapping: the analysis of functional brain images (Elsevier, 2011).

75.   Mantini, D.*, et al.* A signal-processing pipeline for magnetoencephalography resting-

state networks. *Brain Connect* **1**, 49-59 (2011).

76.   Steinwart, I. & Christmann, A. *Support vector machines* (Springer Science &

Business Media, 2008).

77.   Haufe, S.*, et al.* On the interpretation of weight vectors of linear models in

multivariate neuroimaging. *Neuroimage* **87**, 96-110 (2014).

78.   Cichy, R.M., Pantazis, D. & Oliva, A. Resolving human object recognition in space

and time. *Nat Neurosci* **17**, 455-462 (2014).

79.   Hoegholt, N.F.*, et al.* A magnetoencephalography study of first-time mothers

listening to infant cries. *Cereb Cortex* (2022).

80.   Brookes, M.J.*, et al.* Beamformer reconstruction of correlated sources using a

modified source model. *Neuroimage* **34**, 1454-1465 (2007).

81.   Hillebrand, A. & Barnes, G.R. Beamformer analysis of MEG data. *Int Rev Neurobiol*

68, 149-171 (2005).

82.   Huang, M., Mosher, J.C. & Leahy, R. A sensor-weighted overlapping-sphere head

model and exhaustive head model comparison for MEG. *Physics in Medicine & Biology* **44**,

423 (1999).

83.   Nolte, G. The magnetic lead field theorem in the quasi-static approximation and its

use for magnetoencephalography forward calculation in realistic volume conductors. *Phys*

Med Biol **48**, 3637-3652 (2003).

84.   Luckhoo, H.T., Brookes, M.J. & Woolrich, M.W. Multi-session statistics on

beamformed MEG data. *Neuroimage* **95**, 330-335 (2014).

85.   Hall, E.L., Woolrich, M.W., Thomaz, C.E., Morris, P.G. & Brookes, M.J. Using
variance information in magnetoencephalography measures of functional connectivity.
Neuroimage **67**, 203-212 (2013).

86.   Van Veen, B.D., van Drongelen, W., Yuchtman, M. & Suzuki, A. Localization of
brain electrical activity via linearly constrained minimum variance spatial filtering. *IEEE*
Trans Biomed Eng **44**, 867-880 (1997).

87.   Stephan, K.E.*, et al.* Ten simple rules for dynamic causal modeling. *Neuroimage* **49**,
3099-3109 (2010).

88.   David, O.*, et al.* Dynamic causal modeling of evoked responses in EEG and MEG.
Neuroimage **30**, 1255-1272 (2006).

89.   Penny, W.D. Comparing dynamic causal models using AIC, BIC and free energy.
Neuroimage **59**, 319-330 (2012).

90.   Penny, W.D.*, et al.* Comparing families of dynamic causal models. *PLoS Comput Biol*
6, e1000709 (2010).

91.   Bastos, A.M.*, et al.* Canonical microcircuits for predictive coding. *Neuron* **76**, 695-
711 (2012).

92.   Oane, I., Barborica, A. & Mindruta, I.R. Cingulate Cortex: Anatomy, Structural and
Functional Connectivity. *J Clin Neurophysiol* **40**, 482-490 (2023).

93.   Friston, K., Mattout, J., Trujillo-Barreto, N., Ashburner, J. & Penny, W. Variational
free energy and the Laplace approximation. *Neuroimage* **34**, 220-234 (2007).

94.   Rigoux, L., Stephan, K.E., Friston, K.J. & Daunizeau, J. Bayesian model selection for
group studies - revisited. *Neuroimage* **84**, 971-985 (2014).

95.   Daubechies, I. *Ten lectures on wavelets* (SIAM, 1992).

1530