Proximity labeling reveals a new *in vivo* network of interactors for the histone demethylase

KDM5

Matanel Yheskel[1], Simone Sidoli[2], and Julie Secombe[1,3,4]

[1] Department of Genetics, Albert Einstein College of Medicine, Bronx, New York, United States

of America.

[2] Department of Biochemistry, Albert Einstein College of Medicine, Bronx, New York, United

States of America.

[3] Dominick P. Purpura Department of Neuroscience, Albert Einstein College of Medicine, Bronx,

New York, United States of America.

[4] Lead contact and corresponding author: julie.secombe@einsteinmed.edu

Running title: KDM5-interacting proteins identified using proximity labeling

Key words: KDM5, histone demethylase, TurboID, proximity labeling, mass spectrometry,

chromatin modifiers, insulators

Summary

KDM5 family proteins are multi-domain regulators of transcription that when dysregulated

contribute to cancer and intellectual disability. KDM5 proteins can regulate transcription through

their histone demethylase activity in addition to demethylase-independent gene regulatory

functions that remain less characterized. To expand our understanding of the mechanisms that

contribute to KDM5-mediated transcription regulation, we used TurboID proximity labeling to

identify KDM5-interacting proteins. Using *Drosophila melanogaster*, we enriched for biotinylated

proteins from KDM5-TurboID-expressing adult heads, and generated a novel control for DNA-

adjacent background in the form of dCas9:TurboID. We identified both known and novel candidate

proteins, including members of the SWI/SNF and NURF chromatin remodeling complexes, the

non-specific lethal (NSL) complex, Mediator, and several insulator proteins. Combined, our data

shed new light on potential demethylase-independent activities of KDM5. In the context of KDM5

dysregulation, these interactions may play key roles in the alteration of evolutionarily conserved

transcriptional programs implicated in human disorders.

Introduction

Lysine demethylase 5 (KDM5) family proteins are multidomain transcriptional regulators able to recognize and enzymatically modify chromatin.[1,2] The best characterized function of KDM5 proteins is their histone demethylase activity, which cleaves a chromatin mark that is found at most active promoters, trimethylated lysine 4 on histone H3 (H3K4me3).[1,3–6] KDM5 proteins are evolutionarily conserved, with four paralogous genes in mammals encoding KDM5A-KDM5D, while animals with smaller genomes such as nematodes and flies possess a single *kdm5* gene. The importance of KDM5 function is emphasized by the observation that changes to the expression of this family of proteins is associated with two clinical outcomes: cancer and intellectual disability.[7–9] KDM5A and KDM5B are amplified or overexpressed in a range of cancers, including breast, ovarian, skin, and lung.[8,10–13] KDM5A/B appear to play several roles in tumorigenesis, including promoting cell cycle progression and regulating the metabolism of cancer stem cells.[14–16] In contrast to the gain of function seen in cancer cells, loss of function variants in the autosomal paralogs KDM5A, KDM5B, and the X-linked KDM5C have been observed in individuals with intellectual disability (ID).[17–21] KDM5 proteins have an evolutionarily conserved role in regulating critical gene expression programs in neurons as evidenced by morphological and functional neuronal phenotypes in KDM5B and KDM5C knockout mice. [22–24] Similarly, flies and nematodes with *kdm5* mutations display altered neuroanatomical development and/or neurotransmission. [25–28]

KDM5 catalytic function is mediated by the joint activity of the Jumonji N (JmjN) and JmjC domains and is classically thought to result in transcriptional repression. In addition, KDM5 proteins possess other potential gene regulatory domains, including plant homeodomain domain (PHD) motifs that can recognize H3K4me2/3 or H3K4me0, and a potential DNA binding A/T interaction domain (ARID).[29–34] These binding domains likely function in-concert with the histone

demethylase activity of KDM5 by, for example, recruiting it to target promoters through the activity

of individual or combinations of accessory domains. Conversely, non-enzymatic functions of these

domains and/or other motifs of KDM5 that have no currently known function, such as the C_5HC_2

domain, could regulate transcription through distinct mechanisms. There is ample evidence that

KDM5 proteins can regulate transcription independently of their demethylase activity. For

instance, KDM5 is essential for viability in flies in a manner that is independent of its histone

demethylase activity.[33,35] In addition, both demethylase-dependent and independent functions of

KDM5 are critical for *Drosophila* neuronal development and function. [36,37] Interestingly, some

missense alleles of KDM5C observed in individuals with intellectual disability diminish its

enzymatic activity, while others do not.[23,38–40] Similarly, demethylase dependent and independent

activities of KDM5 proteins are likely to be important for their contributions to the etiology of and

spread of cancers.[41,42] For example, KDM5B demethylase-independent functions in breast cancer

promote metastatic potential to the lung.[41,42] Thus, even though KDM5 proteins derive their name

from their enzymatic function, other conserved motifs contribute to their gene regulatory activities,

although these activities remain much less characterized.

Understanding the repertoire of gene regulatory mechanisms utilized by KDM5 family proteins

requires a comprehensive understanding of the proteins they can interact with. Traditional

immunoprecipitation coupled with mass spectrometry (IP-MS) approaches have been used to

identify proteins that form complexes with KDM5 family proteins in both mammals and

Drosophila.[40,43–45] These experiments have revealed several conserved interactions, most notably

with histone deacetylase 1 (HDAC1) and other proteins known to associate with this chromatin

modifier.[46,47] To expand our understanding of the proteins that function with KDM5 to mediate its

gene regulatory activities, we used TurboID-mediated proximity labeling.[48] This has been shown

to be a powerful technique to identify weak or transient interactions that may otherwise be

disturbed during the process of traditional IP experiments.[48–55] This technique takes advantage of

the promiscuous biotin ligase activity of TurboID, which results in the biotinylation of lysine

residues within 10 nm of its active site. When expressed as a chimeric fusion to a protein of

interest, interacting proteins will be biotin-labeled.[56,48] Covalently modified proteins are then

recovered with streptavidin beads and prepared for liquid chromatography-tandem mass

spectrometry (LC-MS/MS). By expressing KDM5 that was N- or C-terminally tagged with TurboID

in vivo, we recovered about half of previously identified interactions in *Drosophila*, and almost all

interactions known to be conserved in mammalian cells, clearly demonstrating the robustness of

this technique. Furthermore, we have discovered a novel interactome for KDM5 that suggests

roles in the function of the SWI/SNF, non-specific lethal (NSL), NURF, and Mediator complexes,

in addition to chromatin insulation.

Results

Chimeric TurboID-KDM5 proteins are functional and broadly biotinylate

To identify KDM5 interactors *in vivo*, we created constructs in which KDM5 was N- or C-terminally

tagged with TurboID to maximally identify proteins that could function with KDM5. Because our

long-term goal is to further develop our *Drosophila* model of KDM5-induced intellectual disability

(ID), we chose to carry out our TurboID studies using adult heads to enrich for neuronal tissue,

using the general workflow shown in Fig 1A. Generating a TurboID system that closely mimics

endogenous *kdm5* expression has been shown to be important for delivering more specific

biotinylation compared to overexpression.[57] Based on our prior generation of a UASp-*kdm5*

transgene that is expressed at approximately endogenous levels when crossed to a range of Gal4

drivers, we generated transgenic flies harboring HA-tagged UASp-TurboID:*kdm5* and UASp-

kdm5:TurboID (Fig. 1B).[58] Therefore, we generated both N-terminal (NT-KDM5) and C-terminal

(CT-KDM5) TurboID fusions of KDM5 to understand the full breadth of its interactome and to

highlight terminus-specific interactions. To test the functionality and expression of the chimeric

KDM5-TurboID proteins, we expressed them ubiquitously in a *kdm5[140]* null mutant background.[33]

Western blot analysis using adult heads showed that NT-KDM5 and CT-KDM5 were expressed

at levels similar to those observed from an endogenously HA-tagged KDM5 (Fig. 1C). Importantly,

N- and C-terminally tagged KDM5 proteins were able to restore viability to $kdm5^{140}$ null mutant

flies, which normally die prior to adulthood (Fig. 1D).[35] Tagging KDM5 with TurboID therefore does

not interfere with its function. Flies in which TurboID-KDM5 was the only source of KDM5

($kdm5^{140}$;Ubi-Gal4>UASp;TurboID:$kdm5$) were used for all subsequent experiments to maximize

the number of interactors identified.

Determining the proper controls to identify the KDM5 proximitome

To confidently identify proteins that function with KDM5, appropriate controls are critical.

Due to the novelty of the technique, there is no standard set of controls for proximity labeling

experiments. Many studies simply enrich over endogenous biotinylation and bead background.[59–

62] Other studies expressed forms of TurboID alone that were localized to the specific cellular

compartment that the protein of interest resided, such as the cellular membrane.[48,49,51,52] As a

non-TurboID-expressing wild-type control with a similar genetic background, we used a fly strain

in which endogenous $kdm5$ is removed and HA-tagged KDM5 is expressed using its endogenous

promoter from a transgene inserted at the same locus as the TurboID constructs

($kdm5^{140}$;$gkdm5^{WT}$). We will refer to these as control animals. We also generated a transgene

able to express nuclear localized, HA-tagged, TurboID alone using the same UAS promoter used

for CT-$kdm5$ and NT-$kdm5$, in an effort to assay general nuclear background (Fig. 1B). To

compare the levels of TurboID alone to TurboID-KDM5 we expressed these transgenes using

Ubi-Gal4 in a wild-type and $kdm5^{140}$ background, respectively. Unsurprisingly, given the small

size of TurboID (36kDa) compared to NT-KDM5 and CT-KDM5 (240kDa), anti-HA western blot

from adult heads showed significantly higher levels of expression for TurboID alone, possibly

creating high levels of background biotinylation in this strain (Fig. 1C). Because biotin is essential

for animal viability and thus included in the standard fly food used for crosses and stock

maintenance, we assessed the ability of all TurboID transgenes to biotinylate proteins when

expressed using Ubi-Gal4 by probing with infrared-conjugated streptavidin. Compared to control

flies, similar levels of biotin-conjugated proteins were observed in heads expressing TurboID, NT-

KDM5 and CT-KDM5, demonstrating their ability to biotinylate *in vivo* (Fig 1C).

To identify proteins preferentially biotinylated by NT-KDM5 and CT-KDM5 compared to

control and TurboID alone, we carried out streptavidin-bead pulldowns in quadruplicate followed

by liquid chromatography and mass spectrometry (LC-MS/MS). This experiment (Experiment 1)

identified a total of 1332 proteins, 476 of which are found in the nucleus where we have previously

shown KDM5 to be localized.[27] Principal component analysis (PCA) of normalized nuclear protein

abundances showed that TurboID alone, NT-KDM5, and CT-KDM5 clustered together, but were

distinct from controls (Fig S1A). We therefore compared the proteins identified in NT-KDM5 and

CT-KDM5 to control heads which revealed enrichment of 172 and 184 proteins, respectively,

using a *p*-value cutoff of 0.05 (Fig S1B, C; Table S1). 136 proteins were commonly enriched by

N- and C-terminally tagged KDM5, suggesting that we can robustly detect proteins in proximity to

KDM5 (Fig. S1D). To assess the quality of our data, we determined how many known KDM5

interactors were identified in our analyses. Fifteen proteins have been established to form a

complex with *Drosophila* KDM5 through IP-MS studies or targeted co-IP experiments (Table S2).

Suggesting the robustness with which the TurboID approach identifies *bona fide* KDM5-

associated proteins, 7 known interactors were identified by NT-KDM5 and 6 by CT-KDM5 (47%

and 40%, respectively). We additionally assessed biotinylated protein enrichment of NT-KDM5

and CT-KDM5 compared to TurboID alone (Fig S1E, F). As expected, based on the higher level

of expression of this protein compared to TurboID-tagged KDM5, a high level of background was

observed in these flies. This resulted in fewer proteins being enriched in NT-KDM5 and CT-KDM5

(32 and 61, respectively), reduced overlap between the datasets, and a reduction in the number

of known interactors identified (Fig S1G, H). Interestingly, TurboID alone appears to show bias in

its biotinylation of nuclear proteins, as comparing TurboID to control revealed significant

enrichment of 199 proteins, a majority of which are involved in chromatin-mediated transcriptional

regulation (Fig. S1I).

Because of concerns related to the use of TurboID alone as a control, we generated a

transgene encoding an enzymatically inactive form of the Cas9 enzyme (dCas9) fused to an HA-

tagged nuclear localized TurboID (UASp-dCas9:TurboID). This dCas9:TurboID fusion protein is

more similar in size to the KDM5 fusion proteins, being 194kDa and 235kDa, respectively,

compared to 36kDa for TurboID alone. In addition, Cas9 and dCas9 can scan the DNA, potentially

making this fusion an appropriate control for chromatin binding proteins such as KDM5 by

restricting biotinylation to DNA-adjacent proteins.[63] Ubi-Gal4-mediated expression of this

transgene revealed that dCas9:TurboID was expressed at similar levels to the KDM5-TurboID

fusion proteins and was able to biotinylate (Fig. 1C). Repeating the proximity labeling experiment,

we carried out triplicate streptavidin-bead pulldowns from heads of control, TurboID,

dCas9:TurboID, NT-KDM5 and CT-KDM5 flies (Experiment 2). MS analyses identified 1,146

proteins across all samples, 203 of which were nuclear. PCA from this second experiment showed

that NT-KDM5 and CT-KDM5 clustered together, indicating that these datasets are more alike to

each other than to any of the controls. Like our first experiment, TurboID alone clustered with NT-

KDM5 and CT-KDM5, and clustered distinctly from control and dCas9:TurboID samples (Fig 2A).

Using these data, we compared NT-KDM5 and CT-KDM5 to control, dCas9:TurboID, and to

TurboID alone (Fig. 2B; Table S3). Proteins enriched in the KDM5 samples compared to control

or TurboID gave similar results to those obtained in the first experiment (Fig 2B). Using control

flies as reference, 82 proteins were identified using NT-KDM5 and 68 for CT-KDM5, with 61 of

these proteins being identified in both datasets (Fig 2C-E). Compared to TurboID alone, only 29

and 24 proteins were enriched for NT-KDM5 and CT-KDM5, with 13 overlapping between the two

datasets (Fig. 2F-H). Importantly, we find comparing NT-KDM5 and CT-KDM5 with

dCas9:TurboID yielded data very similar to that seen using control animals, despite providing a

higher biotinylation background. 66 and 59 proteins were enriched in NT-KDM5 and CT-KDM5,

respectively, with an overlap of 48 proteins (Fig. 2I-K).

To complete our characterization of dCas9:TurboID as a tool to identify enriched DNA-

adjacent proteins, we compared these data to both TurboID alone and to control. Similar to data

from the first experiment comparing TurboID and control, TurboID alone shows a preference for

biotinylating a large number of chromatin-related proteins, even compared to dCas9:TurboID (Fig.

S2A). Comparing dCas9:TurboID to control revealed enrichment in a relatively small number of

proteins that were enriched for transcriptional-regulatory proteins consistent with the ability of

dCas9:TurboID to biotinylate targets while scanning DNA (Fig S2B, C). Confirming the challenges

related to using TurboID alone, the consistency with which proteins were enriched in NT-KDM5

or CT-KDM5 compared to TurboID alone was very low, with very little agreement across

experiments using a p-value cutoff <0.05 or <0.1 (Fig S2D, E). Moreover, the number of previously

identified interactors remained low in experiment two when comparing to TurboID alone, with only

5 and 3 being identified in NT-KDM5 and CT-KDM5, respectively (Fig. S2F). We therefore suggest

that endogenous biotinylation or dCas9:TurboID are superior to TurboID as controls for proximity

labeling experiments where the protein of interest is nuclear-specific and DNA-adjacent.

Proximity labeling identifies new potential KDM5 interacting complexes

To build a high confidence list of proteins which interact with KDM5, we combined data from

experiment 1 comparing NT-KDM5 and CT-KDM5 to control (2 datasets), and experiment 2 in

which NT-KDM5 and CT-KDM5 were compared to control (2 datasets) as well as dCas9:TurboID

(2 datasets). To do this, we began by filtering for enriched nuclear proteins across all six datasets

using a p-value<0.1. We first filtered for proteins identified in at least two of six datasets. Then to

include the possibility of terminus-exclusive interactors, we required that at each terminus proteins

had to be identified in 0 (exclusive to other terminus) or in 2 of 3 datasets (Fig. 3A). Demonstrating

the power of this approach, this includes 7 of the 15 (47%) known *Drosophila* KDM5 interactors.

This ratio increases to 7 of 8 (88%) for those interactors that have been shown in both *Drosophila*

and mammals. In addition, we identified several proteins not previously been found to be

Drosophila KDM5 interactors, but have been purified using KDM5A, KDM5B, KDM5C and/or

KDM5D in mammalian cells. These include the nucleosome remodeler Mi-2, the chromatin

assembly factor 1 (Caf1;Caf1-55), the actyl-lysine binding protein Zymnd8 and the

heterochromatin-associated protein HP1c (Fig 3B). [6,46,47] With these stringent filtering criteria, we

identified a total of 87 proteins (Fig. 3C). These high confidence interactors are highlighted in red

in volcano plots comparing NT-KDM5 and CT-KDM5 to dCas9:TurboID (Fig. 3D, E). All

comparisons are shown in Supplemental Figure 3.

Our proximity-labeling studies using Turbo-KDM5 revealed a broader interactome than

previously described in the literature. To better understand the relationships between the proteins

identified in our study, we generated a protein interaction map using Cytoscape and STRING (Fig.

4A).[64,65] Gene Ontology (GO) analysis shows that many of these proteins have roles in the

regulation of gene expression, chromatin modification, and chromatin remodeling (Fig. 4B). In

addition to confirming the strong link between KDM5 and Sin3/HDAC1-containing complexes,

these analyses also highlighted interactions with new protein complexes. Among these, we find

proteins such as BEAF-32, Chromator (Chro), Putzig (Pzg), and Cp190 that function in regulating

genomic architecture, suggesting a unstudied role for KDM5 in this process.[66–71] To confirm our

MS data, we performed western blots on input and streptavidin pulldowns and probed with

antibodies specific for BEAF-32 and CP190. This confirmed enrichment of these two proteins in

both NT-KDM5 and CT-KDM5 compared to dCas9:TurboID-expressing and control flies (Fig. 4C).

The extensive biotinylation of chromatin-related proteins, such as BEAF-32 and CP190, by

TurboID alone is also evident in this pulldown, as discussed previously. In addition, we identified

proteins critical for forming the transcriptional pre-initiation complex (TPIC), which is consistent

with the promoter-proximal binding of KDM5 proteins across species.[27,46,72–74,74] We analyzed a

recent crystal structure of the human TPIC and found that distinct surfaces interacted with KDM5,

directly revealing the specificity of biotinylation using TurboID-KDM5. Specifically, three adjacent

proteins in the mediator complex (MED1, MED14 and MED17) and three adjacent subunits of

TFIID (Taf4, Taf6, and Taf9) were identified in our analyses (Fig. 4D).[75] This suggests that KDM5

may play a direct role in enhancer-promoter communications that regulate the transcriptional

activity of target genes.

KDM5 and newly identified interactors occupy overlapping genomic binding sites in

***Drosophila* and human cells**

To further explore the relationship between KDM5 and newly identified interactors in

Drosophila, we compared their genomic binding with those of interactors using publicly available

ChIP-seq datasets. We used published KDM5 ChIP-seq data from whole adult flies to interrogate

ChIP-Atlas as a means to identify datasets from any *Drosophila* cell type that significantly overlap

(via permutation 100X).[76] We then overlapped these with our high confidence interactor list to

reveal a total of 27 overlapping datasets. 30 proteins in our interactor list had available data on

ChIP Atlas. (Fig. 5A, B). These overlapping datasets included known interactors such as Sin3A,

in addition to BEAF-32 that we confirmed to be in proximity to KDM5, both of which show that

KDM5 flanks their DNA-binding sites (Fig. 5C, D). Representatives of other newly identified

proteins that are members of several different complexes were also examined, including the DNA

replication-related element factor Dref, the chromatin remodeler IswI, the WD40-repeat protein

will die slowly (wds) and the acetyl-histone binding Brd2/Brd4 ortholog fs(1)h (Fig. 5E-H). The

similar binding profiles of all these proteins can also be seen in representative genome browser

tracks (Fig. 5I)

To assess the extent to which our high confidence KDM5 interactors might be

evolutionarily conserved, we investigated genomic co-occupancy in human cells. We first

converted our 87 *Drosophila* high confidence interactors to their human ortholog(s), which

resulted in a total of 138 proteins due to humans possessing multiple paralogous proteins for

some *Drosophila* proteins (DIOPT v8.0[77] score > 8/15). Due to the strong link between KDM5B

and breast cancer, and the wealth of ChIP-seq datasets available in cell lines derived from this

cancer type, we used KDM5B data from MCF-7 cells for these analyses.[78] Using peaks called

from this ChIP-seq data, we interrogated all available breast cancer cell line datasets again using

ChIP-Atlas (100X permutation). This revealed that 26 candidate interactors had binding profiles

that significantly overlapped with KDM5B binding (Fig. 6A, B). Interestingly, these included the

KDM5 paralogs KDM5A and KDM5C, suggesting that there may be overlapping or redundant

function for these proteins. Similar to our studies using *Drosophila*, some proteins identified were

known interactors, such as SIN3A that is known show similar genomic binding to KDM5B (Fig.

6C).[79] Other proteins overlapped with our genomic binding studies in *Drosophila*, including Brd4

(fs(1)h) and WDR5 (wds), while other proteins were identified because datasets were available in

human cells and not *Drosophila* (Fig. 6 C,D). These included the Mediator subunit MED1 and the

small ubiquitin-like modifier SUMO2 (*Drosophila* Sumo) (Fig. 6E,F). A combined genome browser

emphasizes the colocalization of these proteins with KDM5B (Fig. 6G). Thus, the high confidence

KDM5 interactors identified in *Drosophila* may be important for the function of KDM5B and other

KDM5 paralogs in mammalian cells.

Identified KDM5 interactors are implicated in neurodevelopmental disorders

Based on the association between genetic variants in KDM5A, KDM5B, and KDM5C and

intellectual disability (ID) and autism spectrum disorder (ASD), KDM5-interactors could potentially

be implicated in neurological disorders.[80–85] To examine this in more detail, we used the Simons

Foundation Autism Research Initiative (SFARI) genes as an up-to-date source of genes with

significant causal links to ASD.[86] The 1231 genes in this database are scored based on the level

of confidence of association, with a score of 1 being the strongest link, in addition to whether ASD

occurs as part of a syndrome (S). Using our list of 138 human ortholog-converted KDM5

interactors, we found that 26 of these overlap with SFARI ASD-associated proteins (p=8e-08; Fig.

7A-B). Some of these proteins have clear links to each other, such as TAF4 and TAF6 that are

components of TFIID, while others are associated with numerous other aspects of transcriptional

regulation. To look more broadly into the link between KDM5 interactors and neurodevelopmental

disorders, we used the Developmental Brain Disorder Gene Database (DBD) which is a curated

list of genes implicated in disorders such as ID, ASD, attention deficit hyperactivity disorder

(ADHD) and schizophrenia.[87] Twenty-four human-converted orthologs represented in DBD have

been shown to contribute to ID, ASD, ADHD, and Schizophrenia (Fig. 7C). Unsurprisingly given

the frequency that ID and ASD co-occur, 14 proteins were identified in both datasets, including

KDM5B, KDM5C and BRD4. In addition, 9 ID-associated proteins were identified, including

MED17, the NURD chromatin remodeling complex component GATAD2A and the C-terminal

binding protein (CtBP) transcriptional repressor. Combined, these analyses potentially greatly

expand our understanding of the KDM5-network of proteins and provide new avenues for

investigating the links between KDM5 family proteins and the etiology of neurodevelopmental

disorders. A summary of our KDM5 interaction data highlighting proteins with known roles in

transcriptional regulation is shown in Figure 7D.

Discussion

Here we describe the interactome of *Drosophila* KDM5 in the adult head using TurboID-

mediated proximity labeling. To identify the broadest selection of potential interactors, we N- and

C-terminally TurboID-tagged KDM5 as some interactors were expected to be in proximity to both

termini while others might be terminus specific. Importantly, TurboID-KDM5 chimeric proteins

were functional, as they were able to rescue the lethality caused by a *kdm5^{140}* null allele. Using

NT-KDM5 and CT-KDM5, we performed two experiments to optimize the experimental controls,

as there are none currently established for this relatively new technique. Our study revealed that

expression of TurboID alone led to high background levels of biotinylation, particularly of

chromatin-related proteins. In contrast, using control (endogenous biotinylation and bead

background) and dCas9:TurboID provided similar and more reasonable background to identify

proteins enriched by expression of TurboID-KDM5. Because we carried out two separate MS

experiments, we were able to stringently filter our data to retain only those proteins that were

nuclear localized and showed high reproducibility and rigor. This led to the identification of 87

KDM5 high confidence interactors, 11 of which were previously described in either *Drosophila* or

mammals. Notably, while we refer to proteins identified in our TurboID analyses as interactors,

we acknowledge that proximity labeling does not necessarily detect direct interactions. However,

TurboID biotinylates lysine residues within 10 nm of its active site[56], which is equivalent to about

27 bp of B-DNA (8 bp per 3.4 nm). However unlikely, this short biotinylation radius can result in

the identification of proteins that are nearby but in a distinct complex that is not physically touching

KDM5. While these proteins are not in the same complex(es) as KDM5, they could still function

with KDM5 to regulate gene expression. For simplicity, we will refer TurboID-enriched proteins as

interactors.

Several lines of evidence allow us to have confidence in our described KDM5 interactome.

In addition to the reproducibility with which proteins were detected by NT- and CT-KDM5 across

the two experiments, many of the proteins identified had previously known physical connectivity

with each other. Thus, rather than identifying individual components of complexes, we identified

proteins well known to complex with each other, such as the SWI/SNF and NURF chromatin

remodeling complexes. Interestingly, a functional link between KDM5 and these complexes is

supported by studies in mouse embryonic stem cells which showed that a loss of KDM5B altered

nucleosome position surrounding the transcription start site (TSS), although the mechanism was

not revealed. We also identified the insulator proteins BEAF-32, Chromator, Putzig, and CP190

which complex together.[68,69] Our previous investigation in a fly strain harboring an allele

associated with human intellectual disability identified enrichment of BEAF-32 binding sites at

dysregulated genes.[28] Functionally, it is also notable that mutations in *kdm5*, *BEAF-32*, and *putzig*

all modify position effect variegation (PEV) suggesting that these proteins may function together

to regulate chromatin compaction and/or organization.[88–90] For some TurboID-identified proteins

such as the Mediator complex components (MED1, MED14, and MED17), as well as the TFIID

proteins (Taf4, Taf6, and Taf9), published structural data provides clear support for their link to

KDM5.[75,91] The Mediator proteins and Taf proteins neighbor each other, respectively, in the hTPIC

crystal structure. Moreover, we found that KDM5 interactions could be mapped to distinct surfaces

at the hTPIC providing clear evidence for KDM5 localization with respect to key transcriptional

initiation machinery. This specificity additionally suggests that the enrichment of these

transcriptional preinitiation proteins is not simply due to the binding of KDM5 proteins near the

promoter region of its target genes. If that were the case, then the entire preinitiation complex

would have been identified in our datasets, including TBP and RNA Pol II. Binding specificity in

this context can provide insight into how KDM5 carries out its functions at target gene promoters

to influence gene expression. Although limited by the number of available ChIP-seq datasets

available, corroborating evidence for our interactors also comes from the extensive overlap in

genomic binding observed in *Drosophila* and/or mammalian cells.

KDM5s relationship with other gene regulatory complexes provides insight into the links

between KDM5 dysregulation and human disorders. Many of the interacting complexes identified

in our study have been implicated in tumorigenesis, including NSL and SWI/SNF.[92–95]

Furthermore, like KDM5, MED1 has been implicated as a transcriptional coactivator that mediates

breast cancer metastasis and treatment resistance.[96,97] Identification of KDM5 interactors may

provide insight to novel mechanisms of KDM5-mediated transcriptional regulation which underlie

tumor development and progression. Changes to protein interactions could also contribute to the

intellectual disability seen in individuals with genetic variants in KDM5A, KDM5B, or KDM5C.

Indeed, for mutations that do not alter histone demethylase activity, this may be a contributor to

cognitive dysfunction. Our analyses of KDM5 interactors revealed an enrichment in proteins found

to be altered in neurodevelopmental disorders whose clinical presentations overlap with those

seen for KDM5 genes. KDM5 and interactor mutations could therefore influence

neurodevelopment through common pathways. Altogether, our study suggests that KDM5 likely

functions through numerous transient interactions with interconnected complexes to regulate

gene expression in a context-dependent manner.

ACKNOWLEDGMENTS

We thank members of the Secombe Lab, Melissa Castiglione, Blair Scheider, Michael Rogers,

and Bethany Terry, as well as Michelle Schumacher for their feedback throughout this project and

edits on the manuscript. We also thank the laboratory of Maria Cristina Gambetta for providing

the anti-CP190 antibody and appreciate the availability of stocks from the Bloomington *Drosophila*

Stock Center (NIH P40OD018537). We are additionally grateful to the Cancer Center Support

Grant P30CA013330. This research was supported by the NIH Ruth L. Kirschstein National

Research Service Award F31GM146347 and the Einstein MSTP Training Grant T32GM007288

to M.Y., NIH R01GM112783 and support from the Irma T. Hirschl Trust to J.S. The Sidoli lab

gratefully acknowledges for financial support AFAR (Sagol Network GerOmics award), Deerfield

(Xseed award), Relay Therapeutics, Merck, the NIH Office of the Director (1-S10-OD030286-01),

the Einstein-Mount Sinai Diabetes Research Center, and the Einstein Cancer Center (P30-

CA013330-47).

DECLARATION OF INTERESTS

403   The authors declare no competing interests.

INCLUSION AND DIVERSITY

One or more of the authors of this paper self-identifies as underrepresented ethnic minority in science. One or more of the authors self-identifies as a member of the LGBTQ+ community. One or more of the authors received support from a program designed to increase minority representation in science.

AUTHOR CONTRIBUTIONS

Conceptualization, M.Y., J.S.; Methodology, M.Y., S.S.; Investigation, M.Y. and J.S.; Writing – original draft, J.S. and M.Y., Writing – Reviewing and Editing, J.S., M.Y., and S.S..; Funding acquisition, J.S., M.Y and S.S., Supervision, J.S. and S.S.

	FIGURE LEGENDS

Figure 1: TurboID-tagged KDM5 proteins are functional and biotinylate endogenous

proteins

(A) Schematic of the workflow used for purifying and fingerprinting biotinylated proteins from

adult heads using LC-MS/MS mass spectrometry.

(B) Schematic of the four UASp constructs generated able to express HA-tagged TurboID

alone, dCas9:TurboID, NT-KDM5 and CT-KDM5.

(C) Western blot using adult heads showing levels of expression of KDM5 using anti-HA,

biotinylation using a streptavidin-680 conjugate and the loading control alpha-tubulin.

Genotypes: $kdm5^{140}$; $gkdm5:HA$ (a wild-type strain; Control*)*, Ubi-Gal4/+ ; UASp-

TurboID/+ (TurboID), Ubi-Gal4/+ ; UASp-*dCas9:TurboID* (dCas9:TurboID), $kdm5^{140}$, Ubi-

Gal4/$kdm5^{140}$; UASp-NT- $kdm5$/+ (NT-KDM5) and $kdm5^{140}$, Ubi-Gal4/$kdm5^{140}$; UASp-CT-

$kdm5$ /+ (CT-KDM5).

(D) Rescue of $kdm5^{140}$-induced lethality by ubiquitous expression of UAS-NT-*KDM5* or UAS-

CT-*KDM5* using Ubi-Gal4. Genotype of male and female adult flies shown is $kdm5^{140}$, Ubi-

Gal4/$kdm5^{140}$; UASp-NT- $kdm5$ /+ (NT-KDM5) and $kdm5^{140}$, Ubi-Gal4/$KDM5^{140}$; UASp-

CT- $kdm5$ / + (CT-KDM5).

Figure 2: Identifying candidate KDM5 interactors through TurboID

(A) Principal component analysis of normalized nuclear protein abundances from control (Ctl;

grey), dCas9:TurboID (T-dCas9; green), TurboID (black), NT-KDM5 (red) and CT-KDM5

(blue).

(B) Summary of known *Drosophila* KDM5 interactors, whether the interaction is conserved in

mammals (mouse or human) and the identification of these proteins in experiment 2

compared to control (Ctl) and dCas9:TurboID. Three interactors identified in mammalian

cells but not previously in *Drosophila* are also included (Caf1-55, Mi-2 and Zmynd8).

(C) Volcano plot showing data comparing biotinylated proteins enriched by NT-KDM5 to

443   control.

444   (D) Volcano plot showing data comparing CT-KDM5 to control.

445   (E) Venn diagram showing overlap of the proteins identified using NT-KDM5 and CT-KDM5

446   compared to control (Ctl).

447   (F) Volcano plot showing data comparing NT-KDM5 to TurboID.

448   (G) Volcano plot showing data comparing CT-KDM5 to TurboID.

449   (H) Venn diagram showing overlap of the proteins identified comparing NT-KDM5 and CT-

450   KDM5 to TurboID.

451   (I) Volcano plot showing data comparing NT-KDM5 to dCas9:TurboID.

452   (J) Volcano plot showing data comparing CT-KDM5 to dCas9:TurboID.

453   (K) Venn diagram showing overlap of the proteins identified using NT-KDM5 and CT-KDM5

454   compared to dCas9:TurboID.

455   For all volcano plots shown, known interactors are indicated with text. Red dots indicate

456   significantly enriched proteins ($p<0.05$) and the dotted line on Y-axis indicates $p=0.05$.

458   **Figure 3: Identification of a high confidence KDM5 interactors**

459   (A) Filtering workflow for identification of high confidence KDM5 interactors by combining data

460   from experiments 1 and 2.

461   (B) Summary of known *Drosophila* KDM5 interactors, whether the interaction is conserved in

462   mammals (mouse or human) and the identification of these proteins in experiment 1 and

463   experiment 2 compared to control and dCas9:TurboID. Three interactors identified in

464   mammalian cells but not previously in *Drosophila* are also included (Caf1-55, Mi-2 and

465   Zmynd8).

(C) High confidence interactors based on their identification in experiment 1 (compared to

control) and experiment 2 (compared to control and dCas9:TurboID). Dark red box

indicates enrichment of *p*<0.05, pink indicates *p*<0.1.

(D) Volcano plot showing data comparing NT-KDM5 to dCas9:TurboID with high confidence

KDM5 interactors from panel C indicated with red dots. The dotted line on Y-axis indicates

p=0.05.

(E) Volcano plot showing data comparing CT-KDM5 to dCas9:TurboID with high confidence

KDM5 interactors from panel C indicated with red dots. The dotted line on Y-axis indicates

p=0.05.

Figure 4: TurboID-tagged KDM5 biotinylates proteins involved in several aspects of gene

expression regulation.

(A) STRING analyses of nuclear proteins that were significantly biotinylated by NT-KDM5 and

CT-KDM5. Grey lines indicate known physical interactions between proteins. Darker lines

indicate a higher confidence of interaction. Cytoscape was used to manually cluster

annotated proteins based on their STRING Cluster Enrichment and known functions.

Unconnected notes are not shown.

(B) Gene Ontology Biological Process (GO-BP) analyses of high confidence proteins.

(C) Anti-BEAF-32 and anti-CP190 blot from biotinylated proteins purified from adult heads of

the genotypes shown. Input shows the presence of BEAF-32 and CP190, respectively,

across all samples.

(D) Structure of the human pre-initiation complex (7ENA) showing proteins identified as high

confidence interactors in our TurboID data in red bubbles. DNA (yellow), TBP (pink

bubbles) and RNA polymerase II (cyan bubbles) are also shown.

Figure 5: Proteins enriched in KDM5 TurboID experiments show overlapping genomic binding profiles.

(A) Venn diagram showing overlap between high confidence biotinylated proteins (red) and datasets enriched when comparing published KDM5 ChIP-seq data from whole adult flies to other *Drosophila* ChIP datasets (blue). A total of 27 datasets showed significant overlap using ChIP-atlas. 30 interactors had ChIP-seq datasets in the ChIP Atlas database (green).

(B) Volcano plot showing fold enrichment and p-values of the permutation analyses between KDM5 and 27 ChIP-seq datasets of high confidence interactors.

(C) Genomic binding profiles of Sin3A ChIP from S2 cells and KDM5 showing similar localization.

(D) Genomic binding profiles of BEAF-32 ChIP from S2 cells and KDM5 showing similar localization.

(E) Genomic binding profiles of Dref ChIP from Kc167 cells and KDM5 showing similar localization.

(F) Genomic binding profiles of Iswi ChIP from Kc167 cells and KDM5 showing similar localization.

(G) Genomic binding profiles of wds ChIP from BG3 cells and KDM5 showing similar localization.

(H) Genomic binding profiles of fs(1)h ChIP from Kc167 cells and KDM5 showing similar localization.

(I) Representative genome browser image showing binding of KDM5, Sin3A, BEAF-32, Dref, fs(1)h, Iswi and wds.

Figure 6: Human orthologs of proteins enriched in *Drosophila* KDM5 TurboID experiments show overlapping genomic binding profiles.

(A) Venn diagram showing overlap between human orthologs of *Drosophila* KDM5 high confidence interactors and their enrichment when comparing published KDM5B ChIP-seq data to breast cancer ChIP-seq datasets using ChIP-Atlas. 26 datasets significantly overlapped.

(B) Volcano plot showing fold enrichment and p-values of the permutation analyses between KDM5B and 26 ChIP-seq datasets of human-ortholog converted high confidence interactors.

(C) Genomic binding profiles of KDM5B, SIN3A and BRD4 showing similar genome-wide binding. Binding is shown relative to KDM5B due to the much larger number of SIN3A and BRD4 binding sites in the genome compared to KDM5B.

(D) Genomic binding profiles of WDR5 (ortholog of *Drosophila* wds) ChIP from MCF-10 cells and KDM5B showing similar localization.

(E) Genomic binding profiles of MED1 ChIP from MCF-7 cells and KDM5B showing similar localization.

(F) Genomic binding profiles of SUMO2 (ortholog of *Drosophila* Sumo) ChIP from MCF-7 cells and KDM5B showing similar localization.

(G) Representative genome browser image showing binding of KDM5B, SIN3A, MED1, BRD4, WDR5 and SUMO2.

Figure 7: A subset of KDM5 interactors are implicated in neurodevelopmental disorders

(A) Overlap between human orthologs of *Drosophila* KDM5 interactors and genes associated with ASD using the SFARI database. S indicates syndromic ASD. Scores indicate confidence of causal association, with 1 indicating strongest link.

(B) 26 candidate interacting proteins identified and their SFARI score.

(C) 24 candidate interacting proteins were identified as being implicated in ID, ASD and/or schizophrenia using DBD.

543  (D) Model for KDM5 interactions with key transcriptional proteins that are likely to impact the

544    expression of downstream target genes.

547 **MATERIALS AND METHODS**

548 **Fly strains and care**

549 Fly crosses were maintained at 25°C with 50% humidity and a 12-hour light/dark cycle. Food (per

550 liter) contained 18g yeast, 22g molasses, 80g malt extract, 9g agar, 65 cornmeal, 2.3g methyl

551 para-benzoic acid, 6.35ml propionic acid. The number of male and female animals were equal

552 across the genotypes examined. The $kdm5^{140}$ null allele has been previously described. [35]

554 **Cloning and Transgenesis**

555 The N- and C-terminally TurboID-tagged constructs were generated by cloning the coding region

556 of *kdm5* upstream or downstream of HA:TurboID from pCDNA3-TurboID

557 (RRID:Addgene_107171)[48] in the pUASpattB vector (RRID:DGRC_1358). UASp-HA:TurboID

558 with a NLS was generated by cloning HA:TurboID into the same UASpattB vector. UASp-

559 dCas:HA:Turbo:NLS was made by combining the dCas9 open reading frame from SID3s-dCas9-

560 KRAB (RRID:Addgene_106399) with HA:TurboID:NLS in the pUASpattB vector

561 (RRID:DGRC_1358). All transgenes were generated by injection into y^1 M{RFP[3xP3.PB]

562 GFP[E.3xP3]=vas-int.Dm}ZH-2A w* ; M{3xP3-RFP.attP}ZH-86Fb at BestGene Inc.

564 **Western Blotting**

565 Western analyses were carried out as previously described. [35] Briefly, five 2- to 5-day old adult fly

566 heads were homogenized in 2x NuPAGE LDS sample buffer, sonicated for 10 mins, treated with

567 DTT, run on a 4-12% Bis-Tris 1 mm gel and transferred to a PVDF membrane. The following

568 primary antibodies were used: mouse anti-HA (1:1000, Cell Signaling Technology Cat# 2367,

RRID: AB_10691311), Streptavidin 680 (1:10,000, ThermoFisher, Streptavidin Alexa Fluor 680

conjugate), rabbit anti-alpha-Tubulin (1:5000, Cell Signaling Technology Cat# 2144,

RRID:AB_2210548), mouse anti-BEAF-32 (1:500, DSHB Cat# anti-BEAF, RRID:AB_1553420),

and rabbit anti-CP190 (1:5000, Kaushal et. al 2021[98], non-commercial). Secondary antibodies

used were IRDye® 680RD Donkey anti-Mouse IgG (1:5000; LI-COR Biosciences Cat# 925-

68072, RRID: AB_2814912) and IRDye® 800CW Donkey anti-Rabbit IgG (1:5000; LI-COR

Biosciences Cat# 926-32213, RRID: AB_621848). Blots were scanned and processed using a LI-

COR Odyssey Infrared scanner.

Purifying and identifying proteins using TurboID

Biotinylated Protein Enrichment

2-5 day-old flies were flash frozen in liquid nitrogen and decapitated and a total of ten heads were

used per sample. Heads were homogenized in 250 μL RIPA Buffer (Thermofisher 89901)

supplemented with Halt™ Protease Inhibitor Cocktail (Thermofisher, 78430) and centrifuged at 4

°C for 10 minutes at 15,000XG to remove debris. 100 μL of Pierce™ Streptavidin Magnetic Beads

(Thermofisher, 88817) were washed twice with RIPA and the cleared lysate was added. The

lysate-bead mixture was incubated with rotation at 4 °C overnight. The next day the lysate was

discarded, and beads were washed twice with RIPA, once with 1M KCl, once with 0.1 M Na_2HCO_3,

once with 1 M Urea in 10 mM Tris pH 8.0, and twice again with RIPA. For Western Blot analyses

all RIPA was removed and biotinylated proteins were eluted with 4X NuPAGE™ LDS Sample

Buffer (Invitrogen, NP0007) supplemented with 2 mM biotin and 20 mM DTT.

On-bead protein digestion

Proteins were digested directly on streptavidin beads. 5 mM DTT and 50 mM ammonium

bicarbonate (pH = 8) were added to the solution and left on the bench for about 1 hour for disulfide

bond reduction. Samples were then alkylated with 20 mM iodoacetamide in the dark for 30

minutes. Afterward, 500 ng of trypsin was added to the samples, which were digested at 37 °C

for 18 h. The peptide solution was dried in a vacuum centrifuge.

Sample desalting

597   Prior to mass spectrometry analysis, samples were desalted using a 96-well plate filter (Orochem)

598   packed with 1 mg of Oasis HLB C-18 resin (Waters). Briefly, the samples were resuspended in

599   100 µl of 0.1% TFA and loaded onto the HLB resin, which was previously equilibrated using 100

600   µl of the same buffer. After washing with 100 µl of 0.1% TFA, the samples were eluted with a

601   buffer containing 70 µl of 60% acetonitrile and 0.1% TFA and then dried in a vacuum centrifuge.

LC-MS/MS Acquisition and Analysis

Samples were resuspended in 10 µl of 0.1% TFA and loaded onto a Dionex RSLC Ultimate 300

(Thermo Scientific), coupled online with an Orbitrap Fusion Lumos (Thermo Scientific).

Chromatographic separation was performed with a two-column system, consisting of a C-18 trap

cartridge (300 µm ID, 5 mm length) and a picofrit analytical column (75 µm ID, 25 cm length)

packed in-house with reversed-phase Repro-Sil Pur C18-AQ 3 µm resin. Peptides were separated

using a 90 min gradient from 4-30% buffer B (buffer A: 0.1% formic acid, buffer B: 80% acetonitrile

+ 0.1% formic acid) at a flow rate of 300 nL/min. The mass spectrometer was set to acquire

spectra in a data-dependent acquisition (DDA) mode. Briefly, the full MS scan was set to 300-

1200 m/z in the orbitrap with a resolution of 120,000 (at 200 m/z) and an AGC target of 5×10^{e5}.

MS/MS was performed in the ion trap using the top speed mode (2 secs), an AGC target of 1×10^{e4}

and an HCD collision energy of 35. Raw files were searched using Proteome Discoverer software

(v2.4, Thermo Scientific) using SEQUEST search engine and the UniProt database of Drosophila

melanogaster. The search for total proteome included variable modification of N-terminal

acetylation, and fixed modification of carbamidomethyl cysteine. Trypsin was specified as the

digestive enzyme with up to 2 missed cleavages allowed. Mass tolerance was set to 10 pm for

precursor ions and 0.2 Da for product ions. Peptide and protein false discovery rate was set to

1%. Following the search, data was processed as described by Aguilan et al.[99]. Briefly, proteins

were log2 transformed, normalized by the average value of each sample and missing values were

imputed using a normal distribution 2 standard deviations lower than the mean. Statistical

regulation was assessed using heteroscedastic T-test (if *p*-value < 0.05). Data were assumed to

be Gaussian distributed but this was not formally tested.

Bioinformatic Analyses

The Enrichment Analysis function on ChIP Atlas[76] was used to perform permutation tests.

Experiment types were set to "TFs and others", Cell type class was set to 'All types' and 'Breast'

for Drosophila and Human respectively. A 100X random permutation of each was used as the

control. For *Drosophila*, selected profiles were generated using: KDM5 (SRX1084165), Sin3A

(SRX1158165), BEAF-32 (SRX386677), Dref (SRX170984), Iswi (SRX5346167), wds

(SRX2399656), and fs(1)h (SRX203000). For Human, selected profiles generated using: KDM5B

(SRX3285561), SIN3A (SRX190318), MED1 (SRX673749), WDR5 (SRX3958389), BRD4

(SRX5089551), and SUMO2 (SRX3541112). Deeptools(3.5.1)[100] computeMatrix and

plotHeatmap functions were used to make profiles and heatmaps, and Pygenometracks(3.7)[101]

was used to visualize ChIP-seq tracks.

References

1. Pavlenko, E., Ruengeler, T., Engel, P. & Poepsel, S. Functions and interactions of mammalian

KDM5 demethylases. *Frontiers in genetics* **13**, 906662 (2022).

2. Holowatyj, A., Yang, Z.-Q. & Pile, L. A. Histone lysine demethylases in Drosophila

melanogaster. *Fly (Austin)* **9**, 36–44 (2015).

3. Lee, N. *et al.* The trithorax-group protein Lid is a histone H3 trimethyl-Lys4 demethylase.

Nature Structural & Molecular Biology **14**, 341–343 (2007).

4. Secombe, J., Li, L., Carlos, L. & Eisenman, R. N. The Trithorax group protein Lid is a

trimethyl histone H3K4 demethylase required for dMyc-induced cell growth. *Genes &*

Development **21**, 537–551 (2007).

5. Klose, R. J. *et al.* The retinoblastoma binding protein RBP2 is an H3K4 demethylase. *Cell*

128, 889–900 (2007).

6. Klein, B. J. *et al.* The histone-H3K4-specific demethylase KDM5B binds to its substrate and

product through distinct PHD fingers. *Cell reports* **6**, 325–335 (2014).

7. Yang, G.-J. *et al.* The emerging role of KDM5A in human cancer. *J Hematol Oncol* **14**, 30

(2021).

8. Blair, L. P., Cao, J., Zou, M. R., Sayegh, J. & Yan, Q. Epigenetic regulation by lysine

demethylase 5 (KDM5) enzymes in cancer. *Cancers* **3**, 1383–1404 (2011).

9. Hatch, H. A. M. & Secombe, J. Molecular and cellular events linking variants in the histone

demethylase KDM5C to the intellectual disability disorder Claes-Jensen syndrome. *FEBS J*

(2021) doi:10.1111/febs.16204.

10.   Yamamoto, S. *et al.* JARID1B is a luminal lineage-driving oncogene in breast cancer.

Cancer Cell **25**, 762–777 (2014).

11.   Hinohara, K. *et al.* KDM5 histone demethylase activity links cellular transcriptomic

heterogeneity to therapeutic resistance. *Cancer Cell* **34**, 939-953.e9 (2018).

12.   Horton, J. R. *et al.* Characterization of a linked jumonji domain of the KDM5/JARID1

family of histone H3 lysine 4 demethylases. *The Journal of Biological Chemistry* **291**, 2631–

2646 (2016).

13.   Arifuzzaman, S., Khatun, M. R. & Khatun, R. Emerging of lysine demethylases (KDMs):

From pathophysiological insights to novel therapeutic opportunities. *Biomedicine &*

Pharmacotherapy (Biomedecine & Pharmacotherapie) **129**, 110392 (2020).

14.   He, R. *et al.* H3K4 demethylase KDM5B regulates cancer cell identity and epigenetic

plasticity. *Oncogene* **41**, 2958–2972 (2022).

15.   Xhabija, B. & Kidder, B. L. KDM5B is a master regulator of the H3K4-methylome in

stem cells, development and cancer. *Semin Cancer Biol* **57**, 79–85 (2019).

16.   Roesch, A. *et al.* Overcoming intrinsic multidrug resistance in melanoma by blocking the

mitochondrial respiratory chain of slow-cycling JARID1B(high) cells. *Cancer Cell* **23**, 811–

825 (2013).

17.   Iwase, S. *et al.* The X-linked mental retardation gene SMCX/JARID1C defines a family

of histone H3 lysine 4 demethylases. *Cell* **128**, 1077–1088 (2007).

18.   Hatch, H. A. M., O'Neil, M. H., Marion, R. W., Secombe, J. & Shulman, L. H.

Caregiver-reported characteristics of children diagnosed with pathogenic variants in KDM5C.

American Journal of Medical Genetics. Part A (2021) doi:10.1002/ajmg.a.62381.

19.   Abidi, F. E. *et al.* Mutations in JARID1C are associated with X-linked mental retardation,

short stature and hyperreflexia. *Journal of Medical Genetics* **45**, 787–793 (2008).

20.   Tzschach, A. *et al.* Novel JARID1C/SMCX mutations in patients with X-linked mental

retardation. *Human Mutation* **27**, 389 (2006).

21.   Jensen, L. R. *et al.* Mutations in the JARID1C gene, which is involved in transcriptional

regulation and chromatin remodeling, cause X-linked mental retardation. *American Journal of*

Human Genetics **76**, 227–236 (2005).

22.   Harrington, J., Wheway, G., Willaime-Morawek, S., Gibson, J. & Walters, Z. S.

Pathogenic KDM5B variants in the context of developmental disorders. *Biochim Biophys Acta*

Gene Regul Mech **1865**, 194848 (2022).

23.   Iwase, S. *et al.* A Mouse Model of X-linked Intellectual Disability Associated with

Impaired Removal of Histone Methylation. *Cell Rep* **14**, 1000–1009 (2016).

24.   Scandaglia, M. *et al.* Loss of Kdm5c Causes Spurious Transcription and Prevents the

Fine-Tuning of Activity-Regulated Enhancers in Neurons. *Cell Rep* **21**, 47–59 (2017).

25.   Mariani, L., Lussi, Y. C., Vandamme, J., Riveiro, A. & Salcini, A. E. The H3K4me3/2

histone demethylase RBR-2 controls axon guidance by repressing the actin-remodeling gene

wsp-1. *Development* **143**, 851–863 (2016).

26.   Belalcazar, H. M., Hendricks, E. L., Zamurrad, S., Liebl, F. L. W. & Secombe, J. The

histone demethylase KDM5 is required for synaptic structure and function at the Drosophila

neuromuscular junction. *Cell reports* **34**, 108753 (2021).

27.   Hatch, H. A., Belalcazar, H. M., Marshall, O. J. & Secombe, J. A KDM5-Prospero

transcriptional axis functions during early neurodevelopment to regulate mushroom body

formation. *eLife* **10**, (2021).

28.   Zamurrad, S., Hatch, H. A. M., Drelon, C., Belalcazar, H. M. & Secombe, J. A

Drosophila Model of Intellectual Disability Caused by Mutations in the Histone Demethylase

KDM5. *Cell Rep* **22**, 2359–2369 (2018).

29.   Yao, W., Peng, Y. & Lin, D. The flexible loop L1 of the H3K4 demethylase JARID1B

ARID domain has a crucial role in DNA-binding activity. *Biochem Biophys Res Commun* **396**,

323–328 (2010).

30.   Tu, S. *et al.* The ARID domain of the H3K4 demethylase RBP2 binds to a DNA

CCGCCC motif. *Nat Struct Mol Biol* **15**, 419–421 (2008).

31.   Scibetta, A. G. *et al.* Functional analysis of the transcription repressor PLU-1/JARID1B.

Mol Cell Biol **27**, 7220–7235 (2007).

32.   Villaseñor, R. *et al.* ChromID identifies the protein interactome at chromatin marks. *Nat*

Biotechnol **38**, 728–736 (2020).

33.   Li, L., Greer, C., Eisenman, R. N. & Secombe, J. Essential functions of the histone

demethylase lid. *PLoS Genetics* **6**, e1001221 (2010).

34.   Wang, G. G. *et al.* Haematopoietic malignancies caused by dysregulation of a chromatin-

binding PHD finger. *Nature* **459**, 847–851 (2009).

35.   Drelon, C., Belalcazar, H. M. & Secombe, J. The histone demethylase KDM5 is essential

for larval growth in drosophila. *Genetics* **209**, 773–787 (2018).

36.   Belalcazar, H. M., Hendricks, E. L., Zamurrad, S., Liebl, F. L. W. & Secombe, J. The

histone demethylase KDM5 is required for synaptic structure and function at the Drosophila

neuromuscular junction. *Cell Rep* **34**, 108753 (2021).

37.   Hatch, H. A., Belalcazar, H. M., Marshall, O. J. & Secombe, J. A KDM5-Prospero

transcriptional axis functions during early neurodevelopment to regulate mushroom body

formation. *Elife* **10**, e63886 (2021).

38.   Brookes, E. *et al.* Mutations in the intellectual disability gene KDM5C reduce protein

stability and demethylase activity. *Hum Mol Genet* **24**, 2861–2872 (2015).

39.   Vallianatos, C. N. *et al.* Altered Gene-Regulatory Function of KDM5C by a Novel

Mutation Associated With Autism and Intellectual Disability. *Front Mol Neurosci* **11**, 104

(2018).

40.   Tahiliani, M. *et al.* The histone H3K4 demethylase SMCX links REST target genes to X-

linked mental retardation. *Nature* **447**, 601–605 (2007).

41.   Paroni, G. *et al.* HER2-positive breast-cancer cell lines are sensitive to KDM5 inhibition:

definition of a gene-expression model for the selection of sensitive cases. *Oncogene* **38**, 2675–

2689 (2019).

42.   Cao, J. *et al.* Histone demethylase RBP2 is critical for breast cancer progression and

metastasis. *Cell Rep* **6**, 868–877 (2014).

43.   Moshkin, Y. M. *et al.* Histone chaperones ASF1 and NAP1 differentially modulate

removal of active histone marks by LID-RPD3 complexes during NOTCH silencing. *Mol Cell*

35, 782–793 (2009).

44.   Varier, R. A. *et al.* Recruitment of the Mammalian Histone-modifying EMSY Complex

to Target Genes Is Regulated by ZNF131. *J Biol Chem* **291**, 7313–7324 (2016).

45.   Nishibuchi, G. *et al.* Physical and functional interactions between the histone H3K4

demethylase KDM5A and the nucleosome remodeling and deacetylase (NuRD) complex. *J*

Biol Chem **289**, 28956–28970 (2014).

46.   Ohguchi, H. *et al.* Lysine Demethylase 5A Is Required for MYC-Driven Transcription in

749   Multiple Myeloma. *Blood Cancer Discovery* **2**, 370–387 (2021).

750   47.   Nishibuchi, G. *et al.* Physical and functional interactions between the histone H3K4

751   demethylase KDM5A and the nucleosome remodeling and deacetylase (NuRD) complex. *The*

752   *Journal of Biological Chemistry* **289**, 28956–28970 (2014).

753   48.   Branon, T. C. *et al.* Efficient proximity labeling in living cells and organisms with

754   TurboID. *Nature Biotechnology* **36**, 880–887 (2018).

755   49.   Shokri, L. *et al.* A Comprehensive Drosophila melanogaster Transcription Factor

756   Interactome. *Cell reports* **27**, 955-970.e7 (2019).

757   50.   Uezu, A. *et al.* Identification of an elaborate complex mediating postsynaptic inhibition.

758   *Science* **353**, 1123–1129 (2016).

759   51.   Zhang, B., Zhang, Y. & Liu, J.-L. Highly effective proximate labeling in Drosophila. *G3*

760   *(Bethesda, Md.)* **11**, (2021).

761   52.   Szczesniak, L. M., Bonzerato, C. G. & Wojcikiewicz, R. J. H. Identification of the bok

762   interactome using proximity labeling. *Frontiers in cell and developmental biology* **9**, 689951

763   (2021).

764   53.   Mannix, K. M., Starble, R. M., Kaufman, R. S. & Cooley, L. Proximity labeling reveals

765   novel interactomes in live Drosophila tissue. *Development* **146**, (2019).

766   54.   Sanchez, A. D. *et al.* Proximity labeling reveals non-centrosomal microtubule-organizing

767   center components required for microtubule growth and localization. *Current Biology* **31**,

768   3586-3600.e11 (2021).

55.   Chua, X. Y., Aballo, T., Elnemer, W., Tran, M. & Salomon, A. Quantitative

Interactomics of Lck-TurboID in Living Human T Cells Unveils T Cell Receptor Stimulation-

Induced Proximal Lck Interactors. *Journal of Proteome Research* **20**, 715–726 (2021).

56.   Cho, K. F. *et al.* Proximity labeling in mammalian cells with TurboID and split-TurboID.

Nature Protocols **15**, 3971–3999 (2020).

57.   Stockhammer, A. *et al.* When less is more – Endogenous tagging with TurboID as a tool

to study the native interactome of adaptor protein complexes. *bioRxiv* 2021.11.19.469212

(2022) doi:10.1101/2021.11.19.469212.

58.   Drelon, C., Rogers, M. F., Belalcazar, H. M. & Secombe, J. The histone demethylase

KDM5 controls developmental timing in Drosophila by promoting prothoracic gland

endocycles. *Development* **146**, dev182568 (2019).

59.   Artan, M. *et al.* Interactome analysis of Caenorhabditis elegans synapses by TurboID-

based proximity labeling. *Journal of Biological Chemistry* **297**, 101094 (2021).

60.   Hu, S. *et al.* Identification of mutant p53-specific proteins interaction network using

TurboID-based proximity labeling. *Biochemical and Biophysical Research Communications*

615, 163–171 (2022).

61.   Chevalier, B. *et al.* Differential CFTR-Interactome Proximity Labeling Procedures

Identify Enrichment in Multiple SLC Transporters. *Int J Mol Sci* **23**, 8937 (2022).

62.   Uçkun, E. *et al.* In vivo Profiling of the Alk Proximitome in the Developing Drosophila

Brain. *Journal of Molecular Biology* **433**, 167282 (2021).

63.   Burmistrz, M., Krakowski, K. & Krawczyk-Balska, A. RNA-Targeting CRISPR-Cas

Systems and Their Applications. *Int J Mol Sci* **21**, E1122 (2020).

64.   Szklarczyk, D. *et al.* The STRING database in 2021: customizable protein-protein

792   networks, and functional characterization of user-uploaded gene/measurement sets. *Nucleic*

793   *Acids Res* **49**, D605–D612 (2021).

794   65.   Shannon, P. *et al.* Cytoscape: a software environment for integrated models of

795   biomolecular interaction networks. *Genome Research* **13**, 2498–2504 (2003).

796   66.   Kaushal, A. *et al.* Essential role of Cp190 in physical and regulatory boundary formation.

797   *Science Advances* **8**, eabl8834 (2022).

798   67.   Ramírez, F. *et al.* High-resolution TADs reveal DNA sequences underlying genome

799   organization in flies. *Nature Communications* **9**, 189 (2018).

800   68.   Sabirov, M. *et al.* Mechanisms of CP190 Interaction with Architectural Proteins in

801   Drosophila Melanogaster. *International Journal of Molecular Sciences* **22**, (2021).

802   69.   Melnikova, L. S., Molodina, V. V., Kostyuchenko, M. V., Georgiev, P. G. & Golovnin,

803   A. K. The BEAF-32 Protein Directly Interacts with Z4/putzig and Chriz/Chromator Proteins

804   in Drosophila melanogaster. *Doklady. Biochemistry and biophysics* **498**, 184–189 (2021).

805   70.   Chathoth, K. T. *et al.* The role of insulators and transcription in 3D chromatin

806   organization of flies. *Genome Research* (2022) doi:10.1101/gr.275809.121.

807   71.   Schwartz, Y. B. & Cavalli, G. Three-Dimensional Genome Organization and Function in

808   Drosophila. *Genetics* **205**, 5–24 (2017).

809   72.   He, R. & Kidder, B. L. H3K4 demethylase KDM5B regulates global dynamics of

810   transcription elongation and alternative splicing in embryonic stem cells. *Nucleic Acids*

811   *Research* **45**, 6427–6441 (2017).

812   73.   Hinohara, K. *et al.* KDM5 Histone Demethylase Activity Links Cellular Transcriptomic

813   Heterogeneity to Therapeutic Resistance. *Cancer Cell* **34**, 939-953.e9 (2018).

74.   Liu, X., Greer, C. & Secombe, J. KDM5 interacts with Foxo to modulate cellular levels

of oxidative stress. *PLoS Genetics* **10**, e1004676 (2014).

75.   Chen, X. *et al.* Structures of the human Mediator and Mediator-bound preinitiation

complex. *Science* **372**, eabg0635 (2021).

76.   Zou, Z., Ohta, T., Miura, F. & Oki, S. ChIP-Atlas 2021 update: a data-mining suite for

exploring epigenomic landscapes by fully integrating ChIP-seq, ATAC-seq and Bisulfite-seq

data. *Nucleic Acids Research* **50**, W175–W182 (2022).

77.   Hu, Y. *et al.* An integrative approach to ortholog prediction for disease-focused and other

functional studies. *BMC Bioinformatics* **12**, 357 (2011).

78.   Hinohara, K. *et al.* KDM5 Histone Demethylase Activity Links Cellular Transcriptomic

Heterogeneity to Therapeutic Resistance. *Cancer Cell* **34**, 939-953.e9 (2018).

79.   Spain, M. M., Caruso, J. A., Swaminathan, A. & Pile, L. A. Drosophila SIN3 isoforms

interact with distinct proteins and have unique biological functions. *J Biol Chem* **285**, 27457–

27467 (2010).

80.   Zamurrad, S., Hatch, H. A. M., Drelon, C., Belalcazar, H. M. & Secombe, J. A

drosophila model of intellectual disability caused by mutations in the histone demethylase

KDM5. *Cell reports* **22**, 2359–2369 (2018).

81.   Iwase, S. *et al.* A Mouse Model of X-linked Intellectual Disability Associated with

Impaired Removal of Histone Methylation. *Cell reports* **14**, 1000–1009 (2016).

82.   Poeta, L., Padula, A., Lioi, M. B., van Bokhoven, H. & Miano, M. G. Analysis of a set of

KDM5C regulatory genes mutated in neurodevelopmental disorders identifies temporal

coexpression brain signatures. *Genes* **12**, (2021).

83.   Gonçalves, T. F. *et al.* KDM5C mutational screening among males with intellectual

disability suggestive of X-Linked inheritance and review of the literature. *European Journal*

of Medical Genetics **57**, 138–144 (2014).

84.   Brookes, E. *et al.* Mutations in the intellectual disability gene KDM5C reduce protein

stability and demethylase activity. *Human Molecular Genetics* **24**, 2861–2872 (2015).

85.   Hatch, H. A. M. & Secombe, J. Molecular and cellular events linking variants in the

histone demethylase KDM5C to the intellectual disability disorder Claes-Jensen syndrome.

The FEBS Journal (2021) doi:10.1111/febs.16204.

86.   Abrahams, B. S. *et al.* SFARI Gene 2.0: a community-driven knowledgebase for the

autism spectrum disorders (ASDs). *Mol Autism* **4**, 36 (2013).

87.   Gonzalez-Mantilla, A. J., Moreno-De-Luca, A., Ledbetter, D. H. & Martin, C. L. A

Cross-Disorder Method to Identify Novel Candidate Genes for Developmental Brain

Disorders. *JAMA Psychiatry* **73**, 275–283 (2016).

88.   Eggert, H., Gortchakov, A. & Saumweber, H. Identification of the Drosophila interband-

specific protein Z4 as a DNA-binding zinc-finger protein determining chromosomal structure.

Journal of Cell Science **117**, 4253–4264 (2004).

89.   Shalaby, N. A. *et al.* Systematic discovery of genetic modulation by Jumonji histone

demethylases in Drosophila. *Scientific Reports* **7**, 5240 (2017).

90.   Maharjan, M., McKowen, J. K. & Hart, C. M. Overlapping but Distinct Sequences Play

Roles in the Insulator and Promoter Activities of the Drosophila BEAF-Dependent scs'

Insulator. *Genetics* **215**, 1003–1012 (2020).

91.   Yin, J. & Wang, G. The Mediator complex: a master coordinator of transcription and cell

lineage development. *Development* **141**, 977–987 (2014).

92.   Ferreyra Solari, N. E. *et al.* The NSL Chromatin-Modifying Complex Subunit KANSL2

Regulates Cancer Stem-like Properties in Glioblastoma That Contribute to Tumorigenesis.

Cancer Res **76**, 5383–5394 (2016).

93.   Mittal, P. & Roberts, C. W. M. The SWI/SNF complex in cancer - biology, biomarkers

and therapy. *Nat Rev Clin Oncol* **17**, 435–448 (2020).

94.   Wanior, M., Krämer, A., Knapp, S. & Joerger, A. C. Exploiting vulnerabilities of

SWI/SNF chromatin remodelling complexes for cancer therapy. *Oncogene* **40**, 3637–3654

(2021).

95.   Singh, M. *et al.* Histone Acetyltransferase MOF Orchestrates Outcomes at the Crossroad

of Oncogenesis, DNA Damage Response, Proliferation, and Stem Cell Development. *Mol*

Cell Biol **40**, e00232-20 (2020).

96.   Cao, J. *et al.* Histone demethylase RBP2 is critical for breast cancer progression and

metastasis. *Cell Rep* **6**, 868–877 (2014).

97.   Cui, J. *et al.* Cross-talk between HER2 and MED1 regulates tamoxifen resistance of

human breast cancer cells. *Cancer Res* **72**, 5625–5634 (2012).

98.   Kaushal, A. *et al.* CTCF loss has limited effects on global genome architecture in

Drosophila despite critical regulatory functions. *Nature Communications* **12**, 1011 (2021).

99.   Aguilan, J. T., Kulej, K. & Sidoli, S. Guide for protein fold change and p-value

calculation for non-experts in proteomics. *Mol Omics* **16**, 573–582 (2020).

100.   Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data

analysis. *Nucleic Acids Research* **44**, W160–W165 (2016).

101.   Ramírez, F. *et al.* High-resolution TADs reveal DNA sequences underlying genome

organization in flies. *Nat Commun* **9**, 189 (2018).

Fig1

Fig 2

Fig 3

A

Total Proteins | Exp 1 1332 | Exp 2 1146

Nuclear Proteins | 476 | 203

p<0.1, FC>0 | NT-KDM5: 171 CT-KDM5: 183 | NT-KDM5: 92 (62) CT-KDM5: 78 (56)

In 2+ Datasets | 198

N or C in 2+ Datasets | 87

Final list | High Confidence Interactors

B

Previously Identified Interactors (Drosophila / Mammalian: KDM5A, KDM5B, KDM5C, KDM5D) vs Exp 1 NT-KDMT/Ctl, Exp 1 CT-KDM5/Ctl, Exp 2 NT-KDM5/Ctl, Exp 2 CT-KDM5/Ctl, Exp 2 NT-KDM5/dCas9:TurboID, Exp 2 CT-KDM5/dCas9:TurboID

Rpd3/HDAC1, Sin3A/SIN3A, CG3815/PF1, CG15356/EMSY, CG13367/GATAD1, MRG15, Nap1/NAP1, CtBP/CTBP, Caf1-55/RBBP7, Mi-2/CHD3, HP1c/CBX3, CG1815/Zmynd8, Su(H)/RBPJ, Myc/MYC, Foxo/FOXO, Asf1/ASF1A/B, MCM4, MCM6, Ash2/ASH2L

Identified | Not identified

C High Confidence Interactors

p-value scale: 0, 0.05, 0.1, 1

Row labels: BEAF–32, Brms1, Caf1; Caf1–55, Chro, crp, dom, Dsp1, E(bx), east, fs(1)h, Hcf, HDAC1, IntS1, Kdm5, kis, lola, MED1, MED14, MEP–1, MESR4, mip130, mor, MRG15, p47, polybromo, pzg, Rcd1, row, Sap130, sbb, Sin3A, stwl, Taf4, Top1, Trl, wds, CG13367/GATAD1, CG3815/PF1, CG6340, Iswi, Lam, MED17, Rcc1, Rcd5, Bap55, brm, CG15356, CG7379, Cp190, CtBP, HIPP1, Hsc70–4, Hsp27, Jarid2, Mi–2, Mtor, Nap1, nsl1, Nup153, Nurf–38, Snr1, Srrm1, Sumo, Taf6, Taf9, trr, woc, caz, cg, CG1815/Zmynd8, Dref, His3.3A, HP1c, LBR, Rpt4, Saf–B, upSET, crol, IntS3, Msp300, psq, Rpt6, simj, sls, Ssrp, tsr, Uba2

Column labels: Exp 1 NT-KDMT/Ctl, Exp 1 CT-KDM5/Ctl, Exp2 CT-KDM5/Ctl, Exp 2 CT-KDM5/dCas9:TurboID, Exp 2 NT-KDM5/Ctl, Exp 2 NT-KDM5/dCas9:TurboID

D Exp 2 NT–KDM5/dCas9:TurboID

Up: 66 , Down: 15 , Unchanged: 122

x-axis: Log_2 fold change; y-axis: $-Log_{10} P$

E Exp 2 CT–KDM5/dCas9:TurboID

Up: 59 , Down: 21 , Unchanged: 123

x-axis: Log_2 fold change; y-axis: $-Log_{10} P$

Fig 4

A

NURF
MESR4, Iswi, polybromo, Dref, Nurf-38, E(bx), CtBP, Caf1

Sin3A & SWI/SNF
simj, mor, upSET, brm, Snr1, caz, MEP-1, Brms1, CG3815, Mi-2, CG7379, CG15356, Rpd3, Sin3A, Sap130

DNA-Binding
stwl, sbb, Ssrp, kis, Dsp1, Rcc1, row, His3.3A, woc, mip130, crp, Nap1, CG1815, psq, lola

NSL Complex
nsl1, wds, Bap55, Rcd5, Hcf, MRG15, Rcd1, dom, trr

Transcription Initiation
MED17, MED14, MED1, e(y)1, Taf4, Top1, Taf6

Integrator
IntS1, IntS3

Kdm5

Chromatin Insulators
HIPP1, Cp190, HP1c, Trl, pzg, BEAF-32, fs(1)h, Chro

Nuclear Pore
Nup153, LBR, Lam, Mtor

Other
Hsp27, Uba2, Msp300, Jarid2, east, smt3, sls, Hsc70-4, CG13367, SRm160, Saf-B

● Known Interactor
○ Newly Identified Interactor

B High Confidence Interactor GO–BP

- chromatin organization
- positive regulation of transcription, DNA–templated
- positive regulation of RNA biosynthetic process
- positive regulation of nucleic acid–templated transcription
- positive regulation of macromolecule biosynthetic process
- positive regulation of RNA metabolic process
- chromatin remodeling

GeneRatio: 0.30, 0.35, 0.40

Count: 24, 26, 28, 30, 32, 34

p.adjust: 5.588455e−17, 4.191342e−17, 2.794228e−17, 1.397114e−17, 1.235419e−28

C
Control, TurboID, dCas9:TurboID, NT-KDM5, CT-KDM5

BEAF-32 IP
BEAF-32 Input
CP190 IP
CP190 Input

D
MED1, 14, 17

90 degrees

MED1, 14, 17

RNA Pol II
TBP
DNA

Taf4, 6, 9

human Transcription
Pre-Initiation Complex (TPIC)

Fig5

A High Confidence Interactor vs ChIP Atlas

High Confidence Interactors

Total KDM5 ChIP Atlas Overlaps

57 3 27 194

Datasets Represented

B KDM5 ChIP Overlapped Datasets

C Sin3A KDM5
D BEAF-32 KDM5
E Dref KDM5
F Iswi KDM5
G wds KDM5
H fs(1)h KDM5

I chr2L

Fig6

Fig7

A SFARI Scores

B gene-score

| Bap55/ACTL6B |
| IntS1/INTS1 |
| Taf4/TAF4 |
| Mi-2/CHD3 |
| kis/CHD7 |
| kis/CHD8 |
| trr/KMT2C |
| nsl1/KANSL1 |
| Kdm5/KDM5B |
| Kdm5/KDM5C |
| CG3815/PHF12 |
| brm/SMARCA4 |
| mor/SMARCC2 |
| Sin3A/SIN3A |
| brm/SMARCA2 |
| CG1815/ZMYND8 |
| fs(1)h/BRD4 |
| dom/EP400 |
| CG15356/EMSY |
| Jarid2/JARID2 |
| Kdm5/KDM5A |
| Sin3A/SIN3B |
| Taf6/TAF6 |
| Ssrp/SSRP1 |
| woc/ZMYM2 |
| kis/CHD9 |

C DBD

Drosophila / Human	ID	Autism	ADHD	Schizophrenia
E(bx)/BPTF		■		
fs(1)h/BRD4	■	■		
Mi-2/CHD3	■	■	■	
kis/CHD6	■	■		
kis/CHD7	■	■		
kis/CHD8	■	■	■	■
CtBP/CTBP1		■		
dom/EP400		■		
simj/GATAD2B	■	■		
IntS1/INTS1	■	■	■	
Jarid2/JARID2	■	■		
nsl1/KANSL1	■	■		
Kdm5/KDM5B	■	■	■	■
Kdm5/KDM5C	■	■	■	■
trr/KMT2C	■	■		■
MED17/MED17	■		■	
CG3815/PHF12	■	■	■	
Sin3A/SIN3A	■	■	■	
brm/SMARCA2	■		■	
brm/SMARCA4	■	■	■	
Snr1/SMARCB1	■	■		
mor/SMARCC1	■	■		
mor/SMARCC2	■	■	■	■
woc/ZMYM2	■	■	■	■

D KDM5 Interactors

Nucleus — Transcription Factors · KDM5 · Insulators · MED1,14,17 · RNA Pol II · KDM5 · Gene · Chromatin Modifiers · Taf4, 6, 9

Supplemental Figure 1

Supplemental Figure 2

Supplemental Figure 3

Proximity labeling reveals a new *in vivo* network of interactors for the histone demethylase KDM5

Matanel Yheskel, Simone Sidoli, and Julie Secombe

Supplemental Figure Legenes

Figure S1: KDM5-TurboID studies using control and TurboID alone

(A) Principal components analysis of normalized nuclear protein abundances from control (Ctl), TurboID, NT-KDM5 and CT-KDM5 (experiment 1 data).

(B) Volcano plot showing data comparing NT-KDM5 to control.

(C) Volcano plot showing data comparing CT-KDM5 to control.

(D) Venn diagram showing overlap of the proteins identified using NT-KDM5 and CT-KDM5 compared to control (Ctl).

(E) Volcano plot showing data comparing NT-KDM5 to TurboID.

(F) Volcano plot showing data comparing CT-KDM5 to TurboID.

(G) Venn diagram showing overlap of the proteins identified using NT-KDM5 and CT-KDM5 compared to TurboID.

(H) Summary of known *Drosophila* KDM5 interactors, whether the interaction is conserved in mammals (mouse and/or human) and the identification of these proteins in experiment 1 with all comparisons. Three interactors identified in mammalian cells but not previously in *Drosophila* are also included (Caf1-55, Mi-2 and Zmynd8).

(I) Volcano plot showing data comparing TurboID to control.

For all volcano plots shown, known interactors are indicated with text. Red dots indicate significantly enriched proteins ($p<0.05$) and the dotted line on Y-axis indicates $p=0.05$.

Figure S2: Comparing TurboID alone to dCas9:TurboID

(A) Volcano plot showing data comparing dCas9:TurboID to TurboID alone.

(B) Volcano plot showing data comparing dCas9:TurboID to control.

(C) GO analyses of proteins enriched comparing TurboID to control (Ctl), TurboID to dCas9:TurboID, dCas9:TurboID to TurboID and dCas9:TurboID to control.

(D) UpSet plot showing the number of common interactors identified in experiments in which TurboID was used to compare datasets using $p<0.05$.

(E) UpSet plot showing the number of common interactors identified in experiments in which TurboID was used to compare datasets using $p<0.1$.

(F) Summary of all known interactors identified in all experimental comparisons from experiment 1 and experiment 2.

For all volcano plots shown, red dots indicate significantly enriched proteins ($p<0.05$) and the dotted line on Y-axis indicates $p=0.05$.

Figure S3: Summary of identification of high confidence interactors in each dataset.

(A) Volcano plot showing data comparing NT-KDM5 to control from experiment 1 with high confidence interactors indicated by red dots.

(B) Volcano plot showing data comparing NT-KDM5 to control from experiment 2 with high confidence interactors indicated by red dots.

(C) Volcano plot showing data comparing NT-KDM5 to dCas9:TurboID (experiment 2) with high confidence interactors indicated by red dots.

(D) Volcano plot showing data comparing CT-KDM5 to control from experiment 1 with high confidence interactors indicated by red dots.

(E) Volcano plot showing data comparing CT-KDM5 to control from experiment 2 with high confidence interactors indicated by red dots.

(F) Volcano plot showing data comparing CT-KDM5 to dCas9:TurboID (experiment 2) with

high confidence interactors indicated by red dots.

For all volcano plots shown, known interactors are indicated with text. Red dots indicate

significantly enriched proteins (p-value<0.05) and the dotted line on Y-axis indicates p=0.05.

Supplemental Figure 1

Supplemental Figure 2

Supplemental Figure 3