No evidence for an effect of chronic boat noise on the fitness of reared water fleas

Loïc Prosnier[*1,2], Emilie Rojas[1] and Vincent Médoc[1]

1. Equipe Neuro-Ethologie Sensorielle, Centre de Recherche en Neurosciences de Lyon, INSERM URMS 1028, CNRS UMR 5292, Université Claude Bernard Lyon 1, Université Jean Monnet - Saint-Etienne, 23 rue du Dr Paul Michelon, 42023 Saint-Etienne Cedex 2, France
2. Pôle emploi, 42000 Saint-Etienne, France.

* Corresponding author. Loïc Prosnier, ENES, Université Jean Monnet - St-Etienne, Campus Métare, Bâtiment K. 21, rue du Dr Paul Michelon 42100 Saint-Etienne, FRANCE. lprosnier@gmail.com ORCID: 0000-0001-5576-3601

Abstract

Among the numerous questions about human impacts on ecosystems, there is a growing interest for acoustic pollution. First studies on underwater acoustic pollution focused, and showed effects, on vertebrates' behaviours. Knowledge on the effects on invertebrates is more limited and there is a huge lack concerning zooplankton species, although widely used as bioindicators in chemical pollution. Consequently, it is critical to assess the impact of noise on zooplankton's fitness (survival and fecundity). Here, isolated water fleas, *Daphnia magna*, were reared from birth to death in the presence or absence of motorboat noises. Effects on lifespan and clonal offspring production (e.g., clutch size, number of offspring produced along life) were assessed and chronic exposure to boat noise did not affect *Daphnia*'s fitness. The spectral and temporal features of the sounds could explain the results. This study highlights the importance of integrating noise pollution into ecotoxicological research to understand, and prevent, human impacts on communities.

Keyword: *Daphnia magna*, Acoustic pollution, Boat noise, Fitness, Survival, Reproduction.

1. Introduction

Freshwater ecosystems are vulnerable to many types of anthropogenic pollution (e.g., chemicals, light, radioactivity, nanopollution, sounds) (see for instance Longcore & Rich, 2004; André et al., 2011; Song et al., 2020; Jan et al., 2022) but the most documented to date remain chemical pollutions (e.g., industrial effluents, urban waste, pesticides, drugs) (Truhaut, 1977; Villeneuve & Garcia-Reyero, 2011). Toxicological studies have documented, in a comprehensive and accurate way, the effects of different types of pollutants (e.g., ion, heavy metals, drugs), exposure durations (acute or chronic), intensities (e.g., median Lethal Dose LD_{50}), and the interactions between them and with environmental parameters (temperature, acidity, humidity, etc.). Those results have contributed to the general knowledge allowing the evaluation of other types of pollution.

This study focuses on acoustic – or noise – pollution described as a pervasive and omnipresent pollutant found in all ecosystems (terrestrial, marine, and freshwater) (Shannon et al., 2016; Popper & Hawkins, 2019; Kunc & Schmidt, 2019), and which represents a growing research topic (Williams et al., 2015; Slabbekoorn, 2019). Specifically, this study focuses on the effect of boat traffic, an important source of noise likely to threaten aquatic systems (Rountree et al., 2020; Duarte et al., 2021). Studies on noise pollution have mainly focused on behaviour (Richardson et al., 1985; Duarte et al., 2021). Contrary to many ecotoxicological studies since Truhaut (1977), it remains a gap in understanding how noise pollution affects individual fitness, i.e. survival and fecundity (Francis & Barber, 2013; Read et al., 2014).

Although the effects of noise on large invertebrates, such as decapods or bivalves, have recently received substantial interest (see the reviews of Popper et al. (2001) and Solé et al. (2023)), research largely neglected zooplanktonic invertebrates (Hawkins et al., 2015; Prosnier, 2022), despite their ecological importance and general use as bioindicators in ecotoxicology

(Parmar et al., 2016). Although zooplankton do not possess hearing structures, they present

mechanoreceptors that allow them to detect particle motion, the other component of a sound

with pressure (Gassie et al., 1993; Buskey et al., 2002). For instance, Gassie et al. (1993) found

that marine copepods (*Acartia fossae*) can detect water vibrations, and Buskey et al. (2002)

showed that vibrations can lead to acceleration in individuals of *Acartia spp.*. Marine

zooplankton (e.g., copepods) exposed to acute airguns show a reduction of survival (McCauley

et al., 2017; Fields et al., 2019; Vereide et al., 2023). Copepods also show reduced foraging rate

during boat noise exposure (Kühn et al., 2023). *Chaborus flavicans* larvae, an important

predator of zooplankton, show more body rotations, interpreted as an anti-predatory behaviour,

when exposed to boat noise for the first time (Rojas et al., 2021). These works highlight that

noise could affect the fitness and behaviour of zooplankton species. However long-term

exposure has not been investigated yet except *D. magna* exposed to chronic broadband noise,

that showed a lower speed and surprisingly a higher fitness (Prosnier, Rojas, et al., 2022).

Temporal pattern is an important acoustic parameter of noise (Francis & Barber, 2013) with

chronic exposure at the one hand and acute exposure at the other (Duarte et al., 2021). Chronic

exposure means a continuous or intermittent, regular or random sound (e.g., turbine, boat noise)

whereas acute exposure represents a punctual sound (e.g., airgun) (Nichols et al., 2015;

McCauley et al., 2017). The spectral (i.e., sound level of each frequency) and temporal patterns

of noise are known to affect the behaviour and physiology of organisms in different ways. For

instance fish are more affected by random noise than by continuous or regular noise (Nichols

et al., 2015). These results are interpreted as an ability for vertebrates to habituate to some

predictable long-term noise exposition (Rojas et al., 2021). Consequently, this raises the

question of whether some results with unrealistic noise, such as continuous broadband noise

(Prosnier, Rojas, et al., 2022), could be extrapolated to real situation where organisms are

exposed to boat noise, thus exposed to intermittent random noise (i.e., unpredictable) with high

spectral variability.

The aim of this study was to investigate the effect of chronic exposure to motorboat noise

(intermittent and random noise with spectral variability) on the fitness of the water flea *Daphnia*

magna, a common zooplankton species widely used in ecotoxicology (Ebert, 2022). Previous

studies found no change in their mobility when exposed to acute noise (Sabet et al., 2015, 2019),

whereas prior experiment with chronic exposure to broadband noise (i.e., a continuous noise)

showed alterations in both fitness and behaviour (Prosnier, Rojas, et al., 2022). If arthropods

react like fish, then intermittent boat noises (i.e., unpredictable) should be more impactful than

continuous broadband noise (i.e., predictable). Consequently, motorboat noise should affect the

fitness of *D. magna*. Absence to negative effects are expected according to various chronic and

acute exposure experiments on invertebrates (Solé et al., 2023), but note that a positive effect

was found for *D. magna* chronically exposed to broadband noise (Prosnier, Rojas, et al., 2022).

Material and Methods

1.1. Collection and maintenance of organisms

Daphnia magna had been purchased from Aqualiment (Grand Est, France) and stored in a

20-L rearing aquarium, filled with aged tap water (physiochemical composition is available on

Zenodo repository (Prosnier et al., 2023a)), for one month. They were reared at 18°C under a

12:12 light:dark cycle. *Daphnia magna* were fed, every two days, with 0.05g of algae (i.e.,

736 mJ) with a mix of 80% of *Arthrospira platensis* and 20% of *Aphanizomenon flos-aquae*

(Algo'nergy® Spiruline + Klamath). Note that *A. platensis* is known to have a C:N ratio

between 4 and 8 (Walach et al., 1987; Griffen & Drake, 2009).

1.2. Fecundity and mortality

101   Reproductive success and survival were measured during an experiment similar as done in

102   Prosnier, Rojas, et al. (2022). Gravid *D. magna* were collected from the rearing aquarium and

103   isolated in 50-mL jars containing Volvic® water. Newborns (< 24h) were transferred

Figure 1. Setup. a) Experimental design. In the four tanks blue circles are loudspeakers, dark small circle are microcosms closed with net. b) Summary of all measures from birth to death of clonal individuals in the two treatments.

individually into 150-mL (5.6 x 8.6 cm) glass microcosm, closed with a 0.3-mm mesh tissue

allowing water flows and noise transmission (Fig. 1).

Eighteen glass microcosms were disposed at 20 cm of an underwater loudspeaker in four

150-L rectangular tanks (75 x 60 x 35 cm), filled with 90 L of aged tap water, at 20-22°C and

under a 12:12 light:dark cycle. Silence was broadcasted in the two control tanks and a daily

boat noise playlist (see below) used as treatment was broadcasted in the two other tanks. For

each *D. magna* mother, half of the newborns were assigned to the control and the other half to

the noise treatment, therefore individuals were clones in the two acoustic conditions. Each day,

survival and newborns production were controlled, and when *D. magna* spawned, offspring

were counted and removed. Every two days, individuals were fed with 2 mL of algae (1g/L) –

note that both Griffen & Drake (2009) and Robinson et al. (2013) used a double quantity of *A.*

platensis, but for populations starting with five to eighteen individuals –, and water was changed

once a week. During the first eight days of the experiment (i.e., before the first hatching), dead

D. magna were replaced by new newborns (isolated mothers were maintained in 50-mL jars

during this initial period to be able to initiate new replicates with newborns) to increase the

number of replicates. Experiments were performed with a total of 115 juveniles (58 control and

57 exposed to noise) coming from 25 mothers; almost half of the juveniles in each condition

(26 in control and 23 in noise) reached maturity. The experiment lasted 46 days, from the birth

of the first individual to the death of the last one (the oldest *D. magna* survived 39 days).

Based on daily survival and daily clutch, populational data were analysed using the Euler-

Lotka equation ($\sum f_x m_x e^{-rx} = 1$), with f_x the fecundity at age x, m_x the survival at age x, and

r the intrinsic rate of increase. This equation allows to calculate the reproductive output R_0

($R_0 = \sum f_x m_x$), the generation time G_t $\left(G_t = \frac{\sum x f_x m_x}{\sum f_x m_x}\right)$, and the intrinsic rate of increase r

$\left(r = \frac{\log R_0}{G_T}\right)$ (Leung et al., 2007; Starke et al., 2021).

      1.3. Acoustic treatments

Daphnia magna were exposed to two acoustic treatments (see Rojas et al. (2021) for more

details): a looped 1-h playlist without sound for the control (i.e., only the ambient noise), or

boat noise for the treatment with a playlist including 15 recordings of motorboat sounds

previously made in the Grangent lake (45°45′07.54″N, 4°25′56.47″E, Loire, France). Their

intensity was modulated from 0 to -25 dB Re 1 µPa by 5 dB to create 75 sounds from 103 to

150 dB RMS Re 1 µPa – a naturally-occurring range of noise levels found in lakes (V. Médoc,

Figure 2. Acoustic treatments. a) 24-h temporal sequence of the broadcasted motorboat noises, from 9 a.m. to 6 p.m. Each vertical line represents a boat. b) Soundscapes of the control microcosms recorded in half microcosms with the records of the 15 boat sounds (solid lines), broadcasted at their maximal intensity in the noise tanks, and silence (dotted line). c) Soundscapes in the noise microcosms recorded in half of the microcosms during the broadcast of the15 boat sounds at their maximal intensity (solid lines) and silence (dotted line). d) Spectra in half of the microcosms. Thick lines are means for control (full blue line) and noise treatment (dashed red line) during the 15 boat broadcasts at their maximal intensity, and during the silence period (dotted black line). Shaded areas delimit the min and max Sound-Pressure Level. e) Sound levels in half of the microcosms. Central bars represent the median, boxes the interquartile range, and dots the outliers (> 1.5 times the interquartile range). Coloured dots are the sound levels in control and noise microcosms during the 15 boat broadcasts at their maximal intensity. The four colours (red, yellow, green, blue) correspond to four noise structure spectra that were visually determined (for instance red and yellow have low energy between 200 and 700 or 1000 Hz compared to green and blue boat noises).

pers. obs.) –, that were duplicated to obtain the total of 150 boat noises used in the experiment.

The boat noise playlist was broadcasted from 9 a.m. to 6 p.m. (Fig. 2a). Both playlists (stereo

WAV files) were generated using the Adobe Audition 2020 software (13.0.0.519, Adobe

Systems Inc., Mountain View, CA, USA) and were played back using a Zoom® H4n recorder

connected to an amplifier (DynaVox® CS-PA 1MK), and an underwater loudspeaker UW30

(Electro Voice®).

To check spectra and noise levels in both control and noise microcosms, recordings were

made (Fig. 2b-e) with a Zoom® H4n coupled to a hydrophone (Aquarian Audio H2A-HLR

Hydrophone, frequency response from 10 Hz to 100 kHz) previously calibrated with a

hydrophone (8104, Brüel & Kjær, Naerum, Denmark; sensitivity –205 dB re. 1 V µPa−1;

frequency response from 0.1 Hz to 180 kHz) connected to a sound level meter (Bruël & Kjaer

2238 Mediator, Naerum, Denmark). Emitted boat noises were firstly corrected, using a one-

third octave graphic equalizer (with Adobe Audition 2020), to make their spectrum closer to

those of the original boat sound (Fig. 2). Boat noises were re-recorded only in half of the

microcosms (in each tank) given that they were qualitatively and quantitatively similar due to

the symmetry of the setup and after controlling with a broadband noise (Prosnier, Rojas, et al.,

2022). Note that the playback of boat noise was perceived in the control microcosms with

intensities around 100 dB RMS Re 1 µPa, which was almost comparable to the sound level

during the silence period (Fig. 2e) allowing to neglect noise transmission between the

treatments. Particle motion cannot be measured due to the absence of adequate equipment,

despite its importance for non-hearing species (Nedelec et al., 2016). However, Olivier et al.

(2023) showed that results can still be qualitatively relevant when based solely on sound

pressure level.

 1.4. Statistical analyses

Statistical analyses were performed using R (version 4.2.2) with a significant threshold at

5%. Data allowed to analyse separately the effects on mortality (death age and adult survival)

and fecundity (age at maturity, clutch frequency, mean clutch size, and daily clutch size). The

combination of mortality and fecundity was used as a proxy of fitness and quantified through

total offspring production. Data were also studied at the population scale using the Euler-Lotka

equation (but without statistical analysis due to absence of populational replicates). A survival

analysis (Log-Rank test) was performed to compare survival (death age and adult duration) and

age at maturity (first clutch age) between the control and noise treatments. For the fecundity

parameters, only individuals that clutched at least once (i.e., that reached maturity) were

considered in the analyses. Clutch frequency (i.e., mean time between two clutches) and mean

clutch size were analysed using linear mixed-effects model, with tanks as random effect, thanks

to the normal distribution of the data checked with a Shapiro test. The effect of both noise and

age on daily clutch size was analysed by a type-II analysis of variance, completed with a

pairwise Wilcoxon test between the treatments within each age. To test the effect of noise on

the total number of clutches and offspring along life a generalized linear mixed-effects models

was used, with tanks as random effect and a log function as the link function for Poisson

distribution.

2. Results

177   Chronic boat noise did not affect (Table A1) the survival of *D. magna* (p-value = 0.51, Fig.

178   3a) with a median survival of 4 days for the control and 5 days for the noise treatment. It did

179   not affect the fecundity parameters, with clutch interval around 2.5 days (p-value = 0.24, Fig.

180   3b), mean clutch size around 10 offspring (p-value = 0.74, Fig. 3c) and age at maturity around

181   8 days (p-value = 0.65). Daily clutch size was not influenced by noise (noise: p-value = 0.89,

Figure 3. Effects of noise treatments on *Daphnia magna* survival and fecundity. a) Survival of *D. magna*; b) mean clutch size; c) clutch frequency; d) Cumulative offspring production along life; and e) total number of offspring during lifetime. Numbers in b) are the numbers of *D. magna* for the two treatments for b, c and e. a) Representation according to the Kaplan-Meier method; b-e) central bars represent the median, boxes the interquartile range, and dots the outliers (> 1.5 times the interquartile range). Statistical analysis: dot $P < 0.1$, * $P < 0.05$; ** $P < 0.01$; NS $P > 0.1$. See Table A1 for statistical values.

noise x age: p-value = 0.35, pairwise: p-values > 0.38; Fig. A1), but changed with age (p-value

= 0.003) with larger clutches at intermediate ages. However, taking into account the whole

lifespan, there was an effect on the total number of clutches (p-value = 0.003) with 5 clutches

for the control and 6 clutches for the noise treatment, and a tendency for higher total offspring

production under boat noise exposure (p-value = 0.099, Fig. 3d,e), with on average 60 to 70

newborns.

The populational analysis done with the Lotka-Euler equation confirmed the tendancy on

total offspring production with a reproductive output (R_0) higher for the noise treatment with

63 offspring compared to the 54 offspring in control. Generation time (G_T) was longer in the

noise treatment, with 16.6 days, compared to the 15.2 days in the control. The combination of

both led to an intrinsic rate of increase of 0.25 day^{-1} in the noise treatment compared to 0.26

day^{-1} in the control.

3. Discussion

This study investigated the effect of exposure to chronic boat noise on the fitness of the water

flea *Daphnia magna*. Contrary to what expected, no effect on the survival and fecundity

parameters was observed, which differs from results obtained with exposure to another type of

chronic noise (Prosnier, Rojas, et al., 2022). Such difference in the results might be due to

variations in the temporal and spectral features of the noise. Moreover, as one of the first

experiments on chronic noise, it raises questions on the interactions with other pollutants, and

on the effects on complex communities.

Chronic boat noise had no effect on *Daphnia magna*'s fecundity and survival. These results

are opposite to those on acute and intense exposition of copepods to airgun. McCauley et al.

(2017) observed high mortality for numerous marine zooplankton species including copepods

following the passage of a boat equipped with an airgun. Fields et al. (2019) showed that airgun

exposure leads to increased mortality in *Calanus finmarchicus* within an hour. Our results are

also opposite to those of Prosnier, Rojas, et al. (2022), where an exposition to a continuous

broadband noise leads to a counter-intuitive increase in *D. magna*'s fitness, with higher survival

and greater clutch size. Here, although boat noise had no statistical effect on the survival and

fecundity parameters, there was a tendency for a higher total offspring production with noise,

which would be consistent with the results of Prosnier, Rojas, et al. (2022). However, at a

population scale, the Euler-Lotka equation suggests a tendency for a lower growth when

exposed to boat noise due to a shorter generational time. Such differences in the magnitude of

the effects suggest that response of zooplankton to chronic noise pollution could depend on the

temporal and spectral structure of noise. Indeed, in Prosnier, Rojas, et al. (2022), the broadband

noise broadcasted was continuous and at high level (130 dB RMS re 1 µPa) for all frequencies

(from 0.1 to 20 kHz), whereas, in this study, sounds with a random temporal pattern (a total of

2h of noise *per* day) and different spectra (some boat noises with low intensity between 200

and 800 Hz) were used at various levels (from 108 to 136 dB RMS re 1 µPa). It is already

known that, in hearing vertebrates (i.e., with dedicated organs to detect sound pressure variation

as inner), animals respond differently to chronic noise pollution depending on temporal

variations (continuous, regular, random), spectral variations (i.e., variation in the frequencies),

and if noise is predictable or not (Francis & Barber, 2013). Nichols et al. (2015) showed that

fish were more stressed (higher cortisol concentration) with a higher noise level (from 126 to

141 dB RMS re 1 µPa), and under intermittent random noise (i.e., unpredictable) compared to

continuous noise. However, the review of de Jong et al. (2020) on noise effects on fecundity

revealed that continuous noise with spectral variations, such as boat noise, was more prone to

impact physiological markers (cortisol level, ventilation rates and metabolic rate) and behaviour

(startle and freeze responses, horizontal and vertical avoidances). Another study on zebrafish

larvae showed a negative effect of a continuous white noise on survival and a higher cortisol

concentration (Lara & Vasconcelos, 2021). Thus, it seems that for hearing vertebrates,

depending on the developmental stage and the considered characteristic (fitness, behavior, or

physiology), both temporal and spectral characteristics need to be considered. For zooplankton,

little is known about the importance of noise characteristics. Here, the comparison with Lara &

Vasconcelos (2021) and Prosnier, Rojas, et al. (2022) suggests that negative effects are higher

with continuous noise, contrary what is reported with fishes.

Despite there was no effect on fecundity, it would be necessary to focus on the offspring

coming from mothers exposed to noise. Here, there was no qualitative effect reported during

the current experiments, i.e., all the water fleas produced seemed viable and mobile. More, there

was no increase in the mortality of newborns due to noise, and no effect was reported on the

size of *D. magna* exposed to chronic noise (Prosnier, Rojas, et al., 2022). This would be

consistent with the study of Day et al. (2016) where exposure to air gun did not affect the

embryonic development of the spiny lobster *Jasus edwardsii* (Decapodae). However, airgun

exposition reduces growth and development stage of *Acartia tonsa* nauplii (Vereide et al.,

2023). This aspect seems important as it is known that stress on mother and early stages can

affect daphnia's development (Mittmann et al., 2014; Mushegian et al., 2016) and that effects

can differ across generations (Campos et al., 2016). Consequently, impact studies on noise

should focus on embryonic development and perform multigenerational experiments to

determine the long-term effects of chronic exposure resulting from embryonic misdevelopment

(Mushegian et al., 2016), maternal effects (Radersma et al., 2018), and acclimatation or

adaptation (Ringot et al., 2018; Abdullahi et al., 2022).

An interesting question is to consider the effect of noise as part of cocktails of pollutants. It

is now a common question in ecotoxicology to ask whether stressors (e.g., chemical pollution,

temperature, food quality) act synergistically (Altshuler et al., 2011). For instance, Starke et al.

(2021) showed that food quality impacts more *D. pulicaria* at some higher temperature due to

the increased metabolism. Prosnier et al. (2015) modelled the antagonistic effect of copper and

nutrient enrichment on the *Daphnia* - algae interaction. Regarding noise, Stenton et al. (2022)

showed both synergetic and antagonistic effects of noise and cadmium exposure on Norway

lobster larvae (*Nephrops norvegicus*), depending of the considered parameters (survival,

development, oxidative indicators). Another, McMahon et al. (2017) studied the interactive

effects of light and noise pollutions on a frog-parasite interaction. They showed that light

reduced frog-biting midge (*Corethrella* spp.) abundance at low noise level, whereas there was

no midge at high noise level. It could be also useful to investigate whether all the unwanted

noises produced by many experimental setups (to control light, temperature, oxygenation, and

food) interact with the stressors studied and influence the results. For instance, in the present

study there was a very high mortality in *D. magna* juveniles compared to similar studies (Parisot

et al., 2015; Prosnier, Loeuille, et al., 2022). It suggests suboptimal conditions (i.e., other

stressor than noise), such as a lack of food – for instance Serra et al. (2020) fed *D. magna* daily

for the seven first day –, which may have affected the outcomes through the masking of effects

for example. However, on the other side, suboptimal conditions could make individuals more

prone to be affected by an additional stress like noise. Note that, with the same suboptimal

conditions, Prosnier, Rojas, et al. (2022) obtained a significant difference between the control

and noise treatments. The recent *Larvosonic* system, developed by Olivier et al. (2023), could

be useful to study the impact of noise on zooplankton with a better control of the environmental

conditions.

This study is a first step in our understanding of the importance of noise patterns for

invertebrates (i.e., predictable versus unpredictable noise) in comparison with vertebrates. To

go further, we need more information about noise perception (i.e., mechanoreception and

involved gene) and sensory integration, that could explain the mentioned differences between

vertebrates and invertebrates and seem largely unexplored (Gassie et al., 1993; Popper et al.,

2001). Understanding the various reactions of vertebrates and invertebrates in terms of

behavior, but also in terms of fitness is mandatory to study how noise could affect complex

communities (Francis et al., 2009; Slabbekoorn & Halfwerk, 2009; Slabbekoorn, 2019). For

instance, in a freshwater community, unpredictable noise should affect more fishes, at top trophic levels, than invertebrates. But if there are various effects within zooplankton community, as expected, leading to bottom-up effects. Zooplankton is highly diverse and predatory species might react differently than their zooplanktonic prey. Moreover, in a community, pollutants can alter fitness directly (as in this study) but also indirectly through change in the vulnerability to natural enemies (Read et al., 2014). For instance, noise does not affect frog abundance but reduces that of their parasite (McMahon et al., 2017). The need for more research on invertebrates and fitness impacts, particularly in arthropods, is also true for terrestrial communities (Morley et al., 2014). Thus, a more general overview on the response of invertebrates to anthropogenic noises should be beneficial to mitigate the impacts (Francis & Barber, 2013).

Acknowledgments

The authors would like to thank all the people who contributed to the success of this experiment: Nicolas Boyer and Aurélie Pradeau for *Daphnia* rearing and providing material, Léo Papet, Joël Attia, and Jérémy Rouch for the acoustic treatments and analysis and for providing material, Olivier Valéro for sound recording, Marilyn Beauchaud and Paolo Fonseca for acoustic calibration, Théophile Turco for help in the analyses, and the EYD (ENES Young Discussion) for the useful discussions. Authors also thanks Marie-Agnès Coutellec and the anonymous reviewer for their useful comments.

Note that this article was published as a chapter of "The effects of noise on aquatic life" (Popper et al., 2023): https://doi.org/10.1007/978-3-031-10417-6_129-1 (Prosnier et al., 2023b)

Funding

The authors declare they had no funding for this research and were financially supported by their laboratory.

Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article.

Data, script and code availability

Data, script and code are available on Zenodo. DOI: 10.5281/zenodo.7775919 (Prosnier et al., 2023a)

Supplementary information

Supplementary information is available after the references:

- Appendix: Tables of statistics and supplementary figure

References

Abdullahi M, Zhou J, Dandhapani V, Chaturvedi A, Orsini L (2022) Historical exposure to chemicals reduces tolerance to novel chemical stress in Daphnia (waterflea). *Molecular Ecology*, **31**, 3098–3111. https://doi.org/10.1111/mec.16451

Altshuler I, Demiri B, Xu S, Constantin A, Yan ND, Cristescu ME (2011) An integrated multi-disciplinary approach for studying multiple stressors in freshwater ecosystems: Daphnia as a model organism. *Integrative and Comparative Biology*, **51**, 623–633. https://doi.org/10.1093/icb/icr103

André M, Solé M, Lenoir M, Durfort M, Quero C, Mas A, Lombarte A, van der Schaar M, López-Bejar M, Morell M, Zaugg S, Houégnigan L (2011) Low-frequency sounds induce acoustic trauma in cephalopods. *Frontiers in Ecology and the Environment*, **9**, 489–493. https://doi.org/10.1890/100124

Buskey EJ, Lenz PH, Hartline DK (2002) Escape behavior of planktonic copepods in response to hydrodynamic disturbances: high speed video analysis. *Marine Ecology Progress Series*, **235**, 135–146. https://doi.org/10.3354/meps235135

Campos B, Jordão R, Rivetti C, Lemos MFL, Soares AMVM, Tauler R, Barata C (2016) Two-generational effects of contaminants in Daphnia magna: Effects of offspring quality. *Environmental Toxicology and Chemistry*, **35**, 1470–1477. https://doi.org/10.1002/etc.3290

Day RD, McCauley RD, Fitzgibbon QP, Semmens JM (2016) Seismic air gun exposure during early-stage embryonic development does not negatively affect spiny lobster Jasus edwardsii larvae

(Decapoda:Palinuridae). *Scientific Reports*, **6**, 22723. https://doi.org/10.1038/srep22723

Duarte CM, Chapuis L, Collin SP, Costa DP, Devassy RP, Eguiluz VM, Erbe C, Gordon TAC, Halpern BS,

Harding HR, Havlik MN, Meekan M, Merchant ND, Miksis-Olds JL, Parsons M, Predragovic M, Radford

AN, Radford CA, Simpson SD, Slabbekoorn H, Staaterman E, Van Opzeeland IC, Winderen J, Zhang X,

Juanes F (2021) The soundscape of the Anthropocene ocean. *Science*, **371**, eaba4658.

https://doi.org/10.1126/science.aba4658

Ebert D (2022) Daphnia as a versatile model system in ecology and evolution. *EvoDevo*, **13**, 16.

https://doi.org/10.1186/s13227-022-00199-0

Fields DM, Handegard NO, Dalen J, Eichner C, Malde K, Karlsen Ø, Skiftesvik AB, Durif CMF, Browman HI

(2019) Airgun blasts used in marine seismic surveys have limited effects on mortality, and no sublethal

effects on behaviour or gene expression, in the copepod Calanus finmarchicus (P Ratilal, Ed,). *ICES*

Journal of Marine Science, **76**, 2033–2044. https://doi.org/10.1093/icesjms/fsz126

Francis CD, Barber JR (2013) A framework for understanding noise impacts on wildlife: an urgent conservation

priority. *Frontiers in Ecology and the Environment*, **11**, 305–313. https://doi.org/10.1890/120183

Francis CD, Ortega CP, Cruz A (2009) Noise pollution changes avian communities and species interactions.

Current Biology, **19**, 1415–1419. https://doi.org/10.1016/j.cub.2009.06.052

Gassie D V., Lenz PH, Jeannette Y, Hartline DK (1993) Mechanoreception in zooplankton first antennae:

electrophysiological techniques. *Bulletin of Marine Science*, **53**, 96–105.

Griffen BD, Drake JM (2009) Environment, but not migration rate, influences extinction risk in experimental

metapopulations. *Proceedings of the Royal Society B: Biological Sciences*, **276**, 4363–4371.

https://doi.org/10.1098/rspb.2009.1153

Hawkins AD, Pembroke AE, Popper AN (2015) Information gaps in understanding the effects of noise on fishes

and invertebrates. *Reviews in Fish Biology and Fisheries*, **25**, 39–64. https://doi.org/10.1007/s11160-014-

9369-3

Jan N, Majeed N, Ahmad M, Ahmad Lone W, John R (2022) Nano-pollution: Why it should worry us.

Chemosphere, **302**, 134746. https://doi.org/10.1016/j.chemosphere.2022.134746

de Jong K, Forland TN, Amorim MCP, Rieucau G, Slabbekoorn H, Sivle LD (2020) Predicting the effects of

anthropogenic noise on fish reproduction. *Reviews in Fish Biology and Fisheries*, **30**, 245–268.

https://doi.org/10.1007/s11160-020-09598-9

Kühn S, King F, Heubel KU (2023) Decreased feeding rates of the copepod Acartia tonsa when exposed to

playback harbor traffic noise. *Frontiers in Marine Science*, **10**, 1134792.

https://doi.org/10.3389/fmars.2023.1134792

Kunc HP, Schmidt R (2019) The effects of anthropogenic noise on animals: a meta-analysis. *Biology Letters*, **15**,

20190649. https://doi.org/10.1098/rsbl.2019.0649

Lara RA, Vasconcelos RO (2021) Impact of noise on development, physiological stress and behavioural patterns

in larval zebrafish. *Scientific Reports*, **11**, 6615. https://doi.org/10.1038/s41598-021-85296-1

Leung KMY, Grist EPM, Morley NJ, Morritt D, Crane M (2007) Chronic toxicity of tributyltin to development

and reproduction of the European freshwater snail Lymnaea stagnalis (L.). *Chemosphere*, **66**, 1358–1366.

https://doi.org/10.1016/j.chemosphere.2006.06.051

Longcore T, Rich C (2004) Ecological light pollution. *Frontiers in Ecology and the Environment*, **2**, 191–198.

https://doi.org/10.1890/1540-9295(2004)002[0191:ELP]2.0.CO;2

McCauley RD, Day RD, Swadling KM, Fitzgibbon QP, Watson RA, Semmens JM (2017) Widely used marine

seismic survey air gun operations negatively impact zooplankton. *Nature Ecology & Evolution*, **1**, 0195.

https://doi.org/10.1038/s41559-017-0195

McMahon TA, Rohr JR, Bernal XE (2017) Light and noise pollution interact to disrupt interspecific interactions.

Ecology, **98**, 1290–1299. https://doi.org/10.1002/ecy.1770

Mittmann B, Ungerer P, Klann M, Stollewerk A, Wolff C (2014) Development and staging of the water flea

Daphnia magna (Straus, 1820; Cladocera, Daphniidae) based on morphological landmarks. *EvoDevo*, **5**,

12. https://doi.org/10.1186/2041-9139-5-12

Morley EL, Jones G, Radford AN (2014) The importance of invertebrates when considering the impacts of

anthropogenic noise. *Proceedings of the Royal Society B: Biological Sciences*, **281**, 20132683.

https://doi.org/10.1098/rspb.2013.2683

Mushegian AA, Burcklen E, Schär TMM, Ebert D (2016) Temperature-dependent benefits of bacterial exposure

in embryonic development of Daphnia magna resting eggs. *Journal of Experimental Biology*, **219**, 897–

904. https://doi.org/10.1242/jeb.134759

Nedelec SL, Campbell J, Radford AN, Simpson SD, Merchant ND (2016) Particle motion: the missing link in

underwater acoustic ecology (D Fisher, Ed,). *Methods in Ecology and Evolution*, **7**, 836–842.

https://doi.org/10.1111/2041-210X.12544

Nichols TA, Anderson TW, Širović A (2015) Intermittent noise induces physiological stress in a coastal marine

fish (CA Radford, Ed,). *PLoS ONE*, **10**, e0139157. https://doi.org/10.1371/journal.pone.0139157

Olivier F, Gigot M, Mathias D, Jezequel Y, Meziane T, L'Her C, Chauvaud L, Bonnel J (2023) Assessing the

impacts of anthropogenic sounds on early stages of benthic invertebrates: The "Larvosonic system."

Limnology and Oceanography: Methods, **21**, 53–68. https://doi.org/10.1002/lom3.10527

Parisot F, Bourdineaud J-P, Plaire D, Adam-Guillermin C, Alonzo F (2015) DNA alterations and effects on

growth and reproduction in Daphnia magna during chronic exposure to gamma radiation over three

successive generations. *Aquatic Toxicology*, **163**, 27–36. https://doi.org/10.1016/j.aquatox.2015.03.002

Parmar TK, Rawtani D, Agrawal YK (2016) Bioindicators: the natural indicator of environmental pollution.

Frontiers in Life Science, **9**, 110–118. https://doi.org/10.1080/21553769.2016.1162753

Popper AN, Hawkins AD (2019) An overview of fish bioacoustics and the impacts of anthropogenic sounds on

fishes. *Journal of Fish Biology*, **94**, 692–713. https://doi.org/10.1111/jfb.13948

Popper AN, Salmon M, Horch KW (2001) Acoustic detection and communication by decapod crustaceans.

Journal of Comparative Physiology A: Sensory, Neural, and Behavioral Physiology, **187**, 83–89.

https://doi.org/10.1007/s003590100184

Popper AN, Sisneros J, Hawkins AD, Thomsen F (Eds.) (2023) *The Effects of Noise on Aquatic Life: Principles*

and Practical Considerations. Springer International Publishing, Cham. https://doi.org/10.1007/978-3-

031-10417-6

Prosnier L (2022) How the noises could affect marine food webs ? A lack of zooplankton' s studies.

EcoEvoRxiv, X2NK58. https://doi.org/10.32942/X2NK58

Prosnier L, Loeuille N, Hulot FD, Renault D, Piscart C, Bicocchi B, Deparis M, Lam M, Médoc V (2022)

Parasites make hosts more profitable but less available to predators. *bioRxiv*, 2022.02.08.479552.

https://doi.org/10.1101/2022.02.08.479552

Prosnier L, Loreau M, Hulot FD (2015) Modeling the direct and indirect effects of copper on phytoplankton-

zooplankton interactions. *Aquatic Toxicology*, **162**, 73–81. https://doi.org/10.1016/j.aquatox.2015.03.003

Prosnier L, Rojas E, Médoc V (2023a) Datasets and R source code of manuscript "Chronic boat noise does not

alter the fitness of Daphnia magna." *Dataset on Zenodo*. https://doi.org/10.5281/zenodo.7775919

Prosnier L, Rojas E, Médoc V (2023b) Chronic boat noise does not alter the fitness of Daphnia magna. In: *The*

Effects of Noise on Aquatic Life: Principles and Practical Considerations (eds Popper AN, Sisneros J,

Hawkins AD, Thomsen F), pp. 1–15. Springer International Publishing, Cham.

https://doi.org/10.1007/978-3-031-10417-6_129-1

Prosnier L, Rojas E, Valéro O, Médoc V (2022) Chronic noise unexpectedly increases fitness of a freshwater

zooplankton. *bioRxiv*, 2022.11.19.517212. https://doi.org/10.1101/2022.11.19.517212

Radersma R, Hegg A, Noble DWA, Uller T (2018) Timing of maternal exposure to toxic cyanobacteria and

offspring fitness in Daphnia magna: Implications for the evolution of anticipatory maternal effects.

Ecology and Evolution, **8**, 12727–12736. https://doi.org/10.1002/ece3.4700

Read J, Jones G, Radford AN (2014) Fitness costs as well as benefits are important when considering responses

to anthropogenic noise. *Behavioral Ecology*, **25**, 4–7. https://doi.org/10.1093/beheco/art102

Richardson WJ, Fraker MA, Würsig B, Wells RS (1985) Behaviour of Bowhead Whales Balaena mysticetus

summering in the Beaufort Sea: Reactions to industrial activities. *Biological Conservation*, **32**, 195–230.

https://doi.org/10.1016/0006-3207(85)90111-9

Ringot G, Gasparini J, Wagner M, Cheikh Albassatneh M, Frantz A (2018) More and smaller resting eggs along

a gradient for pollution by metals: dispersal, dormancy and detoxification strategies in Daphnia?

Biological Journal of the Linnean Society, **124**, 11–20. https://doi.org/10.1093/biolinnean/bly026

Robinson JD, Wares JP, Drake JM (2013) Extinction hazards in experimental Daphnia magna populations:

effects of genotype diversity and environmental variation. *Ecology and Evolution*, **3**, 233–243.

https://doi.org/10.1002/ece3.449

Rojas E, Thévenin S, Montes G, Boyer N, Médoc V (2021) From distraction to habituation: Ecological and

behavioural responses of invasive fish to anthropogenic noise. *Freshwater Biology*, **66**, 1606–1618.

https://doi.org/10.1111/fwb.13778

Rountree RA, Juanes F, Bolgan M (2020) Temperate freshwater soundscapes: A cacophony of undescribed

biological sounds now threatened by anthropogenic noise (DM Higgs, Ed,). *PLoS ONE*, **15**, e0221842.

https://doi.org/10.1371/journal.pone.0221842

Sabet SS, Karnagh SA, Azbari FZ (2019) Experimental test of sound and light exposure on water flea swimming

behaviour. *Proceedings of Meetings on Acoustics*, **37**, 010015. https://doi.org/10.1121/2.0001270

Sabet SS, Neo YY, Slabbekoorn H (2015) The effect of temporal variation in sound exposure on swimming and

foraging behaviour of captive zebrafish. *Animal Behaviour*, **107**, 49–60.

https://doi.org/10.1016/j.anbehav.2015.05.022

Serra T, Barcelona A, Pous N, Salvadó V, Colomer J (2020) Synergistic effects of water temperature,

microplastics and ammonium as second and third order stressors on Daphnia magna. *Environmental*

Pollution, **267**, 115439. https://doi.org/10.1016/j.envpol.2020.115439

Shannon G, McKenna MF, Angeloni LM, Crooks KR, Fristrup KM, Brown E, Warner KA, Nelson MD, White

C, Briggs J, McFarland S, Wittemyer G (2016) A synthesis of two decades of research documenting the

effects of noise on wildlife. *Biological Reviews*, **91**, 982–1005. https://doi.org/10.1111/brv.12207

Slabbekoorn H (2019) Noise pollution. *Current Biology*, **29**, R957–R960.

https://doi.org/10.1016/j.cub.2019.07.018

Slabbekoorn H, Halfwerk W (2009) Behavioural ecology: noise annoys at community level. *Current Biology*,

19, R693–R695. https://doi.org/10.1016/j.cub.2009.07.002

Solé M, Kaifu K, Mooney TA, Nedelec SL, Olivier F, Radford AN, Vazzana M, Wale MA, Semmens JM,

Simpson SD, Buscaino G, Hawkins A, Aguilar de Soto N, Akamatsu T, Chauvaud L, Day RD, Fitzgibbon

Q, McCauley RD, André M (2023) Marine invertebrates and noise. *Frontiers in Marine Science*, **10**,

1129057. https://doi.org/10.3389/fmars.2023.1129057

Song Y, Xie L, Lee Y, Brede DA, Lyne F, Kassaye Y, Thaulow J, Caldwell G, Salbu B, Tollefsen KE (2020)

Integrative assessment of low-dose gamma radiation effects on Daphnia magna reproduction: Toxicity

pathway assembly and AOP development. *Science of The Total Environment*, **705**, 135912.

https://doi.org/10.1016/j.scitotenv.2019.135912

Starke CWE, Jones CLC, Burr WS, Frost PC (2021) Interactive effects of water temperature and stoichiometric

food quality on Daphnia pulicaria. *Freshwater Biology*, **66**, 256–265. https://doi.org/10.1111/fwb.13633

Stenton CA, Bolger EL, Michenot M, Dodd JA, Wale MA, Briers RA, Hartl MGJ, Diele K (2022) Effects of pile

driving sound playbacks and cadmium co-exposure on the early life stage development of the Norway

lobster, Nephrops norvegicus. *Marine Pollution Bulletin*, **179**, 113667.

https://doi.org/10.1016/j.marpolbul.2022.113667

Truhaut R (1977) Ecotoxicology: Objectives, Principles and Perspectives. *Ecotoxicology and Environmental*

Safety, **1**, 373–413. https://doi.org/10.1016/B978-0-08-021998-1.50038-5

Vereide EH, Mihaljevic M, Browman HI, Fields DM, Agersted MD, Titelman J, de Jong K (2023) Effects of

airgun discharges used in seismic surveys on development and mortality in nauplii of the copepod Acartia

tonsa. *Environmental Pollution*, 121469. https://doi.org/10.1016/j.envpol.2023.121469

Villeneuve DL, Garcia-Reyero N (2011) Vision & strategy: Predictive ecotoxicology in the 21st century.

Environmental Toxicology and Chemistry, **30**, 1–8. https://doi.org/10.1002/etc.396

Walach MR, Bazin MJ, Pirt SJ, Balyuzi HHM (1987) Computer control of carbon-nitrogen ratio inSpirulina

platensis. *Biotechnology and Bioengineering*, **29**, 520–528. https://doi.org/10.1002/bit.260290417

Williams R, Wright AJ, Ashe E, Blight LK, Bruintjes R, Canessa R, Clark CW, Cullis-Suzuki S, Dakin DT,

Erbe C, Hammond PS, Merchant ND, O'Hara PD, Purser J, Radford AN, Simpson SD, Thomas L, Wale

MA (2015) Impacts of anthropogenic noise on marine life: Publication patterns, new discoveries, and

future directions in research and management. *Ocean & Coastal Management*, **115**, 17–24.

https://doi.org/10.1016/j.ocecoaman.2015.05.021

Appendix: Tables of statistics and supplementary figure

Table A1. Statistical results of chronic boat noises effects on fecundity and mortality of *Daphnia magna* (Fig. 3)

	Mortality		Reproduction			Fitness	
	Survival	Adult duration	Age at maturity	Clutch interval	Mean clutch size	Number of clutches	Number of offspring
df	1	1	1	1	1	1	1
χ²/W	0.4	1.6	0.2	1.4029	-0.54276	3.8197	1.8263
p-value	0.51	0.21	0.65	0.236	0.737	**0.031**	0.099
R2	0.0041	0.03	0.0046	0.03	0.007	0.096	0.327

Figure A1. Effects of noise treatments on *Daphnia magna* daily clutch size. Dots represent the median of the daily clutch size and lines are interquartile ranges. See Fig. 3a,d as complement for the number of individuals at each age. Note that there could be only one clutch for an age (i.e., no interquartile lines) or clutches only for one treatment (i.e., only one point for an age).