A human neural crest model reveals the developmental impact of neuroblastoma-associated chromosomal aberrations

Ingrid M. Saldana-Guerrero[1,2,3,*], Luis F. Montano-Gutierrez[4,*], Katy Boswell[1,2,$], Christoph Hafemeister[4,$], Evon Poon[5,$], Lisa E. Shaw[6,$], Dylan Stavish[1,2,$], Rebecca A. Lea[1], Sara Wernig-Zorc[4], Eva Bozsaky[4], Irfete S. Fetahu[4], Peter Zoescher[4], Ulrike Pötschger[4], Marie Bernkopf[4,7], Andrea Wenninger-Weinzierl[4], Caterina Sturtzel[4], Celine Souilhol[1,2,8], Sophia Tarelli[1,2], Mohamed R. Shoeb[4], Polyxeni Bozatzi[4], Magdalena Rados[4], Maria Guarini[9], Michelle C. Buri[4], Wolfgang Weninger[6], Eva M. Putz[4], Miller Huang[10,11], Ruth Ladenstein[4], Peter W. Andrews[1], Ivana Barbaric[1,2], George D. Cresswell[4], Helen E. Bryant[3], Martin Distel[4], Louis Chesler[5,+], Sabine Taschner-Mandl[4,+], Matthias Farlik[6,+], Anestis Tsakiridis[1,2,#], Florian Halbritter[4,#]

[1] Centre for Stem Cell Biology, School of Biosciences, The University of Sheffield, Sheffield, UK
[2] Neuroscience Institute, The University of Sheffield, Sheffield, UK
[3] Sheffield Institute for Nucleic Acids (SInFoNiA), School of Medicine and Population Health, The University of Sheffield, Sheffield, UK
[4] St. Anna Children's Cancer Research Institute (CCRI), Vienna, Austria
[5] Division of Clinical Studies, The Institute of Cancer Research (ICR) & Royal Marsden NHS Trust, London, UK
[6] Medical University of Vienna, Department of Dermatology, Vienna, Austria
[7] Labdia Labordiagnostik GmbH, Vienna, Austria
[8] Biomolecular Sciences Research Centre, Department of Biosciences and Chemistry, Sheffield Hallam University, Sheffield, UK
[9] CeMM Research Center for Molecular Medicine of the Austrian Academy of Science, Vienna, Austria
[10] Children's Hospital Los Angeles, Cancer and Blood Disease Institutes, and The Saban Research Institute, Los Angeles, CA, USA
[11] Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

*,$,+ These authors contributed equally to this work.
# Equally contributing senior authors.

Correspondence to a.tsakiridis@sheffield.ac.uk (A.T.) or florian.halbritter@ccri.at (F.H.)

Abstract

Early childhood tumours arise from transformed embryonic cells, which often carry large copy number alterations (CNA). However, it remains unclear how CNAs contribute to embryonic tumourigenesis due to a lack of suitable models. Here we employ female human embryonic stem cell (hESC) differentiation and single-cell transcriptome and epigenome analysis to assess the effects of chromosome 17q/1q gains, which are prevalent in the embryonal tumour neuroblastoma (NB). We show that CNAs impair the specification of trunk neural crest (NC) cells and their sympathoadrenal derivatives, the putative cells-of-origin of NB. This effect is exacerbated upon overexpression of *MYCN*, whose amplification co-occurs with CNAs in NB. Moreover, CNAs potentiate the pro-tumourigenic effects of *MYCN* and mutant NC cells resemble NB cells in tumours. These changes correlate with a stepwise aberration of developmental transcription factor networks. Together, our results sketch a mechanistic framework for the CNA-driven initiation of embryonal tumours.

Keywords: neuroblastoma, copy number alterations, tumourigenesis, stem cells, neural crest

Cancers in early childhood are driven by sparse genetic aberrations arising *in utero*, which are thought to lead to defective differentiation and uncontrolled proliferation[1–4]. Most tumours harbour large genomic rearrangements and chromosomal copy number alterations (CNA), which co-occur with mutations in tumour suppressors or tumourigenic transcription factors (TF)[5,6]. The mechanistic interactions between different mutations and early developmental processes are likely foundational drivers of tumour heterogeneity. However, since visible tumours are only detected long after their initiation, early mutation-driven interactions leading to the healthy-to-tumour transition have remained largely intractable.

Neuroblastoma (NB) is the most common extra-cranial solid tumour in infants and an archetypal "developmental cancer"[7–9]. NB tumours are usually found in the adrenal gland or sympathetic ganglia, tissues derived from the trunk neural crest (NC) lineage during embryonic development[10,11], and studies using transgenic animal models and transcriptome analysis have anchored NB tumourigenesis in impaired sympathoadrenal differentiation of trunk NC cells[12–23]. CNAs such as gains of the long arms of chromosomes 17 (chr17q) and 1 (chr1q) have been identified in the majority (up to 65%) of NB tumours[24–28] and their emergence is considered an early tumourigenesis "priming" event[29]. Chr17q/1q gains often co-occur with amplification of the *MYCN* oncogene[24,28,30–33] (at least one CNA in >95% of *MYCN*-amplified tumours[34]), suggesting they may jointly contribute to tumourigenesis. However, despite our advanced understanding of the genetic and developmental origin of NB, it remains unclear to date how CNAs disrupt embryonic cell differentiation and lead to NB initiation.

Here, we used a human embryonic stem cell (hESC)-based model to experimentally dissect the links between NB-associated CNAs, *MYCN* amplification, and tumour initiation. We interrogated the stepwise specification of trunk NC and sympathoadrenal lineages using directed differentiation of isogenic hESC lines with chr17q/1q gains and inducible *MYCN* overexpression. We found that (i) CNAs derail differentiation by potentiating immature NC progenitor phenotypes. Combining CNAs with *MYCN* overexpression completely disrupted normal NC differentiation; (ii) Mutant NC cells acquired tumourigenic hallmarks *in vitro*, the capacity to form tumours in xenografts, and resemble distinct subpopulations of heterogeneous NB tumours; (iii) An extensive re-wiring of chromatin connects the observed transcriptional and functional aberrations with a dysregulated network of developmental TFs. Collectively, our data put forward a CNA-driven distortion of trunk NC and sympathoadrenal differentiation as a priming mechanism for subsequent MYCN-induced tumour initiation.

Results

Differentiation of human embryonic stem cells recapitulates key stages of trunk NC and sympathoadrenal development

To model the initiation stage and cell types relevant to NB tumourigenesis, we turned to an *in vitro* modelling approach. We have previously described an efficient strategy to produce human trunk NC, sympathoadrenal progenitors, and sympathetic neurons from hESCs[35,36]. Our protocol involves treatment with defined cocktails of signalling pathway agonists/antagonists that induce neuromesodermal-potent axial progenitors (NMPs) at day 3 of differentiation (D3)[37], and subsequently steer NMPs toward trunk NC cells (D9) and their sympathoadrenal derivatives (>D14). At D19, the protocol yields catecholamine-producing sympathetic neurons marked by peripherin-expressing axons[35] (**Fig. 1a, Supplementary Fig. 1**).

As a prerequisite for studying the effects of CNAs on trunk NC differentiation, we first needed to define a molecular roadmap of normal hESC differentiation as a control. Therefore, we employed our protocol for the differentiation of karyotypically normal hESCs (H7[38]; 46XX) and performed droplet-based single-cell RNA sequencing (scRNA-seq) at key differentiation stages (D0 ≈ hESCs, D3 ≈ NMPs, D9 ≈ trunk NC, D14 ≈ sympathoadrenal progenitors, D19 ≈ early sympathetic neuron) and intermediate/late time points (D4, D10, D12, D28) to examine the resulting cell populations (up to five replicates each; **Supplementary Table 1**). We obtained 29,857 cells that passed quality control, which we allocated to 14 distinct clusters (C1-C14) (**Fig. 1b; Supplementary Figs. 2a-g**). We bioinformatically annotated these cell clusters using two complementary approaches: (i) by identifying characteristic marker genes (**Fig. 1c; Supplementary Fig. 2h; Supplementary Table 2**), and (ii) by mapping our data to single-cell transcriptomes of trunk NC derivatives in human embryos[15,16] (**Figs. 1d-f, Supplementary Figs. 2i,j**). This strategy identified cells at different stages of trunk NC development, including NMP-like cells (marked by CDX1/2, NKX1-2, and FGF signalling-associated transcripts[37]; cluster C2 in **Fig. 1c, Supplementary Table 2**) and later cell populations of a predominantly trunk axial identity (**Supplementary Fig. 2h**) exhibiting characteristics of Schwann cell precursors (SCP), sympathoblasts (SYM), as well as mesenchymal features (MES). For example, D9 cells split into subpopulations expressing markers of trunk NC/early SCPs (C3; e.g., *SOX10*[16], **Fig. 1c**; weak SCP-like signature, **Fig. 1e**) and sensory neurons (C5; *ONECUT1*[39], **Fig. 1c**; weak SYM-like signature, **Fig. 1f**). At D14, cells started to assume a sympathoadrenal/autonomic progenitor (C8; *ASCL1*) or mesenchymal (C11; *FN1*) identity, and by day D19, we observed three distinct fractions: mature SCP-like cells (C9; *POSTN*[40]; strong SCP signature), autonomic sympathoblasts (C12-C14; *PHOX2A/B*, *ELAVL4*[16,41]; strong SYM signature), and MES-like cells (C11; *COL1A1, FN1*). This is in line with findings showing that trunk NC and SCPs are competent to generate mesenchyme[40,42,43]. Interestingly, we also found cells at the intersection of MES and SYM identity, as observed in mice[39] and NB cell lines[44–48] (**Supplementary Fig. 3; Supplementary Table 3**). After 4 weeks (D28), we also

observed some cells with a partial chromaffin-like cell identity (part of C14; *CHGA*⁺, PMNT⁻) (**Fig.**

1d).

Together, these data confirm that our hESC-based model successfully captures trunk NC and

sympathoadrenal cells as found in embryos during the onset of NB tumourigenesis. Moreover, they

reveal two major developmental branching events: (i) an early commitment of trunk NC toward a

sensory neuron fate; (ii) the late generation of multipotent SCP/sympathoadrenal progenitors, which

subsequently give rise to three distinct cell types: mature SCPs, MES, and SYM.

CNAs and *MYCN* cumulatively disrupt human trunk NC differentiation

Having established a reliable model of trunk NC lineages relevant for NB pathogenesis, we next asked

how chr17q and chr1q gains and their interplay with *MYCN* overexpression, which often co-occurs with

CNAs in NB[24,28,30–34], influence NC development. To this end, we employed two clonal isogenic hESC

lines with NB-associated CNAs that were acquired by H7 hESCs ('WT'; used in **Fig. 1**) as a result of

culture adaptation[49] (**Fig. 2a; Supplementary Fig. 4a**): (i) a gain of chromosome arm 17q11-qter

('17q')[50], and (ii) an additional gain of chr1q in the 17q background ('17q1q'). Whole-exome

sequencing of 17q and 17q1q cells compared to the parental H7 hESCs revealed a small number of

additional mutations and a loss of a small region in chromosome 2 (**Supplementary Figs. 4b;**

Supplementary Tables 4 and 5). For brevity, we labelled the cell lines by their major CNAs, which

overlap regions commonly gained in NB tumours[51] (**Supplementary Fig. 4c**). 17q1q hESCs were

engineered to include a Doxycycline (DOX)-inducible *MYCN* expression cassette to mimic *MYCN*

amplification in a temporally controlled manner ('17q1qMYCN'). DOX treatment of 17q1qMYCN

resulted in robust induction of MYCN, similar to expression levels in MYCN-amplified tumours

(**Supplementary Figs. 4d-f**). In our experiments, we induced *MYCN* overexpression at D5 (when cells

adopt a definitive NC identity[35]) to avoid bias toward central nervous system differentiation, as seen

upon *MYCN* overexpression in earlier pre-NC progenitors[52].

Equipped with these three isogenic 'mutant' hESC lines, we performed differentiation toward

trunk NC and carried out scRNA-seq as described above, yielding a combined dataset comprising

95,766 cells (**Supplementary Table 1**). To assess how differentiation was affected in each mutant cell

line, we first focused on stages D9, D14, and D19 for which we had data from all four experimental

conditions, and bioinformatically mapped the transcriptomes of mutant cells to our reference of normal

trunk NC differentiation (cp. **Fig. 1b**). While many 17q cells intertwined with all WT cell types (~98%

matching the cognate WT stage), fewer 17q1q and 17q1qMYCN cells advanced beyond WT D14 (only

~48% and 22% matched with WT, respectively; **Fig. 2b**). Only ~4% of 17q1qMYCN cells mapped to

mature cell types (**Fig. 2b**). Altogether, at this level of resolution, we found no evidence that 17q

affected differentiation. In contrast, 17q1q and 17q1qMYCN cells matched WT cells of earlier

developmental stages, suggesting impaired differentiation (**Fig. 2c**).

Next, we tested whether the cell types induced from mutant hESCs still truthfully recapitulated

in vivo cell types as seen for WT. Mapping mutant cells onto the same human embryonic adrenal gland

reference[16] identified proportionally fewer SYM- and MES-like cells in 17q1q and 17q1qMYCN (**Figs.**

2d,e). For cells mapped to the respective cell types, we observed a slightly stronger SCP signature in

17q and 17q1q, while the expression of both MES and SYM genes was weaker relative to the WT (**Fig.**

2f). In 17q1qMYCN, the expression of all signatures was weak, suggesting a failure to fully specify the

expected cell types (**Figs. 2d-g**). Consistently, antibody staining for SOX10 and HOXC9 and flow

cytometry revealed depletion of SOX10+ trunk NC cells in 17q1qMYCN cultures (**Fig. 2h;**

Supplementary Fig. 5). The reduced ability of 17q1qMYCN hESCs to differentiate toward trunk NC

derivatives was also reflected by their failure to generate PERIPHERIN-positive neuronal axons

(**Supplementary Fig. 4g**). A similar, albeit milder effect was observed upon DOX-induced *MYCN*

overexpression at later timepoints (**Supplementary Fig. 4h**).

Differential analysis identified 941 (17q vs. WT), 2,039 (17q1q vs. WT), and 5,915

(17q1qMYCN vs. WT) differentially expressed genes (DEGs) at D9 (**Supplementary Table 6**). As

expected, many upregulated genes were located within the known CNAs (41.4% within chr17q for 17q

cells; 18.7% within chr17q and 25.6% within chr1q for 17q1q cells; **Supplementary Fig. 6a**). Pathway

analysis indicated an enrichment of genes related to E2F and MTORC1 signalling components for

DEGs on chr17q (e.g., *BRCA1*, *NME1*), and of apoptosis-related and members of the p53 pathway on

chr1q (e.g., the anti-apoptotic regulator *MCL1*; **Figs. 3a-c; Supplementary Table 7**). Notably, genes

upregulated in 17q1q also include the p53 inhibitor *MDM4*[53] (**Supplementary Table 6**). These

perturbed pathways may contribute to deregulation of expression of genes outside CNAs (e.g.,

upregulation of MYC targets and oxidative phosphorylation, and downregulation of G2-M checkpoint-

related genes in 17q1qMYCN; **Fig. 3a**). To better resolve the molecular impact of each mutation, we

integrated all datasets into a joint projection of WT and aberrant trunk NC differentiation (**Fig. 3d;**

Supplementary Figs. 6b-h; Supplementary Table 8). The strongest changes were found in

17q1qMYCN, which formed disconnected cell clusters not normally produced in our protocol. To

delineate the stepwise alteration of transcriptional programmes, we placed cells from D9 on a spectrum

from WT to 17q1qMYCN by scoring each cell between 0 and 1 based on the fraction of mutant cells

among its gene expression neighbours ("mutation score"; **Fig. 3e**). This allowed us to identify four sets

of genes (D9_1–D9_4) that were correlated with mutations (**Fig. 3f, Supplementary Figs. 7a,b;**

Supplementary Table 9): Gain of CNAs led to a decrease in expression of genes (gene set D9_3, **Fig.**

3f*)* involved in NC development (e.g., *TFAP2B*[54,55]) and gradual induction of genes (D9_4, **Fig. 3f**)

associated with NC/NB cell migration (e.g.. *ZIC2*, *HOXD3*, *GPC3*[56–58]). *MYCN* overexpression in

17q1qMYCN further repressed genes related to NC development (D9_2; e.g., WNT-antagonist

SFRP1[59] and nuclear receptors *NR2F1/2*[60]) and led to upregulation of MYCN targets implicated in NB

(D9_1; e.g., *NME1* on 17q[61]; **Supplementary Table 9**). Interestingly, we had also found *SFRP1* and

NR2F1 to mark the SYM-MES transition state in WT differentiating sympathoadrenal cells (cp.

Supplementary Table 3). Moreover, we found that many of genes that are upregulated in 17q1qMYCN

(D9_1) were also highly expressed in NB tumours (**Supplementary Fig. 7c**).

We further sought to disentangle the relative contributions of *MYCN* overexpression and CNAs

to the observed differentiation block phenotype in 17q1q cells. To this end, we generated additional cell

lines derived from WT and 17q H7 hESCs by equipping each with a DOX-inducible *MYCN* construct

(WTMYCN, 17qMYCN; **Supplementary Fig. 8a**). Moreover, we introduced the same inducible

MYCN expression cassette into a second female hESC cell line[38] (H9) which had independently

acquired chr17q and 1q gains (H9-WT, H9-17q1q, H9-17q1qMYCN). The differentiation trajectories

of these cell lines in the presence and absence of *MYCN* overexpression were interrogated using split-

pool single-cell RNA-seq. To ensure consistency, we also included the previously analysed H7 cell

lines (WT, 17q, 17q1q, 17q1qMYCN) and performed 2-4 replicate experiments. We obtained a total of

45,546 cells (all D9) post-QC and mapped each dataset onto our WT reference, as we had done before

(**Supplementary Fig. 8b**). Starting with gain of chr17q, we found a reduction in cells allocated to

sensory neuronal differentiation (cluster C4 in **Figs. 1b,c**) and instead a slight increase in a transitional

progeny (C7). With chr1q, we also saw an increase of cells in C10. On top of these changes, *MYCN*

overexpression led to most cells allocating to earlier developmental stages including clusters C2 and

C3 (**Supplementary Figs. 8b,c**) – reflecting the differentiation block we had observed before (cp. **Figs.**

2b,c). The observed changes were consistent for derivatives of both parental hESC lines (H7 and H9).

Analysis of marker gene expression associated with the altered cell clusters (C2, C4, C7, C10) in the

different mutant cell lines identified an upregulation of genes like *AZIN1* in all *MYCN*-overexpressing

cells that was not active in their wild-type or CNA-only counterparts (**Supplementary Fig. 8d**).

Conversely, these cells downregulated developmental regulators. For instance, even though MYCN-

overexpressing cells still expressed remnant gene signatures leading them to map to differentiating WT

cell clusters they downregulated genes in developmental pathways like *HHIP* in cluster C7 or *WNT1* in

C10. Moreover, the neurogenic capacity of the mutant trunk NC cells (as reflected by the presence of

PERIPHERIN-positive neuronal axons) was found to be disrupted by *MYCN* overexpression primarily

in the presence of CNAs, with the strongest phenotype being observed in the presence of both chr17q

and chr1q gains (**Supplementary Fig. 8e**), in line with our earlier findings. Collectively, these data

indicate that CNAs potentiate the *MYCN*-driven block to the induction of a trunk NC/sympathoadrenal

identity.

We conclude that NB-associated CNAs alter the differentiation landscape of hESC-derived

trunk NC lineages by promoting transitional progenitor states at the expense of mature sympathoadrenal

cell types. In conjunction with *MYCN* elevation, they block differentiation and trigger atypical

transcriptional programmes incompatible with normal trunk NC development.

Impaired trunk NC differentiation correlates with acquisition of tumourigenic hallmarks

We observed that ectopic *MYCN* induction altered the morphology of cultures by D14 only in the presence of CNAs as cells lost their ability to spread out and form neurites, and 17q1qMYCN cells even formed tight, dome-like colonies (**Fig. 4a**). As this phenomenon is reminiscent of loss of contact inhibition, a cancer hallmark, we next examined whether CNAs/*MYCN* overexpression led to further cellular changes that are typical of tumourigenesis. We first carried out cell cycle analysis of trunk NC cells (D9) generated from each *MYCN*-overexpressing hESC line (from WT/17q/17q1q backgrounds) by monitoring EdU (5-ethynyl-2´-deoxyuridine) incorporation via flow cytometry. We found a significant increase in the proportion of cells in S-phase only when *MYCN* overexpression was combined with CNAs (p = 0.0233 and p = 0.0073 respectively; two-way ANOVA; **Fig. 4b**) indicating altered cell cycle and increased replication similar to NB tumours and cell lines[62–64]. Immunofluorescence analysis of Ki-67 expression further showed that 17q1qMYCN and 17qMYCN cultures exhibited a higher proliferation rate by D14 compared to their CNA-only counterparts (p < 0.0001 and p = 0.0078, respectively; two-way ANOVA; **Fig. 4c**). We next tested how CNAs/*MYCN* influenced colony formation, another hallmark of tumourigenesis. Low-density plating of trunk NC cells (D9) and image analysis showed significantly increased clonogenicity (p = 0.0109; two-way ANOVA) exclusively in 17q1qMYCN cells (**Fig. 4d**). DOX treatment of the unedited parental wild type and chr17q gain cell lines had no effect (**Supplementary Fig. 9a**).

Previous work has indicated that *MYCN* overexpression alone is associated with increased apoptosis in early sympathoadrenal cells[65,66] and can trigger tumourigenesis only in combination with additional mutations[67–69]. Therefore, we also examined apoptosis levels during the transition of D9 trunk NC cells toward the SCP/sympathoblast stage (D14) by assessing cleaved Caspase-3 levels using flow cytometry. We found that *MYCN* overexpression indeed resulted in a higher rate of apoptosis in the WT background, while this increase was reversed in 17q cells (**Supplementary Fig. 9b**). However, this was not the case in cultures derived from *MYCN*-overexpressing cells with chr17q1q gains, which exhibited apoptosis levels similar to their *MYCN*-overexpressing WT counterparts (**Supplementary Figs. 9b,c**). This phenomenon may be linked to the presence of both pro- and anti-apoptotic genes in chromosome arm chr1q (cp. **Fig. 3c**) as well as increased DNA damage (assessed by the presence of γH2AX foci) following *MYCN* overexpression specifically in the 17q1q background (**Supplementary Figs. 9d,e**). Interestingly, we detected lower levels of DNA damage in the absence of MYCN overexpression in 17q and 17q1q trunk NC cells at D9 compared to WT controls suggesting a potential protective effect of 17q/1q gains (**Supplementary Figs. 9d**). Moreover, we investigated whether *MYCN*-overexpressing cells from different backgrounds had acquired additional mutations during differentiation. Whole-exome sequencing analysis at D19 of differentiation did not reveal any new large CNAs and detected only few mutations (<10 mutations with variant allele frequency >0.2 between D0 and D19; **Fig. 4e**, **Supplementary Fig. 9f**; **Supplementary Table 4**). None of the discovered mutations have previously been reported in NB, leading us to conclude that the observed phenotypic changes in 17q1qMYCN

were likely a product of the CNAs and *MYCN* overexpression rather than an expansion of new clonal
cell populations with additional pathognomonic mutations. Despite an increase of proliferation (cp.
Figs. 4b,c), *MYCN* overexpression did not yield more high-frequency mutations during the short
timeframe of our differentiation experiments, consistent with earlier work in human neuroepithelial
stem cells in *vitro* and after xenotransplantation[70] (p = 0.3458, two-sided, paired Wilcoxon test, n = 3
per group; **Supplementary Fig. 9g**).

Finally, we sought to examine the tumourigenic potential of 17q1q-, 17q1qMYCN- and WT-
derived trunk NC (D9 of differentiation) cells *in vivo* by xenografting them into immunodeficient NSG
mice. We first injected aliquots of about 1 million cells subcutaneously into the recipient animals (n =
6 per cell line) and monitored tumour volume over time. After 3-5 weeks with continuous DOX
administration, all 17q1qMYCN-injected mice developed visible tumours at the injection site (**Fig. 5a**).
In contrast, neither WT- nor 17q1q-injected control animals displayed any signs of tumours for up to
16 weeks (**Fig. 5a**). Likewise, orthotopic injection into the adrenal gland (n = 3 mice per condition)
yielded tumour growths visible by magnetic resonance imaging (MRI) after 5 weeks only when *MYCN*
overexpression was induced by DOX in 17q1qMYCN-grafted animals (**Fig. 5b,c**). We found that both
subcutaneous and adrenal xenograft-derived tumours consisted of undifferentiated, small round cells
similar to tumours from transgenic Th-*MYCN* mice[12] (**Supplementary Fig. 10a**). Complementary to
our analysis in mice, we also performed exploratory xenografts of the same cell lines in zebrafish larvae.
To this end, we labelled our 17q1qMYCN and WT cells at D9 with a fluorescent dye (CellTrace Violet)
and injected them into the perivitelline space of zebrafish larvae on day 2 post fertilisation. Consistent
with our results in mice, we found that 17q1qMYCN cells survived longer in zebrafish than WT, which
had diminished in number at day 1 post injection (dpi) and were completely absent at 3 dpi
(**Supplementary Figs. 10b,c**). In contrast, 17q1qMYCN cells survived until 3 dpi, with 16% of larvae
even showing an increase in xenotransplant size. For comparison, injection of cells from a *MYCN*-
amplified NB cell line (SK-N-BE2C-H2B-GFP[71]) resulted in engraftment with subsequent tumour cell
growth in 84% of larvae (**Supplementary Fig. 10d**).

Together, our results demonstrate that CNA-carrying trunk NC cells transit into an
undifferentiated pre-tumourigenic state and acquire altered cellular properties reminiscent of cancer
hallmarks, such as increased proliferation, clonogenic and tumour formation capacity under the
influence of *MYCN* overexpression. Our data also suggest that CNAs enhance the pro-tumourigenic
effects of *MYCN*.

In vitro **differentiation of mutant hESCs captures NB tumour cell heterogeneity**
Given that cells in our *in vitro* model exhibit similarities to NB cells, we asked whether our data could
provide insights into cellular heterogeneity in NB tumours. To this end, we collected scRNA-seq data
from ten *MYCN*-amplified NB tumours from three independent sources[15,17,23]. For each dataset, we
curated *MYCN*+ tumour cells and bioinformatically mapped these to our reference (**Fig. 6a**). For

example, this approach matched most cells from tumour dataset *Jansky_NB14*[15] to clusters C13 and
C14, which correspond to late SYM-like cell states (**Fig. 6b**; cp. **Fig. 1**). A few cells also mapped to
clusters C11 and C12, i.e., cells with MES-like characteristics. The observed transcriptional
heterogeneity was surprising, given that most tumour cells appeared karyotypically homogeneous
(including a chr17q gain) and expressed *MYCN* (**Fig. 6b**).

Extending the *in vitro* reference mapping to all ten tumours portrayed a spectrum of *MYCN*-
amplified cells with a majority C13- or C14-like expression profile, and a subset of cells mapping to
other differentiating trunk NC cell states (**Supplementary Figs. 11a,b**). We observed apparent
differences between studies and tumour types, but to date there is only a limited amount of single-cell
data from NB tumours to robustly interpret such heterogeneity. We therefore sought to interrogate a
large collection of bulk RNA-seq data from NB tumours (SEQC[72,73]). We first intersected the
development-related gene signatures (C1-C14 from **Fig. 1**) with marker genes identified for the tumour
cells that had been mapped to those respective clusters (from all 10 investigated samples; **Fig. 6c;
Supplementary Table 10**) and labelled each refined signature with an asterisk to distinguish it from
the original gene signature (e.g. signature C13* contained genes such as *DLC1* and *RORA;* **Fig. 6c**).
Applying these gene signatures to the NB tumour data, we found that expression signatures C5*
(sensory neuron-like cells) and C13* (differentiating SYM-like cells), jointly separated *MYCN*-
amplified and non-amplified tumours, as well as tumours at different clinical stages (**Fig. 6d**). The C13*
signature effectively stratified tumours with a good and poor survival across the entire cohort even when
corrected for INSS stage, MYCN amplification status, and age (Cox regression analysis with covariates;
Fig. 6e; Supplementary Table 11).

Jointly, these observations demonstrate that our *in vitro* model generates cell types that
transcriptionally resemble different NB cell subpopulations and that it facilitates the systematic
dissection of intra-tumour heterogeneity in NB tumours.

CNAs and *MYCN* disrupt the configuration of NC regulatory circuits during differentiation
NB tumours and cell lines are marked by a 're-wiring' of non-coding regulatory elements (e.g.,
enhancers) giving rise to tumour-specific regulatory circuitries[44,45,74–78]. We therefore hypothesised that
disruption of developmental TFs also underpins the aberrant differentiation observed in our mutant
hESCs (cp. **Figs. 2, 3**) and employed the assay for transposase-accessible chromatin followed by
sequencing (ATAC-seq)[79] to profile chromatin accessibility in the same samples used for scRNA-seq
analysis (n = 51; **Supplementary Table 1**). Chromatin accessibility serves as a proxy for the dynamic
regulatory DNA element activity during differentiation. For instance, the promoters of the hESC
regulator *POU5F1* and trunk NC regulator *SOX10* were accessible only at D0 and D9, respectively
(**Fig. 7a**), while the *PHOX2B* promoter exhibited reduced accessibility in 17q1q and 17q1qMYCN cells
at D19 consistent with impaired differentiation (**Fig. 7b**).

Unsupervised analysis of chromatin patterns on a global level showed that WT and 17q hESCs

changed consistently throughout differentiation (**Fig. 7c**). In contrast, 17q1q and 17q1qMYCN

appeared not to follow the same path as WT in this low-dimensional projection, in line the

differentiation defects observed in our previous analyses (cp. **Figs. 2b,c**). To delineate chromatin

changes in detail, we performed differential accessibility analysis between all differentiation stages per

cell line and between all cell lines at matched stages (**Supplementary Tables 12, 13**). As in our DEG

analysis, we found an increasing number of regions with altered accessibility in 17q (n = 477 regions),

17q1q (n = 2,826), and 17q1qMYCN (n = 6,663; **Fig. 7d**). In total, there were 45,583 regions with

differential accessibility in at least one comparison, which we divided into nine chromatin modules R1-

R9 (**Fig. 7e**). Modules R1-R7 reflect differentiation order, e.g., regions in module R1 were mostly

accessible at D0, and R6 comprises regions accessible at D14 and D19. Most changes observed in

mutant hESC-derivatives fell within these modules (**Supplementary Figs. 12a,b**). 17q1q and

17q1qMYCN cells failed to close chromatin that is usually specific to D9 (R4, R5) and conversely to

open chromatin regions indicative of late sympathoadrenal differentiation (R6, R7; **Supplementary**

Fig. 12c). Additionally, modules R8 and R9 comprised regions with reduced and increased accessibility

in mutant hESC derivatives, respectively, independently of differentiation stage.

We sought to annotate our chromatin modules by looking for overlaps with genomic regions

accessible in human tissues[80–82] (**Supplementary Table 14**). In line with our transcriptome data, we

found a stepwise change toward chromatin resembling differentiated tissues, e.g., neural tissues in R3-

R5 and mesenchyme/stroma in R6/R7 (**Supplementary Figs. 12d-f**). Next, we examined the identity

of genes near the chromatin modules (**Fig. 7f**). For each module, we found enrichments of specific

marker genes identified in our scRNA-seq analysis of WT trunk NC differentiation (i.e., clusters C1-

C14 from **Figs. 1b,c**). For example, chromatin module R7 (accessible in late differentiation stages, lost

in mutants) was linked to clusters C11/C12 (MES-like cells). Next, we examined TF binding motifs in

each module to identify potential upstream regulators (**Fig. 7g**). We found an enrichment of known

regulators of each developmental stage, e.g., TFs associated with trunk NC in R3-R5 (e.g., SOX10) and

with sympathetic neuron development in R6/R7 (e.g., PHOX2A/B)[39]. Moreover, we found enriched

overlaps of modules R2/R4/R5/R8/R9 with super-enhancers identified in mesenchymal NB cell lines

or adrenergic super-enhancers (in the case of R8), depending on the source annotation used[44,45].

Furthermore, R7 and R9 overlapped super-enhancers associated with subsets of NB tumours[75] with

mesenchymal characteristics and with non-*MYCN*-amplified high-risk tumours, respectively (**Fig. 7h**).

No significant overlap was found with super-enhancers specific for *MYCN*-amplified NB. Finally, we

examined the accessibility of each module across a range of NB cell lines (**Supplementary Fig. 12g**).

As expected, we found that modules R1 and R2 (undifferentiated, early embryonic developmental

stages) and modules R4 and R5 (early trunk NC to sympathoadrenal differentiation) were not accessible

in NB cell lines, while modules R6-R8 (late sympathetic neurons and consistently open in mutants)

were often highly accessible in cell lines. Interestingly, R3 (accessible at NMP and NC stage) was

accessible in most NB cell lines examined except in those with mesenchymal characteristics (SK-N-AS

and SHEP; **Supplementary Fig. 12g**). Using data from other studies, we found that R6-R8 were also

accessible in non-NB cell lines and tissues, while R3 was only found accessible in brain tissue

(**Supplementary Fig. 12g**).

Together, our results suggest a systematic reprogramming of chromatin throughout trunk NC

differentiation. In cells with CNAs and *MYCN* overexpression, this orderly reconfiguration of chromatin

was severely disrupted in a manner similar to NB cells, providing a plausible mechanism for the link

between the observed developmental defects and tumour initiation.

CNA/*MYCN*-driven cell identity loss is mediated by sets of developmental transcription factors

Finally, we investigated the links between CNA/*MYCN*-induced changes in chromatin dynamics, gene-

regulatory networks, and the distorted differentiation trajectories observed at the transcriptional level.

In our scRNA-seq analyses, we had recorded a stepwise alteration of expression from WT to

17q1qMYCN at D9 comprising four gene sets: D9_1 – D9_4 (cp. **Figs. 3f**), which revealed *MYCN*-

driven disruptions of early NC and the sensory neuron lineage specification. We hypothesised that these

mutation-linked gene sets were also regulated by distinct TFs and therefore employed an algorithm to

identify TF targets based on correlated expression patterns[83] (**Fig. 8a**). This analysis identified *NR1D1*

and *TFAP4* as putative TF targets of MYCN (**Figs. 8b,c; Supplementary Figs. 13a,b; Supplementary**

Table 15). The nuclear receptor *NR1D1* has been shown to correlate with *MYCN* amplification in NB

patients[84,85] *and TFAP4* inhibition leads to differentiation of *MYCN*-amplified neuroblastoma cells[86,87],

supporting the validity of the inferred target genes.

We intersected the inferred lists of TF targets with the mutation-linked gene sets (D9_1 – D9_4)

and found an enrichment (**Fig. 8d; Supplementary Table 16**) of MYCN, NR1D1, TFAP4, and ZIC2

targets in D9_1 (highly expressed in 17q1qMYCN). Conversely, the gene set D9_2 (expressed in

WT/17q/17q1q) was enriched for targets of TFs expected at this stage of differentiation, e.g.

SOX4/5/10, TFAP2A/B, and nuclear receptors NR2F1/2. The expression of targets of these TFs

increased or decreased along the mutational spectrum, corroborating their association with the

mutations (**Fig. 8e**). While many TF targets switched expression rapidly with *MYCN* overexpression,

others showed a continuous pattern with up-/down-regulation already detectable in 17q and 17q1q, e.g.,

targets of vagal and early NC regulators HOXB3 and CDX2[88] (up), or of sensory neurogenesis regulator

NEUROD1[39] (down). To aid interpretation, we visualised cell-line-specific interactions between TFs

and targets as edges in connected network diagrams (**Fig. 8f; Supplementary Fig. 13c**). These

diagrams showcased the emergence of a new subnetwork of TFs in 17q1qMYCN that centred on

MYCN and incorporated TFs like NR1D1 and TFAP4. In contrast, a subnetwork involving NC-related

TFs such as SOX10 and TFAP2A/B was lost in these cells. Intriguingly, downregulation of TFs linked

to sensory neuronal development (NEUROD1, ONECUT1) was visible already in 17q cells (**Fig. 8f**),

perhaps explaining why sensory neuron-like derivatives were less abundant in 17q cultures (**Fig. 2b**).

In 17q1q, we additionally observed upregulation of TFs related to early posterior NC specification
including HOXB3, LEF1, and CDX2, which was partially reversed (HOXB3) upon *MYCN*
overexpression (**Fig. 8f**). While many of the TFs implicated in these developmental gene-regulatory
networks are not or only weakly expressed in NB tumours (**Supplementary Fig. 14a**), we found that
the targets of MYCN-related TFs (based on our analysis) are highly expressed in *MYCN*-amplified
tumours (**Supplementary Fig. 14b**). Our analysis also revealed that the targets of 17q/1q-related TFs
strongly are expressed in groups of tumours, but we could not determine whether these contained the
corresponding CNAs due incomplete annotations.

In summary, our data suggest a subtle rewiring of gene-regulatory networks in CNA-carrying
hESCs, which may be linked to the depletion of mature sensory NC derivatives and increased early
SCP signature found in our single-cell analyses (cp. **Fig. 2**). Overexpression of *MYCN* resulted in a
switch in favour of known NC-linked TFs downstream of MYCN.

Discussion

Although CNAs are a principal genetic hallmark of paediatric cancers, it has remained difficult to
determine their exact role in tumour initiation due to the lack of suitable human models. In this study,
we used hESCs carrying CNAs that are prevalent in NB (chr17q and chr1q gains). The NC is a transient
embryonic tissue that is inaccessible after birth; therefore, hESC differentiation allowed us to
experimentally study the effects of these mutations on human sympathoadrenal progenitors, the putative
cells-of-origin of NB.

We provide a comprehensive knowledge base of transcriptomic and epigenetic changes in this
model on a temporal (i.e., during differentiation) and a genetic (i.e., with different mutations) axis. Our
data show that chr17q/1q gains impair trunk NC differentiation and potentiate an SCP-like gene
signature. In this aberrant cell state, overexpression of *MYCN* (mimicking *MYCN* amplification
commonly found along with chr17q/chr1q gains in NB tumours) leads to a complete derailment of
sympathoadrenal lineage specification, and a proliferative, tumour-like cellular phenotype that
correlates with the emergence of NB-like tumours *in vivo*. Moreover, chr17q/1q gains were found to
enhance the *MYCN*-driven differentiation block and acquisition of tumourigenic hallmarks such as
proliferation, clonogenicity and resistance to apoptosis. In line with recent studies[29,89], we speculate that
CNAs provide an early selective advantage manifested by the expansion of undifferentiated cells, which
act subsequently as a NB-initiating entity upon a second oncogenic hit such as *MYCN* overexpression.

The accumulation of NB-associated lesions correlated with a failure to reprogram chromatin
during trunk NC differentiation. Upon gain of chr17q1/q, cells lost TFs associated with sensory
differentiation (e.g., *NEUROD1*) and instead upregulated vagal NC TFs like HOXB3 and WNT-related
effector LEF1[88,90]. *MYCN* overexpression on top of these CNAs abolished chromatin states indicative
of sympathoadrenal differentiation, and instead led to the induction of targets of NR1D1, TFAP4, and

other TFs of the reported NB regulatory circruitry[44,45,74–78]. TFAP4 is a well-established downstream effector of MYCN[86,87], and NR1D1 (Rev-erbα) is a circadian rhythm and metabolic regulator, and a downstream effector of MYCN hyperactivity in NB[84,85]. Together with the appearance of sensory-related signatures in NB tumours (C4* and C5*, **Fig. 6**) our "early *MYCN* onset" scenario reveals previously uncharted disruptions of the early sensory NC lineage, which might complement the currently prevailing model of dichotomic mesenchymal/adrenergic heterogeneity in NB[11,44–46,78,91–94]. Thus, our model will facilitate the functional dissection of these TFs via loss-/gain-of-function approaches to decipher their crosstalk with *MYCN*/CNA-driven tumourigenesis.

Complementing earlier studies using cell lines and animal models[12–14,18,19,22], recent single-cell transcriptomic analyses of NB tumours and metastases[15–17,23] corroborated an origin of NB from neuroblastic, SCP-like progenitors, and highlighted intra-tumour heterogeneity comprising subtypes of tumour cells with adrenergic and mesenchymal properties. In our *in vitro* experiments, we also observed cells expressing signatures of both cell types, suggesting that our model could be useful to experimentally investigate the transition between these and other NB-relevant cell types, providing a new scope into their role in therapy resistance[91]. Furthermore, *MYCN* overexpression (in conjunction with chr17q/1q gains) in nascent trunk NC cells was sufficient to drive tumourigenic traits, suggesting that in some cases NB initiation might occur before SCP/neuroblast emergence and that acquisition of an SCP-like identity may also be a consequence of mutations in earlier stages rather than the origin. We also observed that *MYCN* induction resulted in an apparent block of differentiation when activated at other stages. In future, our cell models will provide the means to dissect the specific effects of *MYCN* at different time points and in specific cell types.

In this study we exploited the phenomenon of culture adaptation of hESCs[49], to obtain sets of cell lines with CNAs that are commonly observed in NB in an otherwise largely isogenic background. Our detailed genetic analyses of the used cell lines revealed other mutations that had naturally arisen in these cell lines (**Supplementary Table 4**), including a point mutation in the BCL6-interacting corepressor *BCOR* (*BCOR*^{L1673F}). *BCOR* mutations have been previously observed in human induced pluripotent stem cell cultures[95,96] and NB patients with *BCOR* mutations exhibit a high frequency of CNAs[89]. *BCOR* mutations have also been reported together with CNAs in other cancers, e.g., retinoblastoma[97]. It would be tempting to speculate that *BCOR* dysfunction might facilitate the ability of cells to tolerate the emergence of certain CNAs; however, to date a causal relationship remains to be established.

Our hESC-based model provides a tractable system for analysing tumour initiation events within disease-relevant human embryonic cell-like populations. In this study, we focused on cell-intrinsic transcriptional regulation since our cultures lack tumour-relevant, non-NC cell types (e.g., immune cells or Schwann cells) and do not recapitulate the structural and physical properties of the human tumour micro- and macroenvironment[98–101]. In the future, it will be possible to combine our

system with 3D co-culture approaches with defined cell types or to use biomimetic scaffolds to emulate
cell-cell interactions and extrinsic environmental influences.

In conclusion, this study unravels the developmental effects of NB-associated mutations and
proposes the progressive corruption of gene-regulatory networks by CNAs as an early step toward
tumour initiation by selection of undifferentiated progenitor phenotypes. Transformation is then
triggered by a second hit with *MYCN* overexpression, which tilts cells toward increased proliferation
and formation of aberrant cell types. Our data provide a direct link between CNAs that commonly
emerge in hESC cultures with impaired differentiation and the acquisition of tumourigenic hallmarks,
thus highlighting the importance of rigorous monitoring of such cultures prior to their use in disease
modelling or cell therapy application in line with recent recommendations from the International
Society for Stem Cell Research[49,102].

Online Methods

Human embryonic stem cell (hESC) cell culture and differentiation

Cell lines and cell culture

We employed H7 and H9 hESCs as karyotypically normal, female WT controls[38]. Use of human embryonic cells has been approved by the Human Embryonic Stem Cell UK Steering Committee (SCSC23-29). Their isogenic chr17q counterparts carry a gain in chromosome 17q (region q27q11) via an unbalanced translocation with chromosome 6 (H7) or a gain of 17q via an unbalanced translocation with chromosome 21 with breakpoints at 17q21 and 21p11.2 (H9)[50,103]. The chr17q1q hESC lines were clonally derived, after their spontaneous emergence following the genetic modification of chr17q hESCs. The H7 chr17q1q-MYCN hESC line was generated by introducing a TetOn-PiggyBac plasmid (PB-TRE3G-MYCN, plasmid#104542, Addgene) carrying the wild-type version of the *MYCN* gene[104] while the H9 chr17q1q-MYCN and H7 WT-MYCN and 17q-MYCN hESC lines were produced using a Tet-On "all-in-one" inducible expression cassette containing the TRE3G promoter driving the expression of *MYCN* with a 2A peptide-linked fluorescent reporter (mScarlet) and a pCAG promoter-driven rtTA3G transactivator[105,106]. Plasmids were introduced via nucleofection using either the Neon NxT Electroporation System (Thermo Fisher Scientific) or the Lonza 4D-Nucleofector System. In the case of the latter, the Amaxa 4D-Nucleofector Basic Protocol for Human Stem Cells was employed with the following modification: $2x10^6$ cells were transfected with 2 µg plasmid in 100 µl Nucleocuvettes. All cell lines were tested regularly for mycoplasma and expression of pluripotency markers. Karyotypic analysis was carried out using G-banding (number of cells examined = 20-30). A rapid qPCR assay was also regularly employed to detect the emergence of common CNAs such as chr17q and 1q gains in our hESC lines[107]. hESCs were cultured routinely in feeder-free conditions at 37°C and 5% CO_2 in E8 media[108] complemented with GlutaMax (Cat# 35050061, Thermo Fisher Scientific) on Vitronectin (Cat# A14700, Thermo Fisher Scientific) or on Geltrex LDEV-Free Reduced Growth Factor Basement Membrane Matrix (Cat# A1413202, Thermo Fisher Scientific) as an attachment substrate. All hESC lines described in this manuscript are available upon request and completion of a Material Transfer Agreement.

Differentiation toward trunk neural crest

hESC differentiation toward trunk NC and its derivatives was performed using a modified version of the protocol described previously[35,36]. Briefly, hESCs were harvested using StemPro Accutase Cell Dissociation Reagent (Cat# A1110501, Thermo Fisher Scientific) and plated at 60,000 cells/cm^2 in N2B27 medium supplemented with 20 ng/ml of FGF2 (Cat# 233-FB/CF, R&D) and 4 µM of CHIR 99021 (Cat# 4423, Tocris) and 10 µM of Rock Inhibitor (Y-27632) (Cat# A11001, Generon) in a volume of 300µl/cm^2. The N2B27 medium consisted of 50:50 DMEM F12 (Merck Life Science / Neurobasal medium (Gibco) and 1x N2 supplement (Cat# 17502048, Invitrogen), 1x B27 (Cat#17504044, Invitrogen), 1x GlutaMAX (Cat# 35050061, Thermo Fisher Scientific), 1x MEM Non-essential amino acids (NEAA; Cat#11140050, Thermo Fisher Scientific), 50 µM 2-Mercaptoethanol (Cat# 31350010, Thermo Fisher Scientific). After 24 hours, media was refreshed removing the Rock Inhibitor and cells were cultured for a further 2 days in FGF2/CHIR to generate NMPs (300µl/cm^2). NMPs at D3 were then re-plated at 50,000 cells/cm^2 (H7) or 40,000 cells/cm^2 (H9) in neural crest inducing medium consisting of DMEM/F12, 1x N2 supplement, 1x GlutaMAX, 1x MEM NEAA, the TGF-beta/Activin/Nodal inhibitor SB-431542 (2 µM, Cat# 1614, Tocris), CHIR99021 (1 µM, Cat# 4423, Tocris), BMP4 (15ng/ml, Cat# PHC9534, Thermo Fisher Scientific), the BMP type-I receptor inhibitor DMH-1 (1 µM, Cat# 4126, Tocris), 10 µM of Rock Inhibitor (Y-27632) on Geltrex LDEV-

Free Reduced Growth Factor Basement Membrane Matrix (Cat# A1413202, Thermo Fisher Scientific) in a volume of 300µl/cm^2. 48 hours later (D5), media was replaced removing the Rock Inhibitor. Media was refreshed at D7 and D8 increasing volume to 500µl/cm^2. On D5, the expression of MYCN was induced by supplementing the neural crest media with 100ng/ml (H7-17q1q-MYCN), 200ng/ml (H7 WT-MYCN, 17q-MYCN), or 1000 ng/ml (H9-derived lines) of Doxycycline (Cat# D3447, Merck). On D9, cells were re-plated at 150,000-250,000 cells/cm^2 in plates coated with Geltrex (Thermo Fisher Scientific) in the presence of medium containing BrainPhys (Cat# 05790, Stem Cell Technologies), 1x B27 supplement (Cat# 17504044, Invitrogen), 1x N2 supplement (Cat# 17502048, Invitrogen), 1x MEM NEAA (Cat# 11140050, Thermo Fisher Scientific) and 1x Glutamax (Cat# 35050061, Thermo Fisher Scientific), BMP4 (50 ng/ml, Cat# PHC9534, Thermo Fisher Scientific), recombinant SHH (C24II) (50 ng/ml, Cat# 1845-SH-025, R and D) and purmorphamine (1.5 µM, Cat# SML0868, Sigma) and cultured for 5 days (=D14 of differentiation) in a volume of 250µl/cm^2. Media was refreshed daily. For further sympathetic neuron differentiation, D14 cells were switched into a medium containing BrainPhys neuronal medium (Stem Cell Technologies), 1x B27 supplement (Invitrogen), 1x N2 supplement (Invitrogen), 1x NEAA (Thermo Fisher Scientific) and 1x Glutamax (Thermo Fisher Scientific), NGF (10 ng/ml, Cat#450-01 Peprotech), BDNF (10 ng/ml, Cat# 450-02, Peprotech) and GDNF (10 ng/ml, Cat# 450-10, Peprotech) for a further 5-14 days (volume of 300µl/cm^2 changing media every other day). Volume was increased up to 500µl/cm^2, depending on cell density, after day 17 of differentiation.

Immunostaining

Cells were fixed using 4% PFA (P6148, Sigma-Aldrich) at room temperature for 10 minutes, then washed twice with PBS (without Ca^{2+}, Mg^{2+}) to remove any traces of PFA and permeabilised using a PBS supplemented with 10% FCS, 0.1% BSA and 0.5% Triton X-100 for 10 minutes. Cells were then incubated in blocking buffer (PBS supplemented with 10% FCS and 0.1% BSA) for 1 hour at RT or overnight at 4°C. Primary and secondary antibodies were diluted in the blocking buffer; the former were left overnight at 4°C and the latter for 2 hours at 4°C on an orbital shaker. Samples were washed twice with blocking buffer between the primary and secondary antibodies. Hoechst 33342 (H3570, Invitrogen) was added at a ratio of 1:1000 to the secondary antibodies' mixture to label nuclei in the cells. We used the following primary antibodies SOX10 (D5V9L) (Cell Signalling, 89356S, 1:500); HOXC9 (Abcam, Ab50839,1:50); MYCN (Santa Cruz, SC-53993, 1:100); PHOX2B (Santa Cruz, SC-376997, 1:500); MASH1 (ASCL1) (Abcam, Ab211327, 1:100 or Santa Cruz, SC-374104, 1:500); Ki67 (Abcam, Ab238020, 1:100); PERIPHERIN (Sigma-Aldrich, AB1530, 1:400); TH (Santa Cruz, 25269, 1:500); cleaved Caspase 3 (Asp175) (Cell Signalling, 9661S, 1:400), yH2AX (Cell Signalling, S139/9718S, 1:400). Secondary antibodies: Goat anti-Mouse Affinipure IgG+IgM (H+L) AlexaFluor 647 (Stratech (Jackson ImmunoResearch) 115-605-044-JIR, Polyclonal 1:500); Donkey anti-Rabbit IgG (H+L) Alexa Fluor 488 (Invitrogen, A-21206, 1:1000).

Intracellular flow cytometry staining

Cells were detached and resuspended as single cells using StemPro Accutase Cell Dissociation Reagent (Cat# A1110501, Thermo Fisher Scientific) and then counted. Next, 10 million cells/ml were resuspended in 4% PFA at room temperature for 10 minutes. Then cells were washed once with PBS (without Ca^{2+}, Mg^{2+}) and pelleted at 200g. Cells were resuspended in PBS at 10 million/ml and used for antibody staining. Permeabilisation buffer (0.5% Triton X-100 in PBS with 10% FCS and 0.1% BSA) was added to each sample, followed by incubation at room temperature for 10 minutes. Samples were then washed once with staining buffer (PBS with 10% FCS and 0.1% BSA) and pelleted at 200g. Then samples were resuspended in staining buffer containing pre-diluted primary antibodies: SOX10 (D5V9L) (1:500; 89356S, Cell Signalling); HOXC9 (1:50; Ab50839, Abcam); cleaved Caspase 3

(Asp175) (Cell Signalling, 9661S, 1:400). The samples were left at 4°C on an orbital shaker overnight.
Then, the primary antibodies were removed, and samples were washed two times with staining buffer.
After washings, staining buffer with pre-diluted secondary antibody was added to the samples and
incubated at 4°C for 2 hours. The secondary antibodies used were Goat anti-Mouse Affinipure IgG+IgM
(H+L) AlexaFluor 647 (Stratech (Jackson ImmunoResearch) 115-605-044-JIR, Polyclonal 1:500);
Donkey anti-Rabbit IgG (H+L) Alexa Fluor 488 (Invitrogen, A-21206, 1:1000). Finally, samples were
washed once with staining buffer, resuspended in staining buffer and analysed using a BD FACSJazz
or a CytoFLEX (Beckman Coulter) flow cytometer. A secondary antibody-only sample was used as a
control to set the gating.
595
596   *Cell cycle analysis*
597   The 5-ethynyl-2´-deoxyuridine (EdU) assay was performed following the manufacturer's instructions
598   (Thermo Fisher Scientific, C10633 Alexa Fluor 488). We used 10μM of Edu for a 2-hour incubation.
599   Cells were analysed in the flow cytometer (BD FACSJazz) using the 405 nm laser to detect the
600   Hoechst/DAPI staining and 488 nm to detect the EdU staining.
601
Low-density plating
Day 9 trunk NC cells derived from hESCs as described above were harvested and plated at a density of
500 cells/cm^2 in plates pre-coated with Geltrex LDEV-Free Reduced Growth Factor Basement
Membrane Matrix (Cat# A1413202, Thermo Fisher Scientific) in the presence of DMEM/F12 (Sigma-
Aldrich), 1x N2 supplement, 1x GlutaMAX, 1x MEM NEAA, the TGF-beta/Activin/Nodal inhibitor
SB-431542 (2 μM, Tocris), CHIR99021 (1 μM, Tocris), BMP4 (15ng/ml, Thermo Fisher Scientific),
the BMP type-I receptor inhibitor DMH-1 (1 μM, Tocris) and ROCK inhibitor Y-27632 2HCl (10 μM)
(300µl/cm^2). The culture medium was replaced the following day with medium containing BrainPhys
(Stem Cell Technologies), 1x B27 supplement (Invitrogen), 1x N2 supplement (Invitrogen), 1x NEAA
(Thermo Fisher Scientific) and 1x Glutamax (Thermo Fisher Scientific), BMP4 (50 ng/ml, Thermo
Fisher Scientific), recombinant SHH (C24II) (50 ng/ml, R and D) and Purmorphamine (1.5 μM, Sigma)
(250µl/cm^2). Plates were then incubated at 37°C at 5% CO2. The media was refreshed every 48 hours.
After 5 days of culture, cells were fixed (PFA 4%/10min) and stained with Hoechst 33342 (Cat# H3570,
Invitrogen) for 5 minutes. Colonies were detected using an InCell Analyser 2200 (GE Healthcare) at a
4X magnification. Images were processed using Cell Profiler.
617
DNA damage analysis
DNA damage was measured by assessing the phosphorylation state of the histone H2AX on the SerCells
were fixed and immunostained using the anti-yH2AX as described above at different time points.
Stained cells were imaged using the InCell Analyser 2200 (GE Healthcare) at 40X magnification. Image
analysis was performed using a pipeline in CellProfiler that allowed us to detect the number of foci of
yH2AX antibody per nuclei.
624
Quantitative real-time PCR
RNA extractions were performed using the total RNA purification kit (Norgen Biotek, 17200)
according to the manufacturer's instructions. cDNA synthesis was performed using the High-Capacity
cDNA Reverse Transcription kit (ThermoFisher, 4368814). Quantitative real-time PCR was performed
using PowerUp SYBR master mix (ThermoFisher, A25780) and run on a QuantStudio 12K Flex
(Applied Biosystems).
631
632

Mouse experiments

Cell preparation for xenotransplantation

H7 wild type, 17q1q and 17q1qMYCN hESCs were differentiated up to day 9 following the protocol described above. Cells were harvested using Accutase to create a single cell suspension, counted and resuspended with media containing Matrigel before injection.

Mice and in vivo experiments

All animal experiments were approved by The Institute of Cancer Research Animal Welfare and Ethical Review Body and performed in accordance with the UK Home Office Animals (Scientific Procedures) Act 1986, the UK National Cancer Research Institute guidelines for the welfare of animals in cancer research and the ARRIVE (animal research: reporting in vivo experiments) guidelines. Female NSG mice were obtained from Charles River and enrolled into trial at 6-8 weeks of age. Mice were maintained on a regular diet in a pathogen-free facility on a 12h light/dark cycle with unlimited access to food and water.

Subcutaneous xenograft

One million cells with 50% Matrigel were injected subcutaneously into the right flank of NSG mice (female; 6-8 weeks old) and allowed to establish a murine xenograft model. Studies were terminated when the mean diameter of the tumour reached 15mm. Tumour volumes were measured by Vernier caliper across two perpendicular diameters, and volumes were calculated according to the formula $V=4/3\pi\ [(d1+d2)/4]3$; where d1 and d2 were the two perpendicular diameters. The weight of the mice was measured every 2 days. Mice were fed with either regular diet or DOX diet (chow containing 20 g of DOX per kg of diet) to induce the expression of *MYCN*.

Orthotopic (adrenal)xenograft

100,000 cells with 50% Matrigel were injected into the right adrenal gland of NSG mice (female; 6-8 weeks old) and allowed to establish a murine xenograft model. Detection of xenografted tumours was performed by magnetic resonance imaging (MRI). The weight of the mice was measured every 2 days. Mice were fed with either standard diet or DOX diet (chow containing 20 g of DOX per kg of diet) to induce the expression of *MYCN*. Magnetic resonance images were acquired on a 1 Tesla M3 small animal MRI scanner (Aspect Imaging). Mice were anesthetised using isoflurane delivered via oxygen gas and their core temperature was maintained at 37 °C. Anatomical T2-weighted coronal images were acquired through the mouse abdomen, from which tumour volumes were determined using segmentation from regions of interest (ROI) drawn on each tumour-containing slice using the Horos medical image viewer.

Pathology

Tissue sections were stained with haematoxylin and eosin (H&E) or specific antibodies (MYCN, Merck; Ki67, BD Pharmingen). Immunohistochemistry was performed using standard methods. Briefly, 5μm sections were stained with antibodies, including heat-induced epitope retrieval of specimens using citrate buffer (pH 6) or EDTA buffer.

Zebrafish experiments

Cell preparation for xenotransplantation

Pre-differentiated neural crest cells were frozen on D7 during their *in vitro* differentiation as described above, shipped, and subsequently thawed in DMEM at room temperature. All cells were retrieved in complete neural crest media as described above and plated onto Geltrex-coated wells in the presence of Rock inhibitor (50µM) for 24 hours. 17q1q cells were additionally treated with DOX (100ng/ml) to induce *MYCN* expression. On D8, media were refreshed, and respective DOX treatment was continued but Rock inhibitor was discontinued. On D9, cells were collected for xenografting experiments and labeled with CellTrace™ Violet (Invitrogen, Thermo Fisher Scientific) for imaging. For this, cells were harvested with Accutase (PAN-Biotech) and resuspended at a concentration of $1*10^6$ cells/ml in PBS. CellTrace™ Violet was added to a final concentration of 5 µM for an incubation time of 10 minutes at 37°C in the dark. The cell-staining mixture was filled up with 5 volumes of DMEM supplemented with 10% FBS and the suspension was incubated for 5 min. After gentle centrifugation (5 min, 500 g, 4°C) the collected cells were resuspended in fresh DMEM medium supplemented with 10% FBS and incubated at 37°C for 10 min. Adhering/ clumping cells were separated via a 35 µm cell strainer. The cell number was adjusted to a concentration of 100 cells/ml in PBS. The freshly stained cells were kept on ice until transplantation. SK-N-BE2C-H2B-GFP cells[71] (a kind gift of F. Westermann) were cultured in RPMI 1640 medium with GlutaMAX™ (Cat# 61870044, Thermo Fisher Scientific) supplemented with 10 % (v/v) fetal bovine serum (Cat# F7524500ML, Sigma), 80 units/ml penicillin, 80 µg/ml streptomycin (Cat# 15140122, Thermo Fisher Scientific), 1 nM sodium pyruvate (Cat# P0443100, PAN-Biotech), 25 mM Hepes buffer (PAN-Biotech) and 8 µl/ml G418. For zebrafish xenotransplantations, the GFP-labelled cells were harvested and resuspended in PBS at a density of 10^5/µl as described above.

Zebrafish strains, husbandry, and xenotransplantation

Zebrafish (*Danio rerio*) were reared under standard conditions in a 14 hours / 10 hours light cycle according to the guidelines of the local authorities (Magistratsabteilung MA58 of the municipal administration of Vienna, Austria) under licenses GZ:565304-2014-6 and GZ:534619-2014-4. For xenotransplantation experiments, the pigment mutant strain mitfa[b692/b692]; ednrba[b140/b140] was used. mitfa[b692/b692]; ednrba[b140/b140] embryos raised at 28°C were anaesthetised with Tricaine (0.16 g/l Tricaine (Cat# E1052110G, Sigma-Aldrich), adjusted to pH 7 with 1M Tris pH 9.5, in E3) and xenotransplanted at 2 days post fertilization (dpf) as previously described[109]. For xenotransplantation, a micromanipulator (Cat# M3301R, World Precision Instruments) holding a borosilicate glass capillary (Cat# GB100T-8P, without filament, Science Products) connected to a microinjector (FemtoJet 4i, Eppendorf) was used. Transplantation capillaries were pulled with a needle puller (P-97, Sutter Instruments) and loaded with approximately 5 µl of tumour cell suspension. Cells were injected into the perivitelline space (PVS) of larvae. Visual inspection was carried out at 2 hours post-injection on an Axio Zoom.V16 fluorescence microscope (Zeiss, Jena) and only correctly injected larvae were used in subsequent experiments and further maintained at 34°C.

Automated imaging and quantification

One day post injection (1dpi) and 3dpi xenografted larvae were anaesthetised in 1x Tricaine and embedded in a 96-well ZF plate (Hashimoto Electronic Industry) with 0.5 % ultra-low gelling agarose (Cat# A2576-25G, Sigma-Aldrich) for automated imaging on a high-content imager (Operetta CLS, PerkinElmer). Images were acquired with a 5x air objective. Exposure times for brightfield images was 40ms at 10% LED power. CellTrace Violet was recorded with an excitation of 390-420 nm at 100% LED power and detection at 430-500 nm using an exposure time of 600ms. GFP was excited with 460-

725 490nm and detected at 500-550nm with an exposure time of 400ms. 23 planes with a distance of 25 μm
726 were imaged per field of view of the laterally orientated larvae to cover the whole tumour. Tumour size
727 was quantified with Harmony Software 4.9 (PerkinElmer).
728
729
730 **Ethical use of data**
731
This study did not generate any new genomics data from patients. However, we performed re-analyses
of previously published (sc)RNA-seq and SNP-array data that was previously collected at our
institutions. The collection and research use of human tumour specimen was performed according to
the guidelines of the Council for International Organizations of Medical Sciences (CIOMS) and World
Health Organisation (WHO) and has been approved by the ethics board of the Medical University of
Vienna (Ethikkommission Medizinische Universität Wien; EK2281/2016, 1216/2018, 1754/2022).
Informed consent has been obtained from all patients or parents/guardians/legally authorized
representatives. The age-adapted informed consent for the CCRI Biobank covers the use of left over
materials from medically necessary surgery or biopsy, which after completion of routine diagnostic
procedures is biobanked (EK1853/2016) and available for research purposes, including genetic
analysis, that are further specified in EK1216/2018 and EK1754/2022: to conduct genetic and
transcriptomic analysis and link to clinical data for survival analysis. All data obtained from external
sources derived from studies where informed consent was given for broad research use.
745
746
747 **Whole-exome sequencing**
748
Library generation and sequencing
Genomic DNA (gDNA) from cell lines was isolated using a desalting method and library preparation
was performed with 100ng gDNA and the Enzymatic Fragmentation (undifferentiated hESC lines;
Supplementary Figs. 4b,c) or Enzymatic Fragmentation 2.0 (cells after differentiation; **Fig. 4e,**
Supplementary Figs. 9f,g) kit, and Twist Universal Adapter System (Twist). For whole-exome
sequencing, the libraries were pooled and enriched with the Exome v1.3 and RefSeq (Human
Comprehensive Exome) spike-in capture probes (Twist) according to the manufacturer's protocols.
Libraries were quantified (Qubit 4 Fluorometer) and quality-checked on 4200 TapeStation and 2100
Bioanalyzer automated electrophoresis instruments (Agilent) and diluted before sequencing by the
Biomedical Sequencing Facility at CeMM on an Illumina NovaSeq SP flowcell in 2x100bp paired-end
mode (median coverage 87.2; **Supplementary Table 1**).
760
Variant identification and annotation
Raw reads were processed using the nf-core *sarek*[110,111] WES pipeline version 2.7.2. Variant calling was
performed in a tumour-normal matched mode, with the parental H7 line serving as the matched normal
sample. Three variant callers, *Mutect2*, *Strelka*, and *Manta*[112–114], were employed for comprehensive
variant identification. Resulting VCF files from *Mutect2* and *Strelka* were normalised using *bcftools*
norm (v1.9)[115] and subsequently annotated using the Ensembl *Variant Effect Predictor* (VEP; v99.2)[116].
The identified variants were filtered based on the default quality control measures implemented in each
tool (*FILTER* column in the VCF contains "PASS"). To identify biologically relevant variants a filtering
strategy was applied that was partly inspired by MAPPYACTS[117]: (i) exclude variants for which
GERMQ or STRQ Phred-scaled values are < 30; (ii) exclude variants with a population allele frequency
of over 0.1% (in 1000 Genomes or gnomAD); (iii) only include variants that have any of
"coding_sequence_variant", "frameshift_variant", "incomplete_terminal_codon", "inframe_deletion",

"inframe_insertion", "missense_variant", "protein_altering_variant", "start_lost", "stop_gained",
"stop_lost" as Consequence; (iv) only include variants that have any of IMPACT == HIGH, SIFT ==
"deleterious", PolyPhen == probably_damaging or damaging[118,119]; (v) exclude variants that have a
variant allele frequency <= 5%.

Copy number calling
CNAs were called by Sequenza (version 3.0.0)[120]. GC content was calculated for hg38 using *sequenza-*
utils gc_wiggle. Depth ratio and B-allele frequency information was calculated using *bam2seqz* for each
non-parental cell line using the parental cell line as a normal reference, single nucleotide variants
(SNVs) were considered heterozygous if the allele frequency was in the range 0.4 to 0.6 (*--het* 0.4).
Data was then binned using the *seqz_binning* command. Autosomes and the X chromosome were then
extracted using Sequenza (*sequenza.extract*) and, as the cell lines are not contaminated with normal
cells as is common place in tumour tissue samples, cellularity was tested in a range of 1 to 1.0002 to
ensure a pure solution was produced by Sequenza. Copy number profiles were then plotted using
ComplexHeatmaps[121]. Breakpoints were considered telomeric if they were within 1Mbp of the
beginning or end of the chromosome.

Phylogenetic analysis
Mutations called by Mutect2[122] with the PASS filter and of VARIANT_CLASS SNV as annotated by
VEP[116] that overlapped with the exome target panel without padding were used for phylogenetic
analysis. Mutations were required to have a minimum variant allele frequency (VAF) of 0.2 to ensure
only high frequency clonal mutations were included in the phylogeny. Phylogenetic trees were
constructed using parsimony and the *phangorn* R package[123]. The parsimony ratchet method (*pratchet*)
was used to search for the best tree and the tree was rooted on the parental cell line. Branch lengths
were calculated using the *acctran* function. Distance between tree tips was calculated using the *distTips*
function in the *adephylo* R package[124]. Phylogenetic trees were plotted using *ggtree*[125].

Pre-processing and analysis of NB SNP-array data
SNP-array data from tumour or bone marrow obtained at diagnosis from Austrian cases with INSS stage
4 high-risk NB[51] were re-analysed for chr17 and chr1 CNAs using VARAN-GIE (v0.2.9), yielding 88
samples with CNAs (>10kb) on at least one of these chromosomes. Genomic segments were manually
curated and plotted using *ggplot2*[126] (v3.3.5). The available CNA annotations based on the human
genome reference hg19. Because of this, the breakpoint annotations for our own cell lines have been
brought from hg38 to hg19 using *liftOver* from the R package *rtracklayer* (v1.54.0).

Single-cell RNA sequencing (10x Genomics)

Library generation and sequencing
Single-cell suspensions were barcoded using oligo-conjugated lipids following the MULTI-seq
workflow and frozen live[127] for G1-G13 (note, G2 was removed due to a technical failure), or frozen
live and barcoded after thawing using the CELLPLEX (10x Genomics) workflow for G14-G27. After
thawing cells were stained with DAPI. A maximum of 10,000 live cells per sample were sorted with a
FACS-Aria v3 and pooled in sets of 3 or 4 samples by differentiation stage (from 3-5 independent
replicate differentiation experiments). Each pooled group was processed using the 10X Genomics
Single Cell 3' v3.1 workflow following the manufacturer's instructions. Enriched barcode libraries were
indexed following the MULTI-seq workflow[127]. After quality control, libraries were sequenced on the

Illumina NovaSeq S4 (G1-13) or S2 (G14-27) platform in 2x150bp paired-end mode. **Supplementary**
Table 1 includes an overview of sequencing data and performance metrics.
822

Raw data processing and alignment
Raw sequencing data were processed with the *CellRanger multi* v7.1.0 software (10x Genomics) for
cell-level demultiplexing and alignment to the human reference transcriptome (*refdata-gex-GRCh38-*
2020-A assembly provided by 10x Genomics). Following initial data processing, all subsequent
analyses were performed in R (v4.1.3) using Bioconductor packages and the *Seurat*[128–130] (v4.1.0)
package.
829

Default basic processing
We applied processing of scRNA-seq data in many instances across this manuscript. Unless parameters
are otherwise specified, the default processing of scRNA-seq counts involved the following steps.
Counts were normalised for read depth using Seurat's *SCTransform*[131] v0.3.3 (parameters:
method="glmGamPoi"; *variable.features.n=5000*), followed by *RunPCA* (keeping the top 50
components), and inference of cell neighbourhoods by *FindNeighbors* on the PCA reduction. Finally,
Uniform Manifold Approximation and Projection (UMAP) was performed using Seurat's *RunUMAP*
function with default parameters. Clustering was performed using FindClusters.
838

Quality control
For each dataset, we first assessed technical covariates and characteristic expression profiles separately.
We kept cells with less than 15% mitochondrial UMI counts, and at least 1000 detected genes and
applied basic scRNA-seq processing and clustering of the cells (*SCTransform*[131] v0.3.3, parameters:
method="glmGamPoi"). Cell cycle scoring was calculated as recommended by Seurat and added as a
variable to regress in SCTransform (*vars.to.regress="ccvar"*). We used clusters devoid of markers
and/or characterised by abnormally high mitochondrial expression, to derive a library-specific UMI
count and mitochondrial percentage threshold for high-quality cells (thresholds for counts /
mitochondrial percentage: G1: 3162/10%; G3: 10000/7.5%; G4: 10000/8%; G5: 3162/10%; G6:
10000/8%; G7: 12589/8%; G8: 7943/10%; G9: 7079/10%; G10: 3981/7.5%; G11: 3981/10%; G12:
5012/10%; G13: 10000/10%; G14: 5500/13%; G15: 3500/5%; G16: 3000/8%; G17: 2000/8%; G18:
3500/10%; G19: 1800/6%; G20: 3000/15%; G21: 6000/8%; G22: 5000/6%; G23: 3000/6%; G24:
1500/6%; G25: 3500/5%; G26: 2000/10%; G27: 3000/15%). In addition, empty and doublet droplets
were flagged with *Emptydrops*[132] (v1.14.2; default parameters) and *scDblFinder*[133] (v1.8.0; parameters:
dbr=0.01), respectively. We retained only cells with *Emptydrops* FDR>0.05 and individual scDblscore
cutoffs for the datasets were: G1: 0.01; G3: 0.016; G4: 0.005; G5: 0.005; G6: 0.003; G7: 0.005; G8:
0.005; G9: 0.005; G10: 0.005; G11: 0.005; G12: 0.005; G13: 0.005; G14: 0.005; G15: 0.005; G16:
0.0075; G17: 0.002; G18: 0.007; G19: 0.00375; G20: 0.01; G21: 0.007; G22: 0.007; G23: 0.0125; G24:
0.003; G25: 0.007; G26: 0.005; G27: 0.007.
858

859 *Sample demultiplexing*
860 To demultiplex cells belonging to different pooled samples, we used deMULTIplex2[134] (v1.0.1) with
861 default parameters on each dataset using the tag counts from *CellRanger multi*. All non-singlet cells
862 were filtered out from the dataset.
863

Normalisation, clustering, and marker gene analysis for the main dataset
Raw UMI counts were normalised using Seurat's *SCTransform*[131] (parameters:
variable.features.n=5000, method="glmGamPoi", vars.to.regress="ccvar") to account for
differences in sequencing depth and cell cycle phase (the variable "*ccvar*" variable was calculated as

the difference of S and G2/M scores using Seurat's *CellCycleScoring* method with default parameters).
To integrate data from 3-5 independent differentiation experiments (replicates; **Supplementary Table**
1), we used *scVI*[135] (v0.20.3; parameters: *n_epochs = 50*) using 5000 highly variable features of the
input data with Python 3.11 via reticulate (v1.24). Nearest neighbours were identified using Seurat's
FindNeighbors function (parameters: *k=30*) on the ten scVI components. The same scVI reduction was
used to find a low dimensionality UMAP projection using Seurat's RunUMAP for both the WT-only
(*n.neighbors=50, min.dist=.5, dims=1:8*) and full dataset (*n.neighbors: 30, min.dist=0.4, dims=1:8*
method="umap-learn", metric="correlation"). Clusters on the UMAP projection were defined using
Seurat's *FindClusters* (parameters [full dataset]: *resolution=0.6,* parameters [WT-only]:
resolution=0.4, algorithm=4). Neighbouring clusters that shared functional markers were merged
manually and relabelled to roughly reflect differentiation order. Finally, markers for each cluster were
identified using the *FindAllMarkers2* function (*DElegate*[136] v1.1.0; parameters: *method="deseq",*
min_fc=1, min_rate=0.5, replicate_column="replicate"), with each cluster compared to all the other
cells in the dataset. Genes with an adjusted P-value less than 0.05 were selected as markers.
(**Supplementary Tables 2, 8**). To compare mutant and wild-type cells, we filtered the integrated dataset
to cells from D9 and identified pairwise DEGs ($P_{adj} \leq 0.05$, $|log_2FoldChange| > 0.25$) between each
mutant condition and WT using the *findDE*function (Delegate v1.1.0; parameters:
group_column="condition", method="deseq", replicate_column = "day_rep"). We discarded DEGs
that were not expressed in at least 20% of cells on one side of the comparison. Up- and down-regulated
DEGs on chr1q, on chr17q, and outside either CNA were then tested separately to identify significant
overlaps with MSigDB HALLMARK[137] gene sets using the hypergeometric test implemented in the
hypeR[138] package (v1.10.0). DEGs and enriched pathways are listed in **Supplementary Tables 6 and**
7.
891
Pseudotime trajectory analysis
Pseudotime trajectories were inferred using *Slingshot*[139] (v2.2.0; default parameters) using a filtered
dataset comprising only MES-SYM clusters C11, C12, C13, and C14 (cp. **Fig. 1d; Supplementary**
Fig. 3). The filtered dataset was reprocessed using the basic scRNA-seq processing workflow as
described above and the first two principal components were used to find trajectories between two
extreme clusters. Only one trajectory was found. Genes whose expression was associated with the
trajectories were identified with the generalised additive model and association test as implemented in
tradeSeq[140] (v1.8.0; parameters: *knots=5)*. The top genes with the highest Wald statistic were selected
for reporting (**Supplementary Table 3**). Transcription factors were identified based on the human
transcription factors database[141] in **Supplementary Fig. 3b.**
902
Cross-dataset annotation, label transfer, and signature scores
To map data between scRNA-seq datasets, we employed Seurat's label transfer workflow. Both query
and reference datasets were processed using the default basic scRNA-seq processing workflow as
described above and mapped (*FindTransferAnchors*, *TransferData*, *IntegrateEmbeddings*,
NNTransform, and *MappingScore* functions; default parameters) using the 50 first principal
components of the PCA reduction from both datasets. To visualise cell mappings, we used "glasswork
plots", in which the UMAP of the reference was used to define the coordinates of convex hulls for each
cluster. Query cells mapping to each cluster were plotted at random positions within their cognate
reference cluster hull to mitigate overplotting bias when many cells mapped to a small neighbourhood.
In this study, the following mappings were performed with the same processing and parameters:
1. Human foetal adrenal reference datasets[15,16] mapped onto WT-only (**Figs. 1d-f;**
Supplementary Figs. 2i,j, 3c) and full *in vitro* (**Figs. 2d-f, 3d; Supplementary Fig. 6g**)
scRNA-seq references. Upon obtaining consistent results for both (**Supplementary Fig. 2j**),

the reference provided by Kameneva *et al.* was used throughout the analysis, because of the curated cell type markers they provided (**Supplementary Fig. 2i**). These gene signatures were also quantified with Seurat's *AddModuleScore* function (default parameters) in **Figs. 1e,f, 2f**.

2. Our mutant scRNA-seq data mapped onto the wild-type reference (**Figs. 2b,c**).
3. NB tumour scRNA-seq data mapped onto our WT-only reference (**Figs. 6b,c; Supplementary Fig. 11**). See additional details about these datasets and processing in the section *"Pre-processing and mapping of NB tumour data"* below.
4. Extended data from a split-pool scRNA-seq (Parse Biosciences) mapped to the WT-only dataset (10x Genomics) (**Supplementary Fig. 8b-c**).

Validation of label transfers

WT mappings to adrenal gland references were validated by the presence of relevant markers (**Supplementary Fig. 2i**). Mutant and tumour cell mappings were not strictly curated via markers (i.e., they were allowed to deviate). When analysing markers of mapped mutant and tumour cells, cells with a prediction score of 0.4 or higher were used to minimise ambiguous mappings and maximise marker discovery. Shared markers were consistently found between the query and the cognate cells in the reference, even though their number varied (**Fig. 6c, Supplementary Fig. 8d**).

Visualising label transfers with glasswork plots

To visualise cell mappings, we used "glasswork plots", in which the UMAP of the reference was used to define the coordinates of concave hulls for each cluster (calculated with R package *concaveman* v1.1.0). Query cells mapping to each cluster were plotted at random positions within their mapped target cluster hull to mitigate overplotting. Input cell populations for the plot were downsampled evenly by condition and stage (n = 1000 cells) to avoid sampling effects.

Mutation score analysis

To calculate the mutation score, we focused on days 9, 14 and 19 as they contained samples from all conditions. We encoded each cell's genotype as a number G based on the genetic lineage of hESC lines: $G(WT)=0$, $G(17q)=1$, $G(17q1q)=2$, and $G(17q1qMYCN)=3$. We then calculated the mutation score m as the mean G of the cell's K nearest neighbouring cells ($K=30$) in the scVI reduction's neighbourhood graph (see *"Normalization, clustering, and marker gene analysis"*). Division by 3 yielded a score between 0 and 1. Intuitively, the mutation score of a cell indicates whether a cell phenotypically resembles wild-type cells or cells with a given number of relevant alterations independent of its own genotype. To find genes correlated with the mutation score, we calculated Pearson correlation coefficients with gene expression in three settings: (i) correlation for each gene with m in all cells; (ii) correlation for each gene with m leaving out the 17q1qMYCN cells, to emphasise subtle correlations with CNAs; and (iii) correlation for each gene and the neighbourhood entropy (Shannon entropy of all genotype scores G of the K nearest neighbours), to find genes appearing in mixed regions. All non-duplicate absolute correlations (calculated using R's *cor.test*, parameters: *method="pearson", exact=TRUE*) were subject to Bonferroni correction and ranked. The top-300 correlated genes ($p \leq 0.05$) per differentiation stage (D9, D14, D19) are reported in **Supplementary Table 9**.

Pre-processing and mapping of NB tumour data

We collected scRNA-seq data for tumours with reported *MYCN* amplification from three sources from the stated database or the corresponding authors:
- Three samples (all primary adrenal, 2 male [Dong_T162, Dong_T230], 1 female [Dong_T200; accession GSE137804 [Gene Expression Omnibus])[17],

963   -   three samples (1 primary adrenal, 1 primary intraspinal, 1 relapse/occipital subcutaneous bone
964   metastasis [Jansky_NB14]; 1 female [Jansky_NB08], 2 male [Jansky_NB01, Jansky_NB14];
965   accession EGAS00001004388 [European Genome-Phenome Archive])[15],
966   -   and four samples (all metastatic bone marrow; 3 female [Fetahu_M1, Fetahu_M3, Fetahu_M4],
967   1 male [Fetahu_M2]; accession GSE216176 [Gene Expression Omnibus])[23].

Additional details about each dataset are available from the original research articles. In each dataset,
cells with more than 500 reads per barcode and mitochondrial DNA less than 20% were kept for further
analysis. We then performed an adrenal gland mapping[16] (same workflow as described above) and
discarded cells mapping to the category "HSC_and_immune". This process left us with strong CNA
profiles (see below) at key genomic positions such as chr2p (*MYCN* locus). Cells were then subjected
to default basic scRNA-seq processing (see above) and mapped onto our WT-only reference (see
above).

Inference of CNA profiles from scRNA-seq data
To infer tumour cell CNA profiles from scRNA-seq expression data, we used the *infercnv*[142] R package
(v1.10.1). We first removed cells with less than 500 UMI counts. Then, we created a pan-patient healthy
reference cell population by sampling from each patient 500 cells that we determined to be
HSC/immune cells based on a mapping to a human embryonic adrenal gland reference[16]. For every
patient, we then ran *infercnv* with the non-HSC/immune cells as the main input and the pan-patient
HSC/immune cells as a reference. The *cutoff* parameter was set to 0.1, all other parameters were left at
their default values.

983
Pre-processing and analysis of NB bulk RNA-seq data
We obtained bulk RNA-seq counts and associated metadata from patient-derived NB samples from
three sources: TARGET[24] (phs000467 [Genomic Data Commons]), SEQC[72,73] (GSE49711 [Gene
Expression Omnibus]) and from our institution[98,101,143–149] (labelled "CCRI" in the
figures; GSE94035, GSE147635 and GSE172184 [Gene Expression Omnibus]). Open access
unstranded counts from TARGET patients were obtained directly from the GDC data portal (subsection
TARGET:NBL, phs000467). Counts from the CCRI patients were obtained in-house. Both CCRI and
TARGET datasets were normalised using DESeq2[150] (v1.34.0) and transformed using the variance
stabilizing transformation. A prenormalised $\log_2$ SEQC matrix   was exponentiated, rounded to the
nearest integer, and subjected to variance stabilizing transformation. In all datasets, the names of
relevant marker genes were harmonised manually in case the gene was found with a different name.
Each dataset was analysed separately due to differences in count quantification and normalisation. PCA
projections of the normalised variables revealed mainly biological/clinical variables (and not technical
variables) having major weight in the variance of the datasets. Only NB data collected at diagnosis were
used for our analyses (discarding, e.g., ganglioneuroblastoma and relapse data). To quantify the *in-vitro*
cluster signature strength, we used the intersection of markers found both in our *in-vitro* WT-only
dataset (**Supplementary Table 2**) and in the tumour scRNA-seq datasets (**Supplementary Table 10**).
We then used the function *gsva* (from *GSVA*[151] v1.42.0; parameters: *method="ssgsea"*) to calculate
signature scores for each of the shared cluster signatures.

1003
Survival analysis
We obtained survival data for the SEQC cohort from the original publication[72]. Event-free survival
(EFS) was defined as time from diagnosis to any of the following events: Relapse/progression of
disease, or death due to any cause and secondary malignancies. Patients without events were censored
at last follow-up evaluation. EFS was estimated with Kaplan-Meier estimates and evaluated with Cox-

regression. Cluster signatures (see previous section) were dichotomized using the median-value. The
results of the survival analysis are reported in **Supplementary Table 11**.

1011
1012

1013 **Split-pool single-cell RNA sequencing (Parse Biosciences)**

1014

Library generation and sequencing
Cells were harvested with Accutase to create a single-cell suspension, then were counted using Bio-rad
Tc10 Automated cell counter in the presence of Trypan Blue Stain (Bio-rad). For cell fixation we used
the Evercode Fixation v2 Kit (SKU: ECF 2001, Parse Biosciences, Seattle, USA) as per manufacturer
instructions. A maximum of 5,000 cells per sample were multiplexed using the Evercode WT Mega v2
kit (Parse Biosciences). Three rounds of combinatorial barcoding were performed, and cells were then
pooled and split into 16 sub-libraries (one small 5,000-cell sub-library and 15 large sub-libraries of
32,000 cells each). After DNA amplification and library prep, the small library was sequenced as part
of a larger Illumina NovaSeq S4 flowcell and the 15 large sub-libraries on one dedicated NovaSeq S4
platform in 2x150 bp paired-end mode.

1025

Raw data processing and alignment
Raw sequencing data were processed with the *split-pipe* v1.0.6p software (Parse Biosciences) for cell-
level demultiplexing and alignment to the human reference transcriptome (*refdata-gex-GRCh38-2020-*
A assembly provided by 10x Genomics; parameters: *-m all -c v2*). Following initial data processing, all
subsequent analyses were performed in R (v4.1.3) using Bioconductor packages and the *Seurat*[128–130]
(v4.1.0) package.

1032

Basic processing, quality control, and marker analysis
We applied the *cb_filter_count_matrix* (default parameters) function from *canceRbits* (v0.1.6; default
parameters) to remove cells with high mitochondrial counts (>15%), unusually high/low number of
genes (< 300 genes or z-score of log(genes) not in range (-3, 3)), abnormally high/low number of reads
(z-score of log(transcripts) not in range (-3, 3)), or an abnormal transcript-to-gene ratio (z-score of
residuals of loess fit of "log(genes) ~ log(transcripts)" not in range (-5, 5)), and the *cb_seurat_pipeline*
function (parameters: *seurat_max_pc = 15, metric = "manhattan", k_param = 20, n_neighbors = 40,*
cluster_res = 0.3) to perform a standard Seurat analysis workflow including data normalisation,
dimensionality reduction, and clustering. Subsequently, the data were mapped to our 10x-based WT-
only reference as described above. To identify marker genes for cells mapped to different clusters of
the WT reference (for cells with prediction score >=0.4) we again used DElegate::FindAllMarkers2.
(DElegate v1.1.0; *parameters: method="deseq", min_fc=1, min_rate=0.5,*
replicate_column="replicate"), and kept all genes with adjusted pvalue of 0.05 of less.

1046
1047

1048 **Chromatin accessibility mapping (ATAC-seq)**

1049

Library generation and sequencing
ATAC-seq was performed as described previously[79]. Briefly, 20,000 to 50,000 cells were lysed in the
transposase reaction mix (12.5 μl 2xTD buffer, 2 μl TDE1 [Illumina], 10.25 μl nuclease-free water, and
0.25 μl 1% digitonin [Promega]) for 30 min at 37 °C. Following DNA purification with the MinElute
kit (Qiagen) eluting in 12 μl, 1 μl of eluted DNA was used in a quantitative PCR (qPCR) reaction to
estimate the optimum number of amplification cycles. The remaining 11 μl of each library were
amplified for the number of cycles corresponding to the Cq value (i.e., the cycle number at which

fluorescence has increased above background levels) from the qPCR using custom Nextera primers.
Library amplification was followed by SPRI (Beckman Coulter) size selection to exclude fragments
larger than 1,200 bp. Libraries concentration was measured with a Qubit fluorometer (Life
Technologies), and libraries were quality checked using a 2100 Bioanalyzer (Agilent Technologies).
Libraries were sequenced by the Biomedical Sequencing Facility at CeMM using the Illumina HiSeq
4000 platform in 1x50bp single-end mode. **Supplementary Table 1** includes an overview of the
sequencing data and performance metrics.
1064
Raw data processing, alignment, and quality control
Raw sequencing data were processed using *PEPATAC*[152] (v0.9.5; default parameters) including
alignment to the human genome (*refdata-cell ranger-atac-GRCh38-1.2.0* assembly provided by 10x
Genomics for maximum compatibility with scRNA-seq analyses). Following initial data processing, all
subsequent analyses were performed in R (v4.1.3) using Bioconductor packages and *ggplot2*[126] (v3.3.5)
and *ComplexHeatmap*[121] (v2.10.0) for plotting. After discarding low-quality data (NRF<0.65 or
PBC1<0.7 or PBC2<1 or FRiP<0.025), we removed peaks overlapping blacklisted regions from
ENCODE (http://mitra.stanford.edu/kundaje/akundaje/release/blacklists/hg38-
human/hg38.blacklist.bed.gz) and merged overlapping peaks across all ATAC-seq datasets to create a
common set of consensus genomic regions for subsequent analysis (**Supplementary Table 12**). Next,
we quantified for each input dataset the number of reads overlapping these consensus peaks using
featureCounts[153] (*Rsubread* v2.8.2).
1077
Differential accessibility analysis and chromatin modules
Raw read counts were loaded into DESeq2[150] (v1.34.0; default parameters, design:
~lane+batch+sample_group) for normalization (variance-stabilizing transformation) and differential
analysis. In doing so, we estimated count size factors for normalization excluding regions on
chromosomes with known chromosomal aberrations (i.e., chr1, chr17) to avoid overcompensation due
to differences in global signal strength. We queried all pairwise comparisons of sample groups stratified
by cell line / condition stratified (time-wise differences, e.g., WT-D3 vs. WT-D0) and between
conditions stratified by stage (condition-wise differences, e.g., 17q-D9 vs. WT-D9) and recorded all
significantly differentially accessible regions ($P_{adj} \leq 0.005$, $|\log_2 FoldChange| \geq \log_2(1.5)$; parameters:
pAdjustMethod="BH", lfcThreshold=log2(1.5), independentFiltering=TRUE; **Supplementary Table**
13). To define chromatin regulatory modules, we divided time-wise differences in WT hESCs (n =
41,699 regions) into six chromatin modules (R1-R6) and condition-wise differences (n = 3,914 regions)
into three chromatin modules (R7-R9) by hierarchical clustering using the Ward criterion (parameter:
method = "ward.D2"). To associate ATAC-seq regions with putative target genes, we used the
GenomicRanges[154] package (v1.46.1) to assign each region to all genes (using the *refdata-gex-GRCh38-*
2020-A gene annotation provided by 10x Genomics) with overlapping promoters (transcription start
side) or to distal genes whose promoter within a maximum distance of 250kb whose expression was
significantly correlated with the region's accessibility. To this end, we calculated the correlation
coefficient between normalised read counts in our ATAC-seq data with the normalised read counts in
our matching scRNA-seq data (mean of cells per sample; note, the ATAC-seq was collected from the
same experiments as the first replicate experiments for scRNA-seq). We calculated an empirical false
discovery rate (FDR) by shuffling RNA/ATAC assignments (10 repetitions) and retained associations
with a value ≤ 0.05. Annotated regulatory regions from the analysis of ATAC-seq data are listed in
Supplementary Table 12.

1102

Overlap enrichment analysis for chromatin modules

1103

To characterise the chromatin modules, we interrogated overlaps with genomic regions or associated

genes using the hypergeometric test implemented in the *hypeR*[138] package (v1.10.0) via the *cb_hyper*

function (*canceRbits* v0.1.6; parameters: collapse = FALSE, min_size = 5, max_size = (<75% of the

size of the background dataset)). We looked at three types of overlaps: (a) Annotated reference regions

from the DNase hypersensitivity index[80], from the Cis-element Atlas[81], from the Enhancer Atlas[82], and

NB subgroup-specific super-enhancers[75], which all catalogue regulatory elements active in different

cell or tissue types. (b) Matches to known TF motifs from the *HOCOMOCO* database[155] (v11). Here,

we downloaded motifs from the *HOCOMOCO* website

(*HOCOMOCOv11_full_annotation_HUMAN_mono.tsv*) and used *motifmatchr* (v1.16.0) to scan the

DNA sequences underlying each genomic region for matches. Regions with at least one match to the

motif were recorded as potential binding sites. (c) Marker genes from our scRNA-seq analysis of WT

hESC differentiation (**Fig. 1c**; **Supplementary Table 2**). For this purpose, genomic regions were

associated with genes as described above. In each case, we used the entire set of all analysed genomic

regions as a background for the enrichment analysis, and we considered overlaps with an FDR-corrected

P-value less than 0.005 as significant. For motifs, we find the reported P-values are inflated and

therefore used stricter thresholds: $P_{adj} \leq 0.0000001$, $|\log_2 \text{odds}| > \log_2(2)$. All enrichment results are

reported in **Supplementary Table 14**.

1121

Integration with published ATAC/DNaseI-seq data

1122

To interrogate accessibility of the chromatin modules in existing data from NB cell lines we used fast

gene set enrichment analysis *fgsea* (v1.20.0)[156]. We obtained ready-processed genomic coverage tracks

(*wig* or *bigwig* files) from three studies profiling NB cell lines[75,157,158] (GSE138293, GSE224241,

GSE136279). Additionally, we obtained data from three studies profiling breast[159] (GSE202511) and

lung cancer[160] lines (GSE228832), or human tissue data[161] (https://epigenome.wustl.edu/epimap) as

controls. For studies based on older genome assemblies (GSE138293, GSE224241, GSE136279,

GSE228832, and EpiMap used hg19), we converted our peak coordinates to hg19 using the *liftOver* R

package (v1.18.0). We then used the *GenomicRanges*[154] (v1.46.1) and *plyranges*[162] (v1.14.0) packages

to identify genome segments overlapping our peaks and to aggregate the corresponding mean score

reported in the coverage tracks, which were then used for gene set enrichment analysis via the *cb_fgsea*

function (*canceRbits* v0.1.6; parameters: max_size = Inf).

1134

Identification of transcription factor targets

1135

To identify putative target genes of TFs, we used *GRNboost2*[83] (*arboreto* library v0.1.6, with Python

v3.8.17 via reticulate [v1.24]) to identify genes whose expression could be predicted from the

expression of each TF. We tested all TFs in the *HOCOMOCO* database[155] for which at least one motif

could be identified in our dataset. We found that stronger association values were reported for stem-

cell-related factors, likely because of a proportional overrepresentation of this developmental stage in

our dataset. To alleviate this effect and create more balanced data to build our networks on, we

downsampled our dataset to no more than 500 cells per cluster and took the average importance value

of ten random samples forward for further analysis. Putative targets with high importance values but

without a supporting nearby ATAC-seq peak with a motif matching the respective TF were considered

indirect targets and discarded from the target gene sets. We found that the range of importance values

varied between TFs. We therefore calculated a TF-specific threshold on the importance score to define

target genes. To this end, we ranked importance values and used the *changepoint* package (v2.2.3;

default parameters) to identify the first point at which the mean values of the curve of importance values

changed (disregarding the top 1% highest importance values which often were outliers and disrupted

this analysis). The resulting target gene sets were divided into putative activating and inhibiting interactions by the sign of the Pearson correlation coefficient r of the respective TF-target pairs (using the mean correlation value of the same eight random samples as used for *GRNboost2*). Interactions with $|r|<0.1$ were discarded. To calculate the average expression of target genes in each cell we used only activated targets ($r>0.1$) and the Seurat module score. To identify significant overlaps between target genes and gene sets D9_1 – D9_4 (**Supplementary Table 15**), we used the *hypeR*[138] package (v1.10.0) via the *cb_hyper* function (*canceRbits* v0.1.6; parameters: collapse = FALSE, min_size = 0, max_size = Inf), considering TFs with $P_{adj} \leq 0.05$, $|\log_2 \text{odds}| \geq \log_2(4)$, and frequency $\geq 5\%$ as significant. All target gene sets are reported in **Supplementary Table 15** and all enrichment results in **Supplementary Table 16**.

Gene-regulatory network visualisation

For the visualisation of gene-regulatory networks, we used the *igraph* package (v1.3.1). A directed graph was constructed from edges between genes in the gene sets D9_1, D9_2, D9_3, or D9_4 (**Supplementary Table 9**) and TFs found enriched in the overlap with these genes (**Fig. 8d**). The same automated graph layout (function *layout_with_fr()*) was used to draw mutant-specific network diagrams. To generate mutant-specific networks (**Fig. 8f**), we selected cells derived at D9 and parameterised node colour to indicate the mean scaled expression of the genes in those cells and node size to indicate the mean scaled TF target score (Seurat module score) for TFs or the mean scaled expression for non-TFs. To simplify plots, we only labelled TFs with positive mean scaled expression values (>0.05) and manually aggregated many overlapping values, but all node labels are shown in **Supplementary Fig. 12c**.

Data availability

Code availability

Acknowledgements

We would like to thank the Biomedical Sequencing Facility at the CeMM Research Center for Molecular Medicine of the Austrian Academy of Sciences for assistance with next-generation sequencing, Bettina Brunner-Herglotz (CCRI) for her technical assistance, and Duncan Baker (Sheffield Diagnostic Genetic Services, Sheffield Children's Hospital) for carrying out karyotyping of hESC lines. We would like to acknowledge Yann Jamin and Barbara Martins da Costa (ICR, London) for help with the MRI and animal work, respectively. We are also grateful to Igor Adameyko and Polina

Funding

Funding sources in authorship order: K.B.: Biotechnology and Biological Sciences Research Council (BBSRC) (DTP Studentship, BB/T007222/1); I.S.F.: Austrian Science Fund (FWF) (10.55776/P35072); M.C.B.: Austrian Academy of Sciences (25905); E.M.P.: FWF (10.55776/P32001, 10.55776/P34832); M.H.: National Institutes of Health (R00CA197484); I.B.: UK Regenerative Medicine Platform (MR/R015724/1), Medical Research Council UK (MRC) (MR/X000028/1, MR/X007979/1); M.D.: Austrian Research Promotion Agency (FFG) project 7940628 533 (Danio4Can), Alex's Lemonade Stand Foundation for Childhood Cancer (ALSF) (20-17258); L.C./E.Po.: CRUK Programme Award A28278; L. C.: HEFCE/ICR; S.T.M.: Vienna Science and Technology Fund (WWTF; project LS18-111), FWF (10.55776/P35841B MAPMET), H2020 (project 826494 PRIMAGE); M.F.: ALSF (20-17258); A.T.: BBSRC (New Investigator Research Grant, BB/P000444/1), European Union Horizon 2020 Framework Programme (H2020-EU.1.2.2; project 824070), Children's Cancer and Leukaemia Group/Neuroblastoma UK/Little Princess Trust (CCLGA 2019 28, CCLGA 2020 19, together with H.B.), MRC (MR/V002163/1); F.H.: ALSF (20-17258), FWF (10.55776/TAI454, 10.55776/TAI732, 10.55776/PAT1300223).

Author contributions

I.S.G. and L.M.G. championed the experimental and computational work on this study, respectively. Formal contributions in authorship order (CrediT taxonomy): Conceptualization: A.T., F.H.; Data curation: L.M.G., I.S.G., L.S., K.B., E.Po., S.W.Z., E.B., I.S.F., P.Z., U.P., G.C., S.T.M., M.F., A.T., F.H.; Formal Analysis: L.M.G., I.S.G., C.H., E.Po., S.W.Z., P.Z., U.P., C.St., M.S., G.C., H.B., A.T., F.H.; Funding acquisition: K.B., I.S.F., M.C.B., W.W., P.A., I.B., H.B., M.D., L.C., S.T.M., M.F., A.T., F.H.; Investigation: I.S.G., L.M.G., K.B., E.Po., L.S., D.S., R.Le., E.B., I.S.F., M.B., A.W.W., C.St., C.So., S.T., P.B., M.R., M.G., M.C.B.; Methodology: I.S.G., L.M.G., L.S., E.Po., I.S.F., M.B., P.B., M.G., G.C., H.B., M.D., L.C., S.T.M., M.F., A.T., F.H.; Project administration: A.T., F.H.; Resources: W.W., E.Pu., M.H., R.La., H.B., M.D., L.C., S.T.M, M.F., A.T., F.H.; Software: L.M.G., C.H.; Supervision: E.Pu., G.C., H.B., M.D., L.C., S.T.M., M.F., A.T., F.H.; Visualization: I.S.G., L.M.G., K.B., E.P., C.St., G.C., A.T., F.H.; Writing – original draft: I.S.G., L.M.G., A.T., F.H.; Writing – review & editing: I.S.G., L.M.G., K.B., C.H., E.Po., L.S., I.S.F., C.St., C.So, M.B., P.B., M.G., M.C.B., E.Pu., M.H., P.A., I.B., H.B., G.C., M.D., S.T.M., M.F., A.T., F.H.

Declaration of interests

The authors declare no competing interests.

References

1236   **References**

1237

1.   Marshall, G. M. *et al.* The prenatal origins of cancer. *Nat Rev Cancer* **14**, 277–289 (2014).

2.   Pritchard-Jones, K. Genetics of childhood cancer. *Br Med Bull* **52**, 704–23 (1996).

3.   Scotting, P. J., Walker, D. A. & Perilongo, G. Childhood solid tumours: A developmental
disorder. *Nature Reviews Cancer* vol. 5 481–488 Preprint at https://doi.org/10.1038/nrc1633
(2005).

4.   Maris, J. M. & Denny, C. T. Focus on embryonal malignancies. *Cancer Cell* **2**, 447–50 (2002).

5.   Gröbner, S. N. *et al.* The landscape of genomic alterations across childhood cancers. *Nature*
555, 321–327 (2018).

6.   Ma, X. *et al.* Pan-cancer genome and transcriptome analyses of 1,699 paediatric leukaemias
and solid tumours. *Nature* **555**, 371–376 (2018).

7.   Matthay, K. K. *et al.* Neuroblastoma. *Nat Rev Dis Primers* **2**, 16078 (2016).

8.   Schulte, J. H. & Eggert, A. Neuroblastoma. *Crit Rev Oncog* **20**, 245–270 (2015).

9.   Zeineldin, M., Patel, A. G. & Dyer, M. A. Neuroblastoma: When differentiation goes awry.
Neuron **110**, 2916–2928 (2022).

10.   Nakagawara, A. Neural crest development and neuroblastoma: the genetic and biological
link. *Prog Brain Res* **146**, 231–242 (2004).

11.   Ponzoni, M. *et al.* Recent advances in the developmental origin of neuroblastoma: an
overview. *Journal of Experimental and Clinical Cancer Research* **41**, 1–28 (2022).

12.   Weiss, W. A., Aldape, K., Mohapatra, G., Feuerstein, B. G. & Bishop, J. M. Targeted expression
of MYCN causes neuroblastoma in transgenic mice. *EMBO Journal* **16**, 2985–2995 (1997).

13.   Cohen, M. A. *et al.* Formation of Human Neuroblastoma in Mouse-Human Neural Crest
Chimeras. *Cell Stem Cell* **26**, 579-592.e6 (2020).

14.   Olsen, R. R. *et al.* MYCN induces neuroblastoma in primary neural crest cells. *Oncogene* **36**,
5075–5082 (2017).

15.   Jansky, S. *et al.* Single-cell transcriptomic analyses provide insights into the developmental
origins of neuroblastoma. *Nat Genet* **53**, 683–693 (2021).

16.   Kameneva, P. *et al.* Single-cell transcriptomics of human embryos identifies multiple
sympathoblast lineages with potential implications for neuroblastoma origin. *Nat Genet* **53**,
694–706 (2021).

17.   Dong, R. *et al.* Single-Cell Characterization of Malignant Phenotypes and Developmental
Trajectories of Adrenal Neuroblastoma. *Cancer Cell* **38**, 716-733.e6 (2020).

18.   Hanemaaijer, E. S. *et al.* Single-cell atlas of developing murine adrenal gland reveals relation
of Schwann cell precursor signature to neuroblastoma phenotype. *Proc Natl Acad Sci U S A*
118, (2021).

19.   Kildisiute, G. *et al.* Tumor to normal single-cell mRNA comparisons reveal a pan-
neuroblastoma cancer cell. *Sci Adv* **7**, eabd3311 (2021).

20. de Preter, K. *et al.* Human fetal neuroblast and neuroblastoma transcriptome analysis
confirms neuroblast origin and highlights neuroblastoma candidate genes. *Genome Biol* **7**, 1–
17 (2006).

21. Althoff, K. *et al.* A Cre-conditional MYCN-driven neuroblastoma mouse model as an improved
tool for preclinical studies. *Oncogene* **34**, 3357–3368 (2015).

22. Molenaar, J. J. *et al.* Sequencing of neuroblastoma identifies chromothripsis and defects in
neuritogenesis genes. *Nature* **483**, 589–593 (2012).

23. Fetahu, I. S. *et al.* Single-cell transcriptomics and epigenomics unravel the role of monocytes
in neuroblastoma bone marrow metastasis. *Nat Commun* **14**, 3620 (2023).

24. Pugh, T. J. *et al.* The genetic landscape of high-risk neuroblastoma. *Nat Genet* **45**, 279–284
(2013).

25. Bown, N. *et al.* Gain of chromosome arm 17q and adverse outcome in patients with
neuroblastoma. *N Engl J Med* **340**, 1954–1961 (1999).

26. Gilbert, F. *et al.* Human neuroblastomas and abnormalities of chromosomes 1 and 17. *Cancer
Res* **44**, 5444–9 (1984).

27. Lastowska, M. *et al.* Comprehensive genetic and histopathologic study reveals three types of
neuroblastoma tumors. *J Clin Oncol* **19**, 3080–3090 (2001).

28. Mazzocco, K. *et al.* Genetic abnormalities in adolescents and young adults with
neuroblastoma: A report from the Italian Neuroblastoma Group. *Pediatr Blood Cancer* **62**,
1725–1732 (2015).

29. Körber, V. *et al.* Neuroblastoma arises in early fetal development and its evolutionary
duration predicts outcome. *Nat Genet* **55**, 619–630 (2023).

30. Bogen, D. *et al.* The genetic tumor background is an important determinant for
heterogeneous MYCN-amplified neuroblastoma. *Int J Cancer* **139**, 153–163 (2016).

31. O'Neill, S. *et al.* MYCN amplification and 17q in neuroblastoma: evidence for structural
association. *Genes Chromosomes Cancer* **30**, 87–90 (2001).

32. Huang, M. & Weiss, W. A. Neuroblastoma and MYCN. *Cold Spring Harb Perspect Med* **3**,
(2013).

33. Schleiermacher, G. *et al.* Accumulation of segmental alterations determines progression in
neuroblastoma. *J Clin Oncol* **28**, 3122–3130 (2010).

34. Schleiermacher, G. *et al.* Segmental chromosomal alterations have prognostic impact in
neuroblastoma: a report from the INRG project. *Br J Cancer* **107**, 1418 (2012).

35. Frith, T. J. R. *et al.* Human axial progenitors generate trunk neural crest cells in vitro. *Elife* **7**,
(2018).

36. Frith, T. J. R. & Tsakiridis, A. Efficient Generation of Trunk Neural Crest and Sympathetic
Neurons from Human Pluripotent Stem Cells Via a Neuromesodermal Axial Progenitor
Intermediate. *Curr Protoc Stem Cell Biol* **49**, e81 (2019).

37. Wymeersch, F. J., Wilson, V. & Tsakiridis, A. Understanding axial progenitor biology in vivo
and in vitro. *Development* **148**, (2021).

38. Thomson, J. A. *et al.* Embryonic stem cell lines derived from human blastocysts. *Science* **282**,
1145–7 (1998).

39. Soldatov, R. *et al.* Spatiotemporal structure of cell fate decisions in murine neural crest.
Science (1979) **364**, (2019).

40. Kastriti, M. E. *et al.* Schwann cell precursors represent a neural crest-like state with biased
multipotency. *EMBO J* **41**, e108780 (2022).

41. Pattyn, A., Morin, X., Cremer, H., Goridis, C. & Brunet, J. F. Expression and interactions of the
two closely related homeobox genes Phox2a and Phox2b during neurogenesis. *Development*
124, 4065–4075 (1997).

42. Xie, M. *et al.* Schwann cell precursors contribute to skeletal formation during embryonic
development in mice and zebrafish. *Proc Natl Acad Sci U S A* **116**, 15068–15073 (2019).

43. McGonnell, I. M. & Graham, A. Trunk neural crest has skeletogenic potential. *Curr Biol* **12**,
767–771 (2002).

44. Boeva, V. *et al.* Heterogeneity of neuroblastoma cell identity defined by transcriptional
circuitries. *Nat Genet* **49**, 1408–1413 (2017).

45. Van Groningen, T. *et al.* Neuroblastoma is composed of two super-enhancer-associated
differentiation states. *Nat Genet* **49**, 1261–1266 (2017).

46. van Groningen, T. *et al.* A NOTCH feed-forward loop drives reprogramming from adrenergic
to mesenchymal state in neuroblastoma. *Nat Commun* **10**, 1530 (2019).

47. Ciccarone, V., Spengler, B. A., Meyers, M. B., Biedler, J. L. & Ross, R. A. Phenotypic
diversification in human neuroblastoma cells: expression of distinct neural crest lineages.
Cancer Res **49**, 219–25 (1989).

48. Biedler, J. L., Helson, L. & Spengler, B. A. Morphology and growth, tumorigenicity, and
cytogenetics of human neuroblastoma cells in continuous culture. *Cancer Res* **33**, 2643–52
(1973).

49. Halliwell, J., Barbaric, I. & Andrews, P. W. Acquired genetic changes in human pluripotent
stem cells: origins and consequences. *Nature Reviews Molecular Cell Biology 2020 21:12* **21**,
715–728 (2020).

50. Draper, J. S. *et al.* Recurrent gain of chromosomes 17q and 12 in cultured human embryonic
stem cells. *Nat Biotechnol* **22**, 53–54 (2004).

51. Abbasi, M. R. *et al.* Impact of Disseminated Neuroblastoma Cells on the Identification of the
Relapse-Seeding Clone. *Clin Cancer Res* **23**, 4224–4232 (2017).

52. Kerosuo, L. *et al.* Enhanced expression of MycN/CIP2A drives neural crest toward a neural
stem cell-like fate: Implications for priming of neuroblastoma. *Proc Natl Acad Sci U S A* **115**,
E7351–E7360 (2018).

53. Girish, V. *et al.* Oncogene-like addiction to aneuploidy in human cancers. *Science* **381**,
eadg4521 (2023).

54. Mitchell, P. J., Timmons, P. M., Hébert, J. M., Rigby, P. W. J. & Tjian, R. Transcription factor
AP-2 is expressed in neural crest cell lineages during mouse embryogenesis. *Genes Dev* **5**,
105–119 (1991).

55. Rothstein, M. & Simoes-Costa, M. Heterodimerization of TFAP2 pioneer factors drives
epigenomic remodeling during neural crest specification. *Genome Res* **30**, 35–48 (2020).

56. Howard, A. G. A. & Uribe, R. A. Hox proteins as regulators of extracellular matrix interactions
during neural crest migration. *Differentiation* **128**, 26–32 (2022).

57. Akkermans, O. *et al.* GPC3-Unc5 receptor complex structure and role in cell migration. *Cell*
185, 3931-3949.e26 (2022).

58. TeSlaa, J. J., Keller, A. N., Nyholm, M. K. & Grinblat, Y. Zebrafish Zic2a and Zic2b regulate
neural crest and craniofacial development. *Dev Biol* **380**, 73–86 (2013).

59. Handel, A. E. *et al.* Developmental dynamics of the neural crest–mesenchymal axis in creating
the thymic microenvironment. *Sci Adv* **8**, 9844 (2022).

60. Rada-Iglesias, A. *et al.* Epigenomic annotation of enhancers predicts transcriptional regulators
of human neural crest. *Cell Stem Cell* **11**, 633–648 (2012).

61. Adam, K., Lesperance, J., Hunter, T. & Zage, P. E. The Potential Functional Roles of NME1
Histidine Kinase Activity in Neuroblastoma Pathogenesis. *Int J Mol Sci* **21**, (2020).

62. Ryl, T. *et al.* Cell-Cycle Position of Single MYC-Driven Cancer Cells Dictates Their Susceptibility
to a Chemotherapeutic Drug. *Cell Syst* **5**, 237-250.e8 (2017).

63. Lutz, W. *et al.* Conditional expression of N-myc in human neuroblastoma cells increases
expression of alpha-prothymosin and ornithine decarboxylase and accelerates progression
into S-phase early after mitogenic stimulation of quiescent cells. *Oncogene* **13**, 803–12
(1996).

64. King, D. *et al.* MYCN expression induces replication stress and sensitivity to PARP inhibition in
neuroblastoma. *Oncotarget* **11**, 2141 (2020).

65. Kramer, M., Ribeiro, D., Arsenian-Henriksson, M., Deller, T. & Rohrer, H. Proliferation and
Survival of Embryonic Sympathetic Neuroblasts by MYCN and Activated ALK Signaling. *J
Neurosci* **36**, 10425–10439 (2016).

66. Mobley, B. C. *et al.* Expression of MYCN in Multipotent Sympathoadrenal Progenitors Induces
Proliferation and Neural Differentiation, but Is Not Sufficient for Tumorigenesis. *PLoS One* **10**,
e0133897 (2015).

67. Berry, T. *et al.* The ALK(F1174L) mutation potentiates the oncogenic activity of MYCN in
neuroblastoma. *Cancer Cell* **22**, 117–130 (2012).

68. Olsen, R. R. *et al.* MYCN induces neuroblastoma in primary neural crest cells. *Oncogene* **36**,
5075–5082 (2017).

69. Cohen, M. A. *et al.* Formation of Human Neuroblastoma in Mouse-Human Neural Crest
Chimeras. *Cell Stem Cell* **26**, 579-592.e6 (2020).

70. Huang, M. *et al.* Engineering Genetic Predisposition in Human Neuroepithelial Stem Cells
Recapitulates Medulloblastoma Tumorigenesis. *Cell Stem Cell* **25**, 433-446.e7 (2019).

71. Wrobel, J. K. *et al.* Rapid In Vivo Validation of HDAC Inhibitor-Based Treatments in
Neuroblastoma Zebrafish Xenografts. *Pharmaceuticals (Basel)* **13**, 345 (2020).

72. Su, Z. *et al.* A comprehensive assessment of RNA-seq accuracy, reproducibility and
information content by the Sequencing Quality Control Consortium. *Nat Biotechnol* **32**, 903–
914 (2014).

73. Wang, C. *et al.* The concordance between RNA-seq and microarray data depends on chemical
treatment and transcript abundance. *Nat Biotechnol* **32**, 926–932 (2014).

74. Durbin, A. D. *et al.* EP300 Selectively Controls the Enhancer Landscape of MYCN-Amplified
Neuroblastoma. *Cancer Discov* **12**, 730–751 (2022).

75. Gartlgruber, M. *et al.* Super enhancers define regulatory subtypes and cell identity in
neuroblastoma. *Nat Cancer* **2**, 114–128 (2021).

76. Zeid, R. *et al.* Enhancer invasion shapes MYCN-dependent transcriptional amplification in
neuroblastoma. *Nat Genet* **50**, 515–523 (2018).

77. Jahangiri, L. *et al.* Core regulatory circuitries in defining cancer cell identity across the
malignant spectrum. *Open Biol* **10**, 200121 (2020).

78. Zimmerman, M. W. *et al.* Retinoic acid rewires the adrenergic core regulatory circuitry of
childhood neuroblastoma. *Sci Adv* **7**, (2021).

79. Corces, M. R. *et al.* An improved ATAC-seq protocol reduces background and enables
interrogation of frozen tissues. *Nat Methods* **14**, 959–962 (2017).

80. Meuleman, W. *et al.* Index and biological spectrum of human DNase I hypersensitive sites.
Nature 2020 584:7820 **584**, 244–251 (2020).

81. Zhang, K. *et al.* A single-cell atlas of chromatin accessibility in the human genome. *Cell* **184**,
5985-6001.e19 (2021).

82. Gao, T. & Qian, J. EnhancerAtlas 2.0: an updated resource with enhancer annotation in 586
tissue/cell types across nine species. *Nucleic Acids Res* **48**, D58–D64 (2020).

83. Moerman, T. *et al.* GRNBoost2 and Arboreto: Efficient and scalable inference of gene
regulatory networks. *Bioinformatics* **35**, 2159–2161 (2019).

84. Moreno-Smith, M. *et al.* Restoration of the molecular clock is tumor suppressive in
neuroblastoma. *Nature Communications 2021 12:1* **12**, 1–16 (2021).

85. Altman, B. J. *et al.* MYC Disrupts the Circadian Clock and Metabolism in Cancer Cells. *Cell
Metab* **22**, 1009–1019 (2015).

86. Xue, C. *et al.* MYCN promotes neuroblastoma malignancy by establishing a regulatory circuit
with transcription factor AP4. *Oncotarget* **7**, 54937–54951 (2016).

87. Boboila, S. *et al.* Transcription factor activating protein 4 is synthetically lethal and a master
regulator of MYCN-amplified neuroblastoma. *Oncogene* **37**, 5451–5465 (2018).

88. Ana Paula Azambuja, A., Paula Azambuja, A. & Simoes-Costa, M. The connectome of neural
crest enhancers reveals regulatory features of signaling systems. *Dev Cell* **56**, 1268-1282.e6
(2021).

89. Gundem, G. *et al.* Clonal evolution during metastatic spread in high-risk neuroblastoma. *Nat Genet* **55**, 1022–1033 (2023).

90. Merrill, B. J., Gat, U., DasGupta, R. & Fuchs, E. Tcf3 and Lef1 regulate lineage differentiation of multipotent stem cells in skin. *Genes Dev* **15**, 1688–1705 (2001).

91. Gautier, M., Thirant, C., Delattre, O. & Janoueix-Lerosey, I. Plasticity in neuroblastoma cell identity defines a noradrenergic-to-mesenchymal transition (Nmt). *Cancers (Basel)* **13**, (2021).

92. Thirant, C. *et al.* Reversible transitions between noradrenergic and mesenchymal tumor identities define cell plasticity in neuroblastoma. *Nat Commun* **14**, (2023).

93. Vayani, O. R. *et al.* Adrenergic and mesenchymal signatures are identifiable in cell-free DNA and correlate with metastatic disease burden in children with neuroblastoma. *Pediatr Blood Cancer* **71**, (2024).

94. Durbin, A. D. *et al.* Selective gene dependencies in MYCN-amplified neuroblastoma include the core transcriptional regulatory circuitry. *Nature Genetics* vol. 50 1240–1246 Preprint at https://doi.org/10.1038/s41588-018-0191-z (2018).

95. Rouhani, F. J. *et al.* Substantial somatic genomic variation and selection for BCOR mutations in human induced pluripotent stem cells. *Nature Genetics 2022 54:9* **54**, 1406–1416 (2022).

96. Puigdevall, P., Jerber, J., Danecek, P., Castellano, S. & Kilpinen, H. Somatic mutations alter the differentiation outcomes of iPSC-derived neurons. *Cell Genomics* **3**, 100280 (2023).

97. Aschero, R. *et al.* Recurrent Somatic Chromosomal Abnormalities in Relapsed Extraocular Retinoblastoma. *Cancers 2021, Vol. 13, Page 673* **13**, 673 (2021).

98. Lazic, D. *et al.* Landscape of Bone Marrow Metastasis in Human Neuroblastoma Unraveled by Transcriptomics and Deep Multiplex Imaging. *Cancers (Basel)* **13**, 4311 (2021).

99. Verhoeven, B. M. *et al.* The immune cell atlas of human neuroblastoma. *Cell Rep Med* **3**, 100657 (2022).

100. Costa, A. *et al.* Single-cell transcriptomics reveals shared immunosuppressive landscapes of mouse and human neuroblastoma. *J Immunother Cancer* **10**, (2022).

101. Weiss, T. *et al.* Schwann cell plasticity regulates neuroblastic tumor cell differentiation via epidermal growth factor-like protein 8. *Nat Commun* **12**, (2021).

102. Ludwig, T. E. *et al.* ISSCR standards for the use of human stem cells in basic research. *Stem Cell Reports* **18**, 1744 (2023).

103. Price, C. J. *et al.* Genetically variant human pluripotent stem cells selectively eliminate wild-type counterparts through YAP-mediated cell competition. *Dev Cell* **56**, 2455-2470.e10 (2021).

104. Randolph, L. N., Bao, X., Zhou, C. & Lian, X. An all-in-one, Tet-On 3G inducible PiggyBac system for human pluripotent stem cells and derivatives. *Sci Rep* **7**, (2017).

105. Loew, R., Heinz, N., Hampf, M., Bujard, H. & Gossen, M. Improved Tet-responsive promoters with minimized background expression. *BMC Biotechnol* **10**, 1–13 (2010).

1464   106.   Zhou, X., Vink, M., Klaver, B., Berkhout, B. & Das, A. T. Optimization of the Tet-On system for
1465   regulated gene expression through viral evolution. *Gene Ther* **13**, 1382–1390 (2006).

1466   107.   Laing, O., Halliwell, J. & Barbaric, I. Rapid PCR Assay for Detecting Common Genetic Variants
1467   Arising in Human Pluripotent Stem Cell Cultures. *Curr Protoc Stem Cell Biol* **49**, e83 (2019).

1468   108.   Chen, G. *et al.* Chemically defined conditions for human iPSC derivation and culture. *Nat
1469   Methods* **8**, 424–429 (2011).

1470   109.   Sturtzel, C. *et al.* Refined high-content imaging-based phenotypic drug screening in zebrafish
1471   xenografts. *npj Precision Oncology 2023 7:1* **7**, 1–16 (2023).

1472   110.   Hanssen, F. *et al.* Scalable and efficient DNA sequencing analysis on different compute
1473   infrastructures aiding variant discovery. *bioRxiv* **23**, 2023.07.19.549462 (2023).

1474   111.   Garcia, M. *et al.* Sarek: A portable workflow for whole-genome sequencing analysis of
1475   germline and somatic variants. *F1000Research 2020 9:63* **9**, 63 (2020).

1476   112.   Van der Auwera, G., O'Connor, B. & Safari, an O. M. Company. Genomics in the Cloud: Using
1477   Docker, GATK, and WDL in Terra. *Genomics in the Cloud* 300 (2020).

1478   113.   Kim, S. *et al.* Strelka2: fast and accurate calling of germline and somatic variants. *Nat
1479   Methods* **15**, 591–594 (2018).

1480   114.   Chen, X. *et al.* Manta: rapid detection of structural variants and indels for germline and
1481   cancer sequencing applications. *Bioinformatics* **32**, 1220–1222 (2016).

1482   115.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, (2021).

1483   116.   McLaren, W. *et al.* The Ensembl Variant Effect Predictor. *Genome Biol* **17**, 1–14 (2016).

1484   117.   Berlanga, P. *et al.* The European MAPPYACTS Trial: Precision Medicine Program in Pediatric
1485   and Adolescent Patients with Recurrent Malignancies. *Cancer Discov* **12**, 1266–1281 (2022).

1486   118.   Adzhubei, I., Jordan, D. M. & Sunyaev, S. R. Predicting functional effect of human missense
1487   mutations using PolyPhen-2. *Curr Protoc Hum Genet* **Chapter 7**, (2013).

1488   119.   Ng, P. C. & Henikoff, S. SIFT: Predicting amino acid changes that affect protein function.
1489   *Nucleic Acids Res* **31**, 3812–3814 (2003).

1490   120.   Favero, F. *et al.* Sequenza: allele-specific copy number and mutation profiles from tumor
1491   sequencing data. *Ann Oncol* **26**, 64–70 (2015).

1492   121.   Gu, Z., Eils, R. & Schlesner, M. Complex heatmaps reveal patterns and correlations in
1493   multidimensional genomic data. *Bioinformatics* **32**, 2847–2849 (2016).

1494   122.   McKenna, A. *et al.* The genome analysis toolkit: A MapReduce framework for analyzing next-
1495   generation DNA sequencing data. *Genome Res* **20**, 1297–1303 (2010).

1496   123.   Schliep, K. P. phangorn: phylogenetic analysis in R. *Bioinformatics* **27**, 592–593 (2011).

1497   124.   Jombart, T., Balloux, F. & Dray, S. adephylo: new tools for investigating the phylogenetic
1498   signal in biological traits. *Bioinformatics* **26**, 1907–1909 (2010).

125. Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. Y. ggtree: an r package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods*
Ecol Evol **8**, 28–36 (2017).

126. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. *Use R! series* (Springer, 2016).

127. McGinnis, C. S. *et al.* MULTI-seq: sample multiplexing for single-cell RNA sequencing using
lipid-tagged indices. *Nat Methods* **16**, 619–626 (2019).

128. Stuart, T. *et al.* Comprehensive integration of single-cell data. *Cell* **177**, 1888-1902.e21 (2019).

129. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29
(2021).

130. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol* **36**,
411–420 (2018).

131. Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-seq
data using regularized negative binomial regression. *Genome Biol* **20**, 576827 (2019).

132. Lun, A. T. L. *et al.* EmptyDrops: Distinguishing cells from empty droplets in droplet-based
single-cell RNA sequencing data. *Genome Biol* **20**, 1–9 (2019).

133. Germain, P. L., Robinson, M. D., Lun, A., Garcia Meixide, C. & Macnair, W. Doublet
identification in single-cell sequencing data using *scDblFinder*. *F1000Research 2022 10:979*
10, 979 (2022).

134. Zhu, Q., Conrad, D. N. & Gartner, Z. J. deMULTIplex2: robust sample demultiplexing for
scRNA-seq. *bioRxiv* 2023.04.11.536275 (2023) doi:10.1101/2023.04.11.536275.

135. Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative modeling for single-
cell transcriptomics. *Nature Methods 2018 15:12* **15**, 1053–1058 (2018).

136. Hafemeister, C. & Halbritter, F. Single-cell RNA-seq differential expression tests within a
sample should use pseudo-bulk data of pseudo-replicates. *bioRxiv* 2023.03.28.534443 (2023)
doi:10.1101/2023.03.28.534443.

137. Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set collection.
Cell Syst **1**, 417 (2015).

138. Federico, A. & Monti, S. hypeR: an R package for geneset enrichment workflows.
Bioinformatics **36**, 1307–1308 (2020).

139. Street, K. *et al.* Slingshot: Cell lineage and pseudotime inference for single-cell
transcriptomics. *BMC Genomics* **19**, 477 (2018).

140. van den Berge, K. *et al.* Trajectory-based differential expression analysis for single-cell
sequencing data. *Nat Commun* **11**, 1–13 (2020).

141. Lambert, S. A. *et al.* The Human Transcription Factors. *Cell* **172**, 650–665 (2018).

142. Patel, A. P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in primary
glioblastoma. *Science (1979)* **344**, 1396–1401 (2014).

143.   Rifatbegovic, F. *et al.* Neuroblastoma cells undergo transcriptomic alterations upon
1537   dissemination into the bone marrow and subsequent tumor progression. *Int J Cancer* **142**,
1538   297–307 (2018).

1539   144.   Ladenstein, R. *et al.* [Value of prognostic factors in the Austrian A-NB87 Neuroblastoma
1540   Study]. *Klin Padiatr* **208**, 210–20 (1996).

1541   145.   Fiedler, S. *et al.* Long-Term Outcome and Role of Biology within Risk-Adapted Treatment
1542   Strategies: The Austrian Neuroblastoma Trial A-NB94. *Cancers 2021, Vol. 13, Page 572* **13**,
1543   572 (2021).

1544   146.   Garaventa, A. *et al.* Randomized Trial of Two Induction Therapy Regimens for High-Risk
1545   Neuroblastoma: HR-NBL1.5 International Society of Pediatric Oncology European
1546   Neuroblastoma Group Study. *J Clin Oncol* **39**, 2552–2563 (2021).

1547   147.   Ladenstein, R. *et al.* Investigation of the Role of Dinutuximab Beta-Based Immunotherapy in
1548   the SIOPEN High-Risk Neuroblastoma 1 Trial (HR-NBL1). *Cancers (Basel)* **12**, (2020).

1549   148.   Ladenstein, R. *et al.* Interleukin 2 with anti-GD2 antibody ch14.18/CHO (dinutuximab beta) in
1550   patients with high-risk neuroblastoma (HR-NBL1/SIOPEN): a multicentre, randomised, phase
1551   3 trial. *Lancet Oncol* **19**, 1617–1629 (2018).

1552   149.   Ladenstein, R. *et al.* Busulfan and melphalan versus carboplatin, etoposide, and melphalan as
1553   high-dose chemotherapy for high-risk neuroblastoma (HR-NBL1/SIOPEN): an international,
1554   randomised, multi-arm, open-label, phase 3 trial. *Lancet Oncol* **18**, 500–514 (2017).

1555   150.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
1556   RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

1557   151.   Hänzelmann, S., Castelo, R. & Guinney, J. GSVA: Gene set variation analysis for microarray
1558   and RNA-Seq data. *BMC Bioinformatics* **14**, 1–15 (2013).

1559   152.   Smith, J. P. *et al.* PEPATAC: an optimized pipeline for ATAC-seq data analysis with serial
1560   alignments. *NAR Genom Bioinform* **3**, lqab101 (2021).

1561   153.   Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
1562   assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

1563   154.   Lawrence, M. *et al.* Software for Computing and Annotating Genomic Ranges. *PLoS Comput
1564   Biol* **9**, e1003118 (2013).

1565   155.   Kulakovskiy, I. v. *et al.* HOCOMOCO: Towards a complete collection of transcription factor
1566   binding models for human and mouse via large-scale ChIP-Seq analysis. *Nucleic Acids Res* **46**,
1567   D252–D259 (2018).

1568   156.   Korotkevich, G. *et al.* Fast gene set enrichment analysis. *bioRxiv* 060012 (2021)
1569   doi:10.1101/060012.

1570   157.   Upton, K. *et al.* Epigenomic profiling of neuroblastoma cell lines. *Sci Data* **7**, (2020).

1571   158.   Decaesteker, B. *et al.* SOX11 regulates SWI/SNF complex components as member of the
1572   adrenergic neuroblastoma core regulatory circuitry. *Nature Communications 2023 14:1* **14**, 1–
1573   16 (2023).

159. Xu, D. *et al.* Recapitulation of patient-specific 3D chromatin conformation using machine
learning. *Cell reports methods* **3**, (2023).

160. Hariprakash, J. M. *et al.* Leveraging Tissue-Specific Enhancer-Target Gene Regulatory
Networks Identifies Enhancer Somatic Mutations That Functionally Impact Lung Cancer.
Cancer Res **84**, (2024).

161. Boix, C. A., James, B. T., Park, Y. P., Meuleman, W. & Kellis, M. Regulatory genomic circuitry of
human disease loci by integrative epigenomics. *Nature 2021 590:7845* **590**, 300–307 (2021).

162. Lee, S., Cook, D. & Lawrence, M. Plyranges: A grammar of genomic data transformation.
Genome Biol **20**, (2019).

163. La Manno, G. *et al.* RNA velocity of single cells. *Nature* **560**, 494–498 (2018).

1584
1585

Figures

Figure 1

Figure 1. *In vitro* culture efficiently generates human trunk NC cells and their sympathoadrenal derivatives from hESCs.

a) Diagram depicting the extrinsically supplemented signals employed to direct hESCs toward trunk NC cells and their downstream derivatives, and immunofluorescence analysis of PERIPHERIN protein expression illustrating the generation of sympathetic neurons at D19. Cell nuclei were counterstained using Hoechst 33342.

b) UMAP of scRNA-seq data from wild-type hESCs collected at 9 stages (indicated by different colours) of differentiation to trunk neural crest and sympathoadrenal derivatives. Cells were divided into 14 distinct clusters as indicated by the contours.

c) Heatmap of gene markers for each cluster in **panel b**. Selected genes have been highlighted and UMAPs indicate the expression level of canonical markers for stem (*POU5F1*), neural crest (*SOX10*), mesenchymal (*FN1*), and sympathetic (*PHOX2B*) cells. All marker genes are reported in **Supplementary Table 2**.

d) Cells from D9-D28 of **panel b** labelled by their closest matching cell type from the human embryonic adrenal gland reference[16] via label transfer. Cells in grey could not be verified with markers (**Supplementary Fig. 2i**) or could not be assigned to a single type.

e) Cells from **panel d** coloured by the strength of their SCP marker signature (Seurat module score) in red. A high / medium score distinguishes a cluster of early SCP-like / trunk NC and a late cluster with more mature SCP-like cells.

f) Same as above but visualising simultaneously SYM (orange) and MES (teal) marker signature. Cells with overlapping marker signatures appear in grey tones, with the highest mixture in C12. An early diverging cluster of sensory neuron-like cells has a weak match to the SYM signature. A pseudotime trajectory for the MES-SYM transition in clusters C11-C14 can be found in **Supplementary Figure 3**.

1614   **Abbreviations**: hESC, human embryonic stem cells; D0/3/9/10/12/14/19/28, day 0/3/9/10/12/14/19/28;
1615   UMAP, Uniform Manifold Approximation and Projection; C1-C14, cell clusters; SCP, Schwann cell
1616   precursor; SYM, sympathoblast; MES, mesenchymal.

Figure 2

Figure 2. Copy number alterations and overexpression of *MYCN* impair the specification of trunk
NC derivatives.

a) Scheme depicting the different hESC genetic backgrounds employed and the timing of DOX-
induced MYCN overexpression in the context of our trunk NC differentiation system.
b) scRNA-seq data from mutant cells (17q, 17q1q, 17q1qMYCN at D9, D14, and D19) were
mapped to the wild-type trunk NC reference (illustration on the left side). Glasswork UMAP
plots depicting the destination clusters in the WT reference for cells of the 17q, 17q1q and
17q1qMYCN conditions. Mutant cells are coloured by stage to emphasise mismatches with
WT. Fewer 17q1q and 17q1qMYCN cells map to later differentiation stages, highlighted by
arrows.
c) Alluvial plots comparing the stage at which each mutant cell was harvested versus its
phenotypically closest stage in the WT reference (based on label transfer as also shown in **panel**
b). In each subplot, the top bar indicates the proportion of cells collected at each stage (D9,
D14, D19). The bottom bar indicates the distribution of matching phenotypes in the WT for
that same population of cells, and streams indicate which subpopulations flow into cognate or
non-cognate WT stages. The plots suggest that cells from 17q1q and 17q1qMYCN
progressively mapped to earlier stages compared to WT.
d) Glasswork UMAPs of mapped 17q, 17q1q, and 17q1qMYCN cells (as in **panel b**) coloured by
closest-matching cell type in the human embryonic adrenal gland reference[16]. The category
"other" comprises other cell types in the reference dataset and mappings that could not be
validated by cell type markers (**Supplementary Fig. 2i**).
e) Percentage of cells mapped to each cell type in **panel d** split by cell line.
f) Violin plots indicating the strength of the SCP/SYM/MES (left to right) gene expression
signature (Seurat module score) for cells mapped to the respective cell type, split by cell line.
g) Plot indicating the change in mean expression (colour) and the percentage of cells expressing
the gene (size) for each gene in the signatures from **panel e** relative to WT. WT squares (size
= 1, white) are shown for reference.
h) Flow cytometric analysis of trunk NC markers HOXC9 and SOX10 in D9 cultures obtained
from hESCs marked by the indicated NB-associated lesions.

Abbreviations: WT, wild-type H7 hESCs; D0/3/9/10/12/14/19/28, day 0/3/9/10/12/14/19/28; UMAP,
Uniform Manifold Approximation and Projection; SCP, Schwann cell precursor; SYM, sympathoblast;
MES, mesenchymal.

Figure 3

a

Differential expression and pathway analysis (D9)

	WT vs. 17q	WT vs. 17q1q	WT vs. 17q1qMYCN	
MTORC1 signaling				chr17q
E2F targets				
Mitotic spindle				
P53 pathway				chr1q
Apoptosis				
Oxidative phosphorylation				Not in CNA
MYC targets V1				
MYC targets V2				
Mitotic spindle				
G2M checkpoint				

Gene list overlap (Jaccard index) −0.02 0.02

Up in WT
Up in 17q
Up in 17q1q
Up in 17q1qMYCN

b Genes on chr17q

MTORC1 signaling / E2F targets / Mitotic spindle / WT / 17q / 17q1q / 17q1qMYCN

FDXR ACACA ATP5MC1 NMT1 PSMD12 PSME3 SLC9A3R1 BRCA1 JPT1 KPNA2 LUC7L3 NME1 PPM1D RAD51C TK1 TUBG1 CDC27 CDC42EP4 CEP131 CSNK1D DYNLL2 NF1 SSH2 TBCD TUBD1

c Genes on chr1q

P53 pathway / Apoptosis / WT / 17q / 17q1q / 17q1qMYCN

COQ8A EPHX1 IER5 PHLDA3 PTPN14 RGS16 S100A4 SLC19A2 TNNI1 DAP3 LMNA MCL1 PEA15 PSEN2 ATF3 BTG2 TXNIP

Mean norm.
expression MUT-WT:
≥ 1
0.5
0
−0.5
≤ −1

d

Data integration and joint projection

Condition: WT / 17q / 17q1q / 17q1qMYCN

Label transfer: SCP-like / MES-like / SYM-like / Other / Chromaffin-like

UMAP2 / UMAP1

e

Mutation score m

$m_x = 0.0$

0.66 1.0
0.33
$m_y = 0.8$

0 m 1

D9/D14/D19 only

UMAP2 / UMAP1

f

m rank (D9)

chr1 chr17

D9_1 / D9_2 / D9_3 / D9_4

S100A4 / C1QBP / NPW / MYCN / KCNQ5
NPR3 / NR2F2 / TGFBR3 / EDNRB / NR2F1 / NECTIN3 / SFRP1 / ROBO1 / COL5A2 / ZFHX3 / SOX5 / ROR1 / KLF12 / NCAM1 / BACH2 / GFRA1 / BAZ2B / ACVR2A / FYN / SETBP1 / PRDM12 / TFAP2B / GFRA2 / SOX4 / MDGA1 / ADAM12 / ROBO2 / CADM1 / EBF1 / ISL1
HOXD3 / COL4A6 / GPC3 / APOC1 / ZIC2

Row-scaled expression
4
2
0
−2
−4

Genes:
Receptor
Ligand
TF

Figure 3. Copy number alterations and overexpression of *MYCN* alter the expression of metabolic and developmental pathways.

a) We performed differential expression analysis between WT and derivatives of each mutant hESC line at D9 of differentiation and performed a pathway enrichment analysis to summarise the differentially expressed genes (DEGs). Enrichment was determined by hypergeometric tests (*hypeR*[138], background: all detected genes in our scRNA-seq dataset; $P_{adj} \leq 0.1$) using pathways from MSigDB[137]. The overlap between up- and down-regulated DEGs with the pathway genes is indicated as a positive (green/orange/magenta colour bars) or negative (grey colour) number, respectively. We additionally distinguished between DEGs located on chromosome arms chr17q, chr1q, or anywhere else in the genome to analyse potential direct and indirect effects of CNAs (split from top to bottom). All differentially expressed genes and pathway enrichments are available in **Supplementary Tables 6 and 7**.

b) DEGs located on chromosome arm chr17q from the enriched pathways shown in **panel a**. The heatmap indicates the mean normalised expression difference between each indicated mutant cell line and WT (at D9). The annotation bar on top of the heatmap indicate membership (black colour) of genes in the selection pathways (MSigDB hallmark database).

c) As **panel (b)**, but for DEGs on chr1q and the respective enriched pathways.

d) UMAP of scRNA-seq data from wild-type and mutant hESCs (indicated by colour; cp. **Fig. 2a**) throughout differentiation to trunk neural crest and sympathoadrenal derivatives (top). Bottom: the same dataset coloured by closest-matching cell type in the human embryonic adrenal gland reference[16]. The category "other" comprises other cell types in the reference dataset and mappings that could not be validated by cell type markers (**Supplementary Fig. 2i**).

e) Illustration (top) of the calculation of mutation scores m (k-nearest neighbour (KNN) mutational average) as average score of each cell's neighbours. In this calculation, each neighbour weighs in by its cell line (0 = WT, 1/3 = 17q, 2/3 = 17q1q, 1 = 17q1qMYCN) such that the mutation score allows ordering cells from WT to MYCN mutation. Only cells from D9, D14, and D19 were used, for which data from all conditions were available. The actual scores are shown overlaid on the UMAP from **panel d** (bottom).

f) Heatmap showing the expression of top 300 genes highly correlated to the mutation score m from **panel e** across all cells from D9. Genes have been divided into four groups by hierarchical clustering, and selected TFs, receptors, and ligands are highlighted. All correlated genes are reported in **Supplementary Table 9**. Genes located on chr17q or chr1q are indicated.

Abbreviations: WT, wild-type H7 hESCs; D0/3/9/14/19, day 0/3/9/14/19; UMAP, Uniform Manifold Approximation and Projection; m, mutation score; TF, transcription factor.

Figure 4

Figure 4. Impaired trunk NC specification correlates with acquisition of tumourigenic hallmarks.

a) Representative brightfield images of D14 cultures following differentiation of hESCs with the
indicated genotypes.
b) Flow cytometric analysis of cell cycle in D9 cultures obtained from hESCs marked by the
indicated NB-associated lesions. Top: Representative FACS plots. Bottom: Percentage of cells

found in each of the different stages of the cell cycle (G1, S, G2) corresponding to indicated
NB-associated lesions (n = 3 biological replicates, error bars= standard deviation two-way
ANOVA). P values in comparisons: G1 (17q vs 17qMYCN, p = 0.0266= *; 17q1q vs
17q1qMYCN, p= 0.0153=*), S (17q vs 17qMYCN, p = 0.0233= *; 17q1q vs 17q1qMYCN, p=
0.0073=**). Note only comparisons examining the effect of MYCN overexpression in different
backgrounds are shown.

c)   Immunofluorescence analysis (green) of the expression of the cell proliferation marker KI-67
in D14 (left) cultures obtained from hESCs marked by the indicated NB-associated lesions.
Cell nuclei were counterstained using Hoechst 33342 (blue). Scoring of the percentages of KI-
67-positive cells is also shown (right) (n = 3 biological replicates, error bars= standard
deviation, ordinary two-way ANOVA with Tukey correction). P values in comparisons: 17q vs
17qMYCN, p=0.0078 =**; 17q1q vs. 17q1qMYCN p= 0.0001=***). Note only comparisons
examining the effect of MYCN overexpression in different backgrounds are shown.

d)   Left: Representative brightfield images of cell/colony morphology following a low-density
plating assay using cells marked by the indicated NB-associated lesions after 84 hours. Right:
Comparison of the number of colonies formed by cells marked by the indicated NB-associated
lesions following plating at low density. (n= 3 biological replicates, error bars= SD, Ordinary
Two-way ANOVA test with Tukey correction). P values in comparisons: 17q1q vs.
17q1qMYCN p= 0.0109=*. Note only comparisons examining the effect of MYCN
overexpression in different backgrounds are shown.

e)   Phylogenetic tree indicating the genetic relationship and distance (in number of SNVs detected
by whole-exome sequencing) between different hESC lines before (D0) and after differentiation
(D19). The shape of nodes indicates samples without a *MYCN* overexpression cassette (unfilled
circles), with an expression cassette which has not been activated (filled circles), and with an
activated (by addition of DOX from D5 onwards) *MYCN* overexpression cassette (filled
squares). The colours have been chosen to match those used elsewhere in the paper, without
specific meaning. The plot shows that few additional mutations occurred during differentiation
since the distances between differentiated cell lines and the shared ancestor with the matching
undifferentiated samples are small (<10 mutations). **Supplementary Tables 4 and 5** report
SNVs and CNAs identified in our analyses.

1724   **Abbreviations:** D0/5/14/19, day 0/5/14/19; WT, wild-type H7 hESCs.

Figure 5

Figure 5. hESC-derived trunk NC cells with CNAs form tumours in mice upon *MYCN* overexpression.

    a) Left: Representative images of subcutaneous xenografts of trunk NC cells derived from the indicated cell lines in the presence (17q1qMYCN) and absence (WT, 17q1q) of DOX treatment. Right: Graph showing tumour size growth per mouse corresponding to xenografts of indicated cell lines (n = 6 animals per cell line).

    b) Left: Representative MRI sections of mice at week 5 following xenografting of indicated cell lines in the adrenal gland and DOX treatment regimens. The white lines indicate the tumour perimeter. Right: Graph showing survival of the recipient animals after xenografting (n = 3 animals per cell line).

    c) Summary of mouse xenograft experiments.

Abbreviations: DOX, doxycycline; MRI, magnetic resonance imaging.

Figure 6

1739
1740
Figure 6. Comparison to hESC-based trunk NC differentiation resolves structured heterogeneity
across neuroblastoma tumours.
a)　Schematic overview of the analysis of tumour cells. We curated tumour cells from 10 *MYCN*-
amplified NB samples[15,17,23] from three studies and mapped them onto our reference (cp. **Fig.**
1)[129]. Mapping is represented as tumour cells falling into sectors of the WT *in vitro* reference
(depicted as contours of each cluster in the glasswork plot).
b)　Heatmap depicting gene expression in *MYCN*+ tumour cells of dataset *Jansky_NB14*[15]. Values
are inferCNV[142] copy number estimations per gene, relative to hematopoietic and immune cells
in the sample ordered by genomic position and chromosome (1-22). Cells (one per row) are
shown ordered by the respective cluster in our reference dataset that the tumour cells were
mapped to (C11 to C14) and therein by *MYCN* expression levels (depth-normalised sliding
window average, width = 20 cells). Mappings of other tumours datasets are shown in
Supplementary Fig. 11.
c)　Heatmap showing markers from gene expression signatures C4*, C5*, C9*, C13*, and C14*
(rows, top to bottom) in cells from 10 tumour datasets that were mapped to our in vitro reference
dataset (cp. **panel a**). Each gene expression signature is the intersection of the cluster markers
in our reference dataset (as in Fig. 1) and differentially expressed genes between the respective
tumour cells. No genes were found for C2* and C3*. Markers for tumour cells mapped to C2
and C3 showed no overlap with *in-vitro* cluster markers; thus, only mapped cells are shown.
All genes identified in this analysis are reported in **Supplementary Table 10**.
d)　Scatterplots evaluating the strength of gene expression signatures C5* and C13* (from **panel**
c; calculated using *GSVA*[151]) in NB bulk RNA-seq data from SEQC[72,73]. Each dot corresponds
to one tumour dataset coloured by *MYCN* amplification status (left) or clinical stage (right).
The density of points (kernel density estimate) in each group is indicated in the margins of the
plots.

e) Survival analysis for data from the SEQC cohort stratified by strength of the C13* expression signature. Groups were split by the median. Cox regression adjusted for age-group (< 18 months, 18-60 months, > 60 months), INSS stage 4 (yes / no), and *MYCN* amplification status (yes / no). n = 249 patients per group, or 136 [C13* low] and 47 [C13* high] events. All survival analysis results are reported in **Supplementary Table 11**.

Abbreviations: UMAP, Uniform Manifold Approximation and Projection; EFS, event-free survival.

Figure 7

Figure 7. Differentiation of wild-type and mutant hESCs is associated with epigenetic changes in nine distinct chromatin modules.

a) ATAC-seq read coverage for wild-type hESCs at three example loci. Each area plot reports the normalised read count aggregated per genomic bin (width = 500bp). Multiple semi-transparent area plots are overlaid for each replicate. Genes within each locus are shown on top with thin/thick lines indicating introns/exons. The arrows next to gene names indicate the direction of transcription. Selected peaks have been highlighted manually (dashed red lines).

b) ATAC-seq read coverage of wild-type and mutant hESCs at D19 near the *PHOX2B* locus. Plots as in **panel a**.

c) Principal component analysis of all ATAC-seq datasets, split into four panels by condition. The geometric means of all data belonging to the same stages are connected by arrows to visualise the stepwise chromatin changes during differentiation.

d) Euler diagram visualising the overlap of differentially accessible regions in mutant hESC-derived trunk NC derivatives across all stages compared to WT (*DEseq2*[150]; $P_{adj} \leq 0.005$,

| 1787 | $|\log_2\text{FoldChange}| \geq \log_2(1.5)$). Numbers indicate the total number of regions per cell line |
| 1788 | aggregated over all developmental stages. |

e) Heatmaps showing normalised read counts for all differentially accessible regions (columns) in any pairwise comparison of two stages or conditions (*DEseq2*[150]; $P_{adj} \leq 0.005$, $|\log_2\text{FoldChange}| \geq \log_2(1.5)$; $n_{total} = 45,583$). Regions have been divided into nine non-overlapping modules (R1–R9) by hierarchical clustering. Three annotation columns are shown to the right indicating regions called down- (blue) and up-regulated (red) in each mutant hESC. All regions and differential analysis results are reported in **Supplementary Tables 12 and 13**.

f) Comparison of regions belonging to the nine chromatin modules (from **panel e**) and proximal genes from the cluster markers lists defined in our scRNA-seq analysis (cp. **Fig. 1**). An enrichment analysis for the overlaps was performed using hypergeometric tests (using *hypeR*[138]; background: all genes associated with at lest one ATAC-seq peak) and the sizes and transparency of circles indicates the odds ratio and P-value, respectively. Significant results are indicated with filled circles ($P_{adj} \leq 0.005$). All results are shown in the figure and also reported in **Supplementary Table 14**.

g) Enrichment analysis for overlaps between chromatin modules and known TF motifs (*HOCOMOCO* database[155], v11). The plots are as in **panel f**, with the exception that only overlaps with $P_{adj} \leq 0.0000001$ and $|\log_2\text{FoldChange}| \geq \log_2(2)$ were marked as significant (background: all peaks with at least one motif match). The top results per module are shown and all results are reported in **Supplementary Table 14**.

h) Enrichment analysis of overlaps between regions belonging to the nine chromatin modules and super-enhancers specific to certain NB epigenetic subtypes[44,45,75] (background: all peaks with at least one overlapping region annotated in the super-enhancer analyses). Plots as in **panel f**.

Abbreviations: D0/3/9/14/19, day 0/3/9/14/19; WT, wild-type H7 hESCs; MUT, a "mutant" hESC line (17q, 17q1q, or 17q1qMYCN); R1-R9, chromatin region modules; sig., significant; ADR, adrenergic; MES, mesenchymal; MNA-HR, not *MYCN*-amplified high-risk; MNA-LR, not *MYCN*-amplified low-risk.

Figure 8

Figure 8. Copy number changes facilitate *MYCN*-mediated blockage of differentiation via developmental transcription factor networks.

a) To define putative target genes of TFs, we linked TF motifs identified in ATAC-seq peaks with proximal genes and additionally used the *GRNboost2* algorithm[83] to identify highly correlated TF-target gene candidates based on our scRNA-seq data.

b) Top 2500 targets of MYCN predicted by *GRNboost2*[83]. Putative targets without support in our ATAC-seq data (motif for TF in ≥1 peak near the gene) have been removed. We also calculated the Pearson correlation coefficient (*r*) between each TF and target gene to determine the direction of the putative interaction (*r*>0.1 = "activating", *r*<-0.1 = "inhibitory", others = "marginal"). The top 5 TFs have been highlighted. TF target gene sets are reported in **Supplementary Table 15**.

c) Average expression (Seurat module score) of the MYCN target gene set ("activated" targets from **panel d**) in our integrated scRNA-seq dataset (cp. **Fig. 3d**).

d) Heatmap displaying the percentage of genes in gene sets D9_1 to D9_4 (correlated with mutation score, cp. **Fig. 3e,f**) that overlapped with targets of the indicated TFs (one TF per column). All TF target sets with significant overlaps in at least one comparison are shown (hypergeometric test, *hypeR*[138]; $P_{adj} \leq 0.05$, $|log_2FoldChange| \geq log_2(4)$, frequency ≥ 5%). Enrichment results are also reported in **Supplementary Table 16.**

e) Smoothed line plots evaluating target gene expression (Seurat module score) for selected TFs from **panel f** with increasing mutations (mutation scores from **Figs. 3e,f**). We manually split

the TFs into groups corresponding to target genes losing or gaining expression along the module
score spectrum. The source cell line of each data point is indicated at the bottom.

f) Gene-regulatory network diagrams visualizing putative TF-to-target relations for enriched TF
targets (cp. **panels c-e**). In these diagrams, each node represents a TF or target gene, and each
edge is a link between a TF and a target. We made these networks specific to cells from each
condition (WT, 17q, 17q1q, 17q1qMYCN) by using colour to indicate the mean scaled
expression of each gene in the respective cells at D9 (edges coloured by source TF) and node
size to indicate the mean scaled T_{TF} target score of each TF. Only labels of TFs with positive
scaled expression are shown and selected groups of TFs have been merged for visualisation. A
network diagram with all node labels is shown in **Supplementary Fig. 13c**.

Abbreviations: D9, day 9; R1-R9, chromatin region modules; TF, transcription factor; WT, wild-type
H7 hESCs; sig., significant; r, Pearson correlation coefficient.

1848

1849 **Supplementary figures**

1850

Supplementary Figure 1

1851
1852 **Supplementary Figure 1 (related to Fig. 1). Immunofluorescence analysis of hESCs during trunk**
1853 **NC differentiation.**
1854 Immunofluorescence analysis of the expression of indicated markers at different time points during the
1855 differentiation of hESCs toward trunk NC and its derivatives.
1856 **Abbreviations**: D3/9/14, day 3/9/14.

Supplementary Figure 2

1857

Supplementary Figure 2 (related to Fig. 1). Quality control and reference mapping of single-cell RNA-seq data from wild-type hESC trunk neural crest differentiation.

a-d) UMAP plots showing quality covariates for the wild-type hESC dataset in **Figure 1**.

e-f) Histograms depicting the distribution of read depth (**e**) and number of genes detected (**f**) per cell barcode after quality control filtering.

g) Scatterplots comparing all quality control covariates (shown on **panels a-c**) from the same cell, plotted versus each other, coloured by 10x Genomics run (G1-G27; cp. **Fig. 1b**).

h) Bubble plot indicating the mean expression (colour) and fraction of cells expressing (size) each of the HOX genes per cluster. Genes have been ordered from cranial to sacral axis specification.

i) Side-to-side comparison of cell type marker expression in 200 annotated cells selected at random from the human adrenal gland reference[16] versus the top 200 high-confidence cells mapped to the same cell types in our WT *in vitro* trunk NC dataset. Rows are cell-type marker genes. Columns are cells first divided by cell type (separated with a grey line), then by dataset of origin (adrenal gland: purple, in vitro: orange). Expression values are depth-normalised per

experiment and row-scaled globally. Known stem cell markers were added to trace where the
stem cell population would be spuriously mapped to, in this case kidney, an indicator that cells
not found in the reference may be mapped to kidney. No *in vitro* cells were mapped to
chromaffin or liver identities, leading to the absence of the respective *in vitro* columns. Cells
erroneously mapped, absent from the reference, or lacking relevant cell type markers were
classified as "other" and coloured grey in **Figure 1e**.

j) Alluvial plots comparing the mappings between cells in the *in vitro* dataset compared to two
adrenal gland reference datasets[15,16]. Each "stream" indicates a group of cells that were mapped
to one cell identity in the Jansky et al. (middle) and the Kameneva *et al.* (right) references (also
indicated in colour). For example, cells that were labelled MES in Kameneva et al. (this is the
foundation for the cell type labels used in this paper) also mapped to mesenchymal cells in the
Jansky *et al.* reference, cells that mapped to SYM mapped to cycling neuroblasts and to
neuroblasts. Cells that we labelled as SCPs split into cells that mapped to mesenchymal cells
and late SCPs in Jansky *et al*., consistent with our observations that some of the "SCP-like"
cells in our dataset represent a less mature, early SCP state (see main text).

Abbreviations: D0/3/9/10/12/14/19/28, day 0/3/9/10/12/14/19/28; UMAP, Uniform Manifold
1888 Approximation and Projection; SCP, Schwann cell precursor; SYM, sympathoblast; MES,
1889 mesenchymal.

Supplementary Figure 3

Supplementary Figure 3 (related to Fig. 1). Trajectories connecting gradients of transcriptionally similar cells at different developmental stages.

a) *Slingshot*[139] pseudotime trajectories (top) for wild-type clusters C11-C14. Cells were extracted from the main dataset and reprocessed (see "Basic scRNA-seq processing" in Methods), and trajectories were calculated on the first two scVI components.

b) Heatmap showing the top 400 genes with the strongest association with the trajectory as ranked by *tradeSeq*'s Wald test[140] (based on a *fitGAM* model with 5 knots). Highlighted genes are all the transcription factors found in the association test. **Supplementary Table 3** reports all genes found in this analysis.

c) RNA velocities calculated for the cells in **Figure 1d** using Velocyto[163].

Abbreviations: principal component; SCP, Schwann cell precursor; SYM, sympathoblast; MES, mesenchymal.

Supplementary Figure 4

Supplementary Figure 4 (related to Fig. 2). Genetic and phenotypic characterization of mutant hESC lines.

a) Cytogenetic analysis of the H7-derived hESC lines used in the study.

b) Plots of the depth ratio calculated between each sample and the parental control using Sequenza[120]. CNAs can be seen for chr1q and chr17q, and a small segment which is lost on chr2 (q23.3-24.3). CNA coordinates and SNVs in these cell lines are reported in **Supplementary Tables 4 and 5**.

c) Pile-up of segmental gains (red) and losses (blue) on chromosome 1 (left) and 17 (right) based on SNP array data of 88 NB samples[51] (thin lines) compared to CNAs in our 17q and 17q1q(MYCN) hESC lines (from panel **b**). The X axis corresponds to chromosome position. Each horizontal line depicts all the aberrations found for the same patient. For compatibility

with available annotations, coordinates in this plot are shown with respect to human genome
reference GRCh37/hg19 (while all other analysis in this paper were done using hg38/GRCh38).
d,e) Analysis of MYCN expression at the transcript (**d**) and protein (**e**) level in D9 17q1qMYCN
cultures after DOX treatment at day 5 vs untreated control.
f) Comparison of *MYCN* expression in tumours and our engineered hESCs and their derivatives.
The plots on the left show bulk RNA-seq data from three NB tumour compendia (CCRI, SEQC,
TARGET) divided into cases with and without diagnosed *MYCN* amplification. The plot on the
right ("This study") shows our scRNA-seq data divided by cell line (across all timepoints). The
values in all plots are the $\log_2$ fold changes of over the mean of all non-amplified tumours (left
plots) or all WT cells (right plot).
g) Immunofluorescence analysis of PERIPHERIN expression in D19 cultures following
differentiation of hESCs with the indicated genotypes. Cell nuclei were counterstained using
Hoechst 33342.
h) Immunofluorescence analysis of PERIPHERIN expression in D28 cultures following
differentiation of 17q1q (No DOX) or 17q1qMYCN hESCs following DOX treatment at the
indicated timepoints. Cell nuclei were counterstained using Hoechst 33342.
Abbreviations: WT, wild-type H7 hESCs; CNA, copy number alteration; DOX, Doxycycline.

1933

Supplementary Figure 5

a

b

1934

Supplementary Figure 5 (related to Fig. 2). Plots illustrating the gating strategy for the FACS performed in this manuscript.

1935
1936
1937

a) Secondary-only (negative) gating was set using cells differentiated at the correspondent stage stained only with secondary antibodies (detailed in the methods section). The scatter plots of Forward Scatter (FSC) and Side scatter (SSC) variables were used as the method to identify the main cell population size. Once the main population was identified, SSC and Trigger Pulse Width variables were further used to determine the single-cell population (doublet discrimination). The FITC (488) and Red (647) channels were used to set the baseline of fluorescence (upper panel) and subsequent positive gates (lower panel).

b) Cells were analysed in the flow cytometer (BD FACSJazz) using the 405 nm laser to detect the Hoechst staining and 488 nm to detect the Edu staining.

1947

Supplementary Figure 6

1948
1949
1950 **Supplementary Figure 6 (related to Fig. 3). scRNA-seq analysis of differentiating wild-type and**
1951 **mutant hESCs.**
1952   **a)** Overview of the number of differentially expressed genes (DEGs) in 17q, 17q1q, and
1953   17q1qMYCN cells at D9 of differentiation compared to wild-type. The total number of DEGs
1954   is given (N), and the percentage of those genes that are located on chromosome arms chr17q or
1955   chr1q are indicated. Percentage values >5% have been highlighted (which also correspond to
1956   upregulated DEGs within known CNAs). DEGs are reported in **Supplementary Table 6**.
1957   **b)** Cell clusters defined for the full *in vitro* trunk NC dataset scRNA-seq dataset generated in this
1958   study. Cluster marker genes are reported in **Supplementary Table 8**.
1959   **c-h)** QC covariate plots: reads per cell (**panel ()**, features per cell (**panel d**), % mitochondrial genes
1960   (**panel e**), replicates (one of up to three repeat experiments; **panel f**), developmental stage
1961   (sampling day; **panel g**), and computationally inferred cell cycle stage (**panel h**).

1962 **Abbreviations**: WT, wild-type H7 hESCs; D0/3/9/14/19, day 0/3/9/14/19; UMAP, Uniform Manifold
1963 Approximation and Projection; SCP, Schwann cell precursor; SYM, sympathoblast; MES,
1964 mesenchymal.

1965

Supplementary Figure 7

Supplementary Figure 7 (related to Fig. 3). Differential gene expression in mutant hESC-derived trunk NC and sympathoadrenal cells.

a,b) Heatmaps containing the genes correlated or anti-correlated with the mutation score *m* and for D14 (**a**) and D19 (**b**). Left annotation indicates whether the gene is on chromosome 1q or 17q, respectively. Transcription factors (black), receptors (red) and ligands (blue) have been highlighted. All mutation-score-related genes are reported in **Supplementary Table 9**.

c) Heatmap showing the expression of mutation-score-related genes (**Fig. 3f**, **Supplementary Table 9**) in public bulk RNA-seq data from three NB tumour compendia (SEQC, TARGET, CCRI). The heatmaps display the row-normalised transcript counts per gene and sample. The INSS stage, *MYCN* amplification status, and sex of each sample are indicated by the colour bars on top. Genes in D9_1 that are highly expressed in *MYCN*-amplified tumours are highlighted.

Abbreviations: TF, transcription factor; m rank, mutation score rank.

Supplementary Figure 8

Supplementary Figure 8 (related to Figs. 2,3). Comparison of multiple mutant hESC lines with the WT trunk NC differentiation reference.

a) Analysis of MYCN expression at the transcript (top) and protein (bottom) level in D9 cultures of the indicated genotypes after DOX treatment from day 5 vs untreated control.
b) scRNA-seq data mapped to the wild-type trunk NC reference (from **Fig. 1**). Top: Example glasswork UMAP plots depicting the destination clusters in the WT reference for cells of the WT, 17q, and 17q1q cells (H7 cell line) with and without *MYCN* overexpression. Clusters with an increased (red) and decreased (blue) number of mapped cells are indicated. Bottom:

Heatmap displaying the mapping of all 16 datasets including 1-4 replicates of each cell line / experimental condition. Cells were derived from two parental hESC lines (H7 and H9), with/without gains of chr17q and/or chr1q, and edited with a *MYCN* expression cassette (which can be active [+DOX] or inactive [-DOX]). Clusters with increased/decreased numbers of mapped cells have been highlighted with red/blue boxes, respectively.

c) Barplots summarising the mappings from **panel b** for derivatives of each hESC line (top to bottom). Each bar indicates the proportion of cells (all collected at D9) that mapped to cells in the WT reference of a given developmental stage. The plots suggest that cells with *MYCN* induction mapped to earlier stages compared to WT.

d) Bubble plot showing marker genes of cells mapped to the WT cell clusters associated with the different genetic changes in **panel b** (C2, C4, C7 and C10). Only cells with positive percentages are shown. The size and colour of each circle indicate the percent of cells in the indicated group which express the gene and the average expression, respectively. Some gene expression programmes are affected by specific combinations of mutations and cell types, such as polyamine homeostasis gene *AZIN1* or P53 suppressor *PURPL* in 17qMYCN in C2 or C7.

e) Immunofluorescence analysis of PERIPHERIN (PRPH) expression in D19 cultures following differentiation of *MYCN*-overexpressing hESCs with the indicated genotypes in the presence and absence of DOX. Cell nuclei were counterstained using Hoechst 33342.

Abbreviations: DOX, Doxycycline.

2009
2010
Supplementary Figure 9 (related to Fig. 4). Apoptosis, DNA damage, and mutations in mutant
hESC-derived trunk NC differentiation.

a) Comparison of the number of colonies formed by D14 cells generated from unmodified WT
and 17q cells in the presence and absence of DOX treatment with or without 17q gain (n= 3
biological replicates, error bars= SD, Ordinary One-way ANOVA test with Tukey correction).
No statically significant differences were observed.

b) Time course flow cytometry analysis of cleaved Caspase-3 levels in differentiating trunk NC
cells generated from hESCs of the indicated genotypes in the presence and absence of DOX
treatment from day 5. Error bars= standard deviation, unpaired t-test). P values in comparisons:
WTMYCN (Day 11 DOX vs NO DOX, p = 0.010797 = *; Day 12 DOX vs NO DOX, p=
0.033974 = *), 17qMYCN (Day 12 DOX vs NO DOX, p = 0.034166 = *), 17q1qMYCN (Day
12 DOX vs NO DOX, p = 0.020679 = *; Day 13 DOX vs NO DOX, p = 0.005686 = **).

c) Control comparison of Caspase-3 levels in untreated and DOX-treated WT controls. As in
panel b.

d) Number of γH2AX foci per cell in differentiating cultures corresponding to indicated
timepoints and genotypes in the presence and absence of DOX treatment. FDR values in
comparisons (ANOVA with mixed effect model with negative binomial distribution): FDR ***
< 0.001, ** < 0.01, * < 0.05. Comparisons performed: DOX status within a genotype and day,
genotype vs WT per DOX status and day.

e) Control comparison of the number of γH2AX foci per cell in untreated and DOX-treated
unmodified WT and 17q cells. As in **panel d**.

f) Heatmap containing CNA calls by Sequenza[120] in bins of 1Mbp. Dendrogram represents hierarchical clustering. CNA coordinates are reported in **Supplementary Table 5**.

g) Phylogenetic distance measured in mutations (SNVs) of each sample relative to the parental control using the phylogenetic analysis presented in **Fig. 4e**.

Abbreviations: WT, wild-type H7 hESCs; DOX, Doxycycline.

Supplementary Figure 10

2037
2038
2039 **Supplementary Figure 10 (related to Fig. 5). Mouse and zebrafish xenografts of WT, 17q1q, and**
2040 **17q1qMYCN cells.**

2041 a) Representative sections from individual tumours following xenografting of 17q1qMYCN cells
2042 in the indicated locations and showing haematoxylin and eosin staining (top row), Ki67 (middle
2043 row) and MYCN expression (bottom row). Tumour sections from a Th-*MYCN* NB GEM model
2044 and negative controls are shown.
2045 b) Representative images of zebrafish xenografted with WT or 17q1qMYCN cells labelled with
2046 CellTrace Violet at one day or three days post injection (top and bottom, respectively).

c) Quantification of the area covered by WT or 17q1qMYCN cells in zebrafish xenografts at 1 and 3 dpi. While 17q1qMYCN cells persist, WT cells are not maintained in zebrafish xenografts. Xenografts with WT cells (n = 11), and 17q1qMYCN cells (n = 51).

d) Waterfall plots depicting the change in tumour area for 17q1qMYCN at D9 of differentiation (~ NC stage) and SK-N-BE2C-H2B-GFP[71] cells in zebrafish xenografts from 1dpi to 3dpi.

Abbreviations: dpi, day post injection.

Supplementary Figure 11 (related to Fig. 6). Analysis of *MYCN*-amplified tumour cells and mapping to *in vitro* trunk NC differentiation.

a) *inferCNV*[142] profile heatmaps such as the one in **Fig. 6b** for the remaining 9 tumour datasets[15,17,23] not shown in **Fig. 6**. Each row (tumour cells) and each column (genes, ordered by genomic position), indicate the intensity of the CNA signal relative to non-tumour, HSC/immune cells from the same sample. All samples were curated and processed as described in **Fig. 6** and mapped to our wild-type trunk NC differentiation reference (**Fig. 1**). Cells are ordered first by matching cluster and then by *MYCN* levels within each cluster. Annotation of the chromosomes can be found on top of the heatmap. The tumour type (primary, metastasis, relapse), tissue, chr17q/1q gain status, and sex of each sample are indicated. The asterisk (*) marks cases in which the chr17q/1q gain status is based on the inferred inferCNV copy number profile shown in this figure panel.

b) Bubble plot summarising the mapping of tumour cells to our in vitro trunk NC reference (from **panel a** and **Fig. 6b**). The tumour type (primary, metastasis, relapse), sex, and chr17q/1q gain status are indicated. Ordering by study suggests a study-specific effect; for example, Fetahu samples are all bone marrow metastases and lack mesenchymal C11 cells.

Abbreviations: CNA, copy number alteration; T, primary tumour; M, metastasis; R, relapse; M, male; F, female

Supplementary Figure 12

Supplementary Figure 12 (related to Fig. 7). Chromatin accessibility in differentiating wild-type and mutant hESCs.

a) Pie charts indicating the fraction of differentially accessible regions belonging to chromatin modules (**Fig. 7e**) in mutant hESCs compared to WT (left to right). Up- and down-regulated regions are shown separately (top vs. bottom row). The total number (*n*) of regions in each category is indicated.

b) Barplots indicating the number of up- (positive numbers) and down-regulated (negative numbers) peaks from each comparison of mutant hESCs vs WT, split by chromatin module (R1-R9).

c) Line plots summarizing the dynamics of accessibility per module and cell line throughout differentiation. Each data point indicates the mean normalised read count.

d-f) Enrichment analysis of overlaps between regions belonging to the nine chromatin modules (from left to right) and annotated reference regions from the Regulatory Index[80] (based on DNaseI-seq; **panel d**), Cis-element Atlas[81] (based on scATAC-seq analysis; **panel e**) and the

Enhancer Atlas[82] (based on a meta-analysis of many different data; **panel f**). The size and
transparency of circles indicate the odds ratio and P-value, respectively (hypergeometric test,
hypeR[138]). Significant results are indicated with filled circles ($P_{adj} \leq 0.05$). The top enrichments
per stage have been selected for visualisation (all results are shown in **panel e**) and all results
are reported in **Supplementary Table 14**.

g) Bubble plots showing the outputs of a fast gene set enrichment analysis (*fgsea*[156]) of open
chromatin in external data compared to our chromatin modules. Each sub-panel indicates data
from a different source (from left to right) including three collections of NB cell lines[75,157,158]
(GSE138293, GSE224241, GSE136279), two adult cancer cell lines [159,160] (GSE202511,
GSE228832), and a meta-analysis of human tissue data[161]
(https://epigenome.wustl.edu/epimap). The epigenetic type[44,45] and *MYCN* amplification status
of each NB cell line are indicated.

Abbreviations: WT, wild-type H7 hESCs; R1-R9, chromatin modules identified in **Fig. 7e**; NES,
normalised enrichment score; n.s., not significant.

2102
2103

Supplementary Figure 13 (related to Fig. 8). Analysis of transcription factor target sets and gene-regulatory networks.

  a) Top 2500 targets of selected TFs as predicted by *GRNboost2* algorithm[83] based on our scRNA-seq data. Putative targets without support in our ATAC-seq data (motif for TF in ≥1 peak near the gene) have been removed. We also calculated the Pearson correlation coefficient (*r*) between each TF and target gene to determine the direction of the putative interaction (*r*>0.1, "activating"; *r*<-0.1, "inhibitory"; others, "marginal"). The top TFs in the target lists have been highlighted. TF target gene sets are reported in **Supplementary Table 15**.

  b) Average expression (Seurat module score) of the target gene sets (matching "activating" targets of the TFs in panel **a** in our integrated scRNA-seq dataset (cp. **Fig. 3d**).

  c) Gene-regulatory networks diagram visualising putative TF to target interactions for the genes in gene sets D9_1 to D9_4 (cp. **Fig. 3e,f**) and enriched TF targets (cp. **Fig. 8d**) In these diagrams, each node represents a TF or target gene, and each edge is a link between a TF and a target. Node size is proportional to the mean target score of the indicated TFs (fixed size for non-TF nodes).

Abbreviations: TF, transcription factor; r, Pearson correlation coefficient.

2120

Supplementary Figure 14

2121
2122
Supplementary Figure 14 (related to Fig. 8). Expression of transcription factors and their target
genes in public RNA-seq datasets.

a) Expression of 24 selected TFs (from **Fig. 8e**) in public bulk RNA-seq data from three NB
tumour compendia (SEQC, TARGET, CCRI). The heatmaps display the normalised transcript
counts per gene and sample, colours have been scaled from 0 to the maximum per source dataset
(dark blue to yellow). The INSS stage, *MYCN* amplification status, and sex of each sample are
indicated by the colour bars on top. Some of the examined TFs are not or very weakly expressed
in the investigated samples from tumours.

b) Heatmaps for the same data and TFs as in panel a, but here heatmap values indicate the TF
target score (Seurat module score), a summary of the expression of putative target genes of each
TF in the respective sample. TFs with high activity in 17q1qMYCN cells (the 6 TFs on top) are
active in *MYCN*-amplified samples.

Abbreviations: INSS, International Neuroblastoma Staging System; TF, transcription factor.

Supplementary tables and files

Supplementary Table 1 (related to Figs. 1, 2, 3, 4, 7, 8). scRNA-seq, WES, and ATAC-seq dataset overview

Supplementary Table 2 (related to Fig. 1). scRNA-seq cluster marker genes (WT)

Supplementary Table 3 (related to Fig. 1). scRNA-seq markers of SCP-SYM-MES transition states

Supplementary Table 4 (related to Fig. 2, 4). Whole-exome sequencing SNVs.

Supplementary Table 5 (related to Fig. 2, 4). Whole-exome sequencing CNAs.

Supplementary Table 6 (related to Fig. 3). scRNA-seq MUT vs. WT differentially expressed genes

Supplementary Table 7 (related to Fig. 3). scRNA-seq MUT vs. WT enrichment results

Supplementary Table 8 (related to Fig. 3). scRNA-seq cluster marker genes (WT+MUT)

Supplementary Table 9 (related to Fig. 3). scRNA-seq genes correlated to mutations

Supplementary Table 10 (related to Fig. 6). Tumour-in vitro markers

Supplementary Table 11 (related to Fig. 6). Survival analyses

Supplementary Table 12 (related to Fig. 7). ATAC-seq regions (peaks) and chromatin modules

Supplementary Table 13 (related to Fig. 7). ATAC-seq differential accessibility analysis

Supplementary Table 14 (related to Fig. 7). ATAC-seq chromatin module enrichment results

Supplementary Table 15 (related to Fig. 8). Transcription factor target genes

Supplementary Table 16 (related to Fig. 8). Transcription factor enrichments