Salmonella-induced SIRT1 and SIRT3 are crucial for maintaining the metabolic switch

in bacteria and host for successful pathogenesis

Dipasree Hajra[1], **Raju S. Rajmani,**[3] **Ayushi Devendrasingh Chaudhary**[4,5], **Shashi Kumar**

Gupta,[4,5] **and Dipshikha Chakravortty**[1,2]

[1]Department of Microbiology & Cell Biology, Indian Institute of Science

[2]Adjunct Faculty, School of Biology, Indian Institute of Science Education and Research,
Thiruvananthapuram

[3]Centre of Infectious Disease Research, Indian Institute of Science, Bangalore, India

[4]Pharmacology Division, CSIR-Central Drug Research Institute, Lucknow, India

[5]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, India

Corresponding author: Prof. Dipshikha Chakravortty

Email: dipa@iisc.ac.in

Tel. No. 0091 80 22932842

ABSTRACT

Sirtuins are the major players in host immuno-metabolic regulation. However, the role of

sirtuins in the modulation of the immune metabolism pertaining to Salmonellosis is largely

unknown. Here, our investigation focussed on the role of two important sirtuins, SIRT1 and

SIRT3, shedding light on their impact on intracellular *Salmonella*'s metabolic switch and

pathogenesis establishment. Our study indicated the ability of the live *Salmonella*

Typhimurium to differentially regulate the levels of SIRT1 and SIRT3 for maintaining the

high glycolytic metabolism and low fatty acid metabolism in *Salmonella*. Perturbing SIRT1

or SIRT3 through knockdown or inhibition, resulted in a remarkable shift in the host

metabolism to low fatty acid oxidation and high glycolysis. This switch led to decreased

proliferation of *Salmonella* in the macrophages. Further, *Salmonella*-induced higher levels of

SIRT1 and SIRT3 led to a skewed polarization state of the macrophages from a pro-

inflammatory M1 state toward an immunosuppressive M2 making it more conducive for the

intracellular life of *Salmonella*. Alongside, governing immunological functions by

modulating p65 NF-κB acetylation, SIRT1, and SIRT3 also skew *Salmonella*-induced host

metabolic switch by regulating the acetylation status of HIF-1α and PDHA1. Interestingly,

though knock-down of SIRT1/3 attenuated *Salmonella* proliferation in macrophages, in *in*

vivo mice-model of infection, inhibition or knockdown of SIRT1/3 led to more dissemination

and higher organ burden which can be attributed to enhanced ROS and IL-6 production. Our

study hence reports for the first time that *Salmonella* modulates SIRT1/3 levels to maintain

its own metabolism for successful pathogenesis.

Keywords: SIRT1 and SIRT3, immune regulation, metabolic shift, macrophage

polarization, *Salmonella* metabolism, bacterial dissemination, deacetylation

INTRODUCTION

Sirtuins are NAD+-dependent deacetylases that are present in all forms of life. Sirtuins

comprise a conserved core catalytic domain that removes acetyl moiety from the lysine

residues of proteins in presence of NAD+ as a cofactor [1] giving rise to 2'O-acetyl-ADP-

ribose and free nicotinamide as products[2][3]. Free nicotinamide acts as a non-competitive

inhibitor of sirtuins[4]. They possess variable N terminal and C terminal domains that confer

different subcellular localization, substrate specificity, and functions[5]. Mammals have

seven sirtuins that are responsible for regulating various biological functions such as cell

survival, apoptosis, oxidative stress, metabolism, and inflammation[6][7]. SIRT1,6 and 7

have nuclear localization, SIRT2 is cytoplasmic and SIRT 3,4 and 5 comprise the

mitochondrial SIRTs. In addition to their deacetylase activity, they possess ADP ribosylation

(SIRT1, SIRT4, and SIRT6), desuccinylation and demalonylation (SIRT5), delipoylation

(SIRT4), and demyristoylation and depalmitoylation (SIRT6) enzymatic activities [8].

Previous studies have shown that SIRT1 gets activated in response to acute immune response

and it deacetylates RelA/p65 component of NFκB thereby mediating its proteasomal

degradation[9]. On the other hand, it activates RelB component of NFκB pathway. RelB

causes heterochromatinization of pro-inflammatory genes like TNF-α and IL-β[10]. SIRT1

activates Peroxisome proliferator-activated receptor γ (PPARγ) coactivator-1α (PGC-1α)

mediating a metabolic switch from glycolysis toward fatty acid oxidation. SIRT1-mediated

RelB activation, in turn, activates SIRT3 causing the promotion of mitochondrial

bioenergetics[11]. PGC-1α, a major player in mitochondrial biogenesis, activates SIRT3 [12]

which in turn causes activation of PGC-1α, thereby fuelling a positive feedback loop. SIRT3

accounts for the major mitochondrial deacetylase orchestrating several metabolic processes

such as fatty acid oxidation, promotion of the TCA cycle, and inhibition of ROS

production[13].

Salmonella enterica serovar Typhimurium is a facultative intracellular Gram-negative enteric

pathogen, causing a wide array of infections ranging from self-limiting gastroenteritis to

diarrhea in humans [14]. *Salmonella enterica* serovar Typhi cause systemic infection in

humans with typhoidal symptoms. Recent reports reported incidences of 21 million [15]

typhoid cases and 93 million of non-typhoidal [16] cases round the year. The virulence of

Salmonella is majorly regulated by two pathogenicity islands, namely, SPI-1 and SPI-2. It

uses SPI-1 encoded T3SS and the effector proteins to invade host cells [17]. Inside the

macrophages, they harbour within the *Salmonella* containing vacuoles (SCV) by virtue of its

SPI-2 effectors [18]. Macrophages, dendritic cells and neutrophils are responsible for

successful dissemination throughout the body through the reticulo-endothelial system (RES)

[17].

Macrophages, serving as an intracellular niche for *Salmonella,* exhibit several continua of

polarization states. At the two extreme ends of the spectrum lie the classically polarized M1

macrophages and alternatively activated M2 macrophages. M1 macrophages comprise of the

proinflammatory antimicrobial state producing IL-1β, IL-6, TNF-α,IL-12, IFN-γ cytokines

and exhibit enhanced expression of CD80, CD86 surface markers. The anti-inflammatory M2

macrophages promote bacterial persistence by producing anti-inflammatory cytokines like

IL-10,TGF-β and show increased expression of Arg-1, CD206 surface markers[19][20]. To

sustain the continuous production of proinflammatory cytokines M1 macrophages rely on

glycolysis for their energy requirements. On the other hand, M2 macrophages are fuelled by

enhanced oxidative phosphorylation (OXPHOS) and fatty acid oxidation (FAO)[21].It has

been previously reported that sirtuins mediated attuning of metabolism impact polarization of

macrophages *in vivo*. SIRT1 has the ability to promote the polarization of M2 macrophages

and inhibit inflammation in macrophages of adipose tissue[22][23][24]. SIRT3 suppresses

ROS by deacetylating and activating MnSOD[25].

Several bacteria are known to subvert the host immune system toward an immunosuppressive

state. *Salmonella* or *Mycobacterium* have evolved mechanisms to counteract the M1 state of

the host macrophage. *Salmonella* Typhimurium uses its SPI-2 effectors to inhibit the

recruitment of NADPH oxidase to the SCV, thereby preventing oxidative burst mediated

microbicidal activity[26]. Similarly, *Mycobacterium bovis* bacillus Calmette-Gue´rin

prevents NOS2 recruitment to phagosomes[27]. *Salmonella* Dublin causes inhibition of the

production of pro-inflammatory cytokine like IL-18 and IL-12p70[28].   Moreover,

Mycobacteria inhibits NFκB signalling and IFN-γ mediated downstream pathways[29].

Furthermore, *Yersinia enterocolitica* elicits an M2 response by inducing Arginase-1

expression and TGFβ1 and IL-4production[30]. *Yersinia* TTSS effector LcrV induces an M2

phenotype supposedly by IL-10 production[31].

Since SIRT1 and SIRT3 are the major modulators of the immuno-metabolic paradigm, we

intend to decipher the role of SIRT1 and SIRT3 in influencing host and *Salmonella*

metabolism. This study highlights the role of SIRT1 and SIRT3 in intracellular pathogen

survival by promoting *Salmonella* glycolysis and concomitantly driving host metabolism

towards fatty acid oxidation. Additionally, *Salmonella* trigger an immunosuppressive M2

environment conducive to its intravacuolar proliferation by modulating SIRT1 and SIRT3

levels. Here, we have shown that SIRT1 and SIRT3 knockdown cause decreased M2 surface

marker expression such as CD206 along with increased production of pro-inflammatory

cytokines and ROS generation, together amounting to attenuated bacterial intracellular

proliferation within the infected macrophages. Moreover, SIRT1 mediated p65 NF-κB

deacetylation played a vital role in immune function regulation within the *Salmonella*

infected macrophages with increased interaction of SIRT1 with p65 NF-κB. SIRT1

knockdown or inhibition resulted in hyperacetylation of p65 NF-κB thereby leading to

enhanced pro-inflammatory response in *S*. Typhimurium infected macrophages. Further,

SIRT1 and SIRT3 knockdown or inhibition skewed the *Salmonella*-induced host metabolic

shift by regulating acetylation status of HIF-1α and PDHA. This caused increased host

glycolysis and reduced fatty acid oxidation. However, the *Salmonella* shows the opposing

metabolic profile with increased fatty acid oxidation and reduced glycolysis upon SIRT1 or

SIRT3 inhibition. In the contrary to the macrophages, in *in vivo* mice-model of infection,

SIRT1 and SIRT3 inhibition resulted in increased pathogen loads in organs and triggered

enhanced bacterial dissemination, together leading to increased susceptibility of the mice to

S. Typhimurium infection owing to increased ROS and IL-6 production. To the best of our

knowledge, this is the first report implying the ability of host Sirtuins in impacting

intracellular bacterial metabolism crucial for successful pathogenesis.

RESULTS

Salmonella **modulates SIRT1 and SIRT3 expression along its course of infection in**

macrophages

Upon infection of RAW 264.7 murine macrophages with wildtype *Salmonella* Typhimurium

strain 14028S, we observed an increased expression level of *SIRT 1* and *SIRT3* at initial and

middle phases of infection, precisely at 2hr and 6hr post infection through qPCR (Fig.1, A

and B). The *SIRT1* expression level declined at later phases of infection. On the other hand,

the *SIRT3* transcript levels remained elevated at all time points with respect to uninfected

control with a marked increment at 16hr time point post-infection which subsided at 20hr

post-infection. We even monitored the expression profile of *SIRT1* and *SIRT3* in primary

macrophages like peritoneal macrophages of C57BL/6 mice and observed a similar trend of

elevated expression at initial (2hr), middle (6hr), and late (16hr) time points post-infection

(Fig.1, C and D). In Confocal Laser Scanning Microscopy (CSLM) studies, we observed a

similar increase in SIRT1 and SIRT3 expression at 6hr post infection within the infected

macrophages RAW264.7 macrophages. (Fig.1, E-H). Immunoblotting revealed increased

protein expression of both SIRT1 and SIRT3 at 2hr post-infection in comparison to the

uninfected control (Fig. S1). However, SIRT1 expression exhibits a gradual decline at the late

phase of infection (Fig. S1A, C). In line with the confocal microscopy data, SIRT3

immunoblotting data shows an increased protein expression profile at 6hr and 16hr post-

infection (Fig. S1, B-D). Subsequently, to ascertain whether indeed *Salmonella* could

modulate SIRT1 or SIRT3 expression levels, we evaluated the *SIRT1* and *SIRT3* mRNA and

SIRT1 and SIRT3 protein expression profile within RAW264.7 macrophages upon infection

with wildtype *S*. Typhimurium and SPI-1 (Δ*invC*) (InvC- protein export apparatus) or SPI-2

(Δ*ssaV* and Δ*steE*) (SsaV- Structural component of SPI-2 needle apparatus, SteE- SPI-2

effector protein involved in driving M2 polarization) mutants of *S*. Typhimurium. Our results

depict the ability of wildtype *S*. Typhimurium to induce the expression of both SIRT1 and

SIRT3 within the infected RAW264.7 macrophages. However, infection with either SPI-1 or

SPI-2 mutant abrogates the induction of *SIRT1* whereas only SPI-2 mutants (Δ*ssaV* and Δ*steE*

) and not SPI-1 mutant infection caused reduction in *SIRT3* transcript level expression in the

infected macrophages, implicating the role of SPI-1 and SPI-2 genes in triggering SIRT1 and

SIRT3 in the infected macrophages (Fig. 1, I-J). However, at the protein level, only SPI-2

mutant infection resulted in a predominant decline in SIRT3 expression and a mild reduction

in SIRT1 expression (Fig. S1B). Further, we examined the transcript level profile of *SIRT1*

and *SIRT3* in M1 or M2 polarized RAW264.7 macrophages at 16hr post-infection and we

observed 20-fold and 5-fold increase in *SIRT1* and *SIRT3* expression in M2 polarized infected

macrophages as opposed to 0.5-fold and 0.4 fold downregulation in M1 polarized infected

macrophages (Fig. S1C-D). Thus, an increase in expression profile both at transcript and

protein levels indicates their role in *Salmonella* pathogenesis.

SIRT1 and SIRT3 play crucial role in intracellular bacterial proliferation in infected

murine macrophages

As our previous *SIRT1* and *SIRT3* expression data in the polarized macrophages, hinted at the

role of SIRT1 and SIRT3 in driving polarization of macrophages in the infected

macrophages. We validated the intracellular replication of *S*. Typhimurium within the

infected polarized macrophages. *S*. Typhimurium exhibited increased intracellular fold

proliferation within anti-inflammatory M2 polarized macrophages in comparison to the pro-

inflammatory M1 polarized RAW264.7 macrophages (Fig. S1 E-F). To evaluate the role of

SIRT 1 and SIRT3 in the intracellular proliferation of the bacteria within murine

macrophages, we have undertaken knockdown of SIRT1 and SIRT3 in RAW 264.7 (Fig. S2)

macrophages through PEI-mediated transfection of shRNA plasmids directed against SIRT1

and SIRT3. Post 48hr of transfection, the transfected cells were infected with MOI=10 of

wildtype *S*. Typhimurium and a gentamicin protection assay was performed. Intracellular

proliferation of the bacteria was quantified by plating the cell lysate at 2hr and 16hr post-

infection. *Salmonella* exhibits compromised intracellular survival in SIRT1 and SIRT3

knock-down RAW264.7 macrophages in comparison to the un-transfected and scrambled

controls (Fig.2, A). Further, we have assessed the intracellular proliferation in peritoneal

macrophages isolated from thioglycolate-treated adult C57BL/6 mice post SIRT1 (EX-527)

and SIRT3 (3TYP) inhibitor treatment. SIRT1 or SIRT3 inhibitor-treated macrophages

exhibited attenuated intracellular replication in comparison to the untreated peritoneal

macrophages (Fig,2, B). Together, our results depict the role of SIRT1 and SIRT3 in

controlling the intracellular proliferation of *S*. Typhimurium.

SIRT1 and SIRT3 inhibition contribute to skewed inflammatory host responses upon

Salmonella **infection**

Several reports indicate the role of SIRT1 and SIRT3 in the modulation of host immune

responses pertaining to infection scenarios [35][36][37][38]. Therefore, we intend to check

whether SIRT1 or SIRT3 regulates immune functions in *Salmonella*-infected macrophages.

To delineate the role of SIRT1 and SIRT3 in the modulation of immune responses, we

wished to investigate the production of pro-inflammatory and anti-inflammatory cytokines in

knockdown (KD) RAW 264.7 macrophages upon *S*. Typhimurium infection. Post 48hr

transfection, cells were subjected to wildtype *S*. Typhimurium infection at an MOI of 10. At

the indicated time points, cell-free supernatant was harvested and evaluated for pro-

inflammatory and anti-inflammatory cytokine production by ELISA. Inhibition of both

SIRT1 and SIRT3 increased production of pro-inflammatory cytokine IL-6 significantly at

2hr and 20hr post-infection (Fig.S3, A). Moreover, there was only a significant reduction in

anti-inflammatory IL-10 production upon SIRT1 and SIRT3 KD at 2hr and 20hr post-

infection (Fig.S3, B). Further, we estimated the production of another pro-inflammatory

cytokine, IL-1β at 20hr post-infection under the knockdown condition of SIRT1 and SIRT3

(Fig.S3, C) and we observed heightened IL-1β production under knockdown of SIRT1 and

SIRT3 in comparison to the scrambled infected control. In peritoneal macrophages upon

SIRT1(EX-527-1μM) or SIRT3(3-TYP-1μM) chemical inhibitor treatment, an increase in IL-

6 and IL-1β cytokine levels was observed at 6hr post-infection (Fig.S3, D-E). This indicates

the possible role of SIRT1 and SIRT3 in the regulation of cytokine production upon

Salmonella infection.

Immune functions are an important determinant of macrophage polarization. Since SIRT1

and SIRT3 played an immune-modulatory role in *Salmonella* infection, we investigated

whether *Salmonella* infection is associated with a shift in macrophage polarization status. To

assess the ability of the pathogen to alter the polarization state of the macrophage, we have

undertaken gene expression profiling of various M1 and M2 markers using nanoString

nCounter technology along the course of *Salmonella* Typhimurium infection at the indicated

time points in RAW 264.7 macrophages. A gradual shift from pro-inflammatory M1 toward

the anti-inflammatory M2 state was observed with the progression of *Salmonella* infection.

Along the course of infection, there was a reduction in the expression of M1 markers like

NOS2, *CD40*,*CD86*, *TNF-α*, *Nfkb2*,*IL-6* and a corresponding increase in the expression of the

M2 markers such as *Arg-1*, *CCL-17*, *CD 206*, *IL4ra* with an exception of *Tgf-β* (Fig.3, A).In

order to validate the polarization potency of the pathogen, FACS was performed using a pro-

inflammatory M1 surface marker, CD86 tagged with PE. The data suggests a distinct

decrease in CD86 positive population in the infected sets in comparison to the uninfected and

the fixed dead bacteria control along the course of *S*. Typhimurium infection (Fig.3, B, Fig.

S4, A). Thus, the live pathogen has a propensity to skew the polarization state of the

macrophage toward an anti-inflammatory M2 state to subvert the initial acute inflammatory

response mounted by the host immune system.

To assess the role of SIRT1 and SIRT3 in macrophage polarization, we determined anti-

inflammatory CD206 surface marker profiling of the infected macrophages through flow

cytometry. Knockdown of SIRT1 or SIRT3 in infected RAW 264.7 macrophages resulted in

a reduction in anti-inflammatory CD206 surface marker expression at 16hr post-infection

(Fig. 3, C, Fig. S4, B). Moreover, SIRT1 or SIRT3 knockdown led to enhanced intracellular

ROS generation within the infected macrophages in comparison to the scrambled or the

untransfected control (Fig.S5). Further, the Haematoxylin and Eosin staining of the *S*.

Typhimurium infected mice liver tissue sections depicted exacerbated signs of inflammation

in the SIRT1 (EX-527) or SIRT3 (3TYP) inhibitor-treated cohorts in comparison to the

untreated controls with multiple necrotic foci and increased influx of inflammatory cell

inflates. Moreover, these acute inflammatory signs of the liver sections get ameliorated in the

SIRT1 (SRT1720) activator-treated infected cohort (Fig.8, T-U) (Data described later).

Together, these data suggest the role of SIRT1 and SIRT3 in the modulation of host

inflammatory response. Previous literature reports have shown that SIRT1 physically

interacts with p65 subunit of NF-κB and inhibits transcription by deacetylating p65 at lysine

310 [39]. Moreover, SIRT1-mediated deacetylation of the p65 subunit of the master regulator

of the inflammatory response, NF-κB, results in the reduction of the inflammatory responses

mediated by this transcription factor [40]. To evaluate the immune regulatory mechanism of

SIRT1 in the *S*. Typhimurium (STM) infection scenario, we undertook SIRT1

immunoprecipitation in the infected RAW264.7 macrophages at 16h post-infection alongside

the uninfected macrophages and probed for NF-κB p65 interaction. We observed an

increased interaction of SIRT1 with NF-κB p65 in the infected macrophages in comparison to

the uninfected control (Fig.4, A). Further, the knockdown of SIRT1 resulted in increased

acetylation status of the NF-κB p65 upon infection in comparison to the scrambled, infected

control (Fig.4, B, C). To understand whether the enzymatic domain of SIRT1 possess any

role in mediating this interaction, we carried out NF-κB p65 immunoprecipitation in infected

RAW264.7 macrophages in presence or absence of SIRT1 catalytic chemical inhibitor, EX-

527(1μM) treatment at 16h post-infection. We observed an increased interaction of NF-κB

p65 with SIRT1 in the infected untreated macrophages when compared to the untreated

uninfected control (Fig.4, D). However, the interaction of NF-κB p65 with SIRT1 gets

abrogated under the EX-527 inhibitor treatment in the infected macrophages thereby

implying the function of the catalytic domain in mediating the interaction (Fig.4, D).

Moreover, an increased acetylation status of NF-κB p65 was observed in the EX-527 treated

infected macrophages in comparison to the untreated infected macrophages (Fig. 4 E).

SIRT1 and SIRT3 relieve oxidative stress in infected macrophages and alleviation of the

intracellular ROS generation restores intracellular survival of *S.* Typhimurium within

the SIRT1 or SIRT3 knockdown macrophages

Previous reports have suggested the role of SIRT1 and SIRT3 in oxidative stress conditions.

They are known to act in concert as anti-oxidants by reducing ROS production[41][42][43].

Moreover, enhanced ROS production is also a prototype of the classically activated

macrophages[44][45]. Here, we examined the effect of SIRT1 and SIRT3 knockdown in

intracellular ROS generation in *S.* Typhimurium infected RAW264.7 macrophages through

DCFDA staining in FACS. Results depicted significant enhancement in the production of

ROS at 16hr post infection upon knockdown of SIRT1 or SIRT3 in comparison to

untransfected or scrambled control (Fig.S5, A,B). Upon detection of extracellular ROS

generation through Phenol Red Hydrogen Peroxidase assay, we detected higher ROS

generation upon SIRT3 KD at 6hr post infection and greater ROS production at 16hr and

20hr time points in SIRT1 knockdown macrophages (Fig.S5,C). Therefore, SIRT1 and

SIRT3 play an important role in mitigating the oxidative burst in *Salmonella* infected

macrophages. Our previous findings depicted decreased intracellular burden of *S.*

Typhimurium within the SIRT1 or SIRT3 knockdown macrophages along with increase in

intracellular ROS generation. Therefore, we speculated that the decreased intracellular

proliferation within the SIRT1 or SIRT3 knockdown macrophages might be on account of

increased intracellular ROS production which might lead to increased killing of the

intracellular bacteria. This hypothesis led us to evaluate the intracellular bacterial burden

within the infected SIRT1 or SIRT3 knockdown RAW264.7 macrophages or SIRT1 or

SIRT3 inhibitor treated peritoneal macrophages (isolated from C57BL/6 adult mice post 5[th]

day of thioglycolate injection) in presence of a ROS inhibitor named N-Acetyl Cysteine

(NAC). NAC acts as a scavenger of ROS by antagonizing the activity of proteasome

inhibitors [46]. The attenuated intracellular proliferation of *Salmonella* Typhimurium within

the knockdown or the chemical inhibitor treated macrophages got restored upon 1mM

treatment of ROS scavenger, NAC (Fig.S6). Therefore, intracellular ROS production within

the knockdown murine macrophages is one of the reasons for the attenuated survival of the

bacteria.

SIRT1 and SIRT3 play crucial role in mediating metabolic switch in infected

macrophages

Macrophage polarization is not only governed by immunological changes but also

contributed by metabolic reprogramming[47][24][48]**.** Since previous data suggested

progression of *Salmonella* infection with the shift in polarization state of the macrophage, we

decided to investigate alteration of the metabolic state of the macrophages as macrophage

polarization is governed by immune-metabolic shift. In pursuit of fulfilling such requirement,

we performed gene expression studies of various metabolic genes through nanoString nCounter technology in *S*. Typhimurium infected RAW 264.7 macrophages. Analysis of the gene profile revealed upregulation of genes involved in fatty acid oxidization and tricarboxylic acid cycle and corresponding downregulation of genes involved in glycolysis (Fig.5, A). To validate the findings, we carried out qRTPCR to quantitatively measure the expression of a fatty acid oxidation gene, *PPARδ* in infected RAW 264.7 macrophages. We found that mRNA level was elevated to 2-fold at 2hr and 6hr post-infection. At the late phase of infection, 16hr post infection around 6-fold upregulation in mRNA transcript was noted (Fig.5, B). Lactate estimation assay of *S*. Typhimurium infected RAW264.7 macrophages at the initial time point of 2hr and at the late time point of 16hr post infection revealed decline in lactate (glycolysis end product) production at 16hr in comparison to 2hr post-infection timepoint (Fig. 5C). Together, these results suggest the capability of the pathogen to drive a shift in the metabolic status of the infected macrophages toward fatty acid oxidation. Next, we evaluated the function of SIRT1 and SIRT3 in influencing the metabolic switch in the infected macrophages through qRT PCR with several host fatty acid oxidizing genes (*Acox*, *Hadha, Pdha*, *and AcadL)* and glycolytic gene (*PfkL*) in SIRT1 and SIRT3 knockdown macrophages (Fig. S7, A) and via lactate production assay (Fig. 5, D-E). Lactate estimation assay in SIRT1 and SIRT3 knockdown condition revealed enhanced lactate production at 16hr post infection in comparison to the scrambled control which further authenticates the increased host glycolysis upon SIRT1 and SIRT3 knockdown scenario (Fig. 5, D-E). SIRT1 and SIRT3 knockdown RAW 264.7 infected macrophages showed decreased expression of fatty acid oxidizing genes and increased expression of glycolytic *PfkL* gene in comparison to the scrambled infected control (Fig. S7, A). Similar results were obtained in the infected peritoneal macrophages under the SIRT1 or SIRT3 catalytic inhibitor treatment (Fig. S7, B-D). Moreover, qRT PCR-mediated metabolic gene profiling of liver and spleen isolated from

wildtype *S*. Typhimurium infected macrophages revealed decreased transcript level

expression of murine fatty acid oxidation genes upon treatment with SIRT1 (EX-527) or

SIRT3 (3TYP) catalytic inhibitors which got reversed upon SIRT1 activator (SRT1720)

treatment (Fig. S7, E-F). Moreover, SIRT1 and SIRT3 knockdown or catalytic inhibition in

peritoneal macrophages resulted in increased protein expression of host glycolytic genes such

as Phosphoglycerate kinase (Pgk), Phosphofructokinase (Pfk) with concomitant reduction in

protein expression of TCA cycle gene like Pyruvate dehydrogenase (Pdha1) and fatty acid

oxidation genes such as Hadha and Acox1 (Fig.5, F, Fig. S7D). Fatty acid oxidation assay in

the RAW264.7 macrophages under SIRT1 or SIRT3 knockdown condition or inhibition

treatment revealed significant decrease in fatty acid β oxidation activity of the infected

macrophages in comparison to the scrambled or the untreated control (Fig.5, G). To

investigate whether the host metabolic switch toward increased fatty acid oxidation is being

driven by the pathogen, we performed fatty acid β oxidation activity assay under wildtype *S*.

Typhimurium (STM WT), SPI-1 (Δ*invC*) or SPI-2 (Δ*ssaV* and Δ*steE*) mutants of *S*.

Typhimurium infection condition. We found that the wildtype bacteria with its intact SPI-2

effector secretion apparatus could promote increased host fatty acid β oxidation. In the

contrary, the SPI-2 (Δ*ssaV* and Δ*steE*) mutants of *S*. Typhimurium failed to drive host

metabolic shift towards increased fatty acid oxidation (Fig. 5, H).

Collectively, these data suggest the role of SIRT1 and SIRT3 in mediating the *Salmonella*

induced host metabolic shift in the infected macrophages.

SIRT1 and SIRT3 concomitantly influences *Salmonella* metabolism

Our previous data indicated a shift in host metabolism toward increased fatty acid oxidation

along the course of *Salmonella* Typhimurium infection in murine RAW 264.7 macrophages.

Salmonella Typhimurium drives the metabolism of the infected macrophage toward fatty acid

oxidation. This observation led us to investigate the influence of host metabolic shift on the

metabolic status of the pathogen harboring inside the infected macrophages[49]. We were

intrigued whether increased glucose availability within the fatty-acid oxidizing macrophages

is utilized by the bacteria. Thus, we undertook simultaneous gene expression studies of

various *Salmonella* genes involved in their pathogenesis and metabolism through nanoString

nCounter technology in *S*. Typhimurium infected RAW 264.7 macrophages. The nanoString

gene profile revealed enhanced expression of genes involved in *Salmonella* glycolysis and

glucose uptake such as *pfkA* and *ptsG*, respectively (Fig.6, A). This finding indicates the

ability of the pathogen to drive the metabolic state of the host toward fatty acid oxidation

with corresponding increased glucose utilization by the bacteria favoring their survival inside

the host. qRT PCR results with several bacterial fatty acid oxidizing genes (*fadA,*

fadB,fadL,aceA, aceB) and glycolytic genes (*ptsG*) in knockdown condition of SIRT1 and

SIRT3 in RAW 264.7 macrophages further validated the nanoString gene expression profiles

(Fig.6, B). In scrambled control, *Salmonella* infection progresses with increased *Salmonella*

glycolysis and reduced bacterial fatty acid oxidation. However, knockdown of SIRT1 and

SIRT3 abrogate this bacterial metabolic shift by reducing its glycolysis and by exhibiting

enhanced fatty acid oxidation thereby attenuating pathogen intracellular survival. Similar

observations were obtained from the qPCR data in the infected mice liver and spleen samples

with increased transcript level expression of bacterial fatty acid oxidation genes and

decreased expression of bacterial glycolytic genes upon SIRT1 or SIRT3 inhibitor treatment

(Fig. 6 C, D). Therefore, SIRT1 and SIRT3 driven host metabolic switch potentially

influence the metabolic profile of the intracellular pathogen.

Mechanism behind SIRT1 or SIRT3 mediated metabolic switch

As per our previous findings, SIRT1 or SIRT3 inhibition led to increased host glycolysis and

decline in fatty oxidation in the infected macrophages. HIF1α is a master regulator of

glycolysis in host during stress conditions [50]. Previous reports have suggested HIF1α to be

a target of deacetylation by SIRT1 at Lys 674 which contribute to metabolic reprogramming

in cancer cells. During hypoxia, downregulation of SIRT1 leads to increased acetylation and

activation of HIF1α [51]. Additionally, in CD4+ T cells, ectopic expression of SIRT1

inhibited IL-9 production and glycolysis by negatively regulating HIF1α [52]. To delve into

the mechanism behind SIRT1 mediated modulation of metabolic responses, we assessed the

interaction of SIRT1 with HIF-1α in infected RAW264.7 macrophages. The

immunoprecipitation studies of SIRT1 showed increased interaction of the SIRT1 with

HIF1α in the *S.* Typhimurium infection scenario with respect to the uninfected control (Fig.7,

A). Further, we evaluated the acetylation status of HIF1α in the SIRT1 knockdown status of

the infected macrophages. We found that SIRT1 knockdown showed increased acetylation of

HIF1α in the infected macrophages in comparison to the scrambled infected control at 16hr

post-infection (Fig. 7, B,C). Immunoprecipitation studies under SIRT1 (EX-527) inhibitor

treatment in RAW264.7 macrophages revealed increased acetylation of HIF1-α along with

reduced interaction of HIF-1α with SIRT1, thereby indicating the probable role of the

catalytic domain in influencing the interaction (Fig. 7 D-F). Further, to ascertain the role of

HIF-1α in mediating the glycolytic switch in the infected macrophages, we estimated the

lactate production under SIRT1 and SIRT3 knockdown conditions in the presence or absence

of HIF-1α inhibitor (chetomin)[53]. Our results depicted a decline in lactate production upon

chetomine treatment including under SIRT1 and SIRT3 knockdown conditions (Fig.7G).

Together, our results implicate the role of SIRT1 in governing glycolytic shift in the infected

macrophages by deacetylating HIF1α. Upon SIRT1 knockdown or inhibition, HIF1α gets

hyperacetylated which causes activation of the downstream glycolytic genes. Alternatively,

several key literatures suggest the role of SIRT3 in modulating metabolic programming by

deacetylating several proteins involved in fatty acid oxidation, the tricarboxylic acid cycle

and oxidative phosphorylation [54][55]. PDHA1 (Pyruvate Dehydrogenase E1 subunit alpha)

is a key enzyme linking glycolysis to TCA cycle and oxidative phosphorylation. SIRT3

regulates PDHA1 acetylation by deacetylating PDHA1 at lysine 385 residue, thereby playing

a key role in metabolic reprogramming [55]. PDHA1 acetylation coincides with PDH activity

and increased PDHA1 phosphorylation [55]. Therefore, we investigated the role of SIRT3 in

the modulation of host fatty acid oxidation upon S. Typhimurium infection in RAW264.7

macrophages. To do so, we immunoprecipitated PDHA1 and checked for its interaction with

SIRT3 or SIRT1 under the knockdown condition of SIRT3 or upon SIRT3 inhibitor treatment

(Fig.7 I-J). We observed an increased interaction of PDHA1 with SIRT3 in the infection

scenario in comparison to the uninfected control which gets eventually abolished under the

knockdown condition (Fig.7 I-J) and under the chemical inhibitor treatment of SIRT3 (Fig.7,

K-L) suggesting the role of the SIRT3 in mediating the interaction with PDHA1. Alongside

the decreased interaction of PDHA1 with SIRT3, increased acetylation of PDHA1 was

detected upon SIRT3 inhibitor treatment in infected macrophages (Fig.7, K-L).

SIRT1 or SIRT3 inhibition enhances bacterial burden in mice *in vivo*

6-8 weeks old adult male C57BL/6 mice were treated with SIRT1 inhibitor EX-527, SIRT3

inhibitor 3TYP and SIRT1 activator SRT1720 at a dose of 1mg/kg each via intraperitoneal

injection (every alternate Day) (Fig. 8A). Following the inhibitor treatment, the mice were

orally gavaged with 10^7 CFU of S. Typhimurium 14028S for organ burden evaluation or with

10^8 CFU of wildtype S. Typhimurium for survival studies. On day 5th post-infection, mice

were sacrificed, and the liver, spleen and Mesenteric Lymph Node (MLN) were harvested for

enumeration of the organ burden. The SIRT1 inhibitor, EX-527 and SIRT3 inhibitor, 3TYP-

treated mice cohorts exhibited increased organ loads in liver, spleen and MLN in comparison

to the vehicle control. On the contrary, the SRT1720 treated mice group showed organ

burden comparable to that of the vehicle control (Fig. 8, B). Moreover, the SIRT1 and the

SIRT3 inhibitor-treated mice cohorts died earlier than the vehicle-treated control mice group

or the SIRT1 activator-treated group (Fig. 8, C). Further, the SIRT1 and the SIRT3 inhibitor-

treated mice cohorts showed increased splenic length in comparison to the vehicle-treated

mice group and the SIRT1 activator-treated mice cohort (Fig. 8, D). The increased organ

burden in the EX-527 or 3TYP treated group might be due to increased bacterial

dissemination in blood. To assess bacterial dissemination, blood was collected from infected

mice post-inhibitor treatment at specific days post-infection retro-orbitally and plated onto

Salmonella Shigella (SS) agar plates for bacterial enumeration. Indeed, increased bacterial

dissemination was observed in the blood of mice treated with SIRT1 inhibitor, EX-527 or

SIRT3 inhibitor, 3TYP at day 1-, 2-, 3- ,4- post-infection in comparison to the vehicle-treated

mice (Fig. 8, E). Further, we wanted to examine whether the increased bacterial

dissemination was due to increased ROS production or due to the presence of elevated

inflammatory cytokine levels like IL-6 and IL-1β. In the wildtype C57BL/6 mice treated with

SIRT3 inhibitor 3TYP showed heightened bacterial burden in blood at 5^{th} day post-infection

in comparison to the vehicle control. Nevertheless, the *gp91phox-/-* mice group lacking the

catalytic subunit of NADPH oxidase did not depict significant variation in the bacterial load

among the different mice treatment cohorts (Fig. 8, F). Further, the EX-527 (SIRT1 inhibitor)

and the 3TYP (SIRT3 inhibitor) treated mice possessed elevated levels of serum IL-6

(Fig.8G), IL-1β (Fig. S3F) and showed increased intracellular ROS burden in infected liver

tissues in comparison to the vehicle-treated control and the SRT1720 (SIRT1 activator)

treated mice group (Fig. 8H). The increased mouse serum IL-6 and IL-1β production was in a

similar line with the increased IL-6 or IL-1β cytokine generation in EX-527 or 3TYP treated

peritoneal macrophages under the infection scenario (Fig. S3, D-E). Moreover, estimation of

IL-1β within the infected intestinal ileal sections of the mice revealed increased pro-

inflammatory IL-1β generation in the SIRT1 and SIRT3 inhibitor-treated mice groups in

comparison to the untreated or the SIRT1-activator treated mice cohorts (Fig. S3G).

However, contrary to the *in vitro* studies wherein SIRT1 or SIRT3 knockdown or inhibition

resulted in attenuated intracellular proliferation, here in *in vivo* mouse model of infection, we

observed increased bacterial organ loads owing to increased bacterial dissemination. To

delineate this observation further, we evaluated the bacterial load within splenocytes isolated

from control or inhibitor-treated C57BL/6 mice infected with GFP expressing *S.*

Typhimurium at 5^{th} day post-infection via flow cytometry. We observed heightened bacterial

load in the EX-527 or the 3TYP treated mice cohorts (Fig. 8, I-J). However, when we

enumerated the bacterial count within the F4/80+ macrophage population of the infected

splenocytes, we noticed decreased bacterial loads in the EX-527 or 3TYP -treated mice group

in comparison to the vehicle-treated control group or the SRT-1720 activator-treated group

(Fig. 8, K-L). Further, we evaluated additional splenic populations including CD45+, Ly6C+,

and CD11c+ populations. Our results show that the CD45+ splenic population depicts

increased bacterial loads like that of the total splenic population within the SIRT1 or SIRT3

inhibitor-treated cohorts. However, CD45+ monocytes and Ly6C positive splenic population

exhibit compromised burden within the SIRT1 and SIRT3 inhibitor-treated cohorts.

Moreover, CD11c+ population, CD45+ granulocytes, or lymphocytes show comparable

organ loads to that of the vehicle control or SIRT1 activator-treated mice group (Fig. 8M-S,

Fig. S8). Overall, our data suggest heterogeneous bacterial burden in diverse splenic

populations. This opposing phenotype could be attributed to the increased IL-6 and IL-1β

cytokine storm and elevated ROS production upon the SIRT1 or SIRT3 inhibitor treatment

which in turn resulted in bacterial dissemination *in vivo* and concomitantly restricted the *in*

vitro intracellular proliferation within macrophages. To validate this observation, we

estimated the ROS levels within the liver and spleen tissues harvested from *S.* Typhimurium

infected C57BL/6 mice, treated with specific catalytic inhibitor, activator or vehicle via

DCFDA staining using flow cytometry at 5^{th} day post-infection. We detected escalated levels

of ROS within both the infected liver and spleen tissues of the EX-527 or 3TYP-treated mice

groups in comparison to the vehicle-treated or the SRT1720 treated mice cohorts (Fig. 8, H,

S9). Haematoxylin and eosin staining of the liver sections (harvested at 5^{th} day post-

infection) revealed increased inflammation with multiple areas of severe acute hepatic

necrosis with complete loss of hepatic architecture in the EX-527 and 3-TYP treated liver

samples in comparison to the vehicle-treated control and SRT-1720 treated liver samples

(Fig. 8, T-U). In line with the inhibitor-treated studies, the increased organ loads, and

systemic dissemination driven heightened susceptibility of mice toward *S*. Typhimurium

infection were replicated in *in vivo* SIRT1 and SIRT3 adeno-associated virus serotype 6

(AAV6) mediated knockdown mice model which showed elevated IL-6 production in

comparison to the scrambled control treated mice cohort (Fig. 8, V-X, Fig. S10).

Simultaneously, the haematoxylin-eosin-stained sections of the liver tissues harvested from

the shSIRT1 or shSIRT3 mice cohorts depicted increased pathological scoring with multiple

necrotic areas and severely damaged liver tissue architecture in comparison to the scrambled

mice control (Fig. 8, Y). Altogether, our results implicate the role of SIRT1 and SIRT3 in

controlling *S*. Typhimurium infection *in vivo*.

DISCUSSION

Several studies have confirmed the propensity of *Salmonella* to skew the polarization state of

the infected macrophages toward an immunosuppressive anti-inflammatory state[56][57][58].

We have validated such findings and further elaborated it by depicting the role of SIRT1 and

SIRT3 in the modulation of host immune responses as well as host-bacterial metabolism.

Salmonella Typhimurium infection modulates the expression profile of both SIRT1 and

SIRT3 in the infected macrophages at both mRNA and protein level via its SPI-2 effector.

Downregulation of SIRT1 and SIRT3 through shRNA mediated knockdown resulted in

heightened pro-inflammatory immune responses with increased production of IL-6 cytokine

and decreased surface expression of anti-inflammatory CD206. SIRT1 and SIRT3

downregulation also resulted in increased intracellular ROS production in the infected

macrophages. SIRT1 and SIRT3 knockdown macrophages not only show altered host

immune status but also depicted shift in the metabolic state with increased glycolytic shift.

This altered host metabolism upon SIRT1 and SIRT3 knockdown condition influences the

outcome of infection by regulating the intracellular bacterial metabolism which shows

reduced bacterial glycolysis and increased fatty acid oxidation. All these outcomes account

for attenuated intracellular bacterial proliferation in the SIRT1 and SIRT3 knockdown

macrophages. However, in murine model of infection, SIRT1 or SIRT3 inhibitor treatment

led to increased organ burden and triggered bacterial dissemination (Fig. S11). Overall, our

study highlights the crucial role of SIRT1 and SIRT3 in governing the host immune-

metabolic shift during *Salmonella* infection which in turn is vital for maintaining *Salmonella*

metabolism.

Previous reports have elucidated the role of SIRT1 and SIRT2 pertaining to *Salmonella*

infection. Ganesan et al., have depicted the role of SIRT1 in autophagy in *Salmonella*

infection scenario[59]. Gogoi et al., have demonstrated SIRT2 mediated modulation of

immune responses in dendritic cells[60]. Till date, there is no report highlighting the role of

SIRT3 governing the *Salmonella* pathogenesis. The function of SIRT3 in infection scenario

has been explored quite recently. In *Mycobacterium tuberculosis* infection condition, SIRT3

control mitochondrial function and autophagy[38]. SIRT3 downregulation in *Mycobacterium*

tuberculosis infected macrophages is associated with dysregulated mitochondrial metabolism

and increased cell death[61]. In this study we have explored the role of SIRT1 and SIRT3 in

mediating host immune-metabolic switch in *Salmonella* Typhimurium infected macrophages which further govern intracellular bacterial metabolism and pathogenesis.

Our findings suggest the role of SIRT1 and SIRT3 in mediating polarization of the *Salmonella* infected macrophages toward an anti-inflammatory state. Upon knockdown of SIRT1 and SIRT3 in the infected macrophages we detect robust pro-inflammatory response and oxidative burst. This is in line with the findings by S.Elsela et al., wherein SIRT1 knockout RSV(Respiratory Syncytial Virus) - infected BMDCs showed significant increase in *Il1β*, *Il6* and *Il23* expression and ROS generation in comparison to the wild type RSV-infected BMDCs[35]. Also, Kim et al., showed presence of aggravated inflammatory responses in *Mycobacterium tuberculosis* infected SIRT3-/- BMDMs[38]. This heightened pro-inflammatory cytokine and oxidative burst restrict the intracellular survival of the pathogen as detected by the lower intracellular bacterial burden in the SIRT1 and SIRT3 knockdown murine macrophages. *Salmonella* showed enhanced proliferation in the M2 macrophages owing to the reduced arsenals in terms of pro-inflammatory cytokines and ROS production. Moreover, the M2 macrophages are fuelled by increased fatty acid oxidation and reduced glycolysis[62]. This might facilitate enhanced bacterial proliferation as the host un-utilized intracellular glucose can be readily up taken by the pathogen and used to support its own glycolysis. Similarly, M1 or pro-inflammatory macrophages resort to glycolysis to meet their energy demands[63] thereby limiting the glucose availability for the intracellular pathogen[63][64]. In such condition, bacteria show enhanced fatty acid metabolism to sustain their energy demand[65]. In our study, we found that wild type *S.* Typhimurium infection drive host metabolism toward increased fatty acid oxidation via its SPI-2 effector protein with concomitant increase in the bacterial glycolysis. SIRT1 and SIRT3 inhibition abrogates the metabolic switch and triggers increase in host glycolysis which in turn skew the bacterial metabolism from increased glycolysis to enhanced fatty acid oxidation and reduced

glycolysis. Together, these findings implicate the role of SIRT1 and SIRT3 in reprograming

the host metabolism which in turn affect the intracellular nutrient niche of the pathogen

thereby influencing intracellular *Salmonella* proliferation. However, our *in vivo* findings in

the murine model of infection show increased bacterial burden upon SIRT1 or SIRT3

inhibition. This increased burden could be attributed to increased dissemination from the

macrophages into the bloodstream owing to the increased level of serum IL-6 levels. This is

in concert with previous findings in *Klebsiella pneumoniae* infection in mice wherein

increased inflammatory response upon HIF-1α activation induces bacterial dissemination

[66]. Further correlation analysis of immune responses to *Salmonella* infection revealed that

increased innate immune "cassette" opposes the adaptive immune arm leading to increased

bacterial load [67]. Future studies might explore the host and bacterial interacting partners of

SIRT1 and SIRT3 through mass spectrometry analyses in *Salmonella* infected macrophages

which might hint at the underlying mechanism of their action and regulation. Together, this

study highlights the complex and multifaceted nature of host-pathogen interactions, and the

need for further research to fully understand the role of SIRT1 and SIRT3 in the context of

Salmonella infection.

MATERIALS AND METHODS

Bacterial Strains, and culture conditions

Salmonella enterica serovar Typhimurium (STM) strain ATCC 14028S or ATCC 14028S

constitutively expressing green fluorescent protein (eGFP) or mCherry (RFP) through pFPV

plasmid were used. 4% paraformaldehyde fixed STM (PFA) was used as the killed fixed

bacteria control. The above-mentioned live bacterial strain was grown overnight in LB broth

in 37 ^{0}C at 160 rpm shaking condition in presence or absence of appropriate antibiotic

(Ampiciilin-50µg/ml) (Table-1) after revival of the bacterial strains from glycerol stock

(stored at -80 ^{0}C).

Table:1: List of strains and plasmids used in this study

Strains/Plasmids	Characteristics	Source
Salmonella enterica serovar Typhimurium ATCC strain 14028S (STM)	Wild Type (WT)	Gifted by Prof. M. Hensel
pFPV:GFP	Amp^R	Laboratory Stock
pFPV:mCherry	Amp^R	Laboratory Stock
pLKO.2 : shRNA	Amp^R	Gifted by Prof. Subba Rao Gangi Setty
STM Δ *invC*	Kan^R	Laboratory Stock
STM Δ *ssaV*	Chl^R	Laboratory Stock
STM Δ *steE*	Kan^R	Laboratory Stock

Cell Culture

RAW 264.7 murine macrophages were cultured in DMEM (Lonza) containing 10% FBS

(Gibco) at 37 ^{0}C in a humified incubator with 5% CO_2. Prior to each experiment, cells were

seeded into 24 well or 6 well plate as per requirement at a confluency of 60%.

For macrophage polarization experiments, the seeded macrophages were subjected to

100ng/ml LPS + 20ng/ml IFN-γ treatment for M1 polarization and 20ng/ml IL-4 treatment for M2

polarization for 24 hrs. Post- polarization, the cell supernatant were collected for validation of

polarization status by ELISA and was further subjected to infection protocol.

Peritoneal macrophages were collected in PBS from the peritoneal cavity of C57BL/6 mice

aseptically post thioglycolate treatment using 20G needle and 5ml syringe. Following

centrifugation, cell pellet was resuspended in RPMI-1640 (Lonza) containing 10% heat-

inactivated FBS (Gibco), 100 U/ml penicillin and 100 µg/ml streptomycin and seeded into 6

well- plate. 6hr prior to infection, antibiotic-containing media was replaced with Penicillin-

Streptomycin free RPMI-1640 (Lonza) containing 10% heat-inactivated FBS (Gibco).

Transfection

shRNA mediated knockdown was carried out by PEI mediated transfection protocol. Plasmid

harbouring shRNA in pLKO.2 vector backbone specific to SIRT1 and SIRT3 were used for

transfection. Plasmid harbouring scrambled sequence of shRNA, served as a control, was also

used for transfection. Plasmid DNA was used at a concentration of 500ng and 1μg per well of

a 24-well plate and 6-well plate respectively. Plasmid and PEI were added in 1:2 ratio in

serum free DMEM media and incubated for 20 mins at room temperature. Post incubation,

the DNA: PEI cocktail was added to the seeded RAW 264.7 macrophages. After 6-8hrs of

incubation, serum-free media was replaced with complete media containing 10% FBS. Post

48hr of transfection, transfected cells were either harvested for knockdown confirmation

studies or subjected to infection with STM.

Table 2: List of shRNA used for knockdown

Sirtuins	Construct ID	TRC ID	Sequence (5'---3')
SIRT1	C4	TRCN0000218734	GTACCGGCATGAAGTGCCTCAGATAT TACTCGAGTAATATCTGAGGCACTTC ATGTTTTTTG
SIRT1	C12	TRCN0000018979	CCGGGCAAAGCCTTTCTGAATCTATC TCGAGATAGATTCAGAAAGGCTTTGC TTTTT
SIRT3	E8	TRCN0000038889	CCGGCCCAACGTCACTCACTACTTTC TCGAGAAAGTAGTGAGTGACGTTGG GTTTTTG
SIRT3	E12	TRCN0000038893	CCGGCCACCTGCACAGTCTGCCAAAC

			TCGAGTTTGGCAGACTGTGCAGGTGG TTTTTG

Infection Protocol

Macrophages were infected with stationary-phase bacterial culture with MOI of 10. For synchronization of the infection, tissue culture plates were subjected to centrifugation at 600xg for 5 min and incubated at 37 ^{0}C humified incubator with 5% CO_2 for 25 min. Cells were washed with PBS and were treated with DMEM (Sigma) + 10% FBS (Gibco) containing 100 µg/ml gentamicin for 1 hr. Subsequently, the gentamicin concentration was reduced to 25 µg/ml and maintained until the cells were harvested. For the inhibitor treatment studies, along with 25 µg/ml containing complete media 1µM of SIRT1 (EX-527) inhibitor or SIRT3 (3TYP) or 10mM of N-Acetyl Cysteine (NAC, Sigma) or 50nM of chetomin (Sigma) were added to the cells.

Immunofluorescence confocal microscopic studies

At the specified time points post infection with GFP tagged STM, cells were fixed with 3.5% paraformaldehyde for 15 min. Primary antibody staining was performed with specific primary antibody in the presence of a permeabilizing agent, 0.01% saponin (Sigma) dissolved in 2.5% BSA containing PBS at 4°C for overnight or for 6hr at room temperature (RT). Following this, cells were washed with PBS stained with appropriate secondary antibody tagged with fluorochrome for 1 hr at RT. This was followed by DAPI staining and mounting of the coverslip onto a clean glass slide using the mounting media containing the anti-fade agent. The coverslip sides were sealed with a transparent nail paint. All immunofluorescence images were obtained using Zeiss LSM 710 or Zeiss LSM 880 and were analyzed using ZEN black 2012 software.

Quantitative Real Time PCR

Total RNA was isolated at specific time points post infection by using TRIzol (Takara) as per manufacturer's protocol. Quantification of RNA was performed in Nano Drop (Thermo-Fischer scientific). Quality of isolated RNA was detected by performing 2% agarose gel electrophoresis. 2 µg of RNA was subjected to DNaseI (Thermo Fischer Scientific) treatment at 37°C for 1 hr followed by addition of 0.5M EDTA (final concentration 5mM) and heat inactivation at 65°C for 10 mins. The mRNA was reverse transcribed to cDNA using oligo $(dT)_{18}$ primer, buffer, dNTPs and reverse transcriptase (Takara) as per manufacturer's protocol. The expression profile of target gene was evaluated using specific primers (Table-3) by using SYBR green RT-PCR master mix (Takara) in BioRad Real time PCR instrument. β-actin was used as an internal control for mammalian genes and for bacterial genes 16S rRNA was used. All the reaction was setup in 384 well plate with two replicates for each sample.

Table 3: List of primers

Primer Name	Sequence (5'---3')
SIRT1 FP	ACAAAGTTGACTGTGAAGCTGTAC
SIRT1 RP	GTTCATCAGCTGGGCACCTA
SIRT3 FP	GCTGCTTCTGCGGCTCTATAC
SIRT3 RP	GAAGGACCTTCGACAGACCGT
PPARδ FP	CACAACGCTATCCGCTTTGG
PPARδ RP	ATGCTCCGGGCCTTCTTTTT
βactin FP	CAGCAAGCAGGAGTACGATG
βactin RP	GCAGCTCAGTAACAGTCCG
Acadl FP	CTTGGGAAGAGCAAGCGTACT
Acadl RP	CTGTTCTTTTGTGCCGTAATTCG

Hadha FP	AGCAACACGAATATCACAGGAAG
Hadha RP	AGGCACACCCACCATTTTGG
Acox1 FP	TCGAAGCCAGCGTTACGAG
Acox1 RP	ATCTCCGTCTGGGCGTAGG
Pdha1 FP	TGTCGGTTCCCAGTCCATCA
Pdha1 RP	CGTTTCCTTTTCACAGCACATGA
Pfkl FP	GAACTACGCACACTTGACCAT
Pfkl RP	CTCCAAAACAAAGGTCCTCTGG
ptsG FP	TATCTGGGCTTCTTTGCGGG
ptsG RP	ACCAGGCAACGCTCGATAAA
fadA FP	TCTGGGATTGATGGAGCAGA
fadA RP	AGACCAATACACATCGTCGC
fadB FP	GTCCCCGAAGAGCAGTTAAG
fadB RP	CCAGTTTAGTTTCCGGCAGA
aceA FP	CGATCTGGTATGGTGCGAAA
aceA RP	TTCTGCCAGTTGAAGGATGG
aceB FP	AGCGTTTCAATCAACAGGGT
aceB RP	CCCGTATTTTTACCTGCCGA
Salmonella 16S rRNA FP	GTGAGGTAACGGCTCACCAA
Salmonella 16S rRNA RP	TAACCGCAACACCTTCCTCC

Intracellular proliferation or gentamicin protection assay

Following infection of the transfected cells with STM at an MOI of 10, cells were treated

with DMEM (Sigma) + 10% FBS (Gibco) containing 100 μg/ml gentamicin for 1 hr.

Subsequently, the gentamicin concentration was reduced to 25 µg/ml and maintained until the

specified time point. Post 2hr and 16hr post-infection, cells were lysed in 0.1% triton-X-100.

Lysed cells were serially diluted and plated on *Salmonella-Shigella* (SS) agar to obtain

colony-forming units (cfu). Fold proliferation was calculated as cfu at 16hr divided by cfu at

2hr.

Fold Proliferation= [CFU at 16h]/[CFU at 2h]

Western Blotting

Post appropriate time points of infection, the cells were washed in PBS and subsequently

harvested in PBS. The cell pellets were obtained after centrifugation at 300g for 7 minutes at

4ºC. Cells were lysed in 1X RIPA (10X-0.5M NaCl, 0.5M EDTA pH-8.0, 1M Tris, NP-40,

10% sodium deoxycholate, 10% SDS) buffer containing 10% protease inhibitor cocktail

(Roche) for 30 min on ice. Total protein was estimated using Bradford (Bio-Rad) method of

protein estimation. Protein samples were subjected to 12 % SDS polyacrylamide gel

electrophoresis and then were transferred onto 0.45µm PVDF membrane (18V, 2 hrs). The

membrane was blocked using 5% skim milk in TBST (Tris Buffered Saline containing 0.1%

Tween-20) for 1hr at RT and subsequently probed with appropriate primary antibody (Table-

4) for overnight at 4°C. Following wash in TBST, blot was probed with specific HRP

conjugated secondary antibody for 1hr at RT. The membrane was developed using ECL

(Advansta) and images were captured using ChemiDoc GE healthcare. All densitometric

analysis was performed using ImageJ software.

Table 4: List of Antibodies

Antibody Name	Catalogue Number

SIRT1 Polyclonal Antibody	Cat #PA5-85921 (ThermoFisher Scientific)
SIRT3 Polyclonal Antibody	**Cat#** PA5-13222(ThermoFisher Scientific)
CD 86	**Cat#**12-086282PE(ThermoFisherScientific)
CD 206	**Catalog#** 17-2069-42 (ThermoFisherScientific)
F4/80 Antibody	BD Horizon, Cat-565411
HIF-1α	**sc-13515 SCBT**
NF-κβ p65	CST(D14E12)XP$^{®}$ RabbitmAb #8242
Acetylated-Lysine Antibody	**Cat #9441(Cell Signaling Technology)**
PDHA1	**Catalog No. A13687 (ABclonal)**
HADHA	**Catalog No. A5346 (ABclonal)**
ACOX1	**Catalog No. A8091 (ABclonal)**
PGK	**Catalog No. A12686 (ABclonal)**
PFK	**Catalog No. A5477 (ABclonal)**
β actin	A3854 (Sigma)

Immunoprecipitation

For co-immunoprecipitation, cells were washed with PBS and were lysed in native lysis

buffer containing 1% Nonidet P-40, 20 mM Tris (pH 8), 2 mM EDTA, 150 mM NaCl and

protease inhibitors mixture (Roche Diagnostics) for 30 min at 4°C. Cell debris was removed

by centrifugation at 10,000 rpm for 10 min and the supernatant was treated with the specific

antibody against the protein to be precipitated. Antibody-lysate complexes were

immunoprecipitated using Protein A/G- linked magnetic beads (MagGenome) according to

the manufacturer's protocol. Beads were extensively washed with washing buffer and

denatured at 95 °C for 10 min. Denatured precipitates were subjected to SDS-PAGE (12%

gel) followed by transfer to 0.45 µ PVDF membrane. The membrane was blocked using 5%

skimmed milk in TBST (Tris Buffered Saline containing 0.1% Tween-20) for 1h at room

temperature and eventually probed for the target primary antibodies or Anti-acetylated Lysine

(Ac-K) primary antibody overnight at 4°C. The blot was probed with a specific HRP-

conjugated secondary antibody for 1hr at RT after rigorous washing in TTBS. ECL (BioRad)

was used for detection and images were captured using ChemiDoc GE healthcare.

ELISA

Estimation of cytokines in cell-free supernatant or in mice serum was performed according

to the manufacturer's instructions. Briefly, 96-well ELISA plates (BD Bioscience) were

coated overnight with capture antibody at 4°C. Following day, plates were washed with 0.1%

Tween-20 containing PBS and blocked with 10% FBS for 1 h. Following blocking, wells

were washed and incubated with 100 µL of test samples for 2 h at room temperature.

Subsequently, plates were washed and incubated with detection antibody and enzyme reagent

for 1 h at room temperature (BD Bioscience). TMB (Sigma) was used as a substrate and

reactions were stopped with 2 N H_2SO_4. For the estimation of IL-1β, pre-coated ELISA

(SARD Biosciences) plates were used, and ELISA was performed as per manufacturer's

protocol. Absorbance was measured at 450 nm wavelength in Tecan Plate reader and the

concentration of cytokines were interpolated from a standard curve.

Flow cytometry

After specific time points post infection, cells were washed and harvested in PBS. Following

centrifugation, cell pellet was resuspended in FACS buffer comprised of 1% BSA in PBS.

Blocking was performed with Fc blocker (purified Anti-mouse CD16/CD32, eBioscience)

dissolved in FACS blocking buffer for 30 min on ice. Following a washing step with PBS,

antibody staining was performed with PE-conjugated CD86 antibody or APC-conjugated

CD206 (Thermo Scientific) for 45 min on ice. After washing in PBS, the cell pellet was

resuspended in 1% PFA in PBS. Subsequently, PFA was removed, and cells were

resuspended in FACS buffer and reading was taken in BD FACSVerse instrument. For flow

cytometry studies in mice tissues , the harvested liver or spleen was homogenized into single-

cell suspension post RBC lysis (RBC lysis Buffer, Sigma). The homogenized cell suspension

was washed and resuspended in FACS buffer containing the PE-conjugated Rat anti-mouse

F4/80 antibody (BD Horizon, Cat-565411) or eFluor450-conjugated anti-mouse CD11c

antibody (eBioscience), or FITC-conjugated anti-mouse Ly6C antibody (BD Pharmingen) or

Alexa647-conjugated anti-mouse CD45 antibody (BioLegend). Following staining protocol,

the cells were washed in PBS and the cells were resuspended in FACS buffer and the FACS

protocol was performed either in BD FACSVerse or CytoFLEX flow cytometer (Beckman).

For DCFDA staining, one hour before the indicated time point of infection, cells were

incubated with 10μM DCFDA containing DMEM media at 37°C humidified incubator with

5% CO_2 for 45 min. Post incubation, cells were washed and harvested in PBS. Readings were

measured in BD FACSVerse instrument.

All analysis was done using BD FACSuite software.

Phenol Red-Hydrogen Peroxidase Assay

712 Post 48hr of transfection, cells were infected with STM culture at an MOI of 10. Cells were

713 incubated with Phenol Red solution containing hydrogen peroxidase enzyme (8.5U/ml). At,

714 the designated time points post-infection, the supernatant was collected, and the absorbance

715 was taken at 610 nm in Tecan Plate reader. The exogenously produced H2O2 was quantified

716 using a standard curve of known concentration of H_2O_2.

Gene expression studies by nanoString nCounter technology

Total RNA was isolated at specific time points post infection by using TRIzol (Takara) as

per manufacturer's protocol. Quantification of RNA was performed in Nano Drop (Thermo-

Fischer scientific) and Qubit Bioanalyzer (Agilent 2100 Bioanalyzer). Quality of isolated

RNA was detected by performing 2% agarose gel electrophoresis. Post quality check,

samples were subjected to nanoString nCounter technology (theraCUES). This technology

allows multi-plex, spatially resolved RNA expression quantification with appropriate probes

designed against the target gene.

Lactate Estimation Assay

Cell-supernatant was harvested at the specific time-points post-infection and the lactate

content of the sample was estimated using the Lactate Assay Kit (Sigma, Catalog Number-

MAK064) as per manufacturer's protocol. Briefly, 50µl of the sample was added to each 96

well-plate and each of the well 50µl of the master reaction mix containing 46µl of lactate

assay buffer, 2µl of lactate enzyme mix, and 2µl of lactate probe was added. After the

addition of the master reaction mix to the sample, they were mixed by horizontal shaker or

via pipetting. The plate was incubated for 30 minutes in dark at room temperature. Post

incubation, absorbance was measured at 570nm. The lactate content of the sample was

estimated from the lactate standard curve ranging from 0-10 nmole/µl.

Fatty Acid Oxidation Assay

At 16h post-infection, cell pellets were harvested and stored at-80ºC. The fatty acid oxidation assay protocol was followed as per manufacturer's instructions (AssayGenie Fatty Acid Oxidation (FAO) Assay Kit, Catalogue Code-BR00001). Briefly, cell pellets were lysed using 1X Cell lysis buffer (provided in the kit) and the cell supernatant were obtained post centrifugation of the cell lysate in a cold microfuge at 14,000rpm for 5min. The protein content of the cell supernatants was estimated using Bradford (Bio-Rad) method. 20µl of the protein sample was added to each 96-well plate in duplicate on ice. Each sample was treated with 50µl of control solution and 50µl of reaction solution by swiftly adding one 50µl of control solution to one set of wells and 50µl of reaction solution to the other set of wells. The contents were gently mixed for 10s. The plate is covered and incubated in a 37ºC incubator for 30-60 min (without CO_2). After incubation cherry red colour appears in the wells. The O.D. is measured at 492nm using a plate reader at 30 min,60 min or 120 min. The control well reading was subtracted from the reaction well reading for each sample for each time point. The subtracted reading is used for enzyme activity calculation by considering the incubation time.

Animal Experiment

For all experiments, 6–8 weeks old C57BL/6 or *gp91phox*-/- mice were used. For organ burden analysis, 6 weeks old C57BL/6 or *gp91phox*-/- mice were infected with 10^7 CFU bacteria via oral gavaging. For bacterial enumeration via flow cytometry, mice were infected with 10^7 GFP-expressing bacteria orally. Infected mice were intraperitoneally injected on every other day with either 1 mg/kg body weight of SIRT1 inhibitor EX-527 (Sigma-Aldrich) [32], or SIRT3 inhibitor 3TYP (Selleck Chemical) [33] or SIRT1 activator SRT1720 (Calbiochem, Sigma-Aldrich) [34] or treated with vehicle alone. 5 days post infection, mice

were sacrificed, and bacterial organ load was estimated by plating the tissue homogenates on

SS agar plates. For calculating percentage survival, 6 weeks old C57BL/6 mice were infected

with 10^8 bacteria orally and monitored till fatality. For flow cytometry studies, the harvested

liver or spleen were homogenized into single cell suspension and were subjected to flow

cytometry. The animal experiments were carried out in accordance with the approved

guidelines of the institutional animal ethics committee at the Indian Institute of Science,

Bangalore, India (Registration No: 48/1999/CPCSEA). All procedures involving the use of

animals were performed according to the Institutional Animal Ethics Committee (IAEC)-

approved protocol.

***In vivo* knockdown**

For in vivo knockdown adeno-associated virus serotype 6 (AAV6) was used. AAV6 viruses

were produced in HEK293T cells. Briefly, HEK293T cells were transfected with AAV

plasmid encoding shRNAs targeting SIRT1, SIRT3 and scramble control under U6 promoter

together with helper plasmid using PEI Max 40000 (Polysciences USA). Next day

transfection medium was removed, and fresh culture medium was added. Forty-eight hours

later, medium was collected, and fresh culture medium was added and incubated for next

forty-eight hours. Again, culture medium was collected and cells were harvested by

trypsinization. Collected medium containing secreted AAV viral particles was incubated with

40% PEG 8000 overnight. Precipitated viral particles were then collected by centrifugation

and re-suspended in PBS. Cells containing AAV viral particles were then lysed with citrate

buffer and incubated with 40% PEG 8000 and processed similarly to medium for viral

particle precipitation. Precipitated viral particles were then cleaned with chloroform and

loaded on iodixanol gradient for further cleaning using ultracentrifugation. Cleaned viral

particles were collected from the forty percent gradient and loaded on Amicon 100K cut-off

columns. Purified AAV particles were then titrated by real-time PCR. For infection of AAV6,

the mice were intravenously injected with a volume of 200 μl containing approximately 10^{12}

viral particles harbouring the respective scrambled or shRNA constructs. The mice were

infected at 7^{th} day after injection of the virus with 10^6 cfu units of *S*. Typhimurium orally.

Fifth day post-infection, mice were euthanized, and dissected out for organ harvesting and

blood collection. The knockdown was validated by performing western blotting and qPCR of

the harvested liver tissue (Fig. S10).

Haematoxylin and Eosin Staining

6-8 weeks old C57BL/6 mice were infected with 10^7 bacteria orally. Infected mice were

intraperitoneally injected every alternate with either 1 mg/kg body weight of SIRT1 inhibitor

EX-527, or SIRT3 inhibitor 3TYP or SIRT1 activator SRT1720 or treated with vehicle alone.

5 days post infection, mice were euthanized, and livers were collected and fixed using 3.5%

paraformaldehyde. The fixed liver was then dehydrated using a gradually increasing

concentration of ethanol and embedded in paraffin. 5μm sections were collected on coated

plates. Sections were further rehydrated and then stained with hematoxylin and eosin. Images

were collected in a Leica microscope. **Scoring system:-** according to pathological changes

the tissue sections are scored as 0 for normal pathology, 1 for mild/ minor pathology, 2 for

moderate pathology, and 3 for severe pathological changes.

Statistical analysis

Data were analyzed and graphed using the GraphPad Prism 8 software (San Diego, CA).

Statistical significance was determined by Student's t-test or Two-way ANOVA and

Bonferroni post-t-test to obtain p values. Adjusted p-values below 0.05 are considered

statistically significant. The results are expressed as mean ± SD or SEM of three independent

experiments.

ACKNOWLEDGEMENT

We duly thank Prof. Subba Rao Gangi Setty and Prof. Michael Hensel for providing us with

the shRNA knockdown constructs and the *S.* Typhimurium 14028S strain, respectively.

FUNDING

This work was supported by the DAE SRC fellowship (DAE00195) and DBT-IISc partnership

umbrella program for advanced research in biological sciences and Bioengineering to DC.

Infrastructure support from ICMR (Centre for Advanced Study in Molecular Medicine), DST

(FIST), and UGC (special assistance) is highly acknowledged. DH sincerely acknowledges the

CSIR- SPM fellowship for her financial support. SKG is supported by Ramalingaswami Re-

entry Fellowship BT/RLF/re-entry/14/2019 from DBT, Government of India. The funders

had no role in study design, data collection and analysis, decision to publish, or preparation of

the manuscript.

AUTHOR CONTRIBUTION

DH and DC have conceptualized the study. DH has contributed to the experiment designing,

visualization, methodology, investigation, formal analysis, literature survey, validation,

writing (original draft), reviewing, and editing of the manuscript. RSR has performed the

histopathological scoring of the liver sections and the retro-orbital injection of the adenoviral

constructs in mice. ADC and SKG have prepared and provided AAV6, for *in vivo* mice

knockdown experiments. DC has contributed to the funding acquisition, project

administration, and overall supervision of the work. All authors approved the final version of

the manuscript.

DECLARATION OF INTEREST

The authors are unaware of any conflicting interests and thereby declare no conflict of

interest.

REFERNCES

1. Landry J, Sutton A, Tafrov ST, Heller RC, Stebbins J, Pillus L, et al. The silencing protein SIR2 and its
homologs are NAD-dependent protein deacetylases. Proc Natl Acad Sci U S A. 2000;97(11):5807–11.

2. Jackson MD, Denu JM. Structural identification of 2'- and 3'-O-acetyl-ADP-ribose as novel
metabolites derived from the Sir2 family of beta -NAD+-dependent histone/protein deacetylases. J
Biol Chem. 2002;277(21):18535-18544. doi:10.1074/jbc.M200671200

3. Sauve AA, Celic I, Avalos J, Deng H, Boeke JD, Schramm VL. Chemistry of gene silencing: The
mechanism of NAD+-dependent deacetylation reactions. Biochemistry. 2001;40(51):15456–63.

4. Bitterman KJ, Anderson RM, Cohen HY, Latorre-Esteves M, Sinclair DA. Inhibition of silencing and
accelerated aging by nicotinamide, a putative negative regulator of yeast Sir2 and human SIRT1. J
Biol Chem. 2002;277(47):45099–107.

5. Sanders BD, Jackson B, Marmorstein R. Structural basis for sirtuin function: What we know and
what we don't. Biochim Biophys Acta - Proteins Proteomics 2010;1804(8):1604–16. Available from:
http://dx.doi.org/10.1016/j.bbapap.2009.09.009 10. Lin Z, Fang D. The Roles of SIRT1 in Cancer.
Genes and Cancer. 2013;4(3–4):97– 104.

6. Lin Z, Fang D. The Roles of SIRT1 in Cancer. Genes and Cancer. 2013;4(3–4):97–104.

7. Guarente L. Sirtuins in aging and disease. Cold Spring Harb Symp Quant Biol. 2007;72:483–8.

8. Martínez-Redondo P, Vaquero A. The Diversity of Histone Versus Nonhistone Sirtuin Substrates.
Genes and Cancer. 2013;4(3–4):148–63.

9. Liu TF, Yoza BK, El Gazzar M, Vachharajani VT, McCall CE. NAD+-dependent SIRT1 deacetylase
participates in epigenetic reprogramming during endotoxin tolerance. J Biol Chem.
2011;286(11):9856–64.

10. El Gazzar M, Yoza BK, Hu JYQ, Cousart SL, McCall CE. Epigenetic silencing of tumor necrosis factor
α during endotoxin tolerance. J Biol Chem. 2007;282(37):26857–64.

11. Liu TF, Vachharajani V, Millet P, Bharadwaj MS, Molina AJ, McCall CE. Sequential actions of
SIRT1-RELB-SIRT3 coordinate nuclear-mitochondrial communication during immunometabolic
adaptation to acute inflammation and sepsis. J Biol Chem. 2015;290(1):396–408.

12. Kong X, Wang R, Xue Y, et al. Sirtuin 3, a new target of PGC-1alpha, plays an important role in the
suppression of ROS and mitochondrial biogenesis. PLoS One. 2010;5(7):e11707.

13. van de Ven RAH, Santos D, Haigis MC. Mitochondrial Sirtuins and Molecular Mechanisms of
Aging. Trends Mol Med. 2017;23(4):320–31.

14. Garai, P., D.P. Gnanadhas, and D. Chakravortty, Salmonella enterica serovars Typhimurium and
Typhi as model organisms: revealing paradigm of host-pathogen interactions. Virulence, 2012. 3(4):
p. 377-88

15.Bhutta ZA, Threlfall J. Addressing the global disease burden of typhoid
fever. *JAMA*.2009;302(8):898-899. doi:10.1001/jama.2009.1259

16. Majowicz SE, Musto J, Scallan E, et al. The global burden of nontyphoidal Salmonella
gastroenteritis. Clin Infect Dis. 2010;50(6):882-889. doi:10.1086/650733

17. Haraga, A., M.B. Ohlson, and S.I. Miller, Salmonellae interplay with host cells. Nat Rev Microbiol,
2008. **6**(1): p. 53-66.

18. Hajra, D., A.V. Nair, and D. Chakravortty, An elegant nano-injection machinery for sabotaging the
host: Role of Type III secretion system in virulence of different human and animal pathogenic
bacteria. Phys Life Rev, 2021. **38**: p. 25-54.

19. Lawrence T, Natoli G. Transcriptional regulation of macrophage polarization: Enabling diversity
with identity. Nat Rev Immunol. 2011;11(11):750–61.

20. Martinez FO, Helming L, Gordon S. Alternative Activation of Macrophages: An Immunologic
Functional Perspective. Annu Rev Immunol. 2009;27(1):451–83.

21. Namgaladze D, Brüne B. Fatty acid oxidation is dispensable for human macrophage IL-4-induced
polarization. Biochim Biophys Acta - Mol Cell Biol Lipids 2014;1841(9):1329–35.

22. Hui X, Zhang M, Gu P, Li K, Gao Y, Wu D. Adipocyte SIRT 1 controls systemic insulin sensitivity by
modulating macrophages in adipose tissue. 2017;18(4):645–57.

23. Jia Y, Han S, Li J, et al. IRF8 is the target of SIRT1 for the inflammation response in macrophages.
Innate Immun. 2017;23(2):188-195. doi:10.1177/1753425916683751

24. Kratz M, Coats BR, Hisert KB, Hagman D, Mutskov V, Peris E, et al. Metabolic dysfunction drives a
mechanistically distinct proinflammatory phenotype in adipose tissue macrophages. Cell Metab
2014;20(4):614–25. Available from: http://dx.doi.org/10.1016/j.cmet.2014.08.010

25. Cimen H, Han MJ, Yang Y, Tong Q, Koc H, Koc EC. Regulation of succinate dehydrogenase activity
by SIRT3 in mammalian mitochondria. Biochemistry. 2010;49(2):304–11.

26. Vazquez-Torres A, Xu Y, Jones-Carson J, Holden DW, Lucia SM, Dinauer MC, et al. Salmonella
pathogenicity island 2-dependent evasion of the phagocyte NADPH oxidase. Science (80-).
2000;287(5458):1655–8.

27. Miller BH, Fratti RA, Poschet JF, Timmins GS, Master SS, Burgos M, et al. Mycobacteria Inhibit
Nitric Oxide Synthase Recruitment to Phagosomes during Macrophage Infection. Infect Immun.
2004;72(5):2872–8.

28. Bost KL, Clements JD. Intracellular Salmonella dublin induces substantial secretion of the 40-
kilodalton subunit of interleukin-12 (IL-12) but minimal secretion of IL-12 as a 70-kilodalton protein
in murine macrophages. Infect Immun. 1997;65(8):3186–92.

29. Pathak SK, Basu S, Basu KK, Banerjee A, Pathak S, Bhattacharyya A, et al. Direct extracellular
interaction between the early secreted antigen ESAT-6 of Mycobacterium tuberculosis and TLR2
inhibits TLR signaling in macrophages. Nat Immunol. 2007;8(6):610–8.

30. Tumitan ARP, Monnazzi LGS, Ghiraldi FR, Cilli EM, De Medeiros BMM. Pattern of macrophage
activation in yersinia-resistant and yersinia-susceptible strains of mice. Microbiol Immunol.
2007;51(10):1021–8.

31. Brubaker RR. Interleukin-10 and inhibition of innate immunity to Yersiniae: Roles of Yops and
LcrV (V antigen). Infect Immun. 2003;71(7):3673–81.

32. Morató L, Astori S, Zalachoras I, et al. eNAMPT actions through nucleus accumbens NAD$^+$/SIRT1
link increased adiposity with sociability deficits programmed by peripuberty stress. Sci Adv.
2022;8(9):eabj9109. doi:10.1126/sciadv.abj9109.

33. Hu B, Wang P, Zhang S, et al. HSP70 attenuates compression-induced apoptosis of nucleus
pulposus cells by suppressing mitochondrial fission via upregulating the expression of SIRT3. Exp Mol
Med. 2022;54(3):309-323. doi:10.1038/s12276-022-00745-9.

34. Tan P, Wang M, Zhong A, et al. SRT1720 inhibits the growth of bladder cancer in organoids and
murine models through the SIRT1-HIF axis. *Oncogene*. 2021;40(42):6081-6092. doi:10.1038/s41388-
021-01999-9.

35. Elesela S, Morris SB, Narayanan S, Kumar S, Lombard DB, Lukacs NW. Sirtuin 1 regulates
mitochondrial function and immune homeostasis in respiratory syncytial virus infected dendritic
cells. PLoS Pathog. 2020;16(2):e1008319.

36. Liu G, Bi Y, Xue L, et al. Dendritic cell SIRT1-HIF1α axis programs the differentiation of CD4+ T
cells through IL-12 and TGF-β1. Proc Natl Acad Sci U S A. 2015;112(9):E957-E965.
doi:10.1073/pnas.1420419112.

37. Yang H, Hu J, Chen Y (Jason), Ge B. Role of Sirt1 in innate immune mechanisms against
Mycobacterium tuberculosis via the inhibition of TAK1 activation. Arch Biochem Biophys
2019;667(October 2018):49–58. Available from: https://doi.org/10.1016/j.abb.2019.04.006

38. Kim TS, Jin YB, Kim YS, Kim S, Kim JK, Lee HM, et al. SIRT3 promotes antimycobacterial defenses
by coordinating mitochondrial and autophagic functions Vol. 15, Autophagy. Taylor & Francis; 2019.
1356–1375 p. Available from: https://doi.org/10.1080/15548627.2019.1582743.

39. Yeung F, Hoberg JE, Ramsey CS, et al. Modulation of NF-kappaB-dependent transcription and cell
survival by the SIRT1 deacetylase. EMBO J. 2004;23(12):2369-2380. doi:10.1038/sj.emboj.7600244.

40. Yang H, Zhang W, Pan H, et al. SIRT1 activators suppress inflammatory responses through
promotion of p65 deacetylation and inhibition of NF-κB activity. PLoS One. 2012;7(9):e46364.
doi:10.1371/journal.pone.0046364.

41. Singh CK, Chhabra G, Ndiaye MA, Garcia-peterson LM, Mack NJ, Ahmad N. The Role of Sirtuins in
Antioxidant and Redox Signaling. 2018;28(8):643–61.

42. Kitada M, Ogura Y, Monno I, Koya D. Sirtuins and Type 2 Diabetes: Role in Inflammation,
Oxidative Stress, and Mitochondrial Function. Front Endocrinol (Lausanne). 2019;10:187.

43. Merksamer PI, Liu Y, He W, Hirschey MD, Chen D, Verdin E. The sirtuins, oxidative stress and
aging: an emerging link. Aging (Albany NY). 2013;5(3):144-150. doi:10.18632/aging.100544.

44. Rendra E, Riabov V, Mossel DM, Sevastyanova T, Harmsen MC, Kzhyshkowska J. Immunobiology
Reactive oxygen species (ROS) in macrophage activation and function in diabetes. Immunobiology
2019;224(2):242–53. Available from: https://doi.org/10.1016/j.imbio.2018.11.010

45. Zhou Y, Que KT, Zhang Z, et al. Iron overloaded polarizes macrophage to proinflammation
phenotype through ROS/acetyl-p53 pathway. Cancer Med. 2018;7(8):4012-4022.
doi:10.1002/cam4.1670.

46. Halasi M, Wang M, Chavan TS, Gaponenko V, Hay N, Gartel AL. ROS inhibitor Nacetyl-L-cysteine
antagonizes the activity of proteasome inhibitors. Biochem J. 2013;454(2):201–8.

47.De Santa F, Vitiello L, Torcinaro A, Ferraro E. The Role of Metabolic Remodeling in Macrophage
Polarization and Its Effect on Skeletal Muscle Regeneration. Antioxid Redox Signal.
2019;30(12):1553-1598. doi:10.1089/ars.2017.7420.

48. Galván-Peña S, O'Neill LAJ. Metabolic reprogramming in macrophage polarization. Front
Immunol. 2014;5(AUG):1–6.

49. Taylor SJ, Winter SE. Salmonella finds a way: Metabolic versatility of Salmonella enterica serovar
Typhimurium in diverse host environments. PLoS Pathog 2020;16(6):10–5. Available from:
http://dx.doi.org/10.1371/journal.ppat.1008540

50. Kierans SJ, Taylor CT. Regulation of glycolysis by the hypoxia-inducible factor (HIF): implications
for cellular physiology. J Physiol. 2021;599(1):23-37. doi:10.1113/JP280572.

51. Lim JH, Lee YM, Chun YS, Chen J, Kim JE, Park JW. Sirtuin 1 modulates cellular responses to
hypoxia by deacetylating hypoxia-inducible factor 1alpha. Mol Cell. 2010;38(6):864-878.
doi:10.1016/j.molcel.2010.05.023.

52. Wang Y, Bi Y, Chen X, et al. Histone Deacetylase SIRT1 Negatively Regulates the Differentiation of
Interleukin-9-Producing CD4(+) T Cells. Immunity. 2016;44(6):1337-1349.
doi:10.1016/j.immuni.2016.05.009

53. Viziteu E, Grandmougin C, Goldschmidt H, et al. Chetomin, targeting HIF-1α/p300 complex,
exhibits antitumour activity in multiple myeloma. Br J Cancer. 2016;114(5):519-523.
doi:10.1038/bjc.2016.20

54. Hirschey MD, Shimazu T, Goetzman E, et al. SIRT3 regulates mitochondrial fatty-acid oxidation by
reversible enzyme deacetylation. Nature. 2010;464(7285):121-125. doi:10.1038/nature08778

55. Zhang Y, Wen P, Luo J, et al. Sirtuin 3 regulates mitochondrial protein acetylation and
metabolism in tubular epithelial cells during renal fibrosis. Cell Death Dis. 2021;12(9):847.

56. Panagi I, Jennings E, Zeng J, Günster RA, Stones CD, Mak H, et al. Salmonella Effector SteE
Converts the Mammalian Serine/Threonine Kinase GSK3 into a Tyrosine Kinase to Direct
Macrophage Polarization. Cell Host Microbe. 2020;27(1):41-53.e6.

57. Stapels DAC, Hill PWS, Westermann AJ, Fisher RA, Thurston TL, Saliba AE, et al. Salmonella
persisters undermine host immune defenses during antibiotic treatment. Science.
2018;362(6419):1156–60.

58. Pham THM, Brewer SM, Thurston T, Massis LM, Honeycutt J, Lugo K, et al. Salmonella-Driven
Polarization of Granuloma Macrophages Antagonizes TNFMediated Pathogen Restriction during
Persistent Infection. Cell Host Microbe. 2020;27(1):54-67.e5. Available from:
https://doi.org/10.1016/j.chom.2019.11.011

59. Ganesan R, Hos NJ, Gutierrez S, Fischer J, Stepek JM, Daglidu E, et al. Salmonella Typhimurium
disrupts Sirt1/AMPK checkpoint control of mTOR to impair autophagy. PLoS Pathog. 2017;13(2):1–
22.

60. Gogoi M, Chandra K, Sarikhani M, Ramani R, Sundaresan NR, Chakravortty D. Salmonella escapes
adaptive immune response via SIRT2 mediated modulation of innate immune response in dendritic
cells. PLoS Pathog. 2018;14(11):1–22.

61. Smulan LJ, Martinez N, Kiritsy MC, Kativhu C, Cavallo K, Sassetti CM, et al. Sirtuin 3
downregulation in mycobacterium tuberculosis-infected macrophages reprograms mitochondrial
metabolism and promotes cell death. MBio. 2021;12(1):1–15.

62. Eisele NA, Ruby T, Jacobson A, Manzanillo PS, Cox JS, Lam L, et al. Salmonella require the fatty
acid regulator PPARδ for the establishment of a metabolic environment essential for long-term
persistence. Cell Host Microbe 2013;14(2):171– 82. Available from:
http://dx.doi.org/10.1016/j.chom.2013.07.010

63. Cramer T, Yamanishi Y, Clausen BE, et al. HIF-1alpha is essential for myeloid cell mediated
inflammation. Cell. 2003;112(5):645-657. doi:10.1016/s0092-8674(03)00154-5.

64. Merrill GF, Kurth EJ, Hardie DG, Winder WW. AICA riboside increases AMP activated protein
kinase, fatty acid oxidation, and glucose uptake in rat muscle. Am J Physiol. 1997;273(6):E1107-
E1112. doi:10.1152/ajpendo.1997.273.6.E1107.

65. Reens AL, Nagy TA, Detweiler CS. Salmonella enterica Requires Lipid Metabolism Genes To
Replicate in Proinflammatory Macrophages and Mice. Infect Immun. 2019;88(1):e00776-19

66. Holden VI, Breen P, Houle S, Dozois CM, Bachman MA. Klebsiella pneumoniae Siderophores
Induce Inflammation, Bacterial Dissemination, and HIF-1α Stabilization during Pneumonia. mBio.
2016;7(5):e01397-16. Published 2016 Sep 13. doi:10.1128/mBio.01397-16

67. Hotson AN, Gopinath S, Nicolau M, et al. Coordinate actions of innate immune responses oppose
those of the adaptive immune system during Salmonella infection of mice. Sci Signal.
2016;9(410):ra4. Published 2016 Jan 12. doi:10.1126/scisignal.aaa9303

MAIN FIGURE LEGENDS

Fig.1- *Salmonella* modulates the expression of SIRT1 and SIRT3 along its course of

infection

A-B. Expression studies of *SIRT1* and *SIRT3* through qPCR in RAW 264.7 macrophages.

Data is representative of N=4, n=2. Unpaired two tailed Student's t test was performed to

obtain the p values. (****$p<0.0001$,***$p < 0.001$, ** $p<0.01$, *$p<0.05$)

1011

C-D-Expression studies of *SIRT1* and *SIRT3* through qPCR in peritoneal macrophages

derived from C57BL/6. Data is representative of N=3, n=2. . Unpaired two tailed Student's t

test was performed to obtain the p values. (****$p<0.0001$,***$p < 0.001$, ** $p<0.01$, *$p<0.05$)

E. Representative confocal images of RAW264.7 macrophages exhibiting SIRT1 expression

upon *S*. Typhimurium infection at indicated time points post infection. Data is representative

of N=3, n=80 (microscopic field).

1018

F. Quantitative representation of the expression profile as depicted in the confocal images (E)

in terms of Mean Fluorescence Intensity (MFI). Unpaired two tailed Student's t test was

performed to obtain the p values. (****p<0.0001,***p < 0.001, ** p<0.01)

G. Representative confocal images of RAW264.7 macrophages exhibiting SIRT3 expression

upon *S*. Typhimurium infection at indicated time points post infection. Data is representative

of N=3, n=80 (microscopic field).

H. Quantitative representation of the expression profile as depicted in the confocal images

(G) in terms of Mean Fluorescence Intensity (MFI). Unpaired two tailed Student's t test was

performed to obtain the p values. (****p<0.0001, *** p < 0.001, ** p<0.01)

I. qPCR mediated expression of *SIRT1* in RAW264.7 macrophages upon infection with

wildtype *S*. Typhimurium or SPI-1 ($\Delta invC$)or SPI-2 ($\Delta ssaV$ and $\Delta steE$) *m*utants of *S*.

Typhimurium. Data is representative of N=3,n=3. Unpaired two tailed Student's t test was

performed to obtain the p values. (****p<0.0001,***p < 0.001, ** p<0.01)

J. qPCR mediated expression of *SIRT3* in RAW264.7 macrophages upon infection with

wildtype *S*. Typhimurium or SPI-1 ($\Delta invC$) or SPI-2 ($\Delta ssaV$ and $\Delta steE$) *m*utants of *S*.

Typhimurium. Data is representative of N=3, n=3. Unpaired two tailed Student's t test was

performed to obtain the p values. (****p<0.0001,***p < 0.001, ** p<0.01)

Fig.2- Effect of SIRT1 and SIRT3 knockdown in intracellular bacterial proliferation

within RAW 264.7 and primary murine macrophages.

A. Fold Proliferation of *Salmonella* Typhimurium within RAW 264.7 macrophages in transfected and un-transfected conditions. Data is representative of N=3, n=3. Unpaired two tailed Student's t test was performed to obtain the p values. (** p<0.01, * p<0.05)

B. Fold Proliferation of *Salmonella* Typhimurium within infected peritoneal macrophages isolated from adult male C57BL/6 mice upon SIRT1 (EX-527) or SIRT3 (3TYP) inhibitor treatment. Unpaired two tailed Student's t test was performed to obtain the p values. (** p<0.01, * p<0.05)

Fig.3- *Salmonella* **Typhimurium skews the polarization state of the macrophage toward an immunosuppressive M2 state along the course of infection**

A. nanoString gene expression profiling data of *S*. Typhimurium infected RAW 264.7 macrophages versus uninfected control data sets at 2hr, 6hr and 16hr time points of infection. Data is representative of N=2,n=2.

B. Quantitative representation of flow cytometric analysis of alteration in M1 CD86 positive population in *S*. Typhimurium (STM) infected samples in comparison to uninfected (UI) and Paraformaldehyde Fixed (PFA) bacteria at the indicated time post-infection. Data is representative of N=2, n=3. Two-way ANOVA and Bonferroni post-t-test was used to obtain p values. (*** p < 0.001)

C. Quantitative representation of flow cytometric analysis of M2 surface marker CD206 in *S*. Typhimurium (STM) infected SIRT1 or SIRT3 knockdown RAW264.7 macrophages in comparison to the scrambled control at the indicated time post-infection. UI-Uninfected sample. Data is representative of N=3, n=3. Two-way ANOVA and Bonferroni post-t-test was used to obtain p values. (*** p < 0.001)

Fig.4- SIRT1 mediates modulation of immune functions via deacetylation of p65 subunit of NF-κB in *S*.Typhimurium infected macrophages.

A. Immunoblot depicting p65 NF-κB interaction with SIRT1 post immunoprecipitation of

SIRT1 in uninfected (UI) or *S*. Typhimurium (STM) infected RAW264.7 macrophages at

16hr post-infection. Data is representative of N=3, n=1

B. An immunoblot depicting p65 NF-κB interaction with SIRT1 as well as the p65 NF-κB

acetylation status post immunoprecipitation of p65 (IP: p65) or with control IgG (IP: IgG) in

uninfected (UI) or *S*. Typhimurium (STM) infected RAW264.7 macrophages upon

knockdown with SIRT1 shRNA or scrambled control.

C. Densitometric plot depicting the band intensities of Acetylated p65 over total p65 in blot

B.

D. An immunoblot depicting p65 NF-κB interaction with SIRT1 as well as the p65 NF-κB

acetylation status post immunoprecipitation of p65 (IP: p65) or with control IgG (IP: IgG) in

uninfected (UI) or *S*. Typhimurium (STM) infected RAW264.7 macrophages upon SIRT1

inhibitor (EX-527, 1μM) treatment at 16hr post-infection. UT- Untreated.

E. Densitometric plot depicting the band intensities of Acetylated p65 subunit of NF-κ B over

total p65 NF-κB in blot D.

Fig.5- *Salmonella* Typhimurium drives the metabolism of the infected macrophage

toward fatty acid oxidation

A. Metabolic gene expression data of *S*. Typhimurium infected RAW 264.7 macrophages at

2hr, 6hr and 16hr time points of infection through nanoString. Data is representative of N=2,

n=1.

B. *PPARδ* qPCR expression data in *S*. Typhimurium infected RAW 264.7 macrophages at the

indicated time points of infection. Data is representative of N=3, n=2.

C. Lactate estimation assay of *S*.Typhimurium infected RAW264.7 macrophages at the

initial time point of 2hr and at the late time point of 16hr post infection. Data is representative

of N=3, n=3. Unpaired two-tailed Student's t test was performed to obtain the p values. (**

p<0.01, * p<0.05)

D-E-Lactate estimation assay of *S*.Typhimurium infected RAW264.7 macrophages upon

SIRT1 (D) or SIRT3 (E) knockdown condition at16h post-infection. Data is representative of

N=3 , n=3. Unpaired two-tailed Student's t test was performed to obtain the p values.

F. Immunoblotting of host glycolytic (PGK), TCA cycle (PDHA1) and fatty acid oxidation

(HADHA, ACOX-1) proteins under SIRT1 and SIRT3 knockdown condition of *S*.

Typhimurium infected RAW264.7 macrophages at 16h post-infection.

G. Fatty Acid Oxidation (FAO) Assay of uninfected (UI) and infected (STM) RAW264.7

macrophages under SIRT1 or SIRT3 knockdown or inhibitor treatment. N=2, n=2. (**

p<0.01, * p<0.05)

H. Fatty Acid Oxidation (FAO) Assay of uninfected (UI) and infected RAW264.7

macrophages under infection with wildtype *S*. Typhimurium (STM WT), SPI-1 (Δ*invC*)or

SPI-2 (Δ*ssaV* and Δ*steE*) *m*utants of *S*. Typhimurium. Data is representative of N=2,n=2. (**

p<0.01, * p<0.05)

Fig.6- *Salmonella* Typhimurium infection proceeds with increased glycolysis and

glucose uptake inside the infected RAW 264.7 macrophages

A. *Salmonella* gene expression profiling data of *S*. Typhimurium infected RAW 264.7

macrophages at 2hr, 6hr and 16hr time points of infection through nanoString. SI- *S*.

Typhimurium infected, PI- PFA fixed *S*. Typhimurium infected. SPI-1 genes- *Inv, PrgH* ,

SPI-2 genes- *ssaV,stfF, glk*- glucokinase, *pfkA*-phosphofrucktose kinase A, *ptsG*-

Phosphophenolpyruvate-dependent sugar phosphotransferase system (PTS). Data is

representative of N=2, n=2.

B. qRT PCR gene expression profiling of *Salmonella* metabolic genes within infected

RAW264.7 macrophages in knockdown condition of either SIRT1 or SIRT3 at 6hr post

infection. Data is representative of N=3, n=3.

C-D. qRT PCR gene expression profiling of *Salmonella* metabolic genes within infected

female C57BL/6 mice spleen (C) and liver (D) under SIRT1 or SIRT3 inhibitor treatment

harvested at 5^{th} day post infection of *S*.Typhimurium (10^7 cfu units/animal). Unpaired two-

tailed Student's t test was performed to obtain the p values. Data is representative of N=3,

n=3. (****p<0.0001, *** p < 0.001, ** p<0.01, * p<0.05)

Fig.7- SIRT1 inhibition triggers hyperacetylation of glycolytic master regulator HIF-1α

within *S*. Typhimurium infected macrophages while SIRT3 skews metabolism of *S*.

Typhimurium-infected macrophages via interaction with PDHA1.

A. An immunoblot depicting HIF-1α interaction with SIRT1 post immunoprecipitation of

SIRT1 in uninfected (UI) or *S*. Typhimurium (STM) infected RAW264.7 macrophages at

16hr post-infection. (Derived from same SIRT1 IP blot as in Fig. 4A)

B. Immunoblotting of SIRT1 in SIRT1 knockdown uninfected (UI) or *S*. Typhimurium

(STM) infected RAW 264.7 cells at 16hr post-infection to assess the acetylation status of

HIF-1α.

C . Densitometric plot depicting the band intensities of Acetylated HIF-1α over total HIF-1α

in blot B.

D. An immunoblot depicting HIF-1α interaction with SIRT1 as well as the HIF-1α

acetylation status post immunoprecipitation of HIF-1^{α} (IP:HIF-1α or with control IgG

(IP:IgG) in uninfected (UI) or *S.* Typhimurium (STM) infected RAW264.7 macrophages

upon SIRT1 inhibitor (EX-527, 1μM) treatment at 16hr post-infection. UT-untreated.

E. Densitometric plot depicting the band intensities of Acetylated HIF-1α over total HIF-1α

in blot D.

F. Densitometric plot depicting the band intensities of SIRT1 in blot D.

G. Lactate estimation assay of *S.*Typhimurium infected RAW264.7 macrophages upon

SIRT1 and SIRT3 knockdown condition in the presence of HIF-1α inhibitor, chetomine

(50nM) at16h post-infection. Data is representative of N=3 , n=3. Unpaired two-tailed

Student's t test was performed to obtain the p values. (****p<0.0001, *** p < 0.001, **

p<0.01, * p<0.05)

H. An immunoblot depicting PDHA1 interaction with SIRT1 or SIRT3 post

immunoprecipitation of PDHA1 in uninfected (UI) or *S.* Typhimurium (STM) infected

RAW264.7 macrophages in SIRT3 knockdown condition at 16hr post-infection.

I. Densitometric plot depicting the band intensities of SIRT3 interaction over total input in

blot

J. An immunoblot depicting PDHA1 interaction with SIRT3 post immunoprecipitation of

PDHA1 in uninfected (UI) or *S.* Typhimurium (STM) infected RAW264.7 macrophages at

16hr post-infection under SIRT3 inhibitor (3-TYP, 1μM) treatment at 16hr post-infection.

UT-untreated.

K. Densitometric plot depicting the band intensities of Acetylated PDHA1 over total PDHA1

in blot I.

L. Densitometric plot depicting the band intensities of SIRT3 interaction over total input in

blot I.

Fig.8- Effect of SIRT1 or SIRT3 inhibition on *S.* Typhimurium infected C57BL/6 mice

A**.** The schematic representation of the experimental strategy for studying the effect of SIRT1

and SIRT3 on the *in vivo* pathogenesis of STM (WT).

B. *In-vivo* organ burden of *S.* Typhimurium upon SIRT1 or SIRT3 inhibition in C57BL/6

mice at day 5 post-infection under specified dosage of inhibitor treatment. Data is

representative of N=4, n=8. Mann-Whitney Test was performed to obtain the p

values.(****$p<0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$).

C. Percent survival of *S.* Typhimurium infected C57BL/6 mice upon SIRT1 or SIRT3

inhibitor treatment at a specific dose of inhibitor treatment. Data is representative of N=4,

n=5.

D. Representation of splenic length of *S.* Typhimurium infected spleen tissue harvested from

C57BL/6 mice (males) at day 5 post-infection upon SIRT1 or SIRT3 inhibition at 1mg/kg

dosage. Data is representative of N=4, n=8. (** $p<0.01$, * $p<0.05$).

E. Bacterial load in blood at different days post-infection upon SIRT1 or SIRT3 inhibition at

1mg/kg dosage in C57BL/6 mice (males). Data is representative of N=3, n>5. (** $p<0.01$, *

$p<0.05$) Mann-Whitney Test was performed to obtain the p values. (** $p<0.01$, * $p<0.05$).

F. Bacterial load in blood at day 5 post-infection upon SIRT1 or SIRT3 inhibition at 1mg/kg

dosage in C57BL/6 WT mice (males) and *gp91phox*-/- (males) mice. Data is representative of

N=3, n>5. Mann-Whitney Test was performed to obtain the p values. (** $p<0.01$, * $p<0.05$).

G. Serum IL-6 levels of *S.* Typhimurium infected C57BL/6 WT mice (males) mice treated

with SIRT1(EX-527) or SIRT3 (3TYP) inhibitors or SRT1720 (SIRT1 activator) at 1mg/kg

dosage at 5th day post-infection. Data is representative of N=3, n>5. Unpaired two-tailed

Student's t test was performed to obtain the p values.(** $p<0.01$, * $p<0.05$).

H. Quantitative analysis of percentage population of cells within liver showing DCFDA

positive staining shown in Fig. S9, A. Data is representative of N=3, n>5. Unpaired two-

tailed Student's t- test was performed to obtain the p values. (** $p < 0.01$).

I. Enumeration of GFP positive bacterial cells through flow cytometry in splenic tissues

homogenate harvested from adult male 6-8 week old C57BL/6 mice (subjected to different

chemical treatment-Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1

activator SRT1720 treated at a dose of 1mg/kg) at 5^{th} day post *S.* Typhimurium (expressing

GFP) infection (10^{7} CFU orally). Data is representative of N=3, n>5. Unpaired two-tailed

Student's t test was performed to obtain the p values. (** $p < 0.01$).

J. Quantitative analysis of the percentage population of splenic cells harboring GFP+

bacterial cells shown in I. Unpaired two-tailed Student's t test was performed. (** $p<0.01$, *

$p<0.05$).

K. Enumeration of GFP positive bacterial cells through flow cytometry within F4/80 positive

splenic macrophages present within splenic tissues homogenate harvested from adult male 6-

8 week old C57BL/6 mice (subjected to different chemical treatment-Vehicle treated or

SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1 activator SRT1720 treated aa t dose

of 1mg/kg) at 5^{th} day post *S.* Typhimurium infection. Data is representative of N=3, n>5.

L. Quantitative analysis of percentage population of F4/80 positive macrophage cells

harboring GFP+ bacteria shown in K . Unpaired two-tailed Student's t test was performed to

obtain the p values. Data is representative of N=3, n>5.(* $p < 0.05$).

M-S. Quantitative analysis of different mCherry-*S.*Typhimurium-infected splenic populations

harboring mCherry+bacteria via flow cytometry depicted in Fig. S8. Unpaired two-tailed

Student's t test was performed. (*** $p < 0.001$,** $p<0.01$, * $p<0.05$).

T. Representative image of haematoxylin-eosin-stained liver sections to assess the liver tissue architecture upon *Salmonella* infection at 5[th] days post-infection in different mice cohorts. (UI-Uninfected, STM- *S*. Typhimurium infected, EX-527-SIRT1 inhibitor, 3TYP-SIRT3 inhibitor, SRT1720- SIRT1 activator, Vehicle Control-(PBS containing 0.1% DMSO). Scale bar-50µm. **Scoring system:-** according to pathological changes the tissue sections are scored as 0 for normal pathology, 1 for mild/ minor pathology, 2 for moderate pathology, and 3 for severe pathological changes.

U. Graph representing the histopathological scoring of the liver sections depicted in L. Unpaired two-tailed Student's t test was performed to obtain the p values. (* $p < 0.05$)(****$p<0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$).

V. *In-vivo* organ burden of *S*. Typhimurium upon SIRT1 or SIRT3 adenovirus-mediated *in vivo* knockdown in C57BL/6 mice at day 5 post-infection. Data is representative of N=3, n>3. Mann-Whitney Test was performed to obtain the p values. (****$p<0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$).

W. Serum IL-6 levels of *S*. Typhimurium infected C57BL/6 WT mice (males) upon *in vivo* adenovirus-mediated SIRT1 or SIRT3 knockdown. Data is representative of N=3, n>3. Unpaired two-tailed Student's t test was performed to obtain the p values. (** $p<0.01$, * $p<0.05$).

X. Percent survival of *S*. Typhimurium infected C57BL/6 mice upon *in vivo* adenovirus-mediated SIRT1 or SIRT3 knockdown. Data is representative of N=3, n>3.

Y. Representative image of haematoxylin-eosin-stained liver sections upon *Salmonella* infection at 5[th] days post-infection in different mice cohorts. (Scrambled STM, shSIRT1 STM, shSIRT3 STM). Scale bar-50µm. **Scoring system:-** according to pathological changes

the tissue sections are scored as 0 for normal pathology, 1 for mild/ minor pathology, 2 for

moderate pathology, and 3 for severe pathological changes.

SUPPLEMENTARY FIGURE LEGENDS

Fig.S1- *Salmonella* **modulates the expression of SIRT1 and SIRT3 along its course of**

infection

A. Expression studies of SIRT1 and SIRT3 through immunoblotting in RAW 264.7

macrophages.

B. Representative immunoblot showing SIRT1 and SIRT3 expression at 16hr post-infection

with wildtype *S.* Typhimurium, SPI-1 mutant- Δ*invC* and SPI-2 mutants-Δ*ssaV*, Δ*steE*.

C. *SIRT1* and *SIRT3* transcript level expression in M1 polarized RAW264.7 macrophages at

16h post-infection. Data is representative of N=3, n=3. . Un-paired t-test were performed to

obtain p-values. (****$p<0.0001$,***$p < 0.001$, ** $p<0.01$, *$p<0.05$)

D. Expression profile of *SIRT1* and *SIRT3* in M2 polarized RAW264.7 macrophages at 16h

post-infection. Data is representative of N=3, n=3. Un-paired t-test were performed to obtain

p-values. (****$p<0.0001$,***$p < 0.001$, ** $p<0.01$, *$p<0.05$)

E. Intracellular fold proliferation of *S.* Typhimurium within non-polarized, M1 and M2

polarized RAW264.7 macrophages. Data is representative of N=3, n=3. Un-paired t-test were

performed to obtain p-values. (****$p<0.0001$,***$p < 0.001$, ** $p<0.01$, *$p<0.05$)

F. Validation of M1 and M2 polarized status of RAW264.7 macrophages by estimation of IL-

6 and IL-10 production via ELISA 24h post-polarization. Data is representative of N=3, n=3.

Un-paired t-test were performed to obtain p-values. (****$p<0.0001$,***$p < 0.001$, ** $p<0.01$,

*$p<0.05$)

Fig.S2- **Validation of SIRT1 and SIRT3 knockdown in RAW 264.7 macrophages**

A. Knockdown confirmation of *SIRT1* at transcript level through qPCR.

B. qPCR expression of *SIRT3* in order to validate knockdown efficiency.

C. Immunoblotting of SIRT1 for verification of knockdown. Scr=Scrambled, Mix=transfection with equimolar concentration of both SIRT1 shRNA construct C4 and C12.

D. SIRT3 knockdown validation through western blotting. Scr=Scrambled, Mix=transfection with equimolar concentration of both SIRT3 shRNA construct E8 and E12

Fig. S3- **SIRT1 and SIRT3 modulates cytokine production in *Salmonella* infection scenario**

A. Estimation of pro-inflammatory IL-6 cytokine production upon *S.* Typhimurium infection in SIRT1 and SIRT3 knockdown RAW 264.7 macrophages through ELISA. Data is representative of N=2, n=2. Unpaired two tailed Student's t test was performed to obtain the p values. (** $p<0.01$, * $p<0.05$).

B. Estimation of anti-inflammatory IL-10 cytokine production upon *S.* Typhimurium infection in SIRT1 and SIRT3 knockdown RAW 264.7 macrophages through ELISA. Data is representative of N=2, n=2. Unpaired two tailed Student's t test was performed to obtain the p values. (*** $p < 0.001$, * $p<0.05$)

C. Estimation of IL-1β cytokine production upon *S.* Typhimurium infection in SIRT1 and SIRT3 knockdown RAW 264.7 macrophages through ELISA at 20hr post-infection. Data is representative of N=2, n=2. Unpaired two-tailed Student's t test was performed to obtain the p values. (*** $p < 0.001$, * $p<0.05$) Uninfected (UI) samples values were below the detection level (ND-Not detectable).

D. Estimation of pro-inflammatory IL-6 cytokine production within uninfected or *S.* Typhimurium infected peritoneal macrophages in presence or absence of SIRT1 or SIRT3

inhibitor treatment at 6hr post-infection. Data is representative of N=2, n=2. Unpaired two

tailed Student's t test was performed to obtain the p values. (** $p<0.01$, * $p<0.05$)

E. Estimation of pro-inflammatory IL-1β cytokine production within uninfected or *S.*

Typhimurium infected peritoneal macrophages in presence or absence of SIRT1 or SIRT3

inhibitor treatment at 6hr post-infection. Data is representative of N=2, n=2. Unpaired two

tailed Student's t test was performed to obtain the p values. (** $p<0.01$, * $p<0.05$).

Uninfected (UI) samples values were below the detection level (ND-Not detectable).

F. Serum IL-1β levels of *S.* Typhimurium infected C57BL/6 WT mice (males) upon SIRT1

or SIRT3 inhibitor treatment at 5[th] day post-infection. Data is representative of N=3, n>3.

Unpaired two-tailed Student's t test was performed to obtain the p values. (** $p<0.01$, *

$p<0.05$).

G. Intestinal (ileal) IL-1β levels of *S.* Typhimurium infected C57BL/6 WT mice (males)

upon SIRT1 or SIRT3 inhibitor treatment at 5[th] day post-infection. Data is representative of

N=3, n>3. Unpaired two-tailed Student's t test was performed to obtain the p values. (**

$p<0.01$, * $p<0.05$).

Fig.S4- *Salmonella* Typhimurium skews the polarization state of the macrophage

toward an immunosuppressive M2 state along the course of infection

A- Flow cytometric analysis of alteration in M1 CD86 (PE) positive population in *S.*

Typhimurium (STM) infected samples in comparison to uninfected (UI) and

Paraformaldehyde Fixed (PFA) bacteria at the indicated time post-infection.

B- Flow cytometric analysis of M2 surface marker CD206 (APC) in *S.* Typhimurium (STM)

infected SIRT1 or SIRT3 knockdown RAW264.7 macrophages in comparison to the

scrambled control at the indicated time post-infection. UI-Uninfected sample.

Fig.S5- **SIRT1 and SIRT3 play a role in maintaining ROS balance within** *Salmonella*

infected macrophages.

A. DCFDA fluorescence measured through Flow cytometry in shSIRT1, shSIRT3 or

scrambled shRNA transfected and un-transfected RAW 264.7 macrophages at 16hr post *S.*

Typhimurium infection. Data is representative of N=3, n=2.

B. Quantitative analysis of percentage population of DCFDA positive cells shown in Fig.6,

A. Unpaired two-tailed Student's t test was performed. Data is statistically non-significant.

(** $p<0.01$, * $p<0.05$)

C. Detection of extracellular H_2O_2 production through Phenol Red Hydrogen peroxidase

assay at 610nm absorbance. Two-way ANOVA and Bonferroni post-t-test was used to obtain

p values. (** $p < 0.01$, * $p < 0.05$, ns-not significant) Data is representative of N=2, n=2. (**

$p<0.01$, * $p<0.05$)

Fig.S6- **Effect of SIRT1 and SIRT3 knockdown or inhibition in intracellular bacterial**

proliferation within RAW 264.7 macrophages or peritoneal macrophages in presence of

ROS scavenger N-Acetyl Cysteine (NAC).

A.Fold Proliferation of *Salmonella* Typhimurium within RAW 264.7 macrophages in

transfected and un-transfected conditions in presence or absence of NAC (1mM). Data are

representative of N=3, n=3. Unpaired two tailed Student's t test was performed to obtain the

p values. (****$p<0.0001$,*** $p < 0.001$, ** $p<0.01$, * $p<0.05$)

B.Fold Proliferation of *Salmonella* Typhimurium within peritoneal macrophages isolated

from 6-8 week old adult male C57BL/6 mice upon SIRT1 (EX-527, 1μM) or SIRT3 (3TYP,

1μM) inhibitor treatment in presence or absence of NAC (1mM). Data are representative of

N=3, n=3. Unpaired two tailed Student's t test was performed to obtain the p values.

(****$p<0.0001$,*** $p < 0.001$, ** $p<0.01$, * $p<0.05$)

Fig.S7 SIRT1 and SIRT3 knockdown or inhibitor treatment led to decrease in fatty acid oxidation and promoted host glycolysis in *S.* Typhimurium infected peritoneal macrophages

A-qRT PCR mediated gene expression analysis of several host metabolic genes in SIRT1 or SIRT3 knockdown RAW264.7 macrophages at 6 hr post-infection. Data is representative of N=3,n=3. Unpaired two-tailed Student's t test was performed to obtain the p values. (* $p<0.05$)

B- qRT PCR mediated gene expression analysis of several host metabolic genes in SIRT1 or SIRT3 inhibitor-treated (1μM) peritoneal macrophages isolated from 6-8 week old C57BL/6 male mice liver harvested at 5th day post thioglycolate injection. Data is representative of N=3,n=3. Unpaired two-tailed Student's t test was performed to obtain the p values. (****$p<0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$)

C-Lactate estimation assay of *S.*Typhimurium infected peritoneal macrophages upon SIRT1 or SIRT3 catalytic inhibition at 2h, 6h and 16h post-infection. Data is representative of N=3,n=3. Unpaired two-tailed Student's t test was performed to obtain the p values. (****$p<0.0001$, *** $p < 0.001$, ** $p<0.01$, * $p<0.05$)

D-Immunoblotting of host glycolytic (PGK, PFK) and fatty acid oxidation (HADHA, ACOX-1) proteins under SIRT1 (EX-527) and SIRT3 (3TYP) catalytic inhibitor treated condition in *S.* Typhimurium infected peritoneal macrophages (isolated from thioglycolate treated adult male C57BL/6 mice) at 16h post-infection. Data is representative of N=2, n=1.

E- qRT PCR mediated gene expression profiling of several host metabolic genes in SIRT1 or SIRT3 inhibitor treated 6-8 week old C57BL/6 male mice liver harvested at 5-day post infection of *S.*Typhimurium (10^7 CFU units/animal). Unpaired two-tailed Student's t test was

performed to obtain the p values. Data is representative of N=3, n=3. (****p<0.0001, *** p <

0.001, ** p<0.01, * p<0.05)

F-qRT PCR mediated gene expression profiling of several host metabolic genes in SIRT1 or

SIRT3 inhibitor treated 6-8 week old C57BL/6 male mice spleen harvested at 5 day post-

infection of S.Typhimurium (10^7 CFU units/animal). Unpaired two-tailed Student's t test was

performed to obtain the p values. Data is representative of N=3, n=3. (****p<0.0001, *** p <

0.001, ** p<0.01, * p<0.05)

Fig. S8-Flow cytometric analysis of different splenic populations within mCherry-

labelled S. Typhimurium infected mice cohorts upon SIRT1 or SIRT3 inhibitor

treatment

A-Enumeration of mCherry positive bacterial cells through flow cytometry in splenic tissues

homogenate harvested from adult male 6–8-week-old C57BL/6 mice (subjected to different

chemical treatment-Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1

activator SRT1720 treated at a dose of 1mg/kg) at 5^{th} day post-S. Typhimurium infection (10

^{7} CFU orally).

B-Flow cytometric analysis of eFluor450 conjugated CD11c+ population within single cell

splenic suspension isolated from adult male 6–8-week-old C57BL/6 mice (subjected to

different chemical treatment-Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor

or SIRT1 activator SRT1720 treated at a dose of 1mg/kg) at 5^{th} day post-mCherry-labelled S.

Typhimurium infection (10^7 CFU orally).

C-Flow cytometric analysis of FITC conjugated Ly6C+ population within single cell splenic

suspension isolated from adult male 6–8-week-old C57BL/6 mice (subjected to different

chemical treatment-Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1

activator SRT1720 treated at a dose of 1mg/kg) at 5^{th} day post-mCherry-labelled *S*.

Typhimurium infection (10^7 CFU orally).

D- Flow cytometric analysis of Alexa 647-conjugated CD45+ population within single cell

splenic suspension isolated from adult male 6–8-week-old C57BL/6 mice (subjected to

different chemical treatment-Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor

or SIRT1 activator SRT1720 treated at a dose of 1mg/kg) at 5^{th} day post-mCherry-labelled *S*.

Typhimurium infection (10^7 CFU orally).

E-G-Alexa 647-conjugated CD45+ population (as shown in D) was gated as per granularity

(Side Scatter versus Alexa 647 plot) into P6 population with high granularity (granulocytes),

P5 (monocytes), and P4 population with low granularity (lymphocytes). Flow cytometric

analysis of Alexa 647-conjugated CD45+ (E) P6 (granulocyte), (F) P5 (monocyte), and (G)

P4 (lymphocyte) population within single cell splenic suspension isolated from adult male 6–

8-week-old C57BL/6 mice (subjected to different chemical treatment-Vehicle treated or

SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1 activator SRT1720 treated at a dose

of 1mg/kg) at 5^{th} day post-mCherry-labelled *S*. Typhimurium infection (10^7 CFU orally).

Fig.S9- SIRT1 and SIRT3 inhibition led to enhanced ROS production within

***Salmonella*-infected mice liver and spleen tissues.**

A-DCFDA fluorescence measured through flow cytometry in liver tissues homogenate

harvested from adult male 6-8 week old C57BL/6 mice (subjected to different chemical

treatment- Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1 activator

SRT1720 treated at dose of 1mg/kg) at 5^{th} day post *S*. Typhimurium infection (10^8 CFU

orally). Data is representative of N=3, n=2.

B- DCFDA fluorescence measured through Flow cytometry in splenic tissues homogenate

harvested from adult male 6-8 week old C57BL/6 mice (subjected to different chemical

treatment-Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1 activator

SRT1720 treated at dose of 1mg/kg) at 5^{th} day post *S.* Typhimurium infection. Data is

representative of N=3, n=2.

C- Quantitative analysis of percentage population of splenocytes showing DCFDA positive

staining shown in splenic tissues homogenate harvested from adult male 6-8 week old

C57BL/6 mice as shown in B. Unpaired two-tailed Student's t test was performed to obtain

the p values. (** $p < 0.01$)

Fig. S10 *In vivo* knockdown validation by Western blotting and qPCR in liver

A-Immunoblotting of SIRT1 and SIRT3 in harvested liver samples upon AAV- mediated *in*

vivo knockdown in mice.

B- qPCR mediated gene expression of *SIRT1* in harvested liver samples upon *in vivo* AAV-

mediated gene inhibition in mice. Data is representative of N=3,n=3. Unpaired two-tailed

Student's t test was performed to obtain the p values. (**** $p<0.0001$,*** $p < 0.001$, **

$p<0.01$, * $p<0.05$)

C-qPCR mediated gene expression of *SIRT3* in harvested liver samples upon *in vivo* AAV-

mediated gene inhibition in mice. Data is representative of N=3,n=3.Unpaired two-tailed

Student's t test was performed to obtain the p values. (**** $p<0.0001$,*** $p < 0.001$, **

$p<0.01$, * $p<0.05$)

Fig. S11-Schematic Diagram-In murine macrophages, *Salmonella* infection drives an

immuno-metabolic shift toward an immunosuppressive state with an increase in host fatty

acid oxidation rate by modulating SIRT1 and SIRT3. SIRT1 and SIRT3 regulate the

acetylation status of NFκB p65 subunit, HIF-1α, PDHA1 thereby influencing host

metabolism which in turn impacts the intracellular *Salmonella* metabolism. Inhibition or

knockdown of SIRT1 or SIRT3 triggers increased ROS, pro-inflammatory cytokine burst

with a concomitant decline in host fatty acid oxidation and a corresponding increase in

bacterial fatty acid oxidation together amounting to decreased intracellular bacterial

replication. Contrastingly, in *in vivo* mice model of infection, SIRT1 and SIRT3 cause

increased bacterial dissemination owing to increased ROS, IL-6 and IL-1β production.

1413

1414

1415

MANUSCRIPT_1 TITLE:
Salmonella-induced SIRT1 and SIRT3 are crucial for maintaining the metabolic switch in bacteria and host for successful pathogenesis

MAIN FIGURES

Fig.1

Fig.2

Fig.3

A

B

C

Fig.4

Fig.5

Fig.6

A

Normalized Expression of Bacterial Genes (PI and SI Samples; Average expression for Replicates)

B

Fig.7

Fig.8

I

Vehicle Control

0.07

EX-527(1mg/kg)

0.76

3TYP (1mg/kg)

3.11

SRT (1mg/kg)

0.08

J

T

STM Vehicle Control

Score:1

4X   10X   40X

Slide- 1

Showing minimal/ mild focal inflammatory cell collections in lobular parenchyma

STM SRT1720 (1mg/kg)

Score-1

Slide- 9

Showing minimal/ mild pathology in lobular parenchyma with normal hepatic architecture

STM EX-527(1mg/kg)

Score : 3

C   V
N
INF
N   INF

SLIDE- 4

Showing multiple (small and big areas) severe acute hepatic necrosis (N), with complete loss of hepatic architecture due to loss of hepatic cells, the necrotic area filled with edema and surrounded by inflammatory cells, hepatocytes with cytoplasmic vacuolation (V), inflammatory cells (INF), and damage of the endothelial lining of central vein (C)

STM 3TYP (1mg/kg)

Score-3

INF   N
INF
HPV   HPV
H   V   HPV   HPV
N
HPV   N

Slide- 7

Showing small multiple necrotic foci (N), severe blood hemorrhage (H) congested hepatic portal veins of liver clogged with clumps and inflammatory cell infiltrates (HPV) vacuolar degeneration (V). Multiple Microgranuloma formation with the aggregation of inflammatory cells (INF)

Score-0 UI Vehicle Control

normal pathology with normal hepatic architecture

Score-1 UI EX-527(1mg/kg)

Showing minimal/ mild focal inflammatory cell collections in lobular parenchyma

Score-1 UI 3TYP (1mg/kg)

Showing minimal/ mild focal inflammatory cell collections in lobular parenchyma

Score-0 UI SRT1720 (1mg/kg)

normal pathology with normal hepatic architecture

V

MLN
Liver
Spleen
Blood 5dpi

****

***

y-axis: log (CFU/ml/g wt)

x-axis: Scrambled, shSIRT1, shSIRT3

**

*

y-axis: log (CFU/ml/g wt)

x-axis: Scrambled, shSIRT1, shSIRT3

**

*

y-axis: log (CFU/ml/g wt)

x-axis: Scrambled, shSIRT1, shSIRT3

*

0.0585

y-axis: log (CFU/ml)

x-axis: Scrambled STM, shSIRT1 STM, shSIRT3 STM

W

**

y-axis: IL-6 (pg/ml)

x-axis: UI, STM SCRAMBLED, STM shSIRT1 STM, STM shSIRT3

X

y-axis: Percent Survival

x-axis: Days

- Scrambled STM
- shSIRT1 STM
- shSIRT3 STM

Y

Scrambled STM
Score:0/1

4X   10X   40X

Showing normal hepatic architecture with some area immune cell accumulation

shSIRT1 STM
Score:3

4X   10X   40X

Showing multiple necrotic areas (N), congestion and damage to the endothelial lining of the central vein (C), congestion of the hepatic portal vein (HPV), and severe infiltration of inflammatory immune cells(IC),

shSIRT3 STM
Score:3

4X   10X   40X

Showing multiple necrotic areas (N), congestion of the hepatic portal vein (HPV) and severe infiltration of inflammatory immune cells(IC),