1   **SIRT1 and SIRT3 mediated immune-metabolic switch is crucial for the establishment of**

2   *Salmonella* **pathogenicity of infection.**

3   **Dipasree Hajra**[1]**, Raju S. Rajmani**[3] **and Dipshikha Chakravortty**[1,2]

4   [1]Department of Microbiology & Cell Biology, Indian Institute of Science

5   [2]Adjunct Faculty, School of Biology, Indian Institute of Science Education and Research,
6   Thiruvananthapuram

7   [3]Centre of Infectious Disease Research, Indian Institute of Science, Bangalore, India

8   Corresponding author: Prof. Dipshikha Chakravortty

9   Email: dipa@iisc.ac.in

10   Tel. No. 0091 80 22932842

11

12   **ABSTRACT**

13   *Salmonella* continues to be a threat to the human population by taking a toll on the lives of

14   about 20,000 individuals globally per year. Host sirtuins or NAD+-dependent deacetylases are

15   the major players in host immuno-metabolic regulation. However, the role of sirtuins in the

16   modulation of the immune metabolism pertaining to Salmonellosis is largely unknown. Here,

17   we investigated the role of SIRT1 and SIRT3 in mediating immuno-metabolic switch along the

18   progression of *Salmonella* infection. Gene profiling (Nanostring) data along with flow

19   cytometric analysis indicated the ability of the wildtype *Salmonella* Typhimurium to skew the

20   polarization state of the macrophages from a pro-inflammatory M1 state toward an

21   immunosuppressive M2 state along its course of infection. Further, using shRNA-mediated

22   knockdown studies and catalytic inhibitor treatment, we showed that SIRT1 and SIRT3 play a

23   crucial role in mediating the immuno-metabolic switch that governs macrophage polarization

24   in *Salmonella*-infected macrophages. SIRT1 or SIRT3 knockdown or inhibition led to

25   decreased expression of M2 surface markers such as CD206 and showed increased production

of pro-inflammatory cytokines and ROS generation which together resulted in attenuated

bacterial intracellular proliferation within the infected RAW264.7 macrophages. Alongside

immunological functions by modulating p65 NF-κB acetylation, SIRT1 and SIRT3 knockdown

or inhibition also skew *Salmonella*-induced host metabolic shift towards increased fatty acid

oxidation by regulating acetylation status of HIF-1α and PDHA1 which in turn impacts the

metabolism of *Salmonella* within the infected macrophages. *In vivo* mice-model infection

studies highlighted the role of SIRT1 and SIRT3 in controlling bacterial burden and

dissemination.

Keywords: SIRT1 and SIRT3, immune regulation, metabolic shift, macrophage

polarization, *Salmonella* infection, bacterial dissemination, deacetylation

INTRODUCTION

Sirtuins are NAD+-dependent deacetylases that are present in all forms of life. *Saccharomyces*

cerevisiae serves as the founder of the discovery of Sir2. Sirtuins comprise a conserved core

catalytic domain that removes acetyl moiety from the lysine residues of proteins in presence of

NAD+ as a cofactor [1] giving rise to 2'O-acetyl-ADP-ribose and free nicotinamide as

products[2][3]. Free nicotinamide acts as a non-competitive inhibitor of sirtuins[4]. They

possess variable N terminal and C terminal domains that confer different subcellular

localization, substrate specificity, and functions[5]. Mammals have seven sirtuins that act on

different substrates like histones and transcriptional factors like p53, FOXO, PGC1α, UCP2,

liver X-receptor, and others. Thus, they are responsible for regulating various biological

functions   such   as   cell   survival,   apoptosis,   oxidative   stress,   metabolism,   and

inflammation[6][7]. SIRT1,6 and 7 have nuclear localization, SIRT2 is cytoplasmic in nature

and SIRT 3,4 and 5 comprise the mitochondrial SIRTs. SIRT1 is the most characterized of all

the SIRTs and possesses strong homology with the yeast Sir2 protein. In addition to their

deacetylase activity, they possess ADP ribosylation (SIRT1, SIRT4, and SIRT6),

desuccinylation and demalonylation (SIRT5), delipoylation (SIRT4), and demyristoylation and

depalmitoylation (SIRT6) enzymatic activities [8]. Previous studies have shown that SIRT1-

mediated alteration in gene expression interlinks metabolism and inflammation. SIRT1 gets

activated in response to acute immune response and helps in its resolution. It deacetylates

RelA/p65 component of NFκB thereby mediating its proteasomal degradation[9]. On the other

hand, it activates RelB component of NFκB pathway. RelB causes heterochromatinization of

pro-inflammatory genes like TNF-α and IL-β[10] alongside, it causes activation of the fatty

acid oxidation genes. With the gradual progression toward the adaptive phase of the immune

response, SIRT1 activates Peroxisome proliferator-activated receptor γ (PPARγ) coactivator-

1α (PGC-1α) mediating a metabolic switch from glycolysis toward fatty acid oxidation. SIRT1-

mediated RelB activation, in turn, activates SIRT3 causing the promotion of mitochondrial

bioenergetics[11]. PGC-1α, a major player in mitochondrial biogenesis, activates SIRT3 [12]

which in turn causes activation of PGC-1α, thereby fuelling a positive feedback loop. SIRT3

accounts for the major mitochondrial deacetylase orchestrating several metabolic processes

such as fatty acid oxidation, promotion of the TCA cycle, and inhibition of ROS

production[13].

Salmonella enterica serovar Typhimurium is a facultative intracellular Gram-negative enteric

pathogen, causing a wide array of infections ranging from self-limiting gastroenteritis to

diarrhea in humans [14]. *Salmonella enterica* serovar Typhi cause systemic infection in humans

with typhoidal symptoms. Recent reports reported incidences of 21 million [15] typhoid cases

and 93 million of non-typhoidal [16] cases round the year. The virulence of *Salmonella* is

majorly regulated by two pathogenicity islands, namely, SPI-1 and SPI-2. It uses SPI-1

encoded T3SS and the effector proteins to invade host cells [17]. Inside the macrophages, they

harbour within the SCV by virtue of its SPI-2 effectors [18]. Macrophages, dendritic cells and

neutrophils are responsible for successful dissemination throughout the body through the

reticulo-endothelial system (RES) [17].

Macrophages, serving as an intracellular niche for *Salmonella,* exhibit several continua of

polarization states. At the two extreme ends of the spectrum lie the classically polarized M1

macrophages and alternatively activated M2 macrophages. M1 macrophages comprise of the

proinflammatory antimicrobial state producing IL-1β, IL-6, TNF-α,IL-12, IFN-γ cytokines and

exhibit enhanced expression of CD80, CD86 surface markers. The anti-inflammatory M2

macrophages promote bacterial persistence by producing anti-inflammatory cytokines like IL-

10,TGF-β and show increased expression of Arg-1, CD206 surface markers[19][20]. To sustain

the continuous production of proinflammatory cytokines M1 macrophages rely on glycolysis

for their energy requirements. On the other hand, M2 macrophages are fuelled by enhanced

oxidative phosphorylation (OXPHOS) and fatty acid oxidation (FAO)[21].It has been

previously reported that sirtuins mediated attuning of metabolism impact polarization of

macrophages *in vivo*. SIRT1 has the ability to promote the polarization of M2 macrophages

and inhibit inflammation in macrophages of adipose tissue[22][23][24]. SIRT3 suppresses

ROS by deacetylating and activating MnSOD[25].

Several bacteria are known to subvert the host immune system toward an immunosuppressive

state. Pathogens like *Salmonella* or *Mycobacterium* have evolved mechanisms to counteract

the M1 state of the host macrophage. *Salmonella* Typhimurium uses its SPI-2 effectors to

inhibit the recruitment of NADPH oxidase to the *Salmonella* Containing Vacuoles (SCV),

thereby preventing oxidative burst mediated microbicidal activity[26]. Similarly,

Mycobacterium bovis bacillus Calmette-Gue´rin prevents NOS2 recruitment to

phagosomes[27]. *Salmonella* Dublin causes inhibition of the production of pro-inflammatory

cytokine like IL-18 and IL-12p70[28]. Moreover, *Mycobacteria* inhibits NFκB signalling and

IFN-γ mediated downstream pathways[29]. Furthermore, *Yersinia enterocolitica* elicits an M2

response by inducing Arginase-1 expression and TGFβ1 and IL-4production[30]. *Yersinia*

TTSS effector LcrV induces an M2 phenotype supposedly by IL-10 production[31].

Since SIRT1 and SIRT3 are the major modulators of the immuno-metabolic paradigm

governing the macrophage polarization, we intend to decipher the role of SIRT1 and SIRT3 in

influencing immuno-metabolism in *Salmonella* infection scenario. We show that *Salmonella*

infection modulates SIRT1 and SIRT3 expression along with the progression of infection.

Furthermore, we highlight the role of SIRT1 and SIRT3 in intracellular pathogen survival, ROS

production, immuno-metabolic switch together amounting to the modulation of macrophage

polarization. Here, we have showed that SIRT1 and SIRT3 knockdown triggered decreased

M2 surface marker expression such as CD206 along with increased production of pro-

inflammatory cytokines and ROS generation, together amounting to attenuated bacterial

intracellular proliferation within the infected macrophages. Moreover, SIRT1 mediated p65

NF-κB deacetylation played a vital role in immune function regulation within the *Salmonella*

infected macrophages. Further, we report the ability of the *Salmonella* pathogen to drive the

metabolic state of the host from glycolysis toward increased fatty acid oxidation in RAW264.7

macrophages. SIRT1 and SIRT3 knockdown or inhibition skewed this *Salmonella*-induced

host metabolic shift by regulating acetylation status of HIF-1α and PDHA1 which in turn

impacted the metabolism of *Salmonella* within the infected macrophages. In the contrary, in *in*

vivo mice-model of infection, SIRT1 and SIRT3 inhibition resulted in increased pathogen loads

in organs and triggered enhanced bacterial dissemination, together leading to increased

susceptibility of the mice to *S*. Typhimurium infection.

MATERIALS AND METHODS

Bacterial Strains, and culture conditions

Salmonella enterica serovar Typhimurium (STM) strain ATCC 14028S or ATCC 14028S constitutively expressing green fluorescent protein (eGFP) or mCherry (RFP) through pFPV plasmid were used. 4% paraformaldehyde fixed STM (PFA) was used as the killed fixed bacteria control. The above-mentioned live bacterial strain was grown overnight in LB broth in 37 ^{0}C at 160 rpm shaking condition in presence or absence of appropriate antibiotic (Ampiciilin-50µg/ml) (Table-1) after revival of the bacterial strains from glycerol stock (stored at -80 ^{0}C).

Table:1: List of strains and plasmids used in this study

Strains/Plasmids	Characteristics	Source
Salmonella enterica serovar Typhimurium ATCC strain 14028S	Wild Type (WT)	Gifted by Prof. M. Hensel
pFPV:GFP	Amp^R	Laboratory Stock
pFPV:mCherry	Amp^R	Laboratory Stock
pLKO.2 : shRNA	Amp^R	Gifted by Prof. Subba Rao Gangi Shetty

Cell Culture

RAW 264.7 murine macrophages were cultured in DMEM (Lonza) containing 10% FBS (Gibco) at 37 ^{0}C in a humified incubator with 5% CO_2. Prior to each experiment, cells were seeded into 24 well or 6 well plate as per requirement at a confluency of 60%.

Peritoneal macrophages were collected in PBS from the peritoneal cavity of C57BL/6 mice aseptically post thioglycolate treatment using 20G needle and 5ml syringe. Following centrifugation, cell pellet was resuspended in RPMI-1640 (Lonza) containing 10% heat-inactivated FBS (Gibco), 100 U/ml penicillin and 100 µg/ml streptomycin and seeded into 6

144 well- plate. 6hr prior to infection, antibiotic containing media was replaced with Penicillin-

145 Streptomycin free RPMI-1640 (Lonza) containing 10% heat-inactivated FBS (Gibco).

Transfection

shRNA mediated knockdown was carried out by PEI mediated transfection protocol. Plasmid

harbouring shRNA in pLKO.2 vector backbone specific to SIRT1 and SIRT3 were used for

transfection. Plasmid harbouring scrambled sequence of shRNA, served as a control, was also

used for transfection. Plasmid DNA was used at a concentration of 500ng and 1µg per well of

a 24-well plate and 6-well plate respectively. Plasmid and PEI were added in 1:2 ratio in serum

free DMEM media and incubated for 20 mins at room temperature. Post incubation, the DNA:

PEI cocktail was added to the seeded RAW 264.7 macrophages. After 6-8hrs of incubation,

serum-free media was replaced with complete media containing 10% FBS. Post 48hr of

transfection, transfected cells were either harvested for knockdown confirmation studies or

subjected to infection with STM.

Table 2: List of shRNA used for knockdown

Sirtuins	Construct ID	TRC ID	Sequence (5'---3')
SIRT1	C4	TRCN0000218734	GTACCGGCATGAAGTGCCTCAGATAT TACTCGAGTAATATCTGAGGCACTTC ATGTTTTTTG
SIRT1	C12	TRCN0000018979	CCGGGCAAAGCCTTTCTGAATCTATC TCGAGATAGATTCAGAAAGGCTTTGC TTTTT

SIRT3	E8	TRCN0000038889	CCGGCCCAACGTCACTCACTACTTTCTCGAGAAAGTAGTGAGTGACGTTGGGTTTTTG
SIRT3	E12	TRCN0000038893	CCGGCCACCTGCACAGTCTGCCAAACTCGAGTTTGGCAGACTGTGCAGGTGGTTTTTG

Infection Protocol

Macrophages were infected with stationary-phase bacterial culture with MOI of 10. For

synchronization of the infection, tissue culture plates were subjected to centrifugation at 600xg

for 5 min and incubated at 37 ^{0}C humified incubator with 5% CO_2 for 25 min. Cells were

washed with PBS and were treated with DMEM (Sigma) + 10% FBS (Gibco) containing 100

μg/ml gentamicin for 1 hr. Subsequently, the gentamicin concentration was reduced to 25

μg/ml and maintained until the cells were harvested. For the inhibitor treatment studies, along

with 25 μg/ml containing complete media 1μM of SIRT1 (EX-527) inhibitor or SIRT3 (3TYP)

or 10mM of N-Acetyl Cysteine (NAC) were added to the cells.

Immunofluorescence confocal microscopic studies

At the specified time points post infection with GFP tagged STM, cells were fixed with 3.5%

paraformaldehyde for 15 min. Primary antibody staining was performed with specific primary

antibody in the presence of a permeabilizing agent, 0.01% saponin (Sigma) dissolved in 2.5%

BSA containing PBS at 4°C for overnight or for 6hr at room temperature (RT). Following this,

cells were washed with PBS stained with appropriate secondary antibody tagged with

fluorochrome for 1 hr at RT. This was followed by DAPI staining and mounting of the coverslip

onto a clean glass slide using the mounting media containing the anti-fade agent. The coverslip

sides were sealed with a transparent nail paint. All immunofluorescence images were obtained

using Zeiss LSM 710 or Zeiss LSM 880 and were analyzed using ZEN black 2012 software.

Quantitative Real Time PCR

Total RNA was isolated at specific time points post infection by using TRIzol (Takara) as per

manufacturer's protocol. Quantification of RNA was performed in Nano Drop (Thermo-

Fischer scientific). Quality of isolated RNA was detected by performing 2% agarose gel

electrophoresis. 2 µg of RNA was subjected to DNaseI (Thermo Fischer Scientific) treatment

at 37°C for 1 hr followed by addition of 0.5M EDTA (final concentration 5mM) and heat

inactivation at 65°C for 10 mins. The mRNA was reverse transcribed to cDNA using oligo

$(dT)_{18}$ primer, buffer, dNTPs and reverse transcriptase (Takara) as per manufacturer's protocol.

The expression profile of target gene was evaluated using specific primers (Table-3) by using

SYBR green RT-PCR master mix (Takara) in BioRad Real time PCR instrument. β-actin was

used as an internal control for mammalian genes and for bacterial genes 16S rRNA was used.

All the reaction was setup in 384 well plate with two replicates for each sample.

Table 3: List of primers

Primer Name	Sequence (5'---3')
SIRT1 FP	ACAAAGTTGACTGTGAAGCTGTAC
SIRT1 RP	GTTCATCAGCTGGGCACCTA
SIRT3 FP	GCTGCTTCTGCGGCTCTATAC
SIRT3 RP	GAAGGACCTTCGACAGACCGT
PPARδ FP	CACAACGCTATCCGCTTTGG
PPARδ RP	ATGCTCCGGGCCTTCTTTTT
βactin FP	CAGCAAGCAGGAGTACGATG

βactin RP	GCAGCTCAGTAACAGTCCG
Acadl FP	CTTGGGAAGAGCAAGCGTACT
Acadl RP	CTGTTCTTTTGTGCCGTAATTCG
Hadha FP	AGCAACACGAATATCACAGGAAG
Hadha RP	AGGCACACCCACCATTTTGG
Acox1 FP	TCGAAGCCAGCGTTACGAG
Acox1 RP	ATCTCCGTCTGGGCGTAGG
Pdha1 FP	TGTCGGTTCCCAGTCCATCA
Pdha1 RP	CGTTTCCTTTTCACAGCACATGA
Pfkl FP	GAACTACGCACACTTGACCAT
Pfkl RP	CTCCAAAACAAAGGTCCTCTGG
ptsG	TATCTGGGCTTCTTTGCGGG
ptsG	ACCAGGCAACGCTCGATAAA
fadA FP	TCTGGGATTGATGGAGCAGA
fadA RP	AGACCAATACACATCGTCGC
fadB FP	GTCCCCGAAGAGCAGTTAAG
fadB RP	CCAGTTTAGTTTCCGGCAGA
aceA FP	CGATCTGGTATGGTGCGAAA
aceA RP	TTCTGCCAGTTGAAGGATGG
aceB FP	AGCGTTTCAATCAACAGGGT
aceB RP	CCCGTATTTTTACCTGCCGA
16S rRNA FP	GTGAGGTAACGGCTCACCAA
16S rRNA RP	TAACCGCAACACCTTCCTCC

Intracellular proliferation or gentamicin protection assay

Following infection of the transfected cells with STM at an MOI of 10, cells were treated with DMEM (Sigma) + 10% FBS (Gibco) containing 100 μg/ml gentamicin for 1 hr. Subsequently, the gentamicin concentration was reduced to 25 μg/ml and maintained until the specified time point. Post 2hr and 16hr post-infection, cells were lysed in 0.1% triton-X-100. Lysed cells were serially diluted and plated on *Salmonella-Shigella* (SS) agar to obtain colony-forming units (cfu). Fold proliferation was calculated as cfu at 16hr divided by cfu at 2hr.

Fold Proliferation= [CFU at 16h]/[CFU at 2h]

Western Blotting

Post appropriate time points of infection, the cells were washed in PBS and subsequently harvested in PBS. The cell pellets were obtained after centrifugation at 300g for 7 minutes at 4ºC. Cells were lysed in 1X RIPA (10X-0.5M NaCl, 0.5M EDTA pH-8.0, 1M Tris, NP-40, 10% sodium deoxycholate, 10% SDS) buffer containing 10% protease inhibitor cocktail (Roche) for 30 min on ice. Total protein was estimated using Bradford (Bio-Rad) method of protein estimation. Protein samples were subjected to 12 % SDS polyacrylamide gel electrophoresis and then were transferred onto 0.45µm PVDF membrane (18V, 2 hrs). The membrane was blocked using 5% skim milk in TBST (Tris Buffered Saline containing 0.1% Tween-20) for 1hr at RT and subsequently probed with appropriate primary antibody (Table-4) for overnight at 4°C. Following wash in TBST, blot was probed with specific HRP conjugated secondary antibody for 1hr at RT. The membrane was developed using ECL (Advansta) and images were captured using ChemiDoc GE healthcare. All densitometric analysis was performed using ImageJ software.

216 **Table 4: List of Antibodies**

Antibody Name	Catalogue Number
SIRT1 Polyclonal Antibody	Cat #PA5-85921 (ThermoFisher Scientific)
SIRT3 Polyclonal Antibody	**Cat# PA5-13222(ThermoFisher Scientific)**
CD 86	**Cat#**12-086282PE(ThermoFisherScientific)
CD 206	**Catalog#** 17-2069-42 (ThermoFisherScientific)
F4/80 Antibody	BD Horizon, Cat-565411
HIF-1α	**sc-13515 SCBT**
NF-κβ p65	CST(D14E12)XP® RabbitmAb #8242
Acetylated-Lysine Antibody	**Cat #9441(Cell Signaling Technology)**
PDHA1	**Catalog No. A13687 (ABclonal)**
HADHA	**Catalog No. A5346 (ABclonal)**
ACOX1	**Catalog No. A8091 (ABclonal)**
PGK	**Catalog No. A12686 (ABclonal)**
PFK	**Catalog No. A5477 (ABclonal)**
β actin	A3854 (Sigma)

Immunoprecipitation

For co-immunoprecipitation, cells were washed with PBS and were lysed in native lysis buffer containing 1% Nonidet P-40, 20 mM Tris (pH 8), 2 mM EDTA, 150 mM NaCl and protease inhibitors mixture (Roche Diagnostics) for 30 min at 4°C. Cell debris was removed by centrifugation at 10,000 rpm for 10 min and the supernatant was treated with the specific antibody against the protein to be precipitated. Antibody-lysate complexes were immunoprecipitated using Protein A/G- linked magnetic beads (MagGenome) according to the manufacturer's protocol. Beads were extensively washed with washing buffer and denatured at 95 °C for 10 min. Denatured precipitates were subjected to SDS-PAGE (12% gel) followed by transfer to 0.45 μ PVDF membrane. The membrane was blocked using 5% skimmed milk in TBST (Tris Buffered Saline containing 0.1% Tween-20) for 1h at room temperature and eventually probed for the target primary antibodies or Anti-acetylated Lysine (Ac-K) primary antibody overnight at 4°C. The blot was probed with a specific HRP-conjugated secondary antibody for 1hr at RT after rigorous washing in TTBS. ECL (BioRad) was used for detection and images were captured using ChemiDoc GE healthcare.

ELISA

Estimation of cytokines in cell-free supernatant was performed according to the manufacturer's instructions. Briefly, 96-well ELISA plates (BD Bioscience) were coated overnight with capture antibody at 4°C. Following day, plates were washed with 0.1% Tween-20 containing PBS and blocked with 10% FBS for 1 h. Following blocking, wells were washed and incubated with 100 μL of test samples for 2 h at room temperature. Subsequently, plates were washed and incubated with detection antibody and enzyme reagent for 1 h at room temperature (BD Bioscience). TMB (Sigma) was used as a substrate and reactions were

243   stopped with 2 N H_2SO_4. Absorbance was measured at 450 nm wavelength in Tecan Plate

244   reader and the concentration of cytokines were interpolated from a standard curve.

Flow cytometry

After specific time points post infection, cells were washed and harvested in PBS. Following

centrifugation, cell pellet was resuspended in FACS buffer comprised of 1% BSA in PBS.

Blocking was performed with Fc blocker (purified Anti-mouse CD16/CD32, eBioscience)

dissolved in FACS blocking buffer for 30 min on ice. Following a washing step with PBS,

antibody staining was performed with PE-conjugated CD86 antibody or APC-conjugated

CD206 (Thermo Scientific) for 45 min on ice. After washing in PBS, the cell pellet was

resuspended in 1% PFA in PBS. Subsequently, PFA was removed, and cells were resuspended

in FACS buffer and reading was taken in BD FACSVerse instrument. For flow cytometry

studies, the harvested liver or spleen were homogenized into single cell suspension post RBC

lysis (RBC lysis Buffer, Sigma). The homogenized cell suspension was washed and

resuspended in FACS buffer containing the PE-conjugated Rat anti-mouse F4/80 antibody (BD

Horizon, Cat-565411). Following staining protocol, the cells were washed in PBS and the cells

were resuspended in FACS buffer and the FACS protocol was performed in BD FACSVerse.

For DCFDA staining, one hour before the indicated time point of infection, cells were

incubated with 10μM DCFDA containing DMEM media at 37°C humidified incubator with

5% CO_2 for 45 min. Post incubation, cells were washed and harvested in PBS. Readings were

measured in BD FACSVerse instrument.

All analysis was done using BD FACSuite software.

Phenol Red-Hydrogen Peroxidase Assay

265   Post 48hr of transfection, cells were infected with STM culture at an MOI of 10. Cells were

266   incubated with Phenol Red solution containing hydrogen peroxidase enzyme (8.5U/ml). At,

267   the designated time points post-infection, the supernatant was collected, and the absorbance

268   was taken at 610 nm in Tecan Plate reader. The exogenously produced H2O2 was quantified

269   using a standard curve of known concentration of H_2O_2.

Gene expression studies by nanoString nCounter technology

Total RNA was isolated at specific time points post infection by using TRIzol (Takara) as per

manufacturer's protocol. Quantification of RNA was performed in Nano Drop (Thermo-

Fischer scientific) and Qubit Bioanalyzer (Agilent 2100 Bioanalyzer). Quality of isolated RNA

was detected by performing 2% agarose gel electrophoresis. Post quality check, samples were

subjected to nanoString nCounter technology (theraCUES). This technology allows multi-plex,

spatially resolved RNA expression quantification with appropriate probes designed against the

target gene.

Lactate Estimation Assay

Cell-supernatant was harvested at the specific time-points post-infection and lactate content of

the sample was estimated using the Lactate Assay Kit (Sigma, Catalog Number-MAK064) as

per manufacturer's protocol. Briefly, 50µl of the sample was added to each 96 well-plate and

each of the well 50µl of the master reaction mix containing 46µl of lactate assay buffer, 2µl of

lactate enzyme mix, and 2µl of lactate probe was added. After the addition of the master

reaction mix to the sample, they were mixed by horizontal shaker or via pipetting. The plate

was incubated for 30 minutes in dark at room temperature. Post incubation, absorbance was

measured at 570nm. The lactate content of the sample was estimated from the lactate standard

curve ranging from 0-10 nmole/µl.

Fatty Acid Oxidation Assay

289   At 16h post-infection, cell pellets were harvested and stored at-80ºC. The fatty acid oxidation

290   assay protocol was followed as per manufacturer's instructions (AssayGenie Fatty Acid

Oxidation (FAO) Assay Kit, Catalogue Code-BR00001). Briefly, cell pellets were lysed using 1X Cell lysis buffer (provided in the kit) and the cell supernatant were obtained post centrifugation of the cell lysate in a cold microfuge at 14,000rpm for 5min. The protein content of the cell supernatants was estimated using Bradford (Bio-Rad) method. 20μl of the protein sample was added to each 96-well plate in duplicate on ice. Each sample was treated with 50μl of control solution and 50μl of reaction solution by swiftly adding one 50μl of control solution to one set of wells and 50μl of reaction solution to the other set of wells. The contents were gently mixed for 10s. The plate is covered and incubated in a 37ºC incubator for 30-60 min (without CO_2). After incubation cherry red colour appears in the wells. The O.D. is measured at 492nm using a plate reader at 30 min,60 min or 120 min. The control well reading was subtracted from the reaction well reading for each sample for each time point. The subtracted reading is used for enzyme activity calculation by considering the incubation time.

Animal Experiment

For all experiments, 6–8 weeks old C57BL/6 or *gp91phox*-/- mice were used. For organ burden analysis, 6 weeks old C57BL/6 or *gp91phox*-/- mice were infected with 10^7 CFU bacteria via oral gavaging. For bacterial enumeration via flow cytometry, mice were infected with 10^7 GFP-expressing bacteria orally. Infected mice were intraperitoneally injected on every other day with either 1 mg/kg body weight of SIRT1 inhibitor EX-527 (Sigma-Aldrich) [32], or SIRT3 inhibitor 3TYP (Selleck Chemical) [33] or SIRT1 activator SRT1720 (Calbiochem, Sigma-Aldrich) [34] or treated with vehicle alone. 5 days post infection, mice were sacrificed, and bacterial organ load was estimated by plating the tissue homogenates on SS agar plates. For calculating percentage survival, 6 weeks old C57BL/6 mice were infected with 10^8 bacteria orally and monitored till fatality. For flow cytometry studies, the harvested liver or spleen were homogenized into single cell suspension and were subjected to flow cytometry. The animal experiments were carried out in accordance with the approved guidelines of the institutional

316   animal ethics committee at the Indian Institute of Science, Bangalore, India (Registration No:

317   48/1999/CPCSEA). All procedures involving the use of animals were performed according to

318   the Institutional Animal Ethics Committee (IAEC)-approved protocol.

Haematoxylin and Eosin Staining

6-8 weeks old C57BL/6 mice were infected with 10^7 bacteria orally. Infected mice were

intraperitoneally injected every alternate with either 1 mg/kg body weight of SIRT1 inhibitor

EX-527, or SIRT3 inhibitor 3TYP or SIRT1 activator SRT1720 or treated with vehicle alone.

5 days post infection, mice were euthanized, and livers were collected and fixed using 3.5%

paraformaldehyde. The fixed liver was then dehydrated using a gradually increasing

concentration of ethanol and embedded in paraffin. 5μm sections were collected on coated

plates. Sections were further rehydrated and then stained with hematoxylin and eosin. Images

were collected in a Leica microscope.

Statistical analysis

329   Data were analyzed and graphed using the GraphPad Prism 8 software (San Diego, CA).

330   Statistical significance was determined by Student's t-test or Two way ANOVA and

331   Bonferroni post-t-test to obtain p values. Adjusted p-values below 0.05 are considered

332   statistically significant. The results are expressed as mean ± SD or SEM of three independent

333   experiments.

RESULTS

Salmonella modulates SIRT1 and SIRT3 expression along its course of infection in macrophages

Upon infection of RAW 264.7 murine macrophages with wildtype *Salmonella* Typhimurium strain 14028S, we observed an increased expression level of SIRT 1 and SIRT3 at initial and middle phases of infection, precisely at 2hr and 6hr post infection through qPCR (Fig.1, A and B). The SIRT1 expression level declined at later phases of infection. On the other hand, the SIRT3 transcript levels remained elevated at all time points with respect to uninfected control with marked increment at 16hr time point post infection which subsided at 20hr post infection. We even monitored the expression profile of SIRT1 and SIRT3 in primary macrophages like peritoneal macrophages of C57BL/6 mice and observed a similar trend of elevated expression at initial (2hr) , middle (6hr) and late (16hr) time points post infection (Fig.1, C and D). In Confocal Laser Scanning Microscopy (CSLM) studies, we observed an increase in SIRT1 expression at 2hr post infection followed by a decrease at 6hr and again a rise at 16hr post infection (Fig.1, E and F). SIRT3 showed progressive increase in expression pattern from 2hr to 16hr time points of infection (Fig.1, G and H). Immunoblotting revealed increased protein expression of both SIRT1 and SIRT3 at 2hr post-infection in comparison to the uninfected control (Fig. S1). However, SIRT1 expression exhibits a gradual decline at the late phase of infection (Fig. S1A-B). In line with the confocal microscopy data, SIRT3 immunoblotting data shows an increased protein expression profile at 6hr and 16hr post-infection (Fig. S1C-D). Thus, an increase in expression profile both at transcript and protein levels indicate their role in *Salmonella* pathogenesis.

SIRT1 and SIRT3 play crucial role in intracellular bacterial proliferation in infected murine macrophages

To evaluate the role of SIRT 1 and SIRT3 in intracellular proliferation of the bacteria within

murine macrophages, we have undertaken knock down of SIRT1 and SIRT3 in RAW 264.7

(Fig. S2) macrophages through PEI mediated transfection of shRNA plasmids directed against

SIRT1 and SIRT3. Post 48hr of transfection, the transfected cells were infected with MOI=10

of wildtype *S*. Typhimurium and gentamicin protection assay was performed. Intracellular

proliferation of the bacteria was quantified by plating the cell lysate at 2hr and 16hr post-

infection. *Salmonella* exhibits compromised intracellular survival in SIRT1 and SIRT3 knock-

down RAW264.7 macrophages in comparison to the un-transfected and scrambled controls

(Fig.2, A). Further, we have assessed the intracellular proliferation in peritoneal macrophages

isolated from thioglycolate treated adult C57BL/6 mice post SIRT1 (EX-527) and SIRT3

(3TYP) inhibitor treatment. SIRT1 or SIRT3 inhibitor treated macrophages exhibited

attenuated intracellular replication in comparison to the untreated peritoneal macrophages

(Fig,2, B). Together, our results depict the role of SIRT1 and SIRT3 in controlling intracellular

proliferation of *S*. Typhimurium.

SIRT1 and SIRT3 inhibition contribute to skewed inflammatory host responses upon

Salmonella **infection**

Several reports indicate the role of SIRT1 and SIRT3 in modulation of host immune responses

pertaining to infection scenario[35][36][37][38]. Therefore, we intend to check whether SIRT1

or SIRT3 regulate immune functions in *Salmonella* infected macrophages**.** To delineate the role

of SIRT1 and SIRT3 in the modulation of immune responses, we wished to investigate the

production of pro-inflammatory and anti-inflammatory cytokines in knock down (KD) RAW

264.7 macrophages upon *S.* Typhimurium infection. After following the transfection for 48hr,

cells were subjected to wildtype *S*. Typhimurium infection at an MOI of 10. At the indicated

time points, cell free supernatant was harvested and evaluated for pro-inflammatory and anti-

inflammatory cytokine production by ELISA. Inhibition of both SIRT1 and SIRT3 resulted in

increase in production of pro-inflammatory cytokine IL-6 significantly at 2hr and 16hr post

infection (Fig.S3, A). Nevertheless, there was only significant reduction in anti-inflammatory

IL-10 production upon knockdown of SIRT3 at 2hr and 16hr post infection but knockdown of

SIRT1 did not show any effect on anti-inflammatory cytokine production, IL-10 (Fig.S3, B).

In peritoneal macrophages upon SIRT1(EX-527-1μM) or SIRT3(3-TYP-1μM) chemical

inhibitor treatment, an increase in IL-6 cytokine levels were observed at 6hr post infection

(Fig.S3, C). This indicates the possible role of SIRT1 and SIRT3 in regulation of cytokine

production upon *Salmonella* infection.

Immune functions are an important determinant of macrophage polarization. Since, SIRT1 and

SIRT3 played an immune-modulatory role in *Salmonella* infection, we investigated whether

Salmonella infection is associated with shift in macrophage polarization status. To assess the

ability of the pathogen to alter the polarization state of the macrophage, we have undertaken

gene expression profiling of various M1 and M2 markers using nanoString nCounter

technology along the course of *Salmonella* Typhimurium infection at the indicated time points

in RAW 264.7 macrophages. A gradual shift from pro-inflammatory M1 toward the anti-

inflammatory M2 state was observed with the progression of *Salmonella* infection. With the

progression of infection there was a reduction in the expression of M1 markers like NOS2,

CD40,CD86, TNF-α, Nfkb2,IL-6 and a corresponding increase in the expression of the M2

markers such as Arg-1, CCL-17, CD 206, IL4ra with an exception of Tgf-β (Fig.3, A).In order

to validate the polarization potency of the pathogen, FACS was performed using a pro-

inflammatory M1 surface marker, CD86 tagged with PE. The data suggests a distinct decrease

in CD86 positive population in the infected sets in comparison to the uninfected and the fixed

dead bacteria control along the course of *S.* Typhimurium infection (Fig.3, B,C). Thus, the live

pathogen has a propensity to skew the polarization state of the macrophage toward an anti-

inflammatory M2 state to subvert the initial acute inflammatory response mounted by the host

immune system.

To assess the role of SIRT1 and SIRT3 in macrophage polarization, we determined anti-

inflammatory CD206 surface marker profiling of the infected macrophages through flow

cytometry. Knockdown of SIRT1 or SIRT3 in infected RAW 264.7 macrophages resulted in

reduction in anti-inflammatory CD206 surface marker expression at 16hr post infection (Fig.

3 D, E). Moreover, SIRT1 or SIRT3 knockdown led to enhanced intracellular ROS generation

within the infected macrophages in comparison to the scrambled or the untransfected control

(Fig.S4) which further gets recruited onto the bacteria (Fig. S5) (Data described later).

NanoString gene profiling in knockdown condition of SIRT1 and SIRT3 infected RAW264.7

macrophages indicated decreased levels of M2 markers such as Arg-1, CD206, IL-10, CCL17,

CCL8, IL4ra, and Tgf-β at 6hr post-infection (Fig. S6A) and at 16hr post-infection, SIRT1

knockdown led to increased M1 gene expression such as CD40 and CD86 expression (Fig.

S6B).   Further, the Haematoxylin and Eosin staining of the *S*. Typhimurium infected mice

liver tissue sections depicted exacerbated signs of inflammation in the SIRT1 (EX-527) or

SIRT3 (3TYP) inhibitor treated cohorts in comparison to the untreated controls with multiple

necrotic foci and increased influx of inflammatory cell inflates. Moreover, these acute

inflammatory signs of the liver sections gets ameliorated in the SIRT1 (SRT1720) activator

treated infected cohort (Fig.9 L,M) (Data described later).   Together, these data suggest the

role of SIRT1 and SIRT3 in modulation of host inflammatory response. Previous literature

reports have shown that SIRT1 physically interacts with p65 subunit of NF-κB and inhibits

transcription by deacetylating p65 at lysine 310 [39]. Moreover, SIRT1 mediated deacetylation

of the p65 subunit of the master regulator of the inflammatory response, NF-κB, results in the

reduction of the inflammatory responses mediated by this transcription factor [40]. To evaluate

the immune regulatory mechanism of SIRT1 in the *S*. Typhimurium (STM) infection scenario,

we undertook SIRT1 immunoprecipitation in the infected RAW264.7 macrophages at 16h post

infection alongside the uninfected macrophages and probed for NF-κB p65 interaction. We

observed an increased interaction of SIRT1 with NF-κB p65 in the infected macrophages in

comparison to the uninfected control (Fig.4, A). Further, the knockdown of SIRT1 resulted in

increased acetylation status of the NF-κB p65 upon infection in comparison to the scrambled,

infected control (Fig.4, B-F) and caused reduced interaction of NF-κB p65 with SIRT1 (Fig.

4E,G). To understand whether the enzymatic domain of SIRT1 possess any role in mediating

this interaction, we carried out NF-κB p65 immunoprecipitation in infected RAW264.7

macrophages in presence or absence of SIRT1 catalytic chemical inhibitor, EX-527(1µM)

treatment at 16h post-infection. We observed an increased interaction of NF-κB p65 with

SIRT1 in the infected untreated macrophages when compared to the untreated uninfected

control (Fig.4, H-I). However, the interaction of NF-κB p65 with SIRT1 gets abrogated under

the EX-527 inhibitor treatment in the infected macrophages thereby implying the function of

the catalytic domain in mediating the interaction (Fig.4, J. Moreover, an increased acetylation

status of NF-κB p65 was observed in the EX-527 treated infected macrophages in comparison

to the untreated infected macrophages (Fig. 4 H-I).

SIRT1 and SIRT3 relieve oxidative stress in infected macrophages and alleviation of the

intracellular ROS generation restores intracellular survival of *S*. Typhimurium within

the SIRT1 or SIRT3 knockdown macrophages

Previous reports have suggested the role of SIRT1 and SIRT3 in oxidative stress conditions.

They are known to act in concert as anti-oxidants by reducing ROS production[41][42][43].

Moreover, enhanced ROS production is also a prototype of the classically activated

macrophages[44][45], which lie at one of the extreme end of the spectra of the macrophage

polarization. Here, we examined the effect of SIRT1 and SIRT3 knockdown in intracellular

ROS generation in *S*. Typhimurium infected RAW264.7 macrophages through DCFDA

staining in FACS. Results depicted significant enhancement in the production of ROS at 16hr post infection upon knockdown of SIRT1 or SIRT3 in comparison to untransfected or scrambled control (Fig.S4, A,B). Upon detection of extracellular ROS generation through Phenol Red Hydrogen Peroxidase assay, we detected higher ROS generation upon SIRT3 KD at 6hr post infection and greater ROS production at 16hr and 20hr time points in SIRT1 knockdown macrophages (Fig.S4,C). Moreover, the increased ROS produced upon SIRT1 or SIRT3 knockdown gets targeted to the bacteria within the infected RAW264.7 macrophages (Fig. S5) Therefore, SIRT1 and SIRT3 play an important role in mitigating the oxidative burst in *Salmonella* infected macrophages. Our previous findings depicted decreased intracellular burden of *S.* Typhimurium within the SIRT1 or SIRT3 knockdown macrophages along with increase in intracellular ROS generation. Therefore, we speculated that the decreased intracellular proliferation within the SIRT1 or SIRT3 knockdown macrophages might be on account of increased intracellular ROS production which might lead to increased killing of the intracellular bacteria. This hypothesis led us to evaluate the intracellular bacterial burden within the infected SIRT1 or SIRT3 knockdown RAW264.7 macrophages or SIRT1 or SIRT3 inhibitor treated peritoneal macrophages (isolated from C57BL/6 adult mice post 5[th] day of thioglycolate injection) in presence of a ROS inhibitor named N-Acetyl Cysteine (NAC). NAC acts as a scavenger of ROS by antagonizing the activity of proteasome inhibitors [46]. The attenuated intracellular proliferation of *Salmonella* Typhimurium within the knockdown or the chemical inhibitor treated macrophages got restored upon 1mM treatment of ROS scavenger, NAC (Fig.S7). Therefore, intracellular ROS production within the knockdown murine macrophages is one of the reasons for the attenuated survival of the bacteria.

SIRT1 and SIRT3 play crucial role in mediating metabolic switch in infected macrophages

Macrophage polarization is not only governed by immunological changes but also contributed

by metabolic reprogramming[47][24][48]**.** Since previous data suggested progression of

Salmonella infection with the shift in polarization state of the macrophage, we decided to

investigate alteration of the metabolic state of the macrophages as macrophage polarization is

governed by immune-metabolic shift. In pursuit of fulfilling such requirement, we performed

gene expression studies of various metabolic genes through nanoString nCounter technology

in *S.* Typhimurium infected RAW 264.7 macrophages. Analysis of the gene profile revealed

upregulation of genes involved in fatty acid oxidization and tricarboxylic acid cycle and

corresponding downregulation of genes involved in glycolysis (Fig.5, A). To validate the

findings, we carried out qRTPCR to quantitatively measure the expression of a fatty acid

oxidation gene, PPARδ in infected RAW 264.7 macrophages. We found that mRNA level was

elevated to 2-fold at 2hr and 6hr post infection. At late phase of infection, 16hr post infection

around 6-fold upregulation in mRNA transcript was noted (Fig.5, B). Lactate estimation assay

of *S.* Typhimurium infected RAW264.7 macrophages at the initial time point of 2hr and at the

late time point of 16hr post infection revealed decline in lactate (glycolysis end product)

production at 16hr in comparison to 2hr post-infection timepoint (Fig. 5C). Together, these

results suggest the capability of the pathogen to drive a shift in metabolic status of the infected

macrophages toward fatty acid oxidation. Next, we evaluated the function of SIRT1 and SIRT3

in influencing the metabolic switch in the infected macrophages through qRT PCR with several

host fatty acid oxidizing genes (*Acox*, *Hadha, Pdha*, *and AcadL)* and glycolytic gene (*PfkL*) in

SIRT1 and SIRT3 knockdown macrophages and via lactate production assay. Lactate

estimation assay in SIRT1 and SIRT3 knockdown condition revealed enhanced lactate

production at 16hr post infection in comparison to the scrambled control which further

authenticate the increased host glycolysis upon SIRT1 and SIRT3 knockdown scenario (Fig.

5D,E). SIRT1 and SIRT3 knockdown RAW 264.7 infected macrophages showed decreased

expression of fatty acid oxidizing genes and increased expression of glycolytic *PfkL* gene in

comparison to the scrambled infected control (Fig.5, F). Metabolic gene expression studies in

SIRT1 or SIRT3 knockdown RAW264.7 macrophages via nanoString gene profiling depicted

decreased expression of fatty acid metabolic genes such as *Acadl*, *Pgc1α*, *Pgc1β*, *Pparα*, *Pparδ*,

Pparγ (Fig. S6C). Similar results were obtained in the infected peritoneal macrophages under

the SIRT1 or SIRT3 catalytic inhibitor treatment (Fig. 5 G, H) Moreover, qRT PCR mediated

metabolic gene profiling of liver and spleen isolated from wildtype *S.* Typhimurium infected

macrophages revealed decreased transcript level expression of murine fatty acid oxidation

genes upon treatment with SIRT1 (EX-527) or SIRT3 (3TYP) catalytic inhibitors which got

reversed upon SIRT1 activator (SRT1720) treatment (Fig.5, I, J). Moreover, SIRT1 and SIRT3

knockdown or catalytic inhibition in RAW264.7 macrophages resulted in increased protein

expression of host glycolytic genes such as Phosphoglycerate kinase (Pgk),

Phosphofructokinase (Pfk) with concomitant reduction in protein expression of TCA cycle

gene like Pyruvate dehydrogenase (Pdha1) and fatty acid oxidation genes such as Hadha and

Acox1 (Fig.5 K-M, Fig.S8). Fatty acid oxidation assay in the RAW264.7 macrophages under

SIRT1 or SIRT3 knockdown condition or inhibition treatment revealed significant decrease in

fatty acid β oxidation activity of the infected macrophages in comparison to the scrambled or

the untreated control (Fig.5 N).

Collectively, these data suggest the role of SIRT1 and SIRT3 in mediating the host metabolic

shift in the infected macrophages.

SIRT1 and SIRT3 concomitantly influences *Salmonella* metabolism

Our previous data indicated a shift in host metabolism toward increased fatty acid oxidation

along the course of *Salmonella* Typhimurium infection in murine RAW 264.7 macrophages.

Salmonella Typhimurium drives the metabolism of the infected macrophage toward fatty acid

oxidation. This observation led us to investigate the influence of host metabolic shift on the

metabolic status of the pathogen harboring inside the infected macrophages[49]. We were

intrigued whether increased glucose availability within the fatty-acid oxidizing macrophages

is utilized by the bacteria. Thus, we undertook simultaneous gene expression studies of various

Salmonella genes involved in their pathogenesis and metabolism through nanoString nCounter

technology in *S.* Typhimurium infected RAW 264.7 macrophages. The nanoString gene profile

revealed enhanced expression of genes involved in *Salmonella* glycolysis and glucose uptake

such as *pfkA* and *ptsG*, respectively (Fig.6, A). This finding indicates the ability of the pathogen

to drive the metabolic state of the host toward fatty acid oxidation with corresponding increased

glucose utilization by the bacteria favoring their survival inside the host. qRT PCR results with

several bacterial fatty acid oxidizing genes (*fadA, fadB,fadL,aceA, aceB*) and glycolytic genes

(*ptsG*) in knockdown condition of SIRT1 and SIRT3 in RAW 264.7   macrophages further

validated the nanoString gene expression profiles (Fig.6, B). In scrambled control, *Salmonella*

infection progresses with increased *Salmonella* glycolytic flux and reduced bacterial fatty acid

oxidation. However, knockdown of SIRT1 and SIRT3 abrogate this bacterial metabolic shift

by reducing its glycolytic flux and by exhibiting enhanced fatty acid oxidation thereby

attenuating pathogen intracellular survival. Similar observations were obtained from the qPCR

data in the infected mice liver and spleen samples with increased transcript level expression of

bacterial fatty acid oxidation genes and decreased expression of bacterial glycolytic genes upon

SIRT1 or SIRT3 inhibitor treatment (Fig. 6 C, D). Therefore, SIRT1 and SIRT3 driven host

metabolic switch potentially influence the metabolic profile of the intracellular pathogen.

Mechanism behind SIRT1 or SIRT3 mediated metabolic switch

As per our previous findings, SIRT1 or SIRT3 inhibition led to increased host glycolysis and

decline in fatty oxidation in the infected macrophages. HIF1α is a master regulator of glycolysis

in host during stress conditions [50]. Previous reports have suggested HIF1α to be a target of

deacetylation by SIRT1 at Lys 674 which contribute to metabolic reprogramming in cancer

cells. During hypoxia, downregulation of SIRT1 leads to increased acetylation and activation

of HIF1α [51]. Additionally, in CD4+ T cells, ectopic expression of SIRT1 inhibited IL-9

production and glycolysis by negatively regulating HIF1α [52]. To delve into the mechanism

behind SIRT1 mediated modulation of metabolic responses, we assessed the interaction of

SIRT1 with HIF-1α in infected RAW264.7 macrophages. The immunoprecipitation studies of

SIRT1 showed increased interaction of the SIRT1 with HIF1α in the *S.* Typhimurium infection

scenario with respect to the uninfected control (Fig.7, A). Further, we evaluated the acetylation

status of HIF1α in the SIRT1 knockdown status of the infected macrophages. We found that

SIRT1 knockdown showed increased acetylation of HIF1α in the infected macrophages in

comparison to the scrambled infected control at 16hr post infection (Fig. 7, B,C).

Immunoprecipitation studies under SIRT1 (EX-527) inhibitor treatment in RAW264.7

macrophages revealed increased acetylation of HIF1-α along with reduced interaction of HIF-

1α with SIRT1, thereby indicating the probable role of the catalytic domain in influencing the

interaction (Fig. 7 D-F). Together, our results implicate the role of SIRT1 in governing

glycolytic shift in the infected macrophages by deacetylating HIF1α. Upon SIRT1 knockdown

or inhibition, HIF1α gets hyperacetylated which cause activation of the downstream glycolytic

genes. Alternatively, several key literatures suggest the role of SIRT3 in modulating metabolic

programming by deacetylating several proteins involved in fatty acid oxidation, the

tricarboxylic acid cycle and oxidative phosphorylation [53][54]. PDHA1 (Pyruvate

Dehydrogenase E1 subunit alpha) is a key enzyme linking glycolysis to TCA cycle and

oxidative phosphorylation. SIRT3 regulates PDHA1 acetylation by deacetylating PDHA1 at

lysine 385 residue, thereby playing a key role in metabolic reprogramming [54]. PDHA1

acetylation coincides with PDH activity and increased PDHA1 phosphorylation [54].

Therefore, we investigated the role of SIRT3 in the modulation of the host fatty acid oxidation

upon *S.* Typhimurium infection in RAW264.7 macrophages. To do so, we immunoprecipitated

PDHA1 and checked for its interaction with SIRT3 under knockdown condition of SIRT3 or

upon SIRT3 inhibitor treatment. We observed an increased interaction of PDHA1 with SIRT3

in the infection scenario in comparison to the uninfected control which gets eventually

abolished under the knockdown condition and under the chemical inhibitor treatment of SIRT3

suggesting the role of the SIRT3 in mediating the interaction with PDHA1 (Fig.8, A-D).

SIRT1 or SIRT3 inhibition enhances bacterial burden in mice *in vivo*

6-8 weeks old adult male C57BL/6 mice were treated with SIRT1 inhibitor EX-527, SIRT3

inhibitor 3TYP and SIRT1 activator SRT1720 at a dose of 1mg/kg each via intraperitoneal

injection (every alternate Day) (Fig. 9A). Following the inhibitor treatment, the mice were

orally gavaged with 10^7 cfu of *S.* Typhimurium 14028S for organ burden evaluation or with

10^8 cfu of wildtype *S.* Typhimurium for survival studies. On day 5^{th} post infection, mice were

sacrificed, and the liver, spleen and Mesenteric Lymph Node (MLN) were harvested for

enumeration of the organ burden. The SIRT1 inhibitor, EX-527 and SIRT3 inhibitor, 3TYP

treated mice cohorts exhibited increased organ loads in liver, spleen and MLN in comparison

to the vehicle control. On the contrary, the SRT1720 treated mice group showed organ burden

comparable to that of the vehicle control (Fig. 9, B). Moreover, the SIRT1 and the SIRT3

inhibitor treated mice cohorts died earlier than the vehicle treated control mice group or the the

SIRT1 activator treated group (Fig. 9, C). Further, the SIRT1 and the SIRT3 inhibitor treated

mice cohorts showed increased splenic length in comparison to the vehicle treated mice group

and the SIRT1 activator treated mice cohort (Fig. 9, D). The increased organ burden in the EX-

527 or 3TYP treated group might be due to increased bacterial dissemination in blood. To

assess bacterial dissemination, blood was collected from infected mice post inhibitor treatment

at specific days post infection retro-orbitally and plated onto *Salmonella Shigella* (SS) agar

plates for bacterial enumeration. Indeed, increased bacterial dissemination was observed in the

blood of mice treated with SIRT1 inhibitor, EX-527 or SIRT3 inhibitor, 3TYP at day 1-, 2-, 3-

,4- and 5- day post infection in comparison to the vehicle treated mice (Fig. 9, E). Further, we

wanted to examine whether the increased bacterial dissemination was due to increased ROS

production or due to the presence of elevated inflammatory cytokine levels like IL-6 in the

serum. In the wildtype C57BL/6 mice treated with SIRT3 inhibitor 3TYP showed heightened

bacterial burden in blood at 5^{th} day post infection in comparison to the vehicle control.

Nevertheless, the *gp91phox*-/- mice group did not depict significant variation in the bacterial

load among the different mice treatment cohorts (Fig. 9, F). Further, the EX-527 (SIRT1

inhibitor) and the 3TYP (SIRT3 inhibitor) treated mice possessed elevated levels of serum IL-

6 in comparison to the vehicle treated control and the SRT1720 (SIRT1 activator) treated mice

group (Fig. 9, G). The increased mouse serum IL-6 production was in similar line with the

increased IL-6 cytokine generation in EX-527 or 3TYP treated peritoneal macrophages under

infection scenario (Fig. S3, C). However, contrary to the *in vitro* studies wherein SIRT1 or

SIRT3 knockdown or inhibition resulted in attenuated intracellular proliferation, here in *in vivo*

mouse model of infection, we observed increased bacterial organ loads owing to increased

bacterial dissemination. To delineate this observation further, we evaluated the bacterial load

within splenocytes isolated from control or inhibitor treated C57BL/6 mice infected with GFP

expressing *S.* Typhimurium at 5^{th} day post-infection via flow cytometry. We observed

heightened bacterial load in the EX-527 or the 3TYP treated mice cohorts (Fig. 9, H, I).

However, when we enumerated the bacterial count within the F4/80+ macrophage population

of the infected splenocytes, we noticed decreased bacterial loads in the EX-527 or 3TYP treated

mice group in comparison to the vehicle treated control group or the SRT-1720 activator treated

group (Fig. J, K). This opposing phenotype could be attributed to the increased IL-6 cytokine

storm and elevated ROS production upon the SIRT1 or SIRT3 inhibitor treatment which in

turn resulted in bacterial dissemination *in vivo* and concomitantly restricted the *in vitro*

intracellular proliferation within macrophages. To validate this observation, we estimated the

ROS levels within the liver and spleen tissues harvested from *S.* Typhimurium infected

C57BL/6 mice, treated with specific catalytic inhibitor, activator or vehicle via DCFDA

staining using flow cytometry at 5^{th} day post infection. We detected escalated levels of ROS

within both the infected liver and spleen tissues of the EX-527 or 3TYP treated mice groups in

comparison to the vehicle treated or the SRT1720 treated mice cohorts (Fig. S9). Haematoxylin

and eosin staining of the liver sections (harvested at 5^{th} day post infection) revealed increased

inflammation with multiple areas of severe acute hepatic necrosis with complete loss of hepatic

architecture in the EX-527 and 3-TYP treated liver samples in comparison to the vehicle treated

control and SRT-1720 treated liver samples (Fig. 9, L,M). Altogether, our results implicate the

role of SIRT1 and SIRT3 in controlling *S.* Typhimurium infection *in vivo*.

DISCUSSION

Several studies have confirmed the propensity of *Salmonella* to skew the polarization state of

the infected macrophages toward an immunosuppressive anti-inflammatory state[55][56][57].

We have validated such findings and further elaborated it by depicting the role of SIRT1 and

SIRT3 in the modulation of host immune responses. *Salmonella* Typhimurium infection

modulates the expression profile of both SIRT1 and SIRT3 in the infected macrophages at both

mRNA and protein level. SIRT1 show reduced mRNA and protein expression at late time point

of infection (16hr p.i) in comparison to the initial time point of infection (2hr p.i). On the other

hand, SIRT3 depicts enhanced expression levels even at later points of infection both at the

transcript and protein level. Downregulation of SIRT1 and SIRT3 through shRNA mediated

knockdown resulted in heightened pro-inflammatory immune responses with increased

production of IL-6 cytokine and decreased surface expression of anti-inflammatory CD206.

SIRT1 and SIRT3 downregulation also resulted in increased intracellular ROS production in

the infected macrophages. SIRT1 and SIRT3 knockdown macrophages not only show altered

host immune status but also depicted shift in the metabolic state with increased glycolytic shift.

This altered host metabolism upon SIRT1 and SIRT knockdown condition influence the

outcome of infection by regulating the intracellular bacterial metabolism which show reduced

bacterial glycolysis and increased fatty acid oxidation. All these outcomes account for the

reason for attenuated intracellular bacterial proliferation in the SIRT1 and SIRT3 knockdown

macrophages. However, in murine model of infection, SIRT1 or SIRT3 inhibitor treatment led

to increased organ burden and triggered bacterial dissemination. Overall, our study highlights

the crucial role of SIRT1 and SIRT3 in governing the immune-metabolic shift during

Salmonella infection, and suggests that targeting these proteins may be a potential therapeutic

strategy for treating *Salmonella* infections in the future.

Previous reports have elucidated the role of SIRT1 and SIRT2 pertaining to *Salmonella*

infection. Ganesan et al., have depicted the role of SIRT1 in autophagy in *Salmonella* infection

scenario[58]. Gogoi et al., have demonstrated SIRT2 mediated modulation of immune

responses in dendritic cells[59]. Till date, there is no report highlighting the role of SIRT3

governing the *Salmonella* pathogenesis. The function of SIRT3 in infection scenario has been

explored quite recently. In *Mycobacterium tuberculosis* infection condition, SIRT3 control

mitochondrial function and autophagy[38]. SIRT3 downregulation in *Mycobacterium*

tuberculosis infected macrophages is associated with dysregulated mitochondrial metabolism

and increased cell death[60]. In this study we have explored the role of SIRT1 and SIRT3 in

mediating immune-metabolic switch in *Salmonella* Typhimurium infected macrophages.

Our findings suggest the role of SIRT1 and SIRT3 in mediating polarization of the *Salmonella*

infected macrophages toward an anti-inflammatory state. Upon knockdown of SIRT1 and

SIRT3 in the infected macrophages we detect robust pro-inflammatory response and oxidative

burst. This is in line with the findings by S.Elsela et al., wherein SIRT1 knockout

RSV(Respiratory Syncytial Virus) - infected BMDCs showed significant increase in *Il1β*, *Il6*

and *Il23* expression and ROS generation in comparison to the wild type RSV-infected

BMDCs[35]. Also, Kim et al., showed presence of aggravated inflammatory responses in

Mycobacterium tuberculosis infected SIRT3-/- BMDMs[38]. This heightened pro-

inflammatory cytokine burst and oxidative burst restrict the intracellular survival of the

pathogen as detected by the lower intracellular burden in the SIRT1 and SIRT3 knockdown

murine macrophages. *Salmonella* showed enhanced proliferation in the M2 macrophages

owing to the reduced arsenals in terms of pro-inflammatory cytokines, ROS production and

iNOS generation. Moreover, the M2 macrophages are fuelled by increased fatty acid oxidation

and reduced glycolysis[61]. This might facilitate enhanced bacterial proliferation as the host

un-utilized intracellular glucose can be readily up taken by the pathogen and used to support

its own glycolysis. Similarly, M1 or pro-inflammatory macrophages resort to glycolysis to

meet their energy demands[62] thereby limiting the glucose availability for the intracellular

pathogen[62][63]. In such condition, bacteria show enhanced fatty acid metabolism to sustain

their energy demand[64]. In our study, we found that wild type *S*. Typhimurium infection drive

host metabolism toward increased fatty acid oxidation with concomitant increase in the

bacterial glycolysis. SIRT1 and SIRT3 inhibition abrogates the metabolic switch and triggers

increase in host glycolysis which in turn skew the bacterial metabolism from increased

glycolysis to enhanced fatty acid oxidation and reduced glycolysis. Together, these findings

implicate the role of SIRT1 and SIRT3 in reprograming the host metabolism which in turn

affect the intracellular nutrient niche of the pathogen thereby influencing intracellular

Salmonella proliferation. However, our in vivo findings in the murine model of infection show

increased bacterial burden upon SIRT1 or SIRT3 inhibition. This increased burden could be

attributed to increased dissemination from the macrophages into the bloodstream owing to the

increased level of serum IL-6 levels. This is in concert with previous findings in *Klebsiella*

pneumoniae infection in mice wherein increased inflammatory response upon HIF-1α induces

bacterial dissemination [65]. Further correlation analysis of immune responses to *Salmonella*

infection revealed that increased innate immune "cassette" opposes the adaptive immune arm

leading to increased bacterial load [66]. Future studies might explore the interacting partners

of SIRT1 and SIRT3 through mass spectrometry analyses in *Salmonella* infected macrophages

which might hint at the underlying mechanism of their action and regulation. Together, this

study highlights the complex and multifaceted nature of host-pathogen interactions, and the

need for further research to fully understand the role of SIRT1 and SIRT3 in the context of

Salmonella infection.

AUTHOR CONTRIBUTION

DH and DC have conceptualized the study. DH has contributed to the experiment designing,

visualization, methodology, investigation, formal analysis, literature survey, validation, writing

(original draft), reviewing and editing of the manuscript. RSR has performed the

histopathological scoring of the liver sections. DC has contributed to the funding acquisition,

project administration, and overall supervision of the work. All authors approved the final

version of the manuscript.

DECLARATION OF INTEREST

728 The authors are unaware of any conflicting interests and thereby declare no conflict of interest.

ACKNOWLEDGEMENT

This work was supported by the DAE SRC fellowship (DAE00195) and DBT-IISc partnership

umbrella program for advanced research in biological sciences and Bioengineering to DC.

Infrastructure support from ICMR (Centre for Advanced Study in Molecular Medicine), DST

(FIST), and UGC (special assistance) is highly acknowledged. DH sincerely acknowledges the

CSIR- SPM fellowship for her financial support. The funders had no role in study design, data

collection and analysis, decision to publish, or preparation of the manuscript.

REFERNCES

1. Landry J, Sutton A, Tafrov ST, Heller RC, Stebbins J, Pillus L, et al. The silencing protein SIR2 and its
homologs are NAD-dependent protein deacetylases. Proc Natl Acad Sci U S A. 2000;97(11):5807–11.

2. Jackson MD, Denu JM. Structural identification of 2'- and 3'-O-acetyl-ADP-ribose as novel
metabolites derived from the Sir2 family of beta -NAD+-dependent histone/protein deacetylases. J
Biol Chem. 2002;277(21):18535-18544. doi:10.1074/jbc.M200671200

3. Sauve AA, Celic I, Avalos J, Deng H, Boeke JD, Schramm VL. Chemistry of gene silencing: The
mechanism of NAD+-dependent deacetylation reactions. Biochemistry. 2001;40(51):15456–63.

4. Bitterman KJ, Anderson RM, Cohen HY, Latorre-Esteves M, Sinclair DA. Inhibition of silencing and
accelerated aging by nicotinamide, a putative negative regulator of yeast Sir2 and human SIRT1. J
Biol Chem. 2002;277(47):45099–107.

5. Sanders BD, Jackson B, Marmorstein R. Structural basis for sirtuin function: What we know and
what we don't. Biochim Biophys Acta - Proteins Proteomics 2010;1804(8):1604–16. Available from:
http://dx.doi.org/10.1016/j.bbapap.2009.09.009 10. Lin Z, Fang D. The Roles of SIRT1 in Cancer.
Genes and Cancer. 2013;4(3–4):97– 104.

6. Lin Z, Fang D. The Roles of SIRT1 in Cancer. Genes and Cancer. 2013;4(3–4):97–104.

7. Guarente L. Sirtuins in aging and disease. Cold Spring Harb Symp Quant Biol. 2007;72:483–8.

8. Martínez-Redondo P, Vaquero A. The Diversity of Histone Versus Nonhistone Sirtuin Substrates.
Genes and Cancer. 2013;4(3–4):148–63.

9. Liu TF, Yoza BK, El Gazzar M, Vachharajani VT, McCall CE. NAD+-dependent SIRT1 deacetylase
participates in epigenetic reprogramming during endotoxin tolerance. J Biol Chem.
2011;286(11):9856–64.

10. El Gazzar M, Yoza BK, Hu JYQ, Cousart SL, McCall CE. Epigenetic silencing of tumor necrosis factor
α during endotoxin tolerance. J Biol Chem. 2007;282(37):26857–64.

11. Liu TF, Vachharajani V, Millet P, Bharadwaj MS, Molina AJ, McCall CE. Sequential actions of
SIRT1-RELB-SIRT3 coordinate nuclear-mitochondrial communication during immunometabolic
adaptation to acute inflammation and sepsis. J Biol Chem. 2015;290(1):396–408.

12. Kong X, Wang R, Xue Y, et al. Sirtuin 3, a new target of PGC-1alpha, plays an important role in the
suppression of ROS and mitochondrial biogenesis. PLoS One. 2010;5(7):e11707.

13. van de Ven RAH, Santos D, Haigis MC. Mitochondrial Sirtuins and Molecular Mechanisms of
Aging. Trends Mol Med. 2017;23(4):320–31.

14. Garai, P., D.P. Gnanadhas, and D. Chakravortty, Salmonella enterica serovars Typhimurium and
Typhi as model organisms: revealing paradigm of host-pathogen interactions. Virulence, 2012. **3**(4): p.
377-88

15.Bhutta ZA, Threlfall J. Addressing the global disease burden of typhoid
fever. *JAMA*.2009;302(8):898-899. doi:10.1001/jama.2009.1259

16. Majowicz SE, Musto J, Scallan E, et al. The global burden of nontyphoidal Salmonella
gastroenteritis. Clin Infect Dis. 2010;50(6):882-889. doi:10.1086/650733

17. Haraga, A., M.B. Ohlson, and S.I. Miller, Salmonellae interplay with host cells. Nat Rev Microbiol,
2008. **6**(1): p. 53-66.

18. Hajra, D., A.V. Nair, and D. Chakravortty, An elegant nano-injection machinery for sabotaging the
host: Role of Type III secretion system in virulence of different human and animal pathogenic
bacteria. Phys Life Rev, 2021. **38**: p. 25-54.

19. Lawrence T, Natoli G. Transcriptional regulation of macrophage polarization: Enabling diversity
with identity. Nat Rev Immunol. 2011;11(11):750–61.

20. Martinez FO, Helming L, Gordon S. Alternative Activation of Macrophages: An Immunologic
Functional Perspective. Annu Rev Immunol. 2009;27(1):451–83.

21. Namgaladze D, Brüne B. Fatty acid oxidation is dispensable for human macrophage IL-4-induced
polarization. Biochim Biophys Acta - Mol Cell Biol Lipids 2014;1841(9):1329–35.

22. Hui X, Zhang M, Gu P, Li K, Gao Y, Wu D. Adipocyte SIRT 1 controls systemic insulin sensitivity by
modulating macrophages in adipose tissue. 2017;18(4):645–57.

23. Jia Y, Han S, Li J, et al. IRF8 is the target of SIRT1 for the inflammation response in macrophages.
Innate Immun. 2017;23(2):188-195. doi:10.1177/1753425916683751

24. Kratz M, Coats BR, Hisert KB, Hagman D, Mutskov V, Peris E, et al. Metabolic dysfunction drives a
mechanistically distinct proinflammatory phenotype in adipose tissue macrophages. Cell Metab
2014;20(4):614–25. Available from: http://dx.doi.org/10.1016/j.cmet.2014.08.010

25. Cimen H, Han MJ, Yang Y, Tong Q, Koc H, Koc EC. Regulation of succinate dehydrogenase activity
by SIRT3 in mammalian mitochondria. Biochemistry. 2010;49(2):304–11.

26. Vazquez-Torres A, Xu Y, Jones-Carson J, Holden DW, Lucia SM, Dinauer MC, et al. Salmonella
pathogenicity island 2-dependent evasion of the phagocyte NADPH oxidase. Science (80-).
2000;287(5458):1655–8.

27. Miller BH, Fratti RA, Poschet JF, Timmins GS, Master SS, Burgos M, et al. Mycobacteria Inhibit
Nitric Oxide Synthase Recruitment to Phagosomes during Macrophage Infection. Infect Immun.
2004;72(5):2872–8.

28. Bost KL, Clements JD. Intracellular Salmonella dublin induces substantial secretion of the 40-
kilodalton subunit of interleukin-12 (IL-12) but minimal secretion of IL-12 as a 70-kilodalton protein
in murine macrophages. Infect Immun. 1997;65(8):3186–92.

29. Pathak SK, Basu S, Basu KK, Banerjee A, Pathak S, Bhattacharyya A, et al. Direct extracellular
interaction between the early secreted antigen ESAT-6 of Mycobacterium tuberculosis and TLR2
inhibits TLR signaling in macrophages. Nat Immunol. 2007;8(6):610–8.

30. Tumitan ARP, Monnazzi LGS, Ghiraldi FR, Cilli EM, De Medeiros BMM. Pattern of macrophage
activation in yersinia-resistant and yersinia-susceptible strains of mice. Microbiol Immunol.
2007;51(10):1021–8.

31. Brubaker RR. Interleukin-10 and inhibition of innate immunity to Yersiniae: Roles of Yops and
LcrV (V antigen). Infect Immun. 2003;71(7):3673–81.

32. Morató L, Astori S, Zalachoras I, et al. eNAMPT actions through nucleus accumbens NAD^+/SIRT1
link increased adiposity with sociability deficits programmed by peripuberty stress. Sci Adv.
2022;8(9):eabj9109. doi:10.1126/sciadv.abj9109.

33. Hu B, Wang P, Zhang S, et al. HSP70 attenuates compression-induced apoptosis of nucleus
pulposus cells by suppressing mitochondrial fission via upregulating the expression of SIRT3. Exp Mol
Med. 2022;54(3):309-323. doi:10.1038/s12276-022-00745-9.

34. Tan P, Wang M, Zhong A, et al. SRT1720 inhibits the growth of bladder cancer in organoids and
murine models through the SIRT1-HIF axis. *Oncogene*. 2021;40(42):6081-6092. doi:10.1038/s41388-
021-01999-9.

35. Elesela S, Morris SB, Narayanan S, Kumar S, Lombard DB, Lukacs NW. Sirtuin 1 regulates
mitochondrial function and immune homeostasis in respiratory syncytial virus infected dendritic
cells. PLoS Pathog. 2020;16(2):e1008319.

36. Liu G, Bi Y, Xue L, et al. Dendritic cell SIRT1-HIF1α axis programs the differentiation of CD4+ T
cells through IL-12 and TGF-β1. Proc Natl Acad Sci U S A. 2015;112(9):E957-E965.
doi:10.1073/pnas.1420419112.

37. Yang H, Hu J, Chen Y (Jason), Ge B. Role of Sirt1 in innate immune mechanisms against
Mycobacterium tuberculosis via the inhibition of TAK1 activation. Arch Biochem Biophys
2019;667(October 2018):49–58. Available from: https://doi.org/10.1016/j.abb.2019.04.006

38. Kim TS, Jin YB, Kim YS, Kim S, Kim JK, Lee HM, et al. SIRT3 promotes antimycobacterial defenses
by coordinating mitochondrial and autophagic functions Vol. 15, Autophagy. Taylor & Francis; 2019.
1356–1375 p. Available from: https://doi.org/10.1080/15548627.2019.1582743.

39. Yeung F, Hoberg JE, Ramsey CS, et al. Modulation of NF-kappaB-dependent transcription and cell
survival by the SIRT1 deacetylase. EMBO J. 2004;23(12):2369-2380. doi:10.1038/sj.emboj.7600244.

40. Yang H, Zhang W, Pan H, et al. SIRT1 activators suppress inflammatory responses through
promotion of p65 deacetylation and inhibition of NF-κB activity. PLoS One. 2012;7(9):e46364.
doi:10.1371/journal.pone.0046364.

41. Singh CK, Chhabra G, Ndiaye MA, Garcia-peterson LM, Mack NJ, Ahmad N. The Role of Sirtuins in
Antioxidant and Redox Signaling. 2018;28(8):643–61.

42. Kitada M, Ogura Y, Monno I, Koya D. Sirtuins and Type 2 Diabetes: Role in Inflammation,
Oxidative Stress, and Mitochondrial Function. Front Endocrinol (Lausanne). 2019;10:187.

43. Merksamer PI, Liu Y, He W, Hirschey MD, Chen D, Verdin E. The sirtuins, oxidative stress and
aging: an emerging link. Aging (Albany NY). 2013;5(3):144-150. doi:10.18632/aging.100544.

44. Rendra E, Riabov V, Mossel DM, Sevastyanova T, Harmsen MC, Kzhyshkowska J. Immunobiology
Reactive oxygen species (ROS) in macrophage activation and function in diabetes. Immunobiology
2019;224(2):242–53. Available from: https://doi.org/10.1016/j.imbio.2018.11.010

45. Zhou Y, Que KT, Zhang Z, et al. Iron overloaded polarizes macrophage to proinflammation
phenotype through ROS/acetyl-p53 pathway. Cancer Med. 2018;7(8):4012-4022.
doi:10.1002/cam4.1670.

46. Halasi M, Wang M, Chavan TS, Gaponenko V, Hay N, Gartel AL. ROS inhibitor Nacetyl-L-cysteine
antagonizes the activity of proteasome inhibitors. Biochem J. 2013;454(2):201–8.

47.De Santa F, Vitiello L, Torcinaro A, Ferraro E. The Role of Metabolic Remodeling in Macrophage
Polarization and Its Effect on Skeletal Muscle Regeneration. Antioxid Redox Signal.
2019;30(12):1553-1598. doi:10.1089/ars.2017.7420.

48. Galván-Peña S, O'Neill LAJ. Metabolic reprogramming in macrophage polarization. Front
Immunol. 2014;5(AUG):1–6.

49. Taylor SJ, Winter SE. Salmonella finds a way: Metabolic versatility of Salmonella enterica serovar
Typhimurium in diverse host environments. PLoS Pathog 2020;16(6):10–5. Available from:
http://dx.doi.org/10.1371/journal.ppat.1008540

50. Kierans SJ, Taylor CT. Regulation of glycolysis by the hypoxia-inducible factor (HIF): implications
for cellular physiology. J Physiol. 2021;599(1):23-37. doi:10.1113/JP280572.

51. Lim JH, Lee YM, Chun YS, Chen J, Kim JE, Park JW. Sirtuin 1 modulates cellular responses to
hypoxia by deacetylating hypoxia-inducible factor 1alpha. Mol Cell. 2010;38(6):864-878.
doi:10.1016/j.molcel.2010.05.023.

52. Wang Y, Bi Y, Chen X, et al. Histone Deacetylase SIRT1 Negatively Regulates the Differentiation of
Interleukin-9-Producing CD4(+) T Cells. Immunity. 2016;44(6):1337-1349.
doi:10.1016/j.immuni.2016.05.009

53. Hirschey MD, Shimazu T, Goetzman E, et al. SIRT3 regulates mitochondrial fatty-acid oxidation by
reversible enzyme deacetylation. Nature. 2010;464(7285):121-125. doi:10.1038/nature08778

54. Zhang Y, Wen P, Luo J, et al. Sirtuin 3 regulates mitochondrial protein acetylation and
metabolism in tubular epithelial cells during renal fibrosis. Cell Death Dis. 2021;12(9):847.

55. Panagi I, Jennings E, Zeng J, Günster RA, Stones CD, Mak H, et al. Salmonella Effector SteE
Converts the Mammalian Serine/Threonine Kinase GSK3 into a Tyrosine Kinase to Direct
Macrophage Polarization. Cell Host Microbe. 2020;27(1):41-53.e6.

56. Stapels DAC, Hill PWS, Westermann AJ, Fisher RA, Thurston TL, Saliba AE, et al. Salmonella
persisters undermine host immune defenses during antibiotic treatment. Science.
2018;362(6419):1156–60.

57. Pham THM, Brewer SM, Thurston T, Massis LM, Honeycutt J, Lugo K, et al. Salmonella-Driven
Polarization of Granuloma Macrophages Antagonizes TNFMediated Pathogen Restriction during
Persistent Infection. Cell Host Microbe. 2020;27(1):54-67.e5. Available from:
https://doi.org/10.1016/j.chom.2019.11.011

58. Ganesan R, Hos NJ, Gutierrez S, Fischer J, Stepek JM, Daglidu E, et al. Salmonella Typhimurium
disrupts Sirt1/AMPK checkpoint control of mTOR to impair autophagy. PLoS Pathog. 2017;13(2):1–
22.

59. Gogoi M, Chandra K, Sarikhani M, Ramani R, Sundaresan NR, Chakravortty D. Salmonella escapes
adaptive immune response via SIRT2 mediated modulation of innate immune response in dendritic
cells. PLoS Pathog. 2018;14(11):1–22.

60. Smulan LJ, Martinez N, Kiritsy MC, Kativhu C, Cavallo K, Sassetti CM, et al. Sirtuin 3
downregulation in mycobacterium tuberculosis-infected macrophages reprograms mitochondrial
metabolism and promotes cell death. MBio. 2021;12(1):1–15.

61. Eisele NA, Ruby T, Jacobson A, Manzanillo PS, Cox JS, Lam L, et al. Salmonella require the fatty
acid regulator PPARδ for the establishment of a metabolic environment essential for long-term
persistence. Cell Host Microbe 2013;14(2):171– 82. Available from:
http://dx.doi.org/10.1016/j.chom.2013.07.010

62. Cramer T, Yamanishi Y, Clausen BE, et al. HIF-1alpha is essential for myeloid cell mediated
inflammation. Cell. 2003;112(5):645-657. doi:10.1016/s0092-8674(03)00154-5.

63. Merrill GF, Kurth EJ, Hardie DG, Winder WW. AICA riboside increases AMP activated protein
kinase, fatty acid oxidation, and glucose uptake in rat muscle. Am J Physiol. 1997;273(6):E1107-
E1112. doi:10.1152/ajpendo.1997.273.6.E1107.

64. Reens AL, Nagy TA, Detweiler CS. Salmonella enterica Requires Lipid Metabolism Genes To
Replicate in Proinflammatory Macrophages and Mice. Infect Immun. 2019;88(1):e00776-19

65. Holden VI, Breen P, Houle S, Dozois CM, Bachman MA. Klebsiella pneumoniae Siderophores
Induce Inflammation, Bacterial Dissemination, and HIF-1α Stabilization during Pneumonia. mBio.
2016;7(5):e01397-16. Published 2016 Sep 13. doi:10.1128/mBio.01397-16

66. Hotson AN, Gopinath S, Nicolau M, et al. Coordinate actions of innate immune responses oppose
those of the adaptive immune system during Salmonella infection of mice. Sci Signal.
2016;9(410):ra4. Published 2016 Jan 12. doi:10.1126/scisignal.aaa9303

MANUSCRIPT_1 TITLE:

SIRT1 and SIRT3 mediated immune-metabolic switch is crucial for the establishment of *Salmonella* pathogenicity of infection.

MAIN FIGURES

Fig.1- *Salmonella* modulates the expression of Sirtuin 1 and Sirtuin3 along its course of infection

A-B- Expression studies through qPCR in RAW 264.7 macrophages. Data is representative of N=3, n=2.

C-D-Expression studies through qPCR in peritoneal macrophages derived from C57BL/6. Data is representative of N=2, n=2.

E- Representative confocal images of RAW macrophages exhibiting SIRT1 expression upon *S.* Typhimurium infection at indicated time points post infection. Data is representative of N=2, n=80 (microscopic field).

F-Quantitative representation of the expression profile as depicted in the confocal images (E) in terms of Mean Fluorescence Intensity (MFI). Unpaired two tailed Student's t test was performed to obtain the p values. (*** $p < 0.001$, ** $p < 0.01$)

G- Representative confocal images of RAW macrophages exhibiting SIRT3 expression upon *S.* Typhimurium infection at indicated time points post infection. Data is representative of N=2, n=80 (microscopic field).

H- Quantitative representation of the expression profile as depicted in the confocal images (E) in terms of Mean Fluorescence Intensity (MFI). Unpaired two tailed Student's t test was performed to obtain the p values. (*** $p < 0.001$, ** $p < 0.01$)

Fig.2- Effect of SIRT1 and SIRT3 knockdown in intracellular bacterial proliferation within RAW 264.7 and J774A.1 murine macrophages.

A-Fold Proliferation of *Salmonella* Typhimurium within RAW 264.7 macrophages in transfected and un-transfected conditions. Data is representative of N=3, n=3. Unpaired two tailed Student's t test was performed to obtain the p values. (** $p < 0.01$, * $p < 0.05$)

B- Fold Proliferation of *Salmonella* Typhimurium within infected peritoneal macrophages isolated from adult male C57BL/6 mice upon SIRT1 (EX-527) or SIRT3 (3TYP) inhibitor treatment. Unpaired two tailed Student's t test was performed to obtain the p values. (** $p < 0.01$, * $p < 0.05$)

A

Nanostring

M2 Markers ↑

M1 Markers ↓

B

C

Autofluorescence

Tube_001 - P2

0.02

D

UI SCRAMBLED 2h

Tube_010 - P2

P4

0.06

STM SCRAMBLED 2h

Tube_011 - P2

P4

0.37

STM shSIRT1 2h

Tube_015 - P2

P4

0.05

STM shSIRT3 2h

Tube_019 - P2

P4

0.08

UI SCRAMBLED 16h

Tube_023 - P2

P4

0.10

STM SCRAMBLED 16h

Tube_026 - P2

P4

0.90

STM shSIRT1 16h

Tube_030 - P2

P4

0.32

STM shSIRT3 16h

Tube_034 - P2

P4

0.14

E

%CD206(APC) positive population

ns

*

*

*

*

ns

ns

*

Scrambled UI 2h
Scrambled STM 2h
shSIRT1 STM 2h
shSIRT3 STM 2h
Srambled UI 16h
Scrambled STM 16h
shSIRT1 STM 16h
shSIRT3 STM 16h

Fig.3- *Salmonella* Typhimurium skews the polarization state of the macrophage toward an immunosuppressive M2 state along the course of infection

A-nanoString gene expression profiling data of *S*. Typhimurium infected RAW 264.7 macrophages versus uninfected control data sets at 2hr, 6hr and 16hr time points of infection. Red oval and green oval suggest M2 and M1 markers respectively. Data is representative of N=2,n=1.

B-Flow cytometric analysis of alteration in M1 CD86 positive population in *S*. Typhimurium (STM) infected samples in comparison to uninfected (UI) and Paraformaldehyde Fixed (PFA) bacteria at the indicated time post-infection. Data is representative of N=2, n=3.

C- Quantitative representation of the Flow cytometric analysis depicted in Fig.5, B. Two-way ANOVA and Bonferroni post-t-test was used to obtain p values. (*** $p < 0.001$)

D-Flow cytometric analysis of M2 surface marker CD206 in *S*. Typhimurium (STM) infected SIRT1 or SIRT3 knockdown RAW264.7 macrophages in comparison to the scrambled control at the indicated time post-infection. UI-Uninfected sample. Data is representative of N=3, n=3.

E- Quantitative representation of the Flow cytometric analysis depicted in Fig.5, A. Two-way ANOVA and Bonferroni post-t-test was used to obtain p values. (*** $p < 0.001$)

Fig.4- SIRT1 mediates modulation of immune functions via deacetylation of p65 subunit of NF-κB in *S.*Typhimurium infected macrophages.

A. An immunoblot depicting p65 NF-κB interaction with SIRT1 post immunoprecipitation of SIRT1 in uninfected (UI) or *S.* Typhimurium (STM) infected RAW264.7 macrophages at 16hr post-infection. Data is representative of N=2, n=1

B. Immunoblotting of SIRT1 in SIRT1 knockdown uninfected (UI) or *S.* Typhimurium (STM) infected RAW 264.7 cells at 16hr post-infection to assess the acetylation status of p65 subunit of NF-κB. Data is representative of N=2, n=1.

C-D. Densitometric plot depicting the band intensities of (C) total p65 NF-κB and (D) Acetylated p65 subunit of NF-κ B over total p65 NF-κB in blot B.

E. An immunoblot depicting p65 NF-κB interaction with SIRT1 as well as the p65 NF-κB acetylation status post immunoprecipitation of p65 (IP:p65) or with control IgG (IP:IgG) in uninfected (UI) or *S.* Typhimurium (STM) infected RAW264.7 macrophages upon knockdown with SIRT1 shRNA or scrambled control.

F. Densitometric plot depicting the band intensities of Acetylated p65 over total p65 in blot E.

G. Densitometric plot depicting the band intensities of SIRT1 in blot E.

H. An immunoblot depicting p65 NF-κB interaction with SIRT1 as well as the p65 NF-κB acetylation status post immunoprecipitation of p65 (IP:p65) or with control IgG (IP:IgG) in uninfected (UI) or *S.* Typhimurium (STM) infected RAW264.7 macrophages upon SIRT1 inhibitor (EX-527, 1µM) treatment at 16hr post-infection. UT- Untreated.

I. Densitometric plot depicting the band intensities of Acetylated p65 subunit of NF-κ B over total p65 NF-κB in blot H.

J. Densitometric plot depicting the band intensities of SIRT1 in blot H.

RAW 264.7

A

	SI_2hr	SI_6hr	SI_16hr
Acadl			
PFK2			
PGC-1a			
PPARA			
PPARD			
PPARG			
IDH2			

Fatty acid oxidation ↑

Glycolysis ↓

TCA Cycle ↑

B PPARδ

C RAW

D RAW 264.7

E RAW 264.7

F

RAW264.7

G

Peritoneal Macrophages

Hadha

Fold change/β actin

2.5
2.0
1.5
1.0
0.5
0.0

* *

Untreated UI 16h
EX-527 UI 16h
3TYP UI 16h
Untreated STM 16h
EX-527 STM 16h
3TYP STM 16h

Acox1

Fold change/β actin

2.5
2.0
1.5
1.0
0.5
0.0

*

Untreated UI 16h
EX-527 UI 16h
3TYP UI 16h
Untreated STM 16h
EX-527 STM 16h
3TYP STM 16h

Acadl

Fold change/β actin

2.0
1.5
1.0
0.5
0.0

** **

Untreated UI 16h
EX-527 UI 16h
3TYP UI 16h
Untreated STM 16h
EX-527 STM 16h
3TYP STM 16h

Pdha1

Fold change/β actin

3
2
1
0

* *

Untreated UI 16h
EX-527 UI 16h
3TYP UI 16h
Untreated STM 16h
EX-527 STM 16h
3TYP STM 16h

Pfkl

Fold change/β actin

*** **

Untreated UI 16h
EX-527 UI 16h
3TYP UI 16h
Untreated STM 16h
EX-527 STM 16h
3TYP STM 16h

H

Peritoneal Macrophages

Lactate concentration in ng/ml

** *** *

UT STM 2h
EX-527 STM 2h
3-TYP STM 2h
UT STM 6h
EX-527 STM 6h
3-TYP STM 6h
UT STM 16h
EX-527 STM 16h
3-TYP STM 16h

LIVER

I

Acadl

Fold change/β actin

UI Vehicle control
UI EX-527 (1mg/kg)
UI 3TYP (1mg/kg)
UI SRT1720 (1mg/kg)
STM Vehicle control
STM EX-527 (1mg/kg)
STM 3TYP (1mg/kg)
STM SRT1720 (1mg/kg)

****
****
*

Hadha

Fold change/β actin

UI Vehicle control
UI EX-527 (1mg/kg)
UI 3TYP (1mg/kg)
UI SRT1720 (1mg/kg)
STM Vehicle control
STM EX-527 (1mg/kg)
STM 3TYP (1mg/kg)
STM SRT1720 (1mg/kg)

****
****
**

Acox1

Fold change/β actin

UI Vehicle control
UI EX-527 (1mg/kg)
UI 3TYP (1mg/kg)
UI SRT1720 (1mg/kg)
STM Vehicle control
STM EX-527 (1mg/kg)
STM 3TYP (1mg/kg)
STM SRT1720 (1mg/kg)

****
***
****

Pdha1

Fold change/β actin

UI Vehicle control
UI EX-527 (1mg/kg)
UI 3TYP (1mg/kg)
UI SRT1720 (1mg/kg)
STM Vehicle control
STM EX-527 (1mg/kg)
STM 3TYP (1mg/kg)
STM SRT1720 (1mg/kg)

****
****
****

Pfkl

Fold change/β actin

UI Vehicle control
UI EX-527 (1mg/kg)
UI 3TYP (1mg/kg)
UI SRT1720 (1mg/kg)
STM Vehicle control
STM EX-527 (1mg/kg)
STM 3TYP (1mg/kg)
STM SRT1720 (1mg/kg)

****
*
**

J

Fig.5- *Salmonella* Typhimurium drives the metabolism of the infected macrophage toward fatty acid oxidation

A-Metabolic gene expression data of *S.* Typhimurium infected RAW 264.7 macrophages at 2hr, 6hr and 16hr time points of infection through nanoString. Red oval, green oval and blue oval represent fatty acid oxidizing genes, glycolytic genes and Tricarboxylic acid cycle genes respectively. Data is representative of N=2, n=1.

B-PPARδ qPCR expression data in *S.* Typhimurium infected RAW 264.7 macrophages at the indicated time points of infection. Data is representative of N=3, n=2.

C- Lactate estimation assay of *S.*Typhimurium infected RAW264.7 macrophages at the initial time point of 2hr and at the late time point of 16hr post infection. Data is representative of N=3,n=3. Unpaired two-tailed Student's t test was performed to obtain the p values. Data is representative of N=2, n=2.

D-E-Lactate estimation assay of *S.*Typhimurium infected RAW264.7 macrophages upon SIRT1 (D) or SIRT3 (E) knockdown condition at16h post-infection. Data is representative of N=3,n=3. Unpaired two-tailed Student's t test was performed to obtain the p values. Data is representative of N=2, n=2.

F- qRT PCR mediated gene expression analysis of several host metabolic genes in SIRT1 or SIRT3 knockdown RAW264.7 macrophages at 6 hr post infection. Data is representative of N=2,n=2. Unpaired two-tailed Student's t test was performed to obtain the p values. Data is representative of N=3, n=3.

G-qRT PCR mediated gene expression analysis of several host metabolic genes in SIRT1 or SIRT3 inhibitor-treated (1μM) peritoneal macrophages isolated from 6-8 week old C56BL/6 male mice liver harvested at 5th day post thioglycolate injection. Data is representative of N=2,n=2. Unpaired two-tailed Student's t test was performed to obtain the p values. Data is representative of N=3, n=3.

H- Lactate estimation assay of *S.*Typhimurium infected peritoneal macrophages upon SIRT1 or SIRT3 catalytic inhibition at16h post-infection. Data is representative of N=3,n=3. Unpaired two-tailed Student's t test was performed to obtain the p values. Data is representative of N=2, n=2.

I- qRT PCR mediated gene expression profiling of several host metabolic genes in SIRT1 or SIRT3 inhibitor treated 6-8 week old C56BL/6 male mice liver harvested at 5-day post infection of *S.*Typhimurium (10^7 cfu units/animal). Unpaired two-tailed Student's t test was performed to obtain the p values. Data is representative of N=3, n=3

J- qRT PCR mediated gene expression profiling of several host metabolic genes in SIRT1 or SIRT3 inhibitor treated 6-8 week old C56BL/6 male mice spleen harvested at 5 day post infection of *S.*Typhimurium (10^7 cfu units/animal). Unpaired two-tailed Student's t test was performed to obtain the p values. Data is representative of N=3, n=3.

K. Immunoblotting of host glycolytic (PGK) and TCA cycle (PDHA) proteins in uninfected (UI) or *S.* Typhimurium (STM) infected RAW264.7 macrophages upon SIRT1 enzymatic inhibitor treatment at 16hr post-infection.

L. Densitometric plot depicting the band intensities of proteins in blot K.

M. Immunoblotting of host glycolytic (PGK) and fatty acid oxidation (HADHA, ACOX-1) proteins under SIRT1 and SIRT3 knockdown condition of *S.* Typhimurium infected RAW264.7 macrophages at 16h post-infection.

N-Fatty Acid Oxidation (FAO) Assay of uninfected (UI) and infected (STM) RAW264.7 macrophages under SIRT1 or SIRT3 knockdown or inhibitor treatment. N=2,n=2.

A

RAW264.7

	SI_2hr	SI_6hr	SI_16hr	PI_2hr	PI_6hr	PI_16hr
Inv						
PrgH						
glk						
pfkA						
ptsG						
ssaV						
stfF						

B

fadA

fadB

aceA

aceB

ptsG

C

SPLEEN

D

LIVER

Fig.6- *Salmonella* Typhimurium infection proceeds with increased glycolysis and glucose uptake inside the infected RAW 264.7 macrophages

A-*Salmonella* gene expression profiling data of *S*. Typhimurium infected RAW 264.7 macrophages at 2hr, 6hr and 16hr time points of infection through nanoString. SI- *S*. Typhimurium infected, PI- PFA fixed *S*. Typhimurium infected. SPI-1 genes- *Inv, PrgH* , SPI-2 genes- *ssaV,stfF, glk*- glucokinase, *pfkA*-phosphofrucktose kinase A, *ptsG*- Phosphophenolpyruvate-dependent sugar phosphotransferase system (PTS). Data is representative of N=2, n=1.

B-qRT PCR gene expression profiling of *Salmonella* metabolic genes within infected RAW264.7 macrophages in knockdown condition of either SIRT1 or SIRT3 at 6hr post infection. Data is representative of N=3, n=3.

C- D-qRT PCR gene expression profiling of *Salmonella* metabolic genes within infected female C57BL/6 mice spleen (C) and liver (D) under SIRT1 or SIRT3 inhibitor treatment harvested at 5th day post infection of *S*.Typhimurium (10^7 cfu units/animal). Unpaired two-tailed Student's t test was performed to obtain the p values. Data is representative of N=3, n=3.

Fig.7- SIRT1 inhibition triggers hyperacetylation of glycolytic master regulator HIF-1α within *S.* Typhimurium infected macrophages.

A. An immunoblot depicting HIF-1α interaction with SIRT1 post immunoprecipitation of SIRT1 in uninfected (UI) or *S.* Typhimurium (STM) infected RAW264.7 macrophages at 16hr post-infection. (Derived from same SIRT1 IP blot as in Fig. 4A)

B. Immunoblotting of SIRT1 in SIRT1 knockdown uninfected (UI) or *S.* Typhimurium (STM) infected RAW 264.7 cells at 16hr post-infection to assess the acetylation status of HIF-1α.

C . Densitometric plot depicting the band intensities of Acetylated HIF-1α over total HIF-1α in blot B.

D. An immunoblot depicting HIF-1α interaction with SIRT1 as well as the HIF-1α acetylation status post immunoprecipitation of HIF-1α (IP:HIF-1α or with control IgG (IP:IgG) in uninfected (UI) or *S.* Typhimurium (STM) infected RAW264.7 macrophages upon SIRT1 inhibitor (EX-527, 1µM) treatment at 16hr post-infection. UT-untreated.

E. Densitometric plot depicting the band intensities of Acetylated HIF-1α over total HIF-1α in blot D.

F. Densitometric plot depicting the band intensities of SIRT1 in blot D.

Fig.8- SIRT1 and SIRT3 skews metabolism of *S.* Typhimurium infected macrophages via interaction with PDHA1.

A. An immunoblot depicting PDHA1 interaction with SIRT1 or SIRT3 post immunoprecipitation of PDHA1 in uninfected (UI) or *S.* Typhimurium (STM) infected RAW264.7 macrophages in SIRT3 knockdown condition at 16hr post-infection.

B. Densitometric plot depicting the band intensities of SIRT3 interaction over total input in blot A.

C. An immunoblot depicting PDHA1 interaction with SIRT3 post immunoprecipitation of PDHA1 in uninfected (UI) or *S.* Typhimurium (STM) infected RAW264.7 macrophages at 16hr post-infection under SIRT3 inhibitor (3-TYP, 1μM) treatment at 16hr post-infection. UT-untreated.

D. Densitometric plot depicting the band intensities of Acetylated PDHA1 over total PDHA1 in blot C.

E . Densitometric plot depicting the band intensities of SIRT3 interaction over total input in blot C.

H

EX-527-SIRT1 inhibitor
3TYP-SIRT3 inhibitor
SRT1720-SIRT1 activator

J

Autofluorescence

Vehicle Control STM

K EX-527 (1mg/kg) STM

EX-527 (1mg/kg) STM

SRT 1720 (1mg/kg) STM

L

STM Vehicle Control

Score:1

4X 10X 40X

Slide- 1

Showing minimal/ mild focal inflammatory cell collections in lobular parenchyma

STM SRT1720 (1mg/kg)

Score-1

Slide- 9

Showing minimal/ mild pathology in lobular parenchyma with normal hepatic architecture

STM EX-527(1mg/kg)

Score : 3

C V
N
INF
N

N
INF

SLIDE- 4

Showing multiple (small and big areas) severe acute hepatic necrosis (N), with complete loss of hepatic architecture due to loss of hepatic cells, the necrotic area filled with edema and surrounded by inflammatory cells, hepatocytes with cytoplasmic vacuolation (V), inflammatory cells (INF), and damage of the endothelial lining of central vein (C)

STM 3TYP (1mg/kg)

Score-3

INF
N
INF
HPV
H HPV
V
N

HPV HPV

HPV N

Slide- 7

Showing small multiple necrotic foci (N), severe blood hemorrhage (H) congested hepatic portal veins of liver clogged with clumps and inflammatory cell infiltrates (HPV) vacuolar degeneration (V). Multiple Microgranuloma formation with the aggregation of inflammatory cells (INF)

EX-527-SIRT1 inhibitor
3TYP-SIRT3 inhibitor
SRT1720-SIRT1 activator

Score-0 UI Vehicle Control

4X 10X 40X

normal pathology with normal hepatic architecture

Score-1 UI EX-527(1mg/kg)

Showing minimal/ mild focal inflammatory cell collections in lobular parenchyma

Score-1 UI 3TYP (1mg/kg)

Showing minimal/ mild focal inflammatory cell collections in lobular parenchyma

Score-0 UI SRT1720 (1mg/kg)

normal pathology with normal hepatic architecture

M

Fig.9- Effect of SIRT1 or SIRT3 inhibition on *S*. Typhimurium infected C57BL/6 mice

A-The schematic representation of the experimental strategy for studying the effect of SIRT1 and SIRT3 on the *in vivo* pathogenesis of STM (WT).

B-In-vivo organ burden of *S*. Typhimurium upon SIRT1 or SIRT3 inhibition in C57BL/6 mice at day 5 post infection under specified dosage of inhibitor treatment. Data is representative of N=3, n=3.

C-Survival proportions of *S*. Typhimurium infected C57BL/6 mice upon SIRT1 or SIRT3 inhibitor treatment at a specific dose of inhibitor treatment. Data is representative of N=3, n=3

D- Representation of splenic length of *S*. Typhimurium infected spleen tissue harvested from C57BL/6 mice (males) at day 5 post-infection upon SIRT1 or SIRT3 inhibition at 1mg/kg dosage.

E- Bacterial load in blood at different days post-infection upon SIRT1 or SIRT3 inhibition at 1mg/kg dosage in C57BL/6 mice (males) at day 5 post-infection. Data is representative of N=1, n=1.

F- Bacterial load in blood at day 5 post-infection upon SIRT1 or SIRT3 inhibition at 1mg/kg dosage in C57BL/6 WT mice (males) and gp91phox-/- (males) mice. Data is representative of N=1, n=1.

G- Serum IL-6 levels of *S*. Typhimurium infected C57BL/6 WT mice (males) and gp91phox-/- (males) mice treated with SIRT1(EX-527) or SIRT3 (3TYP) inhibitors or SRT1720 (SIRT1 activator) at 1mg/kg dosage. Data is representative of N=2, n=2.

H-Enumeration of GFP positive bacterial cells through flow cytometry in splenic tissues homogenate harvested from adult male 6-8 week old C56BL/6 mice (subjected to different chemical treatment-Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1 activator SRT1720 treated at a dose of 1mg/kg) at 5th day post *S*. Typhimurium (expressing GFP) infection (10^8 CFU orally). Data is representative of N=2, n=2.

I- Quantitative analysis of the percentage population of splenic cells harboring GFP+ bacterial cells shown in A. Unpaired two-tailed Student's t test was performed.

J- Enumeration of GFP positive bacterial cells through flow cytometry within F4/80 positive splenic macrophages present within splenic tissues homogenate harvested from adult male 6-8 week old C56BL/6 mice (subjected to different chemical treatment-Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1 activator SRT1720 treated aa t dose of 1mg/kg) at 5th day post *S*. Typhimurium infection. Data is representative of N=2, n=2.

K- Quantitative analysis of percentage population of F4/80 positive macrophage cells harboring GFP+ bacteria shown in C. Unpaired two-tailed Student's t test was performed to obtain the p values. (** p < 0.01)

L-Representative image of haematoxylin-eosin stained liver sections to assess the liver tissue architecture upon *Salmonella* infection at 5th days post-infection in different mice cohorts. (UI-Uninfected, STM- *S*. Typhimurium infected, EX-527-SIRT1 inhibitor, 3TYP-SIRT3 inhibitor, SRT1720- SIRT1 activator, Vehicle Control-PBS containing 0.1% DMSO).

M-Graph representing the histopathological scoring of the liver sections depicted in L.

SIRT1 and SIRT3 mediated immune-metabolic switch is crucial for the establishment of *Salmonella* pathogenicity of infection.

SUPPLEMENTARY FIGURES

Fig.S1- *Salmonella* modulates the expression of Sirtuin 1 and Sirtuin 3 at protein level along its course of infection

A-B. Expression studies of SIRT1 (A) and SIRT3 (B) through immunoblotting in RAW 264.7 macrophages. Data is representative of N=3, n=1.

C-D. Quantification of the densitometric plot showing the fold intensities of SIRT1 (C) and SIRT3 (D) protein expression.

Fig.S2- Validation of SIRT1 and SIRT3 knockdown in RAW 264.7 macrophages

A-Knockdown confirmation of SIRT1 at transcript level through qPCR.

B-qPCR expression of SIRT3 in order to validate knockdown efficiency.

C-Immunoblotting of SIRT1 for verification of knockdown. Data is representative of N=3, n=1. (Uns=Un-transfected, Scr=Scrambled, Mix=transfection with equimolar concentration of both SIRT1 shRNA construct C4 and C12)

D-SIRT3 knockdown validation through western blotting. Data is representative of N=3, n=1.(Scr=Scrambled, Mix=transfection with equimolar concentration of both SIRT3 shRNA construct E8 and E12)

E- Densitometric analysis of the immunoblot represented in C.

F- Densitometric analysis of the immunoblot represented in D.

Fig. S3- SIRT1 and SIRT3 modulates cytokine production in *Salmonella* infection scenario

A-Estimation of pro-inflammatory IL-6 cytokine production upon *S.* Typhimurium infection in SIRT1 and SIRT3 knockdown RAW 264.7 macrophages through ELISA. Data is representative of N=2, n=2. Unpaired two tailed Student's t test was performed to obtain the p values. (** p<0.01, * p<0.05).

B- Estimation of anti-inflammatory IL-10 cytokine production upon *S.* Typhimurium infection in SIRT1 and SIRT3 knockdown RAW 264.7 macrophages through ELISA. Data is representative of N=2, n=2. Unpaired two tailed Student's t test was performed to obtain the p values. (*** p < 0.001, * p<0.05)

C-H-Estimation of pro-inflammatory IL-6 cytokine production within uninfected or *S.* Typhimurium infected peritoneal macrophages in presence or absence of SIRT1 or SIRT3 inhibitor treatment . Data is representative of N=2, n=2. Unpaired two tailed Student's t test was performed to obtain the p values. (** p<0.01, * p<0.05)

A

Untransfected STM 16h

Scrambled STM 16h

Uninfected 16h

shSIRT1 STM 16h

shSIRT3 STM 16h

B

Fig.S4- SIRT1 and SIRT3 play a role in maintaining ROS balance within *Salmonella* infected macrophages.

A-DCFDA fluorescence measured through Flow cytometry in shSIRT1, shSIRT3 or scrambled shRNA transfected and un-transfected RAW 264.7 macrophages at 16hr post *S.* Typhimurium infection. Data is representative of N=3, n=2.

B- Quantitative analysis of percentage population of DCFDA positive cells shown in Fig.6, A. Unpaired two-tailed Student's t test was performed. Data is statistically non-significant.

C- Detection of extracellular H_2O_2 production through Phenol Red Hydrogen peroxidase assay at 610nm absorbance. Two-way ANOVA and Bonferroni post-t-test was used to obtain p values. (** $p < 0.01$, * $p < 0.05$, ns-not significant) Data is representative of N=2, n=2.

Fig.S5-Increased ROS burden upon SIRT1 or SIRT3 knockdown gets recruited to the *S.* Typhimurium amounting to its attenuated intracellular proliferation

A. Representative confocal images of SIRT1 knockdown RAW264.7 macrophages exhibiting recruitment of ROS (DCFDA fluorescence) on *S.* Typhimurium at indicated time points post-infection. Data is representative of N=2, n=80 (microscopic field).

B-C. Colocalization coefficient of *S.* Typhimurium with DCFDA in infected SIRT1 knockdown RAW264.7 macrophages at 2hr (B) and 6hr (C) post-infection as depicted in A of Fig.S3. Unpaired two tailed Student's t test was performed to obtain the p values. (*** $p < 0.001$, ** $p<0.01$, * $p<0.05$)

D. Representative confocal images of SIRT3 knockdown RAW264.7 macrophages exhibiting recruitment of ROS (DCFDA fluorescence) on *S.* Typhimurium at indicated time points post-infection. Data is representative of N=2, n=80 (microscopic field).

E-F. Colocalization coefficient of *S.* Typhimurium with DCFDA in infected SIRT3 knockdown RAW264.7 macrophages at 2hr (B) and 6hr (C) post-infection as depicted in A of Fig.S3. Unpaired two tailed Student's t test was performed to obtain the p values. (*** $p < 0.001$, ** $p<0.01$, * $p<0.05$)

Fig.S6 NanoString gene expression profiling data of *S.* Typhimurium infected SIRT1 and SIRT3 knockdown RAW 264.7 macrophages

A-B. nanoString gene expression profiling data of various immunological genes governing M1 and M2 polarization within *S.* Typhimurium infected SIRT1 and SIRT3 knockdown RAW 264.7 macrophages at 6hr (A) and 16hr (B) time points post-infection. Red oval and green oval suggest M2 and M1 markers respectively. Red-High Normalized expression, Blue-Low Normalized expression. Data is representative of N=1, n=1.

C. Metabolic gene expression data of *S.* Typhimurium infected SIRT1 or SIRT3 knockdown RAW 264.7 macrophages at 6hr time points of infection through nanoString. Red-High Normalized expression, Blue-Low Normalized expression. Data is representative of N=1, n=1.

Fig.S7- Effect of SIRT1 and SIRT3 knockdown or inhibition in intracellular bacterial proliferation within RAW 264.7 macrophages or peritoneal macrophages in presence of ROS scavenger N-Acetyl Cysteine (NAC).

A-Fold Proliferation of *Salmonella* Typhimurium within RAW 264.7 macrophages in transfected and un-transfected conditions in presence or absence of NAC (1mM). Data are representative of N=3, n=3. Unpaired two tailed Student's t test was performed to obtain the p values. (* $p<0.05$)

B-Fold Proliferation of *Salmonella* Typhimurium within peritoneal macrophages isolated from 6-8 week old adult male C57BL/6 mice upon SIRT1 (EX-527, 1µM) or SIRT3 (3TYP, 1µM) inhibitor treatment in presence or absence of NAC (1mM). Data are representative of N=3, n=3. Unpaired two tailed Student's t test was performed to obtain the p values. (* $p<0.05$)

A

Peritoneal Macrophages

HADHA
0.29 0.45 0.95 2.37 0.84 0.55

PGK
0.18 0.01 0.01 0.09 1.36 0.4

PFK
0.13 0.05 0.05 0.17 0.35 0.51

β actin

ACOX-1
0.28 0.39 0.52 1.85 0.69 0.38

PDHA1
0.07 0.19 0.41 1.91 0.22 0.20

16h post-infection

Fig.S8 Immunoblotting of host glycolytic (PGK,PFK) and fatty acid oxidation (HADHA, ACOX-1) proteins under SIRT1 (EX-527) and SIRT3 (3TYP) catalytic inhibitor treated condition in *S.* Typhimurium infected peritoneal macrophages (isolated from thioglycolate treated adult male C57BL/6 mice) at 16h post-infection. Data is representative of N=2, n=1.

Fig.S9- SIRT1 and SIRT3 inhibition led to enhanced ROS production within *Salmonella*-infected mice liver and spleen tissues.

A-DCFDA fluorescence measured through flow cytometry in liver tissues homogenate harvested from adult male 6-8 week old C56BL/6 mice (subjected to different chemical treatment-Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1 activator SRT1720 treated at dose of 1mg/kg) at 5th day post *S*. Typhimurium infection (10^8 CFU orally) . Data is representative of N=2, n=2.

B- Quantitative analysis of percentage population of cells showing DCFDA positive staining shown in A. Unpaired two-tailed Student's t test was performed. Data is statistically non-significant.

C- DCFDA fluorescence measured through Flow cytometry in splenic tissues homogenate harvested from adult male 6-8 week old C56BL/6 mice (subjected to different chemical treatment-Vehicle treated or SIRT1 (EX-527) or SIRT3 (3-TYP) inhibitor or SIRT1 activator SRT1720 treated at dose of 1mg/kg) at 5th day post *S*. Typhimurium infection. Data is representative of N=2, n=2.

D- Quantitative analysis of percentage population of cells showing DCFDA positive staining shown in C. Unpaired two-tailed Student's t test was performed to obtain the p values. (** $p < 0.01$)

Graphical Abstract

HIGHLIGHTS

- SIRT1 and SIRT3 promote an immunosuppressive state of the *S.* Typhimurium-infected macrophages.
- SIRT1 together with SIRT3 subverts the host glycolytic switch and concomitantly promotes fatty acid oxidation of the infected macrophages.
- The host immuno-metabolic switch governs intracellular replication of the pathogen within the infected murine macrophages.
- SIRT1 or SIRT3 inhibitor treatment triggered increased bacterial dissemination in *in vivo* mouse model of infection owing to increased inflammatory response.