Transcription bodies regulate gene expression by sequestering CDK9

Martino Ugolini[1,2], Ksenia Kuznetsova[2], Haruka Oda[3], Hiroshi Kimura[3], Nadine L. Vastenhouw[1,2]*

[1]University of Lausanne (UNIL), Center for Integrative Genomics (CIG); Lausanne, Switzerland. [2]Max Planck Institute of Molecular Cell Biology and Genetics (MPI-CBG); Dresden, Germany. [3]Cell Biology Center, Institute of Innovative Research, Tokyo Institute of Technology; Yokohama 226-8503, Japan. *Corresponding author: nadine.vastenhouw@unil.ch

The localization of transcriptional activity in specialized transcription bodies is a hallmark of gene expression in eukaryotic cells. It remains unclear, however, if and how they affect gene expression. Here, we disrupted the formation of two prominent endogenous transcription bodies that mark the onset of zygotic transcription in zebrafish embryos and analysed the effect on gene expression using enriched SLAM-Seq and live-cell imaging. We find that the disruption of transcription bodies results in downregulation of hundreds of genes, providing experimental support for a model in which transcription bodies increase the efficiency of transcription. We also find that a significant number of genes are upregulated, counter to the suggested stimulatory effect of transcription bodies. These upregulated genes have accessible chromatin and are poised to be transcribed in the presence of the two transcription bodies, but they do not go into elongation. Live-cell imaging shows that the disruption of the two large transcription bodies enables these poised genes to be transcribed in ectopic transcription bodies, suggesting that the large transcription bodies sequester a pause release factor. Supporting this hypothesis, we find that CDK9, the kinase that releases paused

polymerase II, is highly enriched in the two large transcription bodies. Importantly, overexpression of CDK9 in wild type embryos results in the formation of ectopic transcription bodies and thus phenocopies the removal of the two large transcription bodies. Taken together, our results show that transcription bodies regulate transcription genome-wide: the accumulation of transcriptional machinery creates a favourable environment for transcription locally, while depriving genes elsewhere in the nucleus from the same machinery.

RNA polymerase II (RNAPII) and transcription factors are often concentrated in specialized transcription bodies [1–10], and it has been suggested that these can bring together multiple genes [11–14]. This has led to the hypothesis that transcription bodies increase the efficiency of transcription by promoting the biomolecular interactions underlying gene expression [15–17]. Analysis of gene expression in artificially induced condensates supports this model [18]. To understand how transcription bodies affect gene expression *in vivo*, however, it is necessary to analyse endogenous transcription bodies. This has been difficult because transcription bodies are often small, short-lived, and numerous. Here, we use the onset of transcription in zebrafish embryos to overcome this problem. In zebrafish embryos, development is initially driven by maternally loaded RNA and protein (Fig. 1a) [19]. Transcription gradually begins during a process called zygotic genome activation (ZGA). Before transcriptional activity can be seen throughout the nucleus, it is confined to two micron-sized transcription bodies (Fig. 1b) [9,20–23], nucleated by the *mir430* locus, which contains multiple copies of the *mir430* gene [21,24]. These bodies are enriched for RNAPII Serine 5 and Serine 2 phosphorylation (Ser5P, Ser2P), which mark the initiating and elongating form of RNAPII, respectively, as well as the transcription factors Nanog and Sox19b [9,20–22].

To investigate the role of transcription bodies in transcription, we generated a fish line in which

the *mir430* locus is deleted [9], resulting in the specific disruption of the two *mir430* transcription

bodies, as observed by the loss of Ser2P signal (Fig. 1c). *miR430* is required for downregulating

maternally loaded transcripts, and its absence results in severe developmental phenotypes [24,25].

Thus, effects on gene expression and development in the *mir430* mutant may be caused by the

loss of the *mir430* transcription bodies, or the loss of the mature microRNA MiR430. To

uncouple these two components and allow us to focus on the first, we supplemented mutant

embryos with the mature form of microRNA MiR430 (Fig. 1d) [24], which restored degradation

of mRNAs containing MiR430 binding sites (Fig. 1e and Extended Data Fig. 1). This rescued

the delay in epiboly (Fig. 1f-g) and the aberrant expression pattern of *goosecoid* (Extended

Data Fig. 2), which are characteristic for the loss of MiR430 [24,25]. Embryos supplemented with

mature microRNAs appeared normal, with the appearance of the Kupffer's vesicle (Fig. 1h),

and overall developmental progression (Fig. 1i and Extended Data Fig. 3) being

indistinguishable from wild type (WT) and heterozygous siblings, and embryos developed into

fertile fish. Thus, the disruption of two large transcription bodies does not result in obvious

developmental defects, and we can use it to analyse the effect of transcription bodies on gene

expression.

To assess the effect of the specific disruption of *mir430* transcription bodies on transcriptional

activity, we developed eSLAM-Seq, which uses a combination of protocols to label, enrich,

and detect nascent transcripts (Extended Data Fig. 4 and Methods), because the large amounts

of maternally loaded RNA in early embryos (Fig. 1a) masks nascent RNAs in total RNA

sequencing approaches. We used eSLAM-Seq to compare gene expression between WT and

mir430$^{-/-}$ + inj MiR430 embryos at the 256-cell stage, when transcriptional activity is largely

restricted to the *mir430* transcription bodies in WT embryos (Fig. 1b). We observed a

significant number of genes that were downregulated in the absence of the *mir430* transcription

bodies (242) and a higher number of genes that were upregulated (716) (Fig. 2a-b and Extended

Data Table 1). Plotting the genomic distribution of genes that were up- and downregulated

showed that both classes of genes are distributed across the genome (Fig. 2c). We conclude

that the absence of two large transcription bodies has a widespread effect on transcription.

The identification of hundreds of up- and downregulated genes in *mir430*$^{-/-}$ + inj MiR430 *vs.*

WT embryos at the 256-cell stage contrasts with the observation that these embryos develop

normally (Fig. 1). To reconcile these two observations, we generated eSLAM-Seq data for WT

and *mir430*$^{-/-}$ + inj MiR430 embryos at the 1024-cell, Oblong, and Sphere stage and compared

it with the data obtained at the 256-cell stage. The expression levels of genes that are

misregulated in *mir430*$^{-/-}$ + inj MiR430 embryos at the 256-cell stage (with a median of 3.5

$\log_2$ fold difference with WT for the upregulated genes, and 2.9 $\log_2$ fold for the downregulated

genes; Fig. 2d) gradually recover during consecutive developmental stages, reaching WT

expression levels at Sphere stage (with a median of 0.03 $\log_2$ fold difference with WT for the

upregulated genes, and 0.07 $\log_2$ fold for the downregulated genes; Fig. 2d). Looking at all

genes, we find that overall fewer genes are misregulated at later stages, from 716 up and 242

down at 256-cell to 51 up and 30 down at Sphere stage (Extended Data Fig. 5). We conclude

that in the absence of *mir430* transcription bodies, gene expression is misregulated, but that

this effect is short-lived as expression levels return to WT levels in a timeframe of a couple of

cell cycles. This likely explains the lack of obvious developmental defects in *mir430*$^{-/-}$ + inj

MiR430 embryos.

We then further analysed the genes that are misregulated at the 256-cell stage. The

downregulation of genes suggests that transcription bodies facilitate transcription of genes

other than *mir430* itself, consistent with work in which transcribed genes were shown to

colocalize [12–14,26], as well as the proposed role of transcription bodies in increasing

transcriptional efficiency [18,27,28]. We hypothesize that the genes that are downregulated in

mir430[-/-] + inj MiR430 embryos localize to the *mir430* transcription bodies in WT embryos

and are positively affected by this localization. The observation that downregulated genes

return to WT expression levels over time suggests that transcription bodies that form during

later stages in *mir430*[-/-] + inj MiR430 embryos (Extended Data Fig. 6) may compensate for the

loss of *mir430* transcription bodies. Here, we focus on the upregulated genes. Their high

number suggest that the *mir430* transcription bodies sequester components of the

transcriptional machinery, preventing genes elsewhere in the nucleus from being transcribed.

To investigate this sequestration model, we characterized the genes that are upregulated in

response to *mir430* transcription body removal. We first asked when these genes are induced

in WT embryos. We analysed their expression at the 1024-cell, Oblong, and Sphere stages, and

compared this to their expression at the 256-cell stage. We found that most of the genes (81%)

are activated during genome activation in WT embryos (Fig. 3a), indicating that loss of the

mir430 transcription bodies specifically activates the expression of genes that are about to be

activated, with 62% (446/716) being advanced by 30 minutes (at the 1024-cell stage in WT,

Fig. 3a). When focusing on the genes that are advanced by 30 minutes, we observe a good

correlation between their degree of upregulation between *mir430*[-/-] + inj MiR430 and WT at

256-cell stage, and between 1024-cell and 256-cell stage in WT (Fig. 3b and Extended Data

Fig. 7). This further supports the notion that the effect on transcription activation is specific.

We then investigated the chromatin accessibility state of the genes that are upregulated. We

analysed an ATAC-Seq dataset we had previously generated in WT embryos [29] and found that

the transcription start sites of genes that are upregulated in the *mir430*[-/-] + inj MiR430 are

highly accessible in WT embryos (Fig. 3c), being as accessible as those of the most highly

expressed genes (Fig. 3c-d), even though they are hardly expressed (Fig. 3e). This suggests

that potentially sequestered factor(s) function downstream of transcription-factor binding and

chromatin opening. Consistent with this, motif analysis of the promoters of upregulated genes

did not identify transcription factor motifs that were both sufficiently enriched compared to

unexpressed genes and present in a sufficiently high fraction of the upregulated genes to

explain our observations (Extended Data Fig. 8). Even motifs of Nanog, Pou5f3 and Sox19b,

transcription factors that have been shown to be required for the activation of transcription

during ZGA in zebrafish [9,29–32] did not fulfil these two requirements. Taken together, our results

suggest that the presence of the *mir430* transcription bodies prevents a significant number of

genes that are poised for activation from going into elongation.

This reminded us of the observation that WT embryos from the 256-cell stage onward, in

addition to the *mir430* transcription bodies in which both transcription initiation and elongation

can be observed, display transcription bodies that are positive for transcription initiation, but

not elongation [9]. We were previously unable to explain this observation but now hypothesized

that the *mir430* transcription bodies might sequester a pause release factor, causing genes in

these transcription bodies to be arrested at the initiation phase. If this were true, these

transcription bodies would be predicted to acquire elongation signals in nuclei of *mir430*[-/-] +

inj MiR430 embryos. To investigate this, we injected 1-cell WT and *mir430*[-/-] + inj MiR430

embryos with fluorescently labelled antigen-binding fragments (Fabs) recognizing RNAPII-

Ser5P (transcription initiation) and RNAPII-Ser2P (transcription elongation), as described

before [9,22,23]. Consistent with our hypothesis, we saw more Ser2P-positive transcription bodies

in *mir430*[-/-] + inj MiR430 than in WT embryos during the 512- and 1024-cell stages (Fig. 3f,

and Extended Data Fig. 9-10). Quantification of initiation-only, initiation/elongation, and

elongation-only transcription bodies confirmed that more transcription bodies go into

productive elongation in *mir430*$^{-/-}$ + inj MiR430 than in WT embryos (with a maximum of 15%

elongation-positive transcription bodies in WT versus 33% in *mir430*$^{-/-}$ + inj MiR430 at the

256-cell stage, and a comparable difference at the 1024-cell stage; Fig. 3g). Thus, the removal

of the *mir430* transcription bodies causes transcription bodies that are arrested in the initiation

phase to transition to the transcription elongation phase.

Our data suggests that transcription bodies sequester a pause release factor. Pause release is

regulated by the pTEFb complex of which CDK9 is the catalytic subunit. We visualized CDK9

in WT embryos by immunofluorescence and found that it is strongly enriched in the two *mir430*

transcription bodies (Fig 4a). If the amount of CDK9 in the nucleus were limiting and its

sequestration in *mir430* transcription bodies would hamper transcription elongation elsewhere

in the nucleus, it would be predicted that overexpression of CDK9 in WT embryos would

induce the formation of ectopic transcription bodies. To directly test this, we compared the

occurrence of ectopic transcription bodies between WT embryos with and without injected

cdk9 mRNA. We visualized RNAPII Ser2P to detect transcription bodies. To distinguish

transcription bodies that are nucleated by the *mir430* locus from ectopic transcription bodies,

we also visualized *miR430* transcripts using Morpholino Visualization of Expression (MOVIE

[21]). At the 128-cell stage, we detected two prominent *mir430* transcription bodies in nuclei of

WT embryos, as well as in WT embryos that were injected with *cdk9* mRNA (Fig. 4b and

Extended Data Fig. 11). In the latter, however, we detected additional transcription bodies.

Because these lack *miR430* MOVIE signal and we do not detect them in uninjected WT

embryos, we consider these to be ectopic transcription bodies. The ectopic bodies are small

and limited in number at the 128- and 256-cell stage, and more prominent at the 512- and 1024-

cell stage (Fig. 4b and Extended Data Fig. 11-14). The abundance of ectopic transcription

bodies in WT embryos in which CDK9 is overexpressed, phenocopies the abundance of ectopic

transcription bodies seen in *mir430^-/-* + inj MiR430 embryos. The quantification of ectopic

transcription shows that at all four examined stages, ectopic transcription bodies are

significantly more abundant in nuclei in which CDK9 is overexpressed (WT + inj *cdk9*

mRNA), or in which the large *mir430* transcription bodies are disrupted (*mir430^-/-* + inj

MiR430), than in WT embryos (Fig. 4c). This supports our hypothesis that CDK9 is

sequestered in *mir430* transcription bodies, limiting CDK9 availability elsewhere in the

nucleus.

Taken together, we show that the removal of the *mir430* transcription bodies results in a

genome-wide misregulation of gene expression. Many genes are downregulated, consistent

with a model where the bodies facilitate transcription by promoting the biomolecular

interactions underlying gene expression [3,16,18]. Unexpectedly, we also observed many genes

that are upregulated upon disruption of the *mir430* transcription bodies. We show that

transcription bodies sequester the pause release factor CDK9, preventing genes that are not

present in the transcription bodies from being transcribed (Extended Data Fig. 15). Factor

sequestration regulating biological processes has been described before [33,34] and a recent study

showed that the loss of the transcription factor Zelda in *Drosophila* embryos caused RNAPII

to relocate from Zelda-containing transcription bodies to the histone locus body [35]. More

generally, work in mouse embryos has shown that CDK9 abundance is an important factor in

the timing of zygotic genome activation[36]. Our work shows that endogenous transcription

bodies play an important role in controlling the availability of transcriptional machinery in the

nucleus.

References (1-36 used in main text)

1. Jackson, D. A., Hassan, A. B., Errington, R. J. & Cook, P. R. Visualization of focal sites of transcription within human nuclei. *EMBO J.* **12**, 1059–1065 (1993).

2. Ghamari, A. *et al.* In vivo live imaging of RNA polymerase II transcription factories in primary cells. *Genes Dev.* **27**, 767–777 (2013).

3. Tsai, A. *et al.* Nuclear microenvironments modulate transcription from low-affinity enhancers. *Elife* **6**, 1–18 (2017).

4. Cho, W. K. *et al.* Mediator and RNA polymerase II clusters associate in transcription-dependent condensates. *Science* **361**, 412–415 (2018).

5. Sabari, B. R. *et al.* Coactivator condensation at super-enhancers links phase separation and gene control. *Science* **361**, eaar3958 (2018).

6. Chong, S. *et al.* Imaging dynamic and selective low-complexity domain interactions that control gene transcription. *Science* **361**, eaar2555 (2018).

7. Boija, A. *et al.* Transcription Factors Activate Genes through the Phase-Separation Capacity of Their Activation Domains. *Cell* **175**, 1842–1855 (2018).

8. Zamudio, A. V. *et al.* Mediator Condensates Localize Signaling Factors to Key Cell Identity Genes. *Mol. Cell* **76**, 753–766 (2019).

9. Kuznetsova, K. *et al.* Nanog organizes transcription bodies. *Curr. Biol.* **33**, 1–10 (2022).

10. Cisse, I. I. *et al.* Real-time dynamics of RNA polymerase II clustering in live human cells. *Science* **341**, 664–667 (2013).

11. Iborra, F. J., Pombo, A., Jackson, D. A. & Cook, P. R. Active RNA polymerases are localized within discrete transcription 'factories' in human nuclei. *J. Cell Sci.* **109**, 1427–1436 (1996).

12. Osborne, C. S. *et al.* Active genes dynamically colocalize to shared sites of ongoing

transcription. *Nat. Genet.* **36**, 1065–1071 (2004).

13. Schoenfelder, S. *et al.* Preferential associations between co-regulated genes reveal a

transcriptional interactome in erythroid cells. *Nat. Genet.* **42**, 53–61 (2010).

14. Fanucchi, S., Shibayama, Y., Burd, S. & Weinberg, M. S. Chromosomal Contact

Permits Transcription between Coregulated Genes. *Cell* **155**, 606–620 (2013).

15. Sutherland, H. & Bickmore, W. A. Transcription factories: Gene expression in unions?

Nat. Rev. Genet. **10**, 457–466 (2009).

16. Hnisz, D., Shrinivas, K., Young, R. A., Chakraborty, A. K. & Sharp, P. A. A Phase

Separation Model for Transcriptional Control. *Cell* **169**, 13–23 (2017).

17. Tsai, A., Galupa, R. & Crocker, J. Robust and efficient gene regulation through

localized nuclear microenvironments. *Development* **147**, dev161430 (2020).

18. Wei, M. T. *et al.* Nucleated transcriptional condensates amplify gene expression. *Nat.*

Cell Biol. **22**, 1187–1196 (2020).

19. Vastenhouw, N. L., Cao, W. X. & Lipshitz, H. D. The maternal-to-zygotic transition

revisited. *Dev.* **146**, dev161471 (2019).

20. Chan, S. H. *et al.* Brd4 and P300 Confer Transcriptional Competency during Zygotic

Genome Activation. *Dev. Cell* **49**, 867–881 (2019).

21. Hadzhiev, Y. *et al.* A cell cycle-coordinated Polymerase II transcription compartment

encompasses gene expression before global genome activation. *Nat. Commun.* **10**, 691

(2019).

22. Sato, Y. *et al.* Histone H3K27 acetylation precedes active transcription during

zebrafish zygotic genome activation as revealed by live-cell analysis. *Dev.* **146**,

dev179127 (2019).

23. Hilbert, L. *et al.* Transcription organizes euchromatin via microphase separation. *Nat.*

Commun. **12**, 1360 (2021).

24.   Giraldez, A. J. *et al.* MicroRNAs regulate brain morphogenesis in zebrafish. *Science*

308, 833–838 (2005).

25.   Liu, Y. *et al.* Genetic Deletion of miR-430 Disrupts Maternal-Zygotic Transition and

Embryonic Body Plan. *Front. Genet.* **11**, 853 (2020).

26.   Monahan, K., Horta, A. & Lomvardas, S. LHX2- and LDB1-mediated trans

interactions regulate olfactory receptor choice. *Nature* **565**, 448–453 (2019).

27.   Ma, L. *et al.* Co-condensation between transcription factor and coactivator p300

modulates transcriptional bursting kinetics. *Mol. Cell* **81**, 1682–1697 (2021).

28.   Sabari, B. R. Biomolecular Condensates and Gene Activation in Development and

Disease. *Dev. Cell* **55**, 84–96 (2020).

29.   Pálfy, M., Schulze, G., Valen, E. & Vastenhouw, N. L. Chromatin accessibility

established by Pou5f3, Sox19b and Nanog primes genes for activity during zebrafish

genome activation. *PLoS Genet.* **16**, e1008546 (2020).

30.   Lee, M. T. *et al.* Nanog, Pou5f1 and SoxB1 activate zygotic gene expression during

the maternal-to-zygotic transition. *Nature* **503**, 360–364 (2013).

31.   Leichsenring, M., Maes, J., Mošsner, R., Driever, W. & Onichtchouk, D. Pou5f1

transcription factor controls zygotic gene activation in vertebrates. *Science* **341**, 1005–

1009 (2013).

32.   Miao, L. *et al.* The landscape of pioneer factor activity reveals the mechanisms of

chromatin reprogramming and genome activation. *Mol. Cell* **82**, 986–1002 (2022).

33.   Klosin, A. *et al.* Phase separation provides a mechanism to reduce noise in cells.

Science **367**, 464–468 (2020).

34.   Harigaya, Y. *et al.* Selective elimination of messenger RNA prevents an incidence of

untimely meiosis. *Nat. 2006 4427098* **442**, 45–50 (2006).

35.   Huang, S. K., Whitney, P. H., Dutta, S., Shvartsman, S. Y. & Rushlow, C. A. Spatial

organization of transcribing loci during early genome activation in Drosophila. *Curr.*

Biol. **31**, 5102–5110 (2021).

36.   Oqani, R. K., Kim, H. R., Diao, Y. F., Park, C. S. & Jin, D. I. The CDK9/Cyclin T1

subunits of P-TEFb in mouse oocytes and preimplantation embryos: A possible role in

embryonic genome activation. *BMC Dev. Biol.* **11**, 1–9 (2011).

37.   Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. & Schilling, T. F. Stages

of embryonic development of the zebrafish. *Dev. Dyn.* **203**, 253–310 (1995).

38.   Truett, G. E. *et al.* Preparation of PCR-quality mouse genomic DNA with hot sodium

hydroxide and tris (HotSHOT). *Biotechniques* **29**, 52–54 (2000).

39.   Hayashi-Takanaka, Y. *et al.* Tracking epigenetic histone modifications in single cells

using Fab-based live endogenous modification labeling. *Nucleic Acids Res.* **39**, 6475–

6488 (2011).

40.   Stasevich, T. J. *et al.* Regulation of RNA polymerase II activation by histone

acetylation in single living cells. *Nature* **516**, 272–275 (2014).

41.   Kimura, H. & Yamagata, K. Visualization of epigenetic modifications in

preimplantation embryos. *Methods Mol. Biol.* **1222**, 127–147 (2015).

42.   Thisse, C. & Thisse, B. High-resolution in situ hybridization to whole-mount zebrafish

embryos. *Nat. Protoc.* **3**, 59–69 (2008).

43.   Heyn, P. *et al.* The earliest transcribed zygotic genes are short, newly evolved, and

different across species. *Cell Rep.* **6**, 285–292 (2014).

44.   Herzog, V. A. *et al.* Thiol-linked alkylation of RNA to assess expression dynamics.

Nat. Methods **14**, 1198–1204 (2017).

45.   Holler, K. *et al.* Spatio-temporal mRNA tracking in the early zebrafish embryo. *Nat.*

Commun. **12**, 3358 (2021).

46.   Bhat, P. *et al.* SLAMseq resolves the kinetics of maternal and zygotic gene expression

in early zebrafish embryogenesis. *bioRxiv* (2022) doi:10.1101/2022.06.01.494399.

47.   Schofield, J. A., Duffy, E. E., Kiefer, L., Sullivan, M. C. & Simon, M. D. TimeLapse-

seq: Adding a temporal dimension to RNA sequencing through nucleoside recoding.

Nat. Methods **15**, 221–225 (2018).

48.   Reichholf, B. *et al.* Time-Resolved Small RNA Sequencing Unravels the Molecular

Principles of MicroRNA Homeostasis. *Mol. Cell* **75**, 756–768 (2019).

49.   Duffy, E. E. *et al.* Tracking Distinct RNA Populations Using Efficient and Reversible

Covalent Chemistry. *Mol. Cell* **59**, 858–866 (2015).

50.   Duffy, E. E., Canzio, D., Maniatis, T. & Simon, M. D. Solid phase chemistry to

covalently and reversibly capture thiolated RNA. *Nucleic Acids Res.* **46**, 6996–7005

(2018).

51.   Sun, W. & Chen, W. Metabolic Labeling of Newly Synthesized RNA with 4sU to in

Parallel Assess RNA Transcription and Decay. in *Methods in molecular biology* vol.

1720 25–34 (Humana Press, New York, NY, 2018).

52.   Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

53.   Zhang, Y., Park, C., Bennett, C., Thornton, M. & Kim, D. Rapid and accurate

alignment of nucleotide conversion sequencing reads with HISAT-3N. *Genome Res.*

31, 1290–1295 (2021).

54.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,

2078–2079 (2009).

55.   Broad Institute. Picard. http://broadinstitute.github.io/picard/ (2022).

56.   Boileau, E., Altmüller, J., Naarmann-de Vries, I. S. & Dieterich, C. A comparison of

metabolic labeling and statistical methods to infer genome-wide dynamics of RNA

turnover. *Brief. Bioinform.* **22**, bbab219 (2021).

57. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

58. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol.* **15**, 1–21 (2014).

59. Bailey, T. L. & Grant, C. E. SEA: Simple Enrichment Analysis of motifs. *bioRxiv*

(2021) doi:10.1101/2021.08.23.457422.

60. Ramírez, F. *et al.* deepTools2: a next generation web server for deep-sequencing data

analysis. *Nucleic Acids Res.* **44**, W160–W165 (2016).

61. R Core Team. R: A Language and Environment for Statistical Computing. (2014).

62. Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*. (Springer International

Publishing, 2016). doi:10.1007/978-3-319-24277-4.

63. Yin, T., Cook, D. & Lawrence, M. ggbio: an R package for extending the grammar of

graphics for genomic data. *Genome Biol.* **13**, 1–14 (2012).

64. Herrgen, L., Schröter, C., Bajard, L. & Oates, A. C. Multiple embryo time-lapse

imaging of zebrafish development. *Methods Mol. Biol.* **546**, 243–254 (2009).

65. Boothe, T. *et al.* A tunable refractive index matching medium for live imaging cells,

tissues and model organisms. *Elife* **6**, e27240 (2017).

66. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

Methods **9**, 676–682 (2012).

67. Katrukha, E. ComDet. https://github.com/ekatrukha/ComDet (2022).

Acknowledgements

We thank the members of the Vastenhouw lab for technical support, helpful feedback, and

stimulating discussions. We thank the Pauli and Ameres labs for technical advice on SLAM-

Seq. We thank Maciej Kerlin, Aleksandar Vještica, and Edlyn Wu for comments on the

manuscript, and the following facilities and services for their support: MPI-CBG - fish facility,

light microscopy facility, and scientific computing, UNIL - genomic technologies facility,

cellular imaging facility and the bioinformatics competence center. M.U. was supported by the

Boehringer Ingelheim Fonds. Research in N.L.V's laboratory was supported by the Max

Planck Society, the University of Lausanne, a European Research Council Consolidator Grant

(101003023), the Volkswagen Foundation (94773), and the German Research Foundation (VA

1209/2-1). Research in H.K.'s laboratory was supported by the Japan Society for the Promotion

of Science KAKENHI (JP18H05527, JP21H04764 and JP20K06484).

Author contributions

Conceptualization: M.U., N.L.V. Methodology: M.U., K.K., H.U. Investigation: M.U., K.K.

Data analysis: M.U. Writing – original draft: M.U. and N.L.V. Writing – reviewing & editing:

all authors. Funding acquisition: M.U., N.L.V., H.K. Supervision: N.L.V.

Competing interests

Authors declare that they have no competing interests.

Data and materials availability

Raw sequencing data will be uploaded on Gene Expression Omnibus (GEO) prior to

publication. Raw imaging data are available upon request. All other data is available in the

main text or the extended data.

Figure 1

Figure 1. Disruption of *mir430* transcription bodies does not substantially impact development. a. Schematic representation of the maternal-to-zygotic transition in zebrafish embryos. Hpf, hours post fertilization. **b.** Visualization of elongating RNAPII (Ser2P) with antigen-binding fragments (Fabs) in WT embryos. Shown are representative micrographs of individual nuclei, extracted from a spinning disk confocal microscopy time lapse. **c.** Visualization of elongating RNAPII (Ser2P) with antigen-binding fragments (Fabs) in WT and *mir430*[-/-] embryos at the 128-cell stage. Shown are representative micrographs of individual

nuclei, extracted from a spinning disk confocal microscopy time lapse. **d.** Schematic

representation of a nucleus in WT, *mir430⁻ᐟ⁻* and *mir430⁻ᐟ⁻* + inj MiR430 embryos. **e.** Rescue of

miR430 activity as assessed by comparing the expression of eGFP encoded by an mRNA with

three perfect target sites for *miR430,* and RFP encoded by an mRNA without such sites, in

different genotypes at 24hpf. Normalized eGFP signal in WT, *mir430⁺ᐟ⁻* and *mir430⁻ᐟ⁻* embryos

without (left) and with (right) injected MiR430. N=3, $27 \leq n \leq 74$. See Extended Data Fig. 1 for

images. **f.** Rescue of *miR430* activity as assessed by epiboly progression. Shown are

representative micrographs of embryos at late epiboly stage in different genotypes. The

misregulation of yolk internalization in *mir430⁻ᐟ⁻* embryos is indicated. **g.** Time at which

epiboly is completed in different genotypes without (left), and with (right) injected MiR430

RNA. N=3, $19 \leq n \leq 53$. **h.** Time at which Kupffer's vesicle appears in different genotypes

without (left), and with (right) injected MiR430. N=3, $14 \leq n \leq 44$. **i.** Larvae at 48 hours post

fertilization for different genotypes. Representative micrographs are shown. The malformation

of trunk morphology and eye, the development of heart oedema, and the appearance of blisters

at the tail tip in *mir430⁻ᐟ⁻* embryos are indicated.

Figure 2

Figure 2. Disruption of *mir430* transcription bodies causes widespread misregulation of gene expression. a. Volcano plot showing up- and downregulated genes in *mir430⁻/⁻* + inj MiR430 *vs.* WT embryos at 256-cell stage. Genes whose coverage is shown in Fig. 2b are shown in black. **b.** Coverage plot of *irx7* (upregulated) and *tmpob* (downregulated) in WT and *mir430⁻/⁻* + inj MiR430. The single strata visualize the labelling degree of the reads. **c.** Distribution of up- and downregulated genes across the genome. The *mir430* locus on chromosome 4 is shown in green and labelled by an asterisk. **d.** Difference in average gene expression between *mir430⁻/⁻* + inj MiR430 and WT embryos across stages for those genes that were identified to be differentially expressed at 256-cell stage.

Figure 3

Figure 3. Loss of *mir430* transcription bodies causes premature gene activation. a. Time

of induction in WT embryos for the 716 genes that are upregulated in *mir430*$^{-/-}$ + inj MiR430

vs. WT embryos (at 256-cell stage). Expression at 256-cell is used as a reference. The genes

are split in four groups based on when they are induced. The percentage of genes in each group

is indicated. **b.** Scatterplot representing the fold change in expression between *mir430*[-/-] + inj

MiR430 and WT embryos at 256-cell on the y-axis, and between 1024-cell and 256-cell in WT

embryos on the x-axis. All upregulated genes are shown in grey, genes induced at 1024-cell

(62%) are shown in blue. **c.** Heatmap of chromatin accessibility at 256-cell of the 716 genes

that are upregulated in *mir430*[-/-] + inj MiR430 embryos compared to WT embryos at 256-cell

stage, and – for comparison – the 716 most expressed genes in WT embryos at 256-cell stage.

Genes are ranked by accessibility. **d.** Graph showing the fraction of promoters of the 716

upregulated (in *mir430*[-/-] + inj MiR430 *vs.* WT at 256-cell stage), and the 716 most expressed

genes (in WT at 256-cell) amongst the x% most accessible promoters at 256-cell in WT

embryos. **e.** Average expression level of the 716 upregulated genes (*mir430*[-/-] + inj MiR430 *vs.*

WT at 256-cell stage), the 716 most expressed genes, and all annotated genes, at the 256-cell

stage, in WT. Outliers are not shown. **f.** Visualization of RNAPII-Ser2P and RNAPII-Ser5P

with antigen-binding fragments (Fabs) in WT and *mir430*[-/-] + inj MiR430 embryos at 512- and

1024-cell stage. Shown are representative micrographs of individual nuclei. See Extended Data

Fig. 9-10 for complete cell cycles. **g.** Quantification of Ser5P-positive/Ser2P-negative (red),

Ser5P-positive/Ser2P-positive   (yellow)   and   Ser5P-negative/Ser2P-positive   (green)

transcription bodies during the cell cycle at 512-cell and 1024-cell, as shown in f. Number

(upper panel) and percentage (lower panel) are shown.

Figure 4. CDK9 sequestration in *mir430* transcription bodies inhibits transcription elongation elsewhere. **a.** Visualization of CDK9 and RNAPII-Ser2P by immunofluorescence in WT nuclei at 256c. Nuclei are labelled with DAPI. **b.** Visualization of RNAPII-Ser2P with antigen-binding fragments (Fabs), and *miR430* RNA with MOVIE in WT, WT + inj *cdk9* mRNA, and *mir430*⁻/⁻ + inj MiR430 embryos across stages. Shown are representative micrographs of individual nuclei, extracted from a spinning disk confocal microscopy time

lapse. See Extended Data Fig. 11-14 for complete cell cycles. **c.** Quantification of ectopic

transcription bodies as shown in b. N=3, $8 \leq n \leq 168$. See Methods for a detailed description of

quantification.

Methods

<u>Zebrafish and molecular biology approaches</u>

Zebrafish husbandry and manipulation

Zebrafish were maintained and raised under standard conditions, and according to Swiss

regulations (canton Vaud, license number VD-H28). Wild type (TLAB) and mutant (*mir430*[-/-]

)[9] embryos were collected immediately upon fertilization and allowed to develop to the desired

stage at 28°C. *mir430*[-/-] embryos were rescued by injecting 2nL of rescue solution (10μM of

MiR430a, MiR430b and MiR430c duplex in 1x siRNA solution (300mM KCl, 30mM HEPES,

1mM MgCl$_2$, brought to pH 7.5 with KOH)), as described before [9,24]. If the chorion was

chemically removed, this was done by incubating embryos right after fertilization in 1.5 mg/mL

Pronase E (Sigma-Aldrich, #107433) for approximately five minutes. Developmental stage

was determined by morphology (according to [37]), the distance between nuclei, as well as cell

size, cell cycle duration, and synchronicity [9]. Fabs were injected into the cell at 0.44ng/embryo

(αRNAPII-Ser2P-Alexa488), or 0.66ng/embryo (αRNAPII-Ser5P-Cy5). Lissamine-labelled

anti-*miR430* morpholino (5'-TCTACCCCAACTTGATAGCACTTTC-3'-Lissamine, Gene

Tools) was injected into the cell at 14 fmole/embryo [21]. s[4]UTP (TriLink, tebu-bio GmbH, #N-

1025-1, diluted to 50mM in 10mM Tris-HCl pH 7.5) was injected into the yolk at

50nM/embryo.

Genotyping

Embryos (live and fixed) were genotyped using the HotSHOT method [38].

mRNA production and injection

eGFP with three perfect *miR430* target sites in its 3'UTR (eGFP-3xPT-miR430b) was inserted

in the pCS2[+] plasmid (Addgene) as described before [24]. The coding sequence of *cdk9*

(ENSDART00000065859.8) was amplified from cDNA using SuperScript III (Invitrogen,

#18080085) and oligodT primers, and inserted in the pCS2$^+$ plasmid (Addgene) using

restriction-enzyme cloning. mRNA was synthesized using the mMessage mMACHINE

Transcription Kit (ThermoFisher, #AM1340) and purified with the RNeasy MinElute Cleanup

Kit (QIAGEN, #74204). mRNA was injected into embryos at the 1-cell stage in the following

amounts: 50pg of eGFP mRNA, and 100pg of RFP or *cdk9* mRNA.

Preparation of antigen-binding fragments (Fabs)

Fluorescently labelled Fabs specific to RNAPII-Ser5P and RNAPII-Ser2P were prepared from

monoclonal antibodies specific to RNAPII-Ser5P and Ser2 phosphorylation [39–41]. Monoclonal

antibodies were digested with Ficin (ThermoFisher Scientific), and Fabs were purified through

protein A-Sepharose columns (Cytiva) to remove Fc and undigested IgG. After passing through

desalting columns (PD MiniTrap G25; Cytiva) to substitute the buffer with PBS, Fabs were

concentrated up to >1 mg/ml using 10 k cut-off filters (Amicon Ultra-0.5 10 k; Merck). Fabs

were conjugated with Alexa Fluor 488 (Sulfodichlorophenol Ester; ThermoFisher Scientific)

or Cy5 (N-hydroxysuccinimide ester monoreactive dye; Cytiv) to yield ~1:1 dye:protein ratio.

After the buffer substitution with PBS, the concentration was adjusted to ~1 mg/ml.

Probe production

A DIG-labeled antisense probe against *goosecoid* (*gsc*) was generated by *in vitro* transcription

from a linearized plasmid using 10x Transcription buffer (Roche, #11465384001), 10x DIG

RNA labelling mix (Roche, #11277073910), T7 RNA polymerase (Roche, #10881775001) and

RNase OUT (Invitrogen, #10777019). Probe was purified with the RNeasy MinElute Cleanup

Kit (QIAGEN, #74204), diluted in Hybridization buffer and stored at -20°C.

Whole mount in situ hybridization

Whole mount *in situ* hybridization (WM-ISH) was performed as described before [42], with some

differences. In brief, embryos were fixed at the desired stage in 4% (w/v) PFA in PBS pH 7.4

overnight at 4°C. Embryos were then manually dechorionated with sharp forceps (Dumont No.

5) in PBS with 0.1% Tween-20 (PBST) and dehydrated by rinsing them in 50% and then 100%

methanol in PBST on a nutator. At this point, embryos could be stored for several weeks at -

20°C. To continue, embryos were gradually rehydrated on a nutator, incubated in Hybridization

buffer (50% formamide, 1.3x SSC pH 5.0, 5mM EDTA, 50μg/mL torula yeast RNA, 0.2%

Tween-20, 0.1% SDS, 100μg/mL Heparin) for 1 hour at 70°C, and incubated with anti-

goosecoid DIG-labelled probe overnight at 70°C. The next day, the probe was washed off with

Hybridization buffer and embryos were incubated for 15 minutes in 50% (v/v) Hybridization

buffer in TBST (25mM Tris-HCl pH 7.8, 137mM NaCl, 2.7mM KCl, 0.1% Tween-20) at 70°C.

After rinsing with TBST, embryos were incubated for 1 hour in TBST and then in Blocking

Buffer (2% blocking reagent in maleic acid buffer pH 7.5 (Roche, #11096176001) and 10%

sheep serum (Sigma Aldrich, #S2263) in TBST) at room temperature on a nutator. anti-DIG

antibody (Roche, #11093274910) was added in a dilution of 1:2.000 and embryos incubated

overnight at 4°C on a nutator. The next day, embryos were washed with TBST and then NTMT

(100mM NaCl, 100mM Tris-HCl pH 9.5, 50mM $MgCl_2$, 0.1% Tween-20), and stained in

NTMT-NBT-BCIP solution (4.5μL/mL of NBT (Roche, #11383213001) and 3.5μL/mL of

BCIP (Roche, #11383221001) in NTMT) in the dark. Once sufficiently stained, embryos were

washed with PBST, and then in 100%, 50% and 30% methanol in PBST at room temperature

on a nutator. Embryos were stored at 4°C in PBST for up to a few days before imaging.

Immunofluorescence

Dechorionated embryos were fixed at the desired stage in 4% (w/v) PFA in PBS pH 7.4

overnight at 4°C. Embryos were then dehydrated by incubating them in 30%, 50% and then

100% methanol in PBS and incubated overnight at -20°C. To continue, embryos were gradually

rehydrated. All following steps were performed while rocking. Embryos were washed multiple

times with PBST (0.8% Triton-X 100 in PBS) and incubated in blocking buffer (10% NGS

(Sigma-Aldrich, #G6767) and 4% BSA (Sigma Aldrich, #A8022) in PBST) for four hours at

room temperature. Primary antibodies were added in 1% NGS in PBST overnight at 4°C (anti-

CDK9, clone C12F7, Cell Signaling Technology, monoclonal IgG antibody generated in

rabbit, 1:200 dilution; anti-RNAPII-Ser2P, clone H5, ab24758, monoclonal IgM antibody

generated in mouse, Abcam, 1:1000 dilution). The next day, embryos were washed with PBST

and then incubated with secondary antibodies (anti-rabbit-IgG AlexaFluor488-conjugated,

Thermo Scientific, #A-21206, 1:500 dilution; anti-mouse-IgM AlexaFluor594-conjugated,

Thermo Scientific, #A-21044, 1:500 dilution) in 1% NGS in PBST overnight at 4°C. Embryos

were then washed with PBST, manually deyolked using sharp forceps (Dumont No. 5), and

mounted in Vectashield Antifade Mounting Medium with DAPI (Vector Laboratories, VC-H-

1200-10) before imaging them on a Spinning Disk microscope.

Enriched SLAM-Sequencing (eSLAM-Seq)

Rationale

Early embryos contain large amounts of maternally loaded RNA [19] which mask the nascent

RNAs in total RNA sequencing approaches. We therefore developed a method to sensitively

detect nascent transcripts. Methods to enrich nascent transcripts have previously been

described. Typically, an uracil/UTP analog (EU/s^{4}UTP) is injected at the 1-cell stage and is

incorporated into newly made RNAs. Such labelled transcripts can then be biotinylated and

enriched by pull-down [20,43]. SLAM-Seq is another approach in which nascent transcripts are

labelled by s[4]UTP incorporation, but here the label is used to convert 4-thio-uracils to

carboxyamido-methylated 4-thio-uracils, which will be misread as a cytosine during reverse

transcription, introducing specific point mutations that enable the identification of reads

derived from labelled transcripts by sequencing [44–46]. We combined these two approaches in

enriched SLAM-Seq (eSLAM-Seq), which is conceptually similar to TT-TimeLapse-Seq and

TT-SLAM-Seq [47,48]. Specifically, we used a combination of s[4]UTP labelling [43] with improved

biotinylation and enrichment chemistry [49,50], followed by SLAM-Seq chemistry.

Sample preparation

Because of the light-sensitive nature of s[4]UTP, all steps until the treatment of RNA with

iodoacetamide (IAA) were performed in the dark. When staging embryos at the

stereomicroscope, for example, we used the lowest light-setting that allowed us to still see the

embryos, and a UV filter was set between the injection plate and the light source to protect the

s[4]UTP. Successive steps were performed protecting the samples from the light by using low

light settings in the laboratory room, by working inside a cardboard box, and by covering the

samples with aluminium foil.

Synchronized s[4]UTP-injected embryos were collected and left to develop to the desired stage.

They were collected in as little 0.3x Danieau's Buffer as possible, homogenized in 500μL

Qiazol (QIAGEN, #79306) with a 5mm stainless steel bead (QIAGEN, #69989) using a

TissueLyser (30 seconds at 30Hz) and stored at -80°C. 100 embryos (256-cell stage) or 50

embryos (1024-cell, Oblong and Sphere stage) were used. Chloroform was added and RNA

was extracted by phase separation, using a MaXtract tube (QIAGEN, #129046), after which it

was precipitated with isopropanol supplemented with 0.2mM DTT and glycogen. The pellet

was washed with 80% ethanol supplemented with 1mM DTT, air-dried, and resuspended in

40μL of 1mM DTT. DNase treatment was performed in 100μL of 1x DNAse reaction buffer

with 4U of TURBO DNase (ThermoFisher, AM2238) for 30 minutes at 37°C. The volume was

brought to 250μL with nuclease-free water and RNA was extracted with 25:24:1 Phenol-

Chloroform-Isoamyl alcohol (Invitrogen, #15593-031) using a MaXtract tube, precipitated

with isopropanol supplemented with 0.2mM DTT, glycogen and sodium acetate. The pellet

was washed with 80% ethanol supplemented with 1mM DTT, air-dried, and resuspended in

40μL of nuclease-free water.

Labelled RNA was biotinylated in 100μL of 20mM HEPES pH 7.4, 1mM EDTA pH 8.0,

10ng/μL MTSEA-biotin-XX (Biotium, #90066) and 20% DMF) by incubation at room

temperature for 30 minutes while rotating. The volume was brought to 250μL with nuclease-

free water and RNA was extracted with 24:1 Chloroform-Isoamyl alcohol (Sigma-Aldrich,

#227056) using a MaXtract tube, and precipitated with isopropanol supplemented with

glycogen and NaCl. The pellet was washed with 80% ethanol, air-dried, and resuspended in

50μL of nuclease-free water.

To enrich for biotinylated RNA, Dynabeads MyOne Streptavidin C1 beads (Invitrogen,

#65001) were used. Per sample, 10μL of beads were washed twice in nuclease-free water and

twice in high-salt wash buffer (100mM Tris-HCl pH 7.4, 10mM EDTA pH 8.0, 1M NaCl,

0.05% Tween-20), incubated for at least 1 hour at room temperature in bead-blocking buffer

(40ng/μL glycogen in high-salt wash buffer) and washed twice with high-salt wash buffer.

RNA was denatured at 65°C for 10 minutes, incubated for 5 minutes on ice and then added to

the blocked beads. After incubation for 15 minutes while rotating at 30 rpm, the supernatant

was removed and beads were washed 3 times with high-salt wash buffer, twice with 5.9M

guanidinium chloride (with 5-minute incubation at room temperature between consecutive

washes), once with TE buffer and three times with TE buffer (with 5-minute incubation at 55°C

between consecutive washes). Bead-bound RNA was finally released by adding 25μL of

freshly made elution buffer (100mM DTT, 20mM HEPES pH 7.4, 1mM EDTA pH 8.0,

100mM NaCl, 0.05% Tween-20) twice. Volume was brought to 100μL with nuclease-free

water and RNA was precipitated with isopropanol supplemented with 0.2mM DTT, glycogen

and NaCl. The pellet was washed with 80% ethanol supplemented with 1mM DTT, air-dried,

and resuspended in 20μL of 1mM DTT.

Iodoacetamide-mediated alkylation was performed in 10mM IAA (freshly dissolved in 100%

ethanol, BioUltra, #I1149), 50mM NaPO$_4$ pH 8.0 and 50% DMSO for 15 minutes at 50°C.

Samples were then transferred on ice and the reaction stopped by quickly adding 1.4μL of 1M

DTT. RNA was precipitated with 100% ethanol supplemented with glycogen and sodium

acetate overnight. The pellet was washed with 80% ethanol, air-dried, and resuspended in

8.5μL of nuclease-free water.

Libraries were prepared with the TruSeq stranded mRNA Kit (Illumina, #20020594) and

TruSeq RNA UD Indexes (Illumina, #20022371). The polyA enrichment step was not

performed. Fragmentation was performed for 1 minute at 94°C. Paired end 150bp sequencing

was performed on an Illumina NovaSeq 6000 System, loading each library on two lanes

(Illumina Reagent Kit v1.5). Sequencing data were demultiplexed with bcl2fastq 2.20.

Quality controls

To assess the integrity of total RNA (Extended Data Fig. 4c, left panel), 1μg/sample was diluted

in 2x RNA Gel Loading Dye (Thermo Scientific, #R0641), denatured for 90 seconds at 85°C,

chilled for 5 minutes on ice and loaded onto an agarose gel alongside RiboRuler High Range

RNA Ladder (ThermoFisher, #SM1823). The presence of two prominent bands (rRNA) are an

indicator of RNA integrity.

To test the specificity of s^{4}UTP biotinylation (Extended Data Fig. 4c, right panel), the RNA

was then transferred from the gel onto a Nylon Hybond-N+ membrane (Amersham,

#RPN203B) by capillary transfer using a Whatman Turboblotter in 20X SSC according to

manufacturer's instructions overnight. After drying, the membrane was crosslinked twice with

120mJ/cm^2 using a UV Stratalinker 1800 (Stratagene). The membrane was then incubated for

20min in Blocking Buffer (10% SDS and 1mM EDTA in 1xPBS pH 7.5), for 1h in 1:10.000

Streptavidin-HRP conjugate (Perkin, #NEL750) in Blocking buffer, 2x for 10min in 10% SDS

in 1xPBS pH 7.5, 2x for 10min in 1% SDS in 1xPBS pH 7.5, and 2x for 10min in 0.1% SDS

in 1xPBS pH 7.5, all while nutating [51]. Finally, the signal was visualized with ECL Western

Blotting Detection Reagents (Amersham, #RPN2106) according to manufacturer's

instructions, and documented with a Fusion FX (Vilber).

<u>Bioinformatic analyses</u>

eSLAM-Seq

To map the raw sequencing data we concatenated Fastq files that came from the same library

but that were run on different lanes with the 'cat' command (GNU core utilities, linux terminal),

trimmed them using Trimmomatic (Paired-end mode, LEADING:3 TRAILING:3

SLIDINGWINDOW:4:15 MINLEN:36) [52], and mapped them using HISAT-3N (--base-change

T,C --repeat --rna-strandness RF --fr --no-discordant) [53], which is a HISAT variant specifically

designed for methods like SLAM-Seq that introduce specific point-mutations. Uniquely

mapped fragments were filtered with Samtools [54] and mate information was fixed with Picard

FixMateInformation (--IGNORE_MISSING_MATES FALSE) [55].

To distinguish between naturally occurring SNPs and SLAM-Seq-specific mismatches, we first

identified naturally occurring SNPs by analysing four IAA-minus samples (one per stage) in

which the iodoacetamide treatment – and therefore the base conversion – was not performed.

The bam files were converted to a suitable format with the Samtools mpileup function and then

SNPs were identified by Varscan pileup2snp (--min-coverage 20 --min-reads2 5 --min-avg-

qual 15 --min-var-freq 0.25 --p-value 0.01). The union of the SNPs that were identified was

used as a consensus list. Then the splbam.py script from pulseR [56] was used to select fragments

with at least one SLAM-Seq-specific mismatch.

To assess gene expression levels of annotated genes, raw read counts were obtained using

FeatureCounts (-p --countReadPairs -B --minOverlap 10 -s 2 -t exon -g gene_id) [57].

Normalization, Principal Component Analysis (PCA) and call for differential expression (alpha

= 0.01 and $\log_2$(Fold Change) threshold = 2) was performed using DESeq2 [58]. When

performing PCA, only the top 500 genes (selected by highest variance in expression levels)

were considered. When counting the number of differentially expressed genes, mitochondrial

and *mir430*-coding genes were excluded.

Motif analysis

To assess the presence and enrichment of specific motifs in promoters of upregulated genes,

we started with the non-redundant JASPAR 2022 Vertebrate core database. From this, only

motifs whose cognate transcription factors are expressed during ZGA (Riboprofiling dataset,

>10rpkm, [30]) were kept. The Nanog motif [29] and Pou5f1 motif [31] were added, because it is

known that these factors are important during zebrafish ZGA. Next, we extracted promoter

sequences (4000bp centred around the TSS) for upregulated genes as well as non-expressed

genes (less than 10 raw reads in total across all samples, background group). Motif enrichment

was then assessed using SEA (MEME Suite, [59]). To estimate depletion of motifs, the

background and input sample were inverted. Only motifs with a P-value < 0.05 and an E-value

≤ 10 were considered.

Promoter accessibility analysis

To analyse promoter accessibility of the upregulated genes in WT, we used an ATAC-Seq

dataset that we previously generated [29]. Replicates were merged and the signal in 4000bp

centred around the TSS was plotted using Deep Tools [60]. Promoter signal was used to rank all

promoters by accessibility and to assess their enrichment amongst the most accessible

promoters. The 716 most expressed (mean TPM) genes (excluding mitochondrial genes and

mir430-coding genes) were used as a comparison.

Graph plotting for eSLAM-Seq

Plots were generated using R version 4.0.5 [61] with the ggplot2 [62] and Ggbio package [63].

<u>Microscopy</u>

Fluorescence microscopy

To determine MiR430-activity using the GFP sensor assay, injected embryos were manually

dechorionated at 24hpf and kept in Petri dishes. Fluorescence signal was measured on an

OLYMPUS SZX16 stereo microscope with a MicroPublisher 5.0 RTV-R-CLR-10C colour

camera (5x magnification, 400ms exposure time).

Bright field imaging

For bright field time lapse imaging, a mold [64] was used to create agarose wells using as little

as possible 1% agarose in 0.3x Danieau's solution in an Ibidi µ-Dish 35mm, high (#81158).

The dish was mounted on a 28°C heated stage (by a Warner automatic temperature controller)

and dechorionated embryos were loaded to the single wells at the 2-cell stage. Movies with 10-

minute time intervals were taken on a Zeiss Axio Observer.Z1 microscope (Zeiss 5x/0.25 Fluar

air objective, Zeiss AxioCam MRm Monochrome CCD camera). A z-stack (400µm in 9 steps)

was taken for each embryo. A lid was used to reduce evaporation of buffer during imaging.

To assess developmental progression of live embryos past somitogenesis, snapshots of

manually dechorionated embryos were taken at 24hpf, 48hpf and 72hpf on a Zeiss Axio

Observer.Z1 microscope (Zeiss 2.5x/0.08 EC Plan-Neofluar air objective, Zeiss AxioCam ICc

1).

The gene expression pattern in fixed embryos after WM-ISH was visualized on a LEICA

M165C stereomicroscope with LEICA MC170 HD colour camera (4x magnification).

Embryos were oriented such that the dorsal region of the embryo was visible.

Spinning Disk microscopy

For imaging on a spinning disk microscope, dechorionated embryos were mounted at 32-cell

stage in small droplets of 1% low-melting agarose (Invitrogen, #16520-050) in 75% v/v 0.3x

Danieau's solution and 25% v/v Optiprep Density Gradient Medium (Sigma Aldrich, #D1556)

[65] on a Ibidi μ-Dish 35mm, high (#81158). Embryos were brought closer to the coverslip

surface by keeping the dish upside down until the agarose solidified (~20 minutes). More

agarose was then added to avoid drying out.

Transcription bodies in live embryos were imaged on a Nikon eclipse Ti2 with a Yokogava

CSU-W1 Spinning Disk unit at 28°C (Okolab temperature control and microscope enclosure).

Multi-position, z-stack (20μm in 41 steps, NIDAQ Piezo Z) time-lapse (2 minutes time

interval) imaging was performed with a Nikon 60x/1.2 Plan Apochromat VC water objective.

Images were captured in parallel on two Photometrics Prime 95B cameras with 100ms

exposure time at 10% laser intensity.

Immunofluorescence signal (CDK9, RNAPII-Ser2P and DAPI) was also imaged on a Nikon

eclipse Ti2 with a Yokogava CSU-W1 Spinning Disk unit. Images were taken with a Nikon

60x/1.2 Plan Apochromat VC water objective. Each channel was captured in series on a

Photometrics Prime 95B camera with 100ms exposure time at 10% laser intensity each.

<u>Image processing and analysis</u>

Software used for image analysis

Microscopy image handling was done using FIJI [66]. Further data processing was carried out

using R version 4.0.5 [61] and plots were generated with the ggplot2 package [62].

Analysis of eGFP reporter and RFP control

To assess *mir430* activity, mean fluorescence of eGFP (targeted by MiR430) and RFP (not

targeted, and used as a normalizer) were measured in a rectangle of equal size in the head

region and a background region outside of the embryo. The signal ratio was then calculated as

follows: $(eGFP_{head} - eGFP_{background}) / (RFP_{head} - RFP_{background})$.

Detection and classification of mir430 and ectopic transcription bodies

We identified transcription bodies by the presence of RNAPII-Ser2P signal. To classify them

into *mir430* and ectopic transcription bodies, we analysed *miR430* MOVIE signal. In case of

overlap between RNAPII-Ser2P and MOVIE signal, bodies were classified as *mir430*

transcription bodies. In case of no overlap between RNAPII-Ser2P and MOVIE signal, bodies

were classified as ectopic transcription bodies (Extended Data Fig. 16). We used FIJI plugin

ComDet [67], which detects spots using consistent intensity thresholding parameters that were

manually set and used for all experiments. We note that frames taken after the disassembly of

the nuclear envelope were not analysed because at that time transcription bodies also

disassembled.

Detection and classification of transcription bodies by transcription state

Nuclei were segmented using the RNAPII-Ser5P channel, and RNAPII-Ser5P and Ser2P

bodies detected using ComDet. This was used to classify transcription bodies into initiating,

initiating-elongating and elongating bodies. We then identified and removed *mir430*

transcription bodies from the analysis based on their morphological features (early appearance

during cell cycle, larger size, growth behaviour) using the RNAPII-Ser2P channel. Here too,

frames taken after the disassembly of the nuclear envelope were not analysed. For each nucleus,

t=0 is set as the first frame at which spots appear, and the last time point is set as the last frame

before prophase. Nuclei in which spots were detected for only one frame were discarded.

<u>Sample size and statistics</u>

A minimum of 3 biological replicates (N) was acquired for each experimental condition.

Lower-case n (n) refers to the number of embryos or nuclei for each experimental condition.

Kruskal-Wallis tests were performed to determine if there is a statistically significant difference

between the three genotypes in the degree of MiR430 activity as determined by the eGFP-

Sensor assay (Fig. 1e), the time of completion of Epiboly (Fig. 1g) and the time of appearance

of Kupffer's vesicle (Fig. 1h). If this test was statistically significant (P-value < 0.05), pairwise

comparisons with Bonferroni correction were performed using a pairwise Wilcoxon rank-sum

test. A comparison was considered significant when adjusted P-value < 0.05. Dotplots were

used to display the data, and adjusted P-values were reported using WT as reference. Red

rhombuses represent the median.

A pairwise Wilcoxon rank-sum test with Bonferroni correction was performed to determine if

the upregulated genes and the top 716 most expressed genes are significantly higher expressed

compared to all genes (based on TPM expression values) (Fig. 3e). A comparison was

considered significant when adjusted P-value < 0.05.

A pairwise Wilcoxon rank-sum test with Bonferroni correction was performed at each stage to

determine if there is a statistically significant difference in the number of detected ectopic

transcription bodies between WT, *mir430*$^{-/-}$ + inj MiR430 and WT + inj *cdk9* mRNA (Fig. 4c).

A comparison was considered significant when adjusted P-value < 0.05.

Extended Data Figure 1

a

b

Extended Data Figure 1. MiR430 activity is restored by injecting MiR430 RNA in *mir430*⁻*/-* embryos. a. Schematic representation of expected eGFP and RFP expression in embryos with active and inactive MiR430 microRNA activity. **b.** Representative micrographs showing eGFP and RFP expression in WT, *mir430*⁺*/-*, *mir430*⁻*/-* and *mir430*⁻*/-* + inj MiR430 embryos at 24hpf.

Extended Data Figure 2

Extended Data Figure 2. Phenotypic analysis of *mir430*[-/-] embryos with and without injected MiR430 (*gsc* expression). Whole mount *in situ* hybridization (WM-ISH) of *Goosecoid (Gsc)* in WT, *mir430*[+/-], *mir430*[-/-] and *mir430*[-/-] + inj MiR430 embryos at Shield stage. Representative micrographs show the embryos in dorsal view. Misexpression (reduced expression levels and reduced anterior extension of expression pattern) of *Gsc* can be observed in *mir430*[-/-] embryos. The fraction in the lower left corner represents the frequency with which the shown phenotype was observed. A schematic representation of the phenotypes is shown below.

Extended Data Figure 3

Extended Data Figure 3. Phenotypic analysis of *mir430⁻ᐟ⁻* embryos with and without injected MiR430 (developmental progress). a. Embryos at 8-Somite stage for WT, *mir430⁺ᐟ⁻*, *mir430⁻ᐟ⁻* and *mir430⁻ᐟ⁻* + inj MiR430 embryos. The reduced body axis extension and the malformation of the optic primordium in *mir430⁻ᐟ⁻* embryos are highlighted. Representative micrographs are shown. **b.** Larvae at 24, 48 and 72 hours post fertilization (hpf) for WT, *mir430⁺ᐟ⁻*, *mir430⁻ᐟ⁻* and *mir430⁻ᐟ⁻* + inj MiR430 embryos. The malformation of trunk morphology and of the eye, the development of heart oedema and the appearance of blisters at the tail tip in *mir430⁻ᐟ⁻* embryos are highlighted. Representative micrographs are shown. The 48hpf stage is also shown in Fig. 1i.

Extended Data Figure 4

Extended Data Figure 4. Establishment of enriched SLAM-Sequencing (eSLAM-Seq) in zebrafish embryos. a. Schematic representation of eSLAM-Seq protocol. See Methods for more detail. **b.** Representative micrographs of larvae that were/were not injected with 50nM/embryo of s⁴UTP at the 1-cell stage. No developmental malformations were observed

upon injection. **c.** Equal amounts of biotinylated and heat-denatured total RNA derived from

injected (+) and uninjected (-) embryos at 1024-cell and Sphere stage were run on an agarose

gel (left panel). The RNA was then transferred on a nylon membrane and biotinylation detected

by Streptavidin-HRP (right panel). Biotinylation was only observed in RNA derived from

injected embryos, showing that biotinylation is specific for s^4UTP-labeled RNA. L: RNA

Ladder. **d.** Box plot representation of mismatch rate in Iodoacetamide- (N=24) and Ethanol-

treated (N=4) samples. As expected, only Iodoacetamide-treated samples show a high level in

A>G (First strand) and T>C (Second strand) mismatch rate. **e.** Principal Component Analysis

(PCA) plot based on the top 500 genes (selected by highest variance in expression levels).

Samples separate according to their developmental stage along the first PC and according to

their experimental condition (WT and *mir430*$^{-/-}$ + inj MiR430) along the second PC. **f.** Box

plot representation of percentage of mitochondrial and nuclear reads, separated by

developmental stage. Most reads are of mitochondrial origin at early ZGA, as shown before [43],

with a gradual increase in the nuclear fraction during later stages.

Extended Data Figure 5

816 **Extended Data Figure 5. The disruption of two large endogenous transcription bodies**

817 **causes short-term misregulation of gene expression.** Volcano plots showing up- and

818 downregulated genes in *mir430*[-/-] + inj MiR430 *vs.* WT embryos at 256-cell **(a)**, 1024-cell **(b)**,

819 Oblong **(c)** and Sphere **(d)** stage. Panel A is also shown in Fig. 2a.

Extended Data Figure 6

823 **Extended Data Figure 6. RNAPII-Ser2P signal at 256-cell, 1024-cell, Oblong and Sphere**

824 **stage in *mir430^{-/-}* + inj MiR430 embryos.** Visualization of RNAPII-Ser2P signal with Fabs.

825 Representative micrographs are shown.

Extended Data Figure 7

Extended Data Figure 7. Correlation between expression level in *mir430^{-/-}* + inj MiR430

***vs.* WT and 1024-cell, Oblong and Sphere *vs.* 256-cell in WT embryos. a.** Scatterplots

representing the fold change in gene expression (log$_2$ scale) between *mir430^{-/-}* + inj MiR430

and WT at 256-cell (y-axis), and the fold change in gene expression ($\log_2$ scale) between 1024-

cell and 256-cell (left), Oblong and 256-cell (central) and Sphere and 256-cell (right) in WT

embryos (x-axis). All upregulated genes are shown in grey, and genes induced at 1024-cell are

highlighted in blue. The left graph is also shown in Fig. 3b. **b-d.** Same as in (A) but genes that

are highlighted represent genes that are induced at Oblong **(b)**, Sphere **(c)**, or not induced

during ZGA **(d)**.

Extended Data Figure 8

Extended Data Figure 8. Transcription factor motif analysis of upregulated genes. Scatter

plot representing the Enrichment Ratio (log$_2$) of transcription factor motifs in the promoters

(TSS ± 2Kb) of upregulated genes compared to the promoters of not expressed genes (y-axis),

and the percentage of promoters of the upregulated genes that have the motif (x-axis). Only

motifs whose corresponding protein is translated during early development [30] were considered,

and only motifs with a p-value < 0.05 and E-value ≤ 10 are shown. Motifs corresponding to

the three pluripotency factors Nanog, Sox19b and Pou5f3 (POU5F1, Pou5f1, Pou5f1::Sox2,

POU5F1B, POU2F1, POU2F1::SOX2) are labelled in red.

Extended Data Figure 9

Extended Data Figure 9. Stalled transcription bodies go into productive transcription in *mir430$^{-/-}$* **+ inj MiR430 embryos (512-cell stage).** Visualization of initiating (Ser5P) and elongating (Ser2P) RNAPII with Fabs in WT **(a)** and *mir430$^{-/-}$* + inj MiR430 **(b)** embryos at 512-cell stage. Shown are representative images of individual nuclei, extracted from a spinning disk confocal microscopy time lapse. The micrographs shown in the main figure are highlighted in green.

Extended Data Figure 10

Extended Data Figure 10. Stalled transcription bodies go into productive transcription in *mir430*[-/-] **+ inj MiR430 embryos (1024-cell stage).** Visualization of initiating (Ser5P) and elongating (Ser2P) RNAPII with Fabs in WT **(a)** and *mir430*[-/-] + inj MiR430 **(b)** embryos at 1024-cell stage. Shown are representative images of individual nuclei, extracted from a spinning disk confocal microscopy time lapse. The micrographs shown in the main figure are highlighted in green.

Extended Data Figure 11

Extended Data Figure 11. RNAPII-Ser2P signal in WT, WT + inj *cdk9* mRNA and *mir430*-/- + inj MiR430 embryos at 128-cell stage. Visualization of RNAPII Ser2P (with Fabs) and *miR430* RNA (with MOVIE) in WT **(a)**, WT + inj *cdk9* mRNA **(b)**, and *mir430*-/- + inj MiR430 **(c)** embryos at 128-cell stage. Shown are representative images of individual nuclei, extracted from a spinning disk confocal microscopy time lapse. The micrographs shown in the main figure are highlighted in green.

Extended Data Figure 12

Extended Data Figure 12. RNAPII-Ser2P signal in WT, WT + inj *cdk9* mRNA and *mir430*$^{-/-}$ + inj MiR430 embryos at 256-cell stage. Visualization of RNAPII Ser2P (with Fabs) and *miR430* RNA (with MOVIE) in WT **(a)**, WT + inj *cdk9* mRNA **(b)**, and *mir430*$^{-/-}$ + inj MiR430 **(c)** embryos at 256-cell stage. Shown are representative images of individual nuclei, extracted from a spinning disk confocal microscopy time lapse. The micrographs shown in the main figure are highlighted in green.

Extended Data Figure 13. RNAPII-Ser2P signal in WT, WT + inj *cdk9* mRNA and *mir430*[-/-] + inj MiR430 embryos at 512-cell stage. Visualization of RNAPII Ser2P (with Fabs) and *miR430* RNA (with MOVIE) in WT **(a)**, WT + inj *cdk9* mRNA **(b)**, and *mir430*[-/-] + inj MiR430 **(c)** embryos at 512-cell stage. Shown are representative images of individual nuclei, extracted from a spinning disk confocal microscopy time lapse. The micrographs shown in the main figure are highlighted in green.

Extended Data Figure 14

Extended Data Figure 14. RNAPII-Ser2P signal in WT, WT + inj *cdk9* mRNA and *mir430*[-/-] + inj MiR430 embryos at 1024-cell stage. Visualization of RNAPII Ser2P (with Fabs) and *miR430* RNA (with MOVIE) in WT **(a)**, WT + inj *cdk9* mRNA **(b)**, and *mir430*[-/-] + inj MiR430 **(c)** embryos at 1024-cell stage. Shown are representative images of individual nuclei, extracted from a spinning disk confocal microscopy time lapse. The micrographs shown in the main figure are highlighted in green.

Extended Data Figure 15

Extended Data Figure 15. Model for the role of transcription bodies in transcription

regulation. In WT nuclei, two large transcription bodies are nucleated by the *mir430* locus.

These facilitate the expression of genes that localize there. They also sequester CDK9 (and

potentially other factors) which is required for pause release, thereby stalling transcription

elsewhere in the nucleus in the initiation state. The specific disruption of the two *mir430*

transcription bodies leads to a redistribution of the transcriptional machinery (among which

CDK9) which results in the downregulation of genes that localize to the *mir430* transcription

bodies and the upregulation of genes elsewhere in the nucleus.

Extended Data Figure 16

	0 min	2 min	4 min	6 min	8 min	10 min	12 min
mir430-seeded transcription bodies			1	2	2	2	
ectopic transcription bodies			0	1	5	3	

Extended Data Figure 16. Classification of *mir430* and ectopic transcription bodies.
Transcription bodies are identified by detecting elongating RNAPII signal. If this overlaps with MOVIE signal (detecting *miR430* RNA) they are classified as *mir430*, and if they do not overlap with MOVIE signal, they are classified as ectopic transcription bodies. See Methods for more detail.

Extended Data Table 1. List of differentially expressed genes in mir430$^{-/-}$ + inj MiR430 *vs.* WT at all stages.