On the origin and evolution of microbial mercury methylation

Heyu Lin[1, *], Edmund R. R. Moody[2], Tom A. Williams[3] and John W. Moreau[1, 4, *, †]

[1]School of Geographical, Atmospheric and Earth Sciences, The University of Melbourne,

Parkville, Victoria 3010, Australia

[2]School of Earth Sciences, University of Bristol, Bristol, BS8 1RJ, United Kingdom

[3]School of Biological Sciences, University of Bristol, Bristol, BS8 1TQ, United Kingdom

[4]School of Geographical and Earth Sciences, University of Glasgow, Glasgow, G12 8RZ,

United Kingdom

KEYWORDS: mercury, methylmercury, evolution, antimicrobial, *hgc* gene, gene loss,

horizontal gene transfer, LUCA

*These authors contributed equally to this work.

†Author for Correspondence: John W. Moreau, School of Geographical and Earth Sciences,

University   of   Glasgow,   Glasgow,   United   Kingdom,   +44   141   330   5461,

john.moreau@glasgow.ac.uk

Abstract

The origin of microbial mercury methylation has long been a mystery. Here we employed

genome-resolved phylogenetic analyses to decipher the evolution of the mercury methylating

gene, *hgcAB*, constrain the ancestral origin of the *hgc* operon, and explain the distribution of

hgc in Bacteria and Archaea. We infer the extent to which vertical inheritance and horizontal

gene transfer have influenced the evolution of mercury methylators and hypothesize that

evolution of this trait bestowed the ability to produce an antimicrobial compound ($MeHg^+$) on

a potentially resource-limited early Earth. We speculate that, in response, the evolution of

$MeHg^+$-detoxifying alkylmercury lyase (encoded by *merB)* reduced a selective advantage for

mercury methylators and resulted in widespread loss of *hgc* in Bacteria and Archaea.

Significance

Neurotoxic methylmercury ($MeHg^+_{(aq)}$) is synthesized from $Hg^{II}_{(aq)}$ in the environment by

microorganisms possessing the gene pair *hgcAB*. Our phylogenetic analyses elucidate the

origin and evolution of the *hgc* operon, and support a hypothesis that mercury methylation

evolved as an antimicrobial production mechanism, possibly from competition for limited

resources on the early Earth. We infer from our analyses that *hgc* has been primarily vertically

inherited in Bacteria and Archaea, with extensive parallel loss, and note that few taxa

possessing *hgc* also possess the gene encoding for $MeHg^+$ demethylation, *merB*. Our findings

support the interpretation that *merB* evolved as a defense mechanism against the evolution of

microbial $Hg^{II}_{(aq)}$ methylation.

Introduction

Microorganisms play key roles in the global mercury cycle. In soils, sediments and natural waters, microbes mediate the enzymatic reductive volatilization of $Hg^{II}_{(aq)}$ to $Hg^0_{(g)}$, the transformation of $Hg^{II}_{(aq)}$ to $MeHg^+_{(aq)}$, the complexation of aqueous $Hg^{II}_{(aq)}$ by bacteriogenic bisulfide, and the adsorption/desorption of $Hg^{II}_{(aq)}$ on/from biogenic metal ox(yhydrox)ides (e.g., Barkay et al. 2003; Barkay & Gu 2022; Barkay & Wagner☐Döbler 2005; Gilmour et al. 2013; Hsu-Kim et al. 2013; Tebo et al. 2004; Wang et al. 2022). Essentially, microbes control or strongly influence the geochemical speciation of mercury in most natural environments today. Microbial mercury methylation in particular has received much attention because $MeHg^+$ is a potent neurotoxin that bioaccumulates across both marine and terrestrial food webs (Kidd et al. 2012).

The origin of microbial mercury methylation has remained a mystery, largely because $MeHg^+$ serves no known biological function. Mercury methylation requires $Hg^{II}_{(aq)}$, and on the early Earth, volcanoes would have emitted abundant $Hg^0_{(g)}$ and some Hg^{II} (Grasby et al. 2019). However, studies have shown that $Hg^0_{(g)}$ can be efficiently oxidized to $Hg^{II}_{(aq)}$ by reactive $Cl^-_{(aq)}$ and $Br^-_{(aq)}$ (e.g., Amyot et al. 2005, Seigneur and Lohman, 2008, Obrist et al. 2010, Wang et al. 2014). Volcanos that emitted $Hg^0_{(g)}$ on the early Earth could also have emitted reactive HCl and BrO (Bobrowski et al. 2003; Guo and Korenaga 2021), providing oxidants to convert $Hg^0_{(g)}$ to $Hg^{II}_{(aq)}$. Recent work supports a dual role for Archean volcanism in $Hg^0_{(g)}$ and halogen emissions, and subsequent oxidation of $Hg^0_{(g)}$ to $Hg^{II}_{(aq)}$ (Zerkle et al. 2021), as well as the possibility of ephemeral periods of atmospheric oxygenation prior to the Great Oxidation Event at ~2.4 billion years ago (Meixnerova et al. 2021). Following this event, $Hg^{II}_{(aq)}$ would have been commonly found as a trace metal in aqueous environments. We assume therefore that $Hg^{II}_{(aq)}$ would have been available on the early Earth for methylation upon evolution of the ancestral *hgc* operon. Intriguingly however, the obligately

anaerobic and "canonical" Hg methylator, *Pseudodesulfovibrio mercurii*, can apparently

oxidize dissolved Hg^0 directly, and methylate the resulting $Hg^{II}_{(aq)}$ (Columbo et al. 2013).

Microbial mercury methylation is currently thought to require the *hgc* operon, most

closely related to the ancient carbon monoxide dehydrogenase/acetyl-CoA synthase gene,

cdh, suggested to have been present in the Last Universal Common Ancestor ("LUCA";

Adam et al. 2018; Parks et al. 2013; Sousa & Martin 2014; Weiss et al. 2016, 2018). The

cobalamin-dependent proteins CdhD and CdhE (hereafter "CdhDE") are present in a highly

conserved operon in genomes enabling methyl transfer in the Wood-Ljungdahl (WL) pathway

for autotrophic carbon assimilation (Svetlitchnaia et al. 2006). While HgcA is also predicted

to be a cobalamin-dependent methyltransferase, it only functions in $Hg^{II}_{(aq)}$ methylation and

does not seem to take part in the WL pathway, as other components of this pathway are

incomplete in characterised *hgcA*-carrying genomes (Date et al. 2019). Similar to *cdh*, *hgc*

seems to be broadly distributed across Bacteria and Archaea representing a range of

environmental habitats and redox potentials (McDaniel et al. 2020; Capo et al. 2022). Some

of these bacteria and archaea have been experimentally confirmed to methylate $Hg^{II}_{(aq)}$, while

others are still considered as "putative methylators" encoding homologous *hgcAB* sequences

that have not yet been experimentally validated. In lieu of cultivated isolates, computational

modelling has been employed to test for consistency between putative HgcAB sequences and

HgcAB structure and functionality in confirmed methylators (Gionfriddo et al. 2016; Lin et

al. 2021).

Many studies of mercury methylation have focused on environmental factors

influencing $Hg^{II}_{(aq)}$ bioavailability for uptake by methylating cells. In addition, previous

efforts to identify the biochemical mechanism of $Hg^{II}_{(aq)}$ methylation have attributed the

process to 1) accidental enzymatic catalysis of a methyl group transfer to $Hg^{II}_{(aq)}$ during either

acetyl co-A formation (Choi et al. 1994) or methionine synthesis (Siciliano & Lean 2002), or

2) an as-yet unidentified pathway (Ekstrom et al. 2003). Here, we focus on understanding the

more elusive origin and evolution of mercury methylating genes, employing genome-resolved

phylogenetic analyses to constrain the ancestry of the *hgc* operon and explain its currently

known distribution in Bacteria and Archaea. We assess the extent to which vertical descent

and horizontal gene transfer have shaped *hgc* evolution, and propose a hypothesis for the

functional origin of microbial mercury methylation as an antimicrobial production

mechanism. We also evaluate a potential evolutionary link between *hgc* and *merB*, the latter

gene encoding for organomercury lyase (MerB), which lyses the C-Hg bond in $MeHg^+$ to

release $Hg^{II}_{(aq)}$ for reductive volatilization (i.e., detoxification) by MerA (Tezuka and

Tonomura 1976).

Results and Discussion

hgc present in LUCA

A dataset containing 478 protein sequences belonging to the protein family PF03599 was

retrieved from UniProt Reference Proteomes database v2022_03 (Chen et al. 2011) at 35%

cutoff (RP35) to study the phylogeny of the protein family (**Supplementary Table S1 & S2**).

Phylogenetic reconstruction of the protein family PF03599 defined three deep-branching

clusters, comprising HgcA, CdhD, and CdhE (**Figure 1A**, for details see **Figure S1**). On the

assumption that CdhD- and CdhE-encoding genes were present in LUCA (Adam et al. 2018;

Sousa & Martin 2014; Weiss et al. 2018), we deduce that *hgcA* was also likely to be present

(i.e., these three protein subfamilies diverged in, or before, LUCA; **Figure 1A**). While

robustly rooting single gene trees is challenging, midpoint, Minvar (Mai et al. 2017) and

minimal ancestor deviation (MAD) (Tria et al. 2017) rooting approaches all supported a root

between CdhD and CdhE+HgcA. In comparison to genes encoding for CdhD and CdhE,

which exhibit relatively wide distributions across extant prokaryotes and broadly congruent

evolutionary histories, *hgcA* exhibits a more restricted distribution across extant taxa. The

latter gene is mainly found in Bacteria, mostly affiliated with Deltaproteobacteria, Firmicutes,

and the FCB group, although with a more restricted distribution than in Archaea (**Figure 1A**).

Based on the rooted protein tree (**Figure 1B**), HgcA has a longer stem than CdhD and CdhE,

which might have resulted from accelerated evolution (e.g., associated with a change in

function) or gene loss (or lineage extinction) in former HgcA-encoding clades.

***hgcA* evolved primarily vertically but with extensive loss**

In our phylogenetic analysis, to increase the number and taxonomic coverage of targets, we

further enlarged the sample size by retrieving HgcA homologs in UniProt Reference

Proteomes database v2022_03 at 75% cutoff (RP75). Two other datasets, including one

containing 700 representative prokaryotic proteomes constructed by Moody et al. (2022) and

another containing several novel *hgc*-carriers published by Lin et al. (2021), were retrieved

and incorporated into the RP75 dataset. After removing redundancies and including sequences

from new phyla the number of HgcA sequences in the dataset was increased from 76 to 169

(**Supplementary Table S1**). The resulting HgcA tree was rooted according to the topology of

the HgcA subtree in **Figure 1**.

Our analysis suggests that *hgcA* genes are separated into two well-supported clades

with high bootstrap support, as shown in **Figure 1A** and **Figure 2** (for details see **Figure S2**).

These two clades correspond to a difference in gene structure: the smaller clade comprises

genes in which *hgcA* and *hgcB* became fused to form *hgcAB*, while the larger contains single

subunit (*hgcA*) genes. Several Asgard archaea, including some Thorarchaeota and

Lokiarchaeota, form a clade in the HgcAB subtree, suggesting that gene fusion occurred prior

to the radiation of Asgard phyla. The fused *hgcAB* of *Pyrococcus furiosus* has been

experimentally shown not to methylate $Hg^{II}_{(aq)}$ (Podar et al. 2015), but the functionality of

other fused-*hgcAB* genes, as well as other members of this clade, is yet to be confirmed. For

example, *Nitrospina*-like *hgcA* sequences have been reported as the dominant *hgcA* homologs

in many marine environments (Gionfriddo et al. 2016; Tada et al. 2020; Villar et al. 2020;

Tada et al. 2021). However, the functionality of these sequences has not yet been

experimentally confirmed, as no known *hgc+* members of this genus are currently held in

isolation. Also, the alphaproteobacterium *Breoghania sp.* L-A4 is the only aerobic *hgcA+*

(fused-*hgcAB+*) microorganism found in this analysis, and its methylation capacity under

aerobic conditions also remains untested.

By contrast, the majority of (non-fused) HgcA sequences form a clade which broadly

follows the universal tree (**Figure 2**). In particular, the deepest split lies between archaeal and

bacterial HgcA homologs. Taken together with the sister relationship of HgcA to CdhD/E,

this finding supports an origin of HgcA in or before LUCA. However, the taxonomic

distribution of HgcA in extant Bacteria is patchy (being found mainly in Firmicutes,

Deltaproteobacteria, and the FCB group). Therefore, if HgcA was indeed present in LUCA,

and its gene tree traces the divergence of Archaea and Bacteria, it must have been lost

subsequently in the other bacterial lineages. While it is undoubtably difficult to resolve the

evolutionary history of ancient single genes with certainty, it is interesting that an ancient

origin for HgcA cannot be excluded based on the inferred tree.

Gene loss is a major force driving microbial genome evolution (Bolotin & Hershberg

2016), reflecting environmental change and/or adaptation (Koskiniemi et al. 2012). So far,

hgcA+ methylators have primarily been reported for anoxic or suboxic environments (Capo et

al. 2022; Lin et al. 2021; Parks et al. 2013). Changes in redox potential that could inhibit

$Hg^{II}_{(aq)}$ methylation may have presented a selective environmental pressure to lose *hgcA*. In

addition, long branches of the HgcA subtree, in comparison to CdhD/E, also suggest a weaker

natural selection for gene maintenance, and greater likelihood of loss through genetic drift, a

common scenario observed for many microorganisms (Bolotin & Hershberg 2016).

A few putative HGT events were inferred for the larger clade of the HgcA tree, e.g.,

Marinimicrobia-HgcA clustered with Euryarchaeota-HgcA in the archaeal cluster, with 98%

coverage and 56% identity by blastp against HgcA from Theionarchaea archaeon DG-70-1,

suggesting a possible lateral acquisition of *hgcA*. This inference is consistent with that of

McDaniel et al. (2020) who also suggested HGT played an important role in HgcA evolution.

Our study differs from that of McDaniel et al. in suggesting that *hgcA* may date back to the

LUCA; this inference is based on the nesting of the HgcA subtree within the broader diversity

of the Cdh family (**Figure 1B**), an aspect of the family's evolutionary history not considered

in that analysis, as well as the recovery in our analyses of a deep split between archaeal and

bacterial sequences in the hgcA subtree, notwithstanding some HGT.

hgcA **and** *hgcB* **co-evolved**

The gene *hgcB* is nearly always located immediately downstream of *hgcA*, with rare cases of

being one gene apart (Gilmour et al. 2018). A lack of either *hgcA* or *hgcB* in a genome is

thought to render the microbial host incapable of $Hg^{II}_{(aq)}$ methylation (Parks et al. 2013).

Therefore, the gene pair *hgcAB* likely evolved together as a conserved operon. Although the

HgcB tree was poorly resolved due to short sequence length (95 amino acids on average), the

overall topologies of the HgcB tree and the HgcA tree were congruent, supporting this

inference (**Figure 3**, for details see **Figure S3**). Nevertheless, several *hgcA*+ genomes did not

carry neighbouring *hgcB* genes, including all *Nitrospina* and a few Deltaproteobacteria and

Firmicutes, potentially because of gene loss during evolution or incomplete transfer events

(i.e., only *hgcA* genes were acquired during the HGT events). Another exception involves a

complete *hgcB* gene present downstream of the fused-*hgcAB* gene in the genome of the

deltaproteobacterial endosymbiont, Delta1. This complete HgcB protein had the longest

branch length of the HgcB tree and was located in a sister cluster of the 'HgcB tail' associated

with the fused-*hgcAB* from the same host, suggesting a possible lateral acquisition of *hgcB*

from closely related taxa.

Phylogenetic distribution of *merB* and its relationship to *hgc*

Microbial detoxification mechanisms for $Hg^{II}_{(aq)}$ and $MeHg^{+}_{(aq)}$, including the enzymes MerA

and MerB, respectively, have been well described (c.f., Robinson and Tuovinen 1984, Barkay

et al. 2010, Barkay & Gu 2022). Recent phylogenetic analyses have revealed a thermophilic

and archaeal origin for the *mer* operon, acquired later by bacteria through multiple

independent transfer events (Christakis et al. 2021) reflective of the environmental advantage

of mercury resistance genes (Boyd & Barkay 2012). In total, 225 MerB sequences were

retrieved from the RP35 dataset (**Supplementary Table S2**). MerB homologs were

distributed across Archaea and Bacteria (**Figure 4**, for details see **Figure S4**), with most

representatives found in Terrabacteria (including Actinobacteria, Firmicutes, and Chloroflexi)

and   Proteobacteria   (including   Alphaproteobacteria,   Betaproteobacteria,   and

Gammaproteobacteria). In contrast, only a small number of archaea, including a few

euryarchaeotes and one Thaumarchaeon, were found to encode MerB. These archaeal MerB

homologs were placed within bacterial MerB clades and were not monophyletic, consistent

with the interpretation of recruitment of *merB* to the *mer* operon from a mesophilic ancestor

(Christakis et al. 2021), which we infer involved HGT from Bacteria to Archaea.

We mapped the presence/absence of *merB* and *hgc* genes onto the species tree (**Figure**

5, for details see **Figure S5**) to investigate their phylogenetic relationships. Intriguingly,

although the distribution of MerB homologs overlapped with HgcAB homologs, few genomes

encode *both* MerB and HgcAB, signifying that mecury methylators generally cannot also

demethylate MeHg$^+$. This apparent polarization in gene distribution suggests a functional

conflict between these two genes (i.e., there is utility in encoding for one or the other function,

but not both). Only a few genomes were found to encode for both MerB and HgcAB; this dual

functionality in *Citrifermentans bemidjiense* Bem was previously recognized and discussed

(Lu et al. 2016). The phylogenetic distribution of these opposing enzymes (**Figure 4 &**

Figure 5) suggests that HgcAB was the earlier of the two to evolve, with MerB originating

more recently and radiating via horizontal gene transfer.

Mercury methylation as antimicrobial synthesis?

The reason why microorganisms evolved the ability to methylate Hg^{II} to MeHg$^+$ has long

been a mystery. Previous studies proposed that this process might reduce susceptibility to

Hg$^{II}_{(aq)}$ toxicity (Trevors 1986). But in fact, the ability to produce MeHg$^+_{(aq)}$ has been shown

not to confer Hg$^{II}_{(aq)}$ resistance in methylating microbes (Gilmour et al. 2011). Previous

studies have suggested that MeHg$^+_{(aq)}$ is more toxic than Hg$^{II}_{(aq)}$ to some microbes (Jonas et

al. 1984; Gilmour et al. 2011), and we hypothesise here that Hg$^{II}_{(aq)}$ methylation may have

evolved effectively as an early antimicrobial against microorganisms without the ability to

metabolize/detoxify this organometallic compound. We note that, in seawater, the higher

proportion of lipophilic MeHg$^+_{(aq)}$ (speciated as MeHgCl$_{(aq)}$) over Hg$^{II}_{(aq)}$ (as HgCl$_{2(aq)}$)

renders the former a more effective toxicant to microbial cells (Mason et al. 1995).

Microorganisms have evolved a wide range of strategies to compete against other microbes

for limited resources (Granato et al. 2019), and our hypothesis that MeHg$^+_{(aq)}$ effectively

originated as an antimicrobial mirrors a similar interpretation for *arsM* encoding for arsenic

methylation (Li et al. 2021). As the *hgcAB* gene pair likely evolved in LUCA (**Figure 1A),**

Hg$^{II}_{(aq)}$ methylation could have stabilised to persist as an early form of antimicrobial

production.

As a defense against $MeHg^+_{(aq)}$, other microorganisms evolved alkylmercury lyase

(MerB) encoded by *merB*, with $Hg^{II}_{(aq)}$ toxicity mitigated via other detoxification systems,

e.g., MerA (Boyd & Barkay 2012), iron-coupled redox reactions (Liu & Wiatrowski 2018),

abiotic reduction (Gu et al. 2011) or other mechanisms (Christakis et al. 2021). Once *merB*

evolved however, *hgcAB* may no longer have offered a selective advantage, and therefore

underwent extensive loss during vertical evolution. Furthermore, we suggest that bacteria

carrying other $MeHg^+_{(aq)}$ detoxification mechanisms might not need to also carry *merB*. For

example, MerB homologs are rarely found in sulfate-reducing bacteria (SRB), as SRB

produce H_2S as a metabolite that reacts with $MeHg^+_{(aq)}$ to form dimethylmercury (DMHg)

which volatilises efficiently from the cell (Baldi et al. 1993). Jonsson et al. (2016) also found

DMHg production resulting from reaction of $MeHg^+_{(aq)}$ with poorly crystalline iron sulfides;

these phases are typically precipitated indirectly by the metabolism of SRB. Other studies

have observed that $MeHg^+_{(aq)}$ is somehow efficiently exported from methylating cells, thus

reducing toxicity to the methylator (Graham et al. 2012; Lin et al. 2015). We recognise,

however, that the mechanisms underpinning $MeHg^+_{(aq)}$ export have not yet been elucidated.

Interestingly, extracellular thiol compounds, such as cysteine, were found to facilitate export

and desorption of $MeHg^+_{(aq)}$ (Lin et al. 2015), possibly due to competitive binding of thiols to

receptors on $MeHg^+_{(aq)}$ transporters. To our knowledge, only two proteins encoded by *mer*

operons, MerC (Sone et al. 2017) and MerE (Sone et al. 2013), have been reported as

potential $MeHg^+_{(aq)}$ transporters, but neither of them is carried by $Hg^{II}_{(aq)}$ methylators.

Our study has revealed an ancient origin for microbial mercury methylation, evolving

from LUCA to radiate extensively throughout the tree of life both vertically, albeit with

extensive loss, and to a lesser extent horizontally. We hypothesise that early mercury

methylating microorganisms may have innovatively transformed a ubiquitous aqueous trace

metal, $Hg^{II}_{(aq)}$, into a highly effective antimicrobial compound. The dual ability of sulfate-

reducing bacteria to methylate mercury and excrete sulfide provides serendipitous protection

against $MeHg^+_{(aq)}$ toxicity. Methylmercury degradation can also be abiotically photo-

catalysed, so it is unlikely that shallow aqueous environments where UV light could penetrate

would have allowed for toxic build-up of $MeHg^+_{(aq)}$. We speculate that anoxic sediments

where volcanogenic mercury and either geogenic or bacteriogenic sulfide were co-present

(such as seafloor sediments near hydrothermal vents) may have provided an ideal

biogeochemical setting for the evolution of microbial mercury methylation. Prior to the

evolution of genomic countermeasures such as *merB*, $MeHg^+_{(aq)}$ concentrations in some

natural environments could have increased to toxic levels beyond those typically found today,

where they are constrained by both biotic and abiotic demethylation mechanisms. We

conclude that microbial $Hg^{II}_{(aq)}$ methylation in extant microorganisms represents the genomic

vestige of an early geosphere-biosphere interaction with major ramifications for Earth's

mercury cycle.

Materials and Methods

Dataset construction

UniProt Reference Proteomes v2022_03 (https://proteininformationresource.org/rps/) at 35%

cutoff (RP35) and 75% cut-off (RP75) were used to establish two datasets. UniProt Reference

Proteomes is a database containing representative proteomes selected from a large proteome

database with more than 30,000 proteomes from UniProtKB (Chen et al. 2011). Every

proteome in the database is comprised of all coding sequences translated from a whole

genome sequence. The RP35 dataset contained 5,535 proteomes and was used to study the

phylogeny of protein family PF03599 (CdhD, CdhE and HgcA) and PF03243 (MerB). The

RP75 dataset contained 17,551 proteomes and was used to study the phylogeny of HgcA

individually. In order to enlarge the diversity and sample size of HgcA sequences, two other

datasets, including one containing 700 representative prokaryotic proteomes constructed by

Moody et al. (2022) and another containing several novel *hgc*-carriers published by Lin et al.

(2021), were retrieved and incorporated into the RP75 dataset, followed by removal of any

redundancies. HgcAB proteins selected for this study including their metadata are listed in

Supplementary Table S1; Other proteins including CdhD, CdhE, and MerB used in this study

are listed in Supplementary Table S2.

Determination of genes of interest

Protein sequences belonging to protein families PF03599 and PF03243 from both datasets

were determined according to annotation by UniProt and also confirmed with hmmsearch

v3.1   (Eddy   1998)   against   curated   HMM   models   provided   by   InterPro

(https://www.ebi.ac.uk/interpro/entry/pfam). HgcA sequences were extracted from the

PF03599 family using hmmsearch against the Hg-MATE database (Capo et al. 2022) and

further determined by the conserved motif N(V/I)WC(A/S). CdhD and CdhE sequences were

extracted from the PF03599 family using blastp v2.11.0 against the experimentally validated

CdhD (Q57577) and CdhE (Q57576) sequences from *Methanocaldococcus jannaschii* JAL-1,

respectively. HgcB sequences were determined by searching for 4Fe4S proteins encoded by

genes adjacent to *hgcA* genes (two open reading frames on either side of *hgcA*) in the genome.

MerB sequences were further confirmed from the PF03243 protein family by hmmsearch

against the MerB database constructed by Christakis et al. (2021).

Reconstruction of phylogenetic trees

Protein sequences belonging to the PF03599 protein family and included in the RP35 dataset

were aligned using MAFFT-linsi v7.453 (Katoh & Standley 2013), resulting in an alignment

containing 478 sequences with 2922 columns. The alignment was used to build a maximum

likelihood (ML) tree using IQ-TREE v2.0.3 (Schmidt et al. 2014) under the best-fitting model

of LG+C50+F+R. Branch supports were estimated with 1000 ultrafast bootstrap (Hoang et al.

2018) replicates (**Figure 1**) The unrooted tree was rooted using three methods: Midpoint-

rooting (Swofford et al. 1996), Minimum Variance (MinVar, Mai et al. 2017), and Minimal

Ancestor Deviation (MAD, Tria et al. 2017), respectively. Similar to the methods described

above, all trees in this study were built using IQ-TREE v2.0.3 (Schmidt et al. 2014) with 1000

ultrafast bootstrap (Hoang et al. 2018) replications. Protein sequences were aligned using

MAFFT (L-INS-i) v7.453 (Katoh & Standley 2013).

To mitigate potential contamination of HgcB tail sequences in the HgcA alignment,

the fused HgcAB proteins were aligned separately and the 'HgcB tail' (last 79 positions) was

manually removed. The resulting 'HgcA head' of the fused HgcAB proteins was aligned with

other HgcA sequences. Finally, aligned 'HgcB tail' sequences were subsequently added back

to the alignment. The first 327 positions of the alignment were poorly aligned, therefore

manually removed. The final alignment contained 169 sequences with 493 columns and was

used to infer a ML tree under the LG+C60+F+G model (**Figure 2**). The tree was rerooted

between the two major groups of HgcA sequences according to the above PF03599 protein

family tree.

HgcB sequences and 'HgcB tail' sequences from the fused HgcAB proteins were

aligned and used to infer a ML tree using a similar method as the PF03599 protein family tree

described above under the LG+C60+F+G model. Phylogenetic congruence between the HgcA

and the HgcB sequences was inferred and visualized using cophylo implemented in the R

package phytools v1.0-3 (Revell 2012) based on the topology of the HgcA and the HgcB ML

trees described above. Nodes of both HgcA and HgcB trees were allowed to be rotated by the

program to optimize vertical matching of tips (**Figure 3**).

In total, 223 MerB sequences were predicted from proteomes in the RP35 dataset.

These sequences were aligned and used to build a ML tree under the best-fit protein model

LG+C40+F+G chosen according to BIC (**Figure 4**).

A species tree of *hgc*+ and *merB*+ proteomes (251 proteomes in total) from the RP35

dataset was reconstructed based on the 27 marker genes proposed by Moody et al. (2022). 27

HMM profiles were created individually based on marker gene alignments and concatenated

using HMMER v3.1. The 27 marker homologs in the 251 proteomes were extracted by

hmmsearch against the concatenated hmm profile with an E-value of 1e-10. The 27 marker

homologs were then aligned respectively and concatenated. Poorly aligned regions were

trimmed using trimAl v1.2 (Silla-Martínez et al. 2009) with parameters "-resoverlap 0.55 -

seqoverlap 60 -automated1". The resulting supermatrix (8024 sites) was used to infer a ML

tree under the model of LG+F+R10 (**Figure 5**). All phylogenetic trees described above were

visualized using iTOL v6 (Letunic & Bork 2021) and refined with Adobe Illustrator (Adobe

Systems Inc., San Jose, CA, USA).

Data availability

All data generated in this study including amino acid alignments and phylogenetic trees are

deposited in Figshare: https://doi.org/10.6084/m9.figshare.21428523

Acknowledgments

We thank Yao-ban Chan and Qiuyi Li at The University of Melbourne for valuable

discussions and advice in the early stage of this study. We also acknowledge helpful

discussions with Caitlin Gionfriddo, Ben Peterson, and Eric Capo. This publication was made

possible in part through support from a Strategic Australian Postgraduate Award (PhD

scholarship) to H.L. from the Environmental Microbiology Research Initiative (EMRI) at The

University of Melbourne (J.W.M.) and laboratory start-up funding from the University of

Glasgow to J.W.M., the Moore Foundation (https://doi.org/10.37807/GBMF9741 to T.A.W.),

the Royal Society (URF\R\201024) to T.A.W., and the John Templeton Foundation (62220 to

T.A.W. and E.R.R.M.). The opinions expressed in this publication are those of the author(s)

and do not necessarily reflect the views of the John Templeton Foundation.

References

Adam PS, Borrel G, Gribaldo S. 2018. Evolutionary history of carbon monoxide dehydrogenase/acetyl-CoA synthase, one of the oldest enzymatic complexes. Proc. Natl. Acad. Sci. 115:E1166–E1173. doi: 10.1073/pnas.1716667115.

Amyot, M., Morel, F.M. and Ariya, P.A., 2005. Dark oxidation of dissolved and liquid elemental mercury in aquatic environments. *Environmental science & technology*, *39*(1), pp.110-114.

Baldi, F., Pepi, M. and Filippelli, M., 1993. Methylmercury resistance in Desulfovibrio desulfuricans strains in relation to methylmercury degradation. *Applied and environmental microbiology*, *59*(8), pp.2479-2485.

Barkay T, Gu B. 2022. Demethylation─the other side of the mercury methylation coin: A critical review. ACS Environ. Au. 2:77–97. doi: 10.1021/acsenvironau.1c00022.

Barkay T, Kritee K, Boyd E, Geesey G. 2010. A thermophilic bacterial origin and subsequent constraints by redox, light and salinity on the evolution of the microbial mercuric reductase. Environ. Microbiol. 12:2904–2917. doi: 10.1111/j.1462-2920.2010.02260.x.

Barkay T, Miller SM, Summers AO. 2003. Bacterial mercury resistance from atoms to ecosystems. FEMS Microbiol Rev. 27:355–384. doi: 10.1016/s0168-6445(03)00046-9.

Barkay T, Wagner⬜Döbler I. 2005. Microbial Transformations of Mercury: Potentials, Challenges, and Achievements in Controlling Mercury Toxicity in the Environment. In: Advances in Applied Microbiology Volume 57.Advances in Applied Microbiology pp. 1–52.

Bobrowski, N., Hönninger, G., Galle, B. and Platt, U., 2003. Detection of bromine monoxide in a volcanic plume. *Nature*, *423*(6937), pp.273-276.

Bolotin E, Hershberg R. 2016. Bacterial intra-species gene loss occurs in a largely clocklike manner mostly within a pool of less conserved and constrained genes. Sci. Rep. 6:35168. doi: 10.1038/srep35168.

Boyd E, Barkay T. 2012. The mercury resistance operon: from an origin in a geothermal environment to an efficient detoxification machine. Front. Microbiol. 3. doi: 10.3389/fmicb.2012.00349.

Capo E et al. 2022. A consensus protocol for the recovery of mercury methylation genes from
metagenomes. Mol. Ecol. Resour. 1–15. doi: 10.1111/1755-0998.13687.

Chen C et al. 2011. Representative proteomes: a stable, scalable and unbiased proteome set
for sequence analysis and functional annotation. PLOS ONE. 6:e18910. doi:
10.1371/journal.pone.0018910.

Choi S-C, Chase T, Bartha R. 1994. Metabolic pathways leading to mercury methylation in
Desulfovibrio desulfuricans LS. Appl. Environ. Microbiol. 60:4072–4077. doi:
10.1128/aem.60.11.4072-4077.1994.

Christakis CA, Barkay T, Boyd ES. 2021. Expanded diversity and phylogeny of mer genes
broadens mercury resistance paradigms and reveals an origin for mera among thermophilic
archaea. Front. Microbiol. 12. doi: 10.3389/fmicb.2021.682605.

Colombo, M.J., Ha, J., Reinfelder, J.R., Barkay, T. and Yee, N., 2013. Anaerobic oxidation of
Hg (0) and methylmercury formation by Desulfovibrio desulfuricans ND132. *Geochimica et
Cosmochimica Acta*, *112*, pp.166-177.
413
Date SS et al. 2019. Kinetics of enzymatic mercury methylation at nanomolar concentrations
catalyzed by HgcAB. Appl Env. Microbiol. doi: 10.1128/AEM.00438-19.

Eddy SR. 1998. Profile hidden Markov models. Bioinformatics. 14:755–763. doi:
10.1093/bioinformatics/14.9.755.

Ekstrom EB, Morel FMM, Benoit JM. 2003. Mercury methylation independent of the acetyl-
coenzyme A pathway in sulfate-reducing bacteria. Appl Env. Microbiol. 69:5414–5422. doi:
10.1128/aem.69.9.5414-5422.2003.

Gilmour CC et al. 2013. Mercury methylation by novel microorganisms from new
environments. Env. Sci Technol. 47:11810–20. doi: 10.1021/es403075t.

Gilmour CC et al. 2011. The sulfate-reducing bacterium *Desulfovibrio desulfuricans* ND132
as a model for understanding bacterial mercury methylation. Appl Env. Microbiol. 77:3938–
51. doi: 10.1128/AEM.02993-10.

Gilmour CC, Bullock AL, McBurney A, Podar M, Elias DA. 2018. Robust mercury
methylation across diverse methanogenic *Archaea*. MBio. 9. doi: 10.1128/mBio.02403-17.

Gionfriddo CM et al. 2016. Microbial mercury methylation in Antarctic sea ice. Nat
Microbiol. 1:16127. doi: 10.1038/nmicrobiol.2016.127.

Graham AM et al. 2012. Detailed Assessment of the Kinetics of Hg-Cell Association, Hg
Methylation, and Methylmercury Degradation in Several Desulfovibrio Species. Appl.
Environ. Microbiol. 78:7337–7346. doi: 10.1128/AEM.01792-12.

Granato ET, Meiller-Legrand TA, Foster KR. 2019. The evolution and ecology of bacterial
warfare. Curr. Biol. 29:R521–R537. doi: 10.1016/j.cub.2019.04.024.

Grasby SE, Them TR, Chen Z, Yin R, Ardakani OH. 2019. Mercury as a proxy for volcanic
emissions in the geologic record. Earth-Sci. Rev. 196:102880. doi:
10.1016/j.earscirev.2019.102880.

Gu B et al. 2011. Mercury reduction and complexation by natural organic matter in anoxic
environments. Proc. Natl. Acad. Sci. 108:1479–1483. doi: 10.1073/pnas.1008747108.

Guo, M. and Korenaga, J., 2021. A halogen budget of the bulk silicate Earth points to a
history of early halogen degassing followed by net regassing. *Proceedings of the National*
Academy of Sciences, *118*(51), p.e2116083118.
443

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. 2018. UFBoot2: Improving
the ultrafast bootstrap approximation. Mol. Biol. Evol. 35:518–522. doi:
10.1093/molbev/msx281.

Hsu-Kim H, Kucharzyk KH, Zhang T, Deshusses MA. 2013. Mechanisms regulating mercury
bioavailability for methylating microorganisms in the aquatic environment: a critical review.
Env. Sci Technol. 47:2441–56. doi: 10.1021/es304370g.

Jonas RB, Gilmour CC, Stoner DL, Weir MM, Tuttle JH. 1984. Comparison of methods to
measure acute metal and organometal toxicity to natural aquatic microbial communities. Appl.
Environ. Microbiol. 47:1005–1011.

Jonsson S, Mazrui NM, Mason RP. 2016. Dimethylmercury Formation Mediated by
Inorganic and Organic Reduced Sulfur Surfaces. Sci. Rep. 6:27958. doi: 10.1038/srep27958.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol. Biol. Evol. 30:772–780. doi:
10.1093/molbev/mst010.

Kidd KA et al. 2012. Biomagnification of mercury through lake trout (*Salvelinus namaycush*)
food webs of lakes with different physical, chemical and biological characteristics. Sci. Total
Environ. 438:135–143. doi: 10.1016/j.scitotenv.2012.08.057.

Koskiniemi S, Sun S, Berg OG, Andersson DI. 2012. Selection-driven gene loss in bacteria.
PLOS Genet. 8:e1002787. doi: 10.1371/journal.pgen.1002787.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. Nucleic Acids Res. 49:W293–W296. doi: 10.1093/nar/gkab301.

Li YP et al. 2021. Antimicrobial Activity of Metals and Metalloids. Annu. Rev. Microbiol.
75:175–197. doi: 10.1146/annurev-micro-032921-123231.

Lin H et al. 2021. Mercury methylation by metabolically versatile and cosmopolitan marine
bacteria. ISME J. 15:1810–1825. doi: 10.1038/s41396-020-00889-4.

Lin H, Lu X, Liang L, Gu B. 2015. Thiol-facilitated cell export and desorption of
methylmercury by anaerobic bacteria. Environ. Sci. Technol. Lett. 2:292–296. doi:
10.1021/acs.estlett.5b00209.

Liu S, Wiatrowski HA. 2018. Reduction of Hg(II) to Hg(0) by biogenic magnetite from two
magnetotactic bacteria. Geomicrobiol. J. 35:198–208. doi: 10.1080/01490451.2017.1362076.

Lu X et al. 2016. Anaerobic mercury methylation and demethylation by *Geobacter*
bemidjiensis Bem. Env. Sci Technol. 50:4366–73. doi: 10.1021/acs.est.6b00401.

Mai U, Sayyari E, Mirarab S. 2017. Minimum variance rooting of phylogenetic trees and
implications for species tree reconstruction. PLOS ONE. 12:e0182238. doi:
10.1371/journal.pone.0182238.

Mason, R.P., Reinfelder, J.R. and Morel, F.M., 1995. Bioaccumulation of mercury and
methylmercury. *Water, Air, and Soil Pollution*, *80*, pp.915-921.
481
McDaniel EA et al. 2020. Expanded phylogenetic diversity and metabolic flexibility of
mercury-methylating microorganisms. mSystems. 5. doi: 10.1128/mSystems.00299-20.

Meixnerová, J., Blum, J.D., Johnson, M.W., Stüeken, E.E., Kipp, M.A., Anbar, A.D. and
Buick, R., 2021. Mercury abundance and isotopic composition indicate subaerial volcanism
prior to the end-Archean "whiff" of oxygen. *Proceedings of the National Academy of*
Sciences, 118(33), p.e2107511118.
488
Moody ER et al. 2022. An estimate of the deepest branches of the tree of life from ancient
vertically evolving genes Perry, GH & Koonin, EV, editors. eLife. 11:e66695. doi:
10.7554/eLife.66695.

Obrist, D., Tas, E., Peleg, M., Matveev, V., Faïn, X., Asaf, D. and Luria, M., 2011. Bromine-
induced oxidation of mercury in the mid-latitude atmosphere. *Nature Geoscience*, *4*(1), pp.22-
26.
495
Parks JM et al. 2013. The genetic basis for bacterial mercury methylation. Science. 339:1332–
5. doi: 10.1126/science.1230667.

Podar M et al. 2015. Global prevalence and distribution of genes and microorganisms
involved in mercury methylation. Sci Adv. 1:e1500675. doi: 10.1126/sciadv.1500675.

Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other
things). Methods Ecol. Evol. 3:217–223. doi: 10.1111/j.2041-210X.2011.00169.x.

Robinson, J.B. and Tuovinen, O., 1984. Mechanisms of microbial resistance and
detoxification of mercury and organomercury compounds: physiological, biochemical, and
genetic analyses. *Microbiological reviews*, *48*(2), pp.95-124.
505
Schmidt HA, Minh BQ, von Haeseler A, Nguyen L-T. 2014. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 32:268–
274. doi: 10.1093/molbev/msu300.

Seigneur C, Lohman K. 2008. Effect of bromine chemistry on the atmospheric mercury cycle.
J. Geophys. Res. Atmospheres. 113. doi: 10.1029/2008JD010262.

Siciliano SD, Lean DRS. 2002. Methyltransferase: An enzyme assay for microbial
methylmercury formation in acidic soils and sediments. Environ. Toxicol. Chem. 21:1184–
1190. doi: 10.1002/etc.5620210610.

Silla-Martínez JM, Capella-Gutiérrez S, Gabaldón T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 25:1972–1973. doi:
10.1093/bioinformatics/btp348.

Sone Y et al. 2017. A novel role of MerC in methylmercury transport and phytoremediation

of methylmercury contamination. Biol. Pharm. Bull. 40:1125–1128. doi: 10.1248/bpb.b17-
00213.

Sone Y et al. 2013. Increase methylmercury accumulation in Arabidopsis thaliana expressing
bacterial broad-spectrum mercury transporter MerE. AMB Express. 3:52. doi: 10.1186/2191-
0855-3-52.

Sousa FL, Martin WF. 2014. Biochemical fossils of the ancient transition from geoenergetics
to bioenergetics in prokaryotic one carbon compound metabolism. Biochim. Biophys. Acta
BBA - Bioenerg. 1837:964–981. doi: 10.1016/j.bbabio.2014.02.001.

Svetlitchnaia T, Svetlitchnyi V, Meyer O, Dobbek H. 2006. Structural insights into
methyltransfer reactions of a corrinoid iron–sulfur protein involved in acetyl-CoA synthesis.
Proc. Natl. Acad. Sci. 103:14331–14336. doi: 10.1073/pnas.0601420103.

Swofford DL, Olsen GJ, Waddell PJ, Hillis DM. 1996. Phylogenetic inference. In: Molecular
systematics.Vol. 15 Sinauer Associates, Inc.: Sunderland, Massachusetts pp. 407–514.

Tada Y, Marumoto K, Takeuchi A. 2020. *Nitrospina*-like bacteria are potential mercury
methylators in the mesopelagic zone in the East China Sea. Front Microbiol. 11. doi:
10.3389/fmicb.2020.01369.

Tada Y, Marumoto K, Takeuchi A, Gralnick JA. 2021. *Nitrospina*-like Bacteria Are
Dominant Potential Mercury Methylators in Both the Oyashio and Kuroshio Regions of the
Western North Pacific. Microbiol. Spectr. 9:e00833-21. doi: doi:10.1128/Spectrum.00833-21.

Tebo BM et al. 2004. Biogenic manganese oxides: properties and mechanisms of formation.
Annu. Rev. Earth Planet. Sci. 32:287–328. doi: 10.1146/annurev.earth.32.101802.120213.

Tezuka, T. and Tonomura, K., 1976. Purification and Properties of an Enzyme Catalyzing the
Splitting of Carbon-Mercury Linkages from Mercury-Resistant Pseudomonas K-62 Strain: 1.
Splitting Enzyme 1. *The Journal of Biochemistry*, *80*(1), pp.79-87.
542
Trevors JT. 1986. Mercury methylation by bacteria. J. Basic Microbiol. 26:499–504. doi:
10.1002/jobm.3620260811.

Tria FDK, Landan G, Dagan T. 2017. Phylogenetic rooting using minimal ancestor deviation.
Nat. Ecol. Evol. 1:1–7. doi: 10.1038/s41559-017-0193.

Villar E, Cabrol L, Heimburger-Boavida LE. 2020. Widespread microbial mercury
methylation genes in the global ocean. Env. Microbiol Rep. doi: 10.1111/1758-2229.12829.

Wang J, Dai J, Chen G, Jiang F. 2022. Role of sulfur biogeochemical cycle in mercury
methylation in estuarine sediments: A review. J Hazard Mater. 423:126964. doi:
https://doi.org/10.1016/j.jhazmat.2021.126964.

Wang, P., Su, S., Xiang, J., You, H., Cao, F., Sun, L., Hu, S. and Zhang, Y., 2014. Catalytic
oxidation of Hg0 by MnOx–CeO2/γ-Al2O3 catalyst at low temperatures. *Chemosphere*, *101*,
pp.49-54
555
Weiss MC et al. 2016. The physiology and habitat of the last universal common ancestor. Nat.
Microbiol. 1:1–8. doi: 10.1038/nmicrobiol.2016.116.

Weiss MC, Preiner M, Xavier JC, Zimorski V, Martin WF. 2018. The last universal common
ancestor between ancient Earth chemistry and the onset of genetics. PLoS Genet.
14:e1007518. doi: 10.1371/journal.pgen.1007518.

Zerkle AL et al. 2021. Sulfur and mercury MIF suggest volcanic contributions to Earth's
atmosphere at 2.7 Ga. doi: 10.7185/geochemlet.2124.

Figure Legends

Figure 1. Phylogenetic tree of the protein family PF03599. (A) Unrooted tree of the protein

family PF03599. The tree was inferred by using the Maximum Likelihood method under

LG+C50+F+R model. This analysis involved 478 amino acid sequences with a total of 2922

positions in the alignments. Different taxonomies are represented by different colors.

Ultrafast bootstrap support values were calculated with 1000 replications, and ultrafast

bootstrap values > 90% are shown by black dots at the nodes. (B) PF03599 tree rooted by the

midpoint. Clades whose average branch length distance to their leaves are below 1.5 are

collapsed with iTOL for better visualization.

Figure 2. Phylogenetic tree of HgcA proteins. The tree was inferred by using the Maximum

Likelihood method under LG+C60+F+G model. This analysis involved alignment of 169

amino acid sequences with a total of 493 positions. Different taxonomies are represented by

different colors. Ultrabootstrap support values were calculated with 1000 replications, and

ultrabootstrap values > 90% are shown by black dots at the nodes. Experimentally validated

functional and non-functional HgcA sequences are labelled by solid and hollow stars. Fused-

HgcAB sequences are indicated by a grey background.

Figure 3. Co-phylogenetic tree of HgcA and HgcB. The phylogeny of HgcA shown on the

left is the same as Figure 2. The phylogeny of HgcB shown on the right was inferred by using

the Maximum Likelihood method under LG+C60+F+G model. Lines between the two trees

connect HgcA (left) and HgcB (right) proteins from the same proteome, illustrating the degree

of congruency in their respective phylogenetic associations. This analysis involved alignment

of 169 amino acid sequences with a total of 322 positions.   Taxonomies of the HgcB

sequences are represented in the same colors as shown in the HgcA phylogeny. Ultrabootstrap

support values were calculated with 1000 replications, and ultrabootstrap values > 90% are

shown by black dots at the nodes. Gray shaded blocks describe fused-HgcAB and 'HgcB tail'

genes in the two trees, respectively. The green dots at the tip of the HgcA tree represent the

corresponding *hgcA* without downstream *hgcB*, and the red triangle represents the

corresponding fused-*hgcAB* with another *hgcB* downstream.

Figure 4. Phylogenetic tree of MerB proteins. The tree was inferred by using the Maximum

Likelihood method under LG+C40+F+G model. This analysis involved alignment of 223

amino acid sequences with a total of 1448 positions. Different taxonomies are represented by

different colors. Ultrabootstrap support values were calculated with 1000 replications, and

ultrabootstrap values > 90% are shown by black dots at the nodes.

Figure 5. Species tree of *hgcA*+ and *merB*+ genomes in the RP35 dataset. The tree was

inferred from a concatenate of 27 marker genes derived from Moody et al. (2022) by using

the Maximum Likelihood method under LG+F+R10 model. This analysis involved alignment

of 251 amino acid sequences with a total of 8024 positions. Different taxonomies are

represented by different colors. The presence of genes *hgcA*/fused *hgcAB*/*hgcB*/*merB* in the

genomes were represented by different symbols and colors in the outer circle. Ultrabootstrap

support values were calculated with 1000 replications, and ultrabootstrap values > 90% are

shown by black dots at the nodes.

Small clade

Fused-HgcAB

Thermoplasmatota   *Methanomassiliicoccus*

Euryarchaeota
FCB group   *Marinimicrobia*

Euryarchaeota   Stenosarchaea group
Methanomicrobia

Archaeal clade

Large superclade

Deltaproteobacteria   *Syntrophaceae*

FCB group   *Bacteroidetes*

Firmicutes   Clostridia
Negativicutes

Deltaproteobacteria   Geobacteraceae
Desulfarculaceae

Deltaproteobacteria   Desulfuromonadaceae
Desulfovibrionaceae
Desulfobulbaceae
Desulfohalobiaceae

Bacterial clade

Tree scale: 1

Asgard   Thorarchaeota
Lokiarchaeota

Asgard   Heimdallarchaeota

Nitrospinae   *Nitrospina*

Euryarchaeota

Terrabacteria   *Actinobacteria*

Alphaproteobacteria   *Breoghania*
Deltaproteobacteria

Asgard

Aminicenantes

Woesebacteria
Thermotogae
Spirochaetes

Euryarchaeota   *Thermococci*

Small clade

Fused-HgcAB

★ Experimentally validated functional

☆ Experimentally validated non-functional

Taxonomy

Archaea
- Euryarchaeota
- Asgard group
- Thermoplasmatota

Bacteria
- Nitrospinae
- Deltaproteobacteria
- Terrabacteria
- PVC group
- Nitrospirae
- Spirochaetes
- FCB group
- Alphaproteobacteria
- Other Bacteria

Rooting Point

Fused HgcAB

HgcA Tree

HgcB Tree

Tree scale: 1

Taxonomy
- Euryarchaeota
- Asgard group
- Thermoplasmatota
- Nitrospinae
- Deltaproteobacteria
- Terrabacteria
- PVC group
- Nitrospirae
- Spirochaetes
- FCB group
- Alphaproteobacteria
- Other Bacteria

— Associations

● *hgcA* without downstream *hgcB*

▲ Fused-*hgcAB* with a downstream *hgcB*

Tree scale: 1

Taxonomy

- Euryarchaeota
- TACK group
- FCB group
- Deltaproteobacteria
- Gammaproteobacteria
- Betaproteobacteria
- Alphaproteobacteria
- Firmicutes
- Chloroflexi
- Actinobacteria
- Acidobacteria
- Other Bacteria