Repurposing the mammalian RNA-binding protein Musashi-1 as an allosteric translation repressor in bacteria

Roswitha Dolcemascolo[1,2,#], María Heras-Hernández[1,#], Lucas Goiriz[1,3,#], Roser Montagud-Martínez[1,2], Alejandro Requena-Menéndez[1], Raúl Ruiz[1], Anna Pérez-Ràfols[4,5], R. Anahí Higuera-Rodríguez[6,7], Guillermo Pérez-Ropero[8,9], Wim F. Vranken[10,11], Tommaso Martelli[4], Wolfgang Kaiser[6], Jos Buijs[8,12], and Guillermo Rodrigo[1,*]

[1]Institute for Integrative Systems Biology (I2SysBio), CSIC – University of Valencia, 46980 Paterna, Spain

[2]Department of Biotechnology, Polytechnic University of Valencia, 46022 Valencia, Spain

[3]Department of Applied Mathematics, Polytechnic University of Valencia, 46022 Valencia, Spain

[4]Giotto Biotech SRL, 50019 Sesto Fiorentino, Italy

[5]Magnetic Resonance Center (CERM), Department of Chemistry Ugo Schiff, Consorzio Interuniversitario Risonanze Magnetiche di Metalloproteine (CIRMMP), University of Florence, 50019 Sesto Fiorentino, Italy

[6]Dynamic Biosensors GmbH, 82152 Planegg, Germany

[7]Department of Physics, Technical University of Munich, 85748 Garching, Germany.

[8]Ridgeview Instruments AB, 75237 Uppsala, Sweden

[9]Department of Chemistry – BMC, Uppsala University, 75123 Uppsala, Sweden.

[10]Structural Biology Brussels, Vrije Universiteit Brussel, 1050 Brussels, Belgium

[11]Interuniversity Institute of Bioinformatics in Brussels, Université Libre de Bruxelles – Vrije Universiteit Brussel, 1050 Brussels, Belgium

[12]Department of Immunology, Genetics, and Pathology, Uppsala University, 75185

26   Uppsala, Sweden

27   #Equal contribution to this work

28   *Correspondence: guillermo.rodrigo@csic.es

30   **Keywords:** Binding kinetics; Dynamic systems; Genetic circuits; Mathematical modeling;

31   Post-transcriptional regulation; RNA recognition motif; Synthetic biology.

ABSTRACT

The RNA recognition motif (RRM) is the most common RNA-binding protein domain identified in nature. However, RRM-containing proteins are only prevalent in eukaryotic phyla, in which they play central regulatory roles. Here, we engineered an orthogonal post-transcriptional control system of gene expression in the bacterium *Escherichia coli* with the mammalian RNA-binding protein Musashi-1, which is a stem cell marker with neurodevelopmental role that contains two canonical RRMs. In the circuit, Musashi-1 is regulated transcriptionally and works as an allosteric translation repressor thanks to a specific interaction with the N-terminal coding region of a messenger RNA and its structural plasticity to respond to fatty acids. We fully characterized the genetic system at the population and single-cell levels showing a significant fold change in reporter expression, and the underlying molecular mechanism by assessing the *in vitro* binding kinetics and *in vivo* functionality of a series of RNA mutants. The dynamic response of the system was well recapitulated by a bottom-up mathematical model. Moreover, we applied the post-transcriptional mechanism engineered with Musashi-1 to specifically regulate a gene within an operon, to implement combinatorial regulation, and to reduce protein expression noise. This work illustrates how RRM-based regulation can be adapted to simple organisms, thereby adding a new regulatory layer in prokaryotes for translation control.

INTRODUCTION

Gene regulation at the post-transcriptional level is pervasive in living organisms of ranging complexity [1-4]. Indeed, the ability to regulate the genetic information flow at different points appears instrumental to maximize the integration of intrinsic and extrinsic signals, which enables an efficient information processing by the organisms. However, the solutions implemented in prokaryotes and eukaryotes greatly differ. In prokaryotes, small RNAs (sRNAs) regulate messenger RNA (mRNA) stability and translation initiation [1], supported by a series of RNA-binding proteins (*e.g.*, Hfq) that act globally [2]. Regulatory proteins of specific scope in these simple organisms mainly operate in the transcriptional layer [5], what is aligned with the models presented in the early times of molecular biology [6]. By contrast, eukaryotes deploy a sizeable number of RNA-binding proteins with a variety of functions [4] that participate in the regulation of mRNA turnover, transport, splicing, and translation in a gene-specific manner and also at a global scale. In animals, in particular, most RNA-binding proteins contain RNA recognition motifs (RRMs) [7]. RRMs are small globular domains of about 90 amino acids that fold into four antiparallel β-strands and two α-helices, which can bind to single-strand RNAs with sufficient affinity and specificity to control biological processes [8]. Yet, while important to attain functional diversity in the post-transcriptional layer in animals, RRMs are not prevalent in all organisms. In fact, the scarcity of RRM-containing proteins in prokaryotes and the often-unknown functional role of those identified by bioinformatic methods [9] question if RRMs can readily work in organisms with much simpler gene expression machinery and intracellular organization. If so, this would raise the potential to use RRM-RNA interactions as an orthogonal layer to engineer gene regulation in prokaryotes.

To address these intriguing questions, we adopted a synthetic biology approach where a specific RRM-containing protein was incorporated in a bacterium in order to engineer a post-transcriptional control module. Synthetic biology has highlighted how living

cells can be (re)programmed through the assembly of independent genetic elements into functional networks for a variety of applications in biotechnology and biomedicine [10]. Yet, synthetic biology can also be used to disentangle natural systems and to probe hypotheses about biological function [11]. In previous work, some proteins with the ability to recognize RNA have been exploited as translation factors in bacteria for a gene-specific regulation [12-14]. The first instance was the tetracycline repressor protein (TetR), which naturally functions as a transcription factor, by means of the selection of synthetic RNA aptamers [12]. The bacteriophage MS2 coat protein (MS2CP) [13] and eukaryotic Pumilio homology domains [14] were also used in synthetic circuits. Alternatively, a wide palette of post-transcriptional control systems based on sRNAs have been developed in recent years to program gene expression in bacteria [15]. Of note, these systems are amenable to be combined with regulatory proteins to attain complex dynamic behaviors [16]. A heterologous RRM-containing protein with definite regulatory activity, in addition to provide empirical evidence on the adaptability of such RNA-binding domains to different genetic backgrounds, would enlarge the synthetic biology toolkit [17], boosting applications in which high orthogonality, expression fine-tuning, and signal integrability are required features. In addition, RRMs can themselves be allosterically regulated, opening up new avenues for post-transcriptional regulation by small molecules.

Still, there are instances of bacterial proteins that regulate translation in a gene-specific manner, such as CsrA to control glycogen biosynthesis [18] or the ribosomal protein S8 to exert self-repression [19]. Besides, it is worth noting that some bacteriophages follow this mechanism to modulate their infection cycle. These are the cases, *e.g.*, of the coat proteins of the phages MS2 (infecting *Escherichia coli*) or PP7 (infecting *Pseudomonas aeruginosa*), which regulate the expression of the cognate phage replicases through protein-RNA interactions [20]. However, one limitation for synthetic biology developments is that such phage proteins are not allosteric. At the post-

transcriptional level, bacteria mostly rely on a large palette of *cis-* and *trans*-acting non-

coding RNAs to either activate or repress protein expression, resulting in the regulation of

translation initiation, mRNA stability, or transcription termination, and even allowing

sensing small molecules [1,15]. Thus, there should be efforts to replicate this functional

versatility with proteins.

In this work, the mammalian RNA-binding protein Musashi-1 (MSI-1) [21] was used

as a translation repressor in the bacterium *E. coli* (**Fig. 1a**). MSI-1 belongs to an

evolutionarily conserved family of RRM-containing proteins, of which a member was first

identified in *Drosophila melanogaster* [22]. MSI-1 contains two RRMs in the N-terminal

region (RRM1 and RRM2) and recognizes the RNA consensus sequence RU_nAGU on the

nanomolar affinity scale [23]. Importantly, MSI-1 can be allosterically inhibited by fatty

acids (in particular, 18-22-carbon ω-9 monounsaturated fatty acids) [24]. In mammals,

MSI-1 is mainly expressed in stem cells of neural and epithelial lineage and plays crucial

roles in differentiation, tumorigenesis, and cell cycle regulation [21]. Notably, MSI-1

regulates Notch signaling by repressing the translation of a key protein in the pathway [23].

Hence, rather than moving genetic elements from simple to complex organisms, as it is

normally done (*e.g.*, the TetR-aptamer module was implemented in simple eukaryotes

[25]), we reversed the path by moving an important mammalian gene (from *Mus musculus*)

to *E. coli*. Some eukaryotic factors have already been implemented in bacteria to regulate

gene expression at different levels [14,26], but the case of RRM-containing proteins has

remained elusive. In the following, we present quantitative experimental and theoretical

results on the response dynamics of a synthetic gene circuit in which MSI-1 works as an

allosteric translation repressor. There, MSI-1 is transcriptionally controlled by the lactose

repressor protein (LacI), and translation regulation by MSI-1 is accomplished by means of

a specific interaction with an mRNA (encoding a reporter protein) that harbors a suitable

binding motif in its N-terminal coding region.

RESULTS

A Musashi protein can down-regulate translation in bacteria. From the amino acid sequence of *M. musculus* MSI-1, we generated a nucleotide sequence with codons optimized for *E. coli* expression. Knowing that the C-terminus of MSI-1 is of low structural complexity [27], we cloned a truncated version of the gene encompassing the first 192 amino acids, which include the two RRMs, to implement our synthetic circuit (**Fig. 1b**). The resulting protein (termed MSI-1*) was expressed from a synthetic PL-based promoter repressed by LacI (termed PLlac) [28] lying in a high copy number plasmid. This allowed controlling the expression of the heterologous RNA-binding protein at the transcriptional level with lactose or isopropyl β-D-1-thiogalactopyranoside (IPTG) in a genetic background over-expressing LacI. As a regulated element, we used the superfolder green fluorescent protein (sfGFP) [29], which was expressed from a constitutive promoter (J23119) lying in a low copy number plasmid (**Suppl. Fig. 1**). An RNA motif obtained by affinity elution-based RNA selection (SELEX) containing two copies of the consensus recognition sequence (*viz.*, GUUAGU and AUUUAGU) [23] was placed in frame after the start codon of *sfGFP*. This motif folds into a stem-loop structure that allows stabilizing the exposure of the recognition sequence to the solvent. In this way, MSI-1* can repress translation by blocking the binding of the ribosome, presumably by imposing a steric hindrance for the 30S ribosomal subunit. This mode of action differs from the natural one in mammals, in which MSI-1 binds to the 3' untranslated region (UTR) of its target mRNA (*Numb*) to repress translation by disrupting the activation function of the poly(A)-binding protein [30]. Here, considering lactose (or IPTG) and oleic acid as the two inputs and sfGFP as the output, MSI-1* being an internal allosteric regulator operating at the post-transcriptional level, an IMPLY gate would model the logic behavior of the resulting circuit (*i.e.*, sfGFP would only turn off with lactose and without oleic acid in the medium).

We first characterized by bulk fluorometry the dose-response curve of the system

using a lactose concentration gradient up to 1 mM. Our data show that MSI-1* down-regulated sfGFP expression by 2.5-fold (**Fig. 1c**). Fitting a Hill equation, we obtained a regulatory coefficient of 99 µM (lactose concentration at which the repression is half of the maximal) and a Hill coefficient of 1.7 (**Suppl. Note 1**). We also observed that IPTG (a synthetic compound) triggered a very similar response. To further inspect the activity of the RNA-binding protein, we filtered out the transcriptional regulatory effect. For that, we expressed the enhanced blue fluorescent protein 2 (eBFP2) [31] from the PLlac promoter to obtain the corresponding dose-response curve with lactose. In this way, eBFP2 expression was a proxy of MSI-1* expression, which allowed representing the transfer function of the engineered regulation (**Fig. 1d**). A Hill equation with no cooperative binding (*i.e.*, Hill coefficient of 1) explained the data with sufficient agreement, suggesting that only one protein interacted with a given mRNA (*i.e.*, each RRM of MSI-1* binds to a consensus sequence repeat, in agreement with a previous structural model [27]). We also measured the cell growth rate for all induction conditions, finding that the values were almost constant. This indicates that the expression of the mammalian protein did not produce a significant burden to the bacterial cell.

In simple terms, protein expression comes from the product of the transcription and translation rates of the gene. Hence, we examined such a decomposition in the case of sfGFP expression regulated by MSI-1*. Of note, the low copy number plasmid harbors an additional transcriptional unit to express the monomeric red fluorescent protein mScarlet [32] from a constitutive promoter (J23119). We then monitored its expression profile with lactose. Assuming that *sfGFP* and *mScarlet* were equally transcribed, as they were expressed from the same promoter, and that the translation rate of *mScarlet* was constant, the product of mScarlet expression and cell growth rate was considered a proxy of the transcription rate of *sfGFP*. Moreover, the ratio of sfGFP and mScarlet expressions was a proxy of the translation rate of *sfGFP* [33]. This served us to represent the dynamics of the

system in a plane defined as translation rate *vs.* transcription rate (termed Crick space [34]), highlighting that the change in sfGFP expression with lactose comes indeed from translation regulation (**Fig. 1e**). A reverse transcription quantitative polymerase chain reaction (RT-qPCR) was used to confirm the preservation of the *sfGFP* mRNA level (**Suppl. Fig. 2**). Finally, to evaluate the heterogeneity of the response within a bacterial population, we performed single-cell measurements of sfGFP expression by flow cytometry. Unimodal distributions able to shift in response to lactose were observed (**Fig. 1f**). Setting a threshold to categorize expression, we found that the percentage of cells in the ON state dropped from 87% to 15% upon addition of 1 mM lactose. In sum, our results show that MSI-1* can regulate translation in a specific manner in *E. coli*, and hence that eukaryotic regulators can be borrowed to be functional elements in prokaryotes.

Mechanistic insight into the engineered regulation based on a protein-RNA interaction. We then introduced a series of point mutations into the SELEX RNA motif to assess their effect over the regulatory activity of the RRM-containing protein (**Fig. 2a**). These mutations change the consensus recognition sequence of at least one repeat. A characterization of all systems revealed that the mutations affected both the maximal level and fold change of sfGFP expression (**Fig. 2b**). Of note, a single point mutation in one repeat leading to RU_nCGU (mutant 1) was quite detrimental for the MSI-1*-based regulation (only 1.4-fold reduction in sfGFP expression). Despite that mutation substantially reduced sfGFP expression in absence of MSI-1*, the presumed repressed state upon addition of lactose did not change much, suggesting the difficulty of the protein for targeting the mutated mRNA. This agrees with the prior observation that, within the consensus sequence, UAG is a minimal core that determines the specific recognition by MSI-1 [35]. A double point mutation changing the minimal cores of the two repeats (UAC rather than UAG; mutant 5) also resulted in a detrimental action, but not to a greater

extent. We also engineered a new reporter system with a minimal RNA motif consisting of

a single copy of the shortest possible consensus sequence (AUAGU), but its

characterization showed no apparent regulation by MSI-1* (**Suppl. Fig. 3**). Taken

together, two copies of the consensus sequence seem necessary for a successful

regulation of protein expression.

To relate the cellular effects with protein-RNA interactions, we obtained a purified

MSI-1* preparation in order to perform *in vitro* binding kinetics assays (**Suppl. Fig. 4**). For

that, a gene coding for a truncated version of the human MSI-1 was expressed from a T7

polymerase promoter in *E. coli*. With respect to the *M. musculus* version, this protein only

differs in one residue of RRM2 (then termed MSI-1$_h$*), which is the subsidiary domain for

RNA recognition (note also that the human and mouse proteins recognize the same

consensus sequence [35]). To avoid the necessity of labelling the molecules of interest

and allow working with very low amounts of protein and RNA, we used the switchSENSE

technology, which allows measuring molecular dynamics on a chip (**Fig. 2c**) [36]. **Fig. 2d**

summarizes the resulting protein-RNA association and dissociation rates (k_{ON} and k_{OFF},

respectively; see also **Suppl. Fig. 5**). In the case of the original RNA motif, we found an

association rate of 1.1 nM^{-1}min^{-1}, which means that a single regulator molecule would take

1-3 min to find its target in the cell, and a residence time of the protein on the RNA of 1.5

min (given by 1/k_{OFF}). Of note, the reported value of k_{ON} is relatively close to the upper limit

imposed by the diffusion rate (~1 nM^{-1}s^{-1}). This fast rate suggests that MSI-1* is able to

find its target mRNA in *E. coli*, competing with ribosomes and ribonucleases, and then

achieve translation regulation. We also found that a single mutation in one of the two UAG

minimal cores (mutants 1 and 2) led to similar association but faster dissociation (almost 4

times faster dissociation), whereas a double mutation affecting the two cores (mutant 5)

disturbed both phases (almost 15 times slower association and 10 times faster

dissociation). The dissociation constant ($K_D = k_{OFF}/k_{ON}$) was 0.62 nM for the original

system, while 87 nM for mutant 5. The switchSENSE technology allowed revealing that

affinity on the subnanomolar scale, refining a previous estimate of 4 nM obtained by gel

shift assays [23]. To contextualize these values, we compared to the binding kinetics of

MS2CP, a phage RNA-binding protein that has evolved in a prokaryotic context and that

we recently exploited to study how expression noise emerges and propagates through

translation regulation [37]. Previous work disclosed an association rate to the cognate

RNA motif of 0.032 $nM^{-1}min^{-1}$ and a residence time of 12 min, leading to a dissociation

constant of 2.6 nM [38]. Thus, MSI-1* would target RNA faster than MS2CP, but once this

happened the phage protein would remain bound longer.

Next, we tried to predict the impact of the mutations on sfGFP expression. On the

one hand, we used an empirical free-energy model (RBS calculator) to obtain an estimate

of the mRNA translation rate from the sequence [39]. However, only a poor correlation (R^2

= 0.16) with the maximal expression level was observed (**Fig. 2e**), suggesting that

additional variables should be considered. For example, it was surprising the higher

expression level in the case of mutant 4, despite a minimal change in the structure of the

RNA motif (**Suppl. Fig. 5a**; we ensured that *sfGFP* was in frame in this case). On the other

hand, when the fold change was correlated with the inverse of the dissociation constant

(1/K_D, *i.e.*, the equilibrium constant) better results were obtained (R^2 = 0.75; **Fig. 2f**).

Mutant 1 is illustrative in this case because, even though a fast association rate was

preserved (1.6 $nM^{-1}min^{-1}$), it displayed a marginal regulatory activity as a result of a shorter

residence time (0.41 min). This indicates that the underlying protein-RNA interaction in the

bacterial circuit was close to thermodynamic equilibrium.

A mathematical model captured the dynamic response of the system. Translation

regulation is more challenging than transcription regulation because mRNA is unstable

compared to DNA, especially in bacteria. In *E. coli*, in particular, the average mRNA half-

life is about 5 min [40]. However, it is possible to derive a common mathematical

framework from which to analyze the dynamics of both regulatory modes (**Fig. 3a**). The

fold change in protein expression is a suitable mesoscopic parameter that is directly

related to the kinetic parameters that characterize the interaction in the cell [41]. Using

mass action kinetics, we obtained a general mathematical description of the fold change

as a function of the regulator concentration (R), the association and dissociation rates, the

leakage fraction of RNA/peptide-chain elongation, and the nucleic acid degradation rate

(**Suppl. Note 2**). To visualize the impact of the different parameters, we represented the

fold change equation as a heatmap. When there is no nucleic acid degradation (DNA), a

linear dependence between the first-order association rate ($k_{ON}R$) and k_{OFF} is established

to maintain a given fold change value (**Fig. 3b**), which would correspond to the case of

transcription regulation. Accordingly, our model converges to the classical description of

fold = $1 + R/K_D$. However, if the nucleic acid degrades quickly (mRNA), the dependence

between the first-order kinetic rates becomes nonlinear (**Fig. 3c**). Indeed, in the case of

translation regulation, it is important to note that when $k_{ON}R$ is lower than the mRNA

degradation rate (*i.e.*, the mRNA is degraded faster than the protein binds), the

functionality is greatly compromised. To overcome this barrier, the regulator needs to be

highly expressed, as MSI-1* is in our system (we estimate $R > 1$ µM with 1 mM lactose).

Furthermore, when the residence time is much longer than the mRNA half-life (*i.e.*, the

mRNA is degraded before the protein unbinds), K_D is not a suitable parameter to

characterize the regulation, which is solely association-dependent, resulting in non-

equilibrium thermodynamics [42]. According to the aforementioned kinetic rates, this would

be the case for MS2CP, but not for MSI-1* (*i.e.*, both k_{ON} and k_{OFF} are instrumental to

describe the regulation exerted by MSI-1*). Furthermore, given the 2.5-fold down-

regulation in our system, we estimated an elongation leakage fraction of 40% (using the

fold change equation in the limit $R \to \infty$). This leakage would come from the ability of

ribosomes to elongate even if MSI-1* is bound and their ability to bind sooner to the *sfGFP*

mRNA due to a conserved transcription-translation coupling mechanism [43].

In addition, we studied the transient response of the gene circuit with lactose, as

both MSI-1* and sfGFP expressions changed with time. For that, we quantified the total

red fluorescence of the cell population (**Fig. 3d**), which is an estimate of the total number

of cells, and the total green fluorescence (**Fig. 3e**), which comes from the composition of

population growth and gene regulation. We developed a bottom-up mathematical model

based on differential equations to predict sfGFP expression in the cell (**Suppl. Note 3**), as

well as a phenomenological model for the bacterial growth (**Suppl. Note 4**). The

parameter values were adjusted with the curves without and with 1 mM lactose. Then, we

used the mathematical model to predict the transient responses for different intermediate

lactose concentrations, finding excellent agreement with the experimental data ($R^2 = 0.98$;

Fig. 3f). We also characterized the time-course response of the circuit with IPTG,

encountering similar results (**Suppl. Fig. 6**). Moreover, to explore the maintenance of the

regulatory behavior when the cell physiology changes, we characterized cells growing in

solid medium with a repurposed LigandTracer technology, which initially was developed to

monitor molecular interactions in real time [44]. In this case, a significant difference in the

total red fluorescence was observed without and with 1 mM IPTG, suggesting that MSI-1*

expression was costly for the cell in these conditions. Besides, the total green

fluorescence of the growing population was recapitulated using the model with a 2.6-fold

down-regulation of cellular sfGFP expression, which is in tune with the results in liquid

medium (**Suppl. Fig. 7**). Subsequently, we analyzed the intracellular response. The time-

dependent ratio of total green and red fluorescence was used as a proxy of sfGFP

expression. A delay in the response is expected because MSI-1* needs to be produced

upon addition of lactose [45]. Nevertheless, our model predicted a faster response than

experimentally observed (**Fig. 3g**). Overall, this quantitative inspection of translation

regulation backs connections between molecular attributes and cellular behavior.

Rational redesign of the targeted transcript to enhance the dynamic range of the

response. The presence of stem-loop structures in the N-terminal coding region

contributes to lower the expression level. The more stable and closer to the start codon,

the greater the impact on expression [46]. We hypothesized that, by destabilizing the RNA

motif for MSI-1 binding, we would obtain an alternative regulatory system with higher

expression levels. Accordingly, a new reporter system was engineered removing three

base pairs from the stem, maintaining the two consensus recognition sequences. An

experimental analysis revealed a 4.9-fold increase of the maximal sfGFP expression level

and a 2.0-fold down-regulation with 1 mM lactose (**Fig. 4a**, redesign 1). We then

investigated the possibility of increasing the dynamic range of the response by placing

three consecutive RNA motifs. However, we did not observe a greater down-regulation

with 1 mM lactose (**Fig. 4a**, redesign 2), suggesting that the additional motifs far away

from the start codon had no effect; what was noticed is an effect on the maximal

expression level.

As a further strategy to enhance the dynamic range of the response, we redesigned

the 5' UTR of *sfGFP* to accommodate two additional RU_nAGU repeats (*viz.*, GUUUAGU

and AUUUAGU) flanking the ribosome binding site (RBS), maintaining the original RNA

motif after the start codon. In this way, MSI-1* can also block the RNA component of the

30S ribosomal subunit. Indeed, this is a widespread post-transcriptional regulatory strategy

in prokaryotes, as it happens *e.g.* with the MS2 phage replicase [20]. It is worth to note

that the new 5' UTR remained unstructured. We characterized by bulk fluorometry the

dose-response curve of this new system, revealing an 8.6-fold down-regulation of sfGFP

expression by MSI-1* (**Fig. 4b**, redesign 3; see also **Suppl. Fig. 8** to appreciate the tight

control of MSI-1* expression with the PLlac promoter). This was a substantial increase in

performance with respect to the 2.5-fold down-regulation of the system shown in **Fig. 1b**.

Fitting a Hill equation, we obtained a regulatory coefficient of 86 µM and a Hill coefficient of

4.5 (**Suppl. Note 1**). While the regulatory coefficient was similar than in the original system

(99 µM), the Hill coefficient was significantly higher (compared to 1.7). Interestingly, an

apparent cooperativity was established between two MSI-1* proteins by binding to

adjacent sites. The dynamics of the system was also represented in the Crick space to

highlight the change in translation rate. At the single-cell level, we found a 91% of ON cells

in the uninduced state that decreased to 5.3% with 1 mM lactose (**Fig. 4c**). Taken

together, our data present MSI-1* as a powerful heterologous translation regulator in

bacteria.

The regulatory activity of a Musashi protein in bacteria can be externally controlled

by a fatty acid. The ability of proteins to respond to small molecules is instrumental for

environmental and metabolic sensing. Previous work revealed that MSI-1 can be

allosterically inhibited by ω-9 monounsaturated fatty acids and, in particular, by oleic acid

[24], an 18-carbon fatty acid naturally found in various animal and plant oils (*e.g.*, olive oil).

Oleic acid binds to the RRM1 domain of MSI-1 and induces a conformational change that

prevents RNA recognition (**Fig. 5a**). To gain insight about the interactions between the

elements of our system, we performed gel electrophoresis mobility shift assays using the

purified MSI-1* protein, the RNA motif as a label-free sRNA molecule, and oleic acid. The

different mobility of the nucleic acids upon binding to proteins and the coincident staining

capacity of nucleic and fatty acids were exploited. We confirmed the MSI-1*-RNA

interaction using a protein concentration gradient in this *in vitro* set up (**Suppl. Fig. 9a**),

and we found that the interaction was completely disrupted in presence of 1 mM oleic acid

(**Fig. 5b**). Furthermore, using an oleic acid concentration gradient, we obtained a half-

maximal effective inhibitory concentration of about 0.5 mM (**Suppl. Fig. 9b**).

Subsequently, we assessed the effect of oleic acid over the regulatory activity of

MSI-1* expressed in *E. coli*. This bacterium has evolved a machinery to uptake fatty acids

from the environment. FadL and FadD are two membrane proteins that act as

transporters, and FadE is the first enzyme that processes the fatty acid via the β-oxidation

cycle [47]. Because of the high turbidity of the cell culture observed in presence of oleic

acid, we characterized the system by single-cell measurements of sfGFP expression by

flow cytometry. In the case of the original system, the percentage of cells in the ON state

increased from 10% (with 1 mM lactose) to 49% upon addition of 20 mM oleic acid (**Fig.**

5c; see the 2D probability-based histograms in **Suppl. Fig. 10**). However, the initial 93% of

ON cells observed in absence of lactose was not recovered. Arguably, oleic acid was

partially degraded once it entered the cell. Nevertheless, the system implemented with the

redesign-3 reporter displayed a better dynamic behavior in response to lactose and oleic

acid. In particular, the percentage of cells in the ON state increased from almost 0 (with 1

mM lactose) to 71% upon addition of 20 mM oleic acid (**Fig. 5d**; see also **Suppl. Fig. 10**).

In addition, we investigated this allosteric regulation by imaging the fluorescence of

bacterial colonies grown in solid medium with different inducers. In stationary phase, FadE

and the rest of oxidative enzymes could be saturated with the fatty acids generated from

the membrane degradation [48], oleic acid then having more time to interact with MSI-1*.

Notably, we found a substantial inhibition of the repressive action of MSI-1* with 20 mM

oleic acid in the case of both systems (**Fig. 5e,f**; see also **Suppl. Fig. 11**). Conclusively,

these results illustrate how the plasticity of RRM-containing proteins (*e.g.*, MSI-1) can be

exploited to engineer, even in simple organisms, gene regulatory circuits that operate in an

integrated way at the transcriptional, translational, and post-translational levels.

Application of a Musashi protein for intra-operon, combinatorial, and noise

regulation. Transcription regulation has been engineered in *E. coli* to end with purposeful

and versatile gene expression programs [26,49]. However, this type of control faces limitations, such as to regulate a specific gene within an operon or to implement a definite combinatorial regulation without a large screening of promoter variants. To show that MSI-1* is instrumental to address these issues and ultimately increase our ability to program gene expression (**Fig. 6a**), a new regulatory circuit was engineered in which *sfGFP* and *mScarlet* were both forming a single transcriptional unit (*i.e.*, bicistronic operon) under a synthetic PL-based promoter regulated by the tetracycline repressor protein (TetR; promoter termed PLtet) [28]. This allowed controlling the expression of both fluorescent proteins at the transcriptional level with anhydrotetracycline (aTC) in a genetic background over-expressing TetR. Furthermore, an RNA motif for MSI-1 binding was placed in front of *sfGFP* (**Fig. 6b**). A characterization by bulk fluorometry using lactose (1 mM) and aTC (100 ng/mL) in a combinatorial way showed the specific regulation of sfGFP expression by MSI-1* and the ability to combine signals exploiting transcription and translation regulation (**Fig. 6c**; implementation with the redesign-3 motif due to its enhanced dynamic range). A NIMPLY gate would model the logic behavior of the resulting circuit (*i.e.*, sfGFP would only turn on with aTC and without lactose in the medium). These data also excluded the possibility that MSI-1* operated transcriptionally as a result of spurious DNA targeting.

In addition, we analyzed how MSI-1* regulated noise in protein expression monitoring green fluorescence in single cells. Inducing the circuit of **Fig. 6b** with 100 ng/mL aTC and 1 mM lactose produced almost the same mean expression level than with an intermediate aTC concentration (15 ng/mL when the implementation was with the original motif and 30 ng/mL when it was with the redesign-3 motif). However, the resulting unimodal distributions displayed different dispersions, lower when MSI-1* was not repressed. The Fano factor (the ratio between variance and mean) [50] was used to quantify the responses, finding reductions of 35% and 65% depending on the implementation (**Fig. 6d**). Furthermore, for the circuit of **Fig. 1b**, we found a 38% lower

Fano factor when inducing with 1 mM lactose and 20 mM oleic acid than with 0.1 mM

lactose, despite having similar mean expression levels (**Suppl. Fig. 12**). Of note, the

response sensitivity was dominated by transcription regulation when the PL-based

promoter was induced with an intermediate concentration of lactose (0.1 mM) or aTC (15-

30 ng/mL). By contrast, the response sensitivity was dominated by translation regulation

when the PL-based promoter was fully induced (1 mM lactose or 100 ng/mL aTC), thereby

controlling the heterogeneity of the response [37]. Overall, these results illustrate the utility

of repurposed mammalian RNA-binding proteins in bacteria for a fine expression control.

DISCUSSION

The successful incorporation of the mammalian MSI-1 protein as a translation factor in *E. coli* highlights, in first place, the versatility of RRM-containing proteins to function as specific post-transcriptional regulators in any living cell, from prokaryotes to eukaryotes. Our data show that the protein-RNA association phase is very fast, which is suitable for regulation even in cellular contexts in which RNA molecules are short-lived, such as in *E. coli* [40]. Nonetheless, it is important to stress that the kinetic parameters *in vivo* might differ from those measured *in vitro* due to off-target bindings and crowding effects [51]. Moreover, our data show that a down-regulation of translation rate up to 8.6-fold can be achieved, with an appropriate design of the target mRNA leader region, and that the engineered cell can sense oleic acid from the environment. Here, the C-terminal low-complexity domain of the native MSI-1 was discarded to create MSI-1* [27], in order to increase solubility, even though this domain might contribute to RNA binding [52]. Further work should be conducted to enhance the fold change of the regulatory module and engineer complex circuits with it.

Interestingly, proteins associated with clustered regularly interspaced short palindromic repeats (CRISPR), which belong to the prokaryotic immune system, contain distorted RRM versions [53]. Some CRISPR proteins might have evolved, for example, from an ancestral RRM-based (palm) polymerase after duplications, fusions, and diversification. Noting that the palm domain indeed presents an RRM-like fold [54], we hypothesize that a boost of functionally diverse RRM-containing proteins took place once the polymerases were confined into the nucleus, as the pressure for efficient replication was relieved in the cytoplasm, which would provide a rationale on the unbalance noticed between eukaryotes and prokaryotes [7,55].

In second place, our results pave the way for engineering more complex circuits in bacteria with plastic and orthogonal RNA-binding proteins, such as MSI-1, capable of

signal multiplexing. Nature is a formidable reservoir of functional genetic material sculpted

by evolution that can be exploited to (re)program specific living cells [10]. However, to

overcome biological barriers, transgenes usually come from related organisms or cognate

parasites, at the cost of limiting the potential engineering. Therefore, efforts to borrow

functional elements from highly diverse organisms are suggestive (*e.g.*, regulatory proteins

from mammals to bacteria), with the ultimate goal of developing industrial or biomedical

applications.

Notably, advances in synthetic biology have pushed the bioproduction of a wide

variety of compounds in bacteria as a result of a better ability to fine tune enzyme

expression [56]. Translation regulation is instrumental to this end because in multiple

cases different enzymes are expressed from the same transcriptional unit (*i.e.*, operon).

Previous work exploited regulatory RNAs for such a tuning [57], but the use of RNA-

binding proteins as translation factors is also appealing. We envision the application of

MSI-1* as a genetic tool for metabolic engineering. The additional use of RNA-binding

proteins able to alter mRNA stability might lead to the implementation of more complex

circuits at the post-transcriptional level. Furthermore, MSI-1* is able to respond to fatty

acids, which are ideal precursors of potential biofuels due to their long hydrocarbon

chains. In particular, biofuel in the form of fatty acid ethyl ester, whose bioproduction in *E.*

coli can be optimized by reengineering the regulation of the β-oxidation cycle with the

allosteric transcription factor FadR [58]. Arguably, MSI-1* might be used in place of or in

combination with FadR for subsequent developments. However, engineering regulatory

circuits for efficient bioproduction is not evident in general as the enzymatic expression

levels may require fine tuning, so systems-level mathematical models need to be

considered for design along with a wide genetic toolkit for implementation [56]. We

anticipate that other animal RRM-containing proteins might be repurposed in *E. coli* as

translation factors. Moreover, protein design might be used to reengineer MSI-1* in order

to respond to new ligands, maintaining high specificity and affinity for a particular RNA

sequence, as previously done with the transcription factor LacI [59].

In addition, the Musashi protein family is of clinical importance, as in humans it is

involved in different neurodegenerative disorders (*e.g.*, Alzheimer's disease) and some

types of cancer [21,60,61]. Therefore, the development of simple genetic systems from

which to test protein mutants, potential target mRNAs, decoying RNA aptamers, and

inhibitory small molecules in a systematic manner is very relevant. Furthermore, isolating

human regulatory elements would help to filter out indirect effects that likely occur in the

natural context. This might lead to new therapeutic opportunities. Nevertheless, one

limitation of using *E. coli* as a chassis is that some post-translational modifications (PTMs)

may be lost, thereby compromising the functionality of the expressed proteins [62].

Fortunately, there are metabolic engineering efforts devoted to implement eukaryotic PTM

pathways in *E. coli*, such as the glycosylation pathway [63].

In conclusion, the functionalization of RRM-containing proteins in bacteria offers

exciting prospects, especially as more information becomes available on how individual

RRM domains bind to precise RNA sequences, interact with further protein domains, and

respond to small molecules through allosteric effects. This work illustrates how synthetic

biology, through the rational assembly of heterologous genes and designer *cis*-regulatory

elements into circuits, is useful to generate knowledge about the application range of a

fundamental type of proteins in nature.

METHODS

Strains, plasmids, and reagents. *E. coli* Dh5α was used for cloning purposes following standard procedures. To express our genetic circuit for functional characterization, *E. coli* MG1655-Z1 cells ($lacI^+$, $tetR^+$) were used. This strain was co-transformed with two plasmids, called pRM1+ (KanR, pSC101-E93R ori; leading to ~230 copies/cell) [64] and pREP6 (CamR, p15A ori; leading to ~15 copies/cell). On the one hand, pRM1+ was obtained by cloning a truncated coding region of the *M. musculus* MSI-1 protein (the first 192 amino acids, containing the two RRMs; UniProt #Q61474; termed MSI-1*). This gene was under the transcriptional control of the inducible promoter PLlac. On the other hand, pREP6 was obtained by cloning the coding region of sfGFP with an RNA sequence motif recognized by MSI-1. The coding region of mScarlet was also present in the plasmid. These two genes were under the control of the constitutive promoter J23119 in two different transcriptional units. In addition to the original RNA sequence motif, five point-mutated sequences were designed and cloned in pREP6. Additional RNA sequence motifs were cloned in front of *sfGFP* for control experiments (the resulting plasmids were named pREP4, pREP4b, pREP4b3x, and pREP7). In particular, pREP4b3x incorporates three RNA motifs in tandem after the start codon, and pREP7 has two RU_nAGU repeats flanking the RBS and a full RNA motif after the start codon. Additional reporter plasmids were constructed using the inducible promoter PLtet to assess the intra-operon regulation, the implementation of combinatorial regulation, and the buffering of expression noise (the resulting plasmids were named pREP6α and pREP7α). Suitable genetic cassettes to obtain the final constructions were synthesized by IDT. **Suppl. Table 1** lists all plasmids used in this work. **Suppl. Table 2** presents the nucleotide sequences of the different genetic elements.

To perform the dynamic assays with LigandTracer (Ridgeview), *E. coli* BL21(DE3) cells ($lacI^+$, $T7pol^+$) were used. This strain was also co-transformed with pRM1+ and

pREP6. To purify a recombinant Musashi protein, *E. coli* BL21-Gold(DE3) cells (*lacI*[+],

T7pol[+]) were used. A truncated coding region of the human MSI-1 protein (the first 200

amino acids; UniProt #O43347; termed MSI-1$_h$*) was cloned under the control of a T7pol

promoter into the plasmid pET29b (KanR, pUC ori).

Luria-Bertani (LB) medium was used for the overnight cultures and M9 minimal

medium (1X M9 minimal salts, 2 mM MgSO$_4$, 0.1 mM CaCl$_2$, 0.05% thiamine, 0.05%

casamino acids, 1% glycerol or 0.4% glucose) for the characterization cultures. M9-

glucose medium was only used for real-time fluorescence quantification in liquid medium

with IPTG. LB-agar was used for real-time fluorescence quantification in solid medium.

Kanamycin and chloramphenicol were used at a concentration of 50 μg/mL and 34 μg/mL,

respectively. Lactose and IPTG were used as the inducers of the system (controlling the

expression of MSI-1* in *E. coli*) at a concentration of 5, 10, 20, 50, 100, 200, 500, or 1000

μM. aTC was also used to induce the modified systems with PLtet at a concentration of 15,

30, or 100 ng/mL. Oleic acid was used as the allosteric inhibitor of MSI-1* at a

concentration of 20 mM in the *in vivo* assays (both in liquid and solid medium). In the *in*

vitro assays, oleic acid was used at a concentration of 0.01, 0.1, 0.2, 0.5, 0.7, 1, 1.5, or 2

mM. It was neutralized with NaOH and used in a medium containing 0.5% tergitol NP-40.

Compounds provided by Merck.

Bulk fluorometry. Cultures (2 mL) inoculated from single colonies (three replicates) were

grown overnight in LB medium with shaking (220 rpm) at 37 ºC. Cultures were then diluted

1:100 in fresh M9 medium (200 μL) with the appropriate inducer (lactose, IPTG, and/or

aTC). The microplate (96 wells, black, clear bottom; Corning) was incubated with shaking

(1300 rpm) at 37 ºC up to 8-10 h (to reach an OD$_{600}$ around 0.5-0.7). At different times, the

microplate was assayed in a Varioskan Lux fluorometer (Thermo) to measure absorbance

(600 nm), green fluorescence (excitation: 485 nm, emission: 535 nm), and red

fluorescence (excitation: 570 nm, emission: 610 nm). To characterize the time-course response of the system, cultures were grown to exponential phase and then diluted before adding the inducer (to minimize the response lag). Mean background values of absorbance and fluorescence, corresponding to M9 medium, were subtracted to correct the signals. Normalized fluorescence was calculated as the slope of the linear regression between fluorescence and absorbance (assuming fluorophore maturation faster than cell doubling time and no proteolytic degradation) [65]. The mean value of normalized fluorescence corresponding to non-transformed cells was then subtracted to obtain a final estimate of expression. In addition, cell growth rate was calculated as the slope of the linear regression between the logarithm of background-subtracted absorbance and time in the exponential phase.

Real-time fluorescence quantification in solid medium. Cultures (2 mL) inoculated from single colonies (three replicates) were grown overnight in LB medium with shaking (220 rpm) at 37 °C. The overnight culture was plated (15 µL) in areas A and D of a MultiDish 2x2 plate (Ridgeview) coated with LB-agar. IPTG was added in areas A and B of the dish at the final concentration of 1 mM. Area C was kept free of cells/inducers as a reference. The dish was then placed in the rotating support of the LigandTracer instrument (Ridgeview) and incubated at 37 °C for 24 h. The fluorescence from sfGFP and mScarlet was quantified with time in the seeded areas of the dish using the BlueGreen (excitation: 488 nm, emission: 535 nm) and OrangeRed (excitation: 568 nm, emission: 620 nm) detectors. The readouts of the opposite parts of the dish were subtracted to correct the signals.

Flow cytometry. Cultures (2 mL) inoculated from single colonies (three replicates) were grown overnight in LB medium with shaking (220 rpm) at 37 °C. Cultures were then diluted

1:100 in fresh LB medium (200 µL) to load a microplate (96 wells, black, clear bottom;

Corning) with the appropriate concentrations of lactose (0, 100, 1000 µM), oleic acid (0, 20

mM), and/or aTC (15, 30, 100 ng/mL). The microplate was then incubated with shaking

(1300 rpm) at 37 °C until cultures reached a sufficient OD_{600}. Cultures (6 µL) were then

diluted in PBS (1 mL). Fluorescence was measured in an LSRFortessa flow cytometer

(BD) using a 488 nm laser and a 530 nm filter for green fluorescence. Events were gated

by using the forward and side scatter signals and compensated (~10^4 events after this

process). The mean value of the autofluorescence of the cells was subtracted to obtain a

final estimate of expression. Data analysis performed with MATLAB (MathWorks).

Purification of a Musashi protein. Cells were grown in LB medium with shaking at 37 °C

until OD_{600} reached 0.6-0.8. Subsequently, the expression of MSI-1_h* was induced with 0.5

mM IPTG. Cells were incubated at 37 °C for 4 h and harvested by centrifugation at 7500

rpm for 15 min at 4 °C. The cell pellet was resuspended in a lysis buffer (50 mM Tris-HCl,

pH 8.0, 500 mM NaCl, 10% glycerol, with protease inhibitor cocktail), ruptured by

sonication, and separated by centrifugation at 30,000 rpm for 35 min at 4 °C. The soluble

fraction was collected and treated with a 5% polyethylenimine solution in order to remove

DNA/RNA attached to the protein. Resuspension of the protein was done in 20 mM Tris-

HCl, pH 9.0, with protease inhibitor cocktail. Soluble protein was filtered with a 0.22 µm

membrane and purified by ion exchange chromatography using an Anion exchange Q FF

16/10 column previously equilibrated in alkaline buffer. The protein was collected on the

flow-through. The protein was filtered and further purified to homogeneity by size exclusion

chromatography using a Hi load 26/60 Superdex 75 pg column previously equilibrated in

alkaline buffer with NaCl. The purified fractions were collected and buffer exchange

chromatography was performed using a HiPrep 26/10 Desalting column previously

equilibrated with the final buffer (20 mM MES, pH 6.0, 100 mM NaCl, 0.5 mM EDTA, with

protease inhibitor cocktail). Purification performed at Giotto.

Binding kinetics assays of protein-RNA interactions. Binding experiments of the

purified MSI-1$_h$* protein against different RNA ligands were performed using the

switchSENSE proximity sensing technology [36,66] and a suitable adapter chip on the

heliX biosensor platform (Dynamic Biosensors). The adapter chip consists of a microfluidic

channel with two gold electrodes functionalized with fluorophore-decorated DNA

nanolevers that serve as linkers between the gold surface and the ligand of interest. A

constant negative voltage is applied to the electrodes to keep the DNA nanolevers in an

upright position. Binding between the injected analyte (MSI-1$_h$*) and the ligand attached to

the sensor surface (RNA) leads to the alteration of the chemical surrounding of the dye,

which results in a fluorescence change. Fluorescence change of the dye in real time

describes the binding kinetics of the molecule of interest. Kinetic experiments consisted of

a protein association phase (5 min) and a dissociation phase (15 min) in which the chip

was rinsed with a buffer (50 mM Tris-HCl, 0.5 mM EDTA, 140 mM NaCl, 0.05% Tween 20,

1 mM TCEP, pH 7.2). A flow rate of 100 µL/min was applied and a sampling rate of 1 Hz

was used.

Six different RNA ligands (original and 5 mutants) were attached to the 5' end of a

generic 48 nt DNA ligand strand, which is part of the DNA linker system on the heliX

adapter chip surface. All oligonucleotides were synthesized by Ella Biotech. The ligand

strand was hybridized with an adapter strand carrying the fluorophore. Different

fluorophores were tested towards their sensitivity for protein-RNA interactions. The green

fluorophore Gb showed the most significant signal change. The other half of the adapter

strand is complementary to a DNA anchor strand, which is pre-attached to the chip

surface. The immobilization of the RNA used a standard functionalization procedure on the

heliX device. Kinetic rate constants and affinities were obtained by fitting the experimental

data with theoretical binding models implemented in the heliOS software (Dynamic Biosensors). Exponential decay models were used. As a negative control to check for unspecific protein-RNA binding, the single-strand RNA sequence CGGCGCCGC was used (without any binding motif). All data were referenced with a blank run and with the negative control.

RT-qPCR. Cultures (2 mL) inoculated from single colonies (three replicates) were grown overnight in LB medium with shaking (220 rpm) at 37 °C. Cultures were then diluted 1:100 in fresh LB medium (2 mL) with the appropriate inducer (lactose) and were grown until OD_{600} reached 0.6-0.8. 500 µL of each culture was mixed with RNAprotect Bacteria Reagent (Qiagen). Subsequently, RNA extraction was carried out with the RNeasy kit (QIAGEN), choosing the enzymatic lysis and proteinase K digestion of bacteria (recommended for Gram-negative bacteria grown in complex media). The eluted RNA sampled were quantified using a NanoDrop spectrophotometer (Thermo).

The TaqPath 1-step RT-qPCR master mix, CG was used. 1 µL of sample was mixed with 500 nM of forward and reverse primers, 250 nM of ssDNA probe, and 5 µL of the master mix for a total volume of 20 µL (adjusted with RNase-free water) in a fast microplate (Applied). Two independent mixes were prepared, one for targeting *sfGFP* and another for the *E. coli* b3500 gene, which was employed as the reference gene. Reactions were performed in a QuantStudio 3 equipment (Thermo) with this protocol: incubation at 25 °C for 2 min for uracil-N glycosylation, followed by 50 °C for 15 min for RT, followed by an inactivation step at 95 °C for 2 min, then followed by 40 cycles of amplification at 95 °C for 3 s and 60 °C for 30 s.

Gel electrophoresis. Mobility shift assays with the purified MSI-1$_h$* protein and its cognate RNA motif were performed. The RNA motif was generated by *in vitro* transcription

with the TranscriptAid T7 high yield transcription kit (Thermo) from a DNA template. It was then purified using the RNA clean and concentrator column (Zymo) and quantified in a NanoDrop spectrophotometer (Thermo). Bovine serum albumin (BSA) was used as a control protein (at 30 µM). Reactions with different combinations of elements were prepared (MSI-1$_h$* at 45 µM, RNA at 11 µM, and oleic acid at 1 mM). Reactions with concentration gradients of MSI-1$_h$* (from 0 to 45 µM) and oleic acid (from 0 to 2 mM) were also performed. Reactions were incubated for 30 min at 37 ⁰C. Reaction volumes were then loaded in 3% agarose gels prepared with 0.5X TBE and stained using RealSafe (Durviz). Gels ran for 45 min at room temperature applying 110 V. The GeneRuler ultra-low range DNA ladder (10-300 bp, Thermo) was used. This staining served to reveal the RNA and oleic acid (free or in complex with the MSI-1$_h$* protein) [67,68]. In addition, gels were soaked for 10 min in the Coomassie blue stain (Fisher) at room temperature with shaking to reveal the proteins. Gels were then soaked in a destaining solution overnight to remove the excess of blue stain. Pictures were taken with the Imager2 gel documentation system (VWR).

Microscopy. LB-agar plates seeded with *E. coli* MG1655-Z1 cells co-transformed with pRM1+ and pREP6 or pREP7 were grown overnight at 37 ⁰C. Lactose (1 mM) and oleic acid (20 mM) were used as supplements. The plates were irradiated with blue light and images were acquired with a 2.8 Mpixel camera with a filter for green fluorescence in a light microscope (Leica MSV269). The commercial software provided by Leica was used to adjust the visualization of the differential fluorescence among plates. The fluorescence intensity of the colonies was quantified with Fiji [69].

Mathematical modeling. On the one hand, Hill equations were used to empirically model sfGFP expression with lactose/IPTG, eBFP2 expression with lactose, and sfGFP

expression with eBFP2 expression (see **Suppl. Note 1** for details). On the other hand, a

system of ordinary differential equations was developed to model the dynamic response of

the synthetic gene circuit from a bottom-up approach. The system accounted for the

intracellular mRNA and protein concentrations, considering a scenario of equilibrium to

model both LacI-DNA and MSI-1*-RNA binding (see **Suppl. Note 3** for details). Parameter

values were obtained by nonlinear fitting against our experimental data.

Molecular visualization *in silico*. The RMM1 of MSI-1 protein structure determined by

nuclear   magnetic   resonance   was   downloaded   from   the   UniProt   database

(www.uniprot.org) [70]. A 3D structure of the RNA motif subsequence involving the two

RU_nAGU repeats was predicted with the RNAComposer software [71]. The oleic acid

molecule was downloaded from the ChemSpider database (www.chemspider.com). All the

molecules were loaded, visualized, colored, trimmed (where necessary), and manually

docked using the open source PyMol software (Schrödinger; pymol.org).

Resources availability. The sequences of all genetic elements used in this work are

presented in the Supplementary Material. Plasmids available upon request to the

corresponding author.

ACKNOWLEDGEMENTS

We thank M. Sattler (TUM) for useful discussions. This work was supported by the grants H2020-MSCA-ITN-2018 #813239 (RNAct) from the European Commission and PGC2018-101410-B-I00 (SYSY-RNA) from the Spanish Ministry of Science and Innovation (co-financed by the European Regional Development Fund). RD, APR, RAHR, and GPR acknowledge each a Marie Curie fellowship. LG was supported by a predoctoral fellowship from the Valencia Regional Government (ACIF/2021/183).

AUTHORS' CONTRIBUTION

GR designed the research. RD, MHH, RMM, and ARM made the genetic constructions and performed the functional characterization experiments under the supervision of GR. APR purified the Musashi-1 protein under the supervision of TM. RAHR performed the binding kinetics assays under the supervision of WK. RD and GPR performed the dynamic analysis under the supervision of JB and GR. LG developed the mathematical models under the supervision of GR. RD, MHH, LG, RR, and GR analyzed the data and prepared the figures with input of RAHR, GPR, WFV, WK, and JB. GR wrote the manuscript with input of RD, MHH, LG, and WFV. All authors revised the manuscript.

COMPETING INTERESTS

RAHR and WFV work for Dynamic Biosensors. GPR and JB work for Ridgeview Instruments.

REFERENCES

1. Waters LS, Storz G (2009) Regulatory RNAs in bacteria. *Cell* 136: 615-628.

2. Holmqvist E, Vogel J (2018) RNA-binding proteins in bacteria. *Nat Rev Microbiol* 16: 601-615.

3. Jonas S, Izaurralde E (2015) Towards a molecular understanding of microRNA-mediated gene silencing. *Nat Rev Genet* 16: 421-433.

4. Glisovic T, Bachorik JL, Yong J, Dreyfuss G (2008) RNA-binding proteins and post-transcriptional gene regulation. *FEBS Lett* 582: 1977-1986.

5. Babu MM, Teichmann SA, Aravind L (2006) Evolutionary dynamics of prokaryotic transcriptional regulatory networks. *J Mol Biol* 358: 614-633.

6. Jacob F, Monod J (1961) Genetic regulatory mechanisms in the synthesis of proteins. *J Mol Biol* 3: 318-356.

7. Maris C, Dominguez C, Allain FHT (2005) The RNA recognition motif, a plastic RNA-binding platform to regulate post-transcriptional gene expression. *FEBS J* 272: 2118-2131.

8. Messias AC, Sattler M (2004) Structural basis of single-stranded RNA recognition. *Acc Chem Res* 37: 279-287.

9. Maruyama K, Sato N, Ohta N (1999) Conservation of structure and cold-regulation of RNA-binding proteins in cyanobacteria: probable convergent evolution with eukaryotic glycine-rich RNA-binding proteins. *Nucleic Acids Res* 27: 2029-2036.

10. Khalil AS, Collins JJ (2010) Synthetic biology: applications come of age. *Nat Rev Genet* 11: 367-379.

11. Bashor CJ, Collins JJ (2018) Understanding biological regulation through synthetic biology. *Annu Rev Biophys* 47: 399-423.

12. Belmont BJ, Niles JC (2010) Engineering a direct and inducible protein-RNA interaction to regulate RNA biology. *ACS Chem Biol* 5: 851-861.

13. Katz N, Cohen R, Solomon O, Kaufmann B, Atar O, Yakhini Z, Goldberg S, Amit R

(2019) Synthetic 5' UTRs can either up- or downregulate expression upon RNA-

binding protein binding. *Cell Syst* 9: 93-106.

14. Cao J, Arha M, Sudrik C, Mukherjee A, Wu X, Kane RS (2015) A universal strategy for

regulating mRNA translation in prokaryotic and eukaryotic cells. *Nucleic Acids Res* 43:

4353-4362.

15. Qi LS, Arkin AP (2014) A versatile framework for microbial engineering using synthetic

non-coding RNAs. *Nat Rev Microbiol* 12: 341-354.

16. Rosado A, Cordero T, Rodrigo G (2018) Binary addition in a living cell based on

riboregulation. *PLoS Genet* 14: e1007548.

17. Shotwell CR, Cleary JD, Berglund JA (2020) The potential of engineered eukaryotic

RNA-binding proteins as molecular tools and therapeutics. *Wiley Interdiscip Rev RNA*

11: e1573.

18. Liu MY, Romeo T (1997) The global regulator CsrA of Escherichia coli is a specific

mRNA-binding protein. *J Bacteriol* 179: 4639-4642.

19. Meyer MM (2018) rRNA mimicry in RNA regulation of gene expression. *Microbiol*

Spectr 6: RWR-0006-2017.

20. Babitzke P, Baker CS, Romeo T (2009) Regulation of translation initiation by RNA

binding proteins. *Annu Rev Microbiol* 63: 27-44.

21. Fox RG, Park FD, Koechlein CS, Kritzik M, Reya T (2015) Musashi signaling in stem

cells and cancer. *Annu Rev Cell Dev Biol* 31: 249-267.

22. Nakamura M, Okano H, Blendy JA, Montell C (1994) Musashi, a neural RNA-binding

protein required for Drosophila adult external sensory organ development. *Neuron* 13:

67-81.

23. Imai T, Tokunaga A, Yoshida T, Hashimoto M, Mikoshiba K, Weinmaster G, Nakafuku

M, Okano H (2001) The neural RNA-binding protein Musashi1 translationally regulates

mammalian numb gene expression by interacting with its mRNA. *Mol Cell Biol* 21: 3888-3900.

24. Clingman CC, Deveau LM, Hay SA, Genga RM, Shandilya SMD, Massi F, Ryder SP (2014) Allosteric inhibition of a stem cell RNA-binding protein by an intermediary metabolite. *eLife* 3: e02848.

25. Ganesan SM, Falla A, Goldfless SJ, Nasamu AS, Niles JC (2016) Synthetic RNA-protein modules integrated with native translation mechanisms to control gene expression in malaria parasites. *Nat Commun* 7: 10727.

26. MacDonald CI, Seamons TR, Emmons JC, Javdan SB, Deans TL (2021) Enhanced regulation of prokaryotic gene expression by a eukaryotic transcriptional activator. *Nat Commun* 12: 4109.

27. Iwaoka R, Nagata T, Tsuda K, Imai T, Okano H, Kobayashi N, Katahira M (2017) Structural insight into the recognition of r(UAG) by Musashi-1 RBD2, and construction of a model of Musashi-1 RBD1-2 bound to the minimum target RNA. *Molecules* 22: 1207.

28. Lutz R, Bujard H (1997) Independent and tight regulation of transcriptional units in Escherichia coli via the LacR/O, the TetR/O and AraC/I1-I2 regulatory elements. *Nucleic Acids Res* 25: 1203-1210.

29. Pédelacq JD, Cabantous S, Tran T, Terwilliger TC, Waldo GS (2006) Engineering and characterization of a superfolder green fluorescent protein. *Nat Biotechnol* 24: 79-88.

30. Kawahara H, Imai T, Imataka H, Tsujimoto M, Matsumoto K, Okano H (2008) Neural RNA-binding protein Musashi1 inhibits translation initiation by competing with eIF4G for PABP. *J Cell Biol* 181: 639-653.

31. Ai HW, Shaner NC, Cheng Z, Tsien RY, Campbell RE (2007) Exploration of new chromophore structures leads to the identification of improved blue fluorescent proteins. *Biochemistry* 46: 5904-5910.

32. Bindels DS, Haarbosch L, Van Weeren L, Postma M, Wiese KE, Mastop M, Aumonier
S, Gotthard G, Royant A, Hink MA, Gadella TWJ Jr (2017) mScarlet: a bright
monomeric red fluorescent protein for cellular imaging. *Nat Methods* 14: 53-56.

33. Klumpp S, Zhang Z, Hwa T (2009) Growth rate-dependent global effects on gene
expression in bacteria. *Cell* 139: 1366-1375.

34. Hausser J, Mayo A, Keren L, Alon U (2019) Central dogma rates and the trade-off
between precision and economy in gene expression. *Nat Commun* 10: 68.

35. Zearfoss NR, Deveau LM, Clingman CC, Schmidt E, Johnson ES, Massi F, Ryder SP
(2014) A conserved three-nucleotide core motif defines Musashi RNA binding
specificity. *J Biol Chem* 289: 35530-35541.

36. Cléry A, Sohier TJ, Welte T, Langer A, Allain FH (2017) switchSENSE: A new
technology to study protein-RNA interactions. *Methods* 118: 137-145.

37. Dolcemascolo R, Goiriz L, Montagud-Martinez R, Rodrigo G (2022) Gene regulation
by a protein translation factor at the single-cell level. *PLoS Comput Biol* 18: e1010087.

38. Buenrostro JD, Araya CL, Chircus LM, Layton CJ, Chang HY, Snyder MP, Greenleaf
WJ (2014) Quantitative analysis of RNA-protein interactions on a massively parallel
array reveals biophysical and evolutionary landscapes. *Nat Biotechnol* 32: 562-568.

39. Salis HM, Mirsky EA, Voigt CA (2009) Automated design of synthetic ribosome binding
sites to control protein expression. *Nat Biotechnol* 27: 946-50.

40. Bernstein JA, Khodursky AB, Lin PH, Lin-Chao S, Cohen SN (2002) Global analysis of
mRNA decay and abundance in Escherichia coli at single-gene resolution using two-
color fluorescent DNA microarrays. *Proc Natl Acad Sci USA* 99: 9697-9702.

41. Garcia HG, Phillips R (2011) Quantitative dissection of the simple repression input-
output function. *Proc Natl Acad Sci USA* 108: 12173-12178.

42. Goiriz L, Rodrigo G (2021) Nonequilibrium thermodynamics of the RNA-RNA
interaction underlying a genetic transposition program. *Phys Rev E* 103: 042410.

43. Kohler R, Mooney RA, Mills DJ, Landick R, Cramer P (2017) Architecture of a

transcribing-translating expressome. *Science* 356: 194-197.

44. Björke H, Andersson K (2006) Measuring the affinity of a radioligand with its receptor

using a rotating cell dish with in situ reference area. *Appl Radiat Isot* 64: 32-37.

45. Rosenfeld N, Alon U (2003) Response delays and the structure of transcription

networks. *J Mol Biol* 329: 645-654.

46. Paulus M, Haslbeck M, Watzele M (2004) RNA stem-loop enhanced expression of

previously non-expressible genes. *Nucleic Acids Res* 32: e78.

47. Fujita Y, Matsuoka H, Hirooka K (2007) Regulation of fatty acid metabolism in

bacteria. *Mol Microbiol* 66: 829-839.

48. Navarro-Llorens JM, Tormo A, Martínez-García E (2010) Stationary phase in gram-

negative bacteria. *FEMS Microbiol Rev* 34: 476-495.

49. Nielsen AAK, Der BS, Shin J, Vaidyanathan P, Paralanov V, Strychalski EA, Ross D,

Densmore D, Voigt CA (2016) Genetic circuit design automation. *Science* 352:

aac7341.

50. Sanchez A, Choubey S, Kondev J (2013) Regulation of noise in gene expression.

Annu Rev Biophys 42: 469-491.

51. Hammar P, Leroy P, Mahmutovic A, Marklund EG, Berg OG, Elf J (2012) The lac

repressor displays facilitated diffusion in living cells. *Science* 336: 1595-1598.

52. Järvelin AI, Noerenberg M, Davis I, Castello A (2016) The new (dis)order in RNA

regulation. *Cell Commun Signal* 14: 9.

53. Koonin EV, Makarova KS (2013) CRISPR-Cas: evolution of an RNA-based adaptive

immunity system in prokaryotes. *RNA Biol* 10: 679-686.

54. Anantharaman V, Iyer LM, Aravind L (2010) Presence of a classical RRM-fold palm

domain in Thg1-type 3'-5'nucleic acid polymerases and the origin of the GGDEF and

CRISPR polymerase domains. *Biol Direct* 5: 43.

55. Koonin EV, Krupovic M, Ishino S, Ishino Y (2020) The replication machinery of LUCA:

common origin of DNA replication and transcription. *BMC Biol* 18: 61.

56. Choi KR, Jang WD, Yang D, Cho JS, Park D, Lee SY (2019) Systems metabolic

engineering strategies: integrating systems and synthetic biology with metabolic

engineering. *Trends Biotechnol* 37: 817-837.

57. Na D, Yoo SM, Chung H, Park H, Park JH, Lee SY (2013) Metabolic engineering of

Escherichia coli using synthetic small regulatory RNAs. *Nat Biotechnol* 31: 170-174.

58. Zhang F, Carothers JM, Keasling JD (2012) Design of a dynamic sensor-regulator

system for production of chemicals and fuels derived from fatty acids. *Nat Biotechnol*

30: 354-359.

59. Taylor ND, Garruss AS, Moretti R, Chan S, Arbing MA, Cascio D, Rogers JK, Isaacs

FJ, Kosuri S, Baker D, Fields S, Church GM, Raman S (2016) Engineering an

allosteric transcription factor to respond to new ligands. *Nat Methods* 13: 177-183.

60. Montalbano M, McAllen S, Puangmalai N, Sengupta U, Bhatt N, Johnson OD, Kharas

MG, Kayed R (2020) RNA-binding proteins Musashi and tau soluble aggregates

initiate nuclear dysfunction. *Nat Commun* 11: 4305.

61. Kang MH, Jeong KJ, Kim WY, Lee HJ, Gong G, Suh N, Gyorffy B, Kim S, Jeong SY,

Mills GB, Park YY (2017) Musashi RNA-binding protein 2 regulates estrogen receptor

1 function in breast cancer. *Oncogene* 36: 1745-1752.

62. Sahdev S, Khattar SK, Saini KS (2008) Production of active eukaryotic proteins

through bacterial expression systems: a review of the existing biotechnology

strategies. *Mol Cell Biochem* 307: 249-264.

63. Valderrama-Rincon JD, Fisher AC, Merritt JH, Fan YY, Reading CA, Chhiba K, Heiss

C, Azadi P, Aebi M, DeLisa MP (2012) An engineered eukaryotic protein glycosylation

pathway in Escherichia coli. *Nat Chem Biol* 8: 434-436.

64. Peterson J, Philips GJ (2008) New pSC101-derivate cloning vectors with elevated

copy numbers. *Plasmid* 59: 193-201.

65. Leveau JH, Lindow SE (2001) Predictive and interpretive simulation of green

fluorescent protein expression in reporter bacteria. *J Bacteriol* 183: 6752-6762.

66. Langer A, Hampel P, Kaiser W, Knezevic J, Welte T, Villa V, Maruyama M, Svejda M,

Jahner S, Fischer F, Strasser R, Rant U (2013) Protein analysis by time-resolved

measurements with an electro-switchable DNA chip. *Nat Commun* 4: 2099.

67. Perea W, Greenbaum NL (2020) Label-free horizontal EMSA for analysis of protein-

RNA interactions. *Anal Biochem* 599: 113736.

68. Fessenden-Raden JM (1972) Effect of fatty acids on the movement and staining of

membrane proteins in polyacrylamide gel electrophoresis. *Biochem Biophys Res*

Commun 46: 1347-1353.

69. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch

S, Rueden C, Saalfeld S, Schmid B, Tinevez JY, White DJ, Hartenstein V, Eliceiri K,

Tomancak P, Cardona A (2012) Fiji: an open-source platform for biological-image

analysis. *Nat Methods* 9: 676-682.

70. Bairoch A, Apweiler R, Wu CH, Barker WC, Boeckmann B, Ferro S, Gasteiger E,

Huang H, Lopez R, Magrane M, Martin MJ, Natale DA, O'Donovan C, Redaschi N,

Yeh LS (2005) The Universal Protein Resource (UniProt). *Nucleic Acids Res* 33 D154-

D159.

71. Popenda M, Szachniuk M, Antczak M, Purzycka KJ, Lukasiak P, Bartol N, Blazewicz

J, Adamiak RW (2012) Automated 3D structure composition for large RNAs. *Nucleic*

Acids Res 40: e112.

 FIGURE LEGENDS

 Figure 1: Musashi-1 can down-regulate translation in bacteria.

a) Overview of the biotechnological development. In mammals, MSI-1 binds to the 3' UTR of its target mRNA to repress translation. Here, the *M. musculus* gene coding for MSI-1 was moved to *E. coli* (transgenesis) to implement a synthetic regulation system at the level of translation. **b)** Schematic of the synthetic gene circuit engineered in *E. coli*. A truncated version of MSI-1 (termed MSI-1*) was expressed from the PLlac promoter to be induced with lactose (or IPTG) in a genetic background over-expressing LacI. sfGFP was used as a reporter expressed from a constitutive promoter (J23119) and under the control of a suitable RNA motif recognized by MSI-1* in the N-terminal coding region of the transcript (*viz.*, located after the start codon). The activity of MSI-1* could in turn be allosterically inhibited by oleic acid. In electronic terms, this circuit implements an IMPLY logic gate. The inset shows the predicted secondary structure of the N-terminal coding region of the reporter mRNA. Within the motif (blue shaded), the consensus recognition sequences (RU$_n$AGU) are bolded and the minimal cores (UAG) are marked in red. System implemented with pRM1+ and pREP6. **c)** Dose-response curve of the system using lactose as inducer (up to 1 mM). MSI-1* down-regulated sfGFP expression by 2.5-fold. The inset shows the dynamic range of the response using lactose or IPTG (1 mM), showing a statistically significant regulation in both cases (Welch's *t*-test, two-tailed *P* < 0.05). **d)** Transfer function of the system (between sfGFP and MSI-1*). The inset shows the dose-response curve of eBFP2 expressed from the PLlac promoter (proxy of MSI-1* expression) with lactose. **e)** Scatter plot of the dynamic response of the system in the Crick space (translation rate *vs.* transcription rate). The dose-response curve of mScarlet expressed from the J23119 promoter with lactose was used to perform the decomposition (vertical line fitted to 48 AU/h). The inset shows the growth rate of the cells for each induction condition (horizontal line fitted to 0.55 h^{-1}). In all cases, points correspond to

experimental data, while solid lines come from adjusted mathematical models. Error bars

correspond to standard deviations (n = 3). **f)** Probability-based histograms of sfGFP

expression from single-cell data for different lactose concentrations, showing a statistically

significant regulation (one-way ANOVA test, $P < 10^{-4}$). The inset shows the percentage of

cells in the ON state (sfGFP expressed), according to a specified threshold, for each

lactose concentration. AU, arbitrary units.

Figure 2: Mechanistic characterization of the Musashi-1-mRNA interaction.

a) Sequences and predicted secondary structures of the different RNA motif variants for MSI-1 binding analyzed in this work. Point-mutations indicated in red. Three-dimensional representations of the RRM1 and RNA motif are also shown. Within the RRM1, the region that recognizes the RNA is shown in blue. **b)** Dynamic range of the response of the different genetic systems using lactose (1 mM), showing a statistically significant regulation in all cases (Welch's t-test, two-tailed $P < 0.05$; although some mutants present a small fold change). **c)** Schematic of the heliX biosensor platform. A double-strand DNA nanolever was immobilized on a gold electrode of the chip. The nanolever carried a fluorophore in one end and the RNA motif for MSI-1 binding in the other. Binding between MSI-1$_h$* (injected analyte) and RNA led to a fluorescence change, whose monitoring in real time served to extract the kinetic constants that characterize the interaction. **d)** Scatter plot of the experimentally-determined kinetic constants of association and dissociation between the protein and the RNA for all systems (original and 5 mutants). Means and deviations calculated in log scale (geometric). **e)** Correlation between the maximal sfGFP expression level (in absence of lactose) and the translation rate predicted with RBS calculator. Linear regression performed. **f)** Correlation between the fold change in sfGFP expression and the dissociation constant (K_D). Deviations calculated by propagation. Linear regression performed ($vs.$ 1/K_D). Blue shaded areas indicate 95% confidence intervals. In all cases, error bars correspond to standard deviations ($n = 3$). AU, arbitrary units.

 Figure 3: A mathematical model captures the dynamic response of the system.

a) Schematics of gene regulation at different levels with proteins that bind to nucleic acids (DNA or RNA). On the left, schematic of transcription regulation (*e.g.*, LacI regulating MSI-1* expression). On the right, schematic of translation regulation (*e.g.*, MSI-1* regulating sfGFP expression). A general mathematical expression (grey shaded) was derived to calculate the fold change in protein expression as a function of the regulator concentration (R), the association and dissociation rates (k_{ON} and k_{OFF}), the elongation leakage fraction (ε), and the nucleic acid degradation rate (δ). **b)** Heatmap of the fold change as a function of $k_{ON}R$ and k_{OFF} (*i.e.*, the first-order kinetic rates that characterize the protein-DNA/RNA interaction) when $\delta = 0$ and $\varepsilon = 0.1$. This would correspond to transcription regulation. **c)** Heatmap of the fold change when $\delta = 0.14$ min^{-1} and $\varepsilon = 0.1$. This would correspond to translation regulation. **d)** Total red fluorescence of the cell population (ΣmScarlet) over time without and with 1 mM lactose. In this case, the cell growth rate was fitted to 0.80 h^{-1}. **e)** Total green fluorescence of the cell population (ΣsfGFP) over time without and with 1 mM lactose. The inset shows the dynamic response for different lactose concentrations. **f)** Correlation between the experimental values of ΣsfGFP at different times and for different lactose concentrations and the predicted values from a mathematical model that accounts for population growth and gene regulation. Data for $t > 2$ h. Linear regression performed. **g)** Ratio of total green and red fluorescence as a proxy of cellular sfGFP expression over time. Ratio not represented at early times due to the high error obtained given the low number of cells present in the culture (grey shaded area). Deviations calculated by propagation. In all cases, points correspond to experimental data, while solid lines come from an adjusted mathematical model. Error bars correspond to standard deviations ($n = 3$). AU, arbitrary units.

Figure 4: mRNA redesign to enhance the down-regulation by Musashi-1.

a) Dynamic range of the response of three redesigned genetic systems using lactose (1 mM), showing a statistically significant regulation in the first and third cases (Welch's t-test, two-tailed $P < 0.05$). The predicted secondary structures of the N-terminal coding regions of the reporter mRNAs are shown on the right. Redesign 1 (red1) was implemented with pREP4b and redesign 2 (red2) with pREP4b3x, which contains three MSI-1 binding sites. These stem-loop structures are less stable than the original one. Redesign 3 (red3) was implemented with pREP7. **b)** Dose-response curve of the redesign-3 system using lactose as inducer (up to 1 mM). MSI-1* down-regulated sfGFP expression by 8.6-fold (Welch's t-test, two-tailed $P < 0.05$). The inset shows the scatter plot of the dynamic response in the Crick space (translation rate $vs.$ transcription rate; vertical line fitted to 27 AU/h). The predicted secondary structure of the N-terminal coding region of the reporter mRNA is shown on the right; the mRNA contains two MSI-1 binding sites (blue shaded). In the 5' UTR, the binding site is formed by two RU_nAGU repeats that flank the RBS without forming secondary structure. In the N-terminal coding region, the binding site is the original one. The minimal cores (UAG) are marked in red. Points correspond to experimental data, while the solid line comes from an adjusted mathematical model. In all cases, error bars correspond to standard deviations ($n = 3$). **c)** Probability-based histograms of sfGFP expression from single-cell data for different lactose concentrations (redesign 3), showing a statistically significant regulation (one-way ANOVA test, $P < 10^{-4}$). The inset shows the percentage of cells in the ON state (sfGFP expressed), according to a specified threshold, for each lactose concentration. AU, arbitrary units.

Figure 5: Oleic acid inhibits the regulatory activity of Musashi-1 in bacteria.

a) Three-dimensional structural schematic of the allosteric regulation. RRM1 of MSI-1 is shown alone, in complex with the RNA motif, and in complex with oleic acid. **b)** Gel electrophoresis mobility shift assay to test the allosteric inhibition of MSI-1* with oleic acid. A purified MSI-1* protein (45 µM), the RNA motif as a label-free sRNA molecule (11 µM), and oleic acid (1 mM) were mixed in a combinatorial way *in vitro*. On the left, nucleic acid-stained gel. On the right, protein-stained gel (Coomassie). The different formed species are indicated. M denotes molecular marker (GeneRuler ultra-low range DNA ladder, 10-300 bp, Thermo). BSA was used as a control. **c, d)** Probability-based histograms of sfGFP expression from single-cell data for different induction conditions (1 mM lactose or 1 mM lactose + 20 mM oleic acid) for the original system (c) and the redesign-3 system (d), showing statistically significant regulation in both cases (one-way ANOVA test, $P < 10^{-4}$). The insets show the percentages of cells in the ON state (sfGFP expressed), according to a specified threshold, for each condition. **e)** On the top, images of *E. coli* colonies harboring pRM1+ and pREP6. Bacteria were seeded in LB-agar plates with suitable inducers (1 mM lactose or 1 mM lactose + 20 mM oleic acid). Fluorescence and bright field images are shown. On the bottom, schematics of the working modes of the synthetic gene circuit according to the different induction conditions. **f)** Images of *E. coli* colonies harboring pRM1+ and pREP7. AU, arbitrary units.

Figure 6: Applications of Musashi-1 for a fine expression control in bacteria.

a) Overview of the regulatory utility of MSI-1*. It could *i)* regulate the expression of a given enzyme belonging to a polycistronic operon for a metabolic pathway control, *ii)* be exploited together with transcription factors to implement combinatorial regulations following the genetic information flow, envisioning biosensing applications, and *iii)* regulate noise in protein expression with the aim of producing cell populations less disperse, especially for bacterial delivery applications in animals. **b)** Schematic of a new synthetic gene circuit engineered in *E. coli*. MSI-1* was always expressed from the PLlac promoter to be induced with lactose in a genetic background over-expressing LacI. sfGFP and mScarlet were used as reporters, both expressed from the PLtet promoter to be induced with aTC in a genetic background over-expressing TetR. In this bicistronic operon, only sfGFP was under the control of a suitable RNA motif recognized by MSI-1* in the leader region of the transcript (original motif or redesign 3). In electronic terms, this circuit implements a NIMPLY logic gate considering sfGFP as the output. System implemented with pRM1+ and pREP6α or pREP7α. **c)** Dynamic range of the response using lactose (1 mM) and aTC (100 ng/mL) in a combinatorial way. aTC significantly activated the expression of the operon (Welch's *t*-test, two-tailed $P < 0.05$) and lactose, through the action of MSI-1*, significantly down-regulated sfGFP expression in a specific way (data for pREP7α; Welch's *t*-test, two-tailed $P < 0.05$). Error bars correspond to standard deviations (n = 3). **d)** Probability-based histograms of sfGFP expression from single-cell data for different inducer concentrations. On the left, pRM1+ and pREP6α with 100 ng/mL aTC and 1 mM lactose (*i*) or 15 ng/mL aTC (*ii*). On the right, pRM1+ and pREP7α with 100 ng/mL aTC and 1 mM lactose (*iii*) or 30 ng/mL aTC (*iv*). The mean expression and Fano factor are shown. AU, arbitrary units.

a

$$\text{fold change} = \frac{k_{ON}R + k_{OFF} + \delta}{\varepsilon k_{ON}R + k_{OFF} + \delta} \approx \begin{cases} 1 + \dfrac{R}{K_D}, & \delta = 0, & \varepsilon = 0 \\ 1 + \dfrac{k_{ON}R}{\delta}, & \delta \gg k_{OFF}, & \varepsilon = 0 \\ \dfrac{1}{\varepsilon}, & R \to \infty \end{cases}$$

R: repressor concentration

b $\delta = 0, \quad \varepsilon = 0.1$

c $\delta = 0.14\ \text{min}^{-1}, \quad \varepsilon = 0.1$

d

e lactose (μM)

f $R^2 = 0.98$

g

a

RRM1 (MSI-1*) → 180° → RRM1–RNA → 180° → RRM1–oleic acid

RNA (motif) · oleic acid · allosterically inhibited protein

b

nucleic acid-stained gel

RNA / MSI-1ₕ* / oleic acid

M   +/−/−   +/+/−   +/+/+   −/+/+   −/−/+   BSA

oleic acid
MSI-1ₕ*-oleic acid
MSI-1ₕ*-RNA
differently folded RNA

300
150
50
20
(bp)

protein-stained gel

RNA / MSI-1ₕ* / oleic acid

M   +/−/−   +/+/−   +/+/+   −/+/+   −/−/+   BSA

MSI-1ₕ*

c

OFF / ON threshold (orig)

probability (%)

sfGFP (AU)

lactose / oleic acid

% cells ON

lactose / oleic acid

d

OFF / ON threshold (red3)

probability (%)

sfGFP (AU)

lactose / oleic acid

% cells ON

lactose / oleic acid

e

none   lactose   lactose, oleic acid

ON   OFF   ON

f

none   lactose   lactose, oleic acid